



# REPRESENTATIONS OF BRAID GROUPS AND GENERALISATIONS

Valerij G. Bardakov, Paolo Bellingeri

► **To cite this version:**

Valerij G. Bardakov, Paolo Bellingeri. REPRESENTATIONS OF BRAID GROUPS AND GENERALISATIONS. 2007. hal-00138067v1

**HAL Id: hal-00138067**

**<https://hal.science/hal-00138067v1>**

Preprint submitted on 23 Mar 2007 (v1), last revised 19 Apr 2007 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# REPRESENTATIONS OF BRAID GROUPS AND GENERALISATIONS

VALERIJ G. BARDAKOV AND PAOLO BELLINGERI

**ABSTRACT.** We define and study extensions of Artin's representation and braid monodromy representation to the case of topological and algebraical generalisations of braid groups. In particular we provide faithful representations of braid groups of oriented surfaces with boundary components as (outer) automorphisms of free groups. We give also similar representations for braid groups of non oriented surfaces with boundary components and we show a representation of braid groups of closed surfaces as outer automorphisms of free groups. Finally, we provide faithful representations of Artin-Tits groups of type  $\mathcal{D}$  as automorphisms of free groups.

## 1. INTRODUCTION

Let  $F_n$  be the free group of rank  $n$  with the set of generators  $\{x_1, x_2, \dots, x_n\}$ . Assume further that  $\text{Aut}(F_n)$  is the automorphism group of  $F_n$ . The Artin braid group  $B_n$  can be represented as a subgroup of  $\text{Aut}(F_n)$ . This representation, due to Artin himself, is defined associating to any generator  $\sigma_i$ , for  $i = 1, 2, \dots, n-1$ , of  $B_n$  the following automorphism of  $F_n$ :

$$\sigma_i : \begin{cases} x_i \mapsto x_i x_{i+1} x_i^{-1}, \\ x_{i+1} \mapsto x_i, \\ x_l \mapsto x_l, \end{cases} \quad l \neq i, i+1.$$

Moreover (see for instance [10, Theorem 5.1]), any automorphism  $\beta$  of  $\text{Aut}(F_n)$  corresponds to an element of  $B_n$  if and only if  $\beta$  satisfies the following conditions:

- i)  $\beta(x_i) = a_i^{-1} x_{s(i)} a_i, \quad 1 \leq i \leq n,$
- ii)  $\beta(x_1 x_2 \dots x_n) = x_1 x_2 \dots x_n,$

where  $s$  is a permutation from the symmetric group  $S_n$  and  $a_i \in F_n$ .

Generalisations of Artin's representation have been provided by Wada [13] and further by Crisp and Paris [5], in order to construct group invariants of oriented links.

Another interesting representation of  $B_n$  is the *braid monodromy representation* of  $B_n$  into  $\text{Aut}(F_{n-1})$  (see last Section), that was proven to be faithful in [6] and [12].

In this paper, we extend Artin's and braid monodromy representations to some generalisations of braid groups.

---

*Date:* March 23, 2007.

*1991 Mathematics Subject Classification.* Primary 20F36; Secondary 20F05, 20F10.

*Key words and phrases.* Braid groups, free groups, automorphisms.

In the case of braid groups of oriented surfaces with boundary components our representations are faithful (Theorem 2). Moreover, in the case of surfaces of genus  $g \geq 1$ , the induced representations of surface braid groups as outer automorphisms hold faithful (Theorem 3).

In the case of closed surfaces, as earlier remarked in [4] for braid groups of the sphere, we cannot extend Artin's representation and we provide a representation in the outer automorphism group of a finitely generated free group.

In the last Section we consider the braid monodromy representation of  $B_n$  in  $\text{Aut}(F_{n-1})$  and we provide a faithful representation of the  $n$ -th Artin-Tits group of type  $\mathcal{D}$  in  $\text{Aut}(F_{n-1})$  (Proposition 10).

**Acknowledgments.** The research of the first author has been supported by the Scientific and Research Council of Turkey (TÜBİTAK). The first author would like to thank Prof. Hasan Gümral, Prof. Vladimir Tolstykh and other members of the Department of Mathematics of Yeditepe University (Istanbul) for their kind hospitality.

## 2. BRAID GROUPS OF ORIENTABLE SURFACES WITH BOUNDARY COMPONENTS

**Surface braids as collections of paths.** Let  $\mathcal{P} = \{p_1, \dots, p_n\}$  be a set of  $n$  distinct points (punctures) in the interior of a surface  $\Sigma$ .

A *geometric braid* on  $\Sigma$  based at  $\mathcal{P}$  is a collection  $(\psi_1, \dots, \psi_n)$  of  $n$  disjoint paths (called *strands*) on  $\Sigma \times [0, 1]$  which run monotonically with  $t \in [0, 1]$  and such that  $\psi_i(0) = (p_i, 0)$  and  $\psi_i(1) \in \mathcal{P} \times \{1\}$ . Two braids are considered to be equivalent if they are isotopic relatively to the base points. The usual product of paths defines a group structure on the equivalence classes of braids. This group, denoted usually by  $B_n(\Sigma)$ , does not depend on the choice of  $\mathcal{P}$  and it is called braid group on  $n$  strands of  $\Sigma$ . The  $n$ th braid group of the disk  $D^2$ ,  $B_n(D^2)$ , is isomorphic to  $B_n$ .

In the following we will denote by  $B_n(\Sigma_{g,p})$  the braid group on  $n$  strands of an orientable surface of genus  $g$  with  $p$  boundary components (we set  $\Sigma_g = \Sigma_{g,0}$ ) and by  $B_n(N_{g,p})$  the braid group on  $n$  strands of a non-orientable surface of genus  $g$  with  $p$  boundary components (we set  $N_g = N_{g,0}$ ).

In this section we will consider an orientable surface  $\Sigma_{g,p}$  of genus  $g \geq 0$  and with  $p > 0$  boundary components. We set also  $n \geq 2$ .

We denote by  $\sigma_1, \dots, \sigma_{n-1}$  the standard generators of the braid group  $B_n$ . Since  $p > 0$ , we can embed a disk in  $\Sigma_{g,p}$  and therefore we can consider  $\sigma_1, \dots, \sigma_{n-1}$  as elements of  $B_n(\Sigma_{g,p})$ . Let also  $a_1, \dots, a_g, b_1, \dots, b_g, z_1, \dots, z_{p-1}$  be the generators of  $\pi_1(\Sigma_{g,p})$ , where  $z_i$ 's denote loops around the holes. Assume that the base point of the fundamental group is the startpoint of the first strand. Then each element  $\gamma \in \pi_1(\Sigma_{g,p})$  determines an element denoted also by  $\gamma$  in  $B_n(\Sigma_{g,p})$ , by considering the braid whose first strand is describing the curve  $\gamma$  and other strands are constant.

Let us set  $x_{2k-1} = a_k$  and  $x_{2k} = b_k$  for  $k = 1, \dots, g$ . According to [1] we have that the group  $B_n(\Sigma_{g,p})$  admits a presentation with generators:

$$\sigma_1, \dots, \sigma_{n-1}, x_1, \dots, x_{2g}, z_1, \dots, z_{p-1},$$

and defining relations:

– Braid relations:

$$\begin{aligned} \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1}, \quad 1 \leq i \leq n-2, \\ \sigma_i \sigma_j &= \sigma_j \sigma_i, \quad |i-j| > 1, \quad 1 \leq i, j \leq n-1, \end{aligned}$$

– Mixed relations:

$$\begin{aligned} \text{(R1)} \quad x_r \sigma_i &= \sigma_i x_r, \quad i \neq 1, \quad 1 \leq r \leq 2g, \\ \text{(R2)} \quad (\sigma_1^{-1} x_r \sigma_1^{-1}) x_r &= x_r (\sigma_1^{-1} x_r \sigma_1^{-1}), \quad 1 \leq r \leq 2g, \\ \text{(R3)} \quad (\sigma_1^{-1} x_s \sigma_1) x_r &= x_r (\sigma_1^{-1} x_s \sigma_1), \quad 1 \leq s < r \leq 2g, \quad (s, r) \neq (2m-1, 2m), \\ \text{(R4)} \quad (\sigma_1^{-1} x_{2m-1} \sigma_1^{-1}) x_{2m} &= x_{2m} (\sigma_1^{-1} x_{2m-1} \sigma_1), \quad 1 \leq m \leq g, \\ \text{(R5)} \quad z_j \sigma_i &= \sigma_i z_j, \quad i \neq 1, \quad 1 \leq j \leq p-1, \\ \text{(R6)} \quad (\sigma_1^{-1} z_j \sigma_1) x_r &= x_r (\sigma_1^{-1} z_j \sigma_1), \quad 1 \leq r \leq 2g, \quad 1 \leq j \neq p-1, \\ \text{(R7)} \quad (\sigma_1^{-1} z_j \sigma_1) z_l &= z_l (\sigma_1^{-1} z_j \sigma_1), \quad 1 \leq j < l \leq p-1, \\ \text{(R8)} \quad (\sigma_1^{-1} z_j \sigma_1^{-1}) z_j &= z_j (\sigma_1^{-1} z_j \sigma_1), \quad 1 \leq j \leq p-1. \end{aligned}$$

Associating to any surface braid the corresponding permutation one obtains a surjective homomorphism

$$\pi : B_n(\Sigma_{g,p}) \longrightarrow S_n,$$

such that  $\pi(\sigma_i) = (i, i+1)$ ,  $i = 1, \dots, n-1$ ,  $\pi(x_r) = \pi(z_j) = e$  for  $1 \leq r \leq 2g$  and  $1 \leq j \leq p-1$ .

In [1] the second author considered the subgroup  $D_n(\Sigma_{g,p}) = \pi^{-1}(S_{n-1})$  and found its generators. We provide a set of defining relations of  $D_n(\Sigma_{g,p})$  using the well-known Reidemeister-Schreier's method (see [8, Chap. 2]).

Let  $M_n = \{m_l \mid 1 \leq l \leq n\}$  be the set defined as follows:

$$m_l = \sigma_{n-1} \dots \sigma_l, \quad l = 1, \dots, n-1, \quad m_n = 1.$$

It is easy to prove (see [1]) that  $|B_n(\Sigma_{g,p}) : D_n(\Sigma_{g,p})| = n$  and that  $M_n$  is a Schreier set of coset representatives of  $D_n(\Sigma_{g,p})$  in  $B_n(\Sigma_{g,p})$ . Define the map  $\bar{\cdot} : B_n(\Sigma_{g,p}) \longrightarrow M_n$  which takes an element  $w \in B_n(\Sigma_{g,p})$  into the representative  $\bar{w}$  from  $M_n$ . The element  $w\bar{w}^{-1}$  belongs to  $D_n(\Sigma_{g,p})$  and, by Theorem 2.7 from [8], the group  $D_n(\Sigma_{g,p})$  is generated by

$$s_{\lambda,a} = \lambda a \cdot (\overline{\lambda a})^{-1},$$

where  $\lambda$  runs over the set  $M_n$  and  $a$  runs over the set of generators of  $B_n(\Sigma_{g,p})$ .

**Case 1.** If  $a \in \{\sigma_1, \dots, \sigma_{n-1}\}$ , then we find the generators

$$\tau_k = \sigma_{n-1} \dots \sigma_{k+1} \sigma_k^2 \sigma_{k+1}^{-1} \dots \sigma_{n-1}^{-1}, \quad k = 1, \dots, n-2, \quad \tau_{n-1} = \sigma_{n-1}^2.$$

**Case 2.** If  $a \in \{x_1, \dots, x_{2g}\}$ , then we find the generators

$$w_r = \sigma_{n-1} \dots \sigma_1 x_r \sigma_1^{-1} \dots \sigma_{n-1}^{-1}, \quad r = 1, \dots, 2g.$$

**Case 3.** If  $a \in \{z_1, \dots, z_{p-1}\}$ , then we find the generators

$$\xi_j = \sigma_{n-1} \dots \sigma_1 z_j \sigma_1^{-1} \dots \sigma_{n-1}^{-1}, \quad j = 1, \dots, p-1.$$

To find defining relations of  $D_n(\Sigma_{g,p})$  we define a rewriting process  $\tau$ . It allows us to rewrite a word which is written in the generators of  $B_n(\Sigma_{g,p})$  and to present an element in  $D_n(\Sigma_{g,p})$  as a word in the generators of  $D_n(\Sigma_{g,p})$ . Let us associate to the reduced word

$$u = a_1^{\varepsilon_1} a_2^{\varepsilon_2} \dots a_\nu^{\varepsilon_\nu}, \quad \varepsilon_l = \pm 1,$$

where

$$a_l \in \{\sigma_1, \sigma_2, \dots, \sigma_{n-1}, x_1, x_2, \dots, x_{2g}, z_1, z_2, \dots, z_{p-1}\},$$

the word

$$\tau(u) = s_{k_1, a_1}^{\varepsilon_1} s_{k_2, a_2}^{\varepsilon_2} \dots s_{k_\nu, a_\nu}^{\varepsilon_\nu}$$

in the generators of  $D_n(\Sigma_{g,p})$ , where  $k_j$  is a representative of the  $(j-1)$ th initial segment of the word  $u$  if  $\varepsilon_j = 1$  and  $k_j$  is a representative of the  $j$ th initial segment of the word  $u$  if  $\varepsilon_j = -1$ . By [8, Theorem 2.9], the group  $D_n(\Sigma_{g,p})$  is defined by relations

$$r_{\mu, \lambda} = \tau(\lambda r_\mu \lambda^{-1}), \quad \lambda \in M_n,$$

where  $r_\mu$  is a defining relation of  $B_n(\Sigma_{g,p})$ .

**Proposition 1.** *The group  $D_n(\Sigma_{g,p})$  admits a presentation with the generators*

$\sigma_1, \dots, \sigma_{n-2}, x_1, \dots, x_{2g}, z_1, \dots, z_{p-1}, \tau_1, \dots, \tau_{n-1}, w_1, \dots, w_{2g}, \xi_1, \dots, \xi_{p-1};$   
and relations:

– *Braid relations*

- (B1)  $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \quad 1 \leq i \leq n-3,$
- (B2)  $\sigma_i \sigma_j = \sigma_j \sigma_i, \quad |i-j| > 1, \quad 1 \leq i, j \leq n-2,$
- (B3)  $\sigma_k^{-1} \tau_l \sigma_k = \tau_l, \quad k \neq l-1, l,$
- (B4)  $\sigma_{l-1}^{-1} \tau_l \sigma_{l-1} = \tau_{l-1},$
- (B5)  $\sigma_l^{-1} \tau_l \sigma_l = \tau_l \tau_{l+1} \tau_l^{-1}, \quad l \neq n-1.$

– *Mixed relations*

- (R1.1)  $x_r \sigma_i = \sigma_i x_r, \quad 2 \leq i \leq n-2, \quad 1 \leq r \leq 2g,$
- (R1.2)  $x_r \tau_i = \tau_i x_r, \quad 2 \leq i \leq n-1,$
- (R1.3)  $w_r \sigma_i = \sigma_i w_r, \quad 1 \leq r \leq 2g, \quad 1 \leq i \leq n-2,$
- (R2.1)  $(\sigma_1^{-1} x_r \sigma_1^{-1}) x_r = x_r (\sigma_1^{-1} x_r \sigma_1^{-1}), \quad 1 \leq r \leq 2g,$
- (R2.2)  $x_r^{-1} w_r x_r = \tau_1^{-1} w_r \tau_1, \quad 1 \leq r \leq 2g,$
- (R2.3)  $x_r^{-1} \tau_1 x_r = \tau_1^{-1} w_r \tau_1 w_r^{-1} \tau_1, \quad 1 \leq r \leq 2g,$
- (R3.1)  $(\sigma_1^{-1} x_s \sigma_1^{-1}) x_r = x_r (\sigma_1^{-1} x_s \sigma_1^{-1}), \quad 1 \leq s < r \leq 2g, \quad (s, r) \neq (2m-1, 2m),$
- (R3.2)  $x_r^{-1} (\tau_1^{-1} w_s \tau_1) x_r = \tau_1^{-1} w_s \tau_1, \quad 1 \leq s < r \leq 2g, \quad (s, r) \neq (2m-1, 2m),$
- (R3.3)  $x_s w_r = w_r x_s, \quad 1 \leq s < r \leq 2g, \quad (s, r) \neq (2m-1, 2m),$

- (R4.1)  $(\sigma_1^{-1}x_{2m-1}\sigma_1^{-1})x_{2m} = x_{2m}(\sigma_1^{-1}x_{2m-1}\sigma_1), 1 \leq m \leq g,$
- (R4.2)  $x_{2m}^{-1}(\tau_1^{-1}w_{2m-1})x_{2m} = \tau_1^{-1}w_{2m-1}\tau_1, 1 \leq m \leq g,$
- (R4.3)  $x_{2m-1}^{-1}w_{2m}x_{2m-1} = \tau_1^{-1}w_{2m}, 1 \leq m \leq g,$
- (R5.1)  $z_j\sigma_i = \sigma_i z_j, 2 \leq i \leq n-2, 1 \leq j \leq p-1,$
- (R5.2)  $z_j\tau_i = \tau_i z_j, 2 \leq i \leq n-1, 1 \leq j \leq p-1,$
- (R5.3)  $\xi_j\sigma_i = \sigma_i \xi_j, 1 \leq j \leq p-1, 1 \leq i \leq n-2,$
- (R6.1)  $(\sigma_1^{-1}z_j\sigma_1)x_r = x_r(\sigma_1^{-1}z_j\sigma_1), 1 \leq r \leq 2g, 1 \leq j \leq p-1,$
- (R6.2)  $x_r^{-1}(\tau_1^{-1}\xi_j\tau_1)x_r = \tau_1^{-1}\xi_j\tau_1, 1 \leq r \leq 2g, 1 \leq j \leq p-1,$
- (R6.3)  $z_jw_r = w_r z_j, 1 \leq r \leq 2g, 1 \leq j \leq p-1,$
- (R7.1)  $(\sigma_1^{-1}z_j\sigma_1)z_l = z_l(\sigma_1^{-1}z_j\sigma_1), 1 \leq j < l \leq p-1,$
- (R7.2)  $z_l^{-1}(\tau_1^{-1}\xi_j\tau_1)z_l = \tau_1^{-1}\xi_j\tau_1, 1 \leq j < l \leq p-1,$
- (R7.3)  $z_j\xi_l = \xi_l z_j, 1 \leq j < l \leq p-1,$
- (R8.1)  $(\sigma_1^{-1}z_j\sigma_1^{-1})z_j = z_j(\sigma_1^{-1}z_j\sigma_1^{-1}), 1 \leq j \leq p-1,$
- (R8.2)  $z_j^{-1}(\tau_1^{-1}\xi_j)z_j = \tau_1^{-1}\xi_j, 1 \leq j \leq p-1,$
- (R8.3)  $z_j^{-1}\xi_j z_j = \tau_1^{-1}\xi_j\tau_1, 1 \leq j \leq p-1.$

The generators  $\sigma_1, \dots, \sigma_{n-2}, x_1, \dots, x_{2g}, z_1, \dots, z_{p-1}$  generate a group isomorphic to  $B_{n-1}(\Sigma_{g,p})$  (see also Remark 3.1 from [1]) and it is easy to see that the relations (B1), (B2), (R1.1), (R2.1),  $\dots$ , (R8.1) are a complete set of relations for  $B_{n-1}(\Sigma_{g,p})$ . From the other relations we can find the following conjugacy formulae:

- (S1)  $\tau_l^{\sigma_k} = \tau_l, k \neq l-1, l,$
- (S2)  $\tau_l^{\sigma_{l-1}} = \tau_{l-1},$
- (S3)  $\tau_l^{\sigma_l} = \tau_{l+1}^{-1}, l \neq n-1,$
- (S4)  $\tau_i^{x_r} = \tau_i, 2 \leq i \leq n-1,$
- (S5)  $w_r^{\sigma_i} = w_r, 1 \leq r \leq 2g, 1 \leq i \leq n-2,$
- (S6)  $w_r^{x_r} = w_r^{\tau_1},$
- (S7)  $\tau_1^{x_r} = \tau_1^{w_r^{-1}\tau_1},$
- (S8)  $w_s^{x_r} = w_s^{[w_r^{-1}, \tau_1]}, 1 \leq s < r \leq 2g, (s, r) \neq (2m-1, 2m),$
- (S9)  $w_r^{x_s} = w_r, 1 \leq s < r \leq 2g, (s, r) \neq (2m-1, 2m),$
- (S10)  $w_{2m-1}^{x_{2m}} = [\tau_1, w_{2m}^{-1}]w_{2m-1}\tau_1,$
- (S11)  $w_{2m}^{x_{2m-1}} = \tau_1^{-1}w_{2m},$
- (S12)  $\tau_i^{z_j} = \tau_i, 2 \leq i \leq n-1, 1 \leq j \leq p-1,$
- (S13)  $\xi_j^{\sigma_i} = \xi_j, 1 \leq i \leq n-2, 1 \leq j \leq p-1,$
- (S14)  $\xi_j^{x_r} = \xi_j^{[w_r^{-1}, \tau_1]}, 1 \leq r \leq 2g, 1 \leq j \leq p-1,$
- (S15)  $w_r^{z_j} = w_r, 1 \leq r \leq 2g, 1 \leq j \leq p-1,$
- (S16)  $\xi_j^{z_l} = \xi_j^{[\xi_l^{-1}, \tau_1]}, 1 \leq j < l \leq p-1,$
- (S17)  $\xi_l^{z_j} = \xi_l, 1 \leq j < l \leq p-1,$
- (S18)  $\tau_1^{z_j} = [\tau_1, \xi_j^{-1}]\tau_1, 1 \leq j \leq p-1,$

(S19)  $\xi_j^{z_j} = \xi_j^{\tau_1}$ ,  $1 \leq j \leq p-1$ ,  
 where  $a^b = a^{-1}ba$  and  $[a, b] = a^{-1}b^{-1}ab$ .

Let  $U_{n-1,g,p}$  be the subgroup of  $D_n(\Sigma_{g,p})$  generated by  $\{\tau_1, \dots, \tau_{n-1}, w_1, \dots, w_{2g}, \xi_1, \dots, \xi_{p-1}\}$ .

**Proposition 2.** *The group  $U_{n-1,g,p}$  is a normal subgroup of  $D_n(\Sigma_{g,p})$  and it is a free group of rank  $n + p + 2g - 2$ .*

*Proof.* The statement was proven in [1, Section.2] using the interpretation of  $U_{n-1,g,p}$  as the fundamental group of the surface  $\Sigma_{g,p}$  with  $n-1$  points removed.  $\square$

From relations (S1)...(S19) we deduce that  $B_{n-1}(\Sigma_{g,p})$  acts on  $U_{n-1,g,p}$  by conjugacy.

**Theorem 1.** *The group  $B_{n-1}(\Sigma_{g,p})$  with  $n \geq 3$ ,  $g \geq 0$  and  $p > 0$  acts by conjugacy on the free group  $U_{n-1,g,p}$ . Therefore we have a representation  $\rho_U : B_{n-1}(\Sigma_{g,p}) \rightarrow \text{Aut}(U_{n-1,g,p})$  defined algebraically as follows:*

– Generators  $\sigma_i$ ,  $i = 1, \dots, n-2$ :

$$\sigma_i : \begin{cases} \tau_i \mapsto \tau_{i+1}^{\tau_i^{-1}} ; \\ \tau_{i+1} \mapsto \tau_i ; \\ \tau_l \mapsto \tau_l, l \neq i, i+1 ; \\ w_r \mapsto w_r, 1 \leq r \leq 2g ; \\ \xi_j \mapsto \xi_j, 1 \leq j \leq p-1. \end{cases}$$

– Generators  $x_r$ ,  $r = 1, \dots, 2g$ :

$$x_r : \begin{cases} \tau_1 \mapsto \tau_1^{w_r^{-1}\tau_1} ; \\ \tau_i \mapsto \tau_i, 2 \leq i \leq n ; \\ w_s \mapsto w_s^{[w_r^{-1}, \tau_1]}, s < r, (s, r) \neq (2m-1, 2m) ; \\ w_{r-1} \mapsto [\tau_1, w_r^{-1}]w_{r-1}\tau_1, \text{ if } r = 2m ; \\ w_r \mapsto w_r^{\tau_1} ; \\ w_s \mapsto w_s, r < s, (r, s) \neq (2m-1, 2m) ; \\ w_{r+1} \mapsto \tau_1^{-1}w_{r+1}, \text{ if } r = 2m-1 ; \\ \xi_j \mapsto \xi_j^{[w_r^{-1}, \tau_1]}, 1 \leq j \leq p-1. \end{cases}$$

– Generators  $z_j$ ,  $j = 1, \dots, p-1$ :

$$z_j : \begin{cases} \tau_1 \mapsto \tau_1^{\xi_j^{-1}\tau_1} ; \\ \tau_i \mapsto \tau_i, 2 \leq i \leq n ; \\ w_r^{z_j} \mapsto w_r, 1 \leq r \leq 2g ; \\ \xi_l^{z_j} \mapsto \xi_l^{[\xi_j^{-1}, \tau_1]}, 1 \leq l < j \leq p-1 ; \\ \xi_j^{z_j} \mapsto \xi_j^{\tau_1} ; \\ \xi_l^{z_j} \mapsto \xi_l, 1 \leq j < l \leq p-1. \end{cases}$$

In the following we outline a proof of the faithfulness of the representation  $\rho_U$  of  $B_{n-1}(\Sigma_{g,p})$  given in Theorem 1 using the interpretation of surface braids as mapping classes.

First, we recall that the mapping class group of a surface  $\Sigma_{g,p}$ , let us denote it by  $\mathcal{M}_{g,p}$ , is the group of isotopy classes of orientation-preserving self-homeomorphisms which fix the boundary components pointwise.

Let  $\mathcal{P} = \{p_1, \dots, p_n\}$  be a set of  $n$  distinct points (*punctures*) in the interior of the surface  $\Sigma_{g,p}$ . The punctured mapping class group of  $\Sigma_{g,p}$  relative to  $\mathcal{P}$  is defined to be the group of isotopy classes of orientation-preserving self-homeomorphisms which fix the boundary components pointwise, and which fix  $\mathcal{P}$  setwise. This group, that we will denote by  $\mathcal{M}_{g,p}^n$ , does not depend on the choice of  $\mathcal{P}$ , but just on its cardinal.

We recall also that a simple closed curve  $C$  is *essential* if either it does not bound a disk or it bounds a disk containing at least two punctures.

Finally, we denote  $T_C$  as a Dehn twist along a simple closed curve  $C$ . Let  $C$  and  $D$  be two simple closed curves bounding an annulus containing only the puncture  $p_j$ . We shall say that the multitwist  $T_C T_D^{-1}$  is a *j-bounding pair braid*, also called *spin map* in [4].

**Surface braids as mapping classes.** Let  $g, p \geq 0$  and let  $\psi_{n,0} : \mathcal{M}_{g,p}^n \rightarrow \mathcal{M}_{g,p}$  be the homomorphism induced by the map which forgets the set  $\mathcal{P}$ . When  $p = 0$ , according to a well-known result of Birman [4, Chapter 4.1], the group  $B_n(\Sigma_g)$  is isomorphic to  $\ker \psi_{n,0}$  if  $g > 1$ . The statement of Birman's theorem concerns the case of closed surfaces, but the proof extends naturally to the case of surfaces with boundary components and the group  $B_n(\Sigma_{g,p})$  is isomorphic to  $\ker \psi_{n,0}$  if  $g \geq 1$  and  $p > 0$ .

Geometrically the correspondence between  $\ker \psi_{n,0}$  and  $B_n(\Sigma_{g,p})$  is realized as follows: given a homeomorphism  $h$  of  $\Sigma_{g,p}$  isotopic to the identity  $Id$  of  $\Sigma_{g,p}$  and fixing boundary components pointwise and  $\mathcal{P}$  setwise, the track of the punctures  $p_1, \dots, p_n$  under an isotopy from  $h$  to  $Id$  is the geometric braid corresponding to the homeomorphism  $h$ .

**Theorem 2.** *Let  $g \geq 0$ ,  $p > 0$  and  $n \geq 3$ . The representation  $\rho_U : B_{n-1}(\Sigma_{g,p}) \rightarrow \text{Aut}(U_{n-1,g,p})$  is faithful.*

*Proof.* First we prove the claim when  $g$  is greater or equal then 1. According to Birman's result we can represent the group  $D_n(\Sigma_{g,p})$  as a (normal) subgroup of  $\mathcal{M}_{g,p}^n$ , more precisely as the subgroup of mapping classes in  $\ker \psi_{n,0}$  sending the  $n$ th puncture into itself (see also Remark 18 from [3]). In particular generators  $\tau_1, \dots, \tau_{n-1}$ ,  $w_1, \dots, w_{2g}$ ,  $\xi_1, \dots, \xi_{p-1}$ , of  $D_n(\Sigma_{g,p})$  (see Proposition 1) correspond to  $n$ -bounding pair braids in  $\mathcal{M}_{g,p}^n$ .

Let  $\psi_{n,n-1} : \mathcal{M}_{g,p}^n \rightarrow \mathcal{M}_{g,p}^{n-1}$  be the homomorphism forgetting the last puncture. Now, let us consider the exact sequence associate to the restriction of  $\psi_{n,n-1}$  to  $D_n(\Sigma_{g,p})$ . The image of  $D_n(\Sigma_{g,p})$  by  $\psi_{n,n-1}$  coincides with  $\ker \psi_{n-1,0}$  which is isomorphic to  $B_{n-1}(\Sigma_{g,p})$ . On the other hand,  $\ker \psi_{n,n-1} \cap D_n(\Sigma_{g,p})$  is the subgroup of  $\mathcal{M}_{g,p}^n$  generated by  $n$ -bounding pair braids  $\tau_1, \dots, \tau_{n-1}$ ,  $w_1, \dots, w_{2g}$ ,  $\xi_1, \dots, \xi_{p-1}$  and therefore it is isomorphic to  $U_{n-1,g,p}$ .

The group  $B_{n-1}(\Sigma_{g,p})$  embeds naturally in  $B_n(\Sigma_{g,p})$  by sending generators of  $B_{n-1}(\Sigma_{g,p})$  in corresponding ones of  $B_n(\Sigma_{g,p})$  and therefore  $B_{n-1}(\Sigma_{g,p})$  can be considered also as a subgroup of  $\mathcal{M}_{g,p}^n$ . Moreover, the action considered in Theorem 1 corresponds to the action by conjugacy of  $B_{n-1}(\Sigma_{g,p})$ , seen as a subgroup of  $\mathcal{M}_{g,p}^n$ , on  $U_{n-1,g,p}$ . Since  $U_{n-1,g,p}$  is a normal subgroup of  $\mathcal{M}_{g,p}^n$  we can define a map  $\Theta : \text{Inn}(\mathcal{M}_{g,p}^n) \rightarrow \text{Aut}(U_{n-1,g,p})$ . We prove that  $\Theta$  is injective and therefore that the action by conjugacy of  $B_{n-1}(\Sigma_{g,p})$  on  $U_{n-1,g,p}$  is



faithful. Let  $g \in \text{Inn}(\mathcal{M}_{g,p}^n)$  such that  $\Theta(g) = 1$  and  $C$  be an essential curve. We can associate to the curve  $C$  another simple closed curve  $D$  such that they bound an annulus containing only the puncture  $p_n$ . The mapping class  $T_C T_D^{-1}$  is then a  $n$ -bounding pair braid.

Since  $\Theta(g) = 1$ , then  $g T_C T_D^{-1} g^{-1} = T_C T_D^{-1}$  and from a simple argument on the index of intersection of curves (see for instance Proposition 2.10 of [2]) one can easily deduce that  $g(C) = C$ . Since  $g(C) = C$  for any essential curve  $C$  it follows that  $g$  is isotopic to the identity (Lemma 5.1 and Theorem 5.3 of [7]). Therefore  $\Theta$  is injective and in particular the representation defined in Theorem 1 is faithful when  $g \geq 1$ .

The only case left is when the genus is equal to zero. We recall that, for  $p > 0$ , the group  $B_n(\Sigma_{0,p})$  is isomorphic to the subgroup  $B_{n+p-1,p-1}$  of  $B_{n+p-1}$  fixing the last  $p-1$  strands. In this case our representation coincides with Artin representation of  $B_{n+p-1}$  in  $\text{Aut}(F_{n+p-1})$  restricted to the subgroup  $B_{n+p-1,p-1}$ .  $\square$

**Corollary 1.** *The group  $B_m(\Sigma_{g,p})$  is residually finite for  $m \geq 1$ ,  $g \geq 0$  and  $p > 0$ .*

*Proof.* Baumslag and Smirnov proved (see for instance [9, Theorem 4.8]) that any finitely presented group which is isomorphic to a subgroup of automorphisms of a free group of finite rank is residually finite. Therefore for  $m > 1$  the statement is a Corollary of Theorem 2. In the case  $m = 1$ , the group  $B_1(\Sigma_{g,p})$  is isomorphic to the fundamental group of  $\Sigma_{g,p}$  which is free and therefore residually finite.  $\square$

Remark also that from Theorem 2 one can derive, using Fox derivatives, a Burau representation for  $B_n(\Sigma_{g,p})$ . Since the restriction on  $B_n$  of such representation coincides with the usual Burau representation of  $B_n$ , the Burau representation for  $B_n(\Sigma_{g,p})$  obtained from Theorem 2 is not faithful for  $n \geq 5$ .

### 3. SURFACE BRAIDS AS OUTER AUTOMORPHISMS OF FREE GROUPS

As recalled in the Introduction, any element  $\beta$  of  $B_n \subset \text{Aut}(F_n)$  fixes the product  $x_1 x_2 \dots x_n$  of generators of  $F_n$ . We prove a similar statement for the group  $B_n(\Sigma_{g,p})$ , for  $p > 0$ .

**Proposition 3.** *Let  $p > 0$  and let  $U_{n-1,g,p}$  be the free group of rank  $n + p + 2g - 2$  defined above. Any element  $\beta$  in  $B_{n-1}(\Sigma_{g,p}) \subset \text{Aut}(U_{n-1,g,p})$  fixes the product*

$$A = \tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1} \xi_1 \xi_2 \dots \xi_{p-1} [w_1^{-1}, w_2] \dots [w_{2g-1}^{-1}, w_{2g}].$$

*Proof.* In order to prove the claim it suffices to verify that any generator of  $B_{n-1}(\Sigma_{g,p})$ , considered as an automorphism of  $U_{n-1,g,p}$ , fixes the element  $A$ .

*Case 1:* generators  $\sigma_i$ ,  $1 \leq i \leq n-2$ . The group  $B_{n-1}$  is a subgroup of  $B_{n-1}(\Sigma_{g,p})$  and by Artin's theorem  $B_{n-1}$  is a subgroup of  $\text{Aut}(F_{n-1})$ ,  $F_{n-1} = \langle \tau_1, \tau_2, \dots, \tau_{n-1} \rangle$ , and fixes the product  $\tau_1 \tau_2 \dots \tau_{n-1}$ . Hence, the generator  $\sigma_i$  also fixes the product  $\tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1}$ .

From Theorem 1 it follows that

$$w_r^{\sigma_i} = w_r, 1 \leq r \leq 2g; \quad \xi_j^{\sigma_i} = \xi_j, 1 \leq j \leq p-1.$$

Hence,  $A^{\sigma_i} = A$ .

*Case 2:* generators  $x_r$ ,  $1 \leq r \leq 2g$ . By Theorem 1 we have

$$(\tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1})^{x_r} = \tau_{n-1}^{-1} \dots \tau_2^{-1} (\tau_1^{-1} w_r \tau_1^{-1} w_r^{-1} \tau_1)$$

and

$$(\xi_1 \dots \xi_{p-1})^{x_r} = (\xi_1 \dots \xi_{p-1})^{[w_r^{-1}, \tau_1]}.$$

Let us consider

$$([w_1^{-1}, w_2] \dots [w_{2g-1}^{-1}, w_{2g}])^{x_r}.$$

We will distinguish two cases:  $r = 2m - 1$  and  $r = 2m$ , where  $m = 1, \dots, g$ . In the first case we have the following equalities:

$$([w_1^{-1}, w_2] \dots [w_{2m-3}^{-1}, w_{2m-2}])^{x_{2m-1}} = ([w_1^{-1}, w_2] \dots [w_{2m-3}^{-1}, w_{2m-2}])^{[w_{2m-1}^{-1}, \tau_1]},$$

$$[w_{2m-1}^{-1}, w_{2m}]^{x_{2m-1}} = [w_{2m-1}^{-\tau_1}, \tau_1^{-1} w_{2m}] = [\tau_1, w_{2m-1}^{-1}] [w_{2m-1}^{-1}, w_{2m}],$$

$$([w_{2m+1}^{-1}, w_{2m+2}] \dots [w_{2g-1}^{-1}, w_{2g}])^{x_{2m-1}} = [w_{2m+1}^{-1}, w_{2m+2}] \dots [w_{2g-1}^{-1}, w_{2g}].$$

In the second case we have the following equalities:

$$([w_1^{-1}, w_2] \dots [w_{2m-3}^{-1}, w_{2m-2}])^{x_{2m}} = ([w_1^{-1}, w_2] \dots [w_{2m-3}^{-1}, w_{2m-2}])^{[w_{2m}^{-1}, \tau_1]},$$

$$[w_{2m-1}^{-1}, w_{2m}]^{x_{2m}} = [\tau_1^{-1} w_{2m-1}^{-1} [w_{2m}^{-1}, \tau_1], w_{2m}] = [\tau_1, w_{2m}^{-1}] [w_{2m-1}^{-1}, w_{2m}],$$

$$([w_{2m+1}^{-1}, w_{2m+2}] \dots [w_{2g-1}^{-1}, w_{2g}])^{x_{2m}} = [w_{2m+1}^{-1}, w_{2m+2}] \dots [w_{2g-1}^{-1}, w_{2g}].$$

In both cases we have

$$([w_1^{-1}, w_2] \dots [w_{2g-1}^{-1}, w_{2g}])^{x_r} = [\tau_1, w_r^{-1}] [w_1^{-1}, w_2] \dots [w_{2g-1}^{-1}, w_{2g}].$$

Hence, the element  $x_r$  acts by conjugation on  $A$  as follows

$$A^{x_r} = \tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1} [w_r^{-1}, \tau_1] (\xi_1 \dots \xi_{p-1})^{[w_r^{-1}, \tau_1]} [\tau_1, w_r^{-1}] [w_1^{-1}, w_2] \dots [w_{2g-1}^{-1}, w_{2g}],$$

and then it is easy to check that  $A^{x_r} = A$ .

*Case 3:* generators  $z_j$ ,  $1 \leq j \leq p - 1$ . By Theorem 1 we have

$$(\tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1})^{z_j} = \tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1} [\xi_j^{-1}, \tau_1],$$

$$(\xi_1 \dots \xi_{p-1})^{z_j} = (\xi_1 \dots \xi_{j-1})^{[\xi_j^{-1}, \tau_1]} \xi_j^{\tau_1} \xi_{j+1} \dots \xi_{p-1} = [\xi_j^{-1}, \tau_1]^{-1} \xi_1 \dots \xi_{p-1},$$

and

$$([w_1^{-1}, w_2] \dots [w_{2g-1}^{-1}, w_{2g}])^{z_j} = [w_1^{-1}, w_2] \dots [w_{2g-1}^{-1}, w_{2g}].$$

Hence,  $A^{z_j} = A$  and the Proposition follows.  $\square$

We recall that  $\rho_U : B_{n-1}(\Sigma_{g,p}) \rightarrow \text{Aut}(U_{n-1,g,p})$  is the representation of  $B_{n-1}(\Sigma_{g,p})$  defined in Theorem 1 and let  $p : \text{Aut}(U_{n-1,g,p}) \rightarrow \text{Out}(U_{n-1,g,p})$  be the canonical projection.

**Theorem 3.** *The representation  $p \circ \rho_U : B_{n-1}(\Sigma_{g,p}) \rightarrow \text{Out}(U_{n-1,g,p})$  is faithful for  $p > 0$  when  $g > 0$  and for  $p > 2$  when  $g = 0$ .*

*Proof.* Since the representation  $\rho_U : B_{n-1}(\Sigma_{g,p}) \rightarrow \text{Aut}(U_{n-1,g,p})$  is faithful (Theorem 2), we can identify  $B_{n-1}(\Sigma_{g,p})$  with  $\rho_U(B_{n-1}(\Sigma_{g,p}))$ .

Now suppose that there exists  $\beta \in B_{n-1}(\Sigma_{g,p}) \cap \text{Inn}(U_{n-1,g,p})$ . From Proposition 3 one deduces that  $\beta$  is a conjugation by a power  $m$  of

$$A = \tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1} \xi_1 \xi_2 \dots \xi_{p-1} [w_1^{-1}, w_2] \dots [w_{2g-1}^{-1}, w_{2g}].$$

Now, let  $g$  be a generator of  $B_{n-1}(\Sigma_{g,p})$ . Since all elements of  $B_{n-1}(\Sigma_{g,p})$  fix the element  $A$  we deduce the following equalities:

$$g^{-1}(\beta(g(x))) = g^{-1}(A^m g(x) A^{-m}) = A^m x A^{-m} = \beta(x),$$

for any  $x \in U_{n-1,g,p}$ . One deduces that  $g^{-1}\beta g = \beta$  for any generator  $g$  of  $B_{n-1}(\Sigma_{g,p})$  and therefore  $\beta$  belongs to the center of  $B_{n-1}(\Sigma_{g,p})$ . Since  $B_{n-1}(\Sigma_{g,p})$  (with  $n \geq 2$ ) has trivial center for  $p > 0$  when  $g > 0$  [11] and for  $p > 2$  when  $g = 0$ , the intersection  $B_{n-1}(\Sigma_{g,p}) \cap \text{Inn}(U_{n-1,g,p})$  is trivial and the claim follows.  $\square$

**Remark 1.** Let  $\phi : B_n \rightarrow \text{Out}(F_n)$  the representation obtained composing Artin representation of  $B_n$  in  $\text{Aut}(F_n)$  with the canonical projection of  $\text{Aut}(F_n)$  in  $\text{Out}(F_n)$ . Such representation is not faithful and it is easy to see that the kernel is the center of  $B_n$ .

#### 4. SURFACE BRAID GROUPS OF NON-ORIENTABLE SURFACES WITH BOUNDARY COMPONENTS

Let  $N_{g,p}$  be a non-orientable surface of genus  $g \geq 1$ , with  $p > 0$  boundary components. Let  $\sigma_1, \dots, \sigma_{n-1}$  be the usual generators of  $B_n$  and  $a_1, \dots, a_g, z_1, \dots, z_{p-1}$  be the usual generators of the fundamental group of  $N_{g,p}$ . As in previous section we can consider  $\sigma_1, \dots, \sigma_{n-1}$  and  $a_1, \dots, a_g, z_1, \dots, z_{p-1}$  as elements of  $B_n(N_{g,p})$ . According to [1] the group  $B_n(N_{g,p})$  admits a presentation with generators:

$$\sigma_1, \dots, \sigma_{n-1}, a_1, \dots, a_g, z_1, \dots, z_{p-1},$$

and relations:

– Braid relations:

$$\begin{aligned} \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1}, \quad 1 \leq i \leq n-2, \\ \sigma_i \sigma_j &= \sigma_j \sigma_i, \quad |i-j| > 1, \quad 1 \leq i, j \leq n-1, \end{aligned}$$

– Mixed relations:

$$\begin{aligned} \text{(R1)} \quad a_r \sigma_i &= \sigma_i a_r, \quad i \neq 1, \quad 1 \leq r \leq g, \\ \text{(R2)} \quad \sigma_1^{-1} a_r \sigma_1^{-1} a_r &= a_r \sigma_1^{-1} a_r \sigma_1, \quad 1 \leq r \leq g, \\ \text{(R3)} \quad (\sigma_1^{-1} a_s \sigma_1) a_r &= a_r (\sigma_1^{-1} a_s \sigma_1), \quad 1 \leq s < r \leq g, \end{aligned}$$

- (R4)  $z_j \sigma_i = \sigma_i z_j, i \neq 1, 1 \leq j \leq p-1,$
- (R5)  $(\sigma_1^{-1} z_i \sigma_1) a_r = a_r (\sigma_1^{-1} z_i \sigma_1), 1 \leq r \leq g, 1 \leq i \leq p-1, n > 1,$
- (R6)  $(\sigma_1^{-1} z_j \sigma_1) z_l = z_l (\sigma_1^{-1} z_j \sigma_1), 1 \leq j < l \leq p-1,$
- (R7)  $(\sigma_1^{-1} z_j \sigma_1^{-1}) z_j = z_j (\sigma_1^{-1} z_j \sigma_1^{-1}), 1 \leq j \leq p-1.$

As in previous section let us consider the natural projection of  $\pi : B_n(N_{g,p}) \longrightarrow S_n$ , which associates to any braid the corresponding permutation. This projection map  $\sigma_i$  in the corresponding transposition and generators  $a_1, \dots, a_g, z_1, \dots, z_{p-1}$  into the identity. As before, let  $D_n(N_{g,p}) = \pi^{-1}(S_{n-1})$  and let  $m_l = \sigma_{n-1} \dots \sigma_l, l = 1, \dots, n-1, m_n = 1$ . The set  $M_n = \{m_l \mid 1 \leq l \leq n\}$  is a Schreier set of coset representatives of  $D_n(N_{g,p})$  in  $B_n(N_{g,p})$  and the group  $D_n(N_{g,p})$  is generated by

$$s_{\lambda,a} = \lambda a \cdot (\overline{\lambda a})^{-1},$$

where  $\lambda$  runs over the set  $M_n$  and  $a$  runs over the set of generators of  $B_n(N_{g,p})$ .

**Case 1.** If  $a \in \{\sigma_1, \dots, \sigma_{n-1}\}$ , then we find the generators

$$\tau_k = \sigma_{n-1} \dots \sigma_{k+1} \sigma_k^2 \sigma_{k+1}^{-1} \dots \sigma_{n-1}^{-1}, k = 1, \dots, n-2, \tau_{n-1} = \sigma_{n-1}^2.$$

**Case 2.** If  $a \in \{a_1, \dots, a_g\}$ , then we find the generators

$$w_r = \sigma_{n-1} \dots \sigma_1 a_r \sigma_1^{-1} \dots \sigma_{n-1}^{-1}, r = 1, \dots, g.$$

**Case 3.** If  $a \in \{z_1, \dots, z_{p-1}\}$ , then we find the generators

$$\xi_j = \sigma_{n-1} \dots \sigma_1 z_j \sigma_1^{-1} \dots \sigma_{n-1}^{-1}, j = 1, \dots, p-1.$$

Using the same argument as in the orientable case one can find the following group presentation for  $D_n(N_{g,p})$ .

**Proposition 4.** *The group  $D_n(N_{g,p}), n \geq 1$  admits a presentation with the generators*

$$\sigma_1, \dots, \sigma_{n-2}, a_1, \dots, a_g, z_1, \dots, z_{p-1}, \tau_1, \dots, \tau_{n-1}, w_1, \dots, w_g, \xi_1, \dots, \xi_{p-1},$$

*and the following relations:*

– *Braid relations:*

- (B1)  $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, 1 \leq i \leq n-3,$
- (B2)  $\sigma_i \sigma_j = \sigma_j \sigma_i, |i-j| > 1, 1 \leq i, j \leq n-2,$
- (B3)  $\sigma_k^{-1} \tau_l \sigma_k = \tau_l, k \neq l-1, l,$
- (B4)  $\sigma_{l-1}^{-1} \tau_l \sigma_{l-1} = \tau_{l-1},$
- (B5)  $\sigma_l^{-1} \tau_l \sigma_l = \tau_l \tau_{l+1} \tau_l^{-1}, l \neq n-1.$

– *Mixed relations:*

- (R1.1)  $a_r \sigma_i = \sigma_i a_r$ ,  $2 \leq i \leq n-2$ ,  $1 \leq r \leq g$ ,
- (R1.2)  $a_r \tau_i = \tau_i a_r$ ,  $2 \leq i \leq n-1$ ,
- (R1.3)  $w_r \sigma_i = \sigma_i w_r$ ,  $1 \leq r \leq g$ ,  $1 \leq i \leq n-2$ ,
- (R2.1)  $(\sigma_1^{-1} a_r \sigma_1^{-1}) a_r = a_r (\sigma_1^{-1} a_r \sigma_1)$ ,  $1 \leq r \leq g$ ,
- (R2.2)  $a_r^{-1} (\tau_1^{-1} w_r) a_r = \tau_1^{-1} w_r \tau_1$ ,  $1 \leq r \leq g$ ,
- (R2.3)  $a_r^{-1} w_r a_r = \tau_1^{-1} w_r$ ,  $1 \leq r \leq g$ ,
- (R3.1)  $(\sigma_1^{-1} a_s \sigma_1) a_r = a_r (\sigma_1^{-1} a_s \sigma_1)$ ,  $1 \leq s < r \leq g$ ,
- (R3.2)  $a_r^{-1} (\tau_1^{-1} w_s \tau_1) a_r = \tau_1^{-1} w_s \tau_1$ ,  $1 \leq s < r \leq g$ ,
- (R3.3)  $a_s w_r = w_r a_s$ ,  $1 \leq s < r \leq g$ ,
- (R4.1)  $z_j \sigma_i = \sigma_i z_j$ ,  $2 \leq i \leq n-2$ ,  $1 \leq j \leq p-1$ ,
- (R4.2)  $z_j \tau_i = \tau_i z_j$ ,  $2 \leq i \leq n-1$ ,  $1 \leq j \leq p-1$ ,
- (R4.3)  $\xi_j \sigma_i = \sigma_i \xi_j$ ,  $1 \leq j \leq p-1$ ,  $1 \leq i \leq n-2$ ,
- (R5.1)  $(\sigma_1^{-1} z_j \sigma_1) a_r = a_r (\sigma_1^{-1} z_j \sigma_1)$ ,  $1 \leq r \leq g$ ,  $1 \leq j \leq p-1$ ,
- (R5.2)  $a_r^{-1} (\tau_1^{-1} \xi_j \tau_1) a_r = \tau_1^{-1} \xi_j \tau_1$ ,  $1 \leq r \leq g$ ,  $1 \leq j \leq p-1$ ,
- (R5.3)  $z_j w_r = w_r z_j$ ,  $1 \leq r \leq g$ ,  $1 \leq j \leq p-1$ ,
- (R6.1)  $(\sigma_1^{-1} z_j \sigma_1) z_l = z_l (\sigma_1^{-1} z_j \sigma_1)$ ,  $1 \leq j < l \leq p-1$ ,
- (R6.2)  $z_l^{-1} (\tau_1^{-1} \xi_j \tau_1) z_l = \tau_1^{-1} \xi_j \tau_1$ ,  $1 \leq j < l \leq p-1$ ,
- (R6.3)  $z_j \xi_l = \xi_l z_j$ ,  $1 \leq j < l \leq p-1$ ,
- (R7.1)  $(\sigma_1^{-1} z_j \sigma_1^{-1}) z_j = z_j (\sigma_1^{-1} z_j \sigma_1^{-1})$ ,  $1 \leq j \leq p-1$ ,
- (R7.2)  $z_j^{-1} (\tau_1^{-1} \xi_j) z_j = \tau_1^{-1} \xi_j$ ,  $1 \leq j \leq p-1$ ,
- (R7.3)  $z_j^{-1} \xi_j z_j = \tau_1^{-1} \xi_j \tau_1$ ,  $1 \leq j \leq p-1$ .

It is easy to see that the relations (B1), (B2), (R1.1), (R2.1),  $\dots$ , (R7.1) are defining relations of  $B_{n-1}(N_{g,p})$ . From the other relations we can find the following conjugacy formulae:

- (S1)  $\tau_l^{\sigma_k} = \tau_l$ ,  $k \neq l-1, l$ ,
- (S2)  $\tau_l^{\sigma_{l-1}} = \tau_{l-1}$ ,
- (S3)  $\tau_l^{\sigma_l} = \tau_{l+1}^{\tau_l^{-1}}$ ,  $l \neq n-1$ ,
- (S4)  $\tau_i^{a_r} = \tau_i$ ,  $2 \leq i \leq n-1$ ,
- (S5)  $w_r^{\sigma_i} = w_r$ ,  $1 \leq r \leq g$ ,  $1 \leq i \leq n-2$ ,
- (S6)  $\tau_1^{a_r} = (\tau_1^{-1})^{w_r^{-1} \tau_1}$ ,
- (S7)  $w_r^{a_r} = \tau_1^{-1} w_r$ ,  $1 \leq r \leq g$ ,
- (S8)  $w_s^{a_r} = w_s^{w_r \tau_1 w_r^{-1} \tau_1}$ ,  $1 \leq s < r \leq g$ ,
- (S9)  $w_r^{a_s} = w_r$ ,  $1 \leq s < r \leq g$ ,
- (S10)  $\tau_i^{z_j} = \tau_i$ ,  $2 \leq i \leq n-1$ ,  $1 \leq j \leq p-1$ ,
- (S11)  $\xi_j^{\sigma_i} = \xi_j$ ,  $1 \leq i \leq n-2$ ,  $1 \leq j \leq p-1$ ,
- (S12)  $\xi_j^{a_r} = \xi_j^{w_r \tau_1 w_r^{-1} \tau_1}$ ,  $1 \leq r \leq g$ ,  $1 \leq j \leq p-1$ ,
- (S13)  $w_r^{z_j} = w_r$ ,  $1 \leq r \leq g$ ,  $1 \leq j \leq p-1$ ,
- (S14)  $\xi_j^{z_l} = \xi_j^{[\xi_l^{-1}, \tau_1]}$ ,  $1 \leq j < l \leq p-1$ ,

- (S15)  $\xi_l^{z_j} = \xi_l, 1 \leq j < l \leq p-1,$   
 (S16)  $\tau_1^{z_j} = [\tau_1, \xi_j^{-1}] \tau_1, 1 \leq j \leq p-1,$   
 (S17)  $\xi_j^{z_j} = \xi_j^{\tau_1}, 1 \leq j \leq p-1.$

Consider the subgroup

$$W_{n,g,p} = \langle \tau_1, \dots, \tau_{n-1}, w_1, \dots, w_g, \xi_1, \dots, \xi_{p-1} \rangle$$

of  $D_{n+1}(N_{g,p})$ . The group  $W_{n,g,p}$  is free (see [1]) and from (S1)–(S17) we see that  $B_n(N_{p,g})$  acts on  $W_{n,g,p}$  by conjugacy.

**Theorem 4.** *The group  $B_n(N_{g,p})$ , for  $n > 1$ ,  $p \geq 1$  and  $g \geq 1$  acts by conjugation on the free group  $W_{n,g,p}$  and the induced representation  $\rho_W : B_n(N_{g,p}) \rightarrow \text{Aut}(W_{n,g,p})$  is defined as follows:*

- Generators  $\sigma_i, i = 1, \dots, n-1$ :

$$\sigma_i : \begin{cases} \tau_i \mapsto \tau_{i+1}^{-1}; \\ \tau_{i+1} \mapsto \tau_i; \\ \tau_l \mapsto \tau_l, l \neq i, i+1; \\ w_r \mapsto w_r, 1 \leq r \leq g; \\ \xi_j \mapsto \xi_j, 1 \leq j \leq p-1. \end{cases}$$

- Generators  $a_r, r = 1, \dots, g$ :

$$a_r : \begin{cases} \tau_1 \mapsto (\tau_1^{-1})^{w_r^{-1}} \tau_1; \\ \tau_i \mapsto \tau_i, 2 \leq i \leq n; \\ w_s \mapsto w_s^{w_r \tau_1 w_r^{-1} \tau_1}, 1 \leq s < r \leq g; \\ w_r \mapsto \tau_1^{-1} w_r; \\ w_s \mapsto w_s, 1 \leq r < s \leq g; \\ \xi_j \mapsto \xi_j^{w_r \tau_1 w_r^{-1} \tau_1}, 1 \leq j \leq p-1. \end{cases}$$

- Generators  $z_j, j = 1, \dots, p-1$ :

$$z_j : \begin{cases} \tau_1 \mapsto [\tau_1, \xi_j^{-1}] \tau_1; \\ \tau_i \mapsto \tau_i, 2 \leq i \leq n; \\ w_r \mapsto w_r, 1 \leq r \leq g; \\ \xi_l \mapsto \xi_l^{[\xi_j^{-1}, \tau_1]}, 1 \leq l < j \leq p-1; \\ \xi_j \mapsto \xi_j^{\tau_1}; \\ \xi_l \mapsto \xi_l, 1 \leq j < l \leq p-1. \end{cases}$$

We don't know if the representation  $\rho_W : B_n(N_{g,p}) \rightarrow \text{Aut}(W_{n,g,p})$  is faithful or not.

**Proposition 5.** *Let  $\beta$  be in  $B_n(N_{g,p})$ . The element  $\rho_W(\beta)$  in  $\text{Aut}(W_{n,g,p})$  fixes the product*

$$A = \tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1} \xi_1 \xi_2 \dots \xi_{p-1} w_1^2 w_2^2 \dots w_g^2.$$

*Proof.* As above it is enough to prove that images of generators of  $B_n(N_{g,p})$  fix  $A$ .

*Case 1:* generators  $\sigma_i$ ,  $1 \leq i \leq n-1$ . The group  $B_n$  is a subgroup of  $B_n(N_{g,p})$  and by Artin's theorem  $B_n$  is a subgroup of  $\text{Aut}(F_n)$ ,  $F_n = \langle \tau_1, \tau_2, \dots, \tau_{n-1} \rangle$  and fixes the product  $\tau_1 \tau_2 \dots \tau_{n-1}$ . Hence,  $\rho_W(\sigma_i)$  also fixes the product  $\tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1}$ .

From Theorem 4 it follows that

$$w_r^{\sigma_i} = w_r, 1 \leq r \leq g; \xi_j^{\sigma_i} = \xi_j, 1 \leq j \leq p-1.$$

Hence,  $A^{\sigma_i} = A$ .

*Case 2:* generators  $a_r$ ,  $1 \leq r \leq g$ . By Theorem 4 we have that

$$(\tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1})^{a_r} = \tau_{n-1}^{-1} \dots \tau_2^{-1} (\tau_1^{w_r^{-1} \tau_1}),$$

and

$$(\xi_1 \dots \xi_{p-1})^{a_r} = (\xi_1 \dots \xi_{p-1})^{w_r \tau_1 w_r^{-1} \tau_1},$$

and

$$(w_1^2 \dots w_g^2)^{a_r} = (w_1^2 \dots w_{r-1}^2)^{w_r \tau_1 w_r^{-1} \tau_1} (\tau_1^{-1} w_r \tau_1^{-1} w_r) (w_{r+1}^2 \dots w_g^2).$$

Hence, the element  $a_r$  acts on  $A$  as follows:

$$A^{a_r} = \tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1} \xi_1 \dots \xi_{p-1} w_1^2 w_2^2 \dots w_g^2,$$

and  $A^{a_r} = A$ .

*Case 3.* Generators  $z_j$ ,  $1 \leq j \leq p-1$ . By Theorem 4 we deduce that

$$(\tau_{n-1}^{-1} \dots \tau_2^{-1} \tau_1^{-1})^{z_j} = \tau_{n-1}^{-1} \dots \tau_2^{-1} (\tau_1^{-1} \xi_j \tau_1^{-1} \xi_j^{-1} \tau_1),$$

$$(\xi_1 \dots \xi_{p-1})^{z_j} = (\xi_1 \dots \xi_{j-1})^{\xi_j \tau_1^{-1} \xi_j^{-1} \tau_1} (\xi_j^{\tau_1}) \xi_{j+1} \dots \xi_{p-1},$$

and

$$(w_1^2 w_2^2 \dots w_g^2)^{z_j} = w_1^2 w_2^2 \dots w_g^2.$$

Hence,  $A^{z_j} = A$ . □

## 5. BRAID GROUPS OF CLOSED SURFACES

Let  $\Sigma_g = \Sigma_{g,0}$  be a closed orientable surface of genus  $g \geq 1$ . The group  $B_n(\Sigma_g)$ ,  $n > 1$  admits a group presentation with generators:

$$\sigma_1, \dots, \sigma_{n-1}, x_1, \dots, x_{2g},$$

and relations:

– Braid relations:

$$\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \quad 1 \leq i \leq n-2,$$

$$\sigma_i \sigma_j = \sigma_j \sigma_i, \quad |i-j| > 1, \quad 1 \leq i, j \leq n-1,$$

– Mixed relations:

$$(R1) \quad x_r \sigma_i = \sigma_i x_r, \quad i \neq 1, \quad 1 \leq r \leq 2g,$$

$$(R2) \quad (\sigma_1^{-1} x_r \sigma_1^{-1}) x_r = x_r (\sigma_1^{-1} x_r \sigma_1^{-1}), \quad 1 \leq r \leq 2g,$$

- (R3)  $(\sigma_1^{-1}x_s\sigma_1)x_r = x_r(\sigma_1^{-1}x_s\sigma_1)$ ,  $1 \leq s < r \leq 2g$ ,  $(s, r) \neq (2m-1, 2m)$ ,  
 (R4)  $(\sigma_1^{-1}x_{2m-1}\sigma_1^{-1})x_{2m} = x_{2m}(\sigma_1^{-1}x_{2m-1}\sigma_1)$ ,  $1 \leq m \leq g$ ,  
 (TR5)  $[x_1^{-1}, x_2][x_3^{-1}, x_4] \dots [x_{2g-1}^{-1}, x_{2g}] = \sigma_1\sigma_2 \dots \sigma_{n-1}^2 \dots \sigma_2\sigma_1$ .

As before, let  $\pi$  be the natural projection of  $B_n(\Sigma_g)$  in  $S_n$ , let  $D_n(\Sigma_g) = \pi^{-1}(S_{n-1})$  and  $m_l = \sigma_{n-1} \dots \sigma_l$ ,  $l = 1, \dots, n-1$ ,  $m_n = 1$ . The set  $M_n = \{m_l \mid 1 \leq l \leq n\}$  is a Schreier set of coset representatives of  $D_n(\Sigma_g)$  in  $B_n(\Sigma_g)$  and  $D_n(\Sigma_g)$  is generated by

$$\sigma_1, \sigma_2, \dots, \sigma_{n-2}, \tau_1, \tau_2, \dots, \tau_{n-1}, x_1, x_2, \dots, x_{2g}, w_1, w_2, \dots, w_{2g},$$

where

$$\tau_k = \sigma_{n-1} \dots \sigma_{k+1} \sigma_k^2 \sigma_{k+1}^{-1} \dots \sigma_{n-1}^{-1}, \quad k = 1, \dots, n-2, \quad \tau_{n-1} = \sigma_{n-1}^2,$$

$$w_r = \sigma_{n-1} \dots \sigma_1 x_r \sigma_1^{-1} \dots \sigma_{n-1}^{-1}, \quad r = 1, \dots, 2g.$$

**Proposition 6.** *The group  $D_n(\Sigma_g)$ ,  $n > 1$ , admits a group presentation with generators:*

$$\sigma_1, \dots, \sigma_{n-2}, x_1, \dots, x_{2g}, \tau_1, \dots, \tau_{n-1}, w_1, \dots, w_{2g},$$

and relations:

– *Braid relations:*

- (B1)  $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$ ,  $1 \leq i \leq n-3$ ,  
 (B2)  $\sigma_i \sigma_j = \sigma_j \sigma_i$ ,  $|i-j| > 1$ ,  $1 \leq i, j \leq n-2$ ,  
 (B3)  $\sigma_k^{-1} \tau_l \sigma_k = \tau_l$ ,  $k \neq l-1, l$ ,  
 (B4)  $\sigma_{l-1}^{-1} \tau_l \sigma_{l-1} = \tau_{l-1}$ ,  
 (B5)  $\sigma_l^{-1} \tau_l \sigma_l = \tau_l \tau_{l+1} \tau_l^{-1}$ ,  $l \neq n-1$ .

– *Mixed relations:*

- (R1.1)  $x_r \sigma_i = \sigma_i x_r$ ,  $2 \leq i \leq n-2$ ,  $1 \leq r \leq 2g$ ,  
 (R1.2)  $x_r \tau_i = \tau_i x_r$ ,  $2 \leq i \leq n-1$ ,  
 (R1.3)  $w_r \sigma_i = \sigma_i w_r$ ,  $1 \leq r \leq 2g$ ,  $1 \leq i \leq n-2$ ,  
 (R2.1)  $(\sigma_1^{-1} x_r \sigma_1^{-1}) x_r = x_r (\sigma_1^{-1} x_r \sigma_1^{-1})$ ,  $1 \leq r \leq 2g$ ,  
 (R2.2)  $x_r^{-1} w_r x_r = \tau_1^{-1} w_r \tau_1$ ,  $1 \leq r \leq 2g$ ,  
 (R2.3)  $x_r^{-1} \tau_1 x_r = \tau_1^{-1} w_r \tau_1 w_r^{-1} \tau_1$ ,  $1 \leq r \leq 2g$ ,  
 (R3.1)  $(\sigma_1^{-1} x_s \sigma_1) x_r = x_r (\sigma_1^{-1} x_s \sigma_1)$ ,  $1 \leq s < r \leq 2g$ ,  $(s, r) \neq (2m-1, 2m)$ ,  
 (R3.2)  $x_r^{-1} (\tau_1^{-1} w_s \tau_1) x_r = \tau_1^{-1} w_s \tau_1$ ,  $1 \leq s < r \leq 2g$ ,  $(s, r) \neq (2m-1, 2m)$ ,  
 (R3.3)  $x_s w_r = w_r x_s$ ,  $1 \leq s < r \leq 2g$ ,  $(s, r) \neq (2m-1, 2m)$ ,  
 (R4.1)  $(\sigma_1^{-1} x_{2m-1} \sigma_1^{-1}) x_{2m} = x_{2m} (\sigma_1^{-1} x_{2m-1} \sigma_1)$ ,  $1 \leq m \leq g$ ,  
 (R4.2)  $x_{2m}^{-1} (\tau_1^{-1} w_{2m-1}) x_{2m} = \tau_1^{-1} w_{2m-1} \tau_1$ ,  $1 \leq m \leq g$ ,  
 (R4.3)  $x_{2m-1}^{-1} w_{2m} x_{2m-1} = \tau_1^{-1} w_{2m}$ ,  $1 \leq m \leq g$ ,  
 (RT.1)  $[x_1^{-1}, x_2][x_3^{-1}, x_4] \dots [x_{2g-1}^{-1}, x_{2g}] = \sigma_1 \sigma_2 \dots \sigma_{n-3} \sigma_{n-2}^2 \sigma_{n-3} \dots \sigma_2 \sigma_1 \tau_1$ ,  
 (RT.2)  $[w_1^{-1}, w_2][w_3^{-1}, w_4] \dots [w_{2g-1}^{-1}, w_{2g}] = \tau_1 \tau_2 \dots \tau_{n-1}$ .

Consider the subgroup  $V_{n-1, g}$  of  $D_n(\Sigma_g)$  generated by  $\{\tau_2, \dots, \tau_{n-1}, w_1, \dots, w_{2g}\}$ . The group  $V_{n-1, g}$  is free and normal in  $D_n(\Sigma_g)$  [1]. Using relation (RT.2) one finds that

$$\tau_1 = [w_1^{-1}, w_2][w_3^{-1}, w_4] \dots [w_{2g-1}^{-1}, w_{2g}] \tau_{n-1}^{-1} \dots \tau_2^{-1}$$



and therefore  $\tau_1 \in V_n$ . Also, consider the subgroup

$$\overline{B}_{n-1}(\Sigma_g) = \langle \sigma_1, \dots, \sigma_{n-2}, x_1, \dots, x_{2g} \rangle$$

of  $D_n(\Sigma_g)$ . The group  $\overline{B}_{n-1}(\Sigma_g)$  is not isomorphic to  $B_{n-1}(\Sigma_g)$ .

In fact, one can prove that the relations (B1), (B2), (R1.1), (R2.1), (R3.1), (R4.1) and (RT.1) are defining relations of  $\overline{B}_{n-1}(\Sigma_g)$  and therefore we can deduce that  $B_{n-1}(\Sigma_g) \simeq \overline{B}_{n-1}(\Sigma_g) / \langle \langle \tau_1 \rangle \rangle$ . From the other relations we can see that  $\overline{B}_{n-1}(\Sigma_g)$  acts on  $V_n$  by conjugacy. Hence, the following theorem holds.

**Proposition 7.** *There exists a representation  $\rho_V : \overline{B}_{n-1}(\Sigma_g) \rightarrow \text{Aut}(V_{n-1,g})$ , induced by the action by conjugacy of  $\overline{B}_{n-1}(\Sigma_g)$  on the free group  $V_{n-1,g} = \langle \tau_2, \dots, \tau_{n-1}, w_1, \dots, w_{2g} \rangle$  given algebraically as follows:*

–Action on generators  $\tau_2, \dots, \tau_{n-1}$ :

$$(S1) \tau_l^{\sigma_k} = \tau_l, k \neq l-1, l,$$

$$(S2) \tau_l^{\sigma_{l-1}} = \tau_{l-1},$$

$$(S3) \tau_l^{\sigma_l} = \tau_{l+1}^{-1}, l \neq n-1,$$

$$(S4) \tau_i^{x_r} = \tau_i, 2 \leq i \leq n-1, 1 \leq r \leq 2g,$$

–Action on generators  $w_1, \dots, w_{2g}$ :

$$(S5) w_r^{\sigma_i} = w_r, 1 \leq r \leq 2g, 1 \leq i \leq n-2,$$

$$(S6) w_r^{x_r} = w_r^{\tau_1}, 1 \leq r \leq 2g,$$

$$(S7) \tau_1^{x_r} = \tau_1^{w_r^{-1}\tau_1}, 1 \leq r \leq 2g,$$

$$(S8) w_s^{x_r} = w_s^{[w_r^{-1}, \tau_1]}, 1 \leq s < r \leq 2g, (s, r) \neq (2m-1, 2m),$$

$$(S9) w_r^{x_s} = w_r, 1 \leq s < r \leq 2g, (s, r) \neq (2m-1, 2m),$$

$$(S10) w_{2m-1}^{x_{2m}} = [\tau_1, w_{2m}^{-1}] w_{2m-1} \tau_1,$$

$$(S11) w_{2m}^{x_{2m-1}} = \tau_1^{-1} w_{2m},$$

where

$$\tau_1 = [w_1^{-1}, w_2][w_3^{-1}, w_4] \dots [w_{2g-1}^{-1}, w_{2g}] \tau_{n-1}^{-1} \dots \tau_2^{-1}.$$

From Proposition 7 we get a representation of  $B_{n-1}(\Sigma_g)$  in  $\text{Aut}(V_{n-1,g}) / \langle \langle t \rangle \rangle$ , where  $t \in \text{Aut}(V_{n-1,g})$  is the conjugation by the word

$$[w_1^{-1}, w_2] \dots [w_{2g-1}^{-1}, w_{2g}] \tau_{n-1}^{-1} \dots \tau_2^{-1}.$$

Therefore, we have the following result:

**Proposition 8.** *There exists a representation  $\rho_{\tilde{V}} : B_{n-1}(\Sigma_g) \rightarrow \text{Out}(V_{n-1,g})$ , given algebraically as follows:*

–Action (up to conjugacy) on generators  $\tau_2, \dots, \tau_{n-1}$ :

$$(S1) \tau_l^{\sigma_k} = \tau_l, k \neq l-1, l,$$

$$(S2) \tau_l^{\sigma_{l-1}} = \tau_{l-1},$$

$$(S3) \tau_l^{\sigma_l} = \tau_{l+1}^{-1}, l \neq n-1,$$

$$(S4) \tau_i^{x_r} = \tau_i, 2 \leq i \leq n-1, 1 \leq r \leq 2g,$$

–Action (up to conjugacy) on generators  $w_1, \dots, w_{2g}$ :

$$(S5) w_r^{\sigma_i} = w_r, 1 \leq r \leq 2g, 1 \leq i \leq n-2,$$

$$(S6) w_r^{x_r} = w_r^{\tau_1}, 1 \leq r \leq 2g,$$

$$(S7) \tau_1^{x_r} = \tau_1^{w_r^{-1}\tau_1}, 1 \leq r \leq 2g,$$

$$(S8) w_s^{x_r} = w_s^{[w_r^{-1}, \tau_1]}, 1 \leq s < r \leq 2g, (s, r) \neq (2m-1, 2m),$$

$$(S9) w_r^{x_s} = w_r, 1 \leq s < r \leq 2g, (s, r) \neq (2m-1, 2m),$$

$$(S10) w_{2m-1}^{x_{2m}} = [\tau_1, w_{2m}^{-1}]w_{2m-1}\tau_1,$$

$$(S11) w_{2m}^{x_{2m-1}} = \tau_1^{-1}w_{2m},$$

where

$$\tau_1 = [w_1^{-1}, w_2][w_3^{-1}, w_4] \dots [w_{2g-1}^{-1}, w_{2g}]\tau_{n-1}^{-1} \dots \tau_2^{-1}.$$

We don't know if the representation  $\rho_{\tilde{V}}$  of  $B_{n-1}(\Sigma_g)$  is faithful or not.

## 6. ARTIN-TITS GROUPS

We recall that classical braid groups are also called Artin-Tits groups of type  $\mathcal{A}$ . More precisely, let  $(W, S)$  be a Coxeter system and let us denote the order of the element  $st$  in  $W$  by  $m_{s,t}$  (for  $s, t \in S$ ). Let  $A(W)$  be the group defined by the following group presentation:

$$A(W) = \langle S \mid \underbrace{st \cdots}_{m_{s,t}} = \underbrace{ts \cdots}_{m_{s,t}} \text{ for any } s \neq t \in S \text{ with } m_{s,t} < +\infty \rangle.$$

The group  $A(W)$  is the Artin-Tits group associated to  $W$ . The group  $A(W)$  is said to be of spherical type if  $W$  is finite. There exists three infinite families of finite Coxeter groups usually denoted respectively as of type  $\mathcal{A}$ ,  $\mathcal{B}$  and  $\mathcal{D}$ . Artin braid groups correspond to the

family of Artin-Tits groups associated to Coxeter groups of type  $\mathcal{A}$  and braid groups of the annulus coincide with Artin-Tits groups associated to Coxeter groups of type  $\mathcal{B}$ . In this Appendix we give a faithful representation of the remaining infinite family of spherical Artin-Tits groups, associated to Coxeter groups of type  $\mathcal{D}$ .

First let us denote by  $A(\mathcal{D}_n)$  the  $n$ -th Artin-Tits group of type  $\mathcal{D}$  with the group presentation provided by its Coxeter graph (see Figure 1), where vertices  $\delta_1, \dots, \delta_n$  are the generators of the group and two generators  $\delta_i, \delta_j$  verify the relation  $\delta_i \delta_j \delta_i = \delta_j \delta_i \delta_j$  if they are related by an edge and elsewhere they commute.

In the following we set  $\lambda_i = (\delta_{i+1} \cdots \delta_3)(\delta_1 \delta_2^{-1})(\delta_{i+1} \cdots \delta_3)^{-1}$  for  $i = 1, \dots, n-1$ .




FIGURE 1.

We denote by  $\pi_D : A(\mathcal{D}_n) \rightarrow B_n$  the epimorphism defined by  $\pi_D(\delta_1) = 1$  and  $\pi_D(\delta_i) = \sigma_{i-1}$  for  $i = 2, 3, \dots, n$ , which admits the section  $s_D : B_n \rightarrow A(\mathcal{D}_n)$  defined by  $s_D(\sigma_i) = \delta_{i+1}$  for  $i = 1, 2, \dots, n-1$ .

**Proposition 9.** [6, Proposition 2.3]

- (1) The representation  $\rho_D : B_n \rightarrow \text{Aut}(F_{n-1})$  given algebraically by:

$$\rho_D(\sigma_1) : \begin{cases} x_1 \mapsto x_1, \\ x_j \mapsto x_1^{-1} x_j, j \neq 1, \end{cases}$$

and for  $2 \leq i \leq n-1$ ,

$$\rho_D(\sigma_i) : \begin{cases} x_{i-1} \mapsto x_i, \\ x_i \mapsto x_i x_{i-1}^{-1} x_i, \\ x_j \mapsto x_j, \end{cases} \quad j \neq i-1, i.$$

is well defined and faithful.

- (2) The group  $A(\mathcal{D}_n)$  is isomorphic to  $F_{n-1} \rtimes_{\rho_D} B_n$ , where the projection on the second factor is  $\pi_D$  and the section  $B_n \rightarrow F_{n-1} \rtimes_{\rho_D} B_n$  is just  $s_D$ .  
(3) In particular  $\ker \pi_D$  is a free group of rank  $n-1$  freely generated by  $\lambda_1, \dots, \lambda_{n-1}$ .

The first item of Proposition 9 was already established in [12] by topological means and the third item is actually proven in the proof of Proposition 2.3 in [6].

From Proposition 9 we can deduce a faithful representation of  $A(\mathcal{D}_n)$  into  $\text{Aut}(F_{n-1})$ .

**Proposition 10.** The representation  $\iota : A(\mathcal{D}_n) \rightarrow \text{Aut}(F_{n-1})$  given algebraically by:

$$\iota(\delta_1) : \begin{cases} x_1 \mapsto x_1, \\ x_j \mapsto x_j x_1^{-1}, j \neq 1, \end{cases}$$

$$\iota(\delta_2) : \begin{cases} x_1 \mapsto x_1, \\ x_j \mapsto x_1^{-1}x_j, j \neq i, \end{cases}$$

and for  $3 \leq i \leq n$ ,

$$\iota(\delta_i) : \begin{cases} x_{i-1} \mapsto x_i, \\ x_i \mapsto x_i x_{i-1}^{-1} x_i, \\ x_j \mapsto x_j, \end{cases} \quad j \neq i-1, i.$$

is well defined and faithful.

*Proof.* By direct calculation one shows easily that  $\iota : A(\mathcal{D}_n) \rightarrow \text{Aut}(F_{n-1})$  is well-defined.

Now let  $\Delta_n$  be the subgroup of  $\text{Aut}(F_{n-1})$  generated by  $\iota(\delta_1), \dots, \iota(\delta_n)$  and let  $\pi_\Delta : \Delta_n \rightarrow B_n$  be the map defined by  $\pi_\Delta(\iota(\delta_1)) = \pi_{\Delta_n}(\iota(\delta_2)) = \sigma_1$  and  $\pi_\Delta(\iota(\delta_i)) = \sigma_{i-1}$  for  $i = 3, \dots, n$ .

From Proposition 9 we deduce that  $\pi_\Delta$  is a well defined epimorphism and that the composition  $\pi_\Delta \circ \iota$  coincides with  $\pi_D$ . Therefore  $\iota$  is injective if and only if its restriction to  $\ker \pi_D$  is injective.

By direct calculation one can verify that  $\iota(\lambda_i)$  is the conjugacy by  $x_i$  for  $i = 1, \dots, n-1$ ; therefore the group generated by  $\iota(\lambda_1), \dots, \iota(\lambda_{n-1})$  is a free group of rank  $n-1$ . Since  $\ker \pi_D$  is freely generated by  $\lambda_1, \dots, \lambda_{n-1}$  (Proposition 9) one deduces that  $\iota$  is injective on  $\ker \pi_D$  and then the claim follows.  $\square$

## REFERENCES

- [1] P. Bellingeri, On presentation of surface braid groups, *J. Algebra* **274** (2004), 543-563.
- [2] P. Bellingeri, On automorphisms of surface braid groups, to appear in *J. Knot Theory and Ramif.*
- [3] P. Bellingeri, S. Gervais and J. Guaschi, Lower central series for surface braids, math.GT/0512155.
- [4] J. S. Birman, Braids, links and mapping class group, Princeton-Tokyo: Univ. press, 1974.
- [5] J. Crisp and L. Paris, Representations of the braid group by automorphisms of groups, invariants of links, and Garside groups. *Pacific J. Math.* **221** (2005), No. 1, 1-27.
- [6] J. Crisp and L. Paris, Artin groups of type B and D, *Adv. Geom.* **5** (2005), 607-636.
- [7] N. V. Ivanov, and J. D. McCarthy, On injective homomorphisms between Teichmüller modular groups, I, *Invent. Math.* **135** (1999), 425-486.
- [8] W. Magnus, A. Karrass, D. Solitar, Combinatorial group theory, Interscience Publishers, New York, 1996.
- [9] R. S. Lindon, P. E. Shupp, Combinatorial group theory, Springer-Verlag, Berlin-Heidelberg-New York, 1977.
- [10] V. Lundsgaard Hansen, Braids and coverings, London Mathematical Society Student Texts 18, Cambridge, 1989.
- [11] L. Paris and D. Rolfsen, Geometric subgroups of surface braid groups, *Ann. Inst. Fourier* **49** (1999), 417-472.
- [12] B. Perron, J. P. Vannier, Groupes de monodromie géométrique des singularités simples, *Math. Ann.* **306** (1996), 231-245.
- [13] M. Wada, Group invariants of links. *Topology* **31** (1992), No. 2, 399-406.

SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK 630090, RUSSIA

*E-mail address:* `bardakov@math.nsc.ru`

UNIVERSITÀ MILANO-BICOCCA, DIPARTIMENTO MATEMATICA E APPLICAZIONI, 20126 MILANO, ITALY

*E-mail address:* `paolo.bellingeri@unimib.it`