



The Veldkamp Space of Two-Qubits

Metod Saniga, Michel R. P. Planat, Petr Pracna, Hans Havlicek

► To cite this version:

Metod Saniga, Michel R. P. Planat, Petr Pracna, Hans Havlicek. The Veldkamp Space of Two-Qubits. 2007. hal-00139548v1

HAL Id: hal-00139548

<https://hal.science/hal-00139548v1>

Preprint submitted on 2 Apr 2007 (v1), last revised 2 Jul 2007 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Veldkamp Space of Two-Qubits

Metod Saniga,¹ Michel Planat,² Petr Pracna³
and
Hans Havlicek⁴

¹Astronomical Institute, Slovak Academy of Sciences, SK-05960 Tatranská Lomnica
Slovak Republic
(msaniga@astro.sk)

²Institut FEMTO-ST, CNRS, Département LPMO, 32 Avenue de l'Observatoire
F-25044 Besançon Cedex, France
(michel.planat@femto-st.fr)

³J. Heyrovský Institute of Physical Chemistry, Academy of Sciences of the Czech Republic
Dolejšková 3, CZ-182 23 Prague 8, Czech Republic
(pracna@jh-inst.cas.cz)

and

⁴Institut für Diskrete Mathematik und Geometrie, Technische Universität Wien
Wiedner Hauptstraße 8–10, A-1040 Vienna, Austria
(havlicek@geometrie.tuwien.ac.at)

Abstract

Given a remarkable representation of the generalized Pauli operators of two-qubits in terms of the points of the generalized quadrangle of order two, $W(2)$, it is shown that specific subsets of these operators can also be associated with the points and lines of the four-dimensional projective space over the Galois field with two elements — the so-called Veldkamp space of $W(2)$. An intriguing novelty is the recognition of (uni- and tri-centric) triads and specific pentads of the Pauli operators in addition to the “classical” subsets answering to geometric hyperplanes of $W(2)$.

MSC Codes: 51Exx, 81R99

PACS Numbers: 02.10.Ox, 02.40.Dr, 03.65.Ca

Keywords: Generalized Quadrangles – Veldkamp Spaces – Pauli Operators of Two-Qubits

1 Introduction

A deeper understanding of the structure of Hilbert spaces of finite dimensions is of utmost importance for quantum information theory. Recently, we made an important step in this respect by demonstrating that the algebra of the generalized Pauli operators on the 2^N -dimensional Hilbert spaces is embodied in the geometry of the symplectic polar space of rank N and order two [1]–[3]. The case of two-qubit operator space, $N = 2$, was scrutinized in very detail [1, 3] by explicitly demonstrating, in different ways, the correspondence between various subsets of the generalized Pauli operators/matrices and the fundamental subgeometries of the associated rank-two polar space — the generalized quadrangle of order two, $W(2)$. In this paper we will reveal another interesting geometry hidden behind the Pauli operators of two-qubits, namely that of the Veldkamp space defined on $W(2)$, $\mathcal{V}(W(2))$.

2 Finite Generalized Quadrangles and Veldkamp spaces

In this section we will briefly highlight the basics of the theory of finite generalized quadrangles [4] and introduce the concept of the Veldkamp space of a point-line incidence geometry [5] to be employed in what follows.

A *finite generalized quadrangle* of order (s, t) , usually denoted $\text{GQ}(s, t)$, is an incidence structure $S = (P, B, I)$, where P and B are disjoint (non-empty) sets of objects, called respectively points and lines, and where I is a symmetric point-line incidence relation satisfying the following axioms [4]: (i) each point is incident with $1 + t$ lines ($t \geq 1$) and two distinct points are incident with at most one line; (ii) each line is incident with $1 + s$ points ($s \geq 1$) and two distinct lines are incident with at most one point; and (iii) if x is a point and L is a line not incident with x , then there exists a unique pair $(y, M) \in P \times B$ for which $xIMyIL$; from these axioms it readily follows that $|P| = (s + 1)(st + 1)$ and $|B| = (t + 1)(st + 1)$. It is obvious that there exists a point-line duality with respect to which each of the axioms is self-dual. Interchanging points and lines in S thus yields a generalized quadrangle S^D of order (t, s) , called the dual of S . If $s = t$, S is said to have order s . A generalized quadrangle of order $(s, 1)$ is called a grid and that of order $(1, t)$ a dual grid. A generalized quadrangle with both $s > 1$ and $t > 1$ is called thick.

Given two points x and y of S one writes $x \sim y$ and says that x and y are collinear if there exists a line L of S incident with both. For any $x \in P$ denote $x^\perp = \{y \in P | y \sim x\}$ and note that $x \in x^\perp$; obviously, $|x^\perp| = 1 + s + st$. Given an arbitrary subset A of P , the *perp*(-set) of A , A^\perp , is defined as $A^\perp = \bigcap \{x^\perp | x \in A\}$ and $A^{\perp\perp} := (A^\perp)^\perp$. A triple of pairwise non-collinear points of S is called a *triad*; given any triad T , a point of T^\perp is called its center and we say that T is *acentric*, *centric* or *unicentric* according as $|T^\perp|$ is, respectively, zero, non-zero or features just a single point. An *ovoid* of a generalized quadrangle S is a set of points of S such that each line of S is incident with exactly one point of the set; hence, each ovoid contains $st + 1$ points.

The concept of crucial importance is a *geometric hyperplane* H of a point-line geometry $\Gamma(P, L)$, which is a proper subset of P such that each line of Γ meets H in one or all points. For $\Gamma = \text{GQ}(s, t)$, one can check that H is one of the following three kinds: (i) the *perp*-set of a point x , x^\perp ; (ii) a (full) subquadrangle of order (s, t') , $t' < t$; and (iii) an ovoid.

Finally, we need to introduce the notion of the *Veldkamp space* of a point-line incidence geometry $\Gamma(P, L)$, $\mathcal{V}(\Gamma)$ [5]. $\mathcal{V}(\Gamma)$ is the space in which (i) a point is a geometric hyperplane of Γ and (ii) a line is the collection H_1H_2 of all geometric hyperplanes H of Γ such that $H_1 \cap H_2 = H_1 \cap H = H_2 \cap H$ or $H = H_i$ ($i = 1, 2$), where H_1 and H_2 are distinct points of $\mathcal{V}(\Gamma)$. If $\Gamma = S$, from the preceding paragraph we learn that the points of $\mathcal{V}(S)$ are, in general, of three different types.

3 The Simplest Thick GQ and its Veldkamp space

The simplest thick GQ is obviously the one with $s = t = 2$, frequently denoted as $W(2)$ and dubbed the “doily.” This quadrangle has a number of interesting representations of which we mention the most important two [4]: one in terms of the points of $PG(3, 2)$ together with the totally isotropic lines with respect to a symplectic polarity, other by the totally singular points and lines of a parabolic quadric in $PG(4, 2)$. From the preceding section we readily get that $W(2)$ is endowed with 15 points/lines, each line contains three points and, dually, each point is on three lines; moreover, it is a self-dual object, i.e., isomorphic to its dual. $W(2)$ features all the three kinds of hyperplanes, of the following cardinalities [4]: 15 *perp*-sets, x^\perp , seven points each; 10 grids (of order $(2, 1)$), nine points each; and six ovoids, five points each — as depicted in Figure 1. The quadrangle exhibits two distinct kinds of triads, viz. *unicentric* and *tricentric*. A point of $W(2)$ is the center of four distinct *unicentric* triads (Figure 2, left); hence, the number of such triads is $4 \times 15 = 60$. *Tricentric* triads always come in “complementary” pairs, one representing the centers of the other, and each such pair is the complement of a grid of $W(2)$ (Figure 2, right); hence, the number of such triads is $2 \times 10 = 20$. A *unicentric* triad is always a subset of an ovoid, which is never the case for a *tricentric* triad; the latter, in graph-combinatorial terms, representing a complete bipartite graph on six vertices. Now, we have enough background information at hand to reveal the structure of the Veldkamp space of our “doily,” $\mathcal{V}(W(2))$.




Figure 1: The three kinds of geometric hyperplanes of $W(2)$. The points of the quadrangle are represented by small circles and its lines are illustrated by the straight segments as well as by the segments of circles; note that not every intersection of two segments counts for a point of the quadrangle. The upper panel shows the points' perp-sets (yellow bullets), the middle panel grids (red bullets) and the bottom panel ovoids (blue bullets); the use of different colouring will become clear later. Each picture — except that in the bottom right-hand corner — stands for five different hyperplanes, the four other being obtained from it by its successive rotations through 72 degrees around the center of the pentagon.




Figure 2: *Left*: — The four distinct unicentric triads (grey bullets) and their common center (black bullet); note that the triads intersect pairwise in a single point and their union covers fully the center's perp-set. *Right*: — A grid (red bullets) and its complement as a disjoint union of two complementary tricentric triads (black and grey bullets); the two triads are also seen to comprise a dual grid (of order $(1, 2)$).

Table 1: A succinct summary of the properties of the five different types of the lines of $\mathcal{V}(W(2))$ in terms of the core-sets and the types of geometric hyperplanes featured by a generic line of a given type. The last column gives the total number of lines per the corresponding type.

Type of Core-Set	Perp-Sets	Grids	Ovoids	#
Single Point	1	0	2	15
Collinear Triple	3	0	0	15
Unicentric Triad	1	1	1	60
Tricentric Triad	3	0	0	20
Pentad	1	2	0	45

From the definition given in Sec. 2, we easily see that $\mathcal{V}(W(2))$ consists of 31 points of which 15 are represented/generated by single-point perp-sets, 10 by grids and six by ovoids. The lines of $\mathcal{V}(W(2))$ feature three points each and are of five distinct types, as illustrated in Figure 3. These types differ from each other in the cardinality and structure of “core-sets,” i.e., the sets of points of $W(2)$ shared by all the three hyperplanes forming a given line. As it is obvious from Figure 3, the lines of the first three types (the first three rows of the figure) have the core-sets of the same cardinality, three, differing from each other only in the structure of these sets as being unicentric triads, tricentric triads and triples of collinear points, respectively. The lines of the fourth type have as core-sets pentads of points, each being a quadruple of points collinear with a given point of $W(2)$, whereas core-sets of the last type’s lines feature just a single point. A much more interesting issue is the composition of the lines. Just a brief look at Figure 3 reveals that only one kind of a geometric hyperplane, namely perp-sets, are present on each line of $\mathcal{V}(W(2))$; grids and ovoids occur only on two kinds of the lines. We also see that the purely homogeneous types are those whose core-sets feature collinear triples and tricentric triads, the most heterogeneous type — the one exhibiting all the three kinds of hyperplanes — being that characterized by unicentric triads. We also notice that there are no lines comprising solely grids and/or solely ovoids, nor the lines featuring only grids and ovoids, which seems to be connected with the fact that the cardinality of a core-set is an odd number. From the properties of $W(2)$ and its triads as discussed above it readily follows that the number of the lines of type one to five is 15, 15, 60, 20 and 45, respectively, totalling 155. All these observations and facts are gathered in Table 1. We conclude this section with the observation that $\mathcal{V}(W(2))$ has the same number of points (31) and lines (155) as $PG(4, 2)$, the four-dimensional projective space over the Galois field of two elements [6]; this is not a coincidence, as the two spaces are, in fact, isomorphic to each other [5].

4 Pauli Operators of Two-Qubits in Light of $\mathcal{V}(W(2))$

As discovered in [1] (see also [3]), the fifteen generalized Pauli operators/matrices associated with the Hilbert space of two-qubits (see, e. g., [7]) can be put into a one-to-one correspondence with the fifteen points of the generalized quadrangle $W(2)$ in such a way that their commutation algebra is completely and uniquely reproduced by the geometry of $W(2)$ in which the concept commuting/non-commuting translates into that of collinear/non-collinear. Given this mapping, it was possible to ascribe a definitive geometrical meaning to sets of three pairwise commuting generalized Pauli operators in terms of lines of $W(2)$ and to other three kinds of distinguished subsets of the operators having their counterparts in geometric hyperplanes of $W(2)$ as shown in Table 2 (see [1, 3] for more details). Yet, $\mathcal{V}(W(2))$ puts this bijection in a different light, in which other three subsets of the Pauli operators come into play, namely those represented by the two types of a triad and by the specific pentads occurring as the core-sets of the lines of $\mathcal{V}(W(2))$ (Table 1). As already mentioned, the role of tricentric triads of the operators has been recognized in disguise of complete bipartite graphs on six vertices [3]. A true novelty here is obviously *unicentric triads* and *pentads* of the generalized Pauli operators as these are all




Figure 3: The five different kinds of the lines of $\mathcal{V}(W(2))$, each being uniquely determined by the properties of its core-set (black bullets). Note that the “yellow” hyperplanes (i. e., perp-sets) occur in each type, and yellow is also the colour of two homogeneous (i. e., endowed with only one kind of a hyperplane) types (2nd and 3rd row). It is also worth mentioning that the cardinality of core-sets is an odd number not exceeding five. The three hyperplanes of any line are always in such relation to each other that their union comprises all the points of $W(2)$.

Table 2: Three kinds of the distinguished subsets of the generalized Pauli operators of two-qubits (PO) viewed as the geometric hyperplanes in the generalized quadrangle of order two (GQ) [1, 3].

PO	set of five mutually non-commuting operators	set of six operators commuting with a given one	nine operators of a Mermin's square
GQ	ovoid	perp-set \ {reference point}	grid

intimately connected with single-point perp-sets; given a point of $W(2)$ (i. e., a generalized Pauli operator of two-qubits), its perp-set fully encompasses *four* unicentric triads (Figure 2, left) and *three* pentads (Figure 3, 4th row) of points/operators. This feature has also a very interesting aspect in connection with the conjecture relating the existence of mutually unbiased bases and finite projective planes raised in [8], because with each point x of $W(2)$ there is associated a projective plane of order two (the Fano plane) whose points are the elements of x^\perp and whose lines are the spans $\{u, v\}^{\perp\perp}$, where $u, v \in x^\perp$ with $u \neq v$ [4].

5 Conclusion

By employing the concept of the Veldkamp space of the generalized quadrangle of order two, we were able to recognize other, on top of those examined in [1]–[3], distinguished subsets of generalized Pauli operators of two-level quantum systems, namely unicentric triads and pentads of them. It may well be that these two kinds of subsets of the two-qubit Pauli operators hold an important key for getting deeper insights into the nature of finite geometries underlying multiple higher-level quantum systems [9], in particular when the dimension of Hilbert space is not a power of a prime [10].

Acknowledgements

This work was partially supported by the Science and Technology Assistance Agency under the contract # APVT-51-012704, the VEGA grant agency projects # 2/6070/26 and # 7012 (all from Slovak Republic), the trans-national ECO-NET project # 12651NJ “Geometries Over Finite Rings and the Properties of Mutually Unbiased Bases” (France) and by the CNRS-SAV Project # 20246 “Projective and Related Geometries for Quantum Information” (France/Slovakia).

References

- [1] Saniga M, Planat M, and Pracna P. Projective ring line encompassing two-qubits. Theor Math Phys 2007, submitted; quant-ph/0611063.
- [2] Saniga M, and Planat M. Multiple qubits and symplectic polar spaces of order two. Adv Studies Theor Phys 2007;1:1–4; quant-ph/0612179.
- [3] Planat M, and Saniga M. On the Pauli graph of N -qudits, J Phys A: Math Theor 2007, submitted; quant-ph/0701211.
- [4] Payne SE, and Thas JA. Finite Generalized Quadrangles. Pitman: Boston–London–Melbourne; 1984.
- [5] Buekenhout F, and Cohen AM. Diagram Geometry (Chapter 10). Springer: New York; 2007 (to appear). Preprints of separate chapters can be found at <http://www.win.tue.nl/~amc/buek/>
- [6] Hirschfeld JWP, and Thas JA. General Galois Geometries. Oxford University Press: Oxford; 1991.
- [7] Lawrence J, Brukner Č, and Zeilinger A. Mutually unbiased binary observable sets on N qubits. Physical Review A 2002;65:032320.

- [8] Saniga M, Planat M, and Rosu H. 4, Mutually unbiased bases and finite projective planes. J Opt B: Quantum Semiclass Opt 2004;6:L19–L20; math-ph/0403057.
- [9] Planat M, and Saniga M. Pauli graph and finite projective lines/geometries. Optics and Optoelectronics 2007, in press; quant-ph/0703154.
- [10] Thas K. The geometry of generalized Pauli operators of N -qudit Hilbert space. Quantum Information and Computation 2007, submitted.