



HAL
open science

Detection of water vapor on Jupiter with the Odin space telescope

T. Cavalié, F. Billebaud, E. Lellouch, N. Biver, M. Dobrijevic, J. Brillet, A. Lecacheux, A. Hjalmarson, A. Sandqvist, U. Frisk, et al.

► To cite this version:

T. Cavalié, F. Billebaud, E. Lellouch, N. Biver, M. Dobrijevic, et al.. Detection of water vapor on Jupiter with the Odin space telescope. Odin Astronomy Workshop in Paris, 16-17 March, 2007, 2007, Paris, France. <hal-00139629>

HAL Id: hal-00139629

<https://hal.science/hal-00139629v1>

Submitted on 2 Apr 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Detection of water vapor on Jupiter with the Odin space telescope

T. Cavalié¹, F. Billebaud¹, E. Lellouch², N. Biver², M. Dobrijévic¹, J. Brillet¹, A. Lecacheux², A. Hjalmarson³, A. Sandqvist⁴, U. Frisk⁵, M. Olberg³ and The Odin Team.

¹Université Bordeaux I, CNRS, OASU, LAB, France ; ²LESIA, Observatoire de Paris, France ; ³Onsala Space Observatory, Sweden ; ⁴Stockholm Observatory, Sweden ; ⁵Swedish Space Corporation, Sweden.

The Infrared Space Observatory (ISO) has detected water vapor in the stratospheres of the giant planets and Titan and CO₂ on Jupiter, Saturn and Neptune (Feuchtgruber et al. 1997, 1999, Lellouch et al. 1997). The presence of the atmospheric cold trap implies an external origin for H₂O (interplanetary dust (IDP), sputtering from the satellites and /or rings, large meteoritic impacts). The sources of water on Jupiter could either be IDP or the Shoemaker-Levy 9 (SL9) comet's impacts (Moses et al. 2000, Lellouch et al. 2002).

The H₂O submillimetric line at 557 GHz was detected by the Submillimeter-Wave Astronomy Satellite (SWAS) in 1999 and 2001 (Bergin et al. 2000, Lellouch et al. 2002), but the vertical profile and the column density derived from the observations are different from the one obtained from ISO measurements (Lellouch et al. 2002). SWAS measurements favored an IDP source, whereas ISO observations favored a SL9 source.

The Swedish sub-millimeter satellite Odin carries out a long lasting monitoring of Jupiter's H₂O (110-101) 557 GHz line, since its launch in 2001. As an example, the high resolution H₂O spectrum obtained on November 8th, 2002, will be presented and discussed here. Both origins have been modeled with our photochemical model (Ollivier et al. 2000, adapted for Jupiter). Spectral analysis shows that a SL9 model gives better fits to the Odin and SWAS data than the IDP model. The column density derived is $(2.7 \pm 0.5) 10^{15} \text{ cm.s}^{-2}$. The initial deposition level of the water by SL9 is found to be at (0.5 ± 0.1) mbar, *i.e.* below the level inferred by Lellouch et al. (2002).

References : Feuchtgruber et al. (1997), *Nature*, 389, 159-162. Feuchtgruber et al. (1999), *The Universe as Seen by ISO. Eds. P. Cox & M. F. Kessler. ESA-SP, 427*, 133. Lellouch et al. (1997), *BAAS*, 29, 992. Moses et al. (2000), *Icarus*, 145, 166-202. Lellouch et al. (2002), *Icarus*, 159, 112-131. Bergin et al. (2000), *ApJ*, 539, L147-L150. Ollivier et al. (2000), *Plan. Space Sci.*, 48, 699-716.