



HAL
open science

Remark on an example by R.Schoen concerning the scalar curvature

Lionel Bérard Bergery, Guy Kass

► **To cite this version:**

Lionel Bérard Bergery, Guy Kass. Remark on an example by R.Schoen concerning the scalar curvature. 2003. hal-00143485

HAL Id: hal-00143485

<https://hal.science/hal-00143485>

Preprint submitted on 25 Apr 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Remarks on an example by R. Schoen concerning the scalar curvature

Lionel Berard Bergery^a and Guy Kass^b

^a Institut Elie Cartan, Université Henri Poincaré Nancy 1, 54506 Vandoeuvre les Nancy, France

^b Laboratoire de Mathématique, Université du Luxembourg, 1511 Luxembourg, Luxembourg

Abstract:

Classification: 53C21, 53A30

Key words:

1. Introduction

Let M be a compact Riemannian manifold of dimension $n \geq 3$. If g is a Riemannian metric on M , we denote by s_g the scalar curvature and by μ_g the canonical measure of the Riemannian manifold (M, g) . Moreover, let $V(g) = \int_M \mu_g$ be the volume, and $S(g) = \int_M s_g \mu_g$ the total scalar curvature of (M, g) . We denote by $[g]$ the conformal class of g :

$$[g] = \{fg/f \text{ smooth positive function on } M\},$$

and by $C(M)$ the set of all conformal classes of Riemannian metrics on M . Let $Riem(M)$ be the set of all Riemannian metrics on M . The Einstein-Hilbert functional $\sigma : Riem(M) \rightarrow \mathbb{R}$ is given by

$$\sigma(g) = \frac{S(g)}{(V(g))^{\frac{n-2}{n}}}.$$

The volume power in this definition is chosen so that σ is homogeneous. Generally the functional σ is neither bounded from above nor below, but its restriction to every conformal class $C \in C(M)$ is bounded from below; the constant

$$\mu_C = \inf_{g \in C} \sigma(g)$$

is called the Yamabe constant of the class C . Moreover, the infimum is always achieved: there exists a metric $g \in C$ such that $\mu_C = \sigma(g)$; such metrics are called Yamabe metrics. Every Yamabe metric g is a solution of the *Yamabe problem*, that is s_g is constant.

The conformal group of (M, g) is the set of diffeomorphisms ϕ of M such that

$$\forall g' \in [g], \phi^* g' \in [g].$$

Let G be a compact subgroup of the conformal group of (M, g) , and let

$$\mu_G = \inf_{g' \in [g], g' \text{ } G\text{-invariant}} \sigma(g').$$

Then there exists a conformal G -invariant metric g' to g such that

$$\sigma(g') = \mu_G,$$

and so, $s_{g'}$ is constant ([Hebey x]).

2. Position of the problem

Let n be an integer ≥ 3 . Let (N, g_N) be a compact connected $n - 1$ dimensional Riemannian manifold. We denote its scalar curvature by s_N and its volume by V_N . We suppose that the scalar curvature of this manifold is constant positive, equal to the one of the canonical sphere S^{n-1} , so we have

$$s_N = (n - 1)(n - 2).$$

On the other hand we consider the circle with length $l > 0$, that is the Riemannian manifold $(S^1(l) = \mathbb{R}/l\mathbb{Z}, dt^2)$, where t denotes the parameter on $S^1(l)$. We take the product manifold $M_l = S^1(l) \times N$ and we consider on M_l the metric g_l given by

$$g_l = dt^2 + g_N.$$

Then we have $V(g_l) = lV_N$, $s_{g_l} = s_N$, and $\sigma(g_l) = s_N(V_N)^{\frac{2}{n}} l^{\frac{2}{n}}$.

Let G_l be the set of isometries of M_l leaving N invariant. Clearly G_l is a compact subgroup of the conformal group of (M_l, g_l) and the set H_l of G_l -invariant metrics that are conformal to g_l is the set of metrics that are conformal to g_l with a ratio depending only on the variable $t \in S^1(l)$:

$$H_l = \{(f \times id_N)g_l/f \in C^\infty(S^1(l), \mathbb{R}_+^*)\}.$$

It follows from the result of Hebey stated in the introduction, that there exists $g' \in H_l$ such that

$$\sigma(g') = \mu_{G_l},$$

and $s_{g'}$ is constant. It follows from the fact that σ is homogeneous that for each $k > 0$ we have

$$\mu_{G_l} = \inf_{g' \in H_l, s_{g'} \equiv k} \sigma(g').$$

We will consider a module of the form $h^{-2} \times id_N$ with $h \in C^\infty(S^1(l), \mathbb{R}_+^*)$ and denote it simply by h^{-2} . Then we have $V(h^{-2}g_l) = V_N \int_0^l h(t)^{-n} dt$, so if $s_{h^{-2}g_l}$ is constant we get

$$\sigma(h^{-2}g_l) = s_{h^{-2}g_l} (V_N)^{\frac{2}{n}} \left(\int_0^l h(t)^{-n} dt \right)^{\frac{2}{n}}.$$

We will determine all the functions h such that $s_{h^{-2}g_l} \equiv n(n-1)$, and study the behavior of the function

$$\mu : l \mapsto \mu_{G_l}.$$

3. Reduction to an ordinary differential equation and resolution

We consider the metrics of the form $h^{-2}g_l$, where h denotes a positive function on $S^1(l)$. That function h may be considered as a function on \mathbb{R} having period l .

We have $h^{-2}g_l = u^{-\frac{4}{n-2}}g_2$, where $u = h^{\frac{2-n}{2}}$ and $g_2 = dt^2 + g_N$. Clearly $s_{g_2} = s_N$. Using (1.161 a)) p.59 in [Besse], we see that the scalar curvatures of $h^{-2}g_l$ and g_2 are related by

$$s_{h^{-2}g_l} = u^{-\frac{4}{n-2}}s_{g_2} - \frac{4(n-1)}{n-2}u^{-\frac{n+2}{n-2}}u''.$$

We have $u' = \frac{2-n}{2}h^{-\frac{n}{2}}h'$ and $u'' = \frac{2-n}{2}h^{-\frac{n+2}{2}}(hh'' - \frac{n}{2}h'^2)$, hence

$$s_{h^{-2}g_l} = h^2s_N + (n-1)(2hh'' - nh'^2).$$

We have $s_N = (n-1)(n-2)$, and we impose the condition $s_{h^{-2}g_l} \equiv n(n-1)$. So we get the following differential equation for the module h :

$$(n-2)h^2 + 2hh'' - nh'^2 = n. \tag{1}$$

Of course a first solution is given by the case where h is constant equal to $\sqrt{\frac{n}{n-2}}$; in this case we have

$$\sigma(h^{-2}g_l) = \sigma(g_l) = (n-1)(n-2)(V_N)^{\frac{2}{n}}l^{\frac{2}{n}}.$$

From now on we will look for non-constant solutions h . After multiplication by h' the differential equation (1) becomes $(n-2)h^2h' + h^{n+1}(h^{-n}h'^2)' = nh'$, that is $(n-2)h^{1-n}h' + (h^{-n}h'^2)' = nh^{-n-1}h'$, and finally

$$(-h^{2-n} + h^{-n}h'^2 + h^{-n})' = 0.$$

So there exists a constant K such that $-h^{2-n} + h^{-n}h'^2 + h^{-n} + K = 0$, that is

$$h'^2 = -Kh^n + h^2 - 1. \quad (2)$$

For $K = 0$ the equation becomes $h'^2 = h^2 - 1$ whose solutions $h(t) = \cosh(t + c)$ are not periodic. For $K \neq 0$ we put

$$F_K(s) = -Ks^n + s^2 - 1.$$

Let $K < 0$. Then the derivative $F_K'(s) = -nKs^{n-1} + 2s$ is positive for $s > 0$, so equation (2) has no periodic solution.

Now let $K > 0$. Then the derivative F_K' vanishes at the point $s_1 = (\frac{2}{nK})^{\frac{1}{n-2}}$. The function F_K increases from -1 to $F_K(s_1)$ on the interval $[0, s_1]$ and decreases from $F_K(s_1)$ to $-\infty$ on the interval $[s_1, +\infty[$. We have

$$F_K(s_1) > 0 \Leftrightarrow K < \frac{2}{n} \left(\frac{n-2}{n} \right)^{\frac{n-2}{2}}.$$

So if $K \geq \frac{2}{n} \left(\frac{n-2}{n} \right)^{\frac{n-2}{2}}$, we have no solution of equation (2). Let $K < \frac{2}{n} \left(\frac{n-2}{n} \right)^{\frac{n-2}{2}}$. Then the function F_K vanishes at two points $0 < s_0 < s_m$, and is positive on the interval $]s_0, s_m[$. Equation (2) is now equivalent to

$$\frac{dh}{\sqrt{F_K(h)}} = \pm dt,$$

that is $\eta_K(h) = \pm t + c$, where

$$\eta_K(h) = \int_{s_0}^h \frac{ds}{\sqrt{F_K(s)}}.$$

The function η_K is well defined on the interval $[s_0, s_m]$ since s_0 and s_m are simple roots of the polynomial F_K . We have $\eta_K'(h) = \frac{1}{\sqrt{F_K(h)}}$, so the function η_K increases from 0 to $\eta_K(s_m)$ on the interval $[s_0, s_m]$ and its derivatives at s_0 and s_m are $+\infty$. Hence η_K admits an inverse function η_K^{-1} which increases from s_0 to s_m on the interval $[0, \eta_K(s_m)]$ and whose derivatives at 0 and $\eta_K(s_m)$ vanish. Equation (2) becomes

$$h(t) = \eta_K^{-1}(\pm t + c).$$

It follows that the positive periodic solutions of equation (2) are the functions $h_{K,c} : t \mapsto h_K(t + c)$ where h_K is the even, $2\eta_K(s_m)$ -periodic function given by $h_K(t) = \eta_K^{-1}(t)$ for $t \in [0, \eta_K(s_m)]$. We are looking for solutions having period l , that is for functions whose

smallest positive period is of the form l/k where k is a positive integer. So the non-constant solutions of our problem are the functions $h_{K,c}$ for which there exists a positive integer k such that

$$\eta_K(s_m) = \frac{l}{2k}. \quad (3)$$

For such solution $h_{K,c}$ we have

$$\begin{aligned} \sigma(h_{K,c}^{-2} g_l) &= n(n-1)(V_N)^{\frac{2}{n}} \left(\int_0^{2k\eta_K(s_m)} h_K(t)^{-n} dt \right)^{\frac{2}{n}} \\ &= n(n-1)(V_N)^{\frac{2}{n}} (2k)^{\frac{2}{n}} \left(\int_0^{\eta_K(s_m)} h_K(t)^{-n} dt \right)^{\frac{2}{n}} \\ &= n(n-1)(V_N)^{\frac{2}{n}} \left(\frac{l}{\eta_K(s_m)} \right)^{\frac{2}{n}} \left(\int_0^{\eta_K(s_m)} \frac{h_K(t)^{-n} h'_K(t)}{\sqrt{F_K(h_K(t))}} dt \right)^{\frac{2}{n}} \\ &= n(n-1)(V_N)^{\frac{2}{n}} \left(\frac{l}{\eta_K(s_m)} \right)^{\frac{2}{n}} \left(\int_{s_0}^{s_m} \frac{t^{-n}}{\sqrt{F_K(t)}} dt \right)^{\frac{2}{n}}. \end{aligned}$$

Let $x = K^{\frac{1}{n-2}} s$, $x_0 = K^{\frac{1}{n-2}} s_0$ and $x_m = K^{\frac{1}{n-2}} s_m$. Then $F_K(s) = K^{-\frac{2}{n-2}} (-x^n + x^2 - K^{\frac{2}{n-2}})$, so we get

$$\eta_K(s_m) = \int_{x_0}^{x_m} \frac{dx}{\sqrt{-x^n + x^2 - K^{\frac{2}{n-2}}}}$$

and

$$\int_{s_0}^{s_m} \frac{t^{-n}}{\sqrt{F_K(t)}} dt = K^{\frac{n}{n-2}} \int_{x_0}^{x_m} \frac{x^{-n} dx}{\sqrt{-x^n + x^2 - K^{\frac{2}{n-2}}}.$$

Clearly x_0 and x_m are the roots of the polynomial $-x^n + x^2 - K^{\frac{2}{n-2}}$, so

$$K = (x_0^2 - x_0^n)^{\frac{n-2}{2}} = (x_m^2 - x_m^n)^{\frac{n-2}{2}}.$$

The function $x_0 \mapsto K$ is an increasing bijection from $]0, r[$ to $]0, \frac{2}{n} \left(\frac{n-2}{n} \right)^{\frac{n-2}{2}} [$, where

$$r = \left(\frac{2}{n} \right)^{\frac{1}{n-2}}.$$

So, the condition $0 < K < \frac{2}{n} \left(\frac{n-2}{n} \right)^{\frac{n-2}{2}}$ is equivalent to $0 < x_0 < r$. For $x_0 \in]0, r[$ we set

$$P_{x_0}(x) = -x^n + x^2 + x_0^n - x_0^2$$

and

$$\varphi(x_0) = \int_{x_0}^{x_m} \frac{dx}{\sqrt{P_{x_0}(x)}}, \quad \psi(x_0) = (x_0^2 - x_0^n)^{\frac{n}{2}} \int_{x_0}^{x_m} \frac{x^{-n} dx}{\sqrt{P_{x_0}(x)}}.$$

Condition (3) becomes

$$\varphi(x_0) = \frac{l}{2k}$$

and we get

$$\sigma(h_{K,c}^{-2} g_l) = n(n-1)(V_N)^{\frac{2}{n}} \left(\frac{\psi(x_0)}{\varphi(x_0)} \right)^{\frac{2}{n}} l^{\frac{2}{n}}.$$

4. Variations of the function φ

We have the factorization

$$P_{x_0}(x) = (x - x_0)(x_m - x)Q_{x_0}(x),$$

with

$$\begin{aligned} Q_{x_0}(x) &= x^{n-2} + (x_m + x_0)x^{n-3} + (x_m^2 + x_m x_0 + x_0^2)x^{n-4} + \dots \\ &+ (x_m^{n-3} + x_m^{n-4}x_0 + \dots + x_0^{n-3})x + (x_m^{n-2} + x_m^{n-3}x_0 + \dots + x_0^{n-2} - 1) \\ &= \sum_{k=1}^{n-1} \frac{x_m^k - x_0^k}{x_m - x_0} x^{n-1-k} - 1. \end{aligned}$$

Let $x = x_0 \cos^2 t + x_m \sin^2 t$. Then $(x - x_0)(x_m - x) = (x_m - x_0)^2 \sin^2 t \cos^2 t$, and $dx = (x_m - x_0) \sin 2t dt$, so we get

$$\varphi(x_0) = 2 \int_0^{\frac{\pi}{2}} \frac{dt}{\sqrt{Q_{x_0}(x_0 \cos^2 t + x_m \sin^2 t)}}.$$

We have $\varphi(r) = \frac{\pi}{\sqrt{Q_r(r)}}$, and $Q_r(r) = [1 + 2 + \dots + (n-1)]r^{n-2} - 1 = n-2$, so

$$\varphi(r) = \frac{\pi}{\sqrt{n-2}}.$$

Using Fatou's lemma we get $\lim_{x_0 \rightarrow 0} \varphi(x_0) \geq 2 \int_0^{\frac{\pi}{2}} \frac{dt}{\sqrt{Q_0(\sin^2 t)}}$; we have $Q_0(x) = x^{n-2} + x^{n-3} + \dots + x = x \frac{1-x^{n-2}}{1-x}$, so $\sqrt{Q_0(\sin^2 t)} = \sin t \frac{\sqrt{1-\sin^{2n-4} t}}{\cos t}$, which shows that

$$\lim_{x_0 \rightarrow 0} \varphi(x_0) = +\infty.$$

The constant term of the polynomial $Q_{x_0}(x)$ is $Q_{x_0}(0) = \frac{P_{x_0}(0)}{-x_0 x_m} > 0$, so all the coefficients of the polynomial $Q_{x_0}(x)$ are positive. Moreover:

Lemma 1 *The coefficients of the polynomial $Q_{x_0}(x)$ are increasing functions of $x_0 \in [0, r]$.*

Proof We have to show that for every $k \in \{2, 3, \dots, n-1\}$, the function

$$F : x_0 \mapsto \frac{x_m^k - x_0^k}{x_m - x_0}$$

is increasing on $[0, r]$. We denote the derivative of x_m as a function of x_0 by x_m' . We have

$$F'(x_0) = \frac{(x_m - x_0)(kx_m^{k-1}x_m' - kx_0^{k-1}) - (x_m^k - x_0^k)(x_m' - 1)}{(x_m - x_0)^2},$$

so

$$F'(x_0) > 0 \iff [(k-1)x_m^k - kx_0x_m^{k-1} + x_0^k]x_m' + [(k-1)x_0^k - kx_0^{k-1}x_m + x_m^k] > 0$$

$$\iff \left[(k-1) - k\frac{x_0}{x_m} + \left(\frac{x_0}{x_m}\right)^k \right] x_m' + \left[(k-1)\left(\frac{x_0}{x_m}\right)^k - k\left(\frac{x_0}{x_m}\right)^{k-1} + 1 \right] > 0.$$

Let $\alpha = \frac{x_0}{x_m}$. It follows from relation $x_0^2 - x_0^n = x_m^2 - x_m^n$ that $x_m^{n-2} = \frac{1-\alpha^2}{1-\alpha^n}$; from this we see that the function $\alpha \mapsto x_0$ is an increasing bijection from $[0, 1]$ to $[0, r]$. The preceding condition becomes

$$[(k-1) - k\alpha + \alpha^k]x_m' + [(k-1)\alpha^k - k\alpha^{k-1} + 1] > 0.$$

The coefficient of x_m' is positive; indeed if we set $u(\alpha) = (k-1) - k\alpha + \alpha^k$, then $u(1) = 0$ and $u'(\alpha) = k(\alpha^{k-1} - 1) < 0$ for $0 < \alpha < 1$. Our condition becomes

$$-x_m' < \frac{(k-1)\alpha^k - k\alpha^{k-1} + 1}{(k-1) - k\alpha + \alpha^k}.$$

We denote the second member by $F_k(\alpha)$ and we will show that for every integer $k \geq 1$ we have

$$\forall \alpha \in]0, 1[, F_{k+1}(\alpha) < F_k(\alpha);$$

then it will be sufficient to show that $-x_m' < F_{n-1}(\alpha)$. We have

$$F_{k+1}(\alpha) < F_k(\alpha) \iff \frac{k\alpha^{k+1} - (k+1)\alpha^k + 1}{k - (k+1)\alpha + \alpha^{k+1}} < \frac{(k-1)\alpha^k - k\alpha^{k-1} + 1}{(k-1) - k\alpha + \alpha^k}$$

$$\Leftrightarrow -\alpha^{2k+1} + \alpha^{2k} + k^2\alpha^{k+2} + (2 - 3k^2)\alpha^{k+1} + (3k^2 - 2)\alpha^k - k^2\alpha^{k-1} - \alpha + 1 > 0$$

$$\Leftrightarrow (1 - \alpha) + \alpha^{2k}(1 - \alpha) - k^2\alpha^{k-1}(1 - \alpha^3) + (3k^2 - 2)\alpha^k(1 - \alpha) > 0$$

$$\Leftrightarrow (1 - \alpha)[(\alpha^k - 1)^2 - k^2\alpha^{k-1}(\alpha - 1)^2] > 0$$

$$\Leftrightarrow (1 - \alpha)^3 \left[\left(\frac{1 - \alpha^k}{1 - \alpha} \right)^2 - k^2\alpha^{k-1} \right] > 0$$

$$\Leftrightarrow \frac{1 - \alpha^k}{1 - \alpha} - k\alpha^{\frac{k-1}{2}} > 0$$

$$\Leftrightarrow 1 - \alpha^k - k\alpha^{\frac{k-1}{2}}(1 - \alpha) > 0 \quad (4)$$

$$\Leftrightarrow 1 - \alpha^k - k\alpha^{\frac{k-1}{2}} + k\alpha^{\frac{k+1}{2}} > 0.$$

Let $f(\alpha) = 1 - \alpha^k - k\alpha^{\frac{k-1}{2}} + k\alpha^{\frac{k+1}{2}}$. Then $f'(\alpha) = -k\alpha^{\frac{k-3}{2}}g(\alpha)$ with $g(\alpha) = \alpha^{\frac{k+1}{2}} + \frac{k-1}{2}\alpha - \frac{k+1}{2}\alpha$; as $g(1) = 0$ and $g'(\alpha) = \frac{k+1}{2}(\alpha^{\frac{k-1}{2}} - 1) < 0$ we have $g(\alpha) > 0$ for $0 < \alpha < 1$, hence $f'(\alpha) < 0$; as $f(1) = 0$ we have finally that $f(\alpha) > 0$ for $0 < \alpha < 1$.

Let us show now that $-x_m' < F_{n-1}(\alpha)$. It follows from relation $x_m^n - x_m^2 = x_0^n - x_0^2$ that the derivative x_m' satisfies $(nx_m^{n-1} - 2x_m)x_m' = nx_0^{n-1} - 2x_0$, so

$$\begin{aligned} x_m' &= \frac{2x_0 - nx_0^{n-1}}{2x_m - nx_m^{n-1}} = \frac{x_0}{x_m} \frac{2 - nx_0^{n-2}}{2 - nx_m^{n-2}} \\ &= \alpha \frac{2 - n\alpha^{n-2} \frac{1-\alpha^2}{1-\alpha^n}}{2 - n \frac{1-\alpha^2}{1-\alpha^n}} \\ &= \alpha \frac{(n-2)\alpha^n - n\alpha^{n-2} + 2}{-2\alpha^n + n\alpha^2 - (n-2)}. \end{aligned}$$

The denominator is negative because $2x_m - nx_m^{n-1} = nx_m(r^{n-2} - x_m^{n-2}) < 0$. Hence we get

$$-x_m' < F_{n-1}(\alpha) \Leftrightarrow \alpha \frac{(n-2)\alpha^n - n\alpha^{n-2} + 2}{2\alpha^n - n\alpha^2 + (n-2)} < \frac{(n-2)\alpha^{n-1} - (n-1)\alpha^{n-2} + 1}{\alpha^{n-1} - (n-1)\alpha + (n-2)}$$

$$\begin{aligned} &\Leftrightarrow \alpha[(n-2)\alpha^n - n\alpha^{n-2} + 2][\alpha^{n-1} - (n-1)\alpha + (n-2)] \\ &- [(n-2)\alpha^{n-1} - (n-1)\alpha^{n-2} + 1][2\alpha^n - n\alpha^2 + (n-2)] < 0 \end{aligned}$$

$$\iff -(n-2)(\alpha-1)^2[1-\alpha^{2n-2}-(n-1)\alpha^{n-2}(1-\alpha^2)] < 0.$$

Comparing with (4) we see that the expression between square brackets is positive, which ends the proof of lemma 1.

Proposition 1 *The function φ is decreasing on the interval $]0, r[$.*

Proof We have

$$\varphi(x_0) = 2 \int_0^{\frac{\pi}{4}} \left(\frac{1}{\sqrt{Q_{x_0}(x_0 \cos^2 t + x_m \sin^2 t)}} + \frac{1}{\sqrt{Q_{x_0}(x_0 \sin^2 t + x_m \cos^2 t)}} \right) dt.$$

Hence it is enough to show that for all $t \in]0, \frac{\pi}{4}[$, the function

$$x_0 \mapsto \frac{1}{\sqrt{Q_{x_0}(x_0 \cos^2 t + x_m \sin^2 t)}} + \frac{1}{\sqrt{Q_{x_0}(x_0 \sin^2 t + x_m \cos^2 t)}}$$

is decreasing on $]0, r[$. For fixed $t \in]0, \frac{\pi}{4}[$ we set

$$\begin{aligned} u(x_0) &= Q_{x_0}(x_0 \cos^2 t + x_m \sin^2 t), \\ v(x_0) &= Q_{x_0}(x_0 \sin^2 t + x_m \cos^2 t). \end{aligned}$$

It follows from the fact that the coefficients of the polynomial $Q_{x_0}(x)$ are positive, that $0 < u < v$. From lemma 1 we have in particular that $x_m' > -1$, so

$$(x_0 \cos^2 t + x_m \sin^2 t)' = \cos^2 t + x_m' \sin^2 t > \cos^2 t - \sin^2 t > 0,$$

and it follows from lemma 1 that $u' > 0$. We will show that $(u+v)' > 0$. Then we get $(uv)' = u(v' + \frac{v}{u}u') > u(v' + u') > 0$, so uv is increasing, and $\frac{2}{\sqrt{uv}}$ is decreasing; moreover

$$\left(\frac{1}{u} + \frac{1}{v} \right)' = -\frac{1}{v^2} \left(\frac{v^2}{u^2} u' + v' \right) < 0,$$

so $\frac{1}{u} + \frac{1}{v}$ is decreasing; now $(\frac{1}{\sqrt{u}} + \frac{1}{\sqrt{v}})^2 = \frac{1}{u} + \frac{1}{v} + \frac{2}{\sqrt{uv}}$ is decreasing, so $\frac{1}{\sqrt{u}} + \frac{1}{\sqrt{v}}$ is decreasing.

Let us show that $(u+v)' > 0$. Let $x_0 = \alpha x_m$, $a = \alpha \cos^2 t + \sin^2 t$, $b = \cos^2 t + \alpha \sin^2 t$. Then

$$u + v = \frac{1}{\sin^2 t \cos^2 t} \frac{(1+\alpha)(\alpha^n - a^n + 1 - b^n)}{(1-\alpha)(1-\alpha^n)} - \frac{1 + \cos(2t)}{\cos^2 t},$$

so it is enough to show that the function

$$f_t : \alpha \mapsto \frac{(1+\alpha)(\alpha^n - a^n + 1 - b^n)}{(1-\alpha)(1-\alpha^n)}$$

is increasing. Straightforward calculation gives the following

$$f'_t(\alpha) = \frac{F_\alpha(t)}{(1-\alpha)^2(1-\alpha^n)^2}$$

where

$$\begin{aligned} F_\alpha(t) &= -2\alpha^{2n} - 2n\alpha^{n+1} + 2n\alpha^{n-1} + 2 \\ &+ (a^n + b^n)(n\alpha^{n+1} + 2\alpha^n - n\alpha^{n-1} - 2) + n(a^{n-1}a' + b^{n-1}b')(-\alpha^{n+2} + \alpha^n + \alpha^2 - 1). \end{aligned}$$

We have to show that $F_\alpha(t) > 0$ for every $\alpha \in]0, 1[$ and every $t \in]0, \frac{\pi}{4}[$. For $t = 0$ we have $a = \alpha, b = 1, a' = 1, b' = 0$, and so

$$F_\alpha(0) = 0.$$

Hence it is enough to show that $F'_\alpha(t) > 0$ for $t \in]0, \frac{\pi}{4}[$. We get

$$F'_\alpha(t) = -n(1-\alpha)^2 \sin(2t)G_\alpha(t)$$

where

$$\begin{aligned} G_\alpha(t) &= [(n-1)\alpha^n + n\alpha^{n-1} + 1](a^{n-1} - b^{n-1}) \\ &+ (n-1)(1+\alpha)(1-\alpha^n)(a^{n-2} \cos^2 t - b^{n-2} \sin^2 t); \end{aligned}$$

but

$$\begin{aligned} a^{n-1} - b^{n-1} &= a^{n-2}(\alpha \cos^2 t + \sin^2 t) - b^{n-2}(\cos^2 t + \alpha \sin^2 t) \\ &= \alpha(a^{n-2} \cos^2 t - b^{n-2} \sin^2 t) + (a^{n-2} \sin^2 t - b^{n-2} \cos^2 t), \end{aligned}$$

so

$$G_\alpha(t) = a^{n-2}(A \cos^2 t + B \sin^2 t) - b^{n-2}(A \sin^2 t + B \cos^2 t),$$

with

$$A = \alpha^n - n\alpha + n - 1, \quad B = (n-1)\alpha^n + n\alpha^{n-1} + 1.$$

We have to show that $G_\alpha(t) < 0$. We have

$$G'_\alpha(t) = \sin(2t) G_{\alpha,1}(t),$$

with

$$\begin{aligned} G_{\alpha,1}(t) &= (B-A)(a^{n-2} + b^{n-2}) \\ &+ (n-2)(1-\alpha)[a^{n-3}(A \cos^2 t + B \sin^2 t) + b^{n-3}(A \sin^2 t + B \cos^2 t)]. \end{aligned}$$

We have

$$G'_{\alpha,1}(t) = (n-2)(1-\alpha) \sin(2t) G_{\alpha,2}(t),$$

with

$$G_{\alpha,2}(t) = 2(B-A)(a^{n-3} - b^{n-3}) \\ + (n-3)(1-\alpha)[a^{n-4}(A \cos^2 t + B \sin^2 t) - b^{n-4}(A \sin^2 t + B \cos^2 t)].$$

For $p \in \{1, 2, \dots, n-1\}$ let

$$G_{\alpha,p}(t) = p(B-A)(a^{n-1-p} + (-1)^{p+1}b^{n-1-p}) \\ + (n-1-p)(1-\alpha)[a^{n-2-p}(A \cos^2 t + B \sin^2 t) + (-1)^{p+1}b^{n-2-p}(A \sin^2 t + B \cos^2 t)].$$

Then

$$G'_{\alpha,p}(t) = (n-1-p)(1-\alpha) \sin(2t) G_{\alpha,p+1}(t).$$

Clearly $G_{\alpha}(\frac{\pi}{4}) = 0$, and for even p we have $G_{\alpha,p}(\frac{\pi}{4}) = 0$. We have the following

Lemma 2 For every $\alpha \in]0, 1[$ we have

- a) If $n = 3$, then $G_{\alpha,1}(0) > 0$.
If $n = 4$, then $G_{\alpha,1}(0) = 0$.
If $n \geq 5$, and if p is odd, $p \leq n-2$, then $G_{\alpha,p}(0) < 0$.
- b) $G_{\alpha,1}(\pi/4) > 0$.
If p is odd, $\frac{n}{2} \leq p \leq n-1$, then $G_{\alpha,p}(\pi/4) < 0$.
If p is odd, $3 \leq p \leq n-1$, then $[G_{\alpha,p}(\pi/4) \geq 0 \Rightarrow G_{\alpha,p-2}(\pi/4) > 0]$.
- c) $G_{\alpha}(0) < 0$.

Proof a) If p is odd, we have

$$G_{\alpha,p}(0) = p(B-A)(a^{n-1-p} + 1) + (n-1-p)(1-\alpha)(a^{n-2-p}A + B) \\ = (n-1)[(p-1)\alpha^{2n-1-p} + (p+1)\alpha^{2n-2-p} - (n-1-p)\alpha^{n+1} + (p-1)\alpha^n + n\alpha^{n-1} \\ - n\alpha^{n-p} - (p-1)\alpha^{n-1-p} + (n-1-p)\alpha^{n-2-p} - (p+1)\alpha - (p-1)],$$

in particular

$$G_{\alpha,1}(0) = \alpha(n-1)[2\alpha^{2n-4} - (n-2)\alpha^n + (n-2)\alpha^{n-4} - 2].$$

For $n = 3$ we get $G_{\alpha,1}(0) = 2(1-\alpha^2)(1-\alpha)^2 > 0$. For $n = 4$ we get $G_{\alpha,1}(0) = 0$. Let now $n \geq 5$. First we show that $G_{\alpha,1}(0) < 0$. We have

$$G_{\alpha,1}(0) = \alpha(n-1)f(\alpha)$$

with

$$f(\alpha) = 2\alpha^{2n-4} - (n-2)\alpha^n + (n-2)\alpha^{n-4} - 2.$$

We get

$$f'(\alpha) = (n-2)\alpha^{n-5}h(\alpha)$$

with

$$h(\alpha) = 4\alpha^n - n\alpha^4 + n - 4.$$

We have $h(1) = 0$ and $h'(\alpha) = 4n\alpha^3(\alpha^{n-4} - 1) < 0$, so $h(\alpha) > 0$ and f is increasing; as $f(1) = 0$, we have $f(\alpha) < 0$ and $G_{\alpha,1}(0) < 0$.

Let $G_{\alpha,p}(0) = (n-1)h_{n,p}(\alpha)$, then

$$h_{n+1,p+1}(\alpha) - h_{n,p}(\alpha) = p u(\alpha) + n v(\alpha) + w(\alpha)$$

with

$$u(\alpha) = \alpha^{2n-p} - \alpha^{2n-2-p} + \alpha^{n+2} - \alpha^n = (\alpha^2 - 1)(\alpha^{2n-2-p} + \alpha^n) < 0,$$

$$v(\alpha) = -\alpha^{n+2} + \alpha^{n+1} + \alpha^n - \alpha^{n-1} = \alpha^{n-1}(1 - \alpha)(\alpha^2 - 1) < 0,$$

$$w(\alpha) = 3\alpha^{2n-1-p} - \alpha^{2n-2-p} + \alpha^{n+2} - \alpha^{n+1} + 2\alpha^n - \alpha^{n-p} - \alpha^{n-1-p} - \alpha - 1$$

$$= (\alpha - 1)(\alpha^{2n-2-p} + \alpha^{n+1}) + (\alpha^{n-1} - 1)(\alpha^{n-p} + \alpha) + (\alpha^n - 1)(\alpha^{n-1-p} + 1) < 0,$$

which implies that $h_{n+1,p+1}(\alpha) < h_{n,p}(\alpha)$. We know that $h_{n,1}(\alpha) = \frac{G_{\alpha,1}(0)}{n-1} < 0$ for all $n \geq 5$, $h_{4,1}(\alpha) = 0$, and we have $h_{4,2}(\alpha) = 4\alpha(\alpha+1)(\alpha^2-1) < 0$, which ends the proof of a).

b) If p is odd, we get

$$G_{\alpha,p}(\pi/4) = (n-1)\frac{(1+\alpha)^{n-1-p}}{2^{n-2-p}}f(\alpha)$$

with

$$f(\alpha) = -(n-2p)\alpha^n + n\alpha^{n-1} - n\alpha + n - 2p.$$

We have

$$f'(\alpha) = n[-(n-2p)\alpha^{n-1} + (n-1)\alpha^{n-2} - 1],$$

$$f''(\alpha) = n(n-1)\alpha^{n-3}[-(n-2p)\alpha + n - 2].$$

For $p = 1$ we have $f''(\alpha) = n(n-1)(n-2)\alpha^{n-3}(1-\alpha) > 0$ and $f'(1) = 0$, so $f' < 0$ on $]0, 1[$; as $f(1) = 0$, we get $f > 0$ on $]0, 1[$, that is $G_{\alpha,1}(\pi/4) > 0$.

For $p = n/2$ (in case where $n/2$ is odd) we have $f'(\alpha) = n[(n-1)\alpha^{n-2} - 1]$, so f' vanishes at $\alpha_0 = (n-2)^{-\frac{1}{n-2}}$ and f is decreasing on $[0, \alpha_0]$ and increasing on $[\alpha_0, 1]$; as $f(0) = f(1) = 0$ it follows that $f < 0$ on $]0, 1[$, so $G_{\alpha,p}(\pi/4) < 0$.

For $p > n/2$ we have $f'' > 0$ on $]0, 1[$; as $f'(0) = -n$ and $f'(1) = 2n(p-1)$, there exists $\alpha_0 \in]0, 1[$ such that f is decreasing on $[0, \alpha_0]$ and increasing on $[\alpha_0, 1]$; as $f(0) = n-2p < 0$ and $f(1) = 0$ it follows that $f < 0$ on $]0, 1[$, so $G_{\alpha,p}(\pi/4) < 0$.

Now let us suppose that $G_{\alpha,p}(\pi/4) \geq 0$. We have

$$f(\alpha) = 2p(\alpha^n - 1) - n\alpha^n + n\alpha^{n-1} - n\alpha + n,$$

and this expression is a strictly increasing function of p , from which we conclude that $G_{\alpha,p-2}(\pi/4) > 0$.

c) We have $G_\alpha(0) = \alpha^{n-2}A - B = \alpha^{2n-2} - (n-1)\alpha^n + (n-1)\alpha^{n-2} - 1$. Denoting this expression by $f(\alpha)$ we get $f'(\alpha) = (n-1)\alpha^{n-3}g(\alpha)$ with $g(\alpha) = 2\alpha^n - n\alpha^2 + n - 2$. As $g(1) = 0$ and $g'(\alpha) = 2n\alpha(\alpha^{n-2} - 1) < 0$, we have $g > 0$ on $]0, 1[$, and f is increasing. As $f(1) = 0$ we have $f < 0$ on $]0, 1[$, so $G_\alpha(0) < 0$.

In order to achieve the proof of proposition 1 we have to show that $G_\alpha(t) < 0$ for all $t \in]0, \pi/4[$, all $\alpha \in]0, 1[$ and all integer $n \geq 3$. We have

$$G_{\alpha,n-1}(t) = (n-1)(B-A)[1 + (-1)^n],$$

and

$$B - A = (n-2)(\alpha^n - 1) + n\alpha(\alpha^{n-2} - 1) < 0.$$

Let us consider first the case $n = 3$. Then $G_{\alpha,2}(t) = 0$, so $G_{\alpha,1}(t)$ is constant. By lemma 2 this constant is positive, so $G_\alpha(t)$ is increasing. As $G_\alpha(\pi/4) = 0$, we have $G_\alpha < 0$ on $]0, \pi/4[$.

Now let $n = 4$. Then $G_{\alpha,3}(t) = 6(B-A) < 0$, so $G_{\alpha,2}(t)$ is decreasing. As $G_{\alpha,2}(\pi/4) = 0$, we have $G_{\alpha,2} > 0$ on $]0, \pi/4[$, so $G_{\alpha,1}(t)$ is increasing. As $G_{\alpha,1}(0) = 0$, we have $G_{\alpha,1} > 0$ on $]0, \pi/4[$, so $G_\alpha(t)$ is increasing. As $G_\alpha(\pi/4) = 0$, we have $G_\alpha < 0$ on $]0, \pi/4[$.

Now let $n \geq 5$, n odd. Then $G_{\alpha,n-1} = 0$, so $G_{\alpha,n-2}(t)$ is constant. As $G_{\alpha,n-2}(0) < 0$, this constant is negative. By lemma 2 there exists an odd integer p_0 (depending on α) such that $G_{\alpha,p}(\pi/4) \geq 0$ and such that for every odd integer p we have

$$p_0 < p \leq n-2 \Rightarrow G_{\alpha,p}(\pi/4) < 0$$

and

$$1 \leq p < p_0 \Rightarrow G_{\alpha,p}(\pi/4) > 0.$$

For every odd integer p such that $p_0 < p \leq n-2$ we then have:

$$G_{\alpha,p+1} : \searrow_0, G_{\alpha,p+1} > 0, G_{\alpha,p} : \nearrow^-, G_{\alpha,p} < 0,$$

so

$$G_{\alpha,p_0+1} : \searrow_0, G_{\alpha,p_0+1} > 0, G_{\alpha,p_0} : - \nearrow^+, G_{\alpha,p_0} : [-, +],$$

and for every odd integer p such that $1 \leq p < p_0$ we have:

$$G_{\alpha,p+1} : \searrow \nearrow^0, G_{\alpha,p+1} : [+,-], G_{\alpha,p} : - \nearrow \searrow_+, G_{\alpha,p} : [-,+],$$

so

$$G_{\alpha,1} : [-,+],$$

and finally

$$G_\alpha : - \searrow \nearrow^0,$$

so $G_\alpha < 0$ on $]0, \pi/4[$.

For $n \geq 5$, n even, we have $G_{\alpha,n-1} = 2(n-1)(B-A) < 0$, and like before we conclude that $G_\alpha < 0$ on $]0, \pi/4[$. ■

5. Study of the function μ

We have shown in the preceding paragraph that the function φ decreases from $+\infty$ to $\frac{\pi}{\sqrt{n-2}}$ on the interval $]0, r[$. The existence condition (3) for a non-constant solution of our problem stated in paragraph 2 is $\varphi(x_0) = \frac{l}{2k}$, (k integer ≥ 1).

For every $l > 0$ our problem has the constant solution $h_0 \equiv \sqrt{\frac{n}{n-2}}$. It satisfies

$$\sigma_0(l) := \sigma(h_0^{-2}g_l) = (n-1)(n-2)(V_N)^{\frac{2}{n}}l^{\frac{2}{n}}.$$

For $l \leq \frac{2\pi}{\sqrt{n-2}}$ this is the only solution, so

$$\mu(l) = (n-1)(n-2)(V_N)^{\frac{2}{n}}l^{\frac{2}{n}}.$$

Let $k \in \mathbb{N}^*$ and $k\frac{2\pi}{\sqrt{n-2}} < l \leq (k+1)\frac{2\pi}{\sqrt{n-2}}$. Then there exist $x_{0,i}(l) \in]0, r[$, $i = 1, 2, \dots, k$, such that $\varphi(x_{0,i}(l)) = \frac{l}{2i}$. The corresponding solutions $h_{l,i}$ satisfy

$$\begin{aligned} \sigma_i(l) &:= \sigma(h_{l,i}^{-2}g_l) = n(n-1)(V_N)^{\frac{2}{n}} \left(\frac{\psi(x_{0,i}(l))}{\varphi(x_{0,i}(l))} \right)^{\frac{2}{n}} l^{\frac{2}{n}} \\ &= n(n-1)(2iV_N)^{\frac{2}{n}} [\psi(x_{0,i}(l))]^{\frac{2}{n}}. \end{aligned}$$

Clearly

$$\mu(l) = \inf_{0 \leq i \leq k} \sigma_i(l).$$

Proposition 2 For $l > \frac{2\pi}{\sqrt{n-2}}$ we have $\mu(l) < \sigma_0(l)$.

Proof It is enough to show that there exists a smooth l -periodic function θ on \mathbb{R} and a (small) real number t such that the function

$$h(x) = 1 + t\theta(x)$$

is positive and satisfies $\sigma(h^{-2}g_l) < \sigma_0(l)$. For small t we have $h^{-n} \sim 1 - nt\theta + \frac{n(n+1)}{2}t^2\theta^2$, so we get (all integrals are taken on the interval $[0, l]$):

$$\int h^{-n} \sim l \left[1 - nt \frac{\int \theta}{l} + \frac{n(n+1)}{2} t^2 \frac{\int \theta^2}{l} \right],$$

and

$$\left(\int h^{-n} \right)^{-\frac{n-2}{n}} \sim l \left[1 - nt \frac{\int \theta}{l} + \frac{n(n+1)}{2} t^2 \frac{\int \theta^2}{l} \right].$$

Moreover

$$\begin{aligned} s_{h^{-2}g_l} &= h^2 s_N + (n-1)(2hh'' - nh'^2) \\ &= (n-1)(n-2)(1 + 2t\theta + t^2\theta^2) + (n-1)[2(1+t\theta)t\theta'' - nt^2\theta'^2] \\ &= (n-1)(n-2) \left[1 + t(2\theta + \frac{2}{n-2}\theta'') + t^2(\theta^2 + \frac{2}{n-2}\theta\theta'' - \frac{n}{n-2}\theta'^2) \right], \end{aligned}$$

so we get

$$s_{h^{-2}g_l} h^{-n} \sim (n-1)(n-2) \left\{ 1 + t \left[(2-n)\theta + \frac{2}{n-2}\theta'' \right] + t^2 \left[\frac{n^2-3n+2}{2}\theta^2 + 2\frac{n-1}{n-2}\theta\theta'' - \frac{n}{n-2}\theta'^2 \right] \right\}$$

and

$$\int s_{h^{-2}g_l} h^{-n} \sim (n-1)(n-2)l \left\{ 1 - (n-2)t \frac{\int \theta}{l} + t^2 \left[\frac{n^2-3n+2}{2} \frac{\int \theta^2}{l} + 2\frac{n-1}{n-2} \frac{\int \theta\theta''}{l} - \frac{n}{n-2} \frac{\int \theta'^2}{l} \right] \right\}.$$

We have $\sigma(h^{-2}g_l) = (V_N)^{\frac{2}{n}} (\int s_{h^{-2}g_l} h^{-n}) (\int h^{-n})^{-\frac{n-2}{n}}$, so

$$\sigma(h^{-2}g_l) \sim \sigma_0(l) [1 + \alpha(\theta)t^2]$$

with

$$\alpha(\theta) = \frac{1}{l^2} \{(n-2)[(f\theta)^2 - lf\theta^2] + lf\theta'^2\}.$$

Let $\theta(x) = \sin(\frac{2\pi}{l}x)$, then $\alpha(\theta) = \frac{2-n}{2} + \frac{2\pi^2}{l^2}$, so $\alpha(\theta) < 0 \iff l > \frac{2\pi}{\sqrt{n-2}}$. Hence this function θ satisfies our conditions. ■

It follows from proposition 2 that for $\frac{2\pi}{\sqrt{n-2}} < l \leq 2\frac{2\pi}{\sqrt{n-2}}$ we have $\mu(l) = \sigma_1(l)$. If $l \rightarrow \frac{2\pi}{\sqrt{n-2}}$, ($l > \frac{2\pi}{\sqrt{n-2}}$), then $x_{0,1}(l) = \varphi^{-1}(\frac{l}{2}) \rightarrow \varphi^{-1}(\frac{\pi}{\sqrt{n-2}}) = r$, and

$$\begin{aligned} \sigma_1(l) &\rightarrow n(n-1)(V_N)^{\frac{2}{n}} 2^{\frac{2}{n}} (\psi(r))^{\frac{2}{n}} \\ &= n(n-1)(V_N)^{\frac{2}{n}} 2^{\frac{2}{n}} (r^2 - r^n) \left(\frac{\pi}{r^n \sqrt{n-2}}\right)^{\frac{2}{n}} \\ &= n(n-1)(V_N)^{\frac{2}{n}} \left(1 - \frac{2}{n}\right) \left(\frac{2\pi}{\sqrt{n-2}}\right)^{\frac{2}{n}} \\ &= (n-1)(n-2)(V_N)^{\frac{2}{n}} \left(\frac{2\pi}{\sqrt{n-2}}\right)^{\frac{2}{n}} \\ &= \sigma_0\left(\frac{2\pi}{\sqrt{n-2}}\right), \end{aligned}$$

which shows that the function μ is continuous at the point $\frac{2\pi}{\sqrt{n-2}}$.

Proposition 3 *The function μ is differentiable at the point $\frac{2\pi}{\sqrt{n-2}}$.*

Proof For $l < \frac{2\pi}{\sqrt{n-2}}$ we have $\mu'(l) = \frac{2(n-1)(n-2)}{n} (V_n)^{\frac{2}{n}} l^{\frac{2-n}{n}}$, so

$$\mu'_{left}\left(\frac{2\pi}{\sqrt{n-2}}\right) = \frac{(n-1)(n-2)}{n} (2V_n)^{\frac{2}{n}} \left(\frac{\pi}{\sqrt{n-2}}\right)^{\frac{2-n}{n}}.$$

For $l > \frac{2\pi}{\sqrt{n-2}}$ we have $\mu(l) = \sigma_1(l) = n(n-1)(2V_n)^{\frac{2}{n}} (\psi(\varphi^{-1}(\frac{l}{2})))^{\frac{2}{n}}$, so

$$\mu'(l) = (n-1)(2V_n)^{\frac{2}{n}} (\psi(\varphi^{-1}(\frac{l}{2})))^{\frac{2-n}{n}} \frac{\psi'(\varphi^{-1}(\frac{l}{2}))}{\varphi'(\varphi^{-1}(\frac{l}{2}))},$$

and

$$\mu'_{right}\left(\frac{2\pi}{\sqrt{n-2}}\right) = (n-1)(2V_n)^{\frac{2}{n}} (\psi(r))^{\frac{2-n}{n}} \lim_{s \rightarrow r} \frac{\psi'(s)}{\varphi'(s)}.$$

From $x_0^2 - x_m^2 = x_0^n - x_m^n$ and $x_0 \neq x_m$ we get $x_0 + x_m = \sum_{i=0}^{n-1} x_0^{n-1-i} x_m^i$, so

$$1 + x'_m = \sum_{i=0}^{n-1} [(n-1-i)x_0^{n-2-i} x_m^i + ix_0^{n-1-i} x_m^{i-1} x'_m],$$

from which we get $1 + x'_m(r) = (n-1)[1 + x'_m(r)]$, so

$$x'_m(r) = -1.$$

Let $x_0 = r - h$, then $x_m = r + h + ah^2 + bh^3 + ch^4 + \dots$. Putting this in relation $x_0^2 - x_m^2 = x_0^n - x_m^n$, we get the values of the constants a, b, c :

$$x_m = r + h - \frac{n-1}{3r}h^2 + \left(\frac{n-1}{3r}\right)^2 h^3 - \frac{(n-1)(19n^2-23n+58)}{540r^3}h^4 + \dots$$

Let $x = x_0 \cos^2 t + x_m \sin^2 t$. Then we get after some calculations

$$Q_{x_0}(x) = (n-2)\left\{1 - \frac{n-1}{3r} \cos 2t \cdot h - \frac{n-1}{36r^2}[12 - 2(n-1) \cos 2t - 3(n-3) \cos^2 2t]h^2\right\} + \dots,$$

$$\frac{1}{\sqrt{Q_{x_0}(x)}} = \frac{1}{\sqrt{n-2}} \left\{1 + \frac{n-1}{6r} \cos 2t \cdot h + \frac{n-1}{72r^2}[3(n+2) - 2(n-1) \cos 2t + 3 \cos 4t]h^2\right\} + \dots,$$

$$\varphi(x_0) = \frac{\pi}{\sqrt{n-2}} \left[1 + \frac{(n-1)(n+2)}{24r^2}h^2\right] + \dots,$$

$$\psi(x_0) = \frac{\pi}{\sqrt{n-2}} \left(\frac{n-2}{n}\right)^{\frac{n}{2}} \left[1 + \frac{(n-1)(n+2)}{24r^2}h^2\right] + \dots,$$

so

$$\begin{aligned} \mu'_{right}\left(\frac{2\pi}{\sqrt{n-2}}\right) &= (n-1)(2V_n)^{\frac{2}{n}} \left(\frac{n-2}{n}\right)^{\frac{2-n}{2}} \left(\frac{\pi}{\sqrt{n-2}}\right)^{\frac{2-n}{n}} \frac{\psi''(r)}{\varphi''(r)} \\ &= (n-1)(2V_n)^{\frac{2}{n}} \left(\frac{n-2}{n}\right)^{\frac{2-n}{2}} \left(\frac{\pi}{\sqrt{n-2}}\right)^{\frac{2-n}{n}} \left(\frac{n-2}{n}\right)^{\frac{n}{2}} \\ &= \frac{(n-1)(n-2)}{n} (2V_n)^{\frac{2}{n}} \left(\frac{\pi}{\sqrt{n-2}}\right)^{\frac{2-n}{n}} \\ &= \mu'_{left}\left(\frac{2\pi}{\sqrt{n-2}}\right). \end{aligned}$$

Proposition 4 For $l > \frac{2\pi}{\sqrt{n-2}}$ we have $\sigma_1(l) < \sigma_0(l)$.

Proof As $\sigma_1(l) = n(n-1)(2V_N)^{\frac{2}{n}}(\psi(x_0))^{\frac{2}{n}}$ and $\sigma_0(l) = (n-1)(n-2)(2V_N)^{\frac{2}{n}}(\varphi(x_0))^{\frac{2}{n}}$, with $x_0 = \varphi^{-1}\left(\frac{l}{2}\right)$, we have to show that for all $x_0 \in]0, r[$,

$$\psi(x_0) < \left(1 - \frac{2}{n}\right)^{\frac{n}{2}} \varphi(x_0).$$

Let us recall that $\varphi(x_0) = \int_{x_0}^{x_m} \frac{dx}{\sqrt{P_{x_0}(x)}}$ and $\psi(x_0) = (x_0^2 - x_0^n)^{\frac{n}{2}} \int_{x_0}^{x_m} \frac{x^{-n} dx}{\sqrt{P_{x_0}(x)}}$. Let

$$s = \sqrt{\frac{x^2}{x_0^2 - x_0^n} - 1}, \quad s_0 = \sqrt{\frac{x_0^{n-2}}{1 - x_0^{n-2}}}, \quad s_m = \sqrt{\frac{x_m^{n-2}}{1 - x_m^{n-2}}};$$
 then

$$\begin{aligned}
\varphi(x_0) &= \int_{s_0}^{s_m} \frac{ds}{(1+s^2)^{\frac{1}{2}} \sqrt{1 - (x_0^2 - x_0^n)^{\frac{n-2}{2}} \frac{(1+s^2)^{\frac{n}{2}}}{s^2}}} \\
&= \sum_{p=0}^{+\infty} \frac{(2p)!}{2^{2p}(p!)^2} (x_0^2 - x_0^n)^{p \frac{n-2}{2}} \int_{s_0}^{s_m} \frac{(1+s^2)^{\frac{pn-1}{2}}}{s^{2p}} ds, \\
\psi(x_0) &= \int_{s_0}^{s_m} \frac{ds}{(1+s^2)^{\frac{n+1}{2}} \sqrt{1 - (x_0^2 - x_0^n)^{\frac{n-2}{2}} \frac{(1+s^2)^{\frac{n}{2}}}{s^2}}} \\
&= \sum_{p=0}^{+\infty} \frac{(2p)!}{2^{2p}(p!)^2} (x_0^2 - x_0^n)^{p \frac{n-2}{2}} \int_{s_0}^{s_m} \frac{(1+s^2)^{\frac{(p-1)n-1}{2}}}{s^{2p}} ds,
\end{aligned}$$

so it is enough to show that for all integer $p \geq 0$ and all $x_0 \in]0, r[$ we have

$$\int_{s_0}^{s_m} \frac{(1+s^2)^{\frac{(p-1)n-1}{2}}}{s^{2p}} ds < \left(1 - \frac{2}{n}\right)^{\frac{n}{2}} \int_{s_0}^{s_m} \frac{(1+s^2)^{\frac{pn-1}{2}}}{s^{2p}} ds.$$

Let

$$F_p(x_0) = \left(1 - \frac{2}{n}\right)^{\frac{n}{2}} \int_{s_0}^{s_m} \frac{(1+s^2)^{\frac{pn-1}{2}}}{s^{2p}} ds - \int_{s_0}^{s_m} \frac{(1+s^2)^{\frac{(p-1)n-1}{2}}}{s^{2p}} ds.$$

If $x_0 = r$, then $s_0 = s_m$, so $F(r) = 0$. Hence it is enough to show that the function F is strictly decreasing on $]0, r[$. we get

$$\frac{2}{n-2} (x_0^2 - x_0^n)^{1 + \frac{p(n-2)}{2}} F'_p(x_0) = \left[\left(1 - \frac{2}{n}\right)^{\frac{n}{2}} - (1 - x_m^{n-2})^{\frac{n}{2}}\right] x_m^{\frac{n}{2}} x'_m - \left[\left(1 - \frac{2}{n}\right)^{\frac{n}{2}} - (1 - x_0^{n-2})^{\frac{n}{2}}\right] x_0^{\frac{n}{2}},$$

so we have to show that

$$x'_m < \frac{x_0^{\frac{n}{2}}}{x_m^{\frac{n}{2}}} \frac{\left(1 - \frac{2}{n}\right)^{\frac{n}{2}} - (1 - x_0^{n-2})^{\frac{n}{2}}}{\left(1 - \frac{2}{n}\right)^{\frac{n}{2}} - (1 - x_m^{n-2})^{\frac{n}{2}}},$$

i.e.

$$\frac{x_0}{x_m} \frac{\left(1 - \frac{2}{n}\right) - (1 - x_0^{n-2})}{\left(1 - \frac{2}{n}\right) - (1 - x_m^{n-2})} < \left(\frac{x_0}{x_m}\right)^{\frac{n}{2}} \frac{\left(1 - \frac{2}{n}\right)^{\frac{n}{2}} - (1 - x_0^{n-2})^{\frac{n}{2}}}{\left(1 - \frac{2}{n}\right)^{\frac{n}{2}} - (1 - x_m^{n-2})^{\frac{n}{2}}}. \quad (5)$$

Let $\alpha = \frac{x_0}{x_m}$ and $u = \sqrt{\frac{n}{n-2} \frac{1-\alpha^{n-2}}{1-\alpha^n}}$. Clearly $u > 1$ and $\alpha u < 1$. Moreover we have $\alpha^{1/2} u < 1$. Inequality (5) becomes

$$\frac{1-u^2}{1-(\alpha u)^2} < \alpha^{\frac{n}{2}-1} \frac{1-u^n}{1-(\alpha u)^n},$$

i.e.

$$\left(\sum_{k=0}^{n-1} (\alpha u)^k \right) (1+u) - \alpha^{\frac{n}{2}-1} (1+\alpha u) \left(\sum_{k=0}^{n-1} u^k \right) > 0. \quad (6)$$

For even n this inequality reads

$$\left(\sum_{k=0}^{\frac{n-2}{2}} [(\alpha u)^k + (\alpha u)^{n-1-k}] \right) (1+u) - \alpha^{\frac{n}{2}-1} (1+\alpha u) \left(\sum_{k=0}^{\frac{n-2}{2}} (u^k + u^{n-1-k}) \right) > 0.$$

For each integer k such that $0 \leq k \leq \frac{n-2}{2}$ we have

$$\begin{aligned} & [(\alpha u)^k + (\alpha u)^{n-1-k}] (1+u) - \alpha^{\frac{n}{2}-1} (1+\alpha u) (u^k + u^{n-1-k}) \\ &= (\alpha u)^k [(1 + (\alpha u)^{n-1-2k}) (1+u) - \alpha^{\frac{n}{2}-1-k} (1+\alpha u) (1+u^{n-1-2k})] \\ &= (\alpha u)^k [(1 - \alpha^{\frac{n}{2}-1-k}) (1 - (\alpha^{1/2} u)^{n-2k}) + u (1 - \alpha^{\frac{n}{2}-k}) (1 - (\alpha^{1/2} u)^{n-2-2k})], \end{aligned}$$

which is positive as $\alpha^{1/2} u < 1$. This proves inequality (6) in the case n even. For odd n this inequality reads

$$\begin{aligned} & \left((\alpha u)^{\frac{n-1}{2}} + \sum_{k=0}^{\frac{n-3}{2}} [(\alpha u)^k + (\alpha u)^{n-1-k}] \right) (1+u) \\ & - \alpha^{\frac{n}{2}-1} (1+\alpha u) \left(u^{\frac{n-1}{2}} + \sum_{k=0}^{\frac{n-3}{2}} (u^k + u^{n-1-k}) \right) > 0, \end{aligned}$$

i.e.

$$\begin{aligned} & (\alpha u)^{\frac{n-1}{2}} (1+u) - \alpha^{\frac{n}{2}-1} (1+\alpha u) u^{\frac{n-1}{2}} \\ & + \sum_{k=0}^{\frac{n-3}{2}} \left((\alpha u)^k [(1 - \alpha^{\frac{n}{2}-1-k}) (1 - (\alpha^{1/2} u)^{n-2k}) + u (1 - \alpha^{\frac{n}{2}-k}) (1 - (\alpha^{1/2} u)^{n-2-2k})] \right) > 0. \end{aligned}$$

The first term is negative:

$$(\alpha u)^{\frac{n-1}{2}} (1+u) - \alpha^{\frac{n}{2}-1} (1+\alpha u) u^{\frac{n-1}{2}} = -\alpha^{\frac{n}{2}-1} u^{\frac{n-1}{2}} (1 - \alpha^{1/2}) (1 - \alpha^{1/2} u),$$

but it is compensated by the first term in the sum:

$$\begin{aligned} & -\alpha^{\frac{n}{2}-1}u^{\frac{n-1}{2}}(1-\alpha^{1/2})(1-\alpha^{1/2}u) + (1-\alpha^{\frac{n}{2}-1})(1-\alpha^{\frac{n}{2}}u^n) \\ & = (1-\alpha^{1/2})(1-\alpha^{1/2}u) \left[\left(\sum_{j=0}^{n-3} \alpha^{j/2} \right) \left(\sum_{j=0}^{n-1} (\alpha^{1/2}u)^j \right) - \alpha^{\frac{n}{2}-1}u^{\frac{n-1}{2}} \right]; \end{aligned}$$

as $\alpha^{\frac{n}{2}-1}u^{\frac{n-1}{2}} = \alpha^{(\frac{n-3}{2})/2}(\alpha^{1/2}u)^{\frac{n-1}{2}}$, the expression between brackets is positive, which ends the proof.

Remark 1 Let k be an integer ≥ 1 and let $l > k\frac{2\pi}{\sqrt{n-2}}$.

There exists a unique $x_{0,k}(l) \in]0, r[$ such that $\varphi(x_{0,k}(l)) = \frac{l}{2k}$, and there exists a unique $x_{0,1}(l/k) \in]0, r[$ such that $\varphi(x_{0,1}(l/k)) = \frac{l}{2k}$, so

$$x_{0,k}(l) = x_{0,1}(l/k).$$

It follows that

$$\sigma_k(l) = k^{\frac{2}{n}}\sigma_1(l/k). \quad (7)$$

By proposition 4 we have $k^{\frac{2}{n}}\sigma_1(l/k) < k^{\frac{2}{n}}\sigma_0(l/k) = \sigma_0(l)$, so

$$\sigma_k(l) < \sigma_0(l). \quad (8)$$

If $l \rightarrow k\frac{2\pi}{\sqrt{n-2}}$, then $x_{0,k}(l) = \varphi^{-1}(\frac{l}{2k}) \rightarrow \varphi^{-1}(\frac{\pi}{\sqrt{n-2}}) = r$, and

$$\sigma_k(l) \rightarrow \sigma_0(k\frac{2\pi}{\sqrt{n-2}}). \quad (9)$$

If $l \rightarrow +\infty$, then $x_{0,k}(l) \rightarrow 0$, and

$$\sigma_k(l) \rightarrow n(n-1)(2kV_N)^{\frac{2}{n}}[\psi(0)]^{\frac{2}{n}}. \quad (9)$$

In particular $\sigma_k(+\infty) = k^{\frac{2}{n}}\sigma_1(+\infty)$.

The following proposition shows that $\psi(0)$ is finite.

Proposition 5 Let $\psi_n(x_0) = (x_0^2 - x_0^n)^{\frac{n}{2}} \int_{x_0}^{x_m} \frac{x^{-n} dx}{\sqrt{P_{x_0}(x)}}$.

If n is even, $n = 2p$, we have

$$\lim_{x_0 \rightarrow 0} \psi_n(x_0) = \frac{2^{2p-2}[(p-1)!]^2}{(2p-1)!}.$$

If n is odd, $n = 2p + 1$, we have

$$\lim_{x_0 \rightarrow 0} \psi_n(x_0) = \frac{(2p)! \pi}{2^{2p+1}(p!)^2}.$$

Proof We write $\psi(x_0)$ in the form

$$\psi(x_0) = (1 - x_0^{n-2})^{\frac{n}{2}} x_0^n \left(\int_{x_0}^{\sqrt{x_0}} \frac{x^{-n} dx}{\sqrt{P_{x_0}(x)}} + \int_{\sqrt{x_0}}^r \frac{x^{-n} dx}{\sqrt{P_{x_0}(x)}} + \int_r^{x_m} \frac{x^{-n} dx}{\sqrt{P_{x_0}(x)}} \right).$$

For $\sqrt{x_0} \leq x \leq r$ we have $x^{-n} \leq x_0^{-n/2}$, and as the function $x \mapsto P_{x_0}(x)$ is increasing on $[0, r]$, $P_{x_0}(x) \geq P_{x_0}(\sqrt{x_0}) = x_0 - x_0^{n/2} - x_0^2 + x_0^n$, so

$$x_0^n \int_{\sqrt{x_0}}^r \frac{x^{-n} dx}{\sqrt{P_{x_0}(x)}} \leq x_0^{\frac{n-1}{2}} \frac{r - \sqrt{x_0}}{\sqrt{1 - x_0^{\frac{n}{2}-1} - x_0 + x_0^{n-1}}},$$

which shows that

$$\lim_{x_0 \rightarrow 0} x_0^n \int_{\sqrt{x_0}}^r \frac{x^{-n} dx}{\sqrt{P_{x_0}(x)}} = 0.$$

As $\int_r^{x_m} \frac{dx}{\sqrt{P_{x_0}(x)}}$ is bounded when $x_0 \rightarrow 0$, we have

$$\lim_{x_0 \rightarrow 0} x_0^n \int_r^{x_m} \frac{x^{-n} dx}{\sqrt{P_{x_0}(x)}} = 0.$$

We have $P_{x_0}(x) = (x^2 - x_0^2)(1 - \frac{x^n - x_0^n}{x^2 - x_0^2})$, and for $x_0 < x < \sqrt{x_0}$,

$$\frac{n}{2} x_0^{n-2} < \frac{x^n - x_0^n}{x^2 - x_0^2} < \frac{x_0^{n/2} - x_0^n}{x_0 - x_0^2},$$

so

$$\frac{1}{\sqrt{1 - \frac{n}{2} x_0^{n-2}}} < \frac{1}{\sqrt{1 - \frac{x^n - x_0^n}{x^2 - x_0^2}}} < \frac{1}{\sqrt{1 - \frac{x_0^{\frac{n}{2}-1} - x_0^{n-1}}{1 - x_0}}}.$$

It follows that

$$\lim_{x_0 \rightarrow 0} \psi(x_0) = \lim_{x_0 \rightarrow 0} x_0^n \int_{x_0}^{\sqrt{x_0}} \frac{x^{-n} dx}{\sqrt{x^2 - x_0^2}}.$$

Let $t = \sqrt{x^2 - x_0^2}$, then

$$\int_{x_0}^{\sqrt{x_0}} \frac{x^{-n} dx}{\sqrt{x^2 - x_0^2}} = \int_0^{\sqrt{x_0 - x_0^2}} \frac{dt}{(t^2 + x_0^2)^{\frac{n+1}{2}}},$$

and standard calculations show the announced results.

Remark 2 The function $n \mapsto \psi_n(0)$ is decreasing, so we have for all integer $n \geq 3$, $\psi_n(0) \leq \psi_n(3) = \pi/4$. Moreover $\lim_{n \rightarrow \infty} \psi_n(0) = 0$.

Proposition 6 *Let $n = 4$. Then*

$$\varphi(x_0) = \frac{\pi\sqrt{2}}{2} \sum_{p=0}^{+\infty} \frac{(4p)!}{2^{6p}(p!)^2(2p)!} (x_m^2 - x_0^2)^{2p},$$

$$\psi(x_0) = \frac{\pi\sqrt{2}}{8} \sum_{p=0}^{+\infty} \frac{(4p)!}{2^{6p}(p!)^2(2p)!} \frac{3}{(4p-3)(4p-1)} (x_m^2 - x_0^2)^{2p},$$

which shows in particular that the function ψ is decreasing.

Proof In this case we have $x_0^2 + x_m^2 = 1$ and $0 < x_0 < r = \frac{1}{\sqrt{2}} < x_m < 1$, so there exists $\theta \in]0, \frac{\pi}{4}[$ such that $x_0 = \sin \theta$ and $x_m = \cos \theta$. Let

$$x^2 = x_0^2 \cos^2 t + x_m^2 \sin^2 t;$$

then $2x dx = (x_m^2 - x_0^2) \sin 2t dt$, $P_{x_0}(x) = (x^2 - x_0^2)(x_m^2 - x^2) = \frac{1}{4}(x_m^2 - x_0^2)^2 \sin^2 2t$, $x^2 = \frac{1}{2}(1 - \cos 2\theta \cos 2t)$, so

$$\begin{aligned} \psi(x_0) &= (x_0^2 - x_m^2)^2 \int_{x_0}^{x_m} \frac{dx}{x^4 \sqrt{P_{x_0}(x)}} \\ &= \frac{\sqrt{2}}{4} \sin^4 2\theta \int_0^{\pi/2} \frac{dt}{(1 - \cos 2\theta \cos 2t)^{5/2}} \\ &= \frac{\sqrt{2}}{8} (1 - \cos^2 2\theta)^2 \int_0^\pi \frac{dt}{(1 - \cos 2\theta \cos t)^{5/2}}. \end{aligned}$$

Let $s = \cos 2\theta = x_m^2 - x_0^2$; then

$$\begin{aligned} \psi(x_0) &= \frac{\sqrt{2}}{8} (1 - s^2)^2 \int_0^\pi \left(\sum_{p=0}^{+\infty} (-1)^p \frac{(2p+4)!}{3 \cdot 2^{2p+2} (p+2)! p!} s^p \cos^p t \right) dt \\ &= \frac{\sqrt{2}}{8} (1 - s^2)^2 \sum_{p=0}^{+\infty} (-1)^p \frac{(2p+4)!}{3 \cdot 2^{2p+2} (p+2)! p!} s^p \int_0^\pi \cos^p t dt \\ &= \frac{\sqrt{2}}{8} (1 - s^2)^2 \sum_{p=0}^{+\infty} \frac{(4p+4)!}{3 \cdot 2^{4p+2} (2p+2)! (2p)!} s^{2p} \int_0^\pi \cos^{2p} t dt \\ &= \frac{\sqrt{2}}{8} (1 - s^2)^2 \sum_{p=0}^{+\infty} \frac{(4p+4)!}{3 \cdot 2^{4p+2} (2p+2)! (2p)!} s^{2p} \frac{(2p)!}{2^{2p} (p!)^2} \pi \\ &= \frac{\pi\sqrt{2}}{8} (1 - s^2)^2 \sum_{p=0}^{+\infty} \frac{(4p+4)!}{3 \cdot 2^{6p+2} (2p+2)! (p!)^2} s^{2p} \end{aligned}$$

$$= \frac{\pi\sqrt{2}}{8} \sum_{p=0}^{+\infty} \frac{(4p)!}{2^{6p}(p!)^2(2p)!} \frac{3}{(4p-3)(4p-1)} s^{2p},$$

and the formula for $\varphi(x_0)$ is obtained in the same way.