



About Brezis-Merle Problem in dimension 2.

Samy Skander Bahoura

► To cite this version:

| Samy Skander Bahoura. About Brezis-Merle Problem in dimension 2.. 2015. hal-00149613v4

HAL Id: hal-00149613

<https://hal.science/hal-00149613v4>

Preprint submitted on 18 Apr 2015 (v4), last revised 3 Aug 2020 (v9)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ABOUT BREZIS-MERLE PROBLEM WITH LIPSCHITZ CONDITION.

SAMY SKANDER BAHOURA

ABSTRACT. We give a blow-up analysis for Brezis-Merle problem with Dirichlet condition. As an application, we have another proof for Brezis-Merle Problem with Lipschitz condition.

Mathematics Subject Classification: 35J60 35B45 35B50

Keywords: blow-up, boundary, Dirichlet condition, a priori estimate, Lipschitz condition.

1. INTRODUCTION AND MAIN RESULTS

We set $\Delta = -(\partial_{11} + \partial_{22})$ on open set Ω of $\mathbb{R}^2$ with a smooth boundary.

We consider the following equation:

$$(P) \begin{cases} \Delta u = V e^u & \text{in } \Omega \subset \mathbb{R}^2, \\ u = 0 & \text{in } \partial\Omega. \end{cases}$$

Here, we assume that:

$$0 \leq V \leq b < +\infty, \quad e^u \in L^1(\Omega) \text{ and } u \in W_0^{1,1}(\Omega).$$

We can see in [6] a nice formulation to this problem (P) in the sens of the distributions. This Problem arises in geometrical and physical problems, see for example [1, 2, 16, 17]. The above equation was studied by many authors, with or without the boundary condition, also for Riemannian surfaces, see [1,18], where one can find some existence and compactness results. In [5] we have the following important Theorem,

Theorem A(Brezis-Merle [5]).*For $(u_i)_i$ and $(V_i)_i$ two sequences of functions relative to (P) with,*

$$0 < a \leq V_i \leq b < +\infty$$

then it holds,

$$\sup_K u_i \leq c,$$

with c depending on a, b, K and Ω .

One can find in [5] an interior estimate if we assume $a = 0$, but we need an assumption on the integral of e^{u_i} , namely, we have:

Theorem B(Brezis-Merle [5]).*For $(u_i)_i$ and $(V_i)_i$ two sequences of functions relative to the problem (P) with,*

$$0 \leq V_i \leq b < +\infty, \text{ and, } \int_{\Omega} e^{u_i} dy \leq C,$$

then it holds;

$$\sup_K u_i \leq c,$$

with c depending on b, C, K and Ω .

When $a = 0$, the boundedness of $\int_{\Omega} e^{u_i}$ is a necessary condition to work on the problem (P) as showed in [5] by the following counterexample.

Theorem C(Brezis-Merle [5]).*There are two sequences $(u_i)_i$ and $(V_i)_i$ of the problem (P) with,*

$$0 \leq V_i \leq b < +\infty, \text{ and, } \int_{\Omega} e^{u_i} dy \leq C,$$

and

$$\sup_{\Omega} u_i \rightarrow +\infty.$$

Note that for the problem (P), by using the Pohozaev identity, we can prove that $\int_{\Omega} e^{u_i}$ is uniformly bounded when $0 < a \leq V_i \leq b < +\infty$ and $\|\nabla V_i\|_{L^\infty} \leq A$ and Ω starshaped, when $a = 0$ and $\nabla \log V_i$ is uniformly bounded, we can bound uniformly $\int_{\Omega} V_i e^{u_i}$. In [14], Ma-Wei have proved that those results stay true for all open sets not necessarily starshaped.

In [8], Chen-Li have proved that if $a = 0$ and $\nabla \log V_i$ is uniformly bounded, then the functions are uniformly bounded near the boundary.

In [8], Chen-Li have proved that if $a = 0$ and $\int_{\Omega} e^{u_i}$ is uniformly bounded and $\nabla \log V_i$ is uniformly bounded, then we have the compactness result directly. Ma-Wei in [14], extend this result in the case where $a > 0$.

If we assume V more regular, we can have another type of estimates, a sup + inf type inequalities. It was proved by Shafrir see [15], that, if $(u_i)_i, (V_i)_i$ are two sequences of functions solutions of the previous equation without assumption on the boundary and, $0 < a \leq V_i \leq b < +\infty$, then we have the following interior estimate:

$$C\left(\frac{a}{b}\right) \sup_K u_i + \inf_{\Omega} u_i \leq c = c(a, b, K, \Omega).$$

One can see in [9] an explicit value of $C\left(\frac{a}{b}\right) = \sqrt{\frac{a}{b}}$. In his proof, Shafrir has used a blow-up function, the Stokes formula and an isoperimetric inequality, see [2]. For Chen-Lin, they have used the blow-up analysis combined with some geometric type inequality for the integral curvature.

Now, if we suppose $(V_i)_i$ uniformly Lipschitzian with A the Lipschitz constant, then, $C(a/b) = 1$ and $c = c(a, b, A, K, \Omega)$, see Brezis-Li-Shafrir [4]. This result was extended for Hölderian sequences $(V_i)_i$ by Chen-Lin, see [9]. Also, one can see in [12], an extension of the Brezis-Li-Shafrir result to compact Riemann surface without boundary. One can see in [13] explicit form, $(8\pi m, m \in \mathbb{N}^*$ exactly), for the numbers in front of the Dirac masses when the solutions blow-up. Here, the notion of isolated blow-up point is used. Also, in [18], we have refined estimates near the isolated blow-up points and the bubbling behavior of the blow-up sequences.

In [7], we have some a priori estimates on the 2 and 3-spheres $\mathbb{S}_2, \mathbb{S}_3$.

Here we give the behavior of the blow-up points on the boundary and a proof of Brezis-Merle Problem with Lipschitz condition.

The Brezis-Merle Problem (see [5]) is:

Problem. Suppose that $V_i \rightarrow V$ in $C^0(\bar{\Omega})$ with $0 \leq V_i \leq b$ for some positive constant b . Also, we consider a sequence of solutions (u_i) of (P) relatively to (V_i) such that,

$$\int_{\Omega} e^{u_i} dx \leq C,$$

is it possible to have:

$$\|u_i\|_{L^\infty} \leq C = C(b, C, V, \Omega)?$$

Here, we give a characterization of the behavior of the blow-up points on the boundary and also, in particular we extend Chen-Li theorem ($\nabla \log V_i$ uniformly bounded). Also, we give a new proof of Chen-Li result (when ∇V_i is uniformly bounded, the method of Chen-Li use some local estimates). For the behavior of the blow-up points on the boundary, the following condition is enough,

$$0 \leq V_i \leq b,$$

The condition $V_i \rightarrow V$ in $C^0(\bar{\Omega})$ is not necessary, but for the proof of the compactness for the Brezis-Merle problem we assume that:

$$\|\nabla V_i\|_{L^\infty} \leq A.$$

We have the following characterization of the behavior of the blow-up points on the boundary.

Theorem 1.1. *Assume that $\max_{\Omega} u_i \rightarrow +\infty$, Where (u_i) are solutions of the problem (P) with:*

$$0 \leq V_i \leq b, \text{ and } \int_{\Omega} e^{u_i} dx \leq C, \forall i,$$

then, after passing to a subsequence, there is a function u , there is a number $N \in \mathbb{N}$ and N points $x_1, \dots, x_N \in \partial\Omega$, such that,

$$\partial_{\nu} u_i \rightarrow \partial_{\nu} u + \sum_{j=1}^N \alpha_j \delta_{x_j}, \quad \alpha_j \geq 4\pi, \text{ weakly in the sense of measure } L^1(\partial\Omega).$$

$$u_i \rightarrow u \text{ in } C_{loc}^1(\bar{\Omega} - \{x_1, \dots, x_N\}).$$

In the following theorem, we have a proof for the global a priori estimate which concern the problem (P). The proof of Chen-Li and Ma-Wei [8,14], use the moving-plane method.

Theorem 1.2. *Assume that (u_i) are solutions of (P) relative to (V_i) with the following conditions:*

$$\|\nabla V_i\|_{L^{\infty}} \leq A \text{ and } \int_{\Omega} e^{u_i} \leq C,$$

We have,

$$\|u_i\|_{L^{\infty}} \leq c(b, A, C, \Omega),$$

2. PROOF OF THE THEOREMS

Proof of theorem 1.1:

We have,

$$\int_{\partial\Omega} \partial_{\nu} u_i d\sigma \leq C,$$

Without loss of generality, we can assume that $\partial_{\nu} u_i > 0$. Thus, (using the weak convergence in the space of Radon measures), we have the existence of a positive Radon measure μ such that,

$$\int_{\partial\Omega} \partial_{\nu} u_i \varphi d\sigma \rightarrow \mu(\varphi), \quad \forall \varphi \in C^0(\partial\Omega).$$

We take an $x_0 \in \partial\Omega$ such that, $\mu(x_0) < 4\pi$. Without loss of generality, we can assume that the following curve, $B(x_0, \epsilon) \cap \partial\Omega := I_\epsilon$ is an interval. (In this case, it is more simple to construct the following test function η_ϵ). We choose a function η_ϵ such that,

$$\begin{cases} \eta_\epsilon \equiv 1, & \text{on } I_\epsilon, \quad 0 < \epsilon < \delta/2, \\ \eta_\epsilon \equiv 0, & \text{outside } I_{2\epsilon}, \\ 0 \leq \eta_\epsilon \leq 1, \\ \|\nabla \eta_\epsilon\|_{L^\infty(I_{2\epsilon})} \leq \frac{C_0(\Omega, x_0)}{\epsilon}. \end{cases}$$

We take a $\tilde{\eta}_\epsilon$ such that,

$$\begin{cases} \Delta \tilde{\eta}_\epsilon = 0 & \text{in } \Omega \subset \mathbb{R}^2, \\ \tilde{\eta}_\epsilon = \eta_\epsilon & \text{in } \partial\Omega. \end{cases}$$

We use the following estimate, see [3, 6, 17],

$$\|\nabla u_i\|_{L^q} \leq C_q, \quad \forall i \text{ and } 1 < q < 2.$$

We deduce from the last estimate that, (u_i) converge weakly in $W_0^{1,q}(\Omega)$, almost everywhere to a function $u \geq 0$ and $\int_\Omega e^u < +\infty$ (by Fatou lemma). Also, V_i weakly converge to a nonnegative function V in L^∞ . The function u is in $W_0^{1,q}(\Omega)$ solution of :

$$\begin{cases} \Delta u = V e^u \in L^1(\Omega) & \text{in } \Omega \subset \mathbb{R}^2, \\ u = 0 & \text{in } \partial\Omega. \end{cases}$$

According to the corollary 1 of Brezis-Merle result, see [5], we have $e^{ku} \in L^1(\Omega)$, $k > 1$. By the elliptic estimates, we have $u \in C^1(\bar{\Omega})$.

We can write,

$$\Delta((u_i - u)\tilde{\eta}_\epsilon) = (V_i e^{u_i} - V e^u)\tilde{\eta}_\epsilon - 2 \langle \nabla(u_i - u), \nabla \tilde{\eta}_\epsilon \rangle. \quad (1)$$

We use the interior estimate of Brezis-Merle, see [5],

Step 1: Estimate of the integral of the first term of the right hand side of (1).

We use the Green formula between $\tilde{\eta}_\epsilon$ and u , we obtain,

$$\int_\Omega V e^u \tilde{\eta}_\epsilon dx = \int_{\partial\Omega} \partial_\nu u \eta_\epsilon \leq 4\epsilon \|\partial_\nu u\|_{L^\infty} = C\epsilon \quad (2)$$

We have,

$$\begin{cases} \Delta u_i = V_i e^{u_i} & \text{in } \Omega \subset \mathbb{R}^2, \\ u_i = 0 & \text{in } \partial\Omega. \end{cases}$$

We use the Green formula between u_i and $\tilde{\eta}_\epsilon$ to have:

$$\int_{\Omega} V_i e^{u_i} \tilde{\eta}_\epsilon dx = \int_{\partial\Omega} \partial_\nu u_i \tilde{\eta}_\epsilon d\sigma \rightarrow \mu(\eta_\epsilon) \leq \mu(I_{2\epsilon}) \leq 4\pi - \epsilon_0, \quad \epsilon_0 > 0 \quad (3)$$

From (2) and (3) we have for all $\epsilon > 0$ there is $i_0 = i_0(\epsilon)$ such that, for $i \geq i_0$,

$$\int_{\Omega} |(V_i e^{u_i} - V e^u) \tilde{\eta}_\epsilon| dx \leq 4\pi - \epsilon_0 + C\epsilon \quad (4)$$

Step 2: Estimate of integral of the second term of the right hand side of (1).

Let $\Sigma_\epsilon = \{x \in \Omega, d(x, \partial\Omega) = \epsilon^3\}$ and $\Omega_{\epsilon^3} = \{x \in \Omega, d(x, \partial\Omega) \geq \epsilon^3\}$, $\epsilon > 0$. Then, for ϵ small enough, Σ_ϵ is hypersurface.

The measure of $\Omega - \Omega_{\epsilon^3}$ is $k_2 \epsilon^3 \leq \mu_L(\Omega - \Omega_{\epsilon^3}) \leq k_1 \epsilon^3$.

Remark: for the unit ball $\bar{B}(0, 1)$, our new manifold is $\bar{B}(0, 1 - \epsilon^3)$.

We write,

$$\int_{\Omega} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx = \int_{\Omega_{\epsilon^3}} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx + \int_{\Omega - \Omega_{\epsilon^3}} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx. \quad (5)$$

Step 2.1: Estimate of $\int_{\Omega - \Omega_{\epsilon^3}} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx$.

First, we know from the elliptic estimates that $\|\nabla \tilde{\eta}_\epsilon\|_{L^\infty} \leq C_1/\epsilon^2$, C_1 depends on Ω

We know that $(|\nabla u_i|)_i$ is bounded in L^q , $1 < q < 2$, we can extract from this sequence a subsequence which converge weakly to $h \in L^q$. But, we know that we have locally the uniform convergence to $|\nabla u|$ (by Brezis-Merle theorem), then, $h = |\nabla u|$ a.e. Let q' be the conjugate of q .

We have, $\forall f \in L^{q'}(\Omega)$

$$\int_{\Omega} |\nabla u_i| f dx \rightarrow \int_{\Omega} |\nabla u| f dx$$

If we take $f = 1_{\Omega - \Omega_{\epsilon^3}}$, we have:

$$\text{for } \epsilon > 0 \exists i_1 = i_1(\epsilon) \in \mathbb{N}, \quad i \geq i_1, \quad \int_{\Omega - \Omega_{\epsilon^3}} |\nabla u_i| \leq \int_{\Omega - \Omega_{\epsilon^3}} |\nabla u| + \epsilon^3.$$

Then, for $i \geq i_1(\epsilon)$,

$$\int_{\Omega - \Omega_{\epsilon^3}} |\nabla u_i| \leq \text{mes}(\Omega - \Omega_{\epsilon^3}) \|\nabla u\|_{L^\infty} + \epsilon^3 = \epsilon^3 (k_1 \|\nabla u\|_{L^\infty} + 1).$$

Thus, we obtain,

$$\int_{\Omega - \Omega_{\epsilon^3}} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx \leq \epsilon C_1 (2k_1 \|\nabla u\|_{L^\infty} + 1) \quad (6)$$

The constant C_1 does not depend on ϵ but on Ω .

Step 2.2: Estimate of $\int_{\Omega_{\epsilon^3}} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx$.

We know that, $\Omega_\epsilon \subset \subset \Omega$, and (because of Brezis-Merle's interior estimates) $u_i \rightarrow u$ in $C^1(\Omega_{\epsilon^3})$. We have,

$$\|\nabla(u_i - u)\|_{L^\infty(\Omega_{\epsilon^3})} \leq \epsilon^3, \text{ for } i \geq i_3 = i_3(\epsilon).$$

We write,

$$\int_{\Omega_{\epsilon^3}} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx \leq \|\nabla(u_i - u)\|_{L^\infty(\Omega_{\epsilon^3})} \|\nabla \tilde{\eta}_\epsilon\|_{L^\infty} \leq C_1 \epsilon \text{ for } i \geq i_3,$$

For $\epsilon > 0$, we have for $i \in \mathbb{N}$, $i \geq \max\{i_1, i_2, i_3\}$,

$$\int_{\Omega} | \langle \nabla(u_i - u) | \nabla \tilde{\eta}_\epsilon \rangle | dx \leq \epsilon C_1 (2k_1 \|\nabla u\|_{L^\infty} + 2) \quad (7)$$

From (4) and (7), we have, for $\epsilon > 0$, there is $i_3 = i_3(\epsilon) \in \mathbb{N}$, $i_3 = \max\{i_0, i_1, i_2\}$ such that,

$$\int_{\Omega} |\Delta[(u_i - u)\tilde{\eta}_\epsilon]| dx \leq 4\pi - \epsilon_0 + \epsilon 2C_1 (2k_1 \|\nabla u\|_{L^\infty} + 2 + C) \quad (8)$$

We choose $\epsilon > 0$ small enough to have a good estimate of (1).

Indeed, we have:

$$\begin{cases} \Delta[(u_i - u)\tilde{\eta}_\epsilon] = g_{i,\epsilon} & \text{in } \Omega \subset \mathbb{R}^2, \\ (u_i - u)\tilde{\eta}_\epsilon = 0 & \text{in } \partial\Omega. \end{cases}$$

with $\|g_{i,\epsilon}\|_{L^1(\Omega)} \leq 4\pi - \epsilon_0$.

We can use Theorem 1 of [5] to conclude that there is $q > 1$ such that:

$$\int_{V_\epsilon(x_0)} e^{q(u_i-u)} dx \leq \int_{\Omega} e^{q(u_i-u)\tilde{\eta}_\epsilon} dx \leq C(\epsilon, \Omega).$$

where, $V_\epsilon(x_0)$ is a neighborhood of x_0 in $\bar{\Omega}$.

Thus, for each $x_0 \in \partial\Omega - \{\bar{x}_1, \dots, \bar{x}_m\}$ there is $\epsilon_{x_0} > 0, q_{x_0} > 1$ such that:

$$\int_{B(x_0, \epsilon_{x_0})} e^{q_{x_0} u_i} dx \leq C, \quad \forall i. \quad (9)$$

By the elliptic estimates, $(u_i \eta)_i$ is uniformly bounded in $W^{2,q_1}(\Omega)$ and also, in $C^1(\bar{\Omega})$. (Here, η is a cutuf function.)

Finally, we have, for some $\epsilon > 0$ small enough,

$$\|u_i\|_{C^{1,\theta}[B(x_0, \epsilon)]} \leq c_3 \quad \forall i.$$

We have proved that, there is a finite number of points $\bar{x}_1, \dots, \bar{x}_m$ such that the squence $(u_i)_i$ is locally uniformly bounded in $\bar{\Omega} - \{\bar{x}_1, \dots, \bar{x}_m\}$.

Proof of theorem 1.2:

The first Pohozaev identity applied around each blow-up point, see for example [14], gives :

$$\int_{\partial\Omega_{x_k}} [(\partial_\nu u_i) \nabla u_i - \frac{1}{2} |\nabla u_i|^2 \nu] dx = \int_{\Omega_{x_k}} \nabla V_i e^{u_i} - \int_{\partial\Omega_{x_k}} V_i e^{u_i} \nu, \quad (10)$$

We use the boundary condition on Ω and the boundedness of u_i and $\partial_j u_i$ outside the x_k , to have:

$$\int_{\partial\Omega} (\partial_\nu u_i)^2 dx \leq c_0(b, A, C, \Omega). \quad (11)$$

Thus we can use the weak convergence in $L^2(\partial\Omega)$ to have a subsequence $\partial_\nu u_i$, such that:

$$\int_{\partial\Omega} \partial_\nu u_i \varphi dx \rightarrow \int_{\partial\Omega} \partial_\nu u \varphi dx, \quad \forall \varphi \in L^2(\partial\Omega), \quad (12)$$

Thus, $\alpha_j = 0, j = 1, \dots, N$ and (u_i) is uniformly bounded.

ACKNOWLEDGEMENT.

The work is supported by YY. Li's Grant. The author would like to thank Pr. YY.Li for his support.

REFERENCES

- [1] T. Aubin. Some Nonlinear Problems in Riemannian Geometry. Springer-Verlag, 1998.
- [2] C. Bandle. Isoperimetric Inequalities and Applications. Pitman, 1980.
- [3] L. Boccardo, T. Gallouet. Nonlinear elliptic and parabolic equations involving measure data. J. Funct. Anal. 87 no 1, (1989), 149-169.
- [4] H. Brezis, YY. Li and I. Shafrir. A sup+inf inequality for some nonlinear elliptic equations involving exponential nonlinearities. J.Funct.Anal.115 (1993) 344-358.
- [5] H. Brezis, F. Merle. Uniform estimates and Blow-up behavior for solutions of $-\Delta u = V(x)e^u$ in two dimension. Commun. in Partial Differential Equations, 16 (8 and 9), 1223-1253(1991).
- [6] H. Brezis, W. A. Strauss. Semi-linear second-order elliptic equations in L^1 . J. Math. Soc. Japan 25 (1973), 565-590.
- [7] Chang, Sun-Yung A, Gursky, Matthew J, Yang, Paul C. Scalar curvature equation on 2- and 3-spheres. Calc. Var. Partial Differential Equations 1 (1993), no. 2, 205-229.
- [8] W. Chen, C. Li. A priori estimates for solutions to nonlinear elliptic equations. Arch. Rational. Mech. Anal. 122 (1993) 145-157.
- [9] C-C. Chen, C-S. Lin. A sharp sup+inf inequality for a nonlinear elliptic equation in $\mathbb{R}^2$. Commun. Anal. Geom. 6, No.1, 1-19 (1998).
- [10] D.G. De Figueiredo, P.L. Lions, R.D. Nussbaum, A priori Estimates and Existence of Positive Solutions of Semilinear Elliptic Equations, J. Math. Pures et Appl., vol 61, 1982, pp.41-63.
- [11] Ding, W, Jost, J, Li, J, Wang, G. The differential equation $\Delta u = 8\pi - 8\pi h e^u$ on a compact Riemann surface. Asian J. Math. 1 (1997), no. 2, 230-248.
- [12] YY. Li. Harnack Type Inequality: the method of moving planes. Commun. Math. Phys. 200,421-444 (1999).
- [13] YY. Li, I. Shafrir. Blow-up analysis for solutions of $-\Delta u = V e^u$ in dimension two. Indiana. Math. J. Vol 3, no 4. (1994). 1255-1270.
- [14] L. Ma, J-C. Wei. Convergence for a Liouville equation. Comment. Math. Helv. 76 (2001) 506-514.
- [15] I. Shafrir. A sup+inf inequality for the equation $-\Delta u = V e^u$. C. R. Acad.Sci. Paris Sér. I Math. 315 (1992), no. 2, 159-164.
- [16] Nagasaki, K, Suzuki, T. Asymptotic analysis for two-dimensional elliptic eigenvalue problems with exponentially dominated nonlinearities. Asymptotic Anal. 3 (1990), no. 2, 173-188.
- [17] Tarantello, G. Multiple condensate solutions for the Chern-Simons-Higgs theory. J. Math. Phys. 37 (1996), no. 8, 3769-3796.
- [18] L. Zhang. Blowup solutions of some nonlinear elliptic equations involving exponential nonlinearities. Comm. Math. Phys. 268 (2006), no. 1, 105-133.

DEPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ PIERRE ET MARIE CURIE, 2 PLACE JUSSIEU, 75005, PARIS, FRANCE.

E-mail address: samybahoura@yahoo.fr