



**HAL**  
open science

## **-trees and laminations for free groups III: Currents and dual -tree metrics**

Thierry Coulbois, Arnaud Hilion, Martin Lustig

► **To cite this version:**

Thierry Coulbois, Arnaud Hilion, Martin Lustig. -trees and laminations for free groups III: Currents and dual -tree metrics. Proceedings of the London Mathematical Society, 2008, 78, pp.755-766. hal-00151896

**HAL Id: hal-00151896**

**<https://hal.science/hal-00151896>**

Submitted on 5 Jun 2007

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ℝ-trees and laminations for free groups III: Currents and dual ℝ-tree metrics

Thierry Coulbois, Arnaud Hilion and Martin Lustig

June 5, 2007

## 1 Introduction

A geodesic lamination  $\mathcal{L}$  on a closed hyperbolic surface  $S$ , when provided with a transverse measure  $\mu$ , gives rise to a “dual ℝ-tree”  $T_\mu$ , together with an action of  $G = \pi_1 S$  on  $T_\mu$  by isometries. A point of  $T_\mu$  corresponds precisely to a leaf of the lift  $\tilde{\mathcal{L}}$  of  $\mathcal{L}$  to the universal covering  $\tilde{S}$  of  $S$ , or to a complementary component of  $\tilde{\mathcal{L}}$  in  $\tilde{S}$ . The  $G$ -action on  $T$  is induced by the  $G$ -action on  $\tilde{S}$  as deck transformations. This construction is well known (see [Mor86]). It is also known [Sko96] that conversely, for every small isometric action of a surface group  $G = \pi_1 S$  on a minimal ℝ-tree  $T$  there exists a “dual” measured lamination  $(\mathcal{L}, \mu)$  on  $S$ , i.e. one has  $T = T_\mu$  up to a  $G$ -equivariant isometry.

This beautiful correspondence has tempted geometers and group theorists to investigate possible generalizations, and the first one arises if one replaces the closed surface by a surface with boundary, and correspondingly the surface group  $G$  by a free group  $F_N$  of finite rank  $N \geq 2$ . A first glimpse of the potential problems can be obtained from two simultaneous but distinct identifications  $F_N \xrightarrow{\cong} \pi_1 S_1$  and  $F_N \xrightarrow{\cong} \pi_1 S_2$ , thus obtaining actions of  $\pi_1 S_1$  on a tree  $T_2$  which are dual to a measured lamination on  $S_2$ , but in general not dual to any measured lamination on the surface  $S_1$ .

Worse, using the index of an ℝ-tree action by  $F_N$  as introduced in [GL95], it is easily seen that for many (perhaps even “most”) small or very small ℝ-trees  $T$  with isometric  $F_N$ -action there is no identification whatsoever of  $F_N$  with the fundamental group of any surface that would make  $T$  dual to a lamination. An example of such trees are the forward limit trees  $T_\alpha$  of

certain irreducible automorphisms with irreducible powers (so called *iwip* automorphisms) of  $F_N$ . Much like pseudo-Anosov surface homeomorphisms, such an iwip automorphism has precisely one forward and one backward limit tree,  $T_\alpha$  and  $T_{\alpha^{-1}}$  respectively, and it induces a North-South dynamics on the space  $\overline{CV}_N$  of projectivized very small  $F_N$ -actions on  $\mathbb{R}$ -trees (see [LL03]). Note that, contrary to the case of pseudo-Anosov homeomorphisms, it is a frequent occurrence for an iwip automorphism  $\alpha$  (see Corollary 5.7 below) that its stretching factor  $\lambda_\alpha$  is different from the stretching factor  $\lambda_{\alpha^{-1}}$  of its inverse.

In [CHL-II] for any  $\mathbb{R}$ -tree  $T$  with isometric  $F_N$ -action, a *dual lamination*  $L(T)$  has been defined, which is the generalization of the geodesic lamination  $\mathfrak{L}$  for a surface tree  $T_\mu$  as discussed above. The goal of the present paper is to investigate the effect of putting an invariant measure  $\mu$  on the dual lamination  $L(T)$ , or, in the proper technical terms, considering a free group current  $\mu$  with support contained in  $L(T)$ . We prove here, if the  $F_N$ -action on  $T$  is very small and has dense orbits, that such a current defines indeed an induced measure on the metric completion  $\overline{T}$  of  $T$ .

In the special case considered above where  $T = T_\mu$  is dual to a measured lamination  $(\mathfrak{L}, \mu)$  on a surface, then the transverse measure  $\mu$  defines indeed a current on  $L(T_\mu)$ , and the induced measure on  $\overline{T}_\mu$  defines a dual distance on  $T_\mu$  which is precisely the same as the original distance on  $T_\mu$  (i.e. the one that comes from the transverse measure  $\mu$  on  $\mathfrak{L}$ ). For arbitrary very small trees  $T$  with dense  $F_N$ -orbits, the measure on  $\overline{T}$  induced by a current  $\mu$  on  $L(T)$  defines also a metric on  $T$ , except that this *dual metric*  $d_\mu$  may in general be in various ways degenerate (compare §5 below). In particular, the dual distance may well be infinite for any two distinct points of  $T$ . Alternatively, it could be zero throughout the interior  $T$  of  $\overline{T}$ .

The main result of this paper is to show that these “exotic” phenomena are not just theoretically possible, but that they actually do occur in important classes of examples.

Let  $\alpha \in \text{Aut}(F_N)$  be an iwip automorphism, let  $T_\alpha$  be the forward limit tree of  $\alpha$ . Then the dual lamination  $L(T_\alpha)$  is uniquely ergodic (see Proposition 5.6): it carries a projectively unique non-trivial current  $\mu$ . In this case the dual metric  $d_\mu$  is simply called the *dual distance*  $d_*$  on  $T_\alpha$  or on  $\overline{T}_\alpha$ .

**Theorem 1.1.** *Let  $\alpha \in \text{Aut}(F_N)$  be an iwip automorphism with  $\lambda_\alpha \neq \lambda_{\alpha^{-1}}$ . Then the dual distance  $d_*$  on the forward limit tree  $T_\alpha$  is zero or infinite throughout  $T_\alpha$ .*

Geodesic currents have been introduced by F. Bonahon for hyperbolic manifolds [Bon86, Bon88]. They turn out to be a powerful tool, and they also admit generalizations to a much larger setting, compare [Fur02]. For free groups and their automorphisms, the first serious application was given in the thesis of M. Bestvina's student R Martin [Mar95]. Recently, I. Kapovich rediscovered currents and studied them systematically, see [Kap04, Kap03]. As Kapovich's papers are very carefully written and very accessible to non-experts, we will review geodesic currents here only briefly and refer for all of the basic detail work to the papers of Kapovich.

The novelty in the setup presented here is the relationship between currents and laminations, which we establish systematically through studying, for any current  $\mu$ , the *support*  $\text{Supp}(\mu)$ . The latter belongs to the space  $\Lambda(F_N)$  of laminations for the free group  $F_N$ , which has been defined and investigated in detail in [CHL-I]. This gives a rather natural map from the space  $\text{Curr}(F_N)$  of currents to the space  $\Lambda(F_N)$ . The space  $\text{Curr}(F_N)$  of currents  $\mu$ , as well as the resulting compact space  $\mathbb{P}\text{Curr}(F_N)$  of projectivized currents  $[\mu]$ , admit a natural action of the group  $\text{Out}(F_N)$  of outer automorphisms of  $F_N$ . The results derived in Proposition 3.1 and in Lemmas 3.2, 3.3 and 3.4 can be summarized as follows:

**Theorem 1.2.** *The map  $\text{Supp} : \text{Curr}(F_N) \rightarrow \Lambda(F_N)$  induces a map*

$$\mathbb{P}\text{Supp} : \mathbb{P}\text{Curr}(F_N) \rightarrow \Lambda(F_N)$$

*which has the following properties:*

1.  $\mathbb{P}\text{Supp}$  is  $\text{Out}(F_N)$ -invariant.
2.  $\mathbb{P}\text{Supp}$  is not injective.
3.  $\mathbb{P}\text{Supp}$  is not surjective. However, every lamination  $L \in \Lambda(F_N)$  possesses a sublamination  $L_0 \subset L$  which belongs to the image of the map  $\mathbb{P}\text{Supp}$ .
4.  $\mathbb{P}\text{Supp}$  is not continuous. However, if  $(\mu_n)_{n \in \mathbb{N}}$  is a sequence of currents which converges to a current  $\mu$ , then the sequence of algebraic laminations  $\mathbb{P}\text{Supp}([\mu_n])$  has at least one accumulation point in  $\Lambda(F_N)$ , and any such accumulation point is a sublamination of  $\mathbb{P}\text{Supp}([\mu])$ .

Summing up, we believe that the results presented in this paper can be interpreted as follows:

On the one hand, the complete correspondence between small  $\mathbb{R}$ -trees and measured laminations, as known from the surface situation, does not fully extend to the world of free groups, very small  $\mathbb{R}$ -tree actions and currents. Unexpected degenerations seem to occur almost as a rule, and much further research is needed before one can speak of a “true understanding”.

On the other hand, the spaces of currents, of  $\mathbb{R}$ -tree actions, and of algebraic laminations for  $F_N$  are naturally related, and although this relationship is more challenging than in the surface case, there is clearly enough interesting structure there to justify further research efforts. A small such further contribution has already been given, in [CHL05], where algebraic laminations were used to characterize  $\mathbb{R}$ -trees up to  $F_N$ -equivariant variations of their metric.

**Acknowledgements.** *This paper originates from a workshop organized at the CIRM in April 05, and it has greatly benefited from the discussions started there and continued around the weekly Marseille seminar “Teichmüller” (partially supported by the FRUMAM).*

## 2 Currents on $F_N$

Let  $\mathcal{A}$  be a basis of the free group  $F_N$  or finite rank  $N \geq 2$ , and let  $F(\mathcal{A})$  denote the set of finite reduced words in  $\mathcal{A}^{\pm 1}$ , which is as usually identified with  $F_N$ . A *geodesic current* for a free group  $F_N$  can be defined in various ways. In particular, there are the following three equivalent ways to introduce them:

**I. Symbolic dynamist’s choice:** Consider the space  $\Sigma_{\mathcal{A}}$  of biinfinite reduced indexed words  $Z = \dots z_{i-1}z_i z_{i+1} \dots$  in  $\mathcal{A}^{\pm 1}$ , provided with the product topology, the shift operator  $\sigma : \Sigma_{\mathcal{A}} \rightarrow \Sigma_{\mathcal{A}}$ , and with the involution  $Z \mapsto Z^{-1}$ , see [CHL-I]. A geodesic current is a non-trivial  $\sigma$ -invariant finite Borel measure  $\mu$  on  $\Sigma_{\mathcal{A}}$ . We also require that  $\mu$  is *symmetric*: the measure is preserved by the involution of  $\Sigma_{\mathcal{A}}$  given by the inversion  $Z \mapsto Z^{-1}$ .

**II. Geometric group theorist’s choice:** Consider the space  $\partial^2 F_N$  of pairs  $(X, Y)$  of boundary points  $X \neq Y \in \partial F_N$ , endowed with the “product” topology, with the canonical diagonal action of  $F_N$ , and with the flip involution

$(X, Y) \mapsto (Y, X)$  as specified in [CHL-I]. A geodesic current is a non-trivial  $F_N$ - and flip-invariant Radon measure  $\mu$  on  $\partial^2 F_N$ , i.e. a Borel measure that is finite on any compact set.

**III. Combinatorist's choice:** A geodesic current is given by a non-zero function  $\mu : F_N = F(\mathcal{A}) \rightarrow \mathbb{R}_{\geq 0}$  with  $\mu(w^{-1}) = \mu(w)$  for all  $w \in F(\mathcal{A})$ , which satisfies the left and the right *Kolmogorov property*. For all reduced words  $w = y_1 \dots y_k \in F(\mathcal{A})$  one has

$$\mu(w) = \sum_{y \in \mathcal{A} \cup \mathcal{A}^{-1} \setminus \{y_k^{-1}\}} \mu(wy) = \sum_{y \in \mathcal{A} \cup \mathcal{A}^{-1} \setminus \{y_1^{-1}\}} \mu(yw).$$

This three viewpoints correspond to the three equivalent definitions given in [CHL-I] of a lamination for the free group  $F_N$ . We assume some familiarity of the reader with these three settings and will freely consider that a lamination is altogether symbolic (viewpoint I), algebraic (viewpoint II) and, a laminary language (viewpoint III). Whenever necessary, we emphasize the particular viewpoint used, by notationally specifying the lamination  $L$  in question as symbolic lamination  $L_{\mathcal{A}}$ , algebraic lamination  $L^2$ , or as laminary language  $\mathcal{L}$  respectively.

For currents, the transition between the three viewpoints is also canonical (see [Kap04]), and we will freely move from one to the other without always notifying the reader. To be specific, the Kolmogorov value  $\mu(w)$  of a reduced word  $w = y_1 \dots y_k \in F(\mathcal{A})$ , from the viewpoint III, is precisely the measure of the *cylinder*

$$C_{\mathcal{A}}(w) = \{\dots z_{i-1} z_i z_{i+1} \dots \mid z_1 = y_1, \dots, z_k = y_k\} \subset \Sigma_{\mathcal{A}}$$

from viewpoint I, and also, corresponding to viewpoint II, equal to the measure of the *algebraic cylinder*  $C_{\mathcal{A}}^2(w) \subset \partial^2 F_N$  given by

$$\{(X, wX') \mid X = x_1 x_2 \dots, X' = x'_1 x'_2 \dots \in \partial F(\mathcal{A}), x_1 \neq y_1, x'_1 \neq y_k^{-1}\}.$$

Note that the algebraic cylinder  $C_{\mathcal{A}}^2(w)$  is the image of the “symbolic” cylinder  $C_{\mathcal{A}}(w)$  under the map  $\Sigma_{\mathcal{A}} \rightarrow \partial^2 F_N$ ,  $Z = Z_- \cdot Z_+ \mapsto (Z_-^{-1}, Z_+)$ .

**Remark 2.1.** The reader should notice that in viewpoints I and III a basis  $\mathcal{A}$  of  $F_N$  is crucially used, while II is “algebraic”. It is very important to remember that basis change induces on the Kolmogorov function a more

complicated operation than just rewriting the given group element in the new basis  $\mathcal{B}$ . The correct transition is given, for any reduced word  $w \in F(\mathcal{B})$ , by decomposing the algebraic cylinder  $C_{\mathcal{B}}^2(w) \subset \partial^2 F_N$  into a finite disjoint union of translates  $u_i C_{\mathcal{A}}^2(v_i)$  of properly chosen algebraic cylinders  $C_{\mathcal{A}}^2(v_i)$ , with  $u_i, v_i \in F(\mathcal{A})$ , and posing:

$$\mu(w) = \sum_i \mu(v_i).$$

Similarly as for laminations (see §1 of [CHL-I]), every element  $w$  of  $F_N \setminus \{1\}$  (or rather, every non-trivial conjugacy class) defines an *integer* current  $\mu_w$ , given (in the language of viewpoint I) as follows: If  $w = u^m$  for the maximal exponent  $m \geq 1$ , then the measure  $\mu_w(C)$  of any measurable set  $C \subset \Sigma_{\mathcal{A}}$  is equal to  $m$  times the number of elements of  $C \cap L_{\mathcal{A}}(u)$ , where  $L_{\mathcal{A}}(u)$  is the finite set of biinfinite words of type  $\dots vv \cdot vv \dots$ , and  $v \in F(\mathcal{A})$  is any of the cyclically reduced words conjugated to  $u$  or to  $u^{-1}$ . Alternatively, (in the language of viewpoint II) the current  $\mu_w$  is given by an  $F_N$ -equivariant Dirac measure  $\mu_w$  on  $\partial^2 F_N$ , defined as follows: For every measurable set  $C^2 \subset \partial^2 F_N$  the value of  $\mu_w(C^2)$  is given by the number of cosets  $g \langle w \rangle \subset F_N$  which contain an element  $v$  that satisfies  $v(w^{-\infty}, w^{\infty}) \in C^2$  or  $v(w^{\infty}, w^{-\infty}) \in C^2$ . A third equivalent definition of  $\mu_w$  (corresponding to viewpoint III) is given by a count of “frequencies”, see [Kap03]. The noteworthy fact that  $\mu_w$  depends only on the element  $w \in F_N$  and not on the word  $w \in F(\mathcal{A})$  is obvious in the second of these definitions, but rather puzzling if one considers only the first or the third.

A current is *rational* if it is a non-negative linear combination of finitely many integer currents.

**Remark 2.2.** The above setup of the concept of currents in its various equivalent forms, together with the canonical identification  $F_N = F(\mathcal{A})$  for any basis  $\mathcal{A}$  of  $F_N$ , provides the ideal means to see very elegantly that many of the classical measure theoretic tools from symbolic dynamics do not depend on the underlying combinatorics of the chosen alphabet, but are rather algebraic in their true nature. Determining the exact point to which ergodic theory tools can be “algebraicized” seems to be a worthy task but goes beyond the scope of this paper.

### 3 The space $\text{Curr}(F_N)$

The set of currents on  $F_N$  will be denoted  $\text{Curr}(F_N)$ . It comes naturally with several interesting structures, which we will discuss briefly in this section. We would like to stress that this space, as well as its projectivization, appears to be a very interesting and useful tool for many fundamental questions about automorphisms of free groups, and we expect that it will play an important role in the future development of this subject.

First, the set  $\text{Curr}(F_N)$  of currents carries the weak topology, which for any basis  $\mathcal{A}$  of  $F_N$  is induced by the canonical embedding of  $\text{Curr}(F_N)$  into the vector space  $\mathbb{R}^{F(\mathcal{A})}$ , given by  $\mu \mapsto (\mu(w))_{w \in F(\mathcal{A})}$ . In particular, a family of currents  $\mu_i$  converges towards a current  $\mu \in \text{Curr}(F_N)$  if and only if  $\mu_i(w)$  converges to  $\mu(w)$  for every  $w \in F(\mathcal{A})$ .

Next, the same formalism as explained in Remark 2.1 for a basis change defines canonically an action by homeomorphisms of  $\text{Out}(F_N)$  on the space  $\text{Curr}(F_N)$ , which is formally given, for any  $\alpha \in \text{Aut}(F_N)$  and any  $\mu \in \text{Curr}(F_N)$ , by  $\alpha_*(\mu)(C) = \mu(\alpha^{-1}(C))$ , for every measurable set  $C \subset \partial^2 F_N$ . This convention defines a left action of  $\text{Out}(F_N)$ :

$$\begin{aligned} \alpha_*(\beta_*(\mu))(C) &= \beta_*(\mu)(\alpha^{-1}(C)) = \mu(\beta^{-1}(\alpha^{-1}(C))) \\ &= \mu((\alpha\beta)^{-1}(C)) = (\alpha\beta)_*(\mu)(C) \end{aligned}$$

For any integer current  $\mu_w$ , with  $w \in F_N \setminus \{1\}$ , this gives (compare [Kap03, Kap04]):

$$\alpha_*(\mu_w) = \mu_{\alpha(w)}$$

Every current  $\mu$  defines naturally a lamination  $L(\mu)$  for the free group  $F_N$ .  $L(\mu)$  can be viewed as an algebraic lamination  $L^2(\mu)$ , i.e. a non-empty subset of  $\partial^2 F_N$  which is closed and invariant under the  $F_N$ -action and the flip involution, compare [CHL-I]. In this setting,  $L^2(\mu) \subset \partial^2 F_N$  is simply the support  $\text{Supp}(\mu)$  of the Borel measure  $\mu$  on  $\partial^2 F_N$ , i.e. the complement of the biggest open set (= the union of all open sets) with measure 0. Alternatively,  $L(\mu)$  is given via its laminary language  $\mathcal{L}(\mu) = \{w \in F(\mathcal{A}) \mid \mu(w) > 0\}$ . We refer to a current  $\mu$  with support contained in a lamination  $L$  simply as *an invariant measure on  $L$* . Alternatively, one says that  $\mu$  is *carried by  $L$* .

A lamination  $L$  which has, up to scalar multiples, only one current  $\mu$  with support  $L(\mu) = L$ , is called *uniquely ergodic*. The simplest examples of non-uniquely ergodic laminations are given by the union  $L$  of two disjoint



laminations  $L_0$  and  $L_1$ , such that  $L_0$  and  $L_1$  are the support of currents  $\mu_0$  and  $\mu_1$  respectively (for example rational laminations  $L_0 = L(a)$  and  $L_1 = L(b)$  for distinct basis elements  $a, b \in \mathcal{A}$ ). For  $0 < \lambda < 1$  one obtains an interval of pairwise projectively distinct currents

$$\mu(\lambda) = \lambda\mu_1 + (1 - \lambda)\mu_0,$$

all with support  $L$ .

**Proposition 3.1.** *Recall that we assume  $N \geq 2$ , and let  $\Lambda(F_N)$  denote the space of laminations for  $F_N$  as introduced in [CHL-I]. The map*

$$\text{Supp} : \text{Curr}(F_N) \rightarrow \Lambda(F_N), \mu \mapsto L(\mu)$$

*is  $\text{Out}(F_N)$ -equivariant, but not continuous and not surjective.*

*Proof.* The  $\text{Out}(F_N)$ -equivariance is a direct consequence of the definition of the action of  $\text{Out}(F_N)$  on  $\text{Curr}(F_N)$  and on  $\Lambda(F_N)$ .

To see that the map  $\text{Supp}$  is non-surjective it suffices to consider the symbolic lamination  $L = L_{\{a,b\}}(Z)$  generated by the biinfinite word  $Z = \dots a a a b \cdot a a a \dots$ . It consists of the  $\sigma$ -orbit of  $Z$  and of the periodic word  $\dots a a \cdot a a \dots$ , as well as of their inverses. However, it is an easy exercise to show that any Kolmogorov function  $\mu$  on the associated laminary language  $\mathcal{L}_{\{a,b\}}(Z)$ , as it takes on values in  $\mathbb{R}_{\geq 0}$  and not in  $\mathbb{R}_{\geq 0} \cup \{\infty\}$ , must associate the value 0 to any word that contains the letter  $b$ , so that all the measure of  $\mu$  will be concentrated on the sublamination  $L_{\{a,b\}}(a)$  of  $L$ .

The fact that the map  $\text{Supp}$  is non-continuous can be seen from the above defined family  $\mu(\lambda)$  of currents with constant support  $L$ , by letting the parameter  $\lambda$  converge inside the open interval  $(0, 1)$  to the value 0 (or 1): For any such  $\lambda$  the support of  $\mu(\lambda)$  is clearly the union  $L_0 \cup L_1$ , while for the limit one gets  $L(\mu(0)) = L_0$  (or  $L(\mu(1)) = L_1$ ).  $\square$

The space  $\text{Curr}(F_N)$  has some additional structures which are not matched by corresponding structures in  $\Lambda(F_N)$ . For example, there is a canonical linear structure on  $\text{Curr}(F_N)$ , given simply by the embedding of  $\text{Curr}(F_N)$  into the real vector space  $\mathbb{R}^{F(\mathcal{A})}$ . Projectivization  $\mu \mapsto [\mu]$  defines the space of *projectivized currents*  $\mathbb{P}\text{Curr}(F_N)$ . Both  $\text{Curr}(F_N)$  and its projectivization are infinite dimensional, but  $\mathbb{P}\text{Curr}(F_N)$  is compact. Clearly, the map  $\text{Supp}$  splits over the projectivization, thus inducing a map  $\mathbb{P}\text{Supp} : \mathbb{P}\text{Curr}(F_N) \rightarrow \Lambda(F_N)$ , which by Proposition 3.1 is  $\text{Out}(F_N)$ -equivariant, non-continuous, and non-surjective. We obtain furthermore

**Lemma 3.2.** *The map  $\mathbb{P}\text{Supp} : \mathbb{P}\text{Curr}(F_N) \rightarrow \Lambda(F_N)$  is non-injective.*

*Proof.* Any non-uniquely ergodic lamination, in particular the above defined family  $\mu(\lambda)$  of currents with constant support  $L$ , shows that the map  $\mathbb{P}\text{Supp}$  is not injective.  $\square$

A second interesting example for the non-continuity of the map  $\text{Supp}$ , other than the one given in the proof of Proposition 3.1, is given by the rational currents  $\frac{1}{n}\mu_{ab^n}$  which converge to  $\mu_b$ , while their support  $L(ab^n)$  converge to the lamination generated by  $\dots bba \cdot bb \dots$  and  $\dots bb \cdot bb \dots$ , which is strictly larger than the lamination  $L(b)$ .

This last example, as also the one given in the proof of Proposition 3.1, indicates that a weaker statement than the continuity might be true for the map  $\text{Supp}$ . Since this will be needed in §5 as an important ingredient for the proof of Proposition 5.6, we formalize it here:

We say that a subset  $\delta$  of  $\Lambda(F_N)$  is *saturated* if  $\delta$  contains with any lamination also all of its sublaminations.

**Lemma 3.3.** *Let  $\delta \subset \Lambda(F_N)$  be a closed saturated subset of laminations. Then the full preimage  $\Delta \subset \mathbb{P}\text{Curr}(F_N)$  of  $\delta$  under the map  $\mathbb{P}\text{Supp}$  is closed.*

*Proof.* We consider a sequence of currents  $\mu_k$  in  $\text{Curr}(F_N)$ , with  $L(\mu_k) \in \delta$  for any  $\mu_k$ . By the compactness of  $\mathbb{P}\text{Curr}(F_N)$  and of  $\Lambda(F_N)$  we can assume, after possibly passing over to a subsequence, that there is a current  $\mu \in \text{Curr}(F_N)$  and a lamination  $L \in \Lambda(F_N)$  with  $[\mu] = \lim_{k \rightarrow \infty} [\mu_k]$  and  $L = \lim_{k \rightarrow \infty} L(\mu_k)$ . By properly normalizing the  $\mu_k$  we can actually assume that  $\mu = \lim_{k \rightarrow \infty} \mu_k$ .

We now fix a basis  $\mathcal{A}$  of  $F_N$  and consider the value of the Kolmogorov function  $\mu(w)$  for any  $w \in F_N \setminus \{1\}$ . If  $\mu(w) > 0$ , then by the topology on  $\text{Curr}(F_N)$ , for any  $\varepsilon$  with  $\mu(w) > \varepsilon > 0$  there is a bound  $k_0$  such that for any  $k \geq k_0$  one has  $|\mu_k(w) - \mu(w)| < \varepsilon$ . This shows for all  $k \geq k_0$  that  $w$  belongs to the laminary language  $\mathcal{L}(\mu_k)$ . But this implies that  $w$  belongs to the laminary language of  $L$ , which shows that  $\mu$  is carried by  $L$ . Since by hypothesis  $\delta$  is closed and saturated, this shows that  $[\mu]$  is contained in  $\Delta$ , so that the latter must be closed.  $\square$

A weaker statement than the surjectivity of the map  $\text{Supp}$  is crucially used in §5, again in the proof of Proposition 5.6:

**Lemma 3.4.** *Every lamination  $L \in \Lambda(F_N)$  contains a sublamination which is the support of some current  $\mu \in \text{Curr}(F_N)$ .*

*Proof.* For some basis  $\mathcal{A}$  of  $F_N$ , let  $Z = \dots z_{i-1}z_i z_{i+1} \dots$  be a leaf of the lamination  $L$ . Let  $Z_n = z_{-n} \dots z_n$  be the central subword of  $Z$  of length  $2n + 1$ .

For every  $n \in \mathbb{N}$  we define a ‘‘counting function’’  $m_n : F(\mathcal{A}) \rightarrow \mathbb{R}_{\geq 0}$ , by setting, for any word  $w$  in  $F(\mathcal{A})$ ,  $m_n(w)$  to be the number of occurrences of  $w$  as subword of  $Z_n$  or of  $Z_n^{-1}$ , divided by  $4n + 2$ . It follows directly that  $m_n$  satisfies the equations that defines the right and the left Kolmogorov property, up to possibly an error of absolute value less than  $\frac{1}{2n+1}$ . The total value of  $m_n$  on the set of words of length 1 is 1, for any  $n \in \mathbb{N}$ . Moreover  $m_n(w)$  is non-zero only for subwords of  $Z$ .

For each word  $w$  in  $F(\mathcal{A})$  we can chose a subsequence of  $(m_n)_{n \in \mathbb{N}}$  whose value at  $w$  converges. By a diagonal argument we get a subsequence that converges pointwise to a limit function  $\mu$  which satisfies the Kolmogorov laws while still having total value 1 on set of words of length 1, so that it is non-zero.

By construction, we have  $m_n(w) = m_n(w^{-1})$  for all  $w \in F(\mathcal{A})$ , so that the same is true for  $\mu$ . Hence  $\mu$  is a current. Its support is contained in the set of subwords of  $Z$  and thus, as a lamination, in  $L$ .  $\square$

A very interesting subspace  $\mathcal{M} \subset \mathbb{P}\text{Curr}(F_N)$  has been introduced by R. Martin in [Mar95] as closure of the  $\text{Out}(F_N)$ -orbit of  $[\mu_a]$ , for any element  $a$  of any basis  $\mathcal{A}$  of  $F_N$ . R. Martin shows that a projectivized integer current  $[\mu_w]$  belongs to  $\mathcal{M}$  if and only if  $w$  is contained in a proper free factor of  $F_N$ . In contrast to the analogous situation for  $\overline{\text{Out}(F_N)L(a)}$  (compare Proposition 8.1 of [CHL-I]), for  $N \geq 3$  it has been shown in [KL06], Theorem B, that  $\mathcal{M}$  is the unique minimal subspace of  $\text{Curr}(F_N)$  which is non-empty, closed and  $\text{Out}(F_N)$ -invariant.

The fact that currents behave somehow more friendly than laminations is underlined by the following fact, proved in R. Martin’s thesis and attributed there to M. Bestvina (compare to Proposition 6.5 of [CHL-I]):

**Proposition 3.5 ([Mar95]).** *The set of projectivized integer currents  $[\mu_w]$ , for any  $w \in F_N$ , is dense in  $\mathbb{P}\text{Curr}(F_N)$ .*

## 4 Geometric currents

A large class of very natural examples for a current  $\mu \in \text{Curr}(F_N)$  is given by any geodesic lamination  $\mathfrak{L} \subset S$ , provided with a transverse measure  $\mu'$ , where  $S$  is a hyperbolic surface with boundary as considered in the section 3 of [CHL-I] and section 6 of [CHL-II]. In this case the measure  $\mu$  on  $\partial^2 F_N$  can be nicely seen geometrically through the canonical identification of  $\partial F_N$  with the space  $\partial \tilde{S}$  of ends of the universal covering  $\tilde{S}$ , which is embedded as subset in the boundary at infinity  $S_\infty^1 = \partial \mathbb{H}^2$ . Two disjoint intervals  $A, B \subset S_\infty^1$ , with intersections  $A' = A \cap \partial \tilde{S}, B' = B \cap \partial \tilde{S}$ , define a measurable set  $A' \times B'$  of  $\partial^2 F_N$ , and the measure  $\mu(A' \times B')$  is precisely given by the measure  $\mu'(\beta)$  of an arc  $\beta$  in  $S$  which is transverse to  $\mathfrak{L}$ , and which lifts to an arc  $\tilde{\beta}$  in  $\tilde{S} \subset \mathbb{H}^2$  that has its two endpoints on the two extremal leaves of  $\tilde{\mathfrak{L}} \subset \tilde{S}$  which bound the set of all leaves of  $\tilde{\mathfrak{L}}$  that have one endpoint in  $A$  and one endpoint in  $B$ .

## 5 The dual metric for $\mathbb{R}$ -trees

In this section we assume familiarity of the reader with the notions of [CHL-II], from which we also import the notation without further explanations.

In the last section we have seen that every transverse measure  $\mu$  on a geodesic lamination  $\mathfrak{L}$  which is contained in a hyperbolic surface  $S$ , with non-empty boundary and with an identification  $\pi_1 S = F_N$ , gives rise to a canonical current in  $\text{Curr}(F_N)$  which we also denote by  $\mu$ . In section 6 of [CHL-II] we have discussed that  $(\mathfrak{L}, \mu)$  determines an  $\mathbb{R}$ -tree  $T_\mu$  with isometric  $F_N$ -action, and that the support of the current  $\mu$  and the dual lamination of  $T_\mu$  are the same: this lamination is precisely the lamination associated to  $\mathfrak{L} \subset S$ .

One of the most intriguing aspects of the relationship between currents and  $\mathbb{R}$ -trees comes from the attempt to extend this correspondence, which for surfaces is almost tautological, to more general  $\mathbb{R}$ -trees  $T$ . Indeed, the goal of this section is to understand better the true nature of the interaction between the metric on  $T$  and an invariant measure  $\mu$  carried by the dual lamination  $L(T)$  as defined in [CHL-II].

In the sequel we consider the dual lamination  $L(T)$  as algebraic lamination  $L^2(T)$ , i.e. a non-empty,  $F_N$ -invariant, flip-invariant and closed subset of

$\partial^2 F_N$ . From [CHL-II] we know that there is a map  $\mathcal{Q}^2 : L^2(T) \rightarrow \overline{T}$  which is  $F_N$ -equivariant and continuous (see Proposition 8.3 of [CHL-II]). Here  $T$  is an element of the boundary  $\partial cv_N$  of the unprojectivized Outer space  $cv_N$ : in particular,  $T$  is a non-trivial  $\mathbb{R}$ -tree with minimal, very small  $F_N$ -action by isometries (see [CHL-II], §2). We also require that the  $F_N$ -orbits of points are dense in  $T$  (“ $T$  has dense orbits”), and we denote by  $\overline{T}$  the metric completion of  $T$ .

**Corollary 5.1.** *For all  $T \in \partial cv_N$  with dense orbits, the map  $\mathcal{Q}^2 : L^2(T) \rightarrow \overline{T}$  is measurable (with respect to the two Borel  $\sigma$ -algebras on  $L^2(T)$  and on  $\overline{T}$ ).*

□

We apply the last corollary in order to define an *extended pseudo-metric*  $d_\mu$  on  $\overline{T}$ , for any current  $\mu$  which is carried by  $L(T)$ . An extended pseudo-metric is just like a metric, except that distinct points  $P, Q$  may have distance 0, positive distance, or distance  $\infty$ .

**Definition 5.2.** Let  $T \in \partial cv_N$  be with dense orbits, and assume that  $\mu \in \text{Curr}(F_N)$  satisfies  $\text{Supp}(\mu) \subset L(T)$ . One then defines, for any  $P, Q \in \overline{T}$ , their  $\mu$ -distance as follows:

$$d_\mu(P, Q) = \mu((\mathcal{Q}^2)^{-1}([P, Q])) \quad [= \mathcal{Q}_*^2(\mu)([P, Q])]$$

Clearly the function  $d_\mu$  is symmetric and, since  $\overline{T}$  is a tree, it satisfies the triangular inequality. For three points  $P, Q, R \in \overline{T}$  with  $Q \in [P, R]$  one has  $d_\mu(P, R) = d_\mu(P, Q) + d_\mu(Q, R)$  unless  $\mu((\mathcal{Q}^2)^{-1}(\{Q\})) > 0$ , which of course can happen (for example if  $Q$  has non-trivial stabilizer which carries all of the support of  $\mu$ ).

We distinguish now three special cases (note that we always assume that  $T$  is a minimal  $\mathbb{R}$ -tree, so that it agrees with its interior): The metric  $d_\mu$  is called *zero throughout  $T$*  if any two points in  $T$  have  $\mu$ -distance 0. It is called *infinite throughout  $T$*  if any two distinct points in  $T$  have  $\mu$ -distance  $\infty$ . It is called *positive throughout  $T$*  if any two distinct points in  $T$  have positive finite  $\mu$ -distance. Otherwise we call the  $\mu$ -distance *mixed*.

A particular case, which is of special importance, is the following:

**Definition 5.3.** An  $\mathbb{R}$ -tree  $T \in \partial cv_N$  is called *dually uniquely ergodic* if the dual lamination  $L(T)$  is uniquely ergodic.

We note that, in the case where  $T$  is dually uniquely ergodic, the  $\mu$ -distance is uniquely determined by  $T$ , up to rescaling. In this case we suppress the measure  $\mu$  and speak simply of the *dual distance*  $d_*$  on  $T$ .

**Conjecture 5.4.** *If  $T$  is dually uniquely ergodic then the dual distance is not mixed.*

We finish this article by proving that the case of dual distances which are infinite or zero throughout the interior does actually exist, and that it occurs in a natural context. We assume from now on a certain familiarity with some of the modern tools for the geometric theory of automorphisms of free groups. Background material and references can be found in [Vog02]. In particular we will use below the following facts and definitions:

**Remark 5.5.** (1) An automorphism  $\alpha$  of  $F_N$  is called *irreducible with irreducible powers (iwip)* if no non-trivial proper free factor of  $F_N$  is mapped by any positive power of  $\alpha$  to a conjugate of itself.

(2) It is known (compare [LL03]) that for every iwip automorphism  $\alpha$  there is, up to  $F_N$ -equivariant homothety, precisely one minimal *forward limit*  $\mathbb{R}$ -tree  $T_\alpha \in \partial cv_N$  which admits a homothety  $H : T_\alpha \rightarrow T_\alpha$  with stretching factor  $\lambda_\alpha > 1$  that twistedly commutes with  $\alpha$ . By this we mean that

$$\alpha(w)H = Hw : T_\alpha \rightarrow T_\alpha$$

holds for every  $w \in F_N$ . Note that both, the map  $H$  as well as the  $F_N$ -action on  $T$ , extend canonically to the metric completion  $\overline{T}_\alpha$ , so that the last statement holds also for  $\overline{T}_\alpha$  instead of  $T_\alpha$ .

(3) In terms of the induced action of  $Out(F_N)$  on the non-projectivized closed Outer space  $\overline{cv}_N$  (see [CHL-II], §9), the equation in (2) can be expressed by stating

$$T\alpha_* = \alpha_*^{-1}T = \lambda_\alpha T,$$

where  $\lambda_\alpha T$  denotes the tree  $T$  rescaled by the factor  $\lambda_\alpha$ .

(4) As a consequence of the equation in (2), the homothety  $H$  satisfies:

$$H\mathcal{Q}^2 = \mathcal{Q}^2\alpha : L^2(T_\alpha) \rightarrow \overline{T}_\alpha.$$

(5) There is no further fixed point of the  $\alpha_*$ -action on  $\overline{CV}_N$  other than the points  $[T_\alpha]$  and  $[T_{\alpha^{-1}}]$  specified above. In [LL03] it is shown that any iwip automorphism has North-South dynamics on  $\overline{CV}_N$ .

(6) One knows from [Mar95], Theorem 30 (again attributed to M. Bestvina) that, if  $\alpha$  is not *geometric*, i.e. induced by a surface homeomorphism  $h : S \xrightarrow{\cong} S$  via some identification  $F_N \cong \pi_1 S$ , then the  $\alpha_*$ -action on  $\mathbb{P}\text{Curr}(F_N)$  possesses precisely two fixed points, an attractive and a repelling one, and that  $\alpha_*$  has a North-South dynamics on  $\mathbb{P}\text{Curr}(F_N)$ .

(7) Let us denote by  $\mu_\alpha \in \text{Curr}(F_N)$  a representative of the attracting fixed point of the  $\alpha_*$ -action on  $\mathbb{P}\text{Curr}(F_N)$ . It satisfies  $\alpha_*(\mu_\alpha) = \lambda_\alpha \mu_\alpha$ , see [Mar95], where  $\lambda_\alpha$  is the stretching factor given in (2).

(8) Following [Mar95], the support of  $\mu_\alpha$  is contained in the so called *legal lamination*  $L_\alpha \in \Lambda(F_N)$ : Its leaves are represented, for any train track representative  $f : \tau \rightarrow \tau$  of  $\alpha$ , by biinfinite legal paths in  $\tau$ , and consequently by non-trivial (in fact: biinfinite) geodesics in  $T_\alpha$  (compare with the *attractive lamination* defined in [BFH97]). In particular, it follows from the alternative definition of the dual lamination,  $L(T) = L_{\mathcal{Q}}(T)$ , given in Theorem 1.1 of [CHL-II], that the two laminations  $L_\alpha$  and  $L(T_\alpha)$  are disjoint.

(9) Any iwip automorphism possesses a train track representative  $f : \tau \rightarrow \tau$  with transition matrix that is primitive. As a consequence, any edge  $e$  of  $\tau$  will have an iterate  $f^k(e)$  which crosses over all other edges. The canonical image in  $T_\alpha$  (under the map  $i : \tilde{\tau} \rightarrow T_\alpha$ , see [LL03]) of any lift of  $f^k(e)$  to the universal covering  $\tilde{\tau}$  is a segment which has the property that the union of its  $F_N$ -translates covers all of  $T_\alpha$ .

**Proposition 5.6.** *For every non-geometric iwip automorphisms  $\alpha \in \text{Aut}(F_N)$ , the forward limit tree  $T_\alpha$  is dually uniquely ergodic.*

*Proof.* From the  $\text{Out}(F_N)$ -equivariance of the map  $\lambda^2 : \partial cv_N \rightarrow \Lambda(F_N)$  in Proposition 9.1 of [CHL-II], together with Remark 5.5 (3) above, it follows that the dual lamination  $L(T_\alpha)$  is fixed by  $\alpha$ . Hence the set  $\Delta(\alpha) \subset \mathbb{P}\text{Curr}(F_N)$ , which consists of all preimages under the map  $\mathbb{P}\text{Supp}$  of the lamination  $L(T_\alpha)$  and any of its sublaminations, is invariant under the action of  $\alpha_*$  (by the equivariance of the maps  $\text{Supp}$  and  $\mathbb{P}\text{Supp}$ , see Proposition 3.1). As the set of all sublaminations of a given lamination is closed, see Proposition 6.4 of [CHL-I], it follows from Lemma 3.3 that  $\Delta(\alpha)$  is closed. Furthermore  $\Delta(\alpha)$  is non-empty, by Lemma 3.4. Thus  $\Delta(\alpha)$  is the non-empty union of closures of  $\alpha_*$ -orbits, so that it must contain the closure of at least one  $\alpha_*$ -orbit in  $\mathbb{P}\text{Curr}(F_N)$ . From the North-South dynamics of the  $\alpha_*$ -action on  $\mathbb{P}\text{Curr}(F_N)$  (Remark 5.5 (6)) it follows that either  $\Delta(\alpha)$  consists of precisely one of the two fixed points  $[\mu_\alpha]$  or  $[\mu_{\alpha^{-1}}]$ , or else it contains both of them.

But according to Remark 5.5 (8) the support of  $\mu_\alpha$  is contained in the legal lamination  $L_\alpha$ , which in turn is disjoint from  $L(T_\alpha)$ . Hence  $[\mu_\alpha]$  is not contained in  $\Delta(\alpha)$ , which proves that the latter consists precisely of the point  $[\mu_{\alpha^{-1}}]$ . This shows that  $L(T_\alpha)$  supports only one (projectivized) current, namely  $[\mu_{\alpha^{-1}}]$ .  $\square$

We can now give the proof of our main result as stated in §1:

*Proof of Theorem 1.1.* From Proposition 5.6 and its proof we know that the forward limit tree  $T_\alpha$  has dual lamination  $L(T_\alpha)$  which carries an (up to homothety) unique current, and that this current is equal to  $\mu_{\alpha^{-1}}$ .

We now calculate, for any  $P, Q \in T_\alpha$  (using Remark 5.5 (4) to get the third, and (7) to get the sixth of the equalities below):

$$d(H(P), H(Q)) = \lambda_\alpha d(P, Q)$$

and

$$\begin{aligned} d_*(H(P), H(Q)) &= \mu_{\alpha^{-1}}((\mathcal{Q}^2)^{-1}([H(P), H(Q)])) \\ &= \mu_{\alpha^{-1}}((H\mathcal{Q}^2\alpha^{-1})^{-1}([H(P), H(Q)])) \\ &= \mu_{\alpha^{-1}}(\alpha((\mathcal{Q}^2)^{-1}([P, Q]))) \\ &= \alpha_*^{-1}(\mu_{\alpha^{-1}}((\mathcal{Q}^2)^{-1}([P, Q]))) \\ &= \lambda_{\alpha^{-1}}\mu_{\alpha^{-1}}((\mathcal{Q}^2)^{-1}([P, Q])) \\ &= \lambda_{\alpha^{-1}}d_*(P, Q) \end{aligned}$$

Assume now that some points  $P \neq Q \in T_\alpha$  have finite dual distance. By iterating  $H$  one finds an interval  $[H^n(P), H^n(Q)]$  with the property that the union of its  $F_N$ -translates covers all of  $T_\alpha$  (compare Remark 5.5 (9)). This implies that any two points in  $T_\alpha$  have finite dual distance. If the dual distance function is furthermore non-zero, by the same argument it follows that any two points have non-zero distance. Thus the dual metric  $d_*$  on  $T_\alpha$  defines a non-trivial  $\mathbb{R}$ -tree  $T_\alpha^*$  with free  $F_N$ -action, and hence, since the equation in Remark 5.5 (2) carries over from  $T_\alpha$  to  $T_\alpha^*$ , the  $\mathbb{R}$ -tree  $T_\alpha^*$  defines a fixed point  $[T_\alpha^*]$  of the  $\alpha_*$ -action on  $\partial CV_N$  (see §2 and §9 of [CHL-II]). By Remark 5.5 (5) the point  $[T_\alpha^*]$  must agree with either  $[T_\alpha]$  or  $[T_{\alpha^{-1}}]$ . But this cannot be because we computed above that the stretching factor of the  $\alpha_*$ -action on  $T_\alpha^*$  is equal to  $\lambda_{\alpha^{-1}}$  and hence bigger than 1 (which rules out  $[T_\alpha^*] = [T_{\alpha^{-1}}]$ ), but different from  $\lambda_\alpha$  (thus ruling out  $[T_\alpha^*] = [T_\alpha]$ , by Remark 5.5 (3)).

Hence the dual metric  $d_*$  must be either zero or infinite throughout  $T_\alpha$ .

$\square$



A concrete example of an automorphism that satisfies the properties stated in Theorem 1.1 as hypotheses is given in [ABHS05] by the automorphism

$$\begin{aligned} a &\mapsto ab \\ b &\mapsto ac \\ c &\mapsto a \end{aligned}$$

of  $F_3$ , which has stretching factor  $1,84\dots$ , while its inverse

$$\begin{aligned} a &\mapsto c \\ b &\mapsto c^{-1}a \\ c &\mapsto c^{-1}b \end{aligned}$$

has stretching factor  $1,39\dots$

An iwip automorphism  $\alpha \in \text{Aut}(F_N)$  is called *parageometric*, if  $\alpha$  is not geometric, but  $T_\alpha$  is a geometric tree (see [GL95, GJLL98]). It has been proved recently in [HM04], see also [Gui04], that in this case the iwip automorphism  $\alpha^{-1}$  is not parageometric, and that its stretching factor  $\lambda_{\alpha^{-1}}$  is strictly smaller than  $\lambda_\alpha$  (compare [Gau05]). A family of such automorphisms, one for any  $N \geq 3$ , has been exhibited and investigated in [JL98]. We summarize:

**Corollary 5.7.** *The dual metric on the forward limit tree of any parageometric iwip automorphism of  $F_N$ , or of its inverse, is always infinite or zero throughout.*

## References

- [ABHS05] P. Arnoux, V. Berthé, A. Hilion, and A. Siegel. Fractal representation of the attractive lamination of an automorphism of the free group. *Ann. Inst. Fourier (Grenoble)* **56**, 2161–2212, 2006
- [BFH97] M. Bestvina, M. Feighn, and M. Handel. Laminations, trees, and irreducible automorphisms of free groups. *Geom. Funct. Anal.* **7**, 215–244, 1997.
- [Bon86] F. Bonahon. Bouts des variétés hyperboliques de dimension 3. *Ann. of Math.* **124**, 71–158, 1986.

- [Bon88] F. Bonahon. The geometry of Teichmüller space via geodesic currents. *Invent. Math.* **92**, 139–162, 1988.
- [CHL05] T. Coulbois, A. Hilion, and M. Lustig. Non-uniquely ergodic  $\mathbb{R}$ -trees are topologically determined by their algebraic lamination. Preprint, 2005. (To appear in *Illinois J. Math.*)
- [CHL-I] T. Coulbois, A. Hilion, and M. Lustig.  $\mathbb{R}$ -trees and laminations for free groups I: Algebraic laminations. ArXiv:math/0609416.
- [CHL-II] T. Coulbois, A. Hilion, and M. Lustig.  $\mathbb{R}$ -trees and laminations for free groups II: The lamination associated to an  $\mathbb{R}$ -tree. ArXiv:math/0702281.
- [Fur02] A. Furman. Coarse-geometric perspective on negatively curved manifolds and groups. In *Rigidity in dynamics and geometry (Cambridge, 2000)*, 149–166. Springer, Berlin, 2002.
- [GJLL98] D. Gaboriau, A. Jaeger, G. Levitt, and M. Lustig. An index for counting fixed points of automorphisms of free groups. *Duke Math. J.* **93**, 425–452, 1998.
- [GL95] D. Gaboriau and G. Levitt. The rank of actions on  $\mathbf{R}$ -trees. *Ann. Sci. École Norm. Sup.* **28**, 549–570, 1995.
- [Gau05] F. Gautero. Combinatorial mapping-torus, branched surfaces and free group automorphisms. Preprint 05.
- [Gui04] V. Guirardel. Core and intersection number for group actions on trees. ArXiv:math/0407206.
- [HM04] M. Handel and L. Mosher. The expansion factors of an outer automorphism and its inverse. ArXiv:math/0410015.
- [JL98] A. Jäger and M. Lustig. Free Group Automorphisms with Many Fixed Points at Infinity. Preprint 1998
- [Kap03] I. Kapovich. The frequency space of a free group. *Internat. J. Alg. Comput.* (special Gaeta Grigorchuk’s 50’s birthday issue), **15**, 939–969, 2005.

- [Kap04] I. Kapovich. Currents on free groups. “Topological and Asymptotic Aspects of Group Theory” (R. Grigorchuk, M. Mihalik, M. Sapir and Z. Sunik, Editors), *AMS Contemporary Mathematics Series*, **394**, 149–176, 2006.
- [KL06] I. Kapovich and M. Lustig, The actions of  $Out(F_n)$  on the boundary of Outer space and on the space of currents: minimal sets and equivariant incompatibility. ArXiv:math/0605548. (To appear in *Ergodic Theory and Dyn. Systems*.)
- [LL03] G. Levitt and M. Lustig. Irreducible automorphisms of  $F_n$  have north-south dynamics on compactified outer space. *J. Inst. Math. Jussieu* **2**, 59–72, 2003.
- [Mar95] R. Martin. *Non-Uniquely Ergodic Foliations of Thin Type, Measured Currents and Automorphisms of Free Groups*. Ph.D.-thesis, UCLA, 1995.
- [Mor86] J. Morgan. Group actions on trees and the compactification of the space of classes of  $SO(n,1)$ -representations. *Topology* **25**, 1–33, 1986.
- [Sko96] R. Skora. Splittings of surfaces. *J. Amer. Math. Soc.* **9**, 605–616, 1996.
- [Vog02] K. Vogtmann. Automorphisms of free groups and outer space. *Geom. Dedicata* **94**, 1–31, 2002.

Thierry Coulbois, Arnaud Hilion and Martin Lustig  
 Mathématiques (LATP)  
 Université Paul Cézanne – Aix-Marseille III  
 av. escadrille Normandie-Niémen  
 13397 Marseille 20  
 France  
 Thierry.Coulbois@univ-cezanne.fr  
 Arnaud.Hilion@univ-cezanne.fr  
 Martin.Lustig@univ-cezanne.fr