



HAL
open science

Finitely generated antisymmetric graph monoids

Friedrich Wehrung, Pere Ara, Francesc Perera

► **To cite this version:**

Friedrich Wehrung, Pere Ara, Francesc Perera. Finitely generated antisymmetric graph monoids. 2007. hal-00156906v1

HAL Id: hal-00156906

<https://hal.science/hal-00156906v1>

Preprint submitted on 25 Jun 2007 (v1), last revised 18 Jun 2008 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FINITELY GENERATED ANTISYMMETRIC GRAPH MONOIDS

PERE ARA, FRANCESC PERERA, AND FRIEDRICH WEHRUNG

ABSTRACT. A *graph monoid* is a commutative monoid for which there is a particularly simple presentation, given in terms of a quiver. Such monoids are known to satisfy various nonstable K-theoretical representability properties for either von Neumann regular rings or C*-algebras. We give a characterization of graph monoids within finitely generated antisymmetric refinement monoids. This characterization is formulated in terms of the *prime elements* of the monoid, and it says that each free prime has at most one free lower cover. We also characterize antisymmetric graph monoids of finite quivers. In particular, the monoid $\mathbb{Z}^\infty = \{0, 1, 2, \dots\} \cup \{\infty\}$ is a graph monoid, but it is not the graph monoid of any finite quiver.

1. INTRODUCTION

Many module-theoretical properties of a ring R can be expressed in terms of the so-called *nonstable K-theory of R* , which can be encoded in the commutative monoid $\mathcal{V}(R)$ defined, in the unital case, as the monoid of all isomorphism types of finitely generated projective right R -modules. Of particular interest is the case where the ring R is *von Neumann regular*, in which case the fundamental, still unsolved, open problem is the characterization problem of all monoids of the form $\mathcal{V}(R)$, published for the first time in [11]. While the original guess—namely, “all conical refinement monoids” (cf. Section 2 for the basic definitions)—got disproved in [16], with a counterexample of size \aleph_2 , the following fundamental question is still open:

Is every countable, conical refinement monoid representable, that is, isomorphic to $\mathcal{V}(R)$, for some von Neumann regular ring R ?

An important positive partial solution was recently obtained by Ara and Brustenga [2], where the authors prove that the representation problem above has a positive solution for the so-called *graph monoids*—in fact, the regular ring solving the problem can be taken an algebra over any given field, see [2, Theorem 4.4]. Graph monoids are a special class of refinement monoids for which there is a particularly simple presentation, given in terms of a row-finite quiver (see Section 3). The graph monoid of a row-finite quiver E is denoted by $M(E)$. We refer the reader to [2] for more information on the problem above and its relationship with the Separativity Problem of [3].

Date: June 20, 2007.

2000 *Mathematics Subject Classification.* Primary 20M14, 06F05. Secondary 16D70, 19A13, 19A49.

Key words and phrases. Monoid; commutative; quiver; graph monoid; separative; refinement; primely generated; primitive; antisymmetric; prime element; free element; regular element; von Neumann regular; C*-algebra.

The first and second authors were partially supported by the DGI and European Regional Development Fund, jointly, through Project MTM2005-00934, and by the Comissionat per Universitats i Recerca de la Generalitat de Catalunya.

For any row-finite quiver E , there is a C^* -algebra $C^*(E)$ associated to it, called the *Cuntz-Krieger graph C^* -algebra* of E . These graph C^* -algebras provide a wide generalization of the ubiquitous *Cuntz algebras* \mathcal{O}_n , introduced by Cuntz in [8]. We refer the reader to [15] for the basic theory of graph C^* -algebras. For any field K , the *Leavitt path K -algebra* of the row-finite quiver E , denoted by $L_K(E)$, has been defined in [1] and [4], as a purely algebraic analogue of the C^* -algebra $C^*(E)$. Indeed, it turns out that $L_{\mathbb{C}}(E)$ can be identified with a dense $*$ -subalgebra of $C^*(E)$. It was proven in [4, Theorem 3.5] that $\mathcal{V}(L_K(E)) \cong M(E)$ for every field K and every row-finite quiver E , and likewise the monoid $M(E)$ is isomorphic to $\mathcal{V}(C^*(E))$ by [4, Theorem 7.1]. The algebras $L_K(E)$ are not in general von Neumann regular, and the main goal of the paper [2] is to build an appropriate von Neumann regular algebra of fractions $Q_K(E)$ of $L_K(E)$ in such a way that the corresponding monoid is not altered: $\mathcal{V}(Q_K(E)) \cong M(E)$; see [2, Theorem 4.4].

Although graph monoids have a simple combinatorial characterization, it is *a priori* difficult to determine whether a given finitely generated monoid (given, say, by generators and relations) is a graph monoid. In this paper we solve that particular problem in the *antisymmetric* case, see Theorem 5.1. Our characterization is formulated in terms of the so-called *prime elements* of our monoid, and it says, for a given finitely generated antisymmetric refinement monoid (we say *primitive monoid*), that *each free prime has at most one free lower cover (among the primes)*. Our main preliminary result is the discovery of a primitive monoid that is not even a *retract* of any graph monoid, see Lemma 4.1. As another surprise, there are finitely generated graph monoids that are not the graph monoid of any *finite* quiver, the simplest of them being $\mathbb{Z}^\infty = \mathbb{Z}^+ \cup \{\infty\}$. A characterization of all antisymmetric graph monoids of finite quivers is given in Theorem 6.1.

2. BASIC CONCEPTS

For elements x and y in a commutative monoid M , we put

$$\begin{aligned} x \leq y &\Leftrightarrow (\exists z)(x + z = y), \\ x < y &\Leftrightarrow (x \leq y \text{ and } y \not\leq x), \\ x \equiv y &\Leftrightarrow (x \leq y \text{ and } y \leq x), \\ x \triangleleft y &\Leftrightarrow x + y = y, \\ x \ll y &\Leftrightarrow x + y \leq y. \end{aligned}$$

An element x of M is

- *free*, if $(n + 1)x \not\leq nx$ for any $n \in \mathbb{Z}^+$;
- *regular*, if $2x \leq x$;
- *idempotent*, if $2x = x$;
- *an atom*, if $x \not\leq 0$ and $x = y + z$ implies that either $y \leq 0$ or $z \leq 0$, for all $y, z \in M$.
- *prime*, if $p \not\leq 0$ and, further, $p \leq x + y$ implies that either $p \leq x$ or $p \leq y$, for all $x, y \in M$.

We denote by $\mathbb{P}(M)$ the set of all prime elements in M . We denote by $\mathbb{P}_{\text{free}}(M)$ (resp., $\mathbb{P}_{\text{reg}}(M)$) the set of all free primes (resp., regular primes) in M . We say that M is

- *conical*, if $x \leq 0$ implies that $x = 0$, for any $x \in M$;
- *antisymmetric*, if its algebraic preordering \leq is antisymmetric;

- *separative*, if $2x = x + y = 2y$ implies that $x = y$, for all $x, y \in M$;
- *strongly separative*, if $2x = x + y$ implies that $x = y$, for all $x, y \in M$;
- *primely generated*, if M is generated, as a monoid, by $\mathbb{P}(M)$. (*This is not equivalent to the definition given in [5], as primes may there be below zero, however, for conical monoids the two definitions are equivalent.*)

For a monoid N and a homomorphism $f: M \rightarrow N$, the *kernel* of f , defined as

$$\ker f = \{(x, y) \in M \times M \mid f(x) = f(y)\},$$

is a monoid congruence of M . A particular sort of congruence is obtained when we start with an *o-ideal* of M , that is, a nonempty subset I of M such that $x + y \in I$ iff $x \in I$ and $y \in I$, for all $x, y \in M$. Namely, the equivalence relation \equiv_I defined on M by the rule

$$x \equiv_I y \iff (\exists u, v \in I)(x + u = y + v), \quad \text{for all } x, y \in M$$

is a monoid congruence of M . We put $M/I = M/\equiv_I$ and we denote by $x_{/I}$ the \equiv_I -equivalence class of any element x of M . Observe that $x_{/I} \leq y_{/I}$ in M/I iff the relation $x \leq_I y$ defined as

$$x \leq_I y \iff (\exists h \in I)(x \leq y + h)$$

holds, for any $x, y \in M$. We shall say that M/I is an *ideal quotient* of M . We denote by

$$M \mid a = \{x \in M \mid (\exists n \in \mathbb{Z}^+)(x \leq na)\}$$

the *o-ideal* generated by an element $a \in M$.

The *antisymmetrisation* of M is the quotient M/\equiv . We put

$$\begin{aligned} L(M, p) &= \{q \in \mathbb{P}(M) \mid q < p \text{ and there is no } r \in \mathbb{P}(M) \text{ with } q < r < p\}, \\ L_{\text{free}}(M, p) &= \{q \in L(M, p) \mid q \text{ is free}\}, \\ L_{\text{reg}}(M, p) &= \{q \in L(M, p) \mid q \text{ is regular}\}, \end{aligned}$$

for any $p \in \mathbb{P}(M)$. We say that M is a *refinement monoid* [9, 17], if for all elements $a_0, a_1, b_0, b_1 \in M$ such that $a_0 + a_1 = b_0 + b_1$, there are elements $c_{i,j} \in M$, for $i, j < 2$, such that $a_i = c_{i,0} + c_{i,1}$ and $b_i = c_{0,i} + c_{1,i}$ for all $i < 2$. It is well-known that every *o-ideal* and every ideal quotient of a refinement monoid is a refinement monoid. It is established in [5, Corollary 6.8] that every finitely generated refinement monoid is primely generated. A monoid is *primitive* [14, Section 3.4], if it is an antisymmetric, primely generated, refinement monoid. For example, $\mathbb{Z}^\infty = \mathbb{Z}^+ \cup \{\infty\}$, endowed with its natural addition, is a primitive monoid. For any prime element p in a refinement monoid M , the map

$$\phi_p: M \rightarrow \mathbb{Z}^\infty, \quad x \mapsto \sup (n \in \mathbb{Z}^+ \mid np \leq x)$$

is a monoid homomorphism from M to \mathbb{Z}^∞ , see [5, Theorem 5.4]. Furthermore, if M is primitive, then the map

$$\phi: M \rightarrow (\mathbb{Z}^\infty)^{\mathbb{P}(M)}, \quad x \mapsto (\phi_p(x) \mid p \in \mathbb{P}(M)) \tag{2.1}$$

is a monoid embedding as well as an order-embedding, see [5, Theorem 5.11] or [17, Corollary 6.14].

We shall need the following lemma.

Lemma 2.1. *Let a, b, c be elements in a refinement monoid M , with c primely generated. If $a + c = b + c$, then there are $x, y \triangleleft c$ such that $a + x = b + y$.*

Proof. By [5, Theorem 4.1], there are $d, a', b', c' \in M$ such that $a = d + a'$, $b = d + b'$, $c = a' + c' = b' + c'$, and $c \leq c'$. Let $h \in M$ such that $c' = c + h$. The elements $x = b' + h$ and $y = a' + h$ are as required. \square

A partially ordered set P is *lower finite*, if the subset $P \downarrow p = \{q \in P \mid q \leq p\}$ is finite, for any $p \in P$. We say that P is a *forest*, if $P \downarrow p$ is a chain, for any $p \in P$.

3. GRAPH MONOIDS

We first recall some definitions from [4]. A *quiver* (in some other references, a *graph*) consists of a ‘vertex set’ E^0 , an ‘edge set’ E^1 , together with maps r and s from E^1 to E^0 describing, respectively, the range and source of edges; so we write $e: s(e) \rightarrow r(e)$, for any $e \in E^1$. We say that $u \in E^0$ *emits edges*, if $s^{-1}\{u\}$ is nonempty. We say that E is *row-finite*, if any $u \in E^0$ emits only finitely many edges, that is, $s^{-1}\{u\}$ is finite. We say that E is *finite*, if both E^0 and E^1 are finite.

The *graph monoid* of a row-finite quiver E , denoted by $M(E)$, is the commutative monoid defined by generators \bar{u} , for $u \in E^0$, and relations

$$\bar{u} = \sum \left(\overline{r(e)} \mid e \in s^{-1}\{u\} \right), \quad \text{for any } u \in E^0. \quad (3.1)$$

Conversely, with any set (alphabet) Σ , any doubly indexed family $(k_{u,v} \mid (u,v) \in \Sigma \times \Sigma)$ of natural numbers such that $\{v \in \Sigma \mid k_{u,v} \neq 0\}$ is finite for any $u \in \Sigma$, and any set of (formal) relations of the form

$$u = \sum (k_{u,v} \cdot v \mid v \in \Sigma) \quad (\text{for all } u \in \Sigma), \quad (3.2)$$

one can associate a row-finite quiver E such that (3.2) is a system of defining relations for $M(E)$: just take $E^0 = \Sigma$ and put $k_{u,v}$ edges with source u and range v , for any $u, v \in \Sigma$. We will say that E is the *quiver associated with the equation system* (3.2).

We shall denote by $\text{Fr}(X)$ the free commutative monoid on X , for any set X ; we identify X with its canonical image in $\text{Fr}(X)$. For a row-finite quiver E and $\alpha, \beta \in \text{Fr}(E^0)$, let $\alpha \rightarrow_1 \beta$ hold, if there are $\gamma \in \text{Fr}(E^0)$ and $x \in E^0$ such that

$$\alpha = \gamma + x \quad \text{and} \quad \beta = \gamma + \sum (r(e) \mid e \in s^{-1}\{x\}).$$

Furthermore, we put $\rightarrow_n = (\rightarrow_1) \circ \dots \circ (\rightarrow_1)$ (n times), for all $n \in \mathbb{Z}^+$, and we denote by \rightarrow the union of all the \rightarrow_n , for $n \in \mathbb{Z}^+$. We denote by $\pi_E: \text{Fr}(E^0) \rightarrow M(E)$ the unique monoid homomorphism such that $\pi_E(x) = \bar{x}$ for all $x \in E^0$. Of course, π_E is surjective, and solving the word problem for $M(E)$ amounts to finding a convenient description of the kernel \sim_E of π_E , defined by

$$\alpha \sim_E \beta \iff \pi_E(\alpha) = \pi_E(\beta), \quad \text{for all } \alpha, \beta \in \text{Fr}(E^0).$$

Such a description is Item (3) of the following lemma, established in [4, Section 4].

Lemma 3.1. *Let E be a row-finite quiver.*

- (1) *The relation \rightarrow is right refining, that is, for all $\alpha_0, \alpha_1, \beta \in \text{Fr}(E^0)$, if $\alpha_0 + \alpha_1 \rightarrow \beta$, then there are $\beta_0, \beta_1 \in \text{Fr}(E^0)$ such that $\alpha_0 \rightarrow \beta_0$, $\alpha_1 \rightarrow \beta_1$, and $\beta = \beta_0 + \beta_1$.*
- (2) *The relation \rightarrow is confluent, that is, for all $\alpha, \beta_0, \beta_1 \in \text{Fr}(E^0)$, if $\alpha \rightarrow \beta_0$ and $\alpha \rightarrow \beta_1$, then there exists $\gamma \in \text{Fr}(E^0)$ such that $\beta_0 \rightarrow \gamma$ and $\beta_1 \rightarrow \gamma$.*
- (3) *For all $\alpha, \beta \in \text{Fr}(E^0)$, $\pi_E(\alpha) = \pi_E(\beta)$ iff there exists $\gamma \in \text{Fr}(E^0)$ such that $\alpha \rightarrow \gamma$ and $\beta \rightarrow \gamma$.*

The following result is established in [4, Proposition 4.4].

Proposition 3.2. *The monoid $M(E)$ is a conical refinement monoid, for any row-finite quiver E .*

In case E is *finite*, $M(E)$ is finitely generated. As finitely generated refinement monoids are primely generated [5, Corollary 6.8], it follows that $M(E)$ is primely generated.

The following few definitions about quivers can be found in [4]. For $u, v \in E^0$, let $u \geq_1 v$ hold, if $v \in r(s^{-1}\{u\})$; denote by \geq the reflexive, transitive closure of \geq_1 . A subset H of E^0 is *hereditary*, if $u \in H$ and $u \geq v$ implies that $v \in H$, for all $u, v \in E^0$. Then we put $H^1 = s^{-1}(H)$ and $E \upharpoonright H = (H, H^1)$, the *restriction of E to H* . As H is hereditary, H^1 is contained in $r^{-1}(H)$. Then we define a quiver, denoted by $E \setminus H$, by $(E \setminus H)^0 = E^0 \setminus H$ and $(E \setminus H)^1 = \{e \in E^1 \mid r(e) \notin H\}$. A subset H of E^0 is *saturated*, if $s^{-1}\{v\} \neq \emptyset$ and $r(s^{-1}\{v\}) \subseteq H$ implies that $v \in H$, for each $v \in E^0$.

A *subquiver* of a quiver F is a pair $E = (E^0, E^1)$ with $E^0 \subseteq F^0$, $E^1 \subseteq F^1$, and $s_F(E^1) \cup r_F(E^1) \subseteq E^0$. Of course, then we denote by s_E and r_E the restrictions of s_F and r_F from E^1 to E^0 , respectively. We say that E is a *complete subquiver* of F , if $s_F^{-1}\{v\} \cap E^1 \neq \emptyset$ implies that $s_F^{-1}\{v\} \subseteq E^1$, for all $v \in E^0$.

A *quiver homomorphism* from a quiver E to a quiver F consists of a pair $f = (f^0, f^1)$ of maps $f^0: E^0 \rightarrow F^0$ and $f^1: E^1 \rightarrow F^1$ such that $r_F \circ f^1 = f^0 \circ r_E$ and $s_F \circ f^1 = f^0 \circ s_E$. We say that f is *complete*, if both f^0 and f^1 are injective and $(f^0(E^0), f^1(E^1))$ is a complete subquiver of F . If F is row-finite and $f: E \rightarrow F$ is a complete quiver embedding, then there exists a unique monoid homomorphism $M(f): M(E) \rightarrow M(F)$ such that $M(f)(\bar{v}) = \overline{f^0(v)}$ for all $v \in E^0$. The assignment $E \mapsto M(E)$, $f \mapsto M(f)$ is a functor.

An easy application of Lemma 3.1 yields the following.

Lemma 3.3. *Let E be a row-finite quiver. Then the following statements hold:*

- (i) *For every hereditary subset H of E^0 , the restriction $E \upharpoonright H$ is a complete subquiver of E , and the canonical homomorphism $M(E \upharpoonright H) \rightarrow M(E)$ is an embedding, whose image is an o -ideal of $M(E)$.*
- (ii) *Conversely, for every o -ideal J of $M(E)$, the set $H = \{u \in E^0 \mid \bar{u} \in J\}$ is a hereditary subset of E^0 , and $J \cong M(E \upharpoonright H)$.*

In the context of Lemma 3.3, we shall identify $M(E \upharpoonright H)$ with its canonical image in $M(E)$. Although the hereditary set H obtained in Lemma 3.3(ii) is saturated, saturation is not required in the proof of Lemma 3.3(i). Further, we observe the following result, established in [4, Lemma 3.1].

Lemma 3.4. *Let E be a row-finite quiver. Then every finite subquiver of E is a subquiver of some finite complete subquiver of E . Consequently, E is a direct limit of finite quivers with complete embeddings.*

As the functor $E \mapsto M(E)$, $f \mapsto M(f)$ preserves direct limits [4, Lemma 3.4], it follows that the graph monoid of any row-finite quiver is a direct limit of graph monoids of finite quivers. Denote by \mathcal{G} the category of all graph monoids of finite quivers with monoid homomorphisms, and by $\overline{\mathcal{G}}$ the category of all commutative monoids that are direct limits of members of \mathcal{G} with monoid homomorphisms. In particular, the graph monoid of any row-finite quiver is an object of $\overline{\mathcal{G}}$.

Observe now that \mathcal{G} is closed under finite direct products (take the disjoint union of the corresponding quivers). Hence, it follows from Corollary 4.2, Remark 4.3, and Lemma 4.4 in [12] that $\overline{\mathcal{G}}$ is closed under direct limits and retracts. The finitely generated monoids from $\overline{\mathcal{G}}$ can be characterized as follows.

Lemma 3.5. *A finitely generated commutative monoid M belongs to $\overline{\mathcal{G}}$ iff it is a retract of some member of \mathcal{G} .*

Proof. As $\overline{\mathcal{G}}$ is closed under retracts, it suffices to prove that if M belongs to $\overline{\mathcal{G}}$, then M is a retract of some member of \mathcal{G} . So let $M = \varinjlim_{i \in I} M_i$, with a directed partially ordered set I , monoids M_i in \mathcal{G} , transition morphisms $f_i^j: M_i \rightarrow M_j$ for $i \leq j$ in I , and limiting morphisms $f_i: M_i \rightarrow M$ for $i \in I$. As M is finitely generated, there exists $i \in I$ such that f_i is surjective. As $M \cong M_i / \ker f_i$ is finitely generated, it is, by Redei's Theorem, finitely presented, thus $\ker f_i$ is a finitely generated monoid congruence of M_i . As $\ker f_i = \bigcup_{j \geq i} \ker f_i^j$ (directed union), there exists $j \geq i$ such that $\ker f_i = \ker f_i^j$. For all $y \in M$, there exists $x \in M_i$ such that $y = f_i(x)$, and then $f_i^j(x)$ does not depend of the choice of $x \in f_i^{-1}\{y\}$; denote it by $e(y)$. Then e is a homomorphism from M to M_j , and $f_j \circ e = \text{id}_M$. Therefore, M is a retract of M_j . \square

We shall also need the following simple observation.

Lemma 3.6. *Both classes \mathcal{G} and $\overline{\mathcal{G}}$ are closed under o-ideals and ideal quotients.*

Proof. Closure of \mathcal{G} under o-ideals follows from Lemma 3.3, while closure of \mathcal{G} under ideal quotients follows from Theorem 5.3 and Lemma 6.6 in [4].

Now we deal with $\overline{\mathcal{G}}$. Any member M of $\overline{\mathcal{G}}$ can be written as a direct limit

$$(M, f_i \mid i \in I) = \varinjlim (M_i, f_i^j \mid i \leq j \text{ in } I),$$

for some directed partially ordered set I , monoids $M_i \in \mathcal{G}$, and monoid homomorphisms $f_i^j: M_i \rightarrow M_j$, $f_i: M_i \rightarrow M$. Let N be an o-ideal of M . The subset $N_i = f_i^{-1}(N)$ is an o-ideal of M_i , for all $i \in I$. We can define g_i^j (resp., g_i) as the restriction of f_i^j from N_i to N_j (resp., from N_i to N), and then it is straightforward to verify that

$$(N, g_i \mid i \in I) = \varinjlim (N_i, g_i^j \mid i \leq j \text{ in } I),$$

and so N belongs to $\overline{\mathcal{G}}$. Furthermore, for all $i \leq j$ in I , there exists a unique monoid homomorphism $h_i^j: M_i/N_i \rightarrow M_j/N_j$ (resp., $h_i: M_i/N_i \rightarrow M/N$) such that $h_i(x/N_i) = f_i(x)/N$ for any $x \in M_i$, and it is straightforward to verify that

$$(M/N, h_i \mid i \in I) = \varinjlim (M_i/N_i, h_i^j \mid i \leq j \text{ in } I),$$

and so M/N belongs to $\overline{\mathcal{G}}$. \square

4. A STRONGLY SEPARATIVE PRIMITIVE MONOID NOT IN $\overline{\mathcal{G}}$

In this section, we denote by M_0 the commutative monoid defined by generators p , a , b and relations $p = p + a = p + b$. So M_0 is a strongly separative, finitely generated, primitive monoid. It can be described as

$$M_0 = (\mathbb{Z}^+ \times \mathbb{Z}^+) \cup \{p, 2p, 3p, \dots\},$$

where $a = (1, 0)$, $b = (0, 1)$, and $p + x = p$ for all $x \in \mathbb{Z}^+ \times \mathbb{Z}^+$.

Lemma 4.1. *The monoid M_0 does not belong to $\overline{\mathcal{G}}$. That is, M_0 is not a direct limit of graph monoids.*

Proof. Suppose that M_0 belongs to $\overline{\mathcal{G}}$. By Lemma 3.5, M_0 is a submonoid of some monoid N in \mathcal{G} with a retraction $\rho: N \rightarrow M_0$. Let E be a finite quiver such that $N \cong M(E)$, with E^0 of minimal cardinality.

As M_0 is conical, $\rho^{-1}\{0\}$ is an o-ideal of N . By Theorem 5.3 and Lemma 6.6 in [4], the monoid $N' = N/\rho^{-1}\{0\}$ belongs to \mathcal{G} , via a subquiver E' of E , with $(E')^0 = E^0 \setminus H$, where $H = \{u \in E^0 \mid \rho(\bar{u}) = 0\}$. As M_0 is also a retract of N' and by the minimality assumption on E , we obtain that $H = \emptyset$, and so $\rho^{-1}\{0\} = \{0\}$. In particular, as both a and b are atoms of M_0 , they are also atoms of N .

Denote by I the o-ideal of M_0 generated by $\{a, b\}$. Hence $J = \rho^{-1}(I)$ is an o-ideal of N . We also denote by ρ the unique monoid homomorphism from $\text{Fr}(E^0)$ to M_0 that sends u to $\rho(\bar{u})$, for all $u \in E^0$.

Claim 1. *Every element of J is cancelable in N .*

Proof of Claim. Let $x + z = y + z$ hold, where $x, y \in N$ and $z \in J$. By Lemma 2.1, there are $u, v \triangleleft z$ such that $x + u = y + v$. As $\rho(u), \rho(v) \triangleleft \rho(z)$ and $\rho(z) \in I$, it follows that $\rho(u) = \rho(v) = 0$, thus $u = v = 0$, and thus $x = y$. \square Claim 1.

As N is a finitely generated graph monoid, it is primely generated. So there are $n \in \mathbb{N}$ and primes q_0, \dots, q_n in N such that $p = \sum_{i=0}^n q_i$. Applying ρ gives $p = \sum_{i=0}^n \rho(q_i)$, thus, up to permutation of the indices and putting $h = \sum_{i=1}^n q_i$, we get $\rho(h) \in I$ and $\rho(q_0) = p$. As $p + a = p$, we get $q_0 + a + h = q_0 + h$, hence, by Claim 1, $q_0 + a = q_0$. Similarly, $q_0 + b = q_0$. Therefore, by keeping the same ρ and by replacing the inclusion map from M_0 into N by the unique homomorphism fixing both a and b and sending p to q_0 , we reduce the problem to the case where $p = q_0$, that is, p is prime in N . Hence there exists $q \in E^0$ such that $\bar{q} \equiv p$. As both a and b are atoms of N , there are $x, y \in E^0$ such that $\bar{x} = a$ and $\bar{y} = b$.

Claim 2. *The inequality $\rho(\bar{u}) \leq p$ holds, for all $u \in E^0$.*

Proof of Claim. The set $H = \{u \in E^0 \mid \rho(\bar{u}) \leq p\}$ is a hereditary subset of E^0 , thus, by Lemma 3.3, the canonical map $j: M(E \upharpoonright H) \rightarrow M(E)$ is a monoid embedding. As p, a , and b are finite sums of elements of H , M_0 is a submonoid of $M(E \upharpoonright H)$, and so the restriction of ρ to $M(E \upharpoonright H)$ defines a retraction from $M(E \upharpoonright H)$ onto M_0 . By the minimality assumption on E , we obtain that $H = E^0$. \square Claim 2.

Now we put $P = \{u \in E^0 \mid \bar{u} = \bar{u} + a = \bar{u} + b\}$. As $\bar{q} \equiv p$ and $p = p + a = p + b$, we obtain that q belongs to P , thus P is nonempty. For $u \in P$, if $\rho(\bar{u}) < p$, then, by Claim 1, \bar{u} is cancelable, a contradiction as $\bar{u} = \bar{u} + a$; hence $\rho(\bar{u}) = p$.

Claim 3. *Every element $u \in P$ emits exactly one edge $e_{(u)}$ such that $r(e_{(u)}) \in P$. Every edge $e \in s^{-1}\{u\} \setminus \{e_{(u)}\}$ satisfies $\overline{r(e)} \in J$.*

Proof of Claim. As $\bar{u} = \bar{u} + a$ and by Lemma 3.1, there exists $\alpha \in \text{Fr}(E^0)$ such that $u \rightarrow \alpha$ and $u + x \rightarrow \alpha$. If u emits no edges, then $\alpha = u$, thus $u + x \rightarrow u$, a contradiction by Lemma 3.1(1); hence u emits edges. From $\bar{u} = \sum (\overline{r(e)} \mid e \in s^{-1}\{u\})$ it follows that $p = \sum (\rho(\overline{r(e)}) \mid e \in s^{-1}\{u\})$. Hence there exists exactly one $e_{(u)} \in s^{-1}\{u\}$ such that $\rho(\overline{r(e_{(u)})}) = p$, and $\rho(\overline{r(e)}) \in I$ for all other $e \in s^{-1}\{u\}$.

If X denotes the set of all those other edges, then $t = \sum (\overline{r(e)} \mid e \in X)$ is cancellable in N and $\bar{u} = \overline{r(e_{(u)})} + t$. As $\bar{u} = \bar{u} + a = \bar{u} + b$, we obtain that $\overline{r(e_{(u)})} = \overline{r(e_{(u)})} + a = \overline{r(e_{(u)})} + b$, so $r(e_{(u)}) \in P$. \square Claim 3.

Now we fix $q_0 \in P$, and we put $e_n = e_{(q_n)}$ and $q_{n+1} = r(e_{(q_n)})$, for every natural number n . So all elements q_n belong to P . As P is finite, there are natural numbers $k < m$ such that $q_k = q_m$. By taking the pair $(m, m - k)$ minimal with respect to the lexicographical ordering, we may assume without loss of generality that $k = 0$, so q_0, \dots, q_{m-1} are pairwise distinct, $q_{m+n} = q_n$, and $e_{m+n} = e_n$, for all $n \in \mathbb{Z}^+$. We put $E_n = s^{-1}\{q_n\} \setminus \{e_n\}$ and $c_n = \sum (\overline{\rho(r(e))} \mid e \in E_n)$, an element of I , for all $n \in \mathbb{Z}^+$. Furthermore, we put $c = \sum_{i < m} c_i$.

Claim 4. *Let $i \in \mathbb{Z}^+$ and $\alpha \in \text{Fr}(E^0)$ such that $q_i \rightarrow \alpha$. Then there are an integer $j \geq i$ and $\beta \in \text{Fr}(E^0)$ such that $\alpha = q_j + \beta$ with $\rho(\beta) = \sum_{i \leq k < j} c_k$.*

Proof of Claim. By induction on l such that $q_i \rightarrow_l \alpha$. For $l = 0$ it is trivial, so suppose the claim established at stage l , and let $q_i \rightarrow_{l+1} \alpha$. So there exists $\alpha' \in \text{Fr}(E^0)$ such that $q_i \rightarrow_l \alpha' \rightarrow_1 \alpha$. By the induction hypothesis, there are $j \geq i$ and $\beta' \in \text{Fr}(E^0)$ such that $\rho(\beta') = \sum_{i \leq k < j} c_k$ and $\alpha' = q_j + \beta'$. By the definition of \rightarrow_1 , either there exists $\gamma \in \text{Fr}(E^0)$ such that $\beta' \rightarrow_1 \gamma$ and then either

$$\alpha = q_j + \gamma$$

or

$$\alpha = q_{j+1} + \sum (r(e) \mid e \in E_j) + \beta'.$$

In the first case, $\rho(\gamma) = \rho(\beta') = \sum_{i \leq k < j} c_k$, so the result holds. In the second case, put $\beta = \sum (r(e) \mid e \in E_j) + \beta'$. Then

$$\rho(\beta) = c_j + \rho(\beta') = \sum_{i \leq k < j+1} c_k,$$

so $j + 1$ and β are as required. \square Claim 4.

Now we can conclude the proof. As $\bar{q}_0 = \bar{q}_0 + a$, there exists $\alpha \in \text{Fr}(E^0)$ such that $q_0 \rightarrow \alpha$ and $q_0 + x \rightarrow \alpha$. The second relation implies the existence of $\alpha', \gamma \in \text{Fr}(E^0)$ such that $q_0 \rightarrow \alpha'$, $x \rightarrow \gamma$, and $\alpha = \alpha' + \gamma$. By Claim 4, there are natural numbers i, j and elements $\beta, \beta' \in \text{Fr}(E^0)$ such that

$$\alpha = q_i + \beta, \quad \alpha' = q_j + \beta', \quad \rho(\beta) = \sum_{k < i} c_k, \quad \text{and} \quad \rho(\beta') = \sum_{k < j} c_k. \quad (4.1)$$

From $\alpha = \alpha' + \gamma$ it follows that $q_i + \beta = q_j + \beta' + \gamma$. As $\rho(\beta)$, $\rho(\beta')$, and $\rho(\gamma)$ belong to I , the elements β , β' , and γ have no component in P , thus $q_i = q_j$, so

$$i \equiv j \pmod{m} \quad \text{and} \quad \beta = \beta' + \gamma. \quad (4.2)$$

From $x \rightarrow \gamma$ it follows that $\rho(\gamma) = a$, thus, applying ρ to the equation in (4.2) and using (4.1), we obtain

$$\sum_{k < i} c_k = a + \sum_{k < j} c_k.$$

Hence, as $I \cong \mathbb{Z}^+ \times \mathbb{Z}^+$ is cancellative, $i > j$ and $\sum_{j \leq k < i} c_k = a$. Furthermore, as $i - j = \ell m$ for some $\ell > 0$ and the sequence $(c_l \mid l \in \mathbb{Z}^+)$ is periodical with period m ,

we get $a = \ell c$. A similar argument gives $b = \ell' c$, for some positive integer ℓ' , which forces $c = 0$, a contradiction as $a = \ell c$. \square

Observe that the monoid M_0 is the antisymmetrisation of the commutative monoid M'_0 defined by generators p, a, b and relation $p = p + a + b$. As $M'_0 = M(E)$ for the quiver E represented by Figure 1, this implies that *the antisymmetrisation of a finitely generated graph monoid is not necessarily a graph monoid*.




FIGURE 1. The quiver E corresponding to $p = p + a + b$.

By Lemmas 3.6 and 4.1, no commutative monoid M such that M_0 is an ideal quotient of an o-ideal of M can belong to $\overline{\mathcal{G}}$. In particular, we obtain the following result.

Theorem 4.2. *Let M be a primely generated refinement monoid with free prime elements p, a, b such that a and b are incomparable in $L(M, p)$. Then M does not belong to $\overline{\mathcal{G}}$. That is, M is not a direct limit of graph monoids.*

Proof. Observe that the assumptions imply that M_0 is isomorphic to the submonoid of M generated by $\{p, a, b\}$, so we may identify those two monoids. The o-ideal $N = M \setminus p$ obviously contains M_0 . As both a and b are prime elements in M , the subset

$$I = \{x \in N \mid a \not\leq x \text{ and } b \not\leq x\}$$

is an o-ideal of N . We shall prove that $M_0 \cong N/I$.

Claim. *For any $x \in N$, there exists $y \in M_0$ such that $x \equiv_I y$.*

Proof of Claim. The subset $N_1 = \{x \in N \mid (\exists y \in M_0)(x \equiv_I y)\}$ is a submonoid of N . We must prove that $x \in N_1$, for any $x \in N$. As M is primely generated, it suffices to consider the case where x is prime, and thus, by the definition of N , $x \leq p$. Obviously we can assume that either $a \leq x$ or $b \leq x$. Now assume, say, that $a \leq x$. If $x < p$, then, as $a \in L(M, p)$, we get $a \equiv x$, so, as a is free, $x = a + y$ for some $y < a$; by the previous case, $y \in N_1$, so $x \in N_1$. The remaining case is where $x \equiv p$. As p is free, there exists $y < p$ such that $x = p + y$. By the previous case, $y \in N_1$, and so $x \in N_1$. \square Claim.

It follows from the claim above that the monoid homomorphism $\varepsilon: M_0 \rightarrow N/I$, $x \mapsto x/I$ is surjective. To prove that it is one-to-one, it suffices to prove the following statements:

The element p/I is free in N/I . Suppose, to the contrary, that $(n+1)p/I \leq np/I$, for some $n \in \mathbb{N}$. This means that there exists $x \in I$ such that $(n+1)p \leq np + x$. By applying the homomorphism $\phi_p: N \rightarrow \mathbb{Z}^\infty$ and using the freeness of p , we obtain that $n+1 \leq n + \phi_p(x)$, thus $p \leq x$, a contradiction as $x \in I$.

By using ϕ_a and ϕ_b instead of ϕ_p , we obtain in a similar manner that *both elements a/I and b/I are free in N/I .*

The elements $a_{/I}$ and $b_{/I}$ are incomparable in N/I . Suppose, say, that $a_{/I} \leq b_{/I}$, that is, there exists $x \in I$ such that $a \leq b + x$. As a is prime and $a \not\leq b$, we get that $a \leq x$, a contradiction as $x \in I$.

Therefore, ε is an isomorphism, and so $M_0 \cong N/I$. By Lemma 4.1, M_0 does not belong to $\overline{\mathfrak{G}}$. Therefore, by Lemma 3.6, neither does M . \square

5. A CHARACTERIZATION OF GRAPH MONOIDS AMONG PRIMITIVE MONOIDS WITH LOWER FINITE SET OF PRIMES

The main goal of the present section is to characterize graph monoids within finitely generated primitive monoids. As every finitely generated primitive monoid has a finite set of primes (which is the smallest generating subset), the following result is slightly more general.

Theorem 5.1. *Let M be a primitive monoid such that $\mathbb{P}(M)$ is lower finite. Then the following statements are equivalent:*

- (i) M is a graph monoid.
- (ii) M is a direct limit of graph monoids.
- (iii) $|\mathbf{L}_{\text{free}}(M, p)| \leq 1$ for each $p \in \mathbb{P}_{\text{free}}(M)$.

Proof. (i) \Rightarrow (ii) is trivial, while (ii) \Rightarrow (iii) follows immediately from Theorem 4.2. It remains to prove the direction (iii) \Rightarrow (i).

So assume that $|\mathbf{L}_{\text{free}}(M, p)| \leq 1$ for each $p \in \mathbb{P}_{\text{free}}(M)$. We shall construct a row-finite quiver E with vertex set

$$E^0 = \mathbb{P}(M) \sqcup \{b_{i,j}^p \mid 0 \leq i < m_p, 0 \leq j, p \in \mathbb{P}_{\text{reg}}(M)\},$$

where $m_p = |\mathbf{L}_{\text{free}}(M, p)|$.

For $p \in \mathbb{P}(M)$ we define elements z_p and w_p in $\text{Fr}(E^0)$ by

$$\begin{aligned} z_p &= \sum (q \mid q \in \mathbb{P}_{\text{reg}}(M), q < p), \\ w_p &= \sum (q \mid q \in \mathbb{P}(M), q < p), \end{aligned}$$

and we consider the following relations:

Case 1: If p is free, put the relation

$$p = p + w_p. \tag{5.1}$$

Case 2: If p is regular and all the elements of $\mathbf{L}(M, p)$ are regular, put the relation

$$p = 2p + \sum (q \mid q \in \mathbf{L}(M, p)). \tag{5.2}$$

Case 3: If p is regular and p_0, \dots, p_{m-1} are the elements of $\mathbf{L}_{\text{free}}(M, p)$, with $m > 0$, put the relations

$$b_{0,0}^p = p, \tag{5.3}$$

$$b_{i,0}^p = 2b_{i,0}^p + b_{i,1}^p + b_{i,2}^p + p_i, \tag{5.4}$$

$$b_{i,1}^p = b_{i,0}^p + 2b_{i,1}^p + b_{i,2}^p. \tag{5.5}$$

Furthermore, define $\alpha: \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ as $\alpha(n) = 2nm + 2$. For $k \in \mathbb{Z}$ set $\epsilon(k) = 1$ if k is even and $\epsilon(k) = 2$ if k is odd, and put $\beta(k) = \lceil \frac{k-1}{2} \rceil$, where $\lceil x \rceil$ denotes the largest integer below x , for any real number x .

For $n \geq 0$ and $0 \leq k \leq 2m - 1$, put the relation

$$b_{i,\alpha(n)+k}^p = b_{i,\alpha(n)+k}^p + \epsilon(k)b_{i,nm+\beta(k)+1}^p + b_{[\beta(k),m-1] \times [\alpha(n)+1,\alpha(n+1)]}^p + z_p, \quad (5.6)$$

where we set

$$b_X^p = \sum_{(i,j) \in X} b_{i,j}^p, \quad \text{for any subset } X \subseteq [0, m-1] \times \mathbb{Z}^+.$$

For example, for $n = 0$, this yields the relations

$$b_{i,2}^p = b_{i,2}^p + b_{i,1}^p + b_{[0,m-1] \times [3,2m+2]}^p + z_p \quad (5.7)$$

$$b_{i,3}^p = b_{i,3}^p + 2b_{i,1}^p + b_{[0,m-1] \times [3,2m+2]}^p + z_p \quad (5.8)$$

...

$$b_{i,2m}^p = b_{i,2m}^p + b_{i,m}^p + b_{[m-1,m-1] \times [3,2m+2]}^p + z_p$$

$$b_{i,2m+1}^p = b_{i,2m+1}^p + 2b_{i,m}^p + b_{[m-1,m-1] \times [3,2m+2]}^p + z_p$$

Let E be the quiver associated with the relations (5.1)–(5.6) above (cf. Section 3), so that $M(E)$ is the commutative monoid defined by generators E^0 and defining relations (5.1)–(5.6). We shall prove that $M(E)$ and M are isomorphic. There is a surjective monoid homomorphism

$$\varphi: M(E) \rightarrow M$$

such that $\varphi(p) = p$ for all $p \in \mathbb{P}(M)$ and $\varphi(b_{i,j}^p) = p$ for all $p \in \mathbb{P}_{\text{reg}}(M)$. To prove that φ is well-defined we need to check that the relations (5.1)–(5.6) are satisfied by the images of E^0 under φ , which is obvious.

It remains to prove that φ is one-to-one. As M is a primitive monoid, the relations $p = p + q$, for $p, q \in \mathbb{P}(M)$ such that $p = p + q$ holds in M , are defining relations of M (see [14, Section 3.5]). As the subset $\mathbb{P}(M) \cup \{b_{i,j}^p \mid p \in \mathbb{P}_{\text{reg}}(M)\}$ generates $M(E)$, it suffices to prove that the following relations hold in $M(E)$:

$$p = p + q, \quad \text{for all } p, q \in \mathbb{P}(M) \text{ with } q < p, \quad (5.9)$$

$$p = b_{i,j}^p = 2p, \quad \text{for all } i < m, \text{ all } j \in \mathbb{Z}^+,$$

$$\text{and all } p \in \mathbb{P}_{\text{reg}}(M), \text{ with } m = |\text{L}_{\text{free}}(p, M)|. \quad (5.10)$$

Strictly speaking, we should write $\bar{p} = \bar{p} + \bar{q}$, and so on, but we shall drop the bars for clarity of notation, choosing instead to specify the monoid where the relations should be verified—in particular, $M(E)$ in the case of (5.9), (5.10). We first prove that $b_{i,j}^p \asymp b_{i',j'}^p$ in $M(E)$, for any pair of indices (i, j) and (i', j') . We argue by induction. Note that (5.4) gives $b_{i,1}^p \leq b_{i,0}^p$ and that (5.5) gives $b_{i,0}^p \leq b_{i,1}^p$, so that we get $b_{i,0}^p \asymp b_{i,1}^p$. Also $b_{i,2}^p \asymp b_{i,0}^p \asymp b_{i,1}^p$, and (5.7) implies $b_{i,1}^p \leq b_{i,2}^p$, hence

$$b_{i,2}^p \asymp b_{i,1}^p \asymp b_{i,0}^p, \quad \text{for all } i < m.$$

Now (5.8) gives $b_{0,3}^p \leq b_{i,3}^p \leq b_{0,3}^p$, whence $b_{i,3}^p \asymp b_{0,3}^p$, for all $i < m$. Using again (5.7), (5.8), we get $b_{i,3}^p \leq b_{i,2}^p$ and $b_{i,1}^p \leq b_{i,3}^p$, so we obtain $b_{i,3}^p \asymp b_{i,1}^p \asymp b_{i,2}^p$, for all $i < m$. It follows that $b_{0,1}^p \asymp b_{i,j}^p$, for all $i < m$ and for all $j < 4$.

Now assume that $b_{i,j}^p \asymp b_{0,1}^p$ for all $i < m$ and all $j < \ell$, with $\ell \geq 4$. We shall check that $b_{i,\ell}^p \asymp b_{0,1}^p$ for all $i < m$. Write $\ell = \alpha(n) + k$ for some $n \geq 0$ and $0 \leq k \leq 2m - 1$. Since $nm + \beta(k) + 1 < \alpha(n) + k = \ell$, we get by induction that

$b_{0,1}^p \asymp b_{i,nm+\beta(k)+1}^p$. Observe also that $b_{i,nm+\beta(k)+1}^p \leq b_{i,\alpha(n)+k}^p$ by (5.6). Now assume that $k > 0$. Then we get from the relation

$$b_{i,\alpha(n)}^p = b_{i,\alpha(n)}^p + b_{i,nm+1}^p + b_{[0,m-1] \times [\alpha(n)+1, \alpha(n+1)]}^p + z_p$$

that $b_{i,\alpha(n)+k}^p \leq b_{i,\alpha(n)}^p$ so that

$$b_{0,1}^p \asymp b_{i,nm+\beta(k)+1}^p \leq b_{i,\alpha(n)+k}^p \leq b_{i,\alpha(n)}^p \asymp b_{0,1}^p.$$

We conclude that $b_{i,\alpha(n)+k}^p \asymp b_{0,1}^p$.

Assume finally that $k = 0$. Then $\ell = \alpha(n)$ with $n \geq 1$, and we get from the relation

$$b_{i,\alpha(n-1)}^p = b_{i,\alpha(n-1)}^p + b_{i,nm-m+1}^p + b_{[0,m-1] \times [\alpha(n-1)+1, \alpha(n)]}^p + z_p$$

that $b_{i,\alpha(n)}^p \leq b_{i,\alpha(n-1)}^p \asymp b_{0,1}^p$ and so

$$b_{0,1}^p \asymp b_{i,nm+1}^p \leq b_{i,\alpha(n)}^p \leq b_{i,\alpha(n-1)}^p \asymp b_{0,1}^p,$$

which proves that $b_{i,\alpha(n)}^p \asymp b_{0,1}^p$.

As M is separative, it embeds into a product of monoids of the form $G \cup \{\infty\}$, for abelian groups G (this follows immediately from Hewitt and Zuckermann's result [7, Theorem 5.59]). Hence, to prove that φ is one-to-one, it is sufficient to establish the following claim.

Claim. *For any abelian group G and any set of elements*

$$\{\tilde{p} \mid p \in \mathbb{P}(M)\} \cup \{\tilde{b}_{i,j}^p \mid p \in \mathbb{P}_{\text{reg}}(M), i < |\mathbb{L}_{\text{free}}(M, p)|, j \in \mathbb{Z}^+\}$$

in $G \cup \{\infty\}$ satisfying the relations (5.1)–(5.6), the relations (5.9) and (5.10) are also satisfied.

Proof of Claim. We prove the claim by induction on the height of p in $\mathbb{P}(M)$. If p is a minimal prime then (5.9) holds vacuously and (5.10) follows from (5.2). Assume now that p is a prime of height $h + 1$ and the result holds for all primes of height at most h . Assume first that p is a free prime. Note that the induction hypothesis together with (5.1) gives us $\tilde{p} = \tilde{p} + \tilde{u}_p$, where $\tilde{u}_p = \sum_{q \in \mathbb{L}(M, p)} \tilde{q}$. If $\mathbb{L}_{\text{free}}(M, p) = \emptyset$, then all the elements of $\mathbb{L}(M, p)$ are regular and so $\tilde{u}_p + \tilde{q} = \tilde{u}_p$ for all $q \in \mathbb{L}(M, p)$ by induction hypothesis. We get that

$$\tilde{p} + \tilde{q} = \tilde{p} + \tilde{u}_p + \tilde{q} = \tilde{p} + \tilde{u}_p = \tilde{p}.$$

This proves (5.9) for all $q \in \mathbb{L}(M, p)$, and thus for all prime $q < p$ by induction hypothesis. Assume now that $\mathbb{L}_{\text{free}}(M, p) \neq \emptyset$. By assumption, $\mathbb{L}_{\text{free}}(M, p) = \{p_*\}$ for some p_* . Since $\tilde{p} = \tilde{p} + \tilde{p}_* + \sum_{q \in \mathbb{L}_{\text{reg}}(M, p)} \tilde{q}$, the induction hypothesis gives again that $\tilde{p} = \tilde{p} + \tilde{q}$ for all $q \in \mathbb{L}_{\text{reg}}(M, p)$, so that $\tilde{p} = \tilde{p} + \sum_{q \in \mathbb{L}_{\text{reg}}(M, p)} \tilde{q}$. From this we get $\tilde{p} + \tilde{p}_* = \tilde{p} + \sum_{q \in \mathbb{L}_{\text{reg}}(M, p)} \tilde{q} + \tilde{p}_* = \tilde{p}$. As before, this gives (5.9) at p .

Assume now that p is a regular prime. The case where $\mathbb{L}_{\text{free}}(M, p) = \emptyset$ is similar to the case above. So assume that $|\mathbb{L}_{\text{free}}(M, p)| = m > 0$. As $\tilde{b}_{i,j}^p \asymp \tilde{b}_{0,0}^p = \tilde{p}$, some $\tilde{b}_{i,j}^p = \infty$ if and only if all $\tilde{b}_{i,j}^p = \infty$. In this case, (5.10) holds trivially. So we can assume that all $\tilde{b}_{i,j}^p$ belong to G . Now (5.4) and (5.5) give

$$\begin{aligned} 0 &= \tilde{b}_{i,0}^p + \tilde{b}_{i,1}^p + \tilde{b}_{i,2}^p + \tilde{p}i \\ 0 &= \tilde{b}_{i,0}^p + \tilde{b}_{i,1}^p + \tilde{b}_{i,2}^p \end{aligned} \tag{5.11}$$

and so $\tilde{p}_i = 0$ for all $i < m$. Furthermore, (5.6) gives that $\tilde{z}_p \in G$, and thus $\tilde{q} \in G$ for all $q \in L_{\text{reg}}(M, p)$; from $\tilde{q} = 2\tilde{q}$ for all such q , together with all $\tilde{p}_i = 0$, we finally get that $\tilde{q} = 0$ for any $q \in L(M, p)$, and thus, by the induction hypothesis, $\tilde{q} = 0$ for any $q < p$ in $\mathbb{P}(M)$.

Now let j be a positive integer. There exists a unique natural number n such that $nm < j \leq (n+1)m$. The integer $k = 2j - 2nm - 2$ lies in the interval $[0, 2m - 2]$, and $j = nm + \beta(k) + 1$. Therefore, by applying (5.6) with the consecutive values k and $k + 1$, we obtain that $\tilde{b}_{i,j}^p = 0$. In particular, $\tilde{b}_{i,1}^p = \tilde{b}_{i,2}^p = 0$, whence, by (5.11), $\tilde{b}_{i,0}^p = 0$, and therefore $\tilde{b}_{i,j}^p = 0$ for all $i < m$ and all $j \in \mathbb{Z}^+$. This concludes the proof of the claim. \square Claim.

This concludes the proof of Theorem 5.1. \square

From Theorem 5.1 we can deduce immediately the following closure result for the class of graph monoids.

Corollary 5.2. *Let M be a primitive monoid with $\mathbb{P}(M)$ lower finite. Then M is a retract of some graph monoid iff M is a graph monoid.*

Observe that even in case M is finitely generated, the quiver constructed in the proof of Theorem 5.1 may not be finite. That in some cases that quiver cannot be made finite will be established in Theorem 6.1.

The analogue of Corollary 5.2 for graph monoids of *finite* quivers does not hold, see Example 6.5.

Not all antisymmetric graph monoids have lower finite set of primes. For example, letting E be the row-finite quiver represented by Figure 2, the monoid $M(E)$ is defined by the generators p_n and the relations $p_n = p_n + p_{n+1}$, for $n \in \mathbb{Z}^+$. As all the p_n s are prime in $M(E)$ and $p_0 > p_1 > p_2 > \dots$, the subset $\mathbb{P}(M(E))$ is not lower finite.




FIGURE 2. A quiver whose graph monoid has no lower finite set of primes.

6. A CHARACTERIZATION OF ANTISYMMETRIC GRAPH MONOIDS OF FINITE QUIVERS

We characterize in this section those antisymmetric finitely generated refinement monoids M which are isomorphic to a graph monoid $M(E)$ for a *finite* quiver E .

Theorem 6.1. *Let M be a finitely generated primitive monoid. Then there exists a finite quiver E such that $M \cong M(E)$ if and only if the set $\mathbb{P}_{\text{reg}}(M)$ of regular primes is a lower subset of $\mathbb{P}(M)$ and $|L_{\text{free}}(M, p)| \leq 1$ for each $p \in \mathbb{P}_{\text{free}}(M)$. Equivalently, the set R of regular elements of M is an *o-ideal* of M and $\mathbb{P}(M/R)$ is a forest.*

One implication follows easily from Theorem 5.1. Indeed, assume that the set $\mathbb{P}_{\text{reg}}(M)$ of regular primes is a lower subset of $\mathbb{P}(M)$ and that $|L_{\text{free}}(M, p)| \leq 1$ for each $p \in \mathbb{P}_{\text{free}}(M)$. Then the quiver E built in the proof of Theorem 5.1 is finite because the case 3 of the proof of Theorem 5.1 does not occur. Thus $M \cong M(E)$ for the finite quiver E .

The other implication will be proved at the end of this section. We start with a crucial observation.

Proposition 6.2. *Let E be a finite quiver. Assume that $M(E)$ is an antisymmetric monoid. Then $\mathbb{P}_{\text{reg}}(M(E))$ is a lower subset of $\mathbb{P}(M(E))$.*

Proof. By [5, Corollary 5.9], each element in $M = M(E)$ is either free or regular.

Observe that if q is a regular prime then $\phi_{q'}(q) \in \{0, \infty\}$ for all $q' \in \mathbb{P}(M)$, according to whether $q' \not\leq q$ or $q' \leq q$ respectively. We proceed by way of contradiction. Let p be a minimal element in $\mathbb{P}_{\text{reg}}(M)$ (with respect to \leq) with the property that $q \leq p$ for some free prime q . Since p is prime, $p = \bar{v}$ for some $v \in E^0$. Let H be the hereditary saturated subset of E^0 generated by v . Then $M(H)$ is the o-ideal generated by p , so that

$$M(H) = M(E) \mid p = \{x \in M(E) \mid x \leq p\},$$

where the latter equality follows from the regularity of p . Replacing E by H , we can assume that $\phi_q(p') = 0$ for every free prime q and for every regular prime p' such that $p' \neq p$. Put

$$\begin{aligned} U &= \{z \in E^0 \mid \bar{z} \neq p\} = \{x_i \mid 1 \leq i \leq m\}, \\ V &= \{z \in E^0 \mid \bar{z} = p\} = \{y_j \mid 1 \leq j \leq n\}. \end{aligned}$$

For $1 \leq i, i' \leq m$ and $1 \leq j, j' \leq n$, set

$$\begin{aligned} \alpha_{i,i'} &= |\{e \in E^1 \mid e: x_i \rightarrow x_{i'}\}|, \\ \beta_{j,j'} &= |\{e \in E^1 \mid e: y_j \rightarrow y_{j'}\}|, \\ \gamma_{i,j} &= |\{e \in E^1 \mid e: y_j \rightarrow x_i\}|. \end{aligned}$$

A presentation of $M(E)$ is obtained in matricial form as follows:

$$X = AX, \quad Y = BY + CX, \quad (6.1)$$

where $X = (x_1, \dots, x_m)^t$, $Y = (y_1, \dots, y_n)^t$, $A = (\alpha_{i,i'})_{1 \leq i, i' \leq m}$, $B = (\beta_{j,j'})_{1 \leq j, j' \leq n}$, and $C = (\gamma_{i,j})_{1 \leq i \leq m, 1 \leq j \leq n}$.

Claim. *For any abelian group G , the only $X \in M_{m \times 1}(G)$ and $Y \in M_{n \times 1}(G)$ that satisfy (6.1) are $X = 0$ and $Y = 0$.*

Proof of Claim. Let M be the submonoid of G generated by $\{x_1, \dots, x_m, y_1, \dots, y_n\}$. There is a unique monoid homomorphism $\psi: M(E) \rightarrow M$ such that $\psi(\bar{x}_i) = x_i$ and $\psi(\bar{y}_j) = y_j$ for all i, j . But since $\bar{x}_i + \bar{y}_j = \bar{y}_j = 2\bar{y}_j$ in $M(E)$, we get that $x_i + y_j = y_j = 2y_j$ in G so that $x_i = y_j = 0$ for all i, j . \square Claim.

We can assume that $\mathbb{P}_{\text{free}}(M) = \{\bar{x}_1, \dots, \bar{x}_k\}$, so that $\bar{x}_1, \dots, \bar{x}_k$ are the different free primes in the collection $\bar{x}_1, \dots, \bar{x}_m$. Consider the “free part” of the homomorphism ϕ of (2.1):

$$\phi_{\text{free}}: M(E) \rightarrow (\mathbb{Z}^\infty)^k, \quad x \mapsto (\phi_{\bar{x}_1}(x), \dots, \phi_{\bar{x}_k}(x)).$$

We can suppose that \bar{x}_1 is a maximal element in $\mathbb{P}_{\text{free}}(M)$ (with respect to \leq). Then $\phi_{\bar{x}_1}(\bar{x}_i) < \infty$ for every free prime \bar{x}_i , and $\phi_{\bar{x}_1}(\bar{x}_j) = 0$ for every regular prime \bar{x}_j , because $\bar{x}_j \neq p$ and $\phi_q(p') = 0$ for every $q \in \mathbb{P}_{\text{free}}(M)$ and every regular prime p' with $p' \neq p$. It follows that $\phi_{\bar{x}_1}(\bar{x}_i) < \infty$ for $i = 1, \dots, m$ and so we obtain that the column matrix

$$X_0 = (\phi_{\bar{x}_1}(\bar{x}_1), \dots, \phi_{\bar{x}_1}(\bar{x}_m))^t \in M_{m \times 1}(\mathbb{Z}^+)$$

satisfies $X_0 = AX_0$. Observe that $\phi_{\bar{x}_1}(\bar{x}_1) = 1$, so that $X_0 \neq 0$. Take $G = \mathbb{Q}$ and any $Y \in M_{n \times 1}(\mathbb{Q})$ such that $Y = BY$. Then $X = 0$ and Y give a solution to (6.1), and so $Y = 0$ by the Claim. Thus $(I - B)Y = 0$ implies $Y = 0$, and so $I - B$ is an invertible matrix in $M_n(\mathbb{Q})$. Consider now the above column matrix $0 \neq X_0 \in M_{m \times 1}(\mathbb{Z}^+)$, and set $Y_0 = (I - B)^{-1}CX_0 \in M_{n \times 1}(\mathbb{Q})$. Then we get a solution (X_0, Y_0) over \mathbb{Q} of the equation (6.1) with $X_0 \neq 0$, a contradiction. \square

Lemma 6.3. *Let N be an o -ideal in a primitive monoid M . Then the following properties hold:*

- (1) $\mathbb{P}(N) = \mathbb{P}(M) \cap N$.
- (2) M/N is a primitive monoid and the canonical map $\pi: M \rightarrow M/N$ induces an \triangleleft -isomorphism from $\mathbb{P}(M) \setminus \mathbb{P}(N)$ onto $\mathbb{P}(M/N)$. Moreover,

$$\phi_{\pi(p)}^{M/N}(\pi(a)) = \phi_p^M(a),$$

for every $p \in \mathbb{P}(M) \setminus \mathbb{P}(N)$ and every $a \in M$.

Proof. (1) Straightforward.

(2) Let us prove that M/N is antisymmetric. The rest is easy. It suffices to prove that $a + x \leq_I a$ implies that $a + x \equiv_I a$, for all $a, x \in M$. Let $h \in I$ such that $a + x \leq a + h$. It follows from [5, Corollary 4.2] that there exists $u \ll a$ such that $x \leq h + u$, and hence there are $h' \leq h$ and $u' \leq u$ such that $x = h' + u'$. From $h' \leq h$ it follows that $h' \in I$, while from $u' \leq u$ it follows that $u' \ll a$. As M is antisymmetric, $u' + a = a$, and thus

$$a + x = (a + u') + h' = a + h' \equiv_I a. \quad \square$$

Corollary 6.4. *Let E be a finite quiver. Assume that $M = M(E)$ is an antisymmetric monoid. Let N be the submonoid of M generated by $\mathbb{P}_{\text{reg}}(M(E))$. Then:*

- (1) N is an o -ideal of M .
- (2) N is the set of regular elements of M .
- (3) Let $H = \{v \in E^0 \mid \bar{v} \in N\}$. Then H is a hereditary saturated subset of E^0 and the quiver $E \setminus H$ satisfies that $M(E \setminus H) = M/N$ is an antisymmetric graph monoid such that all elements are free, and $\mathbb{P}(M/N) = \mathbb{P}_{\text{free}}(M)$.

Proof. (1) Take $p_1, \dots, p_r \in \mathbb{P}_{\text{reg}}(M)$ and $a \in M$ such that $a \leq p_1 + \dots + p_r$. Then $a = q_1 + \dots + q_\ell$, where $q_i \in \mathbb{P}$ so that $q_i \leq p_1 + \dots + p_r$ and, by primeness of q_i , we get $q_i \leq p_j$ for some j . By Proposition 6.2 we get that all q_i are regular, hence a is regular.

(2) If $a \in M$ then, by the decomposition result given in [14, Proposition 3.4.4] or [5, Theorem 5.8], we can write

$$a = p_1 + \dots + p_r + n_1 q_1 + \dots + n_\ell q_\ell,$$

where p_1, \dots, p_r are regular primes, q_1, \dots, q_ℓ are free primes, $p_1, \dots, p_r, q_1, \dots, q_\ell$ are pairwise incomparable, and this expression is unique. If a is a regular element, then $2a = a$ and it follows from the equality

$$2a = p_1 + \dots + p_r + 2n_1 q_1 + \dots + 2n_\ell q_\ell,$$

that all $n_i = 0$, so $a \in N$.

(3) Clear from (1), (2), and Lemma 6.3. By [4, Lemma 6.6] we obtain that $M/N = M(E \setminus H)$ is a graph monoid. \square

We are now ready to complete the proof of Theorem 6.1. Assume that E is a finite quiver such that $M(E)$ is an antisymmetric monoid. Then we get from Corollary 6.4 that the set R of regular elements of $M(E)$ is an \mathfrak{o} -ideal and the monoid $M(E)/R$ is an antisymmetric graph monoid with no regular primes. Theorem 5.1 gives that $\mathbb{P}(M(E)/R)$ is a forest, as desired.

Example 6.5. It follows from Theorem 5.1 that the monoid \mathbb{Z}^∞ is a graph monoid. On the other hand, by Theorem 6.1, \mathbb{Z}^∞ is not the graph monoid of any finite quiver.

Nevertheless, \mathbb{Z}^∞ is a *retract* of the graph monoid of a finite quiver. Indeed, consider the quiver E represented by Figure 3.




FIGURE 3. A quiver whose graph monoid retracts to \mathbb{Z}^∞ .




FIGURE 4. A quiver that represents \mathbb{Z}^∞ .

A presentation of $M(E)$ is given by the two equations

$$a = a + 1, \quad b = 2b + a.$$

As $a + b$ is idempotent and absorbs 1 in $M(E)$, there are unique monoid homomorphisms $\varepsilon: \mathbb{Z}^\infty \rightarrow M(E)$ and $\rho: M(E) \rightarrow \mathbb{Z}^\infty$ such that

$$\begin{aligned} \varepsilon(1) &= 1, & \varepsilon(\infty) &= a + b, \\ \rho(1) &= 1, & \rho(a) = \rho(b) &= \infty. \end{aligned}$$

In particular, $\rho \circ \varepsilon = \text{id}_{\mathbb{Z}^\infty}$, so \mathbb{Z}^∞ is a retract of $M(E)$.

The proof of Theorem 5.1 gives the following infinite presentation for the monoid \mathbb{Z}^∞ :

$$\begin{aligned} b_0 &= 2b_0 + b_1 + b_2 + 1; \\ b_1 &= b_0 + 2b_1 + b_2; \\ b_2 &= b_2 + b_1 + b_3 + b_4; \\ b_3 &= 2b_3 + 2b_1 + b_4; \\ b_4 &= b_4 + b_2 + b_5 + b_6; \\ b_5 &= 2b_5 + 2b_2 + b_6; \\ b_6 &= b_6 + b_3 + b_7 + b_8; \\ &\dots \quad \dots \end{aligned}$$

The corresponding quiver is represented by Figure 4.

7. OPEN PROBLEMS

Problem 1. Is it decidable whether a given finitely generated monoid is isomorphic to the graph monoid of some row-finite (resp., finite) quiver?

As every finitely generated commutative monoid is finitely presented, Problem 1 is well-posed. The two main results of the present paper, Theorems 5.1 and 6.1, solve the analogue of Problem 1 for *antisymmetric* monoids.

Problem 2. Is the retract of a graph monoid always a graph monoid?

By Corollary 5.2, the answer to Problem 2 for primitive monoids with lower finite set of primes is positive. On the other hand, by Example 6.5, its analogue for graph monoids of *finite* quivers fails. Also, observe that the class of graph monoids is not closed under direct limits (with respect to monoid homomorphisms). Indeed, the results of [16, 18] show that there exists a distributive bounded semilattice S that is not representable (i.e., isomorphic to $\mathcal{V}(R)$ for any von Neumann regular ring R). As, by [2, Theorem 4.4], all graph monoids are representable, we see that S cannot be a graph monoid. On the other hand, S , as every distributive semilattice, is a direct limit of finite Boolean semilattices [6, 13], and by Theorem 6.1, every finite distributive semilattice is the graph monoid of a finite quiver, thus representable. This shows that S is a direct limit of graph monoids, without being itself a graph monoid.

REFERENCES

- [1] G. Abrams and G. Aranda Pino, *The Leavitt path algebra of a graph*, J. Algebra **293**, no. 2 (2005), 319–334.
- [2] P. Ara and M. Brustenga, *The regular algebra of a quiver*, J. Algebra **309**, no. 1 (2007), 207–235.
- [3] P. Ara, K. R. Goodearl, K. C. O’Meara, and E. Pardo, *Separative cancellation for projective modules over exchange rings*, Israel J. Math. **105** (1998), 105–137.
- [4] P. Ara, M. A. Moreno, and E. Pardo, *Nonstable K-theory for graph algebras*, Algebr. Represent. Theory **10**, no. 2 (2007), 157–178.
- [5] G. Brookfield, *Cancellation in primely generated refinement monoids*, Algebra Universalis **46**, no. 3 (2001), 342–371.
- [6] S. Bulman-Fleming and K. McDowell, *Flat semilattices*, Proc. Amer. Math. Soc. **72** (1978), 228–232.
- [7] A. H. Clifford and G. B. Preston, “The Algebraic Theory of Semigroups, Vol. I”. Math. Surveys and Monographs **7**, Amer. Math. Soc., Providence, R. I., 1961. xv+224 p.
- [8] J. Cuntz, *Simple C^* -algebras generated by isometries*, Comm. Math. Phys. **57**, no. 2 (1977), 173–185.
- [9] H. Dobbertin, *Refinement monoids, Vaught monoids, and Boolean algebras*, Math. Ann. **265**, no. 4 (1983), 473–487.
- [10] K. R. Goodearl, “Von Neumann Regular Rings”, 2nd ed., Krieger, Malabar, Fl., 1991. xvi+412 p.
- [11] K. R. Goodearl, *Von Neumann regular rings and direct sum decomposition problems*, in: *Abelian Groups and Modules, Padova 1994* (A. Facchini and C. Menini, eds.), Dordrecht (1995) Kluwer, 249–255.
- [12] K. R. Goodearl, E. Pardo, and F. Wehrung, *Semilattices of groups and inductive limits of Cuntz algebras*, J. Reine Angew. Math. **588** (2005), 1–25.
- [13] K. R. Goodearl and F. Wehrung, *Representations of distributive semilattices in ideal lattices of various algebraic structures*, Algebra Universalis **45**, no. 1 (2001), 71–102.
- [14] R. S. Pierce, “Countable Boolean Algebras”, in *Handbook of Boolean Algebras*, Vol. **3**, edited by J. D. Monk with R. Bonnet, Elsevier, 1989, 775–876.
- [15] I. Raeburn, “Graph Algebras”, CBMS Reg. Conf. Ser. Math., Vol. **103**, Amer. Math. Soc., Providence, RI, 2005. vi+113 p.

- [16] F. Wehrung, *Non-measurability properties of interpolation vector spaces*, Israel J. Math. **103** (1998), 177–206.
- [17] F. Wehrung, *The dimension monoid of a lattice*, Algebra Universalis **40**, no. 3 (1998), 247–411.
- [18] F. Wehrung, *A uniform refinement property for congruence lattices*, Proc. Amer. Math. Soc. **127**, no. 2 (1999), 363–370.

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193, BEL-LATERRA (BARCELONA), SPAIN

E-mail address, P. Ara: para@mat.uab.cat

E-mail address, F. Perera: perera@mat.uab.cat

CNRS UMR 6139, UNIVERSITÉ DE CAEN, CAMPUS 2, DÉPARTEMENT DE MATHÉMATIQUES, BP 5186, 14032 CAEN CEDEX, FRANCE

E-mail address, F. Wehrung: wehrung@math.unicaen.fr

URL, F. Wehrung: <http://www.math.unicaen.fr/~wehrung>