HAL open science

Stopped diffusion processes: boundary corrections and overshoot

Emmanuel Gobet, Stéphane Menozzi

To cite this version:

Emmanuel Gobet, Stéphane Menozzi. Stopped diffusion processes: boundary corrections and overshoot. Stochastic Processes and their Applications, 2010, 120 (2), pp.130-162. 10.1016/j.spa.2009.09.014 . hal-00157975v3

HAL Id: hal-00157975

https://hal.science/hal-00157975v3

Submitted on 18 Jun 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stopped diffusion processes: boundary corrections and overshoot

Emmanuel Gobet [a,*], Stéphane Menozzi [b]

[a] *Laboratoire Jean Kuntzmann, Université de Grenoble and CNRS, BP 53, 38041 Grenoble Cedex 9, FRANCE*

[b] *Laboratoire de Probabilités et Modèles Aléatoires, Université Denis Diderot Paris 7, 175 rue de Chevaleret 75013 Paris, FRANCE.*

Abstract

For a stopped diffusion process in a multidimensional time-dependent domain $\mathcal{D}$, we propose and analyse a new procedure consisting in simulating the process with an Euler scheme with step size Δ and stopping it at discrete times $(i\Delta)_{i\in\mathbf{N}^*}$ in a modified domain, whose boundary has been appropriately shifted. The shift is locally in the direction of the inward normal $n(t,x)$ at any point (t,x) on the parabolic boundary of $\mathcal{D}$, and its amplitude is equal to $0.5826(...)|n^*\sigma|(t,x)\sqrt{\Delta}$ where σ stands for the diffusion coefficient of the process. The procedure is thus extremely easy to use. In addition, we prove that the rate of convergence w.r.t. Δ for the associated weak error is higher than without shifting, generalizing previous results by [BGK97] obtained for the one dimensional Brownian motion. For this, we establish in full generality the asymptotics of the triplet exit time/exit position/overshoot for the discretely stopped Euler scheme. Here, the overshoot means the distance to the boundary of the process when it exits the domain. Numerical experiments support these results.

Key words: Stopped diffusion, Time-dependent domain, Brownian overshoot, Boundary sensitivity.
1991 MSC: 60J60, 60H35, 60-08

* Corresponding author
Email addresses: `emmanuel.gobet@imag.fr` (Emmanuel Gobet), `menozzi@math.jussieu.fr` (Stéphane Menozzi).

1 Introduction

1.1 Statement of the problem

We consider a d-dimensional diffusion process whose dynamics is given by

$$X_t = x + \int_0^t b(s, X_s)ds + \int_0^t \sigma(s, X_s)dW_s \tag{1.1}$$

where W is a standard d'-dimensional Brownian motion defined on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t\geq 0}, \mathbb{P})$ satisfying the usual conditions. The mappings b and σ are Lipschitz continuous in space and locally bounded in time, so that (1.1) has a unique strong solution. We consider $(D_t)_{t\geq 0}$, a time-dependent family of smooth bounded domains of $\mathbb{R}^d$, that is also smooth with respect to t (we refer to paragraph 1.5.2 for a precise definition). See Figure 1. For a fixed deterministic time $T > 0$, this defines a time-space domain

$$\mathcal{D} = \bigcup_{0<t<T} \{t\} \times D_t = \{(t, x) : 0 < t < T, x \in D_t\} \subset]0, T[\times \mathbb{R}^d.$$

Cylindrical domains are specific cases of time-dependent domains of the form $\mathcal{D} =]0, T[\times D$, where D is a usual domain of $\mathbb{R}^d$ ($D_t = D$ for any t). Time-dependent domains in dimension $d = 1$ are typically of the form $\mathcal{D} = \{(t, x) : 0 < t < T, \varphi_1(t) < x < \varphi_2(t)\}$ for two functions φ_1 and φ_2 (the time-varying boundaries).



Figure 1. Time space domain and its time-sections.

Now, set $\tau := \inf\{t > 0 : X_t \notin D_t\}$, then $\tau \wedge T$ is the first exit time of $(s, X_s)_s$ from the time-space domain $\mathcal{D}$. Given continuous functions $g, f, k : \bar{\mathcal{D}} \to \mathbb{R}$, we are interested in estimating the quantity

$$\mathbb{E}_x[g(\tau \wedge T, X_{\tau\wedge T})Z_{\tau\wedge T} + \int_0^{\tau\wedge T} Z_s f(s, X_s)ds], Z_s = \exp(-\int_0^s k(r, X_r)dr), \tag{1.2}$$

where as usual $\mathbb{E}_x[.] := \mathbb{E}[.|X_0 = x]$ (resp. $\mathbb{P}_x[.] := \mathbb{P}[.|X_0 = x]$). The approximation of such quantities is a well known issue in finance, since it represents in

this framework the price of a barrier option, see e.g. Andersen and Brotherton-Ratcliffe [ABR96]. These quantities also arise through the Feynman-Kac representation of the solution of a parabolic PDE with Cauchy-Dirichlet boundary conditions, see Costantini *et al.* [CGK06]. They can therefore also be related to problems of heat diffusion in time-dependent domains.

We then choose to approximate the expectation in (1.2) by Monte Carlo simulation. This approach is natural and especially relevant compared to deterministic methods if the dimension d is large. To this end we approximate the diffusion (1.1) by its Euler scheme with time step $\Delta > 0$ and discretization times $(t_i = i\Delta = iT/m)_{i\geq 0}$ ($m \in \mathbf{N}^*$ so that $t_m = T$). For $t \geq 0$, define $\phi(t) = t_i$ for $t_i \leq t < t_{i+1}$ and introduce

$$X_t^\Delta = x + \int_0^t b(\phi(s), X_{\phi(s)}^\Delta)ds + \int_0^t \sigma(\phi(s), X_{\phi(s)}^\Delta)dW_s. \tag{1.3}$$

We now associate to (1.3) the discrete exit time $\tau^\Delta := \inf\{t_i > 0 : X_{t_i}^\Delta \notin D_{t_i}\}$. Approximating the functional $V_\tau := g(\tau \wedge T, X_{\tau\wedge T})Z_{\tau\wedge T} + \int_0^{\tau\wedge T} Z_s f(s, X_s)\, ds$ by

$$\begin{aligned} V_{\tau^\Delta}^\Delta &:= g(\tau^\Delta \wedge T, X_{\tau^\Delta\wedge T}^\Delta)Z_{\tau^\Delta\wedge T}^\Delta + \int_0^{\tau^\Delta\wedge T} Z_{\phi(s)}^\Delta f(\phi(s), X_{\phi(s)}^\Delta)ds \\ &\text{with} \quad Z_t^\Delta = e^{-\int_0^t k(\phi(r), X_{\phi(r)}^\Delta)dr}, \end{aligned}$$

we introduce the quantity

$$\mathrm{Err}(T, \Delta, g, f, k, x) = \mathbb{E}_x[V_{\tau^\Delta}^\Delta - V_\tau] \tag{1.4}$$

that will be referred to as the weak error.

Note that in $V_{\tau^\Delta}^\Delta$, on $\{\tau^\Delta \leq T\}$ g is a.s. not evaluated on the side part $\bigcup_{0\leq t\leq T}\{t\} \times \partial D_t$ of the boundary (g must be understood as a function defined in a neighborhood of the boundary). At first sight, this approximation can seem coarse. Anyhow, it does not affect the convergence rate and really reduces the computational cost with respect to the alternative that would consist in taking the projection on $\partial\mathcal{D}$. It is a commonly observed phenomenon that the error is positive when g is positive (overestimation of $\mathbb{E}_x(V_\tau)$), because we neglect the possible exits between two discrete times: see Boyle and Lau [BL94], Baldi [Bal95], Gobet and Menozzi [GM04]. In addition, it is known that the error is of order $\Delta^{1/2}$: see [GM04] for lower bound results, see [GM07] for upper bounds in the more general case of It processes. But so far, the derivation of an error expansion $\mathbb{E}_x[V_{\tau^\Delta}^\Delta - V_\tau] = C\sqrt{\Delta} + o(\sqrt{\Delta})$ had not been established: this is one of the intermediary results of the current work (see Theorem 4).

Our goal goes beyond this result, by designing a simple and very efficient improved procedure. We propose to stop the Euler scheme at its exit of a



Figure 2. The boundary ∂D_t and the smaller domain D_t^Δ.

smaller domain in order to compensate the underestimation of exits and to achieve an error of order $o(\sqrt{\Delta})$. The smaller domain is defined by its time-section

$$D_t^\Delta = \{x \in D_t : \mathbf{d}(x, \partial D_t) > c_0\sqrt{\Delta}|n^*\sigma(t,x)|\}$$

where $n(t,x)$ is the inward normal vector at the closest point of x on the boundary ∂D_t, see Figures 2 and 3 for details [1] . We shall interpret $|n^*\sigma(t,x)|$ as the noise amplitude along the normal direction to the boundary. The constant c_0 is defined later in (2.1) and equals approximatively 0.5826(...). Thus, the associated exit time of the Euler scheme is given by

$$\hat{\tau}^\Delta = \inf\{t_i > 0 : X_{t_i}^\Delta \notin D_{t_i}^\Delta\} \leq \tau^\Delta.$$

The new Monte Carlo scheme consists in simulating independent realizations of

$$V_{\hat{\tau}^\Delta}^\Delta = g(\hat{\tau}^\Delta \wedge T, X_{\hat{\tau}^\Delta \wedge T}^\Delta) Z_{\hat{\tau}^\Delta \wedge T}^\Delta + \int_0^{\hat{\tau}^\Delta \wedge T} Z_{\phi(s)}^\Delta f(\phi(s), X_{\phi(s)}^\Delta) ds$$

and averaging them out to get an estimator of the required quantity $\mathbb{E}_x(V_\tau)$. Our main result (Theorem 5) is that the asymptotic bias w.r.t. Δ is significantly improved:

$$\mathbb{E}_x[V_{\hat{\tau}^\Delta}^\Delta - V_\tau] = o(\sqrt{\Delta})$$

(instead of $C\sqrt{\Delta} + o(\sqrt{\Delta})$ before). This improvement has been already established in the case of the one-dimensional Brownian motion [BGK97] in the context of computational finance, exploiting heavily the connection with Gaussian random walks and some explicit computations available in the Brownian motion case.

[1] the closest point of x may not be unique for points x far from ∂D_t. But since the above definition of D_t^Δ involves only points close to the boundary, this does not make any difference.

1.2 Contribution of the paper

To achieve the results in the current very general framework, we combine several ingredients (which correspond to the main steps of the proofs).

(1) We first expand the error $\mathbb{E}_x[V^\Delta_{\tau^\Delta} - V_\tau]$ related to the use of the discrete Euler scheme in the domain $\mathcal{D}$. Although this issue deserved many studies in the literature, the expansion results are new. We prove that it relies on the study of the weak convergence of the triplet (exit time, position at exit time, renormalized overshoot at exit time), that is $(\tau^\Delta, X^\Delta_{\tau^\Delta}, \Delta^{-1/2} d(X^\Delta_{\tau^\Delta}, \partial D_{\tau^\Delta}))$, as Δ goes to 0. This weak convergence result is crucial in this work and it is new (see Theorem 3).
Then, combining this with sharp techniques of error analysis, we derive an expansion of the form $\mathrm{Err}(T, \Delta, g, f, k, x) = C\sqrt{\Delta} + o(\Delta)$ in the very general framework of stopped diffusions in time-dependent domains.

(2) Second, we analyse the impact of the boundary shifting, in the continuous time problem (see paragraph 2.3.2). This is related to the differentiability of $\mathbb{E}_x(V_\tau)$ w.r.t. the boundary and it has been addressed in [CGK06]. We apply directly their results. Then, we obtain the global error estimate of the boundary correction procedure (Theorem 5).

We mention that the previous results about the error expansion and correction still hold in the stationary setting, see Section 4, which also seems to be new. A numerical application is discussed in Section 5. Complementary tests are presented in [Gob09], showing that the boundary correction procedure is very generic and seems to work without Markovian property for X. This feature will be investigated in further research.

Let us finally mention that we could also consider the diffusion process discretely stopped: expansion and correction results below would remain the same.

1.3 Comparison with results in literature

Up to now, the behavior of (1.4) had mainly been analysed for cylindrical domains, in the killed case, without source and potential terms (i.e. when the error writes $\mathrm{Err}(T, \Delta, g, 0, 0, x) = \mathbb{E}_x[g(X^\Delta_T)\mathbf{1}_{\tau^\Delta > T}] - \mathbb{E}_x[g(X_T)\mathbf{1}_{\tau > T}]$). Let us first mention the work of Broadie *et al.* [BGK97], who first derived the boundary shifting procedure in the one dimensional geometric Brownian motion setting (Black and Scholes model). In [Gob00] and [GM04], it had been shown that, under some (hypo)ellipticity conditions on the coefficients and some smoothness of the domain and the coefficients, $\mathrm{Err}(T, \Delta, g, 0, 0, x)$ was lower and upper bounded at order 1/2 w.r.t. the time-step Δ. Also, an expan-

sion result for the killed Brownian motion in a cone as well as the associated correction procedure are available in [Men06].

All these works emphasize that the crucial quantity to analyse in order to obtain an expansion is the overshoot above the spatial boundary of the discrete process. In the Brownian one-dimensional framework such analysis goes back to Siegmund [Sie79] and Siegmund and Yuh [SY82]. Also a non linear renewal theory for random walk, i.e. for a curved boundary, had been developed by Siegmund and *al.*, see [Sie85] and references therein, Woodroofe [Woo82] and Zhang [Zha88]. We manage to extend their results to obtain the asymptotic distribution of the overshoot of the Euler scheme, see Sections 2 and 3. Concerning the asymptotics of the overshoot of stochastic processes, let us mention the works of Alsmeyer [Als94] or Fuh and Lai [FL01] for ergodic Markov chains and Doney and Kyprianou for Lvy processes [DK06]. These works are all based on renewal arguments.

Finally, for simulating stopped diffusions we also mention the alternative technique based on Random Walks on Spheres. This method allows to derive a bound for the weak error associated to the approximation of $\mathbb{E}[V_\tau]$ in the elliptic setting for a cylindrical domain, see Milstein [Mil97]. The same approach has also been exploited to obtain some strong error or pathwise bounds for a bounded time-space cylindrical domain, see Milstein and Tretyakov [MT99]. Recently, Deaconu and Lejay [DL06] have developed similar algorithms, but based on random walks on rectangles. However, computationally speaking, our approach is presumably more direct.

1.4 Outline of the paper

Notations and assumptions used throughout the paper are stated in Section 1.5. In Section 2 we give our main results concerning the asymptotics of the overshoot, the error expansion and the boundary correction. These results are proved in Section 3, which is the technical core of the paper. Eventually, Section 4 deals with the stationary extension of our results. We still manage to obtain an expansion and a correction for elliptic PDEs. Some technical results are postponed to the Appendix.

1.5 General notation and assumptions

1.5.1 Miscellaneous

- *Differentiation.* For smooth functions $g(t,x)$, we denote by $\partial_x^\beta g(t,x)$ the

derivative of g w.r.t. x according to the multi-index β, whereas the time-derivative of g is denoted by $\partial_t g(t,x)$. The notation $\nabla g(t,x)$ stands for the usual gradient w.r.t. x (as a row vector) and the Hessian matrix of g (w.r.t. the space variable x) is denoted by $Hg(t,x)$.
The second order linear operator L_t below stands for the infinitesimal generator of the diffusion process X in (1.1) at time t :

$$L_t g(t,x) = \nabla g(t,x) b(t,x) + \frac{1}{2}\mathrm{Tr}(Hg(t,x)[\sigma\sigma^*](t,x)). \tag{1.5}$$

• *Metric.* The Euclidean norm is denoted by $|\cdot|$.
We set $B_d(x,\epsilon)$ for the usual Euclidean d-dimensional open ball with center x and radius ϵ and $\mathbf{d}(x,C)$ for the Euclidean distance of a point x to a closed set C. The r-neighborhood of C is denoted by $V_C(r) = \{x : \mathbf{d}(x,C) \leq r\}$ $(r \geq 0)$.
• *Functions.* For an open set $\mathcal{D}' \subset \mathbb{R} \times \mathbb{R}^d$ and $l \in \mathbf{N}$, $\mathcal{C}^{\lfloor \frac{l}{2} \rfloor, l}(\mathcal{D}')$ (resp. $\mathcal{C}^{\lfloor \frac{l}{2} \rfloor, l}(\overline{\mathcal{D}'})$) is the space of continuous functions f defined on $\mathcal{D}'$ with continuous derivatives $\partial_x^\beta \partial_t^j f$ for $|\beta| + 2j \leq l$ (resp. defined in a neighborhood of $\overline{\mathcal{D}'}$). Also, for $a = l + \theta,\ \theta \in]0,1],\ l \in \mathbf{N}$, we denote by $\mathbf{H}_a(\mathcal{D}')$ (resp. $\mathbf{H}_a(\bar{\mathcal{D}}')$) the Banach space of functions of $\mathcal{C}^{\lfloor \frac{l}{2} \rfloor, l}(\mathcal{D}')$ (resp. $\mathcal{C}^{\lfloor \frac{l}{2} \rfloor, l}(\bar{\mathcal{D}}')$) having l^{th} space derivatives uniformly θ-Hlder continuous and $\lfloor l/2 \rfloor$ time-derivatives uniformly $(a/2 - \lfloor l/2 \rfloor)$-Hlder continuous, see Lieberman [Lie96], p. 46 for details. We may simply write $\mathcal{C}^{\lfloor \frac{l}{2} \rfloor, l}$ or $\mathbf{H}_a$ when $\mathcal{D}' = \mathbb{R} \times \mathbb{R}^d$.
• *Floating constants.* As usual, we use the same symbol C for all finite, non-negative constants which appear in our computations : they may depend on $\mathcal{D}, T, b, \sigma, g, f, k$ but they will not depend on Δ or x. We reserve the notation c for constants also independent of T, g, f and k. Other possible dependences will be explicitly indicated.
In the following $O_{pol}(\Delta)$ (resp. $O(\Delta)$) stands for every quantity $R(\Delta)$ such that, for any $k \in \mathbf{N}$ one has $|R(\Delta)| \leq C_k \Delta^k$ (resp. $|R(\Delta)| \leq C\Delta$) for a constant $C_k > 0$ (uniformly in the starting point x).

1.5.2 Time-space domains

Below, we introduce some usual notations for such domains (see e.g. [Fri64], [Lie96]). In what follows, for any $t \geq 0$, D_t is a non empty bounded domain of $\mathbb{R}^d$, that coincides with the interior of its closure (see [Fri64], Section 3.2). We then define the time-space domain by $\mathcal{D} := \bigcup_{0<t<T} \{t\} \times D_t \subset]0,T[\times \mathbb{R}^d$, see Figure 1.

Regularity assumptions on the domain $\mathcal{D}$ will be formulated in terms of Hölder spaces with *time-space* variables (see [Lie96] p.46 and [Fri64] Section 3.2). Namely, we say that the domain $\mathcal{D}$ is of class $\mathbf{H}_a,\ a \geq 1$ if for every boundary point $(t_0, x_0) \in \bigcup_{0 \leq t \leq T} \{t\} \times \partial D_t$, there exists a neighborhood $]t_0 - \varepsilon_0^2, t_0 + \varepsilon_0^2[\times B_d(x_0, \varepsilon_0)$, an index $1 \leq i \leq d$ and a function $\varphi_0 \in \mathbf{H}_a(]t_0 - \varepsilon_0^2, t_0 +$

$\varepsilon_0^2[\times B_{d-1}((x_0^1, ..., x_0^{i-1}, x_0^{i+1}, ..., x_0^d), \varepsilon_0)$ s.t.

$$\begin{aligned}\Big\{ \cup_{0\le t\le T} \{t\} \times \partial D_t \Big\} \cap \Big\{]t_0 - \varepsilon_0^2, t_0 + \varepsilon_0^2[\times B_d(x_0, \varepsilon_0) \Big\} \\ := \{(t,x) \in (]t_0 - \varepsilon_0^2, t_0 + \varepsilon_0^2[\cap[0,T]) \times B_d(x_0, \varepsilon_0) : \\ x_i = \varphi_0(t, x_1, ..., x_{i-1}, x_{i+1}, ..., x_d)\}.\end{aligned}$$

If $\mathcal{D}$ is of class $\mathbf{H}_2$, all domains D_t, for $t \in [0,T]$, satisfy the uniform interior and exterior sphere condition with the same radius $r_0 > 0$. Moreover, the signed spatial distance F, given by

$$F(t,x) = \begin{cases} -\mathbf{d}(x, \partial D_t), & \text{for } x \in D_t^c,\ \mathbf{d}(x, \partial D_t) \le r_0,\ 0 \le t \le T, \\ \mathbf{d}(x, \partial D_t), & \text{for } x \in D_t,\ \mathbf{d}(x, \partial D_t) \le r_0,\ 0 \le t \le T, \end{cases}$$

belongs to $\mathbf{H}_2\left(\{(t,x) : 0 \le t \le T,\ \mathbf{d}(x, \partial D_t) < r_0\}\right)$ (see [Lie96], Section X.3) and $n(t,x) = [\nabla F]^*(t,x)$ is the unit inward normal vector to D_t at $\pi_{\partial D_t}(x)$ the nearest point to x in ∂D_t (see Figure 3). The function F can be extended as a $\mathbf{H}_2([0,T] \times \mathbb{R}^d)$ function, preserving the sign (see [Lie96], Section X.3).

1.5.3 Diffusion processes stopped at the boundary

We specify the properties of the coefficients (b, σ) in (1.1) with assumption

($\mathbf{A}_\theta$) (with $\theta \in]0,1]$)

1. *Smoothness.* The functions b and σ are in $\mathbf{H}_{1+\theta}$.

2. *Uniform ellipticity.* For some $a_0 > 0$, it holds $\xi^*[\sigma\sigma^*](t,x)\xi \ge a_0|\xi|^2$ for any $(t,x,\xi) \in [0,T] \times \mathbb{R}^d \times \mathbb{R}^d$.

We mention that the additional smoothness of b and σ w.r.t. the time variable is required for the connection with PDEs. We also introduce assumption **($\mathbf{A}_\theta^{'}$)** for which **2.** is replaced by the weaker assumption

2'. *Uniform non characteristic boundary.* For some $r_0 > 0$ there exists $a_0 > 0$ s.t. $\nabla F(t,x)[\sigma\sigma^*](t,x)\nabla F(t,x)^* \ge a_0$ for any $(t,x) \in \bigcup_{0\le t\le T}\{t\} \times V_{\partial D_t}(r_0)$.

The asymptotic results concerning the overshoot hold true under **($\mathbf{A}_\theta^{'}$)**, see Section 2.1. In the following we use the superscript t,x to indicate the usual Markovian dependence, i.e. $\forall s \ge t,\ X_s^{t,x} = x + \int_t^s b(u, X_u^{t,x})du + \int_t^s \sigma(u, X_u^{t,x})dW_u$. Now let

$$\tau^{t,x} := \inf\{s > t : X_s^{t,x} \notin D_s\} \tag{1.6}$$

be the first exit time of $X_s^{t,x}$ from D_s. For functionals of the process X stopped at the exit from $\mathcal{D}$, of the form

$$\begin{aligned} u(t,x) =& \mathbb{E}\Big[g(\tau^{t,x}\wedge T, X^{t,x}_{\tau^{t,x}\wedge T})e^{-\int_t^{\tau^{t,x}\wedge T} k(r,X_r^{t,x})dr} \\ &+ \int_t^{\tau^{t,x}\wedge T} e^{-\int_t^s k(r,X_r^{t,x})dr} f(s,X_s^{t,x})ds\Big], \end{aligned} \tag{1.7}$$

we now recall (see [CGK06]) that the Feynman-Kac representation holds in the time-space domain. Introduce the parabolic boundary $\mathcal{PD} = \partial\mathcal{D}\backslash[\{0\}\times D_0]$.

Proposition 1 [Feynman-Kac's formula and a priori estimates on u]

Assume $(\mathbf{A}_\theta)$, $\mathcal{D}\in\mathbf{H}_1$, $k\in\mathbf{H}_\theta$, $f\in\mathbf{H}_\theta$ *and* $g\in\mathcal{C}^{0,0}$ *with* $\theta\in]0,1[$. *Then, there is a unique solution in* $\mathcal{C}^{1,2}(\mathcal{D})\cap\mathcal{C}^{0,0}(\overline{\mathcal{D}})$ *to*

$$\begin{cases} \partial_t u + L_t u - ku + f = 0 & \text{in } \mathcal{D}, \\ u = g & \text{on } \mathcal{PD}, \end{cases} \tag{1.8}$$

and it is given by (1.7).
In addition, if for some $\theta\in]0,1[$, $\mathcal{D}$ *is of class* $\mathbf{H}_{1+\theta}$, $g\in\mathbf{H}_{1+\theta}$ *then* $u\in\mathbf{H}_{1+\theta}$. *In particular* ∇u *exists and is* θ*-Hlder continuous up to the boundary.*
Eventually, for $\mathcal{D}\in\mathbf{H}_{3+\theta}$, $k,f\in\mathbf{H}_{1+\theta}$, $g\in\mathbf{H}_{3+\theta}$ *satisfying the first order compatibility condition* $(\partial_t + L_T - k)g(T,x) + f(T,x)|_{x\in\partial\mathcal{D}_T} = 0$, *then the function* u *belongs to* $\mathbf{H}_{3+\theta}$.

Proof. The first two existence and uniqueness result for (1.8) are respectively implied by Theorems 5.9 and 5.10 and Theorem 6.45 in Lieberman, [Lie96]. The probabilistic representation is then a usual verification argument, see e.g. Appendix B.1 in [CGK06]. The additional smoothness can be derived from exercise 4.5 Chapter IV in [Lie96] or Theorem 12, Chapter 3 in [Fri64]. □

2 Main Results

2.1 Controls concerning the overshoot

The overshoot is the distance of the discretely killed process to the boundary, when it exits the domain by its side. To be precise, we use F the signed distance function and we consider the quantity $F(t_i, X^\Delta_{t_i})$. It remains positive for $t_i < \tau^\Delta$, and at time $t_i = \tau^\Delta$, it becomes non positive. Additionally, under the ellipticity assumption, the above inequality is strict: $F(\tau^\Delta, X^\Delta_{\tau^\Delta}) < 0$ a.s.. The overshoot is thus defined by $F^-(\tau^\Delta, X^\Delta_{\tau^\Delta})$. Also, since F is in $\mathbf{H}_2$

(and therefore Lipschitz continuous in time and space), it is easy to see that $F^-(\tau^\Delta, X^\Delta_{\tau^\Delta})$ is of order $\sqrt{\Delta}$ (in L_p-norm for instance). Thus, it is natural to study the asymptotics of the rescaled overshoot

$$\Delta^{-1/2} F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}).$$

Adapting the proof of Proposition 6 in [GM04] to our time-dependent context, see also the proof of Proposition 15 for a simpler version, one has the following proposition.

Proposition 2 (Tightness of the overshoot) *Assume* $(\mathbf{A}'_\theta)$*, and that* $\mathcal{D}$ *is of class* $\mathbf{H}_2$*. Then, for some* $c > 0$ *one has*

$$\sup_{\Delta>0, s\in[0,T]} \mathbb{E}_x[\exp(c[\Delta^{-1/2} F^-(s\wedge\tau^\Delta, X^\Delta_{s\wedge\tau^\Delta})]^2)] < +\infty.$$

It is quite plain to prove, by pathwise convergence of X^Δ towards X on compact sets, that $(\tau^\Delta \wedge T, X^\Delta_{\tau^\Delta\wedge T})$ converges in probability to $(\tau\wedge T, X_{\tau\wedge T})$. The next theorem also includes the rescaled overshoot.

Theorem 3 (Joint limit laws associated to the overshoot) *Assume* $(\mathbf{A}'_\theta)$*, and that* $\mathcal{D}$ *is of class* $\mathbf{H}_2$*. Let* φ *be a continuous function with compact support. For all* $t\in[0,T],\ x\in D_0,\ y\geq 0$*,*

$$\mathbb{E}_x[\mathbf{1}_{\tau^\Delta\leq t} Z^\Delta_{\tau^\Delta}\varphi(X^\Delta_{\tau^\Delta})\mathbf{1}_{F^-(\tau^\Delta, X^\Delta_{\tau^\Delta})\geq y\sqrt{\Delta}}] \underset{\Delta\to 0}{\longrightarrow}$$
$$\mathbb{E}_x\Big[\mathbf{1}_{\tau\leq t} Z_\tau\varphi(X_\tau)\Big(1 - H(y/|\nabla F\sigma(\tau, X_\tau)|)\Big)\Big]$$

with $H(y) := (\mathbb{E}_0[s_{\tau^+}])^{-1}\int_0^y \mathbb{P}_0[s_{\tau^+} > z]dz$ *and* $s_0 := 0, \forall n\geq 1, s_n := \sum_{i=1}^n G^i$*, the* G^i *being i.i.d. standard centered normal variables,* $\tau^+ := \inf\{n\geq 0 : s_n > 0\}$*.*

In other words, $(\tau^\Delta, X^\Delta_{\tau^\Delta}, \Delta^{-1/2}F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}))$ weakly converges to $(\tau, X_\tau, |\nabla F\sigma(\tau, X_\tau)|Y)$ where Y is a random variable independent of (τ, X_τ), and which cumulative function is equal to H. Actually, Y has the asymptotic law of the renormalized Brownian overshoot. In the following analysis, the mean of the overshoot is an important quantity and it is worth noting that one has $\mathbb{E}(Y) = \frac{\mathbb{E}_0[s^2_{\tau^+}]}{2\mathbb{E}_0[s_{\tau^+}]} := c_0$. One knows from [Sie79] that

$$c_0 = -\frac{\zeta(1/2)}{\sqrt{2\pi}} = 0.5826... \tag{2.1}$$

The above theorem is the crucial tool in the derivation of our main results. The proof is given in Section 3.1.

2.2 Error expansion and boundary correction

For notational convenience introduce for $x \in D_0$,

$$u(\mathcal{D}) = \mathbb{E}_x(g(\tau \wedge T, X_{\tau\wedge T})Z_{\tau\wedge T} + \int_0^{\tau\wedge T} Z_s f(s, X_s)ds),$$
$$u^\Delta(\mathcal{D}) = \mathbb{E}_x(g(\tau^\Delta \wedge T, X^\Delta_{\tau^\Delta\wedge T})Z^\Delta_{\tau^\Delta\wedge T} + \int_0^{\tau^\Delta\wedge T} Z^\Delta_{\phi(s)} f(\phi(s), X^\Delta_{\phi(s)})ds).$$

Theorem 4 (First order expansion) *Under* $(\mathbf{A}_\theta)$*, for a domain of class* $\mathbf{H}_2$*,* $g \in \mathbf{H}_{1+\theta}$*,* $k, f \in \mathbf{H}_{1+\theta}$ *and for* Δ *small enough*

$$\begin{aligned}&\mathrm{Err}(T, \Delta, g, f, k, x) = u^\Delta(\mathcal{D}) - u(\mathcal{D})\\ &= c_0\sqrt{\Delta}\mathbb{E}_x(\mathbf{1}_{\tau\le T} Z_\tau(\nabla u - \nabla g)(\tau, X_\tau)\cdot\nabla F(\tau, X_\tau)|\nabla F\sigma(\tau, X_\tau)|) + o(\sqrt{\Delta}),\end{aligned}$$

where c_0 *is defined in* (2.1)*.*

Define now a smaller domain $\mathcal{D}^\Delta \subset \mathcal{D}$, which time-section is given by $D^\Delta_t = \{x \in D_t : \mathbf{d}(x, \partial D_t) > c_0\sqrt{\Delta}|\nabla F\sigma(t, x)|\}$, see Figure 2. Introduce the exit time of the Euler scheme from this smaller domain: $\hat\tau^\Delta = \inf\{t_i > 0 : X^\Delta_{t_i} \notin D^\Delta_{t_i}\} \le \tau^\Delta$. The boundary correction procedure consists in simulating

$$g(\hat\tau^\Delta \wedge T, X^\Delta_{\hat\tau^\Delta\wedge T})Z^\Delta_{\hat\tau^\Delta\wedge T} + \int_0^{\hat\tau^\Delta\wedge T} Z^\Delta_{\phi(s)} f(\phi(s), X^\Delta_{\phi(s)})ds. \tag{2.2}$$

As above, we do not compute any projection on the boundary. We denote the expectation of (2.2) by $u^\Delta(\mathcal{D}^\Delta)$. One has:

Theorem 5 (Boundary correction) *Under the assumptions of Theorem 4, if we additionally suppose* $\nabla F(.,.)|\nabla F\sigma(.,.)|$ *is in* $\mathcal{C}^{1,2}$*, then one has:*

$$u^\Delta(\mathcal{D}^\Delta) - u(\mathcal{D}) = o(\sqrt{\Delta}).$$

The additional assumption is due to technical considerations to ensure that the modified domain $\mathcal{D}^\Delta$ is also of class $\mathbf{H}_2$. It is automatically fulfilled for domains of class $\mathcal{C}^3$ and σ in $\mathcal{C}^{1,2}$.

2.3 Proof of Theorems 4 and 5

2.3.1 Error expansion

By usual weak convergence arguments, Theorem 4 is a direct consequence of Proposition 2 (tightness), Theorem 3 (joint limit laws associated to the

overshoot) and Theorem 6 below.

Theorem 6 (First order approximation) *Under the assumptions of Theorem 4, one has*

$$u^\Delta(\mathcal{D}) - u(\mathcal{D}) = o(\sqrt{\Delta}) + \\ \mathbb{E}_x(\mathbf{1}_{\tau^\Delta \le T} Z^\Delta_{\tau^\Delta}(\nabla u - \nabla g)(\tau^\Delta, \pi_{\partial D_{\tau^\Delta}}(X^\Delta_{\tau^\Delta})) \cdot \nabla F(\tau^\Delta, X^\Delta_{\tau^\Delta}) F^-(\tau^\Delta, X^\Delta_{\tau^\Delta})).$$

Remark 7 *In the above statement, we use projections on a non convex set, which needs a clarification. With the notation of Section 1.5.2, introduce* $\tau^{r_0} := \inf\{s > 0 : X^\Delta_s \notin V_{D_s}(r_0)\}$. *For* $s \in [0, \tau^{r_0}]$ *the projection* $\pi_{\bar{D}_s}(X^\Delta_s)$ *is uniquely defined by*

$$\pi_{\bar{D}_s}(X^\Delta_s) = X^\Delta_s + (\nabla F)^*(s, X^\Delta_s) F^-(s, X^\Delta_s), \tag{2.3}$$

see Figure 3. Large deviation arguments (see Lemma 8 below) also give $\mathbb{P}_x[\tau^{r_0} \le \tau^\Delta \le T] = O_{pol}(\Delta)$. *Thus, in the following, for* $s \ge \tau^{r_0}$, $\pi_{\bar{D}_s}(X^\Delta_s)$ *and* $\pi_{\partial D_s}(X^\Delta_s)$ *denote an arbitrary point on* ∂D_s. *This choice yields an exponentially small contribution in our estimates.*



Figure 3. Orthogonal projection $\pi_{\partial D_t}(x)$ of $x \notin D_t$ onto the boundary ∂D_t and the related signed distance $F(t,x)$. Here $F(t,x) < 0$ and $\mathbf{d}(x, \partial D_t) = |F(t,x)| = F^-(t,x)$.

Proof. Denote $e^\Delta := u^\Delta(\mathcal{D}) - u(\mathcal{D})$ the above error. Write now

$$\begin{aligned} e^\Delta =& \mathbb{E}_x[g(\tau^\Delta \wedge T, X^\Delta_{\tau^\Delta \wedge T}) Z^\Delta_{\tau^\Delta \wedge T} - g(\tau^\Delta \wedge T, \pi_{\bar{D}_{\tau^\Delta \wedge T}}(X^\Delta_{\tau^\Delta \wedge T})) Z^\Delta_{\tau^\Delta \wedge T}] \\ &+ \Big\{\mathbb{E}_x[g(\tau^\Delta \wedge T, \pi_{\bar{D}_{\tau^\Delta \wedge T}}(X^\Delta_{\tau^\Delta \wedge T})) Z^\Delta_{\tau^\Delta \wedge T} + \int_0^{\tau^\Delta \wedge T} Z^\Delta_{\phi(s)} f(\phi(s), X^\Delta_{\phi(s)}) ds] \\ &\quad - u(0, X^\Delta_0)\Big\} \\ :=& e^\Delta_1 + e^\Delta_2. \end{aligned}$$

We introduce here the projection for the error analysis. From (2.3) and Proposition 2, a Taylor expansion yields

$$e_1^\Delta = -\mathbb{E}_x[\mathbf{1}_{\tau^\Delta \le T} Z^\Delta_{\tau^\Delta} \nabla g(\tau^\Delta, \pi_{\partial D_{\tau^\Delta}}(X^\Delta_{\tau^\Delta})) \cdot \nabla F(\tau^\Delta, X^\Delta_{\tau^\Delta}) F^-(\tau^\Delta, X^\Delta_{\tau^\Delta})] + O(\Delta^{(1+\theta)/2}). \tag{2.4}$$

In the following, we write $U \overset{\mathbb{E}}{=} V$ (resp $U \overset{\mathbb{E}}{\le} V$) when the equality between U and V holds in mean up to a $O_{pol}(\Delta)$ (resp. $\mathbb{E}_x(U) \le \mathbb{E}_x(V) + O_{pol}(\Delta)$). We also use the notation $U = O(V)$ between two random variables U and V if for a constant C, one has $|U| \le C|V|$. Because $g(\tau^\Delta \wedge T, \pi_{\bar{D}_{\tau^\Delta \wedge T}}(X^\Delta_{\tau^\Delta \wedge T})) = u(\tau^\Delta \wedge T, \pi_{\bar{D}_{\tau^\Delta \wedge T}}(X^\Delta_{\tau^\Delta \wedge T}))$, we can write a telescopic summation:

$$\begin{aligned} e_2^\Delta &\overset{\mathbb{E}}{=} \Big(\sum_{0 \le t_i < \tau^\Delta \wedge T} u(t_{i+1}, \pi_{\bar{D}_{t_{i+1}}}(X^\Delta_{t_{i+1}})) Z^\Delta_{t_{i+1}} \\ &\qquad - u(t_i, \pi_{\bar{D}_{t_i}}(X^\Delta_{t_i})) Z^\Delta_{t_i} + Z^\Delta_{t_i} f(t_i, X^\Delta_{t_i}) \Delta \Big) \mathbf{1}_{\tau^{r_0} > \tau^\Delta \wedge T} \\ &\overset{\mathbb{E}}{=} \Big(\sum_{0 \le t_i < T} \mathbf{1}_{t_i < \tau^\Delta} \Big[u(t_{i+1}, \pi_{\bar{D}_{t_{i+1}}}(X^\Delta_{t_{i+1}})) Z^\Delta_{t_{i+1}} \\ &\qquad - u(t_i, X^\Delta_{t_i}) Z^\Delta_{t_i} + Z^\Delta_{t_i} f(t_i, X^\Delta_{t_i}) \Delta \Big] \Big) \mathbf{1}_{\tau^{r_0} > \tau^\Delta \wedge T} \end{aligned}$$

since for $t_i < \tau^\Delta$, $X^\Delta_{t_i} \in D_{t_i}$ and thus $\pi_{\bar{D}_{t_i}}(X^\Delta_{t_i}) = X^\Delta_{t_i}$. To proceed, the key idea is to introduce on the event $\{t_i < \tau^\Delta\}$, the partition $\{F(t_i, X^\Delta_{t_i}) \in (0, 2\Delta^{\frac{1}{2}(1-\varepsilon)}]\} \cup \{F(t_i, X^\Delta_{t_i}) > 2\Delta^{\frac{1}{2}(1-\varepsilon)}\} := A^\varepsilon_{t_i} \cup (A^\varepsilon_{t_i})^C$, $\varepsilon > 0$. This allows to split the cases for which $X^\Delta_{t_i}$ is close or not to the boundary ∂D_{t_i}. Lemma 8 ensures that $(X^\Delta_s)_{s \in [t_i, t_{i+1}]}$ stayed in $B(X^\Delta_{t_i}, \Delta^{\frac{1}{2}(1-\varepsilon)})$ with a probability exponentially close to one. Then, on $(A^\varepsilon_{t_i})^C$, the smoothness of the domain yields $\mathbf{1}_{(A^\varepsilon_{t_i})^C} \mathbb{P}[X^\Delta_{t_{i+1}} \in D_{t_{i+1}} | \mathcal{F}_{t_i}] = 1 - O(\exp(-c\Delta^{-\varepsilon}))$, see Proposition 19 for a proof of this claim. On the other hand, on $A^\varepsilon_{t_i}$, $X^\Delta_{t_i}$ is sufficiently close to the boundary to make the contribution of the overshoot at time t_{i+1} significant for the error analysis. Write:

$$\begin{aligned} e_2^\Delta &\overset{\mathbb{E}}{=} \Big(\sum_{0 \le t_i < T} \mathbf{1}_{t_i < \tau^\Delta} \Big\{ \mathbf{1}_{A^\varepsilon_{t_i}} \Big[u(t_{i+1}, \pi_{\bar{D}_{t_{i+1}}}(X^\Delta_{t_{i+1}})) Z^\Delta_{t_{i+1}} \\ &\qquad - u(t_i, X^\Delta_{t_i}) Z^\Delta_{t_i} + Z^\Delta_{t_i} f(t_i, X^\Delta_{t_i}) \Delta \Big] \\ &\quad + \mathbf{1}_{(A^\varepsilon_{t_i})^C} \mathbf{1}_{\forall s \in [t_i, t_{i+1}],\ X^\Delta_s \in B(X^\Delta_{t_i}, \Delta^{\frac{1}{2}(1-\varepsilon)})} \Big[u(t_{i+1}, X^\Delta_{t_{i+1}}) Z^\Delta_{t_{i+1}} \\ &\qquad - u(t_i, X^\Delta_{t_i}) Z^\Delta_{t_i} + Z^\Delta_{t_i} f(t_i, X^\Delta_{t_i}) \Delta \Big] \Big\} \Big) \mathbf{1}_{\tau^{r_0} > \tau^\Delta \wedge T} := e_{21}^\Delta + e_{22}^\Delta. \end{aligned} \tag{2.5}$$

Let us first deal with e_{21}^Δ. In our framework, u is $(1+\theta)/2$-Hölder continuous in time and ∇u is θ-Hölder continuous in space on a neighborhood of $\mathcal{D}$. A

Taylor expansion at order one and the equality (2.3) give

$$
\begin{aligned}
e_{21}^{\Delta} &\overset{\mathbb{E}}{=} \Big(\sum_{0\le t_i<T} \mathbf{1}_{t_i<\tau^\Delta} \mathbf{1}_{A_{t_i}^{\varepsilon}} \Big[Z_{t_i}^{\Delta} \nabla u(t_i, X_{t_i}^{\Delta}) \cdot \nabla F(t_{i+1}, X_{t_{i+1}}^{\Delta}) F^{-}(t_{i+1}, X_{t_{i+1}}^{\Delta}) \\
&\quad + O(|F^{-}(t_{i+1}, X_{t_{i+1}}^{\Delta})|^{1+\theta}) + O(|X_{t_{i+1}}^{\Delta} - X_{t_i}^{\Delta}|^{1+\theta}) + O(\Delta^{\frac{1+\theta}{2}}) \Big] \Big) \mathbf{1}_{\tau^{r_0}>\tau^\Delta\wedge T} \\
&\overset{\mathbb{E}}{=} \Big(\mathbf{1}_{\tau^\Delta\le T} Z_{\tau^\Delta}^{\Delta} \nabla u(\tau^\Delta, X_{\tau^\Delta}^{\Delta}) \cdot \nabla F(\tau^\Delta, X_{\tau^\Delta}^{\Delta}) F^{-}(\tau^\Delta, X_{\tau^\Delta}^{\Delta}) \\
&\quad + \sum_{0\le t_i<T} \mathbf{1}_{t_i<\tau^\Delta} \mathbf{1}_{A_{t_i}^{\varepsilon}} \Big[O(|F^{-}(t_{i+1}, X_{t_{i+1}}^{\Delta})|^{1+\theta}) + O(|X_{t_{i+1}}^{\Delta} - X_{t_i}^{\Delta}|^{1+\theta}) \\
&\qquad + O(|X_{t_{i+1}}^{\Delta} - X_{t_i}^{\Delta}|^{\theta} F^{-}(t_{i+1}, X_{t_{i+1}}^{\Delta})) + O(\Delta^{\frac{1+\theta}{2}}) \Big] \Big) \mathbf{1}_{\tau^{r_0}>\tau^\Delta\wedge T}
\end{aligned}
$$

where we used once again Lemma 8 for the last equality. Standard arguments yield $\mathbb{E}[|X_{t_{i+1}}^{\Delta} - X_{t_i}^{\Delta}|^p | \mathcal{F}_{t_i}] = O(\Delta^{\frac{p}{2}})$ for any $p > 0$ and $\mathbb{E}[|F^{-}(t_{i+1}, X_{t_{i+1}}^{\Delta})|^p | \mathcal{F}_{t_i}] = \mathbb{E}[|F^{-}(t_{i+1}, X_{t_{i+1}}^{\Delta}) - F^{-}(t_i, X_{t_i}^{\Delta})|^p | \mathcal{F}_{t_i}] = O(\Delta^{\frac{p}{2}})$ on $\{t_i < \tau^\Delta\}$. Thus, we can now rewrite

$$
\begin{aligned}
e_{21}^{\Delta} &\overset{\mathbb{E}}{=} \Big(\mathbf{1}_{\tau^\Delta\le T} Z_{\tau^\Delta}^{\Delta} \nabla u(\tau^\Delta, \pi_{\partial D_{\tau^\Delta}}(X_{\tau^\Delta}^{\Delta})) \cdot \nabla F(\tau^\Delta, X_{\tau^\Delta}^{\Delta}) F^{-}(\tau^\Delta, X_{\tau^\Delta}^{\Delta}) \Big) \mathbf{1}_{\tau^{r_0}>\tau^\Delta\wedge T} \\
&\quad + e_{211}^{\Delta}, \\
e_{211}^{\Delta} &\overset{\mathbb{E}}{=} \Big(\sum_{0\le t_i<T} \mathbf{1}_{t_i<\tau^\Delta} \mathbf{1}_{A_{t_i}^{\varepsilon}} O(\Delta^{\frac{1+\theta}{2}}) \Big) \mathbf{1}_{\tau^{r_0}>\tau^\Delta\wedge T}.
\end{aligned}
$$

To handle e_{211}^{Δ} the idea is to use the occupation time formula and some sharp estimates concerning the local time of $(F(s, X_s^{\Delta}))_{s\le T\wedge\tau^\Delta}$ in a neighborhood of the boundary. We have

$$
\begin{aligned}
|e_{211}^{\Delta}| &\overset{\mathbb{E}}{\le} C\Delta^{\frac{1+\theta}{2}} \left(\Delta^{-1} \int_0^{T\wedge\tau^\Delta} \mathbf{1}_{F(\phi(t), X_{\phi(t)}^{\Delta}) \in [0, 2\Delta^{1/2(1-\varepsilon)}]} dt \right) \mathbf{1}_{\tau^{r_0}>\tau^\Delta\wedge T} \\
&\overset{\mathbb{E}}{\le} C\Delta^{\frac{1+\theta}{2}} \left(\Delta^{-1} \int_0^{T\wedge\tau^\Delta} \mathbf{1}_{F(t, X_t^{\Delta}) \in [-\Delta^{1/2(1-\varepsilon)}, 3\Delta^{1/2(1-\varepsilon)}]} dt \right) \mathbf{1}_{\tau^{r_0}>\tau^\Delta\wedge T} \\
&\overset{\mathbb{E}}{\le} C\Delta^{\frac{1+\theta}{2}} \left(\Delta^{-1} \int_{-\Delta^{1/2(1-\varepsilon)}}^{3\Delta^{1/2(1-\varepsilon)}} L_{T\wedge\tau^\Delta}^{y}(F(., X_{.}^{\Delta})) dy \right) \mathbf{1}_{\tau^{r_0}>\tau^\Delta\wedge T},
\end{aligned}
$$

where we have used Lemma 8 at the second equality and the uniform ellipticity assumption for the last one. Now an easy adaptation of the proof of Lemma 17 [GM04] to our time-dependent domain framework gives

$$
\mathbb{E}[L_{T\wedge\tau^\Delta}^{y}(F(., X_{.}^{\Delta}))] \le C(|y| + \Delta^{\frac{1}{2}}). \tag{2.6}
$$

Thus, one has $|e_{211}^{\Delta}| \overset{\mathbb{E}}{\le} C\Delta^{\frac{1+\theta}{2} - \frac{\varepsilon}{2}} = o(\Delta^{\frac{1}{2}})$ for ε small enough. Hence, the above estimates and Lemma 8 give

$$
e_{21}^{\Delta} \overset{\mathbb{E}}{=} \Big(\mathbf{1}_{\tau^\Delta\le T} Z_{\tau^\Delta}^{\Delta} \nabla u(\tau^\Delta, \pi_{\partial D_{\tau^\Delta}}(X_{\tau^\Delta}^{\Delta})) \cdot \nabla F(\tau^\Delta, X_{\tau^\Delta}^{\Delta}) F^{-}(\tau^\Delta, X_{\tau^\Delta}^{\Delta}) \Big) + o(\Delta^{\frac{1}{2}}). \tag{2.7}
$$

Let us now turn to e_{22}^{Δ}. If $g \in \mathbf{H}_{3+\theta}$ (which implies $u \in \mathbf{H}_{3+\theta}$ in view of Proposition 1), the term e_{22}^{Δ} can be handled with somehow standard techniques. Namely Taylor like expansions in the spirit of Talay and Tubaro [TT90]. For simplicity we handle e_{22}^{Δ} under the previous smoothness assumption on g and u. The proof under weaker assumptions ($g \in \mathbf{H}_{1+\theta}$), that involves sharp estimates on possibly exploding derivatives of u near the boundary, is postponed to the Appendix. We recall that

$$e_{22}^{\Delta} \stackrel{\mathbb{E}}{=} \Big(\sum_{0 \le t_i < T} \mathbf{1}_{t_i < \tau^{\Delta}} \mathbf{1}_{(A_{t_i}^{\varepsilon})^C} \mathbf{1}_{\forall s \in [t_i, t_{i+1}],\ X_s^{\Delta} \in B(X_{t_i}^{\Delta}, \Delta^{\frac{1}{2}(1-\varepsilon)})} \Big[u(t_{i+1}, X_{t_{i+1}}^{\Delta}) Z_{t_{i+1}}^{\Delta} - u(t_i, X_{t_i}^{\Delta}) Z_{t_i}^{\Delta} + Z_{t_i}^{\Delta} f(t_i, X_{t_i}^{\Delta}) \Delta \Big] \Big) \mathbf{1}_{\tau^{r_0} > \tau^{\Delta} \wedge T}$$

For all $(s, y) \in \mathcal{D}$ introduce the operators $L_{s,y} : \mathcal{C}^{1,2}(\mathcal{D}) \to \mathcal{C}(\mathcal{D})$, $\varphi \mapsto ((t, x) \mapsto L_{s,y}\varphi(t, x) = \nabla\varphi(t, x) b(s, y) + \frac{1}{2}\mathrm{Tr}[H\varphi(t, x)[\sigma\sigma^*](s, y)])$. Recalling that $\partial_t u(t_i, X_{t_i}^{\Delta}) + L_{t_i, X_{t_i}^{\Delta}} u(t_i, X_{t_i}^{\Delta}) - k u(t_i, X_{t_i}^{\Delta}) + f(t_i, X_{t_i}^{\Delta}) = 0$, Itô's formula gives

$$\begin{aligned} e_{22}^{\Delta} \stackrel{\mathbb{E}}{=} \Big(\sum_{0 \le t_i < T} & \mathbf{1}_{t_i < \tau^{\Delta}} \mathbf{1}_{(A_{t_i}^{\varepsilon})^C} \mathbf{1}_{\forall s \in [t_i, t_{i+1}],\ X_s^{\Delta} \in B(X_{t_i}^{\Delta}, \Delta^{\frac{1}{2}(1-\varepsilon)})} \Big[\\ & \int_{t_i}^{t_{i+1}} (Z_s^{\Delta} - Z_{t_i}^{\Delta})(\partial_s + L_{t_i, X_{t_i}^{\Delta}} - k(t_i, X_{t_i}^{\Delta})) u(s, X_s^{\Delta}) ds \\ & + Z_{t_i}^{\Delta} \int_{t_i}^{t_{i+1}} \Big[\Big(\partial_s + L_{t_i, X_{t_i}^{\Delta}} - k(t_i, X_{t_i}^{\Delta}) \Big) u(s, X_s^{\Delta}) \\ & \qquad - \Big(\partial_s + L_{t_i, X_{t_i}^{\Delta}} - k(t_i, X_{t_i}^{\Delta}) \Big) u(t_i, X_{t_i}^{\Delta})) \Big] ds \\ & + M_{t_i, t_{i+1}} \Big] \Big) \mathbf{1}_{\tau^{r_0} > \tau^{\Delta} \wedge T}, \end{aligned} \tag{2.8}$$

where for all $v \in [t_i, t_{i+1}]$, $M_{t_i, v} := \int_{t_i}^{v} Z_s^{\Delta} \nabla u(s, X_s^{\Delta}) \sigma(t_i, X_{t_i}^{\Delta}) dW_s$ is a square-integrable martingale term. Note that in this definition, in whole generality, $M_{t_i, v}$ is not stopped at the exit time $\tau_{t_i} := \inf\{s \ge t_i : X_s^{\Delta} \notin D_s\}$. If $\tau_{t_i} \le t_{i+1}$ (which happens with exponentially small probability on $(A_{t_i}^{\varepsilon})^C$), the term $\nabla u(s, X_s^{\Delta})$, $s \in [\tau_{t_i}, t_{i+1}]$ in $M_{t_i, t_{i+1}}$ has to be understood as the smooth extension of ∇u to the whole space. In particular this extension remains bounded. Now, we derive from Lemma 8

$$\begin{aligned} \mathbb{E}_x \Big[\Big(\sum_{0 \le t_i < T} & \mathbf{1}_{t_i < \tau^{\Delta}} \mathbf{1}_{(A_{t_i}^{\varepsilon})^C} \mathbf{1}_{\forall s \in [t_i, t_{i+1}],\ X_s^{\Delta} \in B(X_{t_i}^{\Delta}, \Delta^{\frac{1}{2}(1-\varepsilon)})} M_{t_i, t_{i+1}} \Big] \Big) \mathbf{1}_{\tau^{r_0} > \tau^{\Delta} \wedge T} \Big] \\ & = \mathbb{E}_x \Big[\sum_{0 \le t_i < T} \mathbf{1}_{t_i < \tau^{\Delta}} \mathbf{1}_{(A_{t_i}^{\varepsilon})^C} M_{t_i, t_{i+1}} \Big] + O_{pol}(\Delta) = O_{pol}(\Delta). \end{aligned}$$

We can thus neglect the contribution of the martingale terms in (2.8). We now develop the other quantities in (2.8) with Taylor integral formulas to derive

$$
\begin{aligned}
&\int_{t_i}^{t_{i+1}} (Z_s^\Delta - Z_{t_i}^\Delta)(\partial_s + L_{t_i, X_{t_i}^\Delta} - k(t_i, X_{t_i}^\Delta))u(s, X_s^\Delta)ds \\
=&O\Big(\Delta^2(|u|_\infty + |\nabla u|_\infty + |\partial_t u|_\infty + |D^2 u|_\infty)\Big), \\
&\int_{t_i}^{t_{i+1}} (\partial_t u(s, X_s^\Delta) - \partial_t u(t_i, X_{t_i}^\Delta))ds \\
=&\int_{t_i}^{t_{i+1}} \nabla\partial_t u(t_i, X_{t_i}^\Delta)\sigma(t_i, X_{t_i}^\Delta)(W_s - W_{t_i})ds \\
&+ O\Big(\Delta^{1+\frac{1+\theta}{2}}[\partial_t u]_{t,\frac{1+\theta}{2}} + \Delta^2|\nabla\partial_t u|_\infty + \Delta \sup_{s\in[t_i,t_{i+1}]} |X_s^\Delta - X_{t_i}^\Delta|^{1+\theta}[\nabla\partial_t u]_{x,\theta}\Big),
\end{aligned}
$$

$$
\begin{aligned}
&\int_{t_i}^{t_{i+1}} (L_{t_i, X_{t_i}^\Delta} u(s, X_s^\Delta) - L_{t_i, X_{t_i}^\Delta} u(t_i, X_{t_i}^\Delta))ds \\
=&\int_{t_i}^{t_{i+1}} \langle H_u(t_i, X_{t_i}^\Delta)\sigma(t_i, X_{t_i}^\Delta)(W_s - W_{t_i}), b(t_i, X_{t_i}^\Delta)\rangle ds \\
&+ \frac{1}{2}\int_{t_i}^{t_{i+1}} \mathrm{Tr}\Big((D^3 u(t_i, X_{t_i}^\Delta)\sigma(t_i, X_{t_i}^\Delta)(W_s - W_{t_i})) \cdot a(t_i, X_{t_i}^\Delta)\Big)ds \\
&+ O\Big(\Delta^2\{|D^2 u|_\infty + |D^3 u|_\infty + |\partial_t\nabla u|_\infty\} + \Delta^{1+\frac{1+\theta}{2}}[D^2 u]_{t,\frac{1+\theta}{2}} \\
&+ \Delta|D^3 u|_\infty \sup_{s\in[t_i,t_{i+1}]} |X_s^\Delta - X_{t_i}^\Delta|^2 + \Delta \sup_{s\in[t_i,t_{i+1}]} |X_s^\Delta - X_{t_i}^\Delta|^{1+\theta}[D^3 u]_{x,\theta}\Big), \\
&k(t_i, X_{t_i}^\Delta)\int_{t_i}^{t_{i+1}} (u(s, X_s^\Delta) - u(t_i, X_{t_i}^\Delta))ds \\
=&k(t_i, X_{t_i}^\Delta)\int_{t_i}^{t_{i+1}} \nabla u(t_i, X_{t_i}^\Delta)\sigma(t_i, X_{t_i}^\Delta)(W_s - W_{t_i})ds \\
&+ O\Big(\Delta^2(|\partial_t u|_\infty + |\nabla u|_\infty) + \Delta|D^2 u|_\infty \sup_{s\in[t_i,t_{i+1}]} |X_s^\Delta - X_{t_i}^\Delta|^2\Big), \qquad (2.9)
\end{aligned}
$$

where $[\cdot]_{t,\alpha}, [\cdot]_{x,\alpha}, \alpha \in (0,1]$ denote respectively the Hölder norms of order α in time and space (see Chapter IV Section 1 p. 46 in [Lie96] for a precise definition).

Hence, bringing together our estimates and exploiting the relations between the spatial and time derivatives for u (through the PDE), from (2.8) and (2.9)

we derive

$$
\begin{aligned}
e_{22}^{\Delta} \overset{\mathbb{E}}{=} \Big(\sum_{0\le t_i<T} & \mathbf{1}_{t_i<\tau^\Delta} \mathbf{1}_{(A_{t_i}^{\varepsilon})^C} \mathbf{1}_{\forall s\in[t_i,t_{i+1}],\ X_s^\Delta\in B(X_{t_i}^\Delta,\Delta^{\frac12(1-\varepsilon)})} \Big[\\
& O\Big(\Delta^2\{1+|u|_\infty+|\nabla u|_\infty+|D^2u|_\infty+|D^3u|_\infty\}\Big) \\
& + O\Big(\Delta^{1+\frac{1+\theta}{2}}\{1+|u|_\infty+|\nabla u|_\infty+|D^2u|_\infty+|D^3u|_\infty+[D^2u]_{t,\frac{1+\theta}{2}}\}\Big) \\
& + O\Big(\Delta \sup_{s\in[t_i,t_{i+1}]} |X_s^\Delta - X_{t_i}^\Delta|^{1+\theta}\{1+|u|_\infty+|\nabla u|_\infty+|D^2u|_\infty+|D^3u|_\infty+[D^3u]_{x,\theta}\}\Big) \\
& + O\Big(\Delta \sup_{s\in[t_i,t_{i+1}]} |X_s^\Delta - X_{t_i}^\Delta|^{2}\{|D^2u|_\infty+|D^3u|_\infty\}\Big) \\
& + \bar{M}_{t_i,t_{i+1}}\Big]\Big) \mathbf{1}_{\tau^{r_0}>\tau^\Delta\wedge T},
\end{aligned}
\tag{2.10}
$$

where $\bar{M}_{t_i,t_{i+1}}$ denotes the sum of the terms involving the Brownian increment $(W_s - W_{t_i})_{s\in[t_i,t_{i+1}]}$ in the above equations (2.9). Under our current assumption, i.e. $u \in \mathbf{H}_{3+\theta}$, all the norms appearing in (2.10) and all the derivatives appearing in the $(\bar{M}_{t_i,t_{i+1}})_{0\le t_i<T}$ are bounded. Hence,

$$
\begin{aligned}
& \mathbf{1}_{t_i<\tau^\Delta} \mathbf{1}_{(A_{t_i}^{\varepsilon})^C} \mathbb{E}[\mathbf{1}_{\forall s\in[t_i,t_{i+1}],\ X_s^\Delta\in B(X_{t_i}^\Delta,\Delta^{\frac12(1-\varepsilon)})} \bar{M}_{t_i,t_{i+1}} \mathbf{1}_{\tau^{r_0}>\tau^\Delta\wedge T} | \mathcal{F}_{t_i}] \\
& \quad = \mathbf{1}_{t_i<\tau^\Delta} \mathbf{1}_{(A_{t_i}^{\varepsilon})^C} \mathbb{E}[\bar{M}_{t_i,t_{i+1}} | \mathcal{F}_{t_i}] + O_{pol}(\Delta) = O_{pol}(\Delta),
\end{aligned}
\tag{2.11}
$$

$$
|e_{22}^{\Delta}| \overset{\mathbb{E}}{\le} C\Delta^{\frac{1+\theta}{2}}. \tag{2.12}
$$

Plug (2.7) and (2.12) into (2.5). The statement is derived from (2.4) and (2.5). We specify in the Appendix how to complete the proof from a sharper version of (2.10) deriving from (2.8), when $g \in \mathbf{H}_{1+\theta}$. □

2.3.2 Boundary Correction

One has

$$
u^\Delta(\mathcal{D}^\Delta) - u(\mathcal{D}) = [u^\Delta(\mathcal{D}^\Delta) - u(\mathcal{D}^\Delta)] + [u(\mathcal{D}^\Delta) - u(\mathcal{D})]. \tag{2.13}
$$

(1) The first contribution in (2.13) has been previously analysed in Theorem 4, except that the domain $\mathcal{D}^\Delta$ depends on Δ. We can show that it is equal to $c_0\sqrt{\Delta}\mathbb{E}(\mathbf{1}_{\tau\le T} Z_\tau (\nabla u - \nabla g)(\tau, X_\tau)\cdot\nabla F(\tau, X_\tau)|\nabla F\sigma(\tau, X_\tau)|) + o(\sqrt{\Delta})$. We briefly sketch the proof of this assertion, which is done in two steps. For this, set $\hat{u}^\Delta = u(\mathcal{D}^\Delta)$ for the solution of the PDE in the domain $\mathcal{D}^\Delta$.

- **Step 1.** It is well known that all PDE estimates depend only on bounds on the derivatives of the level set functions (φ_0) arising in the definition of the time-dependent domains (see section 1.5.2), and on the bounds on the derivatives of data g, f and k. Hence, since $\mathcal{D}^\Delta$ is a small perturbation of class $\mathbf{H}_2$ (because $\nabla F|\nabla F\sigma|$ has this regularity) of the domain $\mathcal{D}$ of class $\mathbf{H}_2$, all PDE estimates on $\hat{u}^\Delta$ remain locally uniform w.r.t. Δ. In addition, $\hat{u}^\Delta$ and its gradient converge uniformly to

u and ∇u. This argumentation allows us to state that the first order approximation theorem holds:

$$u^\Delta(\mathcal{D}^\Delta) - u(\mathcal{D}^\Delta) = o(\sqrt{\Delta}) +$$
$$\mathbb{E}_x(\mathbf{1}_{\hat{\tau}^\Delta \le T} Z^\Delta_{\hat{\tau}^\Delta}(\nabla u - \nabla g)(\hat{\tau}^\Delta, \pi_{\partial D^\Delta_{\hat{\tau}^\Delta}}(X^\Delta_{\hat{\tau}^\Delta})) \cdot \nabla \hat{F}^\Delta(\hat{\tau}^\Delta, X^\Delta_{\hat{\tau}^\Delta})[\hat{F}^\Delta]^-(\hat{\tau}^\Delta, X^\Delta_{\hat{\tau}^\Delta})),$$

where $\hat{F}^\Delta$ and $\hat{\tau}^\Delta$ are respectively the signed distance to the side of $\mathcal{D}^\Delta$ and the related discrete exit time.

- **Step 2.** The second step is to prove that the analogous version of Theorem 3 holds, with $\hat{\tau}^\Delta$ instead of τ^Δ. Actually, a careful reading of its proof shows that it is indeed the case, without modification.

(2) Finally, the last term in (2.13) is related to the sensitivity of a Dirichlet problem with respect to the domain. By an application of Theorem 2.2 in [CGK06] with $\Theta(t,x) = -c_0 \nabla F(t,x)|\nabla F \sigma(t,x)|$ (in $\mathcal{C}^{1,2}$), one gets that this contribution equals

$$-c_0\sqrt{\Delta}\mathbb{E}(\mathbf{1}_{\tau \le T} Z_\tau (\nabla u - \nabla g)(\tau, X_\tau) \cdot \nabla F(\tau, X_\tau)|\nabla F \sigma(\tau, X_\tau)|) + o(\sqrt{\Delta}).$$

This proves that the new procedure has an error $o(\sqrt{\Delta})$. □

3 Technical results concerning the overshoot

This section is devoted to the proof of Theorem 3. We first state some useful auxiliary results.

Lemma 8 (Bernstein's inequality) *Assume* $(\mathbf{A}_\theta\text{-}\mathbf{1})$. *Consider two stopping times S, S' upper bounded by T with $0 \le S' - S \le \Theta \le T$. Then for any $p \ge 1$, there are some constants $c > 0$ and $C := C((\mathbf{A}_\theta\text{-}\mathbf{1})\ , T)$, such that for any $\eta \ge 0$, one has a.s:*

$$\mathbb{P}[\sup_{t\in[S,S']} |X^\Delta_t - X^\Delta_S| \ge \eta \,\Big|\, \mathcal{F}_S] \le C \exp\left(-c\frac{\eta^2}{\Theta}\right),$$
$$\mathbb{E}[\sup_{t\in[S,S']} |X^\Delta_t - X^\Delta_S|^p \,\Big|\, \mathcal{F}_S] \le C\Theta^{p/2}.$$

For a proof of the first inequality we refer to Chapter 3, §3 in [RY99]. The last inequality easily follows from the first one or from the BDG inequalities.

Lemma 9 (Convergence of exit time) *Assume* $(\mathbf{A}'_\theta)$ *and that the domain is of class* $\mathbf{H}_2$. *The following convergences hold in probability:*

(1) $\lim_{\Delta\to 0} \tau^\Delta \wedge T = \tau \wedge T$;
(2) $\lim_{\Delta\to 0} X^\Delta_{\tau^\Delta\wedge T} = X_{\tau\wedge T}$;

(3) $\lim_{\Delta\to 0}\sup_{t\le T}|X^{\Delta}_{\phi(t)}-X_t|=0$.

The proof of the first two assertions in the case of space-time domain is analogous to the case of cylindrical domain (see [GM05]) and thus left to the reader. The last convergence is standard.

The following results are key tools to prove Theorem 3. A similar version is proved in [Sie79], but here, we additionally prove the uniform convergence.

Lemma 10 (Asymptotic independence of the overshoot and the discrete exit time). *Let W be a standard one dimensional BM. Put $x>0$ and consider the domain $\mathcal{D}:=]0,T[\times]-\infty,x[$. With the notation of Section 2, for any $\varepsilon>0$ we have*

$$\lim_{\Delta\longrightarrow 0}\sup_{t\in[0,T],y\ge 0,x\ge\Delta^{1/2-\varepsilon}}\Big|\mathbb{P}_0[\tau^{\Delta}\le t,(W_{\tau^{\Delta}}-x)\le y\sqrt{\Delta}]-\mathbb{P}_0[\tau\le t]H(y)\Big|=0. \tag{3.1}$$

If the Euler scheme starts close to the boundary at a small distance d, its discrete exit likely occurs after a time roughly equal to d^2. This feature is quantified in the above lemma.

Lemma 11 *Assume $(\mathbf{A}'_\theta)$, and that the domain is of class $\mathbf{H}_2$. Let $0<\beta<\alpha<1/2$. For all $\eta>0$, there exists $C:=C_\eta>0$ s.t. for Δ small enough, $\forall s\in\Delta\mathbb{N}\cap[0,T]$ and $\forall x\in V_{\partial D_s}(\Delta^{\alpha})\cap D_s$, one has*

$$\mathbb{P}[\tau^{\Delta}\wedge T\ge\Delta^{2\beta}|X^{\Delta}_s=x]\le C(\Delta^{\alpha-\beta-\eta}+\Delta^{\beta}),$$

where $\tau^{\Delta}:=\inf\{t_i>s:X^{\Delta}_{t_i}\notin D_{t_i}\}$.

Lemma 12 *Assume $(\mathbf{A}'_\theta)$, and that the domain is of class $\mathbf{H}_2$. There exists $C>0$, such that $\forall s\in\Delta\mathbb{N}\cap[0,T]$, $\forall x\in D_s$, $\forall t\in[s,T]$ and $\forall b\ge a\ge 0$, one has*

$$\mathbb{P}[\tau^{\Delta}\le t,\Delta^{-1/2}F^{-}(\tau^{\Delta},X^{\Delta}_{\tau^{\Delta}})\in[a,b]|X^{\Delta}_s=x]\le C\Big((b-a)+\Delta^{1/4}\Big)$$

where τ^{Δ} is shifted as in the previous lemma.

The proof of these three lemmas is postponed to Section 3.2.

We mention that if $\sigma\sigma^*$ is uniformly elliptic, Lemma 12 is valid without the $\Delta^{1/4}$ (see the proof for details). In that case, it means that the law of the renormalized overshoot is absolutely continuous w.r.t. the Lebesgue measure on $\mathbb{R}^+$, with a bounded density. This is also true at the limit, in view of Theorem 3.

Consider first the case $\mathcal{D} =]0,T[\times D$ where D is a half space. The theorem in the case of BM is then a direct consequence of Lemma 10. Now to deal with the Euler scheme, we introduce a first neighborhood whose distance to the boundary goes to 0 with Δ at a speed lower than $\Delta^{1/2}$ (below, the speed is tuned by a parameter α, see Figure 4). The characteristic exit time for a starting point in this neighborhood is short (Lemma 11), thus the diffusion coefficients are somehow constant and we are almost in the BM framework. Also, a second localization w.r.t. to the hitting time of this neighborhood guarantees that up to a rescaling we are far enough from the boundary to apply the renewal arguments needed for the asymptotic law of the overshoot (this is tuned by another parameter ε, see Figure 4).
For a more general time-space domain of class $\mathbf{H}_2$ two additional tools are used: a time-space change of chart and a local half space approximation of the domain by some tangent hyperplane.
For notational convenience, we assume from now on that the time-section domains $(D_t)_{t\in[0,T]}$ are convex so that $\pi_{\partial D_t}$ is always uniquely defined on D_t^c. To handle the case of general $\mathbf{H}_2$ domains, an additional localization procedure similar to the one of Theorem 6 is needed. We leave it to the reader.
For the sake of clarity, we also assume $k \equiv 0$ ($Z \equiv 1$). This is an easy simplification since owing to Lemma 9, $Z^{\Delta}_{\tau^{\Delta}\wedge T}$ converges to $Z_{\tau\wedge T}$ in L_1.

Step 1: preliminary localization. For $\alpha < 1/2$ specified later on, define $\tau_{\Delta^\alpha} := \inf\{t_i > 0 : F(t_i, X^{\Delta}_{t_i}) \le \Delta^\alpha\} \le \tau^{\Delta}$. We aim at studying the convergence of

$$\Psi_\Delta(t,x,y) := \mathbb{E}_x[\mathbf{1}_{\tau^\Delta \le t, F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) \ge y\sqrt{\Delta}} \varphi(X^\Delta_{\tau^\Delta})]$$

and for this, we define for all $0 \le s \le t < T$ ($s \in \Delta\mathbb{N}$), $(\tilde{x}, y) \in \mathbb{R}^d \times \mathbb{R}^+$

$$\begin{aligned}\Psi_\Delta(s,t,\tilde{x},y) :=& \mathbb{P}[\tau^\Delta \le t, F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) \ge y\sqrt{\Delta} | X^\Delta_s = \tilde{x}],\\ A(t,\alpha,\varepsilon) :=& \{\tau_{\Delta^\alpha} < \tau^\Delta, \tau_{\Delta^\alpha} < t, F(\tau_{\Delta^\alpha}, X^\Delta_{\tau_{\Delta^\alpha}}) \ge \Delta^{1/2-\varepsilon}\}.\end{aligned}$$

Here, ε is a fixed parameter in $]0,1/2[$, such that $\alpha < 1/2 - \varepsilon$ (take $\varepsilon = (\alpha + 1/2)/2$ for instance).
In the definition of Ψ_Δ, τ^Δ has to be understood as the shifted exit time $\inf\{t_i > s : X^\Delta_{t_i} \notin D_{t_i}\}$. By Lemma 8, $\mathbb{P}_x[\tau^\Delta = \tau_{\Delta^\alpha} \le t] + \mathbb{P}_x[\tau_{\Delta^\alpha} < t, F(\tau_{\Delta^\alpha}, X^\Delta_{\tau_{\Delta^\alpha}}) < \Delta^{1/2-\varepsilon}] = O_{pol}(\Delta)$ using $\alpha < 1/2 - \varepsilon$. Hence,

$$\begin{aligned}\Psi_\Delta(t,x,y) =& \mathbb{E}_x[\mathbf{1}_{A(t,\alpha,\varepsilon), F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) \ge y\sqrt{\Delta}} \varphi(X^\Delta_{\tau^\Delta}) \mathbf{1}_{\tau^\Delta \le t}] + O_{pol}(\Delta)\\ =& \mathbb{E}_x[\mathbf{1}_{A(t,\alpha,\varepsilon), F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) \ge y\sqrt{\Delta}} (\varphi(X^\Delta_{\tau^\Delta}) - \varphi(X^\Delta_{\tau_{\Delta^\alpha}})) \mathbf{1}_{\tau^\Delta \le t}]\\ &+ \mathbb{E}_x[\mathbf{1}_{A(t,\alpha,\varepsilon)} \varphi(X^\Delta_{\tau_{\Delta^\alpha}}) \Psi_\Delta(\tau_{\Delta^\alpha}, t, X^\Delta_{\tau_{\Delta^\alpha}}, y)] + O_{pol}(\Delta).\end{aligned}$$

The first term in the right hand side above converges to 0, using the conver-

gence in probability of $|X^{\Delta}_{\tau^{\Delta}\wedge T} - X^{\Delta}_{\tau_{\Delta^\alpha}\wedge T}|$ to 0 (analogously to Lemma 9). This gives

$$\Psi_\Delta(t,x,y) = \mathbb{E}_x[\mathbf{1}_{A(t,\alpha,\varepsilon)}\varphi(X^{\Delta}_{\tau_{\Delta^\alpha}})\Psi_\Delta(\tau_{\Delta^\alpha}, t, X^{\Delta}_{\tau_{\Delta^\alpha}}, y)] + o(1). \tag{3.2}$$



Figure 4. The two localization neighborhoods with $\alpha < \frac{1}{2} - \varepsilon$.

Let us comment again these two localisations. That with Δ^α enables us to freeze the coefficients of the Euler scheme, because the exit time is likely close to the initial time. That with $\Delta^{1/2-\varepsilon}$ ensures that it starts far enough from the boundary to induce the limiting behavior of the overshoot. This right balance regarding the distance of the initial point to the boundary is crucial. The final choice of α (and thus ε) depends on the regularity θ of the coefficients b and σ.
Now, it remains to study the convergence of $\Psi_\Delta(.)$.

Step 2: diffusion with frozen coefficients. Denote $\tau_{\Delta^\alpha} := \tilde{s}$, $X^{\Delta}_{\tau_{\Delta^\alpha}} := \tilde{x}$. Conditionally to $\mathcal{F}_{\tilde{s}}$, introduce now the one dimensional process $(Y_s)_{s\geq\tilde{s}}$, $Y_s = F(\tilde{s},\tilde{x}) + (\nabla F\sigma)(\tilde{s},\tilde{x})(W_s - W_{\tilde{s}})$. Note that we do not take into account the drift part in the frozen process. From the next localization procedure, it yields a negligible term. Since Y has constant coefficients, we apply below Lemma 10 to handle the overshoot of Y w.r.t. $\mathbb{R}^{+*}$. Define $\tau^{\Delta,Y} := \inf\{t_i > \tilde{s} : Y_{t_i} \leq 0\}$ and rewrite

$$\begin{aligned}\Psi_\Delta(\tilde{s},t,\tilde{x},y) &:= \Psi^C_\Delta(\tilde{s},t,\tilde{x},y) + R_\Delta(\tilde{s},t,\tilde{x},y), \\ \Psi^C_\Delta(\tilde{s},t,\tilde{x},y) &:= \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^{\Delta,Y} \leq t, (Y_{\tau^{\Delta,Y}})^- \geq y\sqrt{\Delta}].\end{aligned} \tag{3.3}$$

From **($\mathbf{A}'_\theta$-2')** that guarantees that Y has a non degenerate variance and Lemma 10, one gets

$$\sup_{(\tilde{s},\tilde{x})\in\mathcal{A}^{\alpha,\varepsilon}} |\Psi^C_\Delta(\tilde{s},t,\tilde{x},y) - \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^{\Delta,Y} \leq t](1 - H(y/|(\nabla F\sigma)(\tilde{s},\tilde{x})|))| \underset{\Delta\to 0}{\longrightarrow} 0,$$

where $\mathcal{A}^{\alpha,\varepsilon} := \{(t,x) : 0 \leq t \leq T, x \in V_{\partial D_t(\Delta^\alpha)} \backslash V_{\partial D_t}(\Delta^{1/2-\varepsilon})\}$. Plug now this identity in (3.3) to obtain with the same uniformity

$$\Psi_\Delta(\tilde{s},t,\tilde{x},y) = \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^{\Delta,Y} \leq t](1 - H(y/|(\nabla F\sigma)(\tilde{s},\tilde{x})|)) + R_\Delta(\tilde{s},t,\tilde{x},y) + o(1). \tag{3.4}$$

Step 3: control of the rests. We now show that $R_\Delta(\tilde{s},t,\tilde{x},y) = o(1)$ where the rest is still uniform for $(\tilde{s},\tilde{x}) \in \mathcal{A}^{\alpha,\varepsilon}$. This part is long and technical. First,

decomposing the space using the events $\{\tau^\Delta = \tau^{\Delta,Y}\}$, $\{F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) \geq y\sqrt{\Delta}\}$, $\{(Y_{\tau^{\Delta,Y}})^- \geq y\sqrt{\Delta}\}$ and their complementary events, write:

$$\begin{aligned}|R_\Delta(\tilde{s},t,\tilde{x},y)| &\leq R^1_\Delta(\tilde{s},t,\tilde{x})\\ &+ \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^\Delta \leq t, F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) \geq y\sqrt{\Delta}, (Y_{\tau^{\Delta,Y}})^- < y\sqrt{\Delta}, \tau^\Delta = \tau^{\Delta,Y}]\\ &+ \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^\Delta \leq t, F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) < y\sqrt{\Delta}, (Y_{\tau^{\Delta,Y}})^- \geq y\sqrt{\Delta}, \tau^\Delta = \tau^{\Delta,Y}] \quad (3.5)\end{aligned}$$

with $R^1_\Delta(\tilde{s},t,\tilde{x}) \leq \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^\Delta \leq t, \tau^\Delta \neq \tau^{\Delta,Y}] + \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^{\Delta,Y} \leq t, \tau^\Delta \neq \tau^{\Delta,Y}] := (R^{11}_\Delta + R^{12}_\Delta)(\tilde{s},t,\tilde{x})$. Let y_Δ be a given positive function of the time-step s.t. $y_\Delta \underset{\Delta\to 0}{\to} 0$ specified later on.
On the event $\{\tau^\Delta = \tau^{\Delta,Y}, |Y_{\tau^{\Delta,Y}} - F(\tau^{\Delta,Y}, X^\Delta_{\tau^{\Delta,Y}})| \leq y_\Delta\sqrt{\Delta}\}$, the conditions $F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) \geq y\sqrt{\Delta}$ and $(Y_{\tau^{\Delta,Y}})^- < y\sqrt{\Delta}$ imply $\Delta^{-1/2}(Y_{\tau^{\Delta,Y}})^- \in [y - y_\Delta, y)$. Similarly, $(Y_{\tau^{\Delta,Y}})^- \geq y\sqrt{\Delta}$ and $F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) < y\sqrt{\Delta}$ imply $\Delta^{-1/2}(Y_{\tau^{\Delta,Y}})^- \in [y, y + y_\Delta)$. Hence, by setting

$$\begin{aligned}R^2_\Delta(\tilde{s},t,\tilde{x}) &:= 2\mathbb{P}_{\tilde{s},\tilde{x}}[\tau^{\Delta,Y} \leq t, \tau^\Delta = \tau^{\Delta,Y}, |Y_{\tau^{\Delta,Y}} - F(\tau^{\Delta,Y}, X^\Delta_{\tau^{\Delta,Y}})| > y_\Delta\sqrt{\Delta}],\\ R^3_\Delta(\tilde{s},t,\tilde{x},y) &:= \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^{\Delta,Y} \leq t, \Delta^{-1/2}(Y_{\tau^{\Delta,Y}})^- \in [y - y_\Delta, y + y_\Delta), \tau^\Delta = \tau^{\Delta,Y}],\end{aligned}$$

we obtain $R_\Delta(\tilde{s},t,\tilde{x},y)| \leq (R^1_\Delta + R^2_\Delta)(\tilde{s},t,\tilde{x}) + R^3_\Delta(\tilde{s},t,\tilde{x},y)$.
Term $R^3_\Delta(\tilde{s},t,\tilde{x},y)$. From Lemma 12 applied to the process with frozen coefficients, one gets

$$R^3_\Delta(\tilde{s},t,\tilde{x},y) \leq C(y_\Delta + \Delta^{1/4}). \quad (3.6)$$

Term $R^2_\Delta(\tilde{s},t,\tilde{x})$. Let us explain the leading ideas of the estimates below. Usually, it is easy to prove inequalities like $|Y_t - F(t, X^\Delta_t)|_{L_2} = O(\Delta^{1/2})$ (for a fixed t), but this not enough to control R^2_Δ. To achieve our goal, we take advantage of the fact that the time t is the stopping time $\tau^{\Delta,Y}$ which is likely close to $\tilde{s}$. Thus, $Y_{\tau^{\Delta,Y}} - F(\tau^{\Delta,Y}, X^\Delta_{\tau^{\Delta,Y}})$ should be much smaller that $\Delta^{1/2}$ in L_2-norm.
Introduce for $0 < \beta < \alpha < 1/2$, $\tau_{\Delta^\beta} := \inf\{s > \tilde{s} : |X^\Delta_s - \tilde{x}| \geq \Delta^\beta\} \wedge (\tilde{s} + \Delta^\delta)$, $\delta := 2\beta + \gamma$, $\gamma > 0$. Clearly, one has

$$\begin{aligned}|R^2_\Delta(\tilde{s},t,\tilde{x})| \leq& 2\mathbb{P}_{\tilde{s},\tilde{x}}\Big[\tau^{\Delta,Y} \leq t, \tau^\Delta = \tau^{\Delta,Y}, \tau^\Delta < \tau_{\Delta^\beta},\\ &\qquad |Y_{\tau^{\Delta,Y}} - F(\tau^{\Delta,Y}, X^\Delta_{\tau^{\Delta,Y}})| > y_\Delta\sqrt{\Delta}\Big] + 2\mathbb{P}_{\tilde{s},\tilde{x}}[\tau^\Delta \geq \tau_{\Delta^\beta}, \tau^\Delta \leq t]\\ :=& (R^{21}_\Delta + R^{22}_\Delta)(\tilde{s},t,\tilde{x}).\end{aligned}$$

Let us first deal with $R^{21}_\Delta(\tilde{s},t,\tilde{x})$. By the Markov inequality, one has

$$R^{21}_\Delta(\tilde{s},t,\tilde{x}) \leq 2\Delta^{-1}y_\Delta^{-2}\mathbb{E}_{\tilde{s},\tilde{x}}\Big[\mathbf{1}_{\tau^\Delta < \tau_{\Delta^\beta}, \tau^{\Delta,Y} \leq t, \tau^\Delta = \tau^{\Delta,Y}}|Y_{\tau^{\Delta,Y}} - F(\tau^{\Delta,Y}, X^\Delta_{\tau^{\Delta,Y}})|^2\Big]. \quad (3.7)$$

Note that since $\mathcal{D}$ is of class $\mathbf{H}_2$, F has the same regularity, i.e. it is uniformly Lipschitz continuous in time, its first space derivatives are uniformly Lipschitz

continuous in space and 1/2-Hlder continuous in time. Thus, assuming up to a regularization procedure that $F \in C^{1,2}([0,T]\times\mathbb{R}^d)$, It's formula yields for all $t \geq \tilde{s}$,

$$\begin{aligned}
F(t,X_t^\Delta) =& F(\tilde{s},\tilde{x}) + \int_{\tilde{s}}^t \nabla F(s,X_s^\Delta)dX_s^\Delta \\
&+ \int_{\tilde{s}}^t \Big(\partial_s F(s,X_s^\Delta) + \frac{1}{2}\mathrm{Tr}(H_F(s,X_s^\Delta)\sigma\sigma^*(\phi(s),X_{\phi(s)}^\Delta))\Big)ds \\
:=& F(\tilde{s},\tilde{x}) + \int_{\tilde{s}}^t \nabla F(s,X_s^\Delta)\sigma(\phi(s),X_{\phi(s)}^\Delta)dW_s + R_F^\Delta(\tilde{s},t,\tilde{x}) \qquad (3.8)\\
=& Y_t + R_F^\Delta(\tilde{s},t,\tilde{x}) + \int_{\tilde{s}}^t \Big(\nabla F(s,X_s^\Delta)\sigma(\phi(s),X_{\phi(s)}^\Delta) - [\nabla F\sigma](\tilde{s},\tilde{x})\Big)dW_s.
\end{aligned}$$

From **($\mathbf{A}_\theta'$-1)** and the assumptions on $\mathcal{D}$ one derives $|R_F^\Delta|(\tilde{s},t,\tilde{x}) \leq C(t-\tilde{s})$. Thus, for any given stopping time $U \in [\tilde{s},\tau_{\Delta^\beta}]$, the working assumptions (i.e. smoothness of σ, F) and standard computations yield

$$\mathbb{E}[|F(U,X_U^\Delta) - Y_U|^2] \leq C(\Delta^{2\beta+\delta} + \Delta^{\delta(1+\theta)}).$$

From (3.7) and the above control with $U = \tau^{\Delta,Y} \wedge \tau_{\Delta^\beta}$, one obtains

$$R_\Delta^{21}(\tilde{s},t,\tilde{x}) \leq C y_\Delta^{-2}\Delta^{-1}(\Delta^{2\beta+\delta} + \Delta^{\delta(1+\theta)}). \qquad (3.9)$$

Let us now control $R_\Delta^{22}(\tilde{s},t,\tilde{x})$. From Lemmas 8 and 11, for any $\eta > 0$ we write

$$\begin{aligned}
R_\Delta^{22}(\tilde{s},t,\tilde{x}) &\leq \mathbb{P}_{\tilde{s},\tilde{x}}[\tau_{\Delta^\beta} < \tilde{s} + \Delta^\delta] + \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^\Delta \wedge t \geq \tilde{s} + \Delta^\delta] \\
&\leq C_\eta\Big(\exp\Big(-c\Delta^{2\beta-\delta}\Big) + \Delta^{\alpha-\eta-\delta/2} + \Delta^{\delta/2}\Big). \qquad (3.10)
\end{aligned}$$

Take now $\alpha = \frac{1+\frac{\theta}{2}}{2(1+\theta)} < 1/2$, $\eta = \frac{\theta}{16(\theta+1)}$, $\gamma = \frac{1}{8(1+\theta)}$, $y_\Delta = \Delta^{\theta/16}$. Check that for $\delta = 2\beta + \gamma = 2\alpha - 4\eta$, one has $\delta = \frac{1+\theta/4}{1+\theta}$, $\beta = \frac{7/8+\theta/4}{2(1+\theta)} < \alpha$, $3\eta < \alpha$. Thus, $R_\Delta^{22}(\tilde{s},t,\tilde{x}) = O(\Delta^\eta)$. In addition, $y_\Delta^{-2}\Delta^{\delta(1+\theta)-1} = \Delta^{\theta/8}, y_\Delta^{-2}\Delta^{2\beta+\delta-1} = O(\Delta^{1/(8(1+\theta))})$. Hence, from (3.9) and (3.10)

$$R_\Delta^2(\tilde{s},t,\tilde{x}) \leq C\Big(\Delta^{1/(8(1+\theta))} + \Delta^{\theta/8} + \Delta^{\theta/(16(\theta+1))}\Big) \leq C\Delta^{\theta/32}. \qquad (3.11)$$

<u>*Term* $R_\Delta^1(\tilde{s},t,\tilde{x})$.</u> We give an upper bound for $R_\Delta^{11}(\tilde{s},t,\tilde{x})$. The term $R_\Delta^{12}(\tilde{s},t,\tilde{x})$ can be handled in the same way. From the previous control on $R_\Delta^{22}(\tilde{s},t,\tilde{x})$ and for the previous parameters, one gets

$$\begin{aligned}
R_\Delta^{11}(\tilde{s},t,\tilde{x}) =& \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^\Delta \leq t, \tau^\Delta \neq \tau^{\Delta,Y}, \tau^\Delta < \tau_{\Delta^\beta}] + O(\Delta^\eta) \\
=& \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^\Delta \leq t, \tau^\Delta > \tau^{\Delta,Y}, \tau^\Delta < \tau_{\Delta^\beta}] \\
&+ \mathbb{P}_{\tilde{s},\tilde{x}}[\tau^\Delta \leq t, \tau^\Delta < \tau^{\Delta,Y}, \tau^\Delta < \tau_{\Delta^\beta}] + O(\Delta^\eta).
\end{aligned}$$

Then, splitting the first probability according to $\Delta^{-1/2}(Y_{\tau^{\Delta,Y}})^- \leq y_\Delta$ or not,

and the second one according to $\Delta^{-1/2}F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) \le y_\Delta$ or not, we obtain

$$\begin{aligned}
&R^{11}_\Delta(\tilde{s},t,\tilde{x})\\
\le&\Big(\mathbb{P}_{\tilde{s},\tilde{x}}[\tau^{\Delta,Y} \le t, \Delta^{-1/2}(Y_{\tau^{\Delta,Y}})^- \le y_\Delta]\\
&+\mathbb{P}_{\tilde{s},\tilde{x}}[\tau^\Delta \le t, \tau^\Delta > \tau^{\Delta,Y}, \tau^\Delta < \tau_{\Delta^\beta}, \Delta^{-1/2}|Y_{\tau^{\Delta,Y}} - F(\tau^{\Delta,Y}, X^\Delta_{\tau^{\Delta,Y}})| \ge y_\Delta]\Big)\\
+&\Big(\mathbb{P}_{\tilde{s},\tilde{x}}[\tau^\Delta \le t, \tau^\Delta < \tau^{\Delta,Y}, \tau^\Delta < \tau_{\Delta^\beta}, \Delta^{-1/2}|Y_{\tau^\Delta} - F(\tau^\Delta, X^\Delta_{\tau^\Delta})| \ge y_\Delta]\\
&+\mathbb{P}_{\tilde{s},\tilde{x}}[\tau^\Delta \le t, \Delta^{-1/2}F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) \le y_\Delta]\Big) + C\Delta^\eta,
\end{aligned}$$

for the previous function $(y_\Delta)_{\Delta>0}$. Since we could obtain the same type of bound for $R^{12}_\Delta(\tilde{s},t,\tilde{x})$, from Lemma 12 and following the computations that gave (3.9) we derive for the previous set of parameters

$$R^1_\Delta(\tilde{s},t,\tilde{x}) \le C(y_\Delta^{-2}\Delta^{-1}(\Delta^{2\beta+\delta} + \Delta^{\delta(1+\theta)}) + \Delta^\eta + y_\Delta + \Delta^{1/4}) \le C\Delta^{\theta/32}. \tag{3.12}$$

From (3.12), (3.11), (3.6) we finally obtain $R_\Delta(\tilde{s},t,\tilde{x},y) = O(\Delta^{\theta/32}) = o(1)$. The rest is uniform w.r.t. $(\tilde{s},\tilde{x},y) \in \mathcal{A}^{\alpha,\varepsilon} \times \mathbb{R}^+$.

Step 4. Final step. Plug the previous results in (3.4). We derive from (3.2)

$$\begin{aligned}
\Psi_\Delta(t,x,y) = \mathbb{E}_x[&\mathbf{1}_{A(t,\alpha,\varepsilon)}\varphi(X^\Delta_{\tau_{\Delta^\alpha}})\\
&\times \mathbb{P}_{\tau_{\Delta^\alpha},X^\Delta_{\tau_{\Delta^\alpha}}}[\tau^{\Delta,Y} \le t](1 - H(y/|\nabla F\sigma(\tau_{\Delta^\alpha}, X^\Delta_{\tau_{\Delta^\alpha}})|))] + o(1).
\end{aligned}$$

Moreover, note that taking $y = 0$ in the previous controls gives immediately

$$\mathbb{P}_{\tilde{s},\tilde{x}}(\tau^{\Delta,Y} \le t) - \mathbb{P}_{\tilde{s},\tilde{x}}(\tau^\Delta \le t) = o(1)$$

uniformly in $(\tilde{s},\tilde{x}) \in \mathcal{A}^{\alpha,\varepsilon}$. Thus, we finally obtain

$$\Psi_\Delta(t,x,y) = \mathbb{E}_x[\mathbf{1}_{A(t,\alpha,\varepsilon)}\varphi(X^\Delta_{\tau_{\Delta^\alpha}})\mathbf{1}_{\tau^\Delta \le t}(1 - H(y/|\nabla F\sigma(\tau_{\Delta^\alpha}, X^\Delta_{\tau_{\Delta^\alpha}})|))] + o(1).$$

Under continuity arguments as in step 1 (localization), we eventually get

$$\Psi_\Delta(t,x,y) = \mathbb{E}_x[\mathbf{1}_{\tau^\Delta \le t}\varphi(X^\Delta_{\tau^\Delta})(1 - H(y/|\nabla F\sigma(\tau^\Delta, X^\Delta_{\tau^\Delta})|))] + o(1).$$

We complete the proof using Lemma 9:

$$\Psi_\Delta(t,x,y) \underset{\Delta\to 0}{\to} \mathbb{E}_x[\mathbf{1}_{\tau \le t}\varphi(X_\tau)(1 - H(y/|\nabla F\sigma(\tau, X_\tau)|))].$$

□

3.2 Proof of Lemmas 10, 11 and 12

Proof of Lemma 10. We shall insist on the dependence of the exit times with respect to x, by setting $\tau^\Delta := \inf\{t_i = i\Delta > 0 : W_{t_i} \geq x\} := \tau_x^\Delta$ and analogously for $\tau = \tau_x$. Our proof relies on the following convergence (see equation (19) in Siegmund [Sie79]): if we set (for any $y, z \geq 0$)

$$D(z, y) = \mathbb{P}_0[W_{\tau_z^\Delta} - z \leq y\sqrt{\Delta}] - H(y),$$

then

$$\lim_{z\Delta^{-1/2} \to +\infty} |D(z, y)| = 0.$$

Using the monotonicity and the uniform continuity of $H(y)$, Dini's Theorem yields that the above limit is actually uniform with respect to $y \geq 0$. It follows

$$\sup_{y\geq 0,\ z\in[\Delta^{1/2-\varepsilon/3},\infty)} |D(z, y)| \underset{\Delta\to 0}{\to} 0. \tag{3.13}$$

Additionnally, we have

$$\sup_{x\geq 0,\ t\in[\Delta^{1-4\varepsilon/3},T]} |\mathbb{P}_0(\tau_x^\Delta > t) - \mathbb{P}_0(\tau_x > t)| \underset{\Delta\to 0}{\to} 0. \tag{3.14}$$

To prove this, we apply Theorem 3.4 in [Avi07] which states that

$$\sup_{x\in\mathbb{R}} \mathbb{E}|\mathbf{1}_{M<x} - \mathbf{1}_{\hat{M}<x}| \leq 3(\sup_{m\in\mathbb{R}} f_M(m)\|M - \hat{M}\|_{L_p})^{\frac{p}{p+1}}$$

for any $p > 0$ and for any random variables M and $\hat{M}$, such that M has a bounded density $f_M(.)$. Now, consider $M = \sup_{s\leq t} W_s$ and $\hat{M} = \sup_{s=i\Delta\leq t} W_s$. The density of M is bounded by $2/\sqrt{2\pi t}$. On the other hand, Lemma 6 in [AGP95] gives $\|M - \hat{M}\|_{L_p} \leq C_p(T)\Delta^{1/2}$. Hence, we get for $t \geq \Delta^{1-4\varepsilon/3}$,

$$|\mathbb{P}_0(\tau_x^\Delta > t) - \mathbb{P}_0(\tau_x > t)| \leq \mathbb{E}|\mathbf{1}_{\hat{M}<x} - \mathbf{1}_{M<x}| \leq C_p(T)\Delta^{\frac{2\varepsilon p}{3(p+1)}},$$

which leads to (3.14).

We can now proceed to the proof of Lemma 10, assuming that $x \geq \Delta^{1/2-\varepsilon}$. First, note that if $x/\sqrt{t} \geq \Delta^{-\varepsilon/3} \to +\infty$ as $\Delta \to 0$, $\mathbb{P}_0(\tau_x^\Delta \leq t)$ and $\mathbb{P}_0(\tau_x \leq t)$ are both $O_{pol}(\Delta)$. Thus, the difference in Lemma 10 converges to 0 as $\Delta \to 0$.

Suppose now that $x/\sqrt{t} \leq \Delta^{-\varepsilon/3}$, hence $\sqrt{t} \geq x\Delta^{\varepsilon/3} \geq \Delta^{1/2-2\varepsilon/3}$, and write for $t \in \Delta\mathbf{N}^*$

$$P := \mathbb{P}_0[\tau_x^\Delta > t, W_{\tau_x^\Delta} - x \leq y\sqrt{\Delta}] = \int_0^{+\infty} q_t^{x,\Delta}(0, x - z)\mathbb{P}_0[W_{\tau_z^\Delta} - z \leq y\sqrt{\Delta}]dz$$

where $q_t^{x,\Delta}(.,.)$ denotes the transition density of the Brownian motion discretely killed at level x. Introduce the partition $\mathbb{R}^+ = [0, \Delta^{1/2-\varepsilon/3}) \cup [\Delta^{1/2-\varepsilon/3}, +\infty)$.

Then,

$$P = R + \int_{\Delta^{1/2-\varepsilon/3}}^{+\infty} q_t^{x,\Delta}(0, x-z)D(z,y)dz + \mathbb{P}_0[\tau_x^\Delta > t]H(y)$$

where $|R| \leq 2\mathbb{P}_0[W_t \in [x-\Delta^{1/2-\varepsilon/3}, x]] \leq \frac{2}{\sqrt{2\pi t}}\Delta^{1/2-\varepsilon/3} \leq \frac{2}{\sqrt{2\pi}}\Delta^{\varepsilon/3}$ since $\sqrt{t} \geq \Delta^{1/2-2\varepsilon/3}$. Finally, taking advantage of the estimates (3.13) and (3.14) readily completes our proof. □

Proof of Lemma 11. We take $s = 0$ for notational simplicity. Introduce $\tau_{\Delta^\beta} := \inf\{t \geq 0 : X_t^\Delta \notin V_{\partial D_t}(\Delta^\beta)\}$ and for $\gamma > 0$ write from Lemma 8 and the notation of (3.8) (up to the same regularization procedure concerning F)

$$\begin{aligned}\mathbb{P}_x[\tau^\Delta \wedge T \geq \Delta^{2\beta}] =& \mathbb{P}_x[\inf_{0\leq i\leq \Delta^{2\beta-1}} \Big(F(0,x) + \int_0^{t_i} \nabla F(s, X_s^\Delta)\sigma(\phi(s), X_{\phi(s)}^\Delta)dW_s \\ &+ R_F^\Delta(0,t_i,x)\Big) \geq 0, \tau_{\Delta^\beta} \geq \Delta^{2\beta+\gamma}] + O_{pol}(\Delta) := Q,\end{aligned}$$

where under the assumptions of the Lemma, $|R_F^\Delta(0,t_i,x)| \leq Ct_i$ and $F(0,x) \leq \Delta^\alpha$. For a given $r > 0$, consider the event $\mathcal{A}_r = \{\exists s \leq T : |X_s^\Delta - X_{\phi(s)}^\Delta| \geq r\}$ where the increments of X^Δ between two close times are large: by Lemma 8, it has an exponentially small probability. Hence, if we set

$$M_u := \int_0^u \nabla F(s, X_s^\Delta)\sigma(\phi(s), X_{\phi(s)}^\Delta)dW_s := B_{\langle M\rangle_u},\ \tilde{t}_i = \langle M\rangle_{t_i},$$

B is a standard Brownian motion (on a possibly enlarged probability space) owing to the Dambis, Dubbins-Schwarz Theorem, cf. Theorem V.1.7 in [RY99]. In addition, the above time change is strictly increasing on the set $\mathcal{A}_r^c$ and $\langle M\rangle_t - \langle M\rangle_s \geq (t-s)a_0/2$ $(t \geq s)$ up to taking r small enough, because **($\mathbf{A}_\alpha^{'}$-2)** is in force. It readily follows that

$$\begin{aligned}Q \leq& \mathbb{P}_x[\inf_{0\leq i\leq \Delta^{2\beta+\gamma-1}}(M_{t_i} + Ct_i) \geq -\Delta^\alpha, \tau_{\Delta^\beta} \geq \Delta^{2\beta+\gamma}] + O_{pol}(\Delta) \\ \leq& \mathbb{P}_x[\inf_{0\leq i\leq \Delta^{2\beta+\gamma-1}}(B_{\tilde{t}_i} + 2Ca_0^{-1}\tilde{t}_i) \geq -\Delta^\alpha, \tau_{\Delta^\beta} \geq \Delta^{2\beta+\gamma}, \mathcal{A}_r^c] + O_{pol}(\Delta) \\ \leq& \mathbb{P}_x[\inf_{0\leq i\leq \Delta^{2\beta+\gamma-1}}(B_{\tilde{t}_i} + 2Ca_0^{-1}\tilde{t}_i) \geq -\Delta^\alpha, \tau_{\Delta^\beta} \geq \Delta^{2\beta+\gamma}, \\ &\inf_{0\leq s\leq \langle M\rangle_{\Delta^{2\beta+\gamma}}}(B_s + 2Ca_0^{-1}s) \leq -\Delta^{\alpha-\zeta}, \mathcal{A}_r^c] + O_{pol}(\Delta) \\ &+ \mathbb{P}_x[\tau_{\Delta^\beta} \geq \Delta^{2\beta+\gamma}, \inf_{0\leq s\leq \langle M\rangle_{\Delta^{2\beta+\gamma}}}(B_s + 2Ca_0^{-1}s) \geq -\Delta^{\alpha-\zeta}, \mathcal{A}_r^c],\end{aligned}$$

for $\zeta > 0$. Thus, from Lemma 8 and standard controls

$$\begin{aligned}Q \leq& \mathbb{P}_x[\exists i : 0 \leq i \leq \Delta^{2\beta+\gamma-1}, \sup_{s\in[\tilde{t}_i,\tilde{t}_{i+1}]}|B_s - B_{\tilde{t}_i} + 2Ca_0^{-1}(s-\tilde{t}_i)| \geq \Delta^{\alpha-\zeta} - \Delta^\alpha, \\ &\tau_{\Delta^\beta} \geq \Delta^{2\beta+\gamma}] + \mathbb{P}_x[\inf_{0\leq s\leq a_0\Delta^{2\beta+\gamma}/2} B_s \geq -\Delta^{\alpha-\zeta} - C\Delta^{2\beta+\gamma}] + O_{pol}(\Delta) \\ \leq& O_{pol}(\Delta) + C(\Delta^{\alpha-\zeta-\beta-\gamma/2} + \Delta^{\beta+\gamma/2}).\end{aligned}$$

Choose now γ, ζ s.t. $(\zeta + \frac{\gamma}{2}) = \eta > 0$. The proof is complete. $\square$

Proof of Lemma 12. Taking also $s = 0$ for notational convenience, we write

$$\begin{aligned} P :=& \mathbb{P}_x[\tau^\Delta \le t, \Delta^{-1/2} F^-(\tau^\Delta, X^\Delta_{\tau^\Delta}) \in [a,b]] \le O_{pol}(\Delta) \\ &+ \sum_{i=1}^{\lfloor t/\Delta \rfloor} \mathbb{E}_x[\mathbf{1}_{\tau^\Delta > t_{i-1}, X^\Delta_{t_{i-1}} \in V_{\partial D_{t_{i-1}}}(r_0)} \mathbb{P}_{\mathcal{F}_{t_{i-1}}}[\Delta^{-1/2} F^-(t_i, X^\Delta_{t_i}) \in [a,b]]] \end{aligned} \tag{3.15}$$

using Lemma 8 for the last identity.
A Taylor formula gives: $F(t_i, X^\Delta_{t_i}) = F(t_{i-1}, X^\Delta_{t_{i-1}}) + \Sigma_{t_{i-1}}(W_{t_i} - W_{t_{i-1}}) + R^\Delta_{t_{i-1},t_i} := \mathcal{N}_{t_{i-1}} + R^\Delta_{t_{i-1},t_i}$ where $\Sigma_{t_{i-1}} = \nabla F \sigma(t_{i-1}, X^\Delta_{t_{i-1}})$, $\mathbb{E}_{\mathcal{F}_{t_{i-1}}}[|R^\Delta_{t_{i-1},t_i}|^2] \le C\Delta^2$. Conditionally to $\mathcal{F}_{t_{i-1}}$, $\mathcal{N}_{t_{i-1}}$ has a Gaussian distribution $\mathcal{N}(F(t_{i-1}, X^\Delta_{t_{i-1}}), \|\Sigma_{t_{i-1}}\|^2 \Delta)$.
In addition, on the event $X^\Delta_{t_{i-1}} \in V_{\partial D_{t_{i-1}}(r_0)}$, $\|\Sigma_{t_{i-1}}\|^2 \Delta \ge a_0 \Delta$ and we obtain

$$\begin{aligned} Q_{i-1} :=& \mathbb{P}_{\mathcal{F}_{t_{i-1}}}[F^-(t_i, X^\Delta_{t_i}) \in [a\Delta^{1/2}, b\Delta^{1/2}]] \\ =& \mathbb{P}_{\mathcal{F}_{t_{i-1}}}[(\mathcal{N}_{t_{i-1}} + R^\Delta_{t_{i-1},t_i})^- \in [a\Delta^{1/2}, b\Delta^{1/2}]] \\ \le& \mathbb{P}_{\mathcal{F}_{t_{i-1}}}[\mathcal{N}_{t_{i-1}} \in [-b\Delta^{1/2} - \Delta^{3/4}, -a\Delta^{1/2} + \Delta^{3/4}]] \\ &+ \mathbb{P}_{\mathcal{F}_{t_{i-1}}}[|R^\Delta_{t_{i-1},t_i}| \ge \Delta^{3/4}, X^\Delta_{t_i} \notin \mathcal{D}_{t_i}] \\ \le& \mathbb{P}_{\mathcal{F}_{t_{i-1}}}[\mathcal{N}_{t_{i-1}} \in [-\Delta^{1/2}(b + \Delta^{1/4}), -\Delta^{1/2}(a - \Delta^{1/4})]] \\ &+ C\Delta^{1/4} \exp\left(-c \frac{d(X^\Delta_{t_{i-1}}, \partial D_{t_{i-1}})^2}{\Delta}\right) \end{aligned}$$

using the Cauchy-Schwarz inequality and Lemma 8 for the last inequality. Hence, we derive from (3.15)

$$\begin{aligned} P \le& \sum_{i=1}^{\lfloor t/\Delta \rfloor} \mathbb{E}_x[\mathbf{1}_{\tau^\Delta > t_{i-1}, X^\Delta_{t_{i-1}} \in V_{\partial D_{t_{i-1}}}(r_0)} \left(C\Delta^{1/4} \exp\left(-c \frac{d(X^\Delta_{t_{i-1}}, \partial D_{t_{i-1}})^2}{\Delta}\right) \right. \\ &\left. + \int_{-\Delta^{1/2}(b+\Delta^{1/4})}^{-\Delta^{1/2}(a-\Delta^{1/4})} \exp\left(-\frac{(y - F(t_{i-1}, X^\Delta_{t_{i-1}}))^2}{2\|\Sigma_{t_{i-1}}\|^2 \Delta}\right) \frac{dy}{(2\pi\Delta)^{1/2} \|\Sigma_{t_{i-1}}\|} \right)] + O_{pol}(\Delta). \end{aligned}$$

We now upper bound the above integral on the event $\{\tau^\Delta > t_{i-1}\} \subset \{F(t_{i-1}, X^\Delta_{t_{i-1}}) > 0\}$.

- If $y \le 0$, clearly one has $(y - F(t_{i-1}, X^\Delta_{t_{i-1}}))^2 \ge F^2(t_{i-1}, X^\Delta_{t_{i-1}})$.
- If $y \in (0, [\Delta^{1/2}(\Delta^{1/4} - a)]_+)$, one has $(y - F(t_{i-1}, X^\Delta_{t_{i-1}}))^2 \ge \frac{1}{2} F^2(t_{i-1}, X^\Delta_{t_{i-1}}) - y^2 \ge \frac{1}{2} F^2(t_{i-1}, X^\Delta_{t_{i-1}}) - \Delta^{3/2}$.

Thus, we obtain that P is bounded by

$$C(b - a + \Delta^{1/4}) \sum_{i=1}^{\lfloor t/\Delta \rfloor} \mathbb{E}_x[\mathbf{1}_{\tau^\Delta > t_{i-1}, X^\Delta_{t_{i-1}} \in V_{\partial D_{t_{i-1}}}(r_0)} \exp(-c \frac{F^2(t_{i-1}, X^\Delta_{t_{i-1}})}{\Delta})] + O_{pol}(\Delta).$$

The end of the proof is now achieved by standard computations done in [GM04] p. 212 to 217. We only mention the main steps and refer for the details to the above reference. First, we replace the discrete sum on i by a continuous integral, then we apply the occupation time formula to the distance process $(F(s, X_s^\Delta))_{s\leq\tau^\Delta}$ using the non characteristic boundary condition, as in the proof of Theorem 6:

$$\begin{aligned} P \leq & C\frac{(b-a+\Delta^{1/4})}{\Delta}\int_0^t \mathbb{E}_x[\mathbf{1}_{\tau^\Delta>s, X_s^\Delta\in V_{\partial D_s}(r_0)}\exp(-c\frac{F^2(s,X_s^\Delta)}{\Delta})]ds + O_{pol}(\Delta) \\ \leq & C\frac{(b-a+\Delta^{1/4})}{\Delta}\int_{-r_0}^{r_0}\exp(-c\frac{y^2}{\Delta})\mathbb{E}_x[L^y_{t\wedge\tau^\Delta}(F(.,X_.^\Delta))]dy + O_{pol}(\Delta). \end{aligned}$$

Then, we use (2.6) to obtain $P \leq C((b-a+\Delta^{1/4})$ which is our claim. □

Remark 13 *Finally, we mention that if $\sigma\sigma^*$ is uniformly elliptic, the rest $R^\Delta_{t_{i-1},t_i}$ can be avoided and the result can be stated without the contribution $\Delta^{1/4}$. Indeed, we can directly exploit that the Euler scheme has conditionally a non degenerate Gaussian distribution and usual changes of chart associated to a parametrization of the boundary (see e.g. [Gob00]) give the expected result.*

4 Extension to the stationary case

4.1 Framework

In this section we assume that the coefficients in (1.1) are time independent and that the mappings b, σ are uniformly Lipschitz continuous, i.e. $(X_t)_{t\geq 0}$ is the unique strong solution of

$$X_t = x + \int_0^t b(X_s)ds + \int_0^t \sigma(X_s)dW_s, t \geq 0,\ x \in \mathbb{R}^d.$$

For a bounded domain $D \subset \mathbb{R}^d$, and given functions $f, g, k : \bar{D} \to \mathbb{R}$, we are interested in estimating

$$u(x) := \mathbb{E}_x[g(X_\tau)Z_\tau + \int_0^\tau f(X_s)Z_s ds],\ Z_s = \exp(-\int_0^s k(X_r)dr), \tag{4.1}$$

where $\tau := \inf\{t > 0 : X_t \notin D\}$.

Adapting freely the previous notations for Hlder spaces to the elliptic setting, introduce for $\theta \in]0, 1]$:

($\mathbf{A}_\theta$) 1. *Smoothness of the coefficients.* $b, \sigma \in \mathbf{H}_{1+\theta}$.

2. *Uniform ellipticity.* For some $a_0 > 0$, $\forall (x,\xi) \in \mathbb{R}^d \times \mathbb{R}^d$, $\xi^* \sigma\sigma^*(x)\xi \geq a_0|\xi|^2$.

(D) *Smoothness of the domain.* The bounded domain D is of class $\mathbf{H}_2$.

($\mathbf{C}_\theta$) *Other coefficients.* The boundary data $g \in \mathbf{H}_{1+\theta}$, $f, k \in \mathbf{H}_{1+\theta}$ and $k \geq 0$.

Note that under **($\mathbf{A}_\theta$)** and since D is bounded, Lemma 3.1 Chapter III of [Fre85] yields $\sup_{x\in\bar{D}} \mathbb{E}_x[\tau] < \infty$. Thus, (4.1) is well defined under our current assumptions.

From Theorem 6.13, the final notes of Chapter 6 in [GT98] and Theorem 2.1 Chapter II in Freidlin [Fre85], the Feynman-Kac representation in our elliptic setting writes

Proposition 14 (Elliptic Feynman-Kac's formula and estimates) *Assume* **($\mathbf{A}_\theta$)**, **(D)**, **($\mathbf{C}_\theta$)** *are in force. Then, there is a unique solution in* $\mathbf{H}_{1+\theta} \cap \mathcal{C}^2(D)$ *to*

$$\begin{cases} Lu - ku + f = 0, \text{ in } D, \\ u|_{\partial D} = g, \end{cases} \tag{4.2}$$

(where L stands for the infinitesimal generator of X) and the solution is given by (4.1).

In the following we denote by $F(x)$ the signed spatial distance to the boundary ∂D. Under **(D)**, D satisfies the exterior and interior uniform sphere condition with radius $r_0 > 0$ and $F \in \mathbf{H}_2(V_{\partial D}(r_0))$ where $V_{\partial D}(r_0) := \{x \in \mathbb{R}^d : \mathbf{d}(x, \partial D) \leq r_0\}$. Also, F can be extended to a $\mathbf{H}_2$ function preserving the sign. For more details on the distance function, we refer to Appendix 14.6 in [GT98].

4.2 Tools and results

Below, we keep the previous notations concerning the Euler scheme. We also use the symbol C for nonnegative constants that may depend on D, b, σ, g, f, k but not on Δ or x. We reserve the notation c for constants also independent of D, g, f, k.

We recall a known result from Gobet and Maire [GM05] (Theorem 4.2) which provides an uniform bound for the p-th moment of τ^Δ:

$$\forall p \geq 1, \ \limsup_{\Delta\to 0} \sup_{x\in\bar{D}} \mathbb{E}_x[(\tau^\Delta)^p] < \infty. \tag{4.3}$$

Let us now state the main results of Section 2 in our current framework.

Proposition 15 (Tightness of the overshoot) *Assume* **($\mathbf{A}_\theta$-2)**, *and that D is of class* $\mathbf{H}_2$. *Then, for some $c > 0$,*

$$\sup_{\Delta>0} \mathbb{E}_x[\exp(c[\Delta^{-1/2}F^-(X^\Delta_{\tau^\Delta})]^2)] < +\infty.$$

From the proof of Theorem 3 and the estimate (4.3) we derive:

Theorem 16 (Joint limit laws associated to the overshoot) *Assume* **($\mathbf{A}_\theta$)**, *and that D is of class* $\mathbf{H}_2$. *Let φ be a continuous function with compact support. With the notation of Theorem 3, for all $x \in D,\ y \geq 0$,*

$$\mathbb{E}_x[Z^\Delta_{\tau^\Delta}\varphi(X^\Delta_{\tau^\Delta})\mathbf{1}_{F^-(X^\Delta_{\tau^\Delta})\geq y\sqrt{\Delta}}] \underset{\Delta\to 0}{\longrightarrow} \mathbb{E}_x\Big[Z_\tau\varphi(X_\tau)\Big(1 - H(y/|\nabla F\sigma(X_\tau)|)\Big)\Big].$$

4.3 Error expansion and boundary correction

For notational convenience introduce for $x \in D$,

$$u(D) = \mathbb{E}_x(g(X_\tau)Z_\tau + \int_0^\tau Z_s f(X_s)ds),$$
$$u^\Delta(D) = \mathbb{E}_x(g(X^\Delta_{\tau^\Delta})Z^\Delta_{\tau^\Delta} + \int_0^{\tau^\Delta} Z^\Delta_{\phi(s)} f(X^\Delta_{\phi(s)})ds).$$

The second quantity is well defined owing to (4.3).

Theorem 17 (First order expansion) *Under* **($\mathbf{A}_\theta$)**, **(D)**, **($\mathbf{C}_\theta$)**, *for Δ small enough and with the notation of Theorem 4*

$$\begin{aligned}&\mathrm{Err}(\Delta, g, f, k, x) = u^\Delta(D) - u(D)\\ &= c_0\sqrt{\Delta}\mathbb{E}_x(Z_\tau(\nabla u - \nabla g)(X_\tau)\cdot\nabla F(X_\tau)|\nabla F\sigma(X_\tau)|) + o(\sqrt{\Delta}).\end{aligned}$$

Define now $D^\Delta = \{x \in D : \mathbf{d}(x, \partial D) > c_0\sqrt{\Delta}|\nabla F\sigma(x)|\}$. Introduce $\hat\tau^\Delta = \inf\{t_i > 0 : X^\Delta_{t_i} \in D^\Delta\}$. Set

$$u^\Delta(D^\Delta) = \mathbb{E}_x[g(X^\Delta_{\hat\tau^\Delta})Z^\Delta_{\hat\tau^\Delta} + \int_0^{\hat\tau^\Delta} Z^\Delta_{\phi(s)} f(X^\Delta_{\phi(s)})ds].$$

One has:

Theorem 18 (Boundary correction) *Under* **($\mathbf{A}_\theta$)**, **(D)**, **($\mathbf{C}_\theta$)** *and assuming additionally $\nabla F(.)|\nabla F\sigma(.)|$ is in $\mathcal{C}^2$, then for Δ small enough one has*

$$u^\Delta(D^\Delta) - u(D) = o(\sqrt{\Delta}).$$

Note carefully that all the constants appearing in the error analysis for the parabolic case have at most linear growth w.r.t the fixed final time T. Estimate (4.3) allows to control uniformly the integrability of these constants in our current framework. Thus, since the arguments remain the same, we only give below sketches of the proofs.

Proof of Proposition 15. It is sufficient to prove that there exist constants $\tilde{c} > 0$ and C s.t. $\forall A \geq 0,\ \sup_{\Delta>0} \mathbb{P}_x[F^-(X^\Delta_{\tau^\Delta}) \geq A\Delta^{1/2}] \leq C\exp(-\tilde{c}A^2)$. Then any choice of $c < \tilde{c}$ is valid. For $x \in D$, we write

$$\begin{aligned} P &:= \mathbb{P}_x[F^-(X^\Delta_{\tau^\Delta}) \geq A\Delta^{1/2}] \\ &= \sum_{i\in\mathbf{N}^*} \mathbb{E}[\mathbf{1}_{\tau^\Delta > t_{i-1}} \mathbf{1}_{\tau^\Delta_{t_{i-1}} < t_i} \mathbb{P}[F^-(X^\Delta_{t_i}) \geq A\Delta^{1/2} | \mathcal{F}_{\tau^\Delta_{t_{i-1}}}]] \end{aligned}$$

where $\tau^\Delta_{t_{i-1}} := \inf\{s \geq t_{i-1} : X^\Delta_s \notin D\}$. From Lemma 8, we get

$$P \leq C\exp(-\tilde{c}A^2) \sum_{i\in\mathbf{N}^*} \mathbb{P}[\tau^\Delta > t_{i-1}, \tau^\Delta_{t_{i-1}} < t_i].$$

Lemma 16 from [GM04] remains valid under our current assumptions and yields

$$P \leq C\exp(-\tilde{c}A^2) \sum_{i\in\mathbf{N}^*} \mathbb{E}[\mathbf{1}_{\tau^\Delta > t_{i-1}} (\mathbb{P}[X^\Delta_{t_i} \notin D | \mathcal{F}_{t_{i-1}}] + O_{pol}(\Delta))].$$

On the one hand, $\sum_{i\in\mathbf{N}^*} \mathbf{1}_{\tau^\Delta > t_{i-1}} \mathbf{1}_{X^\Delta_{t_i} \notin D} = \mathbf{1}_{\tau^\Delta < \infty} = 1$ owing to (4.3). On the other hand, we have $\sum_{i\in\mathbf{N}^*} \mathbb{P}_x[\tau^\Delta > t_{i-1}] = \Delta^{-1}\mathbb{E}_x[\tau^\Delta] \leq C/\Delta$ using (4.3) again. Finally, we obtain that $P \leq C\exp(-\tilde{c}A^2)$ which concludes the proof. □

Proof of Theorem 17. Similarly to the proof of Theorem 6 we suppose first that $u \in \mathbf{H}_{3+\theta}$. The general case can be deduced as in the parabolic case using suitable Schauder estimates, given in the final notes of Chapter 6 in [GT98], see also our Appendix.

In this simplified setting, keeping the notations introduced in the proof of Theorem 6, we obtain

$$\begin{aligned} \mathrm{Err}(\Delta, g, f, k, x) \overset{\mathbb{E}}{=} Z^\Delta_{\tau^\Delta}(\nabla u - \nabla g)(\pi_{\partial D}(X^\Delta_{\tau^\Delta}))\nabla F(X^\Delta_{\tau^\Delta})F^-(X^\Delta_{\tau^\Delta}) \\ + \Big(\sum_{i\in\mathbf{N}} \mathbf{1}_{t_i < \tau^\Delta} \Big[\mathbf{1}_{A^\varepsilon_{t_i}} O(\Delta^{\frac{1+\theta}{2}}) \qquad (4.4) \\ + \mathbf{1}_{(A^\varepsilon_{t_i})^C} \mathbf{1}_{\forall s\in[t_i,t_{i+1}], X^\Delta_s \in B(X^\Delta_{t_i}, \Delta^{\frac{1}{2}(1-\varepsilon)})} (u(X^\Delta_{t_{i+1}})Z^\Delta_{t_{i+1}} - u(X^\Delta_{t_i})Z^\Delta_{t_i} \\ + Z^\Delta_{t_i} f(X^\Delta_{t_i})\Delta)\Big]\Big)\mathbf{1}_{\tau^{r_0} > \tau^\Delta}. \qquad (4.5) \end{aligned}$$

Since the constant in (2.6) depends linearly on time, the contribution associated to the remainder (4.4) can be bounded by $C\Delta^{\frac{3+\theta-\varepsilon}{2}} \times(\Delta^{-1}\mathbb{E}_x[\tau^\Delta])$. From (4.3), this quantity is a $O(\Delta^{\frac{1+\theta-\varepsilon}{2}}) = o(\Delta^{\frac{1}{2}})$ for ε small enough. Similarly to (2.10) the term (4.5) can be bounded by

$$\begin{aligned}
&\mathbb{E}\Big[\Big(\sum_{i\in\mathbf{N}} \mathbf{1}_{t_i<\tau^\Delta}\mathbf{1}_{(A^\varepsilon_{t_i})^C} O(\Delta^2\{1+|u|_\infty+|\nabla u|_\infty+|D^2u|_\infty+|D^3u|_\infty\} \\
&\qquad\qquad + \Delta^{\frac{3+\theta}{2}}[D^3u]_{x,\theta})\Big)\Big] + O_{pol}(\Delta) \\
&\leq C\Delta^{\frac{1+\theta}{2}}\mathbb{E}[\tau^\Delta] = o(\Delta^{1/2}).
\end{aligned}$$

We eventually derive the result as in Section 2. □

Theorem 18 can be proved as Theorem 5, using a sensitivity result analogous to Theorem 2.2 in [CGK06] for elliptic problems, see e.g. Simon [Sim80]. We skip the details.

5 Numerical results

The numerical behavior of the correction of Theorem 5 had already been illustrated for the killed case in Section 3 of [Men06]. Additional tests are presented in [Gob09]. We now focus on the stopped case with the following example. Take $d = 3$ and introduce the following diffusion process

$$\begin{aligned}
dX_t &= b(X_t)dt + \sigma(X_t)dW_t,\ \forall x\in\mathbb{R}^3,\ b(x) = (x_2\ x_3\ x_1)^*, \\
\sigma(x) &= \begin{pmatrix} (1+|x_3|)^{1/2} & 0 & 0 \\ \frac{1}{2}(1+|x_1|)^{1/2} & \left(\frac{3}{4}\right)^{1/2}(1+|x_1|)^{1/2} & 0 \\ 0 & \frac{1}{2}(1+|x_2|)^{1/2} & \left(\frac{3}{4}\right)^{1/2}(1+|x_2|)^{1/2} \end{pmatrix},
\end{aligned} \tag{5.1}$$

and X_0 to be specified later on. Set $D = B(0,2)$. We consider an elliptic problem. Starting from a given function $u(x) = x_1x_2x_3$ defined on $\bar{D}$, we derive the PDE of type (4.2) associated to (5.1) satisfied by u by taking $g = u|_{\partial D}$, setting $f = -Lu$ where L stands for the infinitesimal generator of X in (5.1) and $k = 0$. One can easily check that $-f(x) = x_2^2x_3 + x_3^2x_1 + x_1^2x_2 + \frac{1}{2}[x_3(1+|x_1|)^{1/2}(1+|x_3|)^{1/2} + x_1\left(\frac{3}{4}\right)^{1/2}(1+|x_1|)^{1/2}(1+|x_2|)^{1/2}]$. Thus we have an explicit expression for the solution of (4.2).

For x_0 s.t. $(x_0^i)_{1\leq i\leq 3} \in \{-.7,-.3,.3,.7\}$, we take $N_{MC} = 10^6$ sample paths for the Monte Carlo simulation and let Δ vary in $\{.01,.05,.1\}$. For all the computations, the size of the 95% confidence interval always varies in $[1.5\times$

$10^{-3}, 2 \times 10^{-3}]$. For the absolute value of the absolute and relative errors over the $3 \times 4^3 = 192$ points of the spatial grid, we report the results in Table 1. These results for the correction seem to indicate that the remainder $o(\Delta^{1/2})$ in Theorem 18 is actually a $O(\Delta)$. This will concern further research.

Δ	Without correction	In the corrected domain
.1	0.169 (199%)	0.0220 (24.4%)
.05	0.114 (133%)	0.0115 (13.1%)
.01	0.0471 (54.7%)	0.0026 (2.98%)

Table 1
Supremum of the absolute error for the Euler scheme (relative error in % in parenthesis)

In Tables 2 and 3, we also report the results obtained for the spatial points $x_0 = (-.7, .3, .7)$ and $x_0 = (-.7, .7, -.7)$.

Δ	Without correction	In the corrected domain
.1	-.0913+/- .0019	-.1477 +/- .0016
.05	-.1051 +/- .0018	-.1465+/- .0016
.01	-.1282 +/- .0017	-.1476+/- .0016

Table 2
Estimated value at $x_0 = (-.7, .3, .7)$ (with 95% confidence interval). True value $u(x_0) = -.147$.

Δ	Without correction	In the corrected domain
.1	.5368 +/- .0019	.3866 +/- .0016
.05	.4648 +/- .0018	.3634 +/- .0016
.01	.3851 +/- .0016	.3473 +/- .0016

Table 3
Estimated value at $x_0 = (-.7, .7, -.7)$ (with 95% confidence interval). True value $u(x_0) = .343$.

Eventually, for the Monte Carlo method, taking $x_0 = (-.7, .3, .7)$ and the previous values of Δ, in Figure 5 we plot $-\log(\mathrm{Err}_{MC})$ in function of $-\log(\Delta)$, where $\mathrm{Err}_{MC} := \left\{ \frac{1}{MC} \sum_{i=1}^{MC} \left(g(X^{\Delta,i}_{\tau^{\Delta,i}}) + \int_0^{\tau^{\Delta,i}} f(X^{\Delta,i}_{\phi(s)}) ds \right) \right\} - u(x_0)$. The curve is quite close to a right line with slope 1/2 as it should from Theorem 17.



Figure 5. Error for the Monte Carlo method (without correction) as a function of Δ, in logarithmic scales. Evaluation at $x_0 = (-.7, .3, .7)$.

6 Conclusion

We have proposed and analysed a boundary correction procedure to simulate stopped/killed diffusion processes. This is valid for non-stationary and stationary problems, in time-dependent or time-independent domains. The resulting scheme is elementary to implement and its numerical accuracy is very good in our experiments. The proof relies on new asymptotic results regarding the renormalized overshoots.

To conclude, we note that the boundary correction procedure is very generic and could be at least formally extended to general It processes of the form $dX_t = b_t dt + \sigma_t dW_t$. In that case, the smaller domain would be defined ω by ω replacing $\nabla F(t,x)\sigma(t,x)$ by $\nabla F(t,X_t)\sigma_t$. Even if our current proof relies on Markovian properties, we conjecture that the correction should once again give a $o(\sqrt{\Delta})$ independently of the Markovian structure. Numerical tests in [Gob09] support this conjecture, which will be addressed mathematically in further research.

A Proof of Theorem 6 in the general setting

In this section, we detail how the proof of Section 2 has to be modified under the assumptions of Theorem 4, i.e. for $g \in \mathbf{H}_{1+\theta}$ and without compatibility condition so that $u \in \mathbf{H}_{1+\theta}$. Actually, u is smooth inside the domain but high order derivatives may explode close to the boundary. These features have to be accurately quantified to show that the induced singularities are integrable.

Introduce the parabolic distance $\mathbf{pd}$: for $(s,x),(t,y)\in\bar{\mathcal{D}}$, $\mathbf{pd}((s,x),(t,y)) = \max(|s-t|^{1/2},|x-y|)$. We also denote for a closed set $\mathcal{A}\in\bar{\mathcal{D}}$ and $(s,x)\in\mathcal{D}$, $\mathbf{pd}((s,x),\mathcal{A})$ the parabolic distance of (s,x) to $\mathcal{A}$. Note that $\mathbf{pd}((s,x),\mathcal{PD}\cap\{v\geq s\})\geq\min(F(s,x),\sqrt{T-s})$, so that we obtain the easy inequality:

$$\frac{1}{\mathbf{pd}((s,x),\mathcal{PD}\cap\{v\geq s\})}\leq\frac{1}{F(s,x)}+\frac{1}{\sqrt{T-s}}. \tag{A.1}$$

Under our current assumptions, for some constant $C>0$, we have

$$|D^2u(s,x)|+|D^3u(s,x)|\leq C\mathbf{pd}((s,x),\mathcal{PD}\cap\{v\geq s\})^{-2}; \tag{A.2}$$

$$\text{for } (t,y)\neq(s,x),\quad \frac{|D^3u(s,x)-D^3u(t,y)|}{\mathbf{pd}((s,x),(t,y))^{\theta}}$$
$$\leq C[\mathbf{pd}((s,x),\mathcal{PD}\cap\{v\geq s\})\wedge\mathbf{pd}((t,y),\mathcal{PD}\cap\{v\geq t\})]^{-2-\theta}; \tag{A.3}$$

$$\text{for } t\neq s,\quad \frac{|D^2u(s,x)-D^2u(t,x)|}{|t-s|^{(1+\theta)/2}}$$
$$\leq C[\mathbf{pd}((s,x),\mathcal{PD}\cap\{v\geq s\})\wedge\mathbf{pd}((t,x),\mathcal{PD}\cap\{v\geq t\})]^{-2-\theta}. \tag{A.4}$$

The above constant C is uniform w.r.t. $(s,x)\in\mathcal{D},(t,y)\in\mathcal{D}$ or $(t,x)\in\mathcal{D}$. These inequalities are obtained with the interior Schauder estimates for the PDEs satisfied by the partial derivatives $(\partial_{x_i}u)_{1\leq i\leq d}$, see Theorem 4.9 in [Lie96].

We first state an important proposition for the error analysis with possibly explosive controls as in (A.2)-(A.3)-(A.4) for the derivatives. Namely, under our current regularity assumptions, in order to perform a Taylor expansion we have to work with interior points located in small balls, which distance to the boundary is uniformly bounded from below within the ball. The next proposition states that this is the case if the ball centers are "far enough" from the side of $\mathcal{D}$.

Proposition 19 *Assume $\mathcal{D}\in\mathbf{H}_2$ and take $\varepsilon\in]0,1[$. For all $(t,x)\in\bar{\mathcal{D}}\cap V_{\partial\mathcal{D}}(r_0/2)\backslash V_{\partial\mathcal{D}}(2\Delta^{1/2(1-\varepsilon)})$ (r_0 is defined in Section 1.5.2), one has for $\forall y\in B(x,\Delta^{1/2(1-\varepsilon)})$ and $s\in[t,t+\Delta]$*

$$F(s,y)\geq\frac{1}{4}F(t,x)$$

for Δ small enough (uniformly in t,x,s,y). In particular, y belongs to D_s.

Proof. Since $F\in\mathbf{H}_2$, one has

$$F(s,y)\geq F(t,x)-C\Delta+\langle\nabla F(t,x),y-x\rangle-C\Delta^{1-\varepsilon}.$$

The norm of $\nabla F(t,x)$ equals 1, since $\nabla F(t,x)$ is the unit inward normal vector at the closest point of x on ∂D_t. Therefore, for Δ small enough and using $\frac{1}{2}F(t,x) \geq \Delta^{\frac{1}{2}(1-\varepsilon)}$, we have

$$F(s,y) \geq F(t,x) - \frac{3}{2}\Delta^{\frac{1}{2}(1-\varepsilon)} \geq \frac{1}{4}F(t,x),$$

which is the expected inequality. □

We are now in a position to deduce useful local upper bounds for the derivatives of u and their Hölder-norms, under the assumptions of Theorem 6.

Corollary 20 *Take $\varepsilon \in]0,1[$. There exists a constant $C > 0$ such that for Δ small enough, for all $(t,x) \in \bar{\mathcal{D}} \backslash V_{\partial\mathcal{D}}(2\Delta^{1/2(1-\varepsilon)})$, for all $(y,z) \in B(x, \Delta^{1/2(1-\varepsilon)})$ and $(r,s) \in [t, t+\Delta]$, we have*

$$|D^2u(s,y)| + |D^3u(s,y)| \leq \frac{C}{F^2(t,x)} + \frac{C}{T-t}; \tag{A.5}$$

$$\text{for } y \neq z, \quad \frac{|D^3u(s,y) - D^3u(s,z)|}{|y-z|^\theta} \leq \frac{C}{F^{2+\theta}(t,x)} + \frac{C}{(T-t)^{1+\theta/2}}; \tag{A.6}$$

$$\text{for } r \neq s, \quad \frac{|D^2u(r,y) - D^2u(s,y)|}{|r-s|^{(1+\theta)/2}} \leq \frac{C}{F^{2+\theta}(t,x)} + \frac{C}{(T-t)^{1+\theta/2}}. \tag{A.7}$$

Proof. Note that if $(t,x) \in \bar{\mathcal{D}} \backslash V_{\partial\mathcal{D}}(2\Delta^{1/2(1-\varepsilon)})$, we have $T - t \geq 4\Delta^{1-\varepsilon}$.
Estimate (A.5). In view of (A.2) and (A.1), the upper bound of $|D^2u(s,y)| + |D^3u(s,y)|$ is equal to $\frac{C}{F^2(s,y)} + \frac{C}{T-s}$. On the one hand, by easy computations, we prove

$$\frac{1}{T-s} \leq \frac{1}{T-t}\frac{T-t}{T-t-\Delta} \leq \frac{1}{T-t}\frac{1}{1-\Delta^\varepsilon/4} \leq \frac{C}{T-t}$$

for Δ small enough. On the other hand, we have

$$\frac{1}{F(s,y)} \leq \frac{C}{F(t,x)}.$$

Indeed, if x is far from D_t (and thus y far from D_s), both terms $F(s,y)$ and $F(t,x)$ are bounded from above and from below. In the other case when $(t,x) \in \bar{\mathcal{D}} \cap V_{\partial\mathcal{D}}(r_0/2) \backslash V_{\partial\mathcal{D}}(2\Delta^{1/2(1-\varepsilon)})$, Proposition 19 yields $\frac{F(s,y)}{F(t,x)} \geq \frac{1}{4}$. Therefore, the upper bound (A.5) readily follows.
Estimates (A.6) and (A.7). They are proved following the same arguments, the details of which are left to the reader. □

Recall from the previous proof of Theorem 6 that the main term to analyse is

$$
\begin{aligned}
e_{22}^{\Delta} &\overset{\mathbb{E}}{=} \Big(\sum_{0\le t_i<T} \mathbf{1}_{t_i<\tau^{\Delta}} \mathbf{1}_{(A_{t_i}^{\varepsilon})^C} \mathbf{1}_{\forall s\in[t_i,t_{i+1}],\ X_s^{\Delta}\in B(X_{t_i}^{\Delta},\Delta^{\frac{1}{2}(1-\varepsilon)})} \Big[u(t_{i+1},X_{t_{i+1}}^{\Delta})Z_{t_{i+1}}^{\Delta} \\
&\qquad -u(t_i,X_{t_i}^{\Delta})Z_{t_i}^{\Delta} + Z_{t_i}^{\Delta} f(t_i,X_{t_i}^{\Delta})\Delta \Big]\Big) \mathbf{1}_{\tau^{r_0}>\tau^{\Delta}\wedge T} \\
&= \Big(\sum_{0\le t_i<T-4\Delta^{1-\varepsilon}} \cdots \Big) \mathbf{1}_{\tau^{r_0}>\tau^{\Delta}\wedge T} + \Big(\sum_{T-4\Delta^{1-\varepsilon}\le t_i<T} \cdots \Big) \mathbf{1}_{\tau^{r_0}>\tau^{\Delta}\wedge T} := e_{221}^{\Delta} + e_{222}^{\Delta},
\end{aligned}
$$

where we have just splitted the summation on t_i.

Control of e_{221}^{Δ}. The idea is to perform a stochastic expansion of $u(t_{i+1},X_{t_{i+1}}^{\Delta})Z_{t_{i+1}}^{\Delta} - u(t_i,X_{t_i}^{\Delta})Z_{t_i}^{\Delta} + Z_{t_i}^{\Delta} f(t_i,X_{t_i}^{\Delta})\Delta$ as in (2.8). Under our current assumptions, the difference comes from the high-order derivatives that are no more uniformly bounded or uniformly Hölder but only locally, with local estimates given in Corollary 20. Thus, following the same computations that have led to (2.10), we obtain

$$
\begin{aligned}
e_{221}^{\Delta} \overset{\mathbb{E}}{=} \Big(&\sum_{0\le t_i<T-4\Delta^{1-\varepsilon}} \mathbf{1}_{t_i<\tau^{\Delta}} \mathbf{1}_{(A_{t_i}^{\varepsilon})^C} \mathbf{1}_{\forall s\in[t_i,t_{i+1}],\ X_s^{\Delta}\in B(X_{t_i}^{\Delta},\Delta^{\frac{1}{2}(1-\varepsilon)})} \Big[\\
&O\Big((\Delta^2 + \Delta|X_s^{\Delta} - X_{t_i}^{\Delta}|^2)\big(\frac{1}{F^2(t_i,X_{t_i}^{\Delta})} + \frac{1}{T-t_i}\big)\Big) \\
&+ O\Big((\Delta^{1+\frac{1+\theta}{2}} + \Delta|X_s^{\Delta} - X_{t_i}^{\Delta}|^{1+\theta})\big(\frac{1}{F^{2+\theta}(t_i,X_{t_i}^{\Delta})} + \frac{1}{(T-t_i)^{1+\theta/2}}\big)\Big) \\
&+ \bar{M}_{t_i,t_{i+1}} \Big]\Big) \mathbf{1}_{\tau^{r_0}>\tau^{\Delta}\wedge T}. \qquad \text{(A.8)}
\end{aligned}
$$

The derivatives appearing in $(\bar{M}_{t_i,t_{i+1}})_{0\le t_i<T}$ (see equations (2.9) and (2.10)) are controlled by (A.5) on $(A_{t_i}^{\varepsilon})^C$. The control of (2.11) remains valid for the $(\bar{M}_{t_i,t_{i+1}})_{0\le t_i<T}$ that yields a negligeable contribution. It follows that

$$
\begin{aligned}
|e_{221}^{\Delta}| \overset{\mathbb{E}}{\le} C\Delta^{\frac{1+\theta}{2}} \Big(&\sum_{0\le t_i<T-4\Delta^{1-\varepsilon}} \mathbf{1}_{t_i<\tau^{\Delta}} \mathbf{1}_{F(t_i,X_{t_i}^{\Delta})\ge 2\Delta^{\frac{1}{2}(1-\varepsilon)}} \Delta \Big[\\
&\frac{1}{F^{2+\theta}(t_i,X_{t_i}^{\Delta})} + \frac{1}{(T-t_i)^{1+\theta/2}} \Big]\Big) \mathbf{1}_{\tau^{r_0}>\tau^{\Delta}\wedge T}.
\end{aligned}
$$

Standard computations show that

$$
\Delta^{\frac{1+\theta}{2}} \sum_{0\le t_i<T-4\Delta^{1-\varepsilon}} \frac{\Delta}{(T-t_i)^{1+\theta/2}} \le \Delta^{\frac{1+\theta}{2}} \int_0^{T-4\Delta^{1-\varepsilon}+\Delta} \frac{dt}{(T-t)^{1+\theta/2}} = O(\Delta^{\frac{1}{2}+\frac{\theta\varepsilon}{2}}),
$$

which implies

$$
|e_{221}^{\Delta}| \overset{\mathbb{E}}{\le} C\Delta^{\frac{1+\theta}{2}} \left(\int_0^{T\wedge\tau^{\Delta}} \mathbf{1}_{F(\phi(t),X_{\phi(t)}^{\Delta})\in[2\Delta^{1/2(1-\varepsilon)},r_0/2]} F(\phi(t),X_{\phi(t)}^{\Delta})^{-2-\theta} dt \right) + O(\Delta^{\frac{1}{2}+\frac{\theta\varepsilon}{2}}).
$$

Adapting the previous analysis of Section 2 for the term e^{Δ}_{211}, we get

$$
\begin{aligned}
|e^{\Delta}_{221}| &\overset{\mathbb{E}}{\leq} C\Delta^{\frac{1+\theta}{2}}\left(\int_0^{T\wedge\tau^{\Delta}} \mathbf{1}_{F(t,X^{\Delta}_t)\in[\Delta^{1/2(1-\varepsilon)},3r_0/4]} F(t,X^{\Delta}_t)^{-2-\theta}dt\right) + O(\Delta^{\frac{1}{2}+\frac{\theta\varepsilon}{2}}) \\
&\overset{\mathbb{E}}{\leq} C\Delta^{\frac{1+\theta}{2}}\left(\int_{\Delta^{1/2(1-\varepsilon)}}^{3r_0/4} y^{-2-\theta} L^{y}_{T\wedge\tau^{\Delta}}(F(.,X^{\Delta}_{.}))dy\right) + O(\Delta^{\frac{1}{2}+\frac{\theta\varepsilon}{2}}),
\end{aligned}
$$

using Lemma 8 for the last but one inequality, and the occupation time formula for $F(t,X^{\Delta}_t)$ for the last one (recall that σ is uniformly elliptic).
Finally using (2.6), one gets

$$
|e^{\Delta}_{221}| \leq C\Delta^{\frac{1+\theta}{2}}\left(\int_{\Delta^{1/2(1-\varepsilon)}}^{3r_0/4} y^{-2-\theta}(y+\Delta^{1/2})dy\right) + O(\Delta^{\frac{1}{2}+\frac{\theta\varepsilon}{2}}) \leq C\Delta^{\frac{1}{2}+\frac{\theta\varepsilon}{2}} = o(\Delta^{1/2}).
$$

Control of e^{Δ}_{222}. Apply a Taylor formula with integral rest at order one in space. The θ-Hölder continuity in space of ∇u and the $(1+\theta)/2$-Hölder continuity in time of u directly give a contribution in $O(\Delta^{1/2+\theta/2-\varepsilon}) = o(\Delta^{1/2})$ for ε small enough. This completes the proof. □

References

[ABR96] L. Andersen and R. Brotherton-Ratcliffe. Exact exotics. *Risk*, 9:85–89, 1996.

[AGP95] S. Asmussen, P. Glynn, and J. Pitman. Discretization error in simulation of one-dimensional reflecting Brownian motion. *Ann. Appl. Probab.*, 5(4):875–896, 1995.

[Als94] G. Alsmeyer. On the Markov renewal theorem. *Stoch. Proc. Appl.*, 50(1):37–56, 1994.

[Avi07] R. Avikainen. Convergence rates for approximations of functionals of SDEs. *Submitted, available on http://arxiv.org/abs/0712.3635*, 2007.

[Bal95] P. Baldi. Exact asymptotics for the probability of exit from a domain and applications to simulation. *Ann. Prob.*, 23(4):1644–1670, 1995.

[BGK97] M. Broadie, P. Glasserman, and S. Kou. A continuity correction for discrete barrier options. *Mathematical Finance*, 7:325–349, 1997.

[BL94] P.P. Boyle and S.H. Lau. Bumping up against the barrier with the binomial method. *Jour. of Derivat.*, 1:6–14, 1994.

[CGK06] C. Costantini, E. Gobet, and N. El Karoui. Boundary sensitivities for diffusion processes in time dependent domains. *Appl. Math. Optim.*, 54–2:159–187, 2006.

[DK06] R. A. Doney and A. E. Kyprianou. Overshoots and undershoots of Lévy processes. *Ann. Appl. Probab.*, 16–1:91–106, 2006.

[DL06] M. Deaconu and A. Lejay. A random walk on rectangles algorithm. *Methodology And Computing In Applied Probability*, 8(1):135–151, 2006.

[FL01] C.D. Fuh and T.L. Lai. Asymptotic expansions in multidimensional Markov renewal theory and first passage times for Markov random walks. *Adv. in Appl. Probab.*, 33(3):652–673, 2001.

[Fre85] M. Freidlin. *Functional integration and Partial differential equations.* Annals of Mathematics studies, Princeton University Press, 1985.

[Fri64] A. Friedman. *Partial differential equations of parabolic type.* Prentice-Hall, 1964.

[GM04] E. Gobet and S. Menozzi. Exact approximation rate of killed hypoelliptic diffusions using the discrete Euler scheme. *Stoch. Proc. and Appl.*, 112:210–223, 2004.

[GM05] E. Gobet and S. Maire. Sequential Control Variates for Functionals of Markov Processes. *Siam. Journal of Num. Analysis*, 43-3:1256–1275, 2005.

[GM07] E. Gobet and S. Menozzi. Discrete sampling of functionals of Itô processes. *Séminaire de Probabilités*, XL:355–375, 2007.

[Gob00] E. Gobet. Euler schemes for the weak approximation of killed diffusion. *Stoch. Proc. Appl.*, 87:167–197, 2000.

[Gob09] E. Gobet. *Handbook of Numerical Analysis, Vol. XV, Special Volume: Mathematical Modeling and Numerical Methods in Finance*, chapter Advanced Monte Carlo methods for barrier and related exotic options, pages 497–528. Elsevier, Netherlands: North-Holland, 2009.

[GT98] D. Gilbarg and N.S. Trudinger. *Elliptic partial differential equations of second order.* Springer Verlag, 1998.

[Lie96] G.M. Lieberman. *Second Order parabolic differential equations, 1st edn.* World Scientfic, River Edge, NJ, 1996.

[Men06] S. Menozzi. Improved simulation for the killed Brownian motion in a cone. *Siam Jour. Num. Anal.*, 44-6:2610–2632, 2006.

[Mil97] G. N. Milstein. Weak approximation of a diffusion process in a bounded domain. *Stoch. Stoch. Reports*, 64:211–233, 1997.

[MT99] G. N. Milstein and M. V. Tretyakov. Simulation of a space-time bounded diffusion. *Ann. Appl. Prob.*, 9–3:732–779, 1999.

[RY99] D. Revuz and M. Yor. *Continuous martingales and Brownian motion. 3rd ed.* Grundlehren der Mathematischen Wissenschaften. 293. Berlin: Springer, 1999.

[Sie79] D. Siegmund. Corrected diffusion approximations in certain random walk problems. *Adv. in Appl. Probab.*, 11(4):701–719, 1979.

[Sie85] D. Siegmund. *Sequential Analysis.* Springer, 1985.

[Sim80] J Simon. Differentiation with respect to the domain in boundary value problems. *Numer. Funct. Anal. Optim.*, 7–8:649–687, 1980.

[SY82] D. Siegmund and Y.S. Yuh. Brownian approximations for first passage probabilities. *Z. Wahrsch. verw. Gebiete*, 59:239–248, 1982.

[TT90] D. Talay and L. Tubaro. Expansion of the global error for numerical schemes solving stochastic differential equations. *Stoch. Anal. and App.*, 8-4:94–120, 1990.

[Woo82] M. Woodroofe. *Nonlinear renewal theory in sequential analysis.* SIAM, 1982.

[Zha88] C.H. Zhang. A nonlinear renewal theory. *Ann. Prob.*, 16-2:793–824, 1988.