



# Heisenberg Uncertainty Principle for the q-Bessel Fourier transform

Lazhar Dhaouadi

## ► To cite this version:

| Lazhar Dhaouadi. Heisenberg Uncertainty Principle for the q-Bessel Fourier transform. 2007. hal-00161302v1

HAL Id: hal-00161302

<https://hal.science/hal-00161302v1>

Preprint submitted on 10 Jul 2007 (v1), last revised 1 Jul 2008 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Heisenberg Uncertainty Principle for the $q$ -Bessel Fourier transform

Lazhar Dhaouadi \*

## Abstract

In this paper we give a  $q$ -analogue of The Heisenberg Uncertainty Principle for the  $q$ -Bessel Fourier transform:

$$\mathcal{F}_{q,v}f(x) = c_{q,v} \int_0^\infty f(t)j_v(xt, q^2)t^{2v+1}d_q t,$$

where  $j_v(x, q)$  is the normalized Hahn-Exton  $q$ -Bessel function.

## 1 The $q$ -Fourier Bessel transform

Throughout this paper we consider  $0 < q < 1$  and we adopt the standard conventional notations of [2]. We put

$$\mathbb{R}_q^+ = \{q^n, \quad n \in \mathbb{Z}\},$$

and for complex  $a$

$$(a; q)_0 = 1, \quad (a; q)_n = \prod_{i=0}^{n-1} (1 - aq^i), \quad n = 1 \dots \infty.$$

Jackson's  $q$ -integral (see [3]) in the interval  $[0, \infty[$  is defined by

$$\int_0^\infty f(x)d_q x = (1 - q) \sum_{n=-\infty}^{\infty} q^n f(q^n).$$

---

\*Institut Préparatoire aux Etudes d'Ingénieur de Bizerte (Université du 7 novembre Carthage). Route Menzel Abderrahmene Bizerte, 7021 Zarzouna, Tunisia.  
E-mail lazhardhaouadi@yahoo.fr

We introduce the following functional spaces  $\mathcal{L}_{q,p,v}$  of even functions  $f$  defined on  $\mathbb{R}_q^+$  such that

$$\|f\|_{q,p,v} = \left[ \int_0^\infty |f(x)|^p x^{2v+1} d_q x \right] < \infty.$$

The normalized Hahn-Exton q-Bessel function of order  $v > -1$  (see [5]) is defined by

$$j_v(z, q) = \sum_{n=0}^{\infty} (-1)^n \frac{q^{\frac{n(n-1)}{2}}}{(q, q)_n (q^{v+1}, q)_n} z^n,$$

and the q-Bessel Fourier transform  $\mathcal{F}_{q,v}$  introduced in [1,4] as follow

$$\mathcal{F}_{q,v} f(x) = c_{q,v} \int_0^\infty f(t) j_v(xt, q^2) t^{2v+1} d_q t,$$

where

$$c_{q,v} = \frac{1}{1-q} \frac{(q^{2v+2}, q^2)_\infty}{(q^2, q^2)_\infty}.$$

Define the  $q$ -Bessel translation operator as follows:

$$T_{q,x}^v f(y) = c_{q,v} \int_0^\infty \mathcal{F}_{q,v}(f)(t) j_v(xt, q^2) j_v(yt, q^2) t^{2v+1} d_q t, \quad \forall x, y \in \mathbb{R}_q^+, \forall f \in \mathcal{L}_{q,v,1}.$$

The  $q$ -convolution product of both functions  $f, g \in \mathcal{L}_{q,v,1}$  is defined by

$$f *_q g(x) = c_{q,v} \int_0^\infty T_{q,x}^v f(y) g(y) y^{2v+1} d_q y.$$

The followings results was proved in [1].

**Proposition 1** *For any function  $f \in \mathcal{L}_{q,v,1}$  we have*

$$\int_0^\infty T_{q,x}^v f(y) y^{2v+1} d_q y = \int_0^\infty f(y) y^{2v+1} d_q y.$$

**Proposition 2** *Given two functions  $f, g \in \mathcal{L}_{q,v,1}$  then*

$$f *_q g \in \mathcal{L}_{q,v,1},$$

and

$$\mathcal{F}_{q,v}(f *_q g) = \mathcal{F}_{q,v}(f) \mathcal{F}_{q,v}(g).$$

### Theorem 1

1. The  $q$ -Bessel Fourier transform  $\mathcal{F}_{q,v}$  possess an extension

$$U : \mathcal{L}_{q,2,v} \rightarrow \mathcal{L}_{q,2,v}.$$

2. For  $f \in \mathcal{L}_{q,2,v}$ , we have

$$\|U(f)\|_{q,2,v} = \|f\|_{q,2,v}.$$

3. The application  $U$  is bijective and

$$U^{-1} = U.$$

**Proposition 3** Given  $1 < p \leq 2$  and  $\frac{1}{p} + \frac{1}{\bar{p}} = 1$ . If  $f \in \mathcal{L}_{q,p,v}$ , then

$$\mathcal{F}_{q,v}(f) \in \mathcal{L}_{\bar{p},2,v}$$

and

$$\|\mathcal{F}_{q,v}(f)\|_{q,\bar{p},v} \leq B_{q,v}^{(\frac{2}{p}-1)} \|f\|_{q,p,v},$$

where

$$B_{q,v} = \frac{1}{1-q} \frac{(-q^2; q^2)_\infty (-q^{2v+2}; q^2)_\infty}{(q^2; q^2)_\infty}.$$

The  $q$ -exponential function is defined by

$$e(z, q) = \sum_{n=0}^{\infty} \frac{z^n}{(q, q)_n} = \frac{1}{(z; q)_\infty}, \quad |z| < 1.$$

**Proposition 4** The  $q$ -Gauss kernel

$$G^v(x, t, q^2) = \frac{(-q^{2v+2}t, -q^{-2v}/t; q^2)_\infty}{(-t, -q^2/t; q^2)_\infty} e\left(-\frac{q^{-2v}}{t}x^2, q^2\right),$$

satisfying

$$\mathcal{F}_{q,v} \{e(-ty^2, q^2)\}(x) = G^v(x, t, q^2),$$

and for all function  $f \in \mathcal{L}_{q,2,v}$

$$\lim_{n \rightarrow \infty} \|c_{q,v} G^v(x, q^{2n}, q^2) *_q f - f\|_{q,2,v} = 0.$$

## 2 Uncertainty Principale

The following theorem is Jensens inequality and is crucial for the proof of our main result.

**Theorem 2** *Let  $\mu$  be a probability measure on  $\mathbb{R}_q^+$ , If  $f$  is a positive function in  $\mathcal{L}_{q,v,1}$  and  $g$  is convex on  $\mathbb{R}^+$  then*

$$g\left(\int_0^\infty f d\mu(x)\right) \leq \int_0^\infty g \circ f(x) d\mu(x).$$

**Proof.** There exists  $c \in \mathbb{R}$  such that for all  $x, y$  in a convex set  $\mathbb{R}^+$  it holds

$$g(x) \geq g(y) + c(x - y).$$

Let  $t = \int_0^\infty f(x) d\mu(x)$ . Then  $t \in \mathbb{R}^+$ . Now let  $x \in \mathbb{R}_q^+$  then

$$g(f(x)) \geq g(t) + c(f(x) - t).$$

Integrating both sides and using the special value of  $t$  gives

$$\int_0^\infty g(f(x)) d\mu(x) \geq \int_0^\infty [g(t) + c(f(x) - t)] d\mu(x) = g(t).$$

This finish the proof. ■

Now we are in a position to state and prove the q-analogue of the Uncertainty Principale.

**Theorem 3** *Given  $a, b > 0$  and let*

$$\sigma_a = \int_0^\infty \exp(-|x|^a) x^{2v+1} d_q x,$$

*then for all  $c, d \in \mathbb{R}_q^+$  satisfying*

$$0 < B_{q,v} \frac{\sigma_a \sigma_b}{(cd)^{2v+2}} < 1,$$

*the following hold for any function  $f \in \mathcal{L}_{q,2,v}$*

$$c^a \|x^{a/2} f\|_{q,v,2}^2 + d^b \|x^{b/2} \mathcal{F}_{q,v} f\|_{q,v,2}^2 \geq -\log \left( B_{q,v} \frac{\sigma_a \sigma_b}{(cd)^{2v+2}} \right) \|f\|_{q,v,2}^2.$$

**Proof.** For a positive function  $\phi \in \mathcal{L}_{q,v,1}$  with  $\|\phi\|_{q,v,1} = 1$  define the entropy of  $\phi$  to be

$$E(\phi) = \int_0^\infty \phi(x) \log \phi(x) x^{2v+1} d_q x,$$

and for some  $a > 0$  defining

$$M_a(\phi) = \left( \int_0^\infty |x|^a \phi(x) x^{2v+1} d_q x \right)^{\frac{1}{a}}.$$

In the following assume that  $E(\phi)$  and  $M_a(\phi)$  exist. For a given  $c \in \mathbb{R}_q^+$  let

$$d\gamma(x) = k_c^{-1} \exp(-|cx|^a) x^{2v+1} d_q x \text{ where } k_c = \frac{\sigma_a}{c^{2v+2}}.$$

Then  $d\gamma(x)$  is a probability measure. Defining

$$\psi(x) = k_c \exp(|cx|^a) \phi(x),$$

we get

$$\int_0^\infty \psi(x) d\gamma(x) = 1.$$

According to the fact that  $t \mapsto t \log t$  is convex on  $\mathbb{R}^+$ , so Jensen's inequality gives

$$0 = \int_0^\infty \psi(x) d\gamma(x) \log \left[ \int_0^\infty \psi(x) d\gamma(x) \right] \leq \int_0^\infty \psi(x) \log \psi(x) d\gamma(x).$$

$$\begin{aligned} 0 &\leq \int_0^\infty \phi(x) \log [k_c \exp(|cx|^a) \phi(x)] \cdot x^{2v+1} d_q x \\ &= \int_0^\infty \phi(x) [\log k_c + |cx|^a + \log \phi(x)] x^{2v+1} d_q x. \end{aligned}$$

$$0 \leq \log k_c + c^a \int_0^\infty |x|^a \phi(x) x^{2v+1} d_q x + \int_0^\infty \phi(x) \log \phi(x) x^{2v+1} d_q x.$$

In the end

$$0 \leq \log k_c + c^a M_a(\phi) + E(\phi). \quad (1)$$

For  $f \in \mathcal{L}_{q,v,2}$  with  $\|f\|_{q,v,2} = 1$  we will show that

$$E(|f|^2) + E(|\mathcal{F}_{q,v}f|^2) \leq \log B_{q,v}. \quad (2)$$

Start by assuming that

$$f \in \mathcal{L}_{q,v,1} \cap \mathcal{L}_{q,v,2}.$$

Then  $f$  will be in  $\mathcal{L}_{q,v,p}$  for  $1 < p < 2$ . With

$$\frac{1}{p} + \frac{1}{\bar{p}} = 1,$$

Hausdorff-Young's inequality (Proposition 3) tells us that  $\mathcal{F}_{q,v}f$  is in  $\mathcal{L}_{q,v,\bar{p}}$ . So we can define the functions

$$A(p) = \int_0^\infty |f(x)|^p dx \quad \text{and} \quad B(\bar{p}) = \int_0^\infty |\mathcal{F}_{q,v}f(x)|^{\bar{p}} x^{2v+1} d_q x.$$

Now define

$$\begin{aligned} C(p) &= \log \|\mathcal{F}_{q,v}f\|_{\bar{p}} - \log \left( B_{q,v}^{\frac{2}{\bar{p}}-1} \|f\|_p \right) \\ &= \frac{1}{\bar{p}} \log B(\bar{p}) - \frac{1}{p} \log A(p) - \left( \frac{2}{p} - 1 \right) \log B_{q,v}. \end{aligned}$$

By Hausdorff-Young's inequality

$$C(p) \leq 0, \quad \text{for } 1 < p < 2,$$

and by Parsevals equality (Theorem 1)

$$C(2) = 0.$$

In the other hand

$$C'(p) = \frac{\bar{p}'}{\bar{p}} \frac{B'(\bar{p})}{B(\bar{p})} - \frac{\bar{p}'}{\bar{p}^2} \log B(\bar{p}) - \frac{p'}{p} \frac{A'(p)}{A(p)} + \frac{p'}{p^2} \log A(p) + \frac{2}{p^2} \log B_{q,v}.$$

The derivative of  $\bar{p}$  with respect to  $p$  is

$$\bar{p}' = -\frac{1}{(p-1)^2}.$$

For a given  $x > 0$  the function  $p \mapsto \frac{x^p - x^2}{p-2}$  is increasing and

$$\lim_{p \rightarrow 2^-} \frac{x^p - x^2}{p-2} = x^2 \log x.$$

So

$$C'(2^-) = \lim_{p \rightarrow 2^-} \frac{C(p) - C(2)}{p-2} = -\frac{1}{2} [A'(2^-) + B'(2^+)] + \frac{1}{2} \log B_{q,v},$$

where

$$A'(2^-) = \lim_{p \rightarrow 2^-} \frac{A(p) - A(2)}{p-2} = E(|f|^2),$$

$$B'(2^+) = \lim_{\bar{p} \rightarrow 2^+} \frac{B(\bar{p}) - B(2)}{\bar{p}-2} = E(|\mathcal{F}_{q,v} f|^2).$$

Therefore

$$C'(2^-) \geq 0 \Rightarrow A'(2^-) + B'(2^+) - \log B_{q,v} \leq 0,$$

and then

$$E(|f|^2) + E(|\mathcal{F}_{q,v} f|^2) \leq \log B_{q,v}. \quad (3)$$

If  $\|f\|_{q,v,2} \neq 1$  then by replacing  $f$  by  $\frac{f}{\|f\|_2}$  the inequality (3) become

$$E(|f|^2) + E(|\mathcal{F}_{q,v} f|^2) \leq \|f\|_2^2 \log B_{q,v} + 2 \log \|f\|_{q,v,2}. \quad (4)$$

Let us now assume that  $f \in \mathcal{L}_{q,v,2}$  and that  $E(|f|^2)$  and  $E(|\mathcal{F}_{q,v} f|^2)$  are defined, and then approximate  $f$  by functions in  $\mathcal{L}_{q,v,1} \cap \mathcal{L}_{q,v,2}$ . Let

$$h_n(x) = c_{q,v} e(-q^{2n}x^2, q^2).$$

The following holds by (4)

$$E(|h_n f|^2) + E(|\mathcal{F}_{q,v} h_n *_q \mathcal{F}_{q,v} f|^2) \leq \|h_n f\|_2^2 \log B_{q,v} + 2 \log \|h_n f\|_{q,v,2}, \quad (5)$$

and by Proposition 2 and Proposition 4

$$\lim_{n \rightarrow \infty} \|h_n f - f\|_{q,v,2} = \lim_{n \rightarrow \infty} \|\mathcal{F}_{q,v} h_n *_q \mathcal{F}_{q,v} f - \mathcal{F}_{q,v} f\|_{q,v,2} = 0,$$

which implies

$$\lim_{n \rightarrow \infty} E(|h_n f|^2) = E(|f|^2). \quad (6)$$

The functions

$$\phi_1(z) = |z|^2 \log^+ |z| \text{ and } \phi_2(z) = |z|^2 \left( -\log^- |z| + \frac{3}{2} \right),$$

are convex on  $\mathbb{C}$ , where

$$\log^+ x = \max \{0, \log x\} \text{ and } \log^- x = \min \{0, \log x\}.$$

Since  $\mathcal{F}_{q,v} h_n \geq 0$  and by proposition 1

$$\int_0^\infty T_{q,x} \mathcal{F}_{q,v} h_n(t) t^{2v+1} d_q t = \int_0^\infty \mathcal{F}_{q,v} h_n(t) t^{2v+1} d_q t = 1,$$

for a given  $x \in \mathbb{R}_q^+$  the following holds by Jensen's Inequality

$$\begin{aligned} \phi_i(\mathcal{F}_{q,v} h_n *_q \mathcal{F}_{q,v} f)(x) &\leq \int_0^\infty \phi_i(\mathcal{F}_{q,v} f)(t) T_{q,x} \mathcal{F}_{q,v} h_n(t) t^{2v+1} d_q t \\ &= \mathcal{F}_{q,v} h_n *_q \phi_i(\mathcal{F}_{q,v} f)(x). \end{aligned}$$

It follows that

$$\mathcal{F}_{q,v} h_n *_q \phi_i(\mathcal{F}_{q,v} f) \in \mathcal{L}_{q,v,1}$$

since  $\phi_i(\mathcal{F}_{q,v} f)$  is in  $\mathcal{L}_{q,v,1}$  and therefore

$$\phi_i(\mathcal{F}_{q,v} h_n *_q \mathcal{F}_{q,v} f) \in \mathcal{L}_{q,v,1}.$$

On the other hand

$$\mathcal{F}_{q,v} h_n *_q \mathcal{F}_{q,v} f \rightarrow \mathcal{F}_{q,v} f.$$

The Dominated convergence theorem gives

$$\lim_{n \rightarrow \infty} \int_0^\infty \phi_i(\mathcal{F}_{q,v} h_n *_q \mathcal{F}_{q,v} f)(x) x^{2v+1} d_q x = \int_0^\infty \phi_i(\mathcal{F}_{q,v} f) x^{2v+1} d_q x.$$

It also holds that

$$\begin{aligned} E(|\mathcal{F}_{q,v} f|^2) &= 2 \int_0^\infty \phi_1(\mathcal{F}_{q,v} f) x^{2v+1} d_q \\ &\quad - 2 \int_0^\infty \phi_2(\mathcal{F}_{q,v} f) x^{2v+1} d_q + 3 \|\mathcal{F}_{q,v} f\|_{q,v,2}^2, \end{aligned}$$

and

$$\begin{aligned} E(|\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f|^2) &= 2 \int_0^\infty \phi_1(\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f) x^{2v+1} d_q x \\ &\quad - 2 \int_0^\infty \phi_2(\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f) x^{2v+1} d_q x \\ &\quad + 3 \|\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f\|_{q,v,2}^2. \end{aligned}$$

Then

$$\lim_{n \rightarrow \infty} E(|\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f|^2) = E(|\mathcal{F}_{q,v}f|^2). \quad (7)$$

With (5) and the limits (6) and (7) we complete the proof of (2).

For a given  $b > 0$  and  $c, d \in \mathbb{R}_q^+$  such that  $0 < B_{q,v}k_c k_d < 1$  and by (1) we can write

$$-E(|f|^2) \leq \log k_c + c^a \|x^{a/2} f\|_{q,v,2}^2.$$

$$-E(|\mathcal{F}_{q,v}f|^2) \leq \log k_d + d^b \|x^{b/2} \mathcal{F}_{q,v}f\|_{q,v,2}^2.$$

Which implies with (2)

$$\begin{aligned} -\log B_{q,v} &\leq -E(|f|^2) - E(|\mathcal{F}_{q,v}f|^2) \\ &\leq \log k_c k_d + c^a \|x^{a/2} f\|_{q,v,2}^2 + d^b \|x^{b/2} \mathcal{F}_{q,v}f\|_{q,v,2}^2. \end{aligned}$$

By replacing  $f$  by  $\frac{f}{\|f\|_{q,v,2}}$  we get

$$c^a \|x^{a/2} f\|_{q,v,2}^2 + d^b \|x^{b/2} \mathcal{F}_{q,v}f\|_{q,v,2}^2 \geq -\log(B_{q,v}k_c k_d) \|f\|_{q,v,2}^2.$$

This finish the proof. ■

**Corollary 1** *There exist  $k > 0$  such that for any function  $f \in \mathcal{L}_{q,v,2}$  we have*

$$\|xf\|_{q,v,2}^2 \|\mathcal{F}_{q,v}f\|_{q,v,2}^2 \geq k \|f\|_{q,v,2}^2.$$

**Proof.** Let  $a = b = 2$  then by Theorem 3

$$\|xf\|_{q,v,2}^2 + \|\mathcal{F}_{q,v}f\|_{q,v,2}^2 \geq -\frac{1}{c^2} \log \left( B_{q,v} \frac{\sigma_2^2}{c^{4(v+1)}} \right) \|f\|_{q,v,2}^4.$$

Now put

$$f_t(x) = f(tx), \quad t \in \mathbb{R}_q^+,$$

then

$$\mathcal{F}_{q,v} f_t(x) = \frac{1}{t^{2v+2}} \mathcal{F}_{q,v} f(x/t), \quad \|x \mathcal{F}_{q,v} f_t\|_{q,v,2}^2 = \frac{1}{t^{2v}} \|\mathcal{F}_{q,v} f\|_{q,v,2}^2,$$

and

$$\|f_t\|_{q,v,2}^2 = \frac{1}{t^{2v+2}} \|f\|_{q,v,2}^2, \quad \|x f_t\|_{q,v,2}^2 = \frac{1}{t^{2v+4}} \|x f\|_{q,v,2}^2,$$

which gives

$$t^4 \|\mathcal{F}_{q,v} f\|_{q,v,2}^2 + t^2 \frac{1}{c^2} \log \left( B_{q,v} \frac{\sigma_2^2}{c^{4(v+1)}} \right) \|f\|_{q,v,2}^2 + \|x f\|_{q,v,2}^2 \geq 0,$$

and then

$$\|x f\|_{q,v,2}^2 \|\mathcal{F}_{q,v} f\|_{q,v,2}^2 \geq \frac{1}{c^4} \log \left( B_{q,v} \frac{\sigma_2^2}{c^{4(v+1)}} \right)^2 \|f\|_{q,v,2}^4.$$

We choose  $k$  as follows

$$k = \sup_{c \in \mathbb{R}_q^+} \frac{1}{c^4} \log \left( B_{q,v} \frac{\sigma_2^2}{c^{4(v+1)}} \right)^2$$

Which leads to the result. ■

## References

- [1] L. Dhaouadi, A. Fitouhi and J. El Kamel, Inequalities in q-Fourier Analysis, Journal of Inequalities in Pure and Applied Mathematics, Volume 7, Issue 5, Article 171, 2006.
- [2] G. Gasper and M. Rahman, Basic hypergeometric series, Encyclopedia of mathematics and its applications 35, Cambridge university press, 1990.
- [3] F. H. Jackson, On a q-Definite Integrals, Quarterly Journal of Pure and Application Mathematics 41, 1910, 193-203.
- [4] T. H. Koornwinder and R. F. Swarttouw, On q-Analogues of the Hankel and Fourier Transform, Trans. A. M. S. 1992, 333, 445-461.
- [5] R. F. Swarttouw, The Hahn-Exton q-Bessel functions PhD Thesis The Technical University of Delft ((1992)).