



HAL
open science

Heisenberg Uncertainty Principle for the q -Bessel Fourier transform

Lazhar Dhaouadi

► **To cite this version:**

Lazhar Dhaouadi. Heisenberg Uncertainty Principle for the q -Bessel Fourier transform. 2007. hal-00161302v2

HAL Id: hal-00161302

<https://hal.science/hal-00161302v2>

Preprint submitted on 1 Jul 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Heisenberg Uncertainty Principle for the q -Bessel Fourier transform

Lazhar Dhaouadi *

Abstract

In this paper we use an I.I. Hirschman-W. Beckner entropy argument to give an uncertainty inequality for the q -Bessel Fourier transform:

$$\mathcal{F}_{q,v}f(x) = c_{q,v} \int_0^\infty f(t)j_v(xt, q^2)t^{2v+1}d_qt,$$

where $j_v(x, q)$ is the normalized Hahn-Exton q -Bessel function.

1 Introduction

I.I.Hirschman-W. Beckner entropy argument is one further variant of Heisenberg's uncertainty principle.

Let \widehat{f} be the Fourier transform of f defined by

$$\widehat{f}(x) = \int_{-\infty}^{\infty} f(y)e^{2i\pi xy}f(y)dy, \quad x \in \mathbb{R}.$$

If $f \in L^2(\mathbb{R})$ with L^2 -norm $\|f\|_2 = 1$, then by Plancherel's theorem $\|\widehat{f}\|_2 = 1$, so that $|f(x)|^2$ and $|\widehat{f}(x)|^2$ are probability frequency functions. The variance of a probability frequency g is defined by

$$V[g] = \int_{\mathbb{R}} x^2g(x)dx.$$

The Heisenberg uncertainty principle can be stated as follows

$$V[|f|^2]V[|\widehat{f}|^2] \geq \frac{1}{16\pi^2}. \quad (1)$$

*Institut Préparatoire aux Etudes d'Ingénieur de Bizerte (Université du 7 novembre Carthage). Route Menzel Abderrahmene Bizerte, 7021 Zarzouna, Tunisia. E-mail lazhardhaouadi@yahoo.fr

If g is a probability frequency function, then the entropy of g is defined by

$$E(g) = \int_{\mathbb{R}} g(x) \log(x) dx.$$

With f as above, Hirschman [10] proved that

$$E(|f|^2) + E(|\widehat{f}|^2) \leq 0. \quad (2)$$

By an inequality of Shannon and Weaver it follows that (2) implies (1). Using the Babenko-Beckner inequality

$$\|\widehat{f}\|_{p'} \leq A(p) \|f\|_p, \quad 1 < p < 2, \quad A(p) = \left[p^{1/p} (p')^{-1/p'} \right]^{1/2},$$

in Hirschman's proof of (2) another uncertainty inequality is deduced. For more detail the reader can consult [8,10,11].

In this paper we use I.I. Hirschman entropy argument de give an uncertainty inequality for the q -Bessel Fourier transform (also called q -Hankel transform).

Note that other versions of the Heisenberg uncertainty principle for the q -Fourier transform have recently appeared in the literature [1,2,6]. There are some differences of the results cited above and our result:

- In [1] the uncertainty inequality is established for the q -cosine and q -sine transform but here is established for the q -Bessel transform.
- In [2] the uncertainty inequality is for the q^2 -Fourier transform but here is for the q -Hankel transform.
- In [6] the uncertainty inequality is established for functions in q -Schwartz space. In this paper the uncertainty inequality is established for functions in $\mathcal{L}_{q,2,v}$ space.

The inequality discuss here is a quantitative uncertainty principles which give an information about how a function and its q -Bessel Fourier transform relate. A qualitative uncertainty principles give an information about how a function (and its Fourier transform) behave under certain circumstances. A classical qualitative uncertainty principle called Hardy's theorem. In [4,7] a q -version of the Hardy's theorem for the q -Bessel Fourier transform was established.

In the end, our objective is to develop a coherent harmonic analysis attached to the q -Bessel operator

$$\Delta_{q,v} f(x) = \frac{1}{x^2} [f(q^{-1}x) - (1 + q^{2v})f(x) + q^{2v}f(qx)].$$

Thus, this paper is an opportunity to implement the arguments of the q -Bessel Fourier analysis proved before, as the Plancherel formula, the positivity of the q -translation operator, the q -convolution product, the q -Gauss kernel...

2 The q -Bessel Fourier transform

In the following we will always assume $0 < q < 1$ and $v > -1$. We denote by

$$\mathbb{R}_q = \{\pm q^n, n \in \mathbb{Z}\}, \quad \mathbb{R}_q^+ = \{q^n, n \in \mathbb{Z}\}.$$

For more informations on the q -series theory the reader can see the references [9,12,14] and the references [3,5,13] about the q -bessel Fourier analysis. Also for details of the proofs of the following results in this section can be found in [3].

Definition 1 *The q -Bessel operator is defined as follows*

$$\Delta_{q,v}f(x) = \frac{1}{x^2} [f(q^{-1}x) - (1 + q^{2v})f(x) + q^{2v}f(qx)].$$

Definition 2 *The normalized q -Bessel function of Hahn-Exton is defined by*

$$j_v(x, q^2) = \sum_{n=0}^{\infty} (-1)^n \frac{q^{n(n+1)}}{(q^{2v+2}, q^2)_n (q^2, q^2)_n} x^{2n}.$$

Proposition 1 *The function*

$$x \mapsto j_v(\lambda x, q^2)$$

is the eigenfunction of the operator $\Delta_{q,v}$ associated with the eigenvalue $-\lambda^2$.

Definition 3 *The q -Jackson integral of a function f defined on \mathbb{R}_q is*

$$\int_0^{\infty} f(t) d_q t = (1 - q) \sum_{n \in \mathbb{Z}} q^n f(q^n).$$

Definition 4 *We denote by $\mathcal{L}_{q,p,v}$ the space of even functions f defined on \mathbb{R}_q such that*

$$\|f\|_{q,p,v} = \left[\int_0^{\infty} |f(x)|^p x^{2v+1} d_q x \right]^{1/p} < \infty.$$

Definition 5 We denote by $\mathcal{C}_{q,0}$ the space of even functions defined on \mathbb{R}_q tending to 0 as $x \rightarrow \pm\infty$ and continuous at 0 equipped with the topology of uniform convergence. The space $\mathcal{C}_{q,0}$ is complete with respect to the norm

$$\|f\|_{q,\infty} = \sup_{x \in \mathbb{R}_q} |f(x)|.$$

Definition 6 The q -Bessel Fourier transform $\mathcal{F}_{q,v}$ (also called q -Hankel transform) is defined by

$$\mathcal{F}_{q,v}f(x) = c_{q,v} \int_0^\infty f(t)j_v(xt, q^2)t^{2v+1}d_qt, \quad \forall x \in \mathbb{R}_q.$$

where

$$c_{q,v} = \frac{1}{1-q} \frac{(q^{2v+2}; q^2)_\infty}{(q^2; q^2)_\infty}.$$

Proposition 2 Let $f \in \mathcal{L}_{q,1,v}$ then $\mathcal{F}_{q,v}f$ existe and $\mathcal{F}_{q,v}f \in \mathcal{C}_{q,0}$.

Definition 7 The q -translation operator is given as follows

$$T_{q,x}^v f(y) = c_{q,v} \int_0^\infty \mathcal{F}_{q,v}f(t)j_v(yt, q^2)j_v(xt, q^2)t^{2v+1}d_qt \quad \forall f \in \mathcal{L}_{q,1,v}.$$

Definition 8 The operator $T_{q,x}^v$ is said positive if $T_{q,x}^v f \geq 0$ when $f \geq 0$ for all $x \in \mathbb{R}_q$. We denote by Q_v the domain of positivity of $T_{q,x}^v$

$$Q_v = \{q \in]0, 1[, \quad T_{q,x}^v \text{ is positive}\}.$$

In the following we assume that $q \in Q_v$.

Proposition 3 If $f \in \mathcal{L}_{q,1,v}$ then

$$\int_0^\infty T_{q,x}^v f(y)y^{2v+1}d_qy = \int_0^\infty f(y)y^{2v+1}d_qy.$$

Definition 9 The q -convolution product is defined as follows

$$f *_q g(x) = c_{q,v} \int_0^\infty T_{q,x}^v f(y)g(y)y^{2v+1}d_qy.$$

Proposition 4 Let $f, g \in \mathcal{L}_{q,1,v}$ then $f *_q g \in \mathcal{L}_{q,1,v}$ and we have

$$\mathcal{F}_{q,v}(f *_q g) = \mathcal{F}_{q,v}(g) \times \mathcal{F}_{q,v}(f).$$

Proposition 5 Let $f \in \mathcal{L}_{q,1,v}$ and $g \in \mathcal{L}_{q,2,v}$ then $f *_q g \in \mathcal{L}_{q,2,v}$ and we have

$$\mathcal{F}_{q,v}(f *_q g) = \mathcal{F}_{q,v}(f) \times \mathcal{F}_{q,v}(g).$$

Theorem 1 The q -Bessel Fourier transform $\mathcal{F}_{q,v}$ satisfies

1. $\mathcal{F}_{q,v}$ sends $\mathcal{L}_{q,2,v}$ to $\mathcal{L}_{q,2,v}$.
2. For $f \in \mathcal{L}_{q,2,v}$, we have

$$\|\mathcal{F}_{q,v}(f)\|_{q,2,v} = \|f\|_{q,2,v}.$$

3. The operator $\mathcal{F}_{q,v} : \mathcal{L}_{q,2,v} \rightarrow \mathcal{L}_{q,2,v}$ is bijective and

$$\mathcal{F}_{q,v}^{-1} = \mathcal{F}_{q,v}.$$

Proposition 6 Given $1 < p \leq 2$ and $\frac{1}{p} + \frac{1}{\bar{p}} = 1$. If $f \in \mathcal{L}_{q,p,v}$ then

$$\mathcal{F}_{q,v}(f) \in \mathcal{L}_{q,\bar{p},2,v}$$

and

$$\|\mathcal{F}_{q,v}(f)\|_{q,\bar{p},v} \leq B_{q,v}^{\left(\frac{2}{p}-1\right)} \|f\|_{q,p,v},$$

where

$$B_{q,v} = \frac{1}{1-q} \frac{(-q^2; q^2)_\infty (-q^{2v+2}; q^2)_\infty}{(q^2; q^2)_\infty}.$$

Definition 10 The q -exponential function is defined by

$$e(z, q) = \sum_{n=0}^{\infty} \frac{z^n}{(q, q)_n} = \frac{1}{(z; q)_\infty}, \quad |z| < 1.$$

Proposition 7 The q -Gauss kernel

$$G^v(x, t^2, q^2) = \frac{(-q^{2v+2}t^2, -q^{-2v}/t^2; q^2)_\infty}{(-t^2, -q^2/t^2; q^2)_\infty} e\left(-\frac{q^{-2v}}{t^2}x^2, q^2\right), \quad \forall x, t \in \mathbb{R}_q^+$$

satisfies

$$\mathcal{F}_{q,v} \{e(-t^2y^2, q^2)\} (x) = G^v(x, t^2, q^2),$$

and for all function $f \in \mathcal{L}_{q,2,v}$

$$\lim_{n \rightarrow \infty} \|G^v(x, q^{2n}, q^2) *_q f - f\|_{q,2,v} = 0.$$

3 Uncertainty Principle

The following Lemma are crucial for the proof of our main result. First we enunciate the Jensens inequality

Lemma 1 *Let γ be a probability measure on \mathbb{R}_q^+ . Let g be a convex function on a subset I of \mathbb{R} . If $\psi : \mathbb{R}_q^+ \rightarrow I$ satisfies*

$$\int_0^\infty \psi(u) d\gamma(u) \in I,$$

then we have

$$g\left(\int_0^\infty \psi(x) d\gamma(x)\right) \leq \int_0^\infty g \circ \psi(x) d\gamma(x).$$

Proof. Let

$$t = \int_0^\infty \psi(u) d\gamma(u).$$

There exist $c \in \mathbb{R}$ such that for all $y \in I$ it holds

$$g(y) \geq g(t) + c(y - t).$$

Now let $y = \psi(x)$ we obtain

$$g(\psi(x)) \geq g(t) + c(\psi(x) - t).$$

Integrating both sides and using the special value of t gives

$$\int_0^\infty g(\psi(x)) d\gamma(x) \geq \int_0^\infty [g(t) + c(\psi(x) - t)] d\gamma(x) = g(t).$$

This finish the proof. ■

Lemma 2 *Let f be an even function defined on \mathbb{R}_q . Assume $\psi : \mathbb{R} \rightarrow \mathbb{R}_+$ is a convex function and $\psi \circ f \in \mathcal{L}_{q,1,v}$. If ϱ_n is a sequence of non-negative function in $\mathcal{L}_{q,1,v}$ such that*

$$\mathcal{F}_{q,v}(\varrho_n)(0) = c_{q,v} \int_0^\infty \varrho_n(x) x^{2v+1} d_q x = 1$$

*and $\varrho_n *_q f \rightarrow f$ then $\psi \circ (\varrho_n *_q f)$ is in $\mathcal{L}_{q,1,v}$ and*

$$\lim_{n \rightarrow \infty} \int_0^\infty \psi \circ (\varrho_n *_q f)(x) x^{2v+1} d_q x = \int_0^\infty \psi \circ f(x) x^{2v+1} d_q x.$$

Proof. For a given x and by Proposition 3 we have

$$c_{q,v} \int_0^\infty T_{q,x}^v \varrho_n(y) y^{2v+1} d_q y = 1$$

From the positivity of $T_{q,x}^v$ we see that

$$c_{q,v} T_{q,x}^v \varrho_n(y) y^{2v+1} d_q y$$

is a probability measure on \mathbb{R}_q^+ . The following holds by Jensens Inequality

$$\begin{aligned} \psi \circ (\varrho_n *_q f)(x) &= \psi \left[c_{q,v} \int_0^\infty f(y) T_{q,x}^v \varrho_n(y) y^{2v+1} d_q y \right] \\ &\leq c_{q,v} \int_0^\infty \psi \circ f(y) T_{q,x}^v \varrho_n(y) y^{2v+1} d_q y \\ &= \varrho_n *_q \psi \circ f(x). \end{aligned}$$

By the use of the Fatou's Lemma and Proposition 4 we obtain

$$\begin{aligned} &\int_0^\infty \psi \circ f(x) x^{2v+1} d_q x \\ &= \int_0^\infty \liminf_{n \rightarrow \infty} \psi \circ (\varrho_n *_q f)(x) x^{2v+1} d_q x \\ &\leq \liminf_{n \rightarrow \infty} \int_0^\infty \psi \circ (\varrho_n *_q f)(x) x^{2v+1} d_q x \\ &\leq \limsup_{n \rightarrow \infty} \int_0^\infty \psi \circ (\varrho_n *_q f)(x) x^{2v+1} d_q x \\ &\leq \lim_{n \rightarrow \infty} \int_0^\infty \varrho_n *_q \psi \circ f(x) x^{2v+1} d_q x \\ &= \frac{1}{c_{q,v}} \lim_{n \rightarrow \infty} \mathcal{F}_{q,v}(\varrho_n)(0) \times \mathcal{F}_{q,v}(\psi \circ f)(0) \\ &= \int_0^\infty \psi \circ f(x) x^{2v+1} d_q x. \end{aligned}$$

This finish the proof. ■

Definition 11 For a positive function ϕ define the entropy of ϕ to be

$$E(\phi) = \int_0^\infty \phi(x) \log \phi(x) x^{2v+1} d_q x.$$

$E(\phi)$ can have values in $[-\infty, \infty]$.

Remark 1 For a given $c \in \mathbb{R}_q^+$ let

$$d\gamma(x) = k_c^{-1} \exp(-|cx|^a) x^{2v+1} d_q x$$

where

$$\sigma_a = \int_0^\infty \exp(-|x|^a) x^{2v+1} d_q x, \quad k_c = \frac{\sigma_a}{c^{2v+2}}.$$

Then $d\gamma(x)$ is a probability measure on \mathbb{R}_q^+ .

Lemma 3 Let $a > 0$. For a positive function $\phi \in \mathcal{L}_{q,1,v}$ such that

$$\|\phi\|_{q,1,v} = 1$$

and

$$M_a(\phi) = \left(\int_0^\infty |x|^a \phi(x) x^{2v+1} d_q x \right)^{\frac{1}{a}}$$

is finite, we have

$$-E(\phi) \leq \log k_c + c^a M_a^a(\phi). \quad (3)$$

Proof. Indeed, defining

$$\psi(x) = k_c \exp(|cx|^a) \phi(x),$$

From Remark 1 we see that

$$\int_0^\infty \psi(x) d\gamma(x) = 1.$$

According to the fact that $g : t \mapsto t \log t$ is convex on \mathbb{R}_+^* , so Jensen's inequality gives

$$g \left[\int_0^\infty \psi(x) d\gamma(x) \right] \leq \int_0^\infty g \circ \psi(x) d\gamma(x).$$

Hence,

$$0 = \left[\int_0^\infty \psi(x) d\gamma(x) \right] \log \left[\int_0^\infty \psi(x) d\gamma(x) \right] \leq \int_0^\infty \psi(x) \log \psi(x) d\gamma(x).$$

This implies

$$\begin{aligned} 0 &\leq \int_0^\infty \phi(x) \log [k_c \exp(|cx|^a) \phi(x)] x^{2v+1} d_q x \\ &= \int_0^\infty \phi(x) [\log k_c + |cx|^a + \log \phi(x)] x^{2v+1} d_q x. \end{aligned}$$

$$0 \leq \log k_c + c^a \int_0^\infty |x|^a \phi(x) x^{2v+1} d_q x + \int_0^\infty \phi(x) \log \phi(x) x^{2v+1} d_q x.$$

In the end

$$0 \leq \log k_c + c^a M_a^a(\phi) + E(\phi).$$

This finish the proof. ■

Lemma 4 *Let $f \in \mathcal{L}_{q,1,v} \cap \mathcal{L}_{q,2,v}$ then we have*

$$E(|f|^2) + E(|\mathcal{F}_{q,v} f|^2) \leq 2 \|f\|_{q,v,2}^2 \log(B_{q,v} \|f\|_{q,v,2}^2). \quad (4)$$

Proof. Hölder inequality implies that f will be in $\mathcal{L}_{q,p,v}$ for $1 < p \leq 2$. With

$$\frac{1}{p} + \frac{1}{\bar{p}} = 1,$$

Hausdorff-Young's inequality (Proposition 6) tells us that $\mathcal{F}_{q,v} f$ is in $\mathcal{L}_{q,\bar{p},v}$. So we can define the functions

$$A(p) = \int_0^\infty |f(x)|^p d_q x \quad \text{and} \quad B(\bar{p}) = \int_0^\infty |\mathcal{F}_{q,v} f(x)|^{\bar{p}} x^{2v+1} d_q x.$$

Now define

$$\begin{aligned} C(p) &= \log \|\mathcal{F}_{q,v} f\|_{q,\bar{p},v} - \log \left(B_{q,v}^{\frac{2}{p}-1} \|f\|_{q,p,v} \right) \\ &= \frac{1}{\bar{p}} \log B(\bar{p}) - \frac{1}{p} \log A(p) - \left(\frac{2}{p} - 1 \right) \log B_{q,v}. \end{aligned}$$

By Hausdorff-Young's inequality

$$C(p) \leq 0, \quad \text{for } 1 < p < 2,$$

and by Plancherel equality (Theorem 1 part 2)

$$C(2) = 0.$$

Then

$$C'(2^-) \geq 0.$$

On the other hand for $1 < p < 2$ we have

$$C'(p) = \frac{\bar{p}'}{\bar{p}} \frac{B'(\bar{p})}{B(\bar{p})} - \frac{\bar{p}'}{\bar{p}^2} \log B(\bar{p}) - \frac{1}{p} \frac{A'(p)}{A(p)} + \frac{1}{p^2} \log A(p) + \frac{2}{p^2} \log B_{q,v}.$$

The derivative of \bar{p} with respect to p is

$$\bar{p}' = -\frac{1}{(p-1)^2}.$$

For a given $x > 0$ we have

$$\lim_{p \rightarrow 2} \frac{x^p - x^2}{p-2} = x^2 \log x.$$

Then

$$A'(2^-) = \lim_{p \rightarrow 2^-} \frac{A(p) - A(2)}{p-2} = \frac{1}{2} E(|f|^2),$$

$$B'(2^+) = \lim_{\bar{p} \rightarrow 2^+} \frac{B(\bar{p}) - B(2)}{\bar{p}-2} = \frac{1}{2} E(|\mathcal{F}_{q,v} f|^2).$$

Since

$$p \mapsto \frac{x^p - x^2}{p-2}$$

is an increasing function, the exchange of the signs limit and integral is valid sense. On the other hand

$$\lim_{p \rightarrow 2^-} A(p) = \|f\|_{q,v,2}^2, \quad \lim_{\bar{p} \rightarrow 2^+} B(\bar{p}) = \|\mathcal{F}_{q,v} f\|_{q,v,2}^2 = \|f\|_{q,v,2}^2.$$

So

$$C'(2^-) = \lim_{p \rightarrow 2^-} \frac{C(p) - C(2)}{p-2} = -\frac{1}{2\|f\|_{q,v,2}^2} [A'(2^-) + B'(2^+)] + \frac{1}{2} \log(B_{q,v} \|f\|_{q,v,2}^2).$$

Therefore

$$A'(2^-) + B'(2^+) - \|f\|_{q,v,2}^2 \log(B_{q,v} \|f\|_{q,v,2}^2) \leq 0,$$

and then

$$E(|f|^2) + E(|\mathcal{F}_{q,v} f|^2) \leq 2\|f\|_{q,v,2}^2 \log(B_{q,v} \|f\|_{q,v,2}^2).$$

This finish the proof. ■

Lemma 5 *Let $f \in \mathcal{L}_{q,2,v}$ then we have*

$$E(|f|^2) + E(|\mathcal{F}_{q,v} f|^2) \leq 2\|f\|_{q,v,2}^2 \log(B_{q,v} \|f\|_{q,v,2}^2). \quad (5)$$

Proof. Assume that $E(|f|^2)$ and $E(|\mathcal{F}_{q,v}f|^2)$ are defined and then approximate f by functions in $\mathcal{L}_{q,1,v} \cap \mathcal{L}_{q,2,v}$. Let

$$h_n(x) = e(-q^{2n}x^2, q^2).$$

The function h_n is in $\mathcal{L}_{q,2,v}$ then $h_n f \in \mathcal{L}_{q,1,v}$. On the other hand $h_n \in \mathcal{C}_{q,0}$ then $h_n f \in \mathcal{L}_{q,2,v}$. We obtain

$$h_n f \in \mathcal{L}_{q,1,v} \cap \mathcal{L}_{q,2,v}.$$

The following holds by (2)

$$E(|h_n f|^2) + E(|\mathcal{F}_{q,v}(h_n f)|^2) \leq 2\|h_n f\|_{q,2,v}^2 \log(B_{q,v}\|h_n f\|_{q,2,v}^2). \quad (6)$$

One can see by the Lebesgue Dominated Convergence Theorem that

$$\lim_{n \rightarrow \infty} \|h_n f\|_{q,2,v} = \|f\|_{q,2,v} \quad (7)$$

and

$$\lim_{n \rightarrow \infty} E(|h_n f|^2) = E(|f|^2). \quad (8)$$

By the use of Proposition 5 and the inversion formula (Theorem 1 part 3) we see that

$$\mathcal{F}_{q,v}(h_n f) = \mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f.$$

We will prove that

$$\lim_{n \rightarrow \infty} E(|\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f|^2) = E(|\mathcal{F}_{q,v}f|^2).$$

The functions

$$\phi_1(x) = x^2 \log^+ |x| \quad \text{and} \quad \phi_2(x) = x^2 \left(-\log^- |x| + \frac{3}{2} \right),$$

are convex on \mathbb{R} , where

$$\log^+ x = \max\{0, \log x\} \quad \text{and} \quad \log^- x = \min\{0, \log x\}.$$

Note that

$$2\phi_1(x) - 2\phi_2(x) + 3x^2 = x^2 \log |x|^2.$$

Since

- From the inversion formula we see that

$$c_{q,v} \int_0^\infty \mathcal{F}_{q,v}h_n(t)t^{2v+1}d_q t = h_n(0) = 1.$$

- The function $\mathcal{F}_{q,v}h_n \geq 0$.
- The functions ϕ_i are convex on \mathbb{R} .
- $E(\mathcal{F}_{q,v}f)$ is finite then $\phi_i(\mathcal{F}_{q,v}f)$ is in $\mathcal{L}_{q,1,v}$.
- From Proposition 7 we have

$$\lim_{n \rightarrow \infty} \mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f(x) = \mathcal{F}_{q,v}f(x)$$

we deduce that $\mathcal{F}_{q,v}h_n$ and ϕ_i satisfy the conditions of Lemma 2. Then we obtain

$$\lim_{n \rightarrow \infty} \int_0^\infty \phi_i \circ (\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f)(x) x^{2v+1} d_q x = \int_0^\infty \phi_i \circ (\mathcal{F}_{q,v}f)(x) x^{2v+1} d_q x, \quad i = 1, 2.$$

It also hold

$$\begin{aligned} E(|\mathcal{F}_{q,v}f|^2) &= 2 \int_0^\infty \phi_1(\mathcal{F}_{q,v}f) x^{2v+1} d_q x \\ &\quad - 2 \int_0^\infty \phi_2(\mathcal{F}_{q,v}f) x^{2v+1} d_q x + 3 \|\mathcal{F}_{q,v}f\|_{q,2,v}^2, \end{aligned}$$

and

$$\begin{aligned} E(|\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f|^2) &= 2 \int_0^\infty \phi_1(\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f) x^{2v+1} d_q x \\ &\quad - 2 \int_0^\infty \phi_2(\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f) x^{2v+1} d_q x \\ &\quad + 3 \|\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f\|_{q,2,v}^2. \end{aligned}$$

Then

$$\lim_{n \rightarrow \infty} E(|\mathcal{F}_{q,v}h_n *_q \mathcal{F}_{q,v}f|^2) = E(|\mathcal{F}_{q,v}f|^2). \quad (9)$$

With (6) and the limits (7), (8) and (9) we complete the proof of (5).

Note that these limits also hold in the case where $E(|f|^2)$ and $E(|\mathcal{F}_{q,v}f|^2)$ are ∞ or $-\infty$. ■

Now we are in position to state and prove the uncertainty inequality for the q -Bessel Fourier transform.

Theorem 2 *Given $a, b > 0$. Then for all $c, d \in \mathbb{R}_q^+$ satisfying*

$$0 < B_{q,v}^2 \frac{\sigma_a \sigma_b}{(cd)^{2v+2}} < 1,$$

the following hold for any function $f \in \mathcal{L}_{q,2,v}$

$$c^a \left\| x^{a/2} f \right\|_{q,2,v}^2 + d^b \left\| x^{b/2} \mathcal{F}_{q,v}f \right\|_{q,2,v}^2 \geq -\log \left(B_{q,v}^2 \frac{\sigma_a \sigma_b}{(cd)^{2v+2}} \right) \|f\|_{q,2,v}^2.$$

Proof. Assume that $\|f\|_{q,2,v} = 1$. By (3) we can write

$$\begin{aligned} -E(|f|^2) &\leq \log k_c + c^a \left\| x^{a/2} f \right\|_{q,2,v}^2 \\ -E\left(|\mathcal{F}_{q,v} f|^2\right) &\leq \log k_d + d^b \left\| x^{b/2} \mathcal{F}_{q,v} f \right\|_{q,2,v}^2. \end{aligned}$$

Which implies with (5)

$$\begin{aligned} -2 \log B_{q,v} &\leq -E\left(|f|^2\right) - E\left(|\mathcal{F}_{q,v} f|^2\right) \\ &\leq \log(k_c k_d) + c^a \left\| x^{a/2} f \right\|_{q,2,v}^2 + d^b \left\| x^{b/2} \mathcal{F}_{q,v} f \right\|_{q,2,v}^2. \end{aligned}$$

By replacing f by $\frac{f}{\|f\|_{q,2,v}}$ we get

$$c^a \left\| x^{a/2} f \right\|_{q,2,v}^2 + d^b \left\| x^{b/2} \mathcal{F}_{q,v} f \right\|_{q,2,v}^2 \geq -\log(B_{q,v}^2 k_c k_d) \|f\|_{q,2,v}^2.$$

This finish the proof. ■

Corollary 1 *There exist $k > 0$ such that for any function $f \in \mathcal{L}_{q,2,v}$ we have*

$$\|x f\|_{q,2,v} \|x \mathcal{F}_{q,v} f\|_{q,2,v} \geq k \|f\|_{q,2,v}^2.$$

Proof. Let $a = b = 2$ and $c = d$ then by Theorem 3

$$\|x f\|_{q,2,v}^2 + \|x \mathcal{F}_{q,v} f\|_{q,2,v}^2 \geq -\frac{1}{c^2} \log\left(B_{q,v}^2 \frac{\sigma_2^2}{c^{4(v+1)}}\right) \|f\|_{q,2,v}^2,$$

where

$$0 < \left(B_{q,v}^2 \frac{\sigma_2^2}{c^{4(v+1)}}\right) < 1.$$

Now put

$$f_t(x) = f(tx), \quad t \in \mathbb{R}_q^+,$$

then

$$\mathcal{F}_{q,v} f_t(x) = \frac{1}{t^{2v+2}} \mathcal{F}_{q,v} f(x/t), \quad \|x \mathcal{F}_{q,v} f_t\|_{q,2,v}^2 = \frac{1}{t^{2v}} \|\mathcal{F}_{q,v} f\|_{q,2,v}^2,$$

and

$$\|f_t\|_{q,2,v}^2 = \frac{1}{t^{2v+2}} \|f\|_{q,2,v}^2, \quad \|x f_t\|_{q,2,v}^2 = \frac{1}{t^{2v+4}} \|x f\|_{q,2,v}^2,$$

which gives

$$t^4 \|x\mathcal{F}_{q,v}f\|_{q,2,v}^2 + t^2 \frac{1}{c^2} \log \left(B_{q,v}^2 \frac{\sigma_2^2}{c^{4(v+1)}} \right) \|f\|_{q,v,2}^2 + \|xf\|_{q,2,v}^2 \geq 0,$$

and then

$$\|xf\|_{q,2,v} \|x\mathcal{F}_{q,v}f\|_{q,2,v} \geq \psi(c) \|f\|_{q,2,v}^2.$$

where

$$\psi(c) = \frac{v+1}{[\sigma_2 B_{q,v}]^{\frac{1}{v+1}}} |z_c \log(z_c)|, \quad z_c = \frac{[\sigma_2 B_{q,v}]^{\frac{1}{v+1}}}{c^2}, \quad 0 < z_c < 1.$$

One can see that

$$\sup_{0 < z_c < 1} \psi(c) = \psi(q^\alpha), \quad \alpha = \frac{\log[\sigma_2 B_{q,v}]}{2(1+v)\log q} + \frac{1}{2\log q}.$$

Let

$$n_1 = \lfloor \alpha \rfloor, \quad n_2 = \lceil \alpha \rceil,$$

where $\lfloor \cdot \rfloor$ and $\lceil \cdot \rceil$ are respectively the floor and ceiling functions. Now the constant k is given as follows

$$k = \psi(q^{n_1}), \quad \text{if } \lceil \alpha \rceil \geq \alpha - \frac{1}{2\log q}$$

and

$$k = \max\{\psi(q^{n_1}), \psi(q^{n_2})\}, \quad \text{if } \lceil \alpha \rceil < \alpha - \frac{1}{2\log q}.$$

This finish the proof. ■

References

- [1] N. Bettaibi, A. Fitouhi and W. Binous, Uncertainty principles for the q-trigonometric Fourier transforms, *Math. Sci. Res. J.* 11 (2007).
- [2] N. Bettaibi, Uncertainty principles in q^2 -analogue Fourier analysis, *Math. Sci. Res. J.* 11 (2007).
- [3] L. Dhaouadi, A. Fitouhi and J. El Kamel, Inequalities in q-Fourier Analysis, *Journal of Inequalities in Pure and Applied Mathematics*, Volume 7, Issue 5, Article 171, 2006.

- [4] L. Dhaouadi, Hardy's theorem for the q -Bessel Fourier transform, arXiv: 0707.2346 v1 [math.CA].
- [5] A. Fitouhi, M.Hamza and F. Bouzeffour, The $q - j_\alpha$ Bessel function J. Appr. Theory. 115,144-166(2002).
- [6] A. Fitouhi, N. Bettaibi, W. Binous and H.B. Elmonser, Uncertainty principles for the basic Bessel transform, Ramanujan J., in press.
- [7] A. Fitouhi, N. Bettaibi and R. Bettaieb, On Hardy's inequality for symmetric integral transforms and analogous, Appl. Math. Comput. 198 (2008).
- [8] G.B. Folland and A. Sitaram. The uncertainty principle: a mathematical survey. J. Fourier Anal. Appl., 3(3):207–238, 1997.
- [9] G. Gasper and M. Rahman, Basic hypergeometric series, Encyclopedia of mathematics and its applications 35, Cambridge university press, 1990.
- [10] I.I. Hirschman, Jr. A note on entropy. Amer. J. Math., 79:152156, 1957.
- [11] H.P. Heinig, and M. Smith, Extensions of the Heisenberg-Weyl inequality. Internat. J. Math. Math. Sci. 9 (1986), no. 1, 185–192.
- [12] F. H. Jackson, On a q -Definite Integrals, Quarterly Journal of Pure and Application Mathematics 41, 1910, 193-203.
- [13] T. H. Koornwinder and R. F. Swarttouw, On q -Analogues of the Hankel and Fourier Transform, Trans. A. M. S. 1992, 333, 445-461.
- [14] R. F. Swarttouw, The Hahn-Exton q -Bessel functions PhD Thesis The Technical University of Delft (1992).