



HAL
open science

On the Graf's addition theorem for Hahn Extton q-Bessel function

Lazhar Dhaouadi, Ahmed Fitouhi

► **To cite this version:**

Lazhar Dhaouadi, Ahmed Fitouhi. On the Graf's addition theorem for Hahn Extton q-Bessel function. Constructive Approximation, 2011, 34, pp.453-472. 10.1007/s00365-011-9132-0 . hal-00161528v2

HAL Id: hal-00161528

<https://hal.science/hal-00161528v2>

Submitted on 16 Sep 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the Graf's addition theorem for Hahn Extton q -Bessel function

Lazhar Dhaouadi* Ahmed Fitouhi†

Abstract

In this paper we study the positivity of the generalized q -translation associated with the q -Bessel Hahn Extton function which is deduced by a new formulation of the Graf's addition formula related to this function.

1 Introduction and Preliminaries

It is well known that the generalized translation operator T associated with the Bessel function of the first kind is positive in the sense if $f > 0$, then $Tf > 0$. This property is easily seen when we write Tf as an integral representation with a kernel involving the area of some triangle ([1],[9]) and has several applications in many mathematical fields such that hypergroup structure, heat equation....

In 2002, the appropriated q -generalized translation for the q -Bessel Hahn Extton function was founded [3] and the problem of its positivity asked. In literature we meet many attempts to show this property in particular case, nerveless they are no definitive response at to day. In [2] the authors prove that for the q -cosinus , the correspondent q -translation operator is positive if $q \in [0, q_0]$ for some q_0 . In this work and owing a new formulation of the Graf's addition theorem [7] we give an affirmative answer about this theme by a technic involving some inclusion of sets.

To make this work self containing, we begin by the following preliminaries. Throughout this paper we consider $0 < q < 1$ and we adopt the standard conventional notations of [4]. For complex a We put

*IPEB, 7021 Zarzouna, Bizerte,Tunisia. E-mail lazhardhaouadi@yahoo.fr

†Faculté des sciences de Tunis, 1060 Tunis, Tunisia. E-mail ahmed.fitouhi@fst.rnu.tn

$$(a; q)_0 = 1, \quad (a; q)_n = \prod_{i=0}^{n-1} (1 - aq^i), \quad n = 1 \dots \infty.$$

Jackson's q -integral (see [5]) over the interval $[0, \infty[$ is defined by

$$\int_0^\infty f(x) d_q x = (1 - q) \sum_{n=-\infty}^\infty q^n f(q^n).$$

We denote by

$$\mathbb{R}_q^+ = \{q^n, \quad n \in \mathbb{Z}\},$$

and we consider $\mathcal{L}_{q,p,v}$ the space of even functions f defined on \mathbb{R}_q^+ such that

$$\|f\|_{q,p,v} = \left[\int_0^\infty |f(x)|^p x^{2v+1} d_q x \right]^{1/p} < \infty.$$

The q -Bessel function of third kind and of order v , called also Hahn-Exton function, is defined by the q -hypergeometric function (see [8])

$$J_v(x, q) = \frac{(q^{v+1}; q)_\infty}{(q; q)_\infty} x^v {}_1\phi_1(0, q^{v+1}, q, qx^2), \quad \Re(v) > -1,$$

and has a normalized form is given by

$$j_v(x, q) = \frac{(q; q)_\infty}{(q^{v+1}; q)_\infty} x^{-v} J_v(x, q) = {}_1\phi_1(0, q^{v+1}, q, qx^2) = \sum_{n=0}^\infty (-1)^n \frac{q^{\frac{n(n+1)}{2}}}{(q; q)_n (q^{v+1}; q)_n} x^{2n}.$$

It's an entire analytic function in z .

In ([2],[6]) the q -Bessel Fourier transform $\mathcal{F}_{q,v}$ and its properties were studied in great detail, it is defined as follow

$$\mathcal{F}_{q,v} f(x) = c_{q,v} \int_0^\infty f(t) j_v(xt, q^2) t^{2v+1} d_q t,$$

where

$$c_{q,v} = \frac{1}{1 - q} \frac{(q^{2v+2}, q^2)_\infty}{(q^2, q^2)_\infty}.$$

There is many way to define the q -Bessel translation operator [2],[3]. One of them can be enounced for suitable function f as follows:

$$T_{q,x}^v f(y) = c_{q,v} \int_0^\infty \mathcal{F}_{q,v}(f)(t) j_v(xt, q^2) j_v(yt, q^2) t^{2v+1} d_q t, \quad \forall x, y \in \mathbb{R}_q^+, \forall f \in \mathcal{L}_{q,1,v}.$$

Now we say that $T_{q,x}^v$ is positive if $T_{q,x}^v f \geq 0$ for $f \geq 0$.
Let us putting the domain of positivity of $T_{q,x}^v$ by

$$Q_v = \{q \in]0, 1[, \quad T_{q,x}^v \text{ is positive for all } x \in \mathbb{R}_q^+\}.$$

Trough the result of [2], Q_v is not empty. So for $q \in Q_v$ the q -convolution product of the two functions f and $g \in \mathcal{L}_{q,1,v}$ is defined by

$$f *_q g(x) = c_{q,v} \int_0^\infty T_{q,x}^v f(y)g(y)y^{2v+1}d_q y.$$

To close this section we present the following results proved in [2] which will be used in the remainder.

Proposition 1

$$|j_v(q^n, q^2)| \leq \frac{(-q^2; q^2)_\infty (-q^{2v+2}; q^2)_\infty}{(q^{2v+2}; q^2)_\infty} \begin{cases} 1 & \text{if } n \geq 0 \\ q^{n^2 - (2v+1)n} & \text{if } n < 0 \end{cases} .$$

Theorem 1 *The operator $\mathcal{F}_{q,v}$ satisfies*

1. For all functions $f \in \mathcal{L}_{q,1,v}$,

$$\mathcal{F}_{q,v}^2 f(x) = f(x), \quad \forall x \in \mathbb{R}_q^+.$$

2. If $f \in \mathcal{L}_{q,1,v}$ and $\mathcal{F}_{q,v} f \in \mathcal{L}_{q,1,v}$ then

$$\|\mathcal{F}_{q,v} f\|_{q,v,2} = \|f\|_{q,v,2}.$$

2 The Graf's addition formula

The Graf's addition formula for Hahn-Exton q -Bessel function proved by H.T.Koelink and F. Swarttouw [7] plays a central role here . It can be stated as follows.

$$\begin{aligned} & J_v(Rq^{1/2(y+z+v)}, q)J_{x-v}(q^{1/2z}, q) \\ &= \sum_{k \in \mathbb{Z}} J_k(Rq^{1/2(x+y+k)}, q)J_{v+k}(Rq^{1/2(y+k+v)}, q)J_x(q^{1/2(z-k)}, q); \end{aligned}$$

and it is valid when $z \in \mathbb{Z}$ and $R, x, y, v \in \mathbb{C}$ satisfying

$$|R|^2 q^{1+\Re(x)+\Re(y)} < 1, \quad \Re(x) > -1, \quad R \neq 0.$$

This formula has originally been derived for $v, x, y \in \mathbb{Z}$, $R > 0$ by the interpretation of the Hahn-Exton q -Bessel function as matrix elements of irreducible unitary representation of the quantum group of plane motions. If we replace q by q^2 and R by q^r in the previous formula we get:

$$\begin{aligned} & J_v(q^{y+z+v+r}, q^2) J_{x-v}(q^z, q^2) \\ &= \sum_{k \in \mathbb{Z}} J_k(q^{x+y+k+r}, q^2) J_{v+k}(q^{y+k+v+r}, q^2) J_x(q^{z-k}, q^2), \end{aligned}$$

and put

$$m = y + z + v + r,$$

so

$$x + y + k + r = m + k + x - z - v,$$

$$y + k + v + r = m + k - z,$$

and we have

$$\begin{aligned} & J_v(q^m, q^2) J_{x-v}(q^z, q^2) \\ &= \sum_{k \in \mathbb{Z}} J_k(q^{m+k+x-z-v}, q^2) J_{v+k}(q^{m+k-z}, q^2) J_x(q^{z-k}, q^2). \end{aligned}$$

This last formula is valid for $z \in \mathbb{Z}$ and $r, x, y, v \in \mathbb{C}$ satisfying

$$1 + 2\Re(r) + \Re(x) + \Re(y) = 1 + \Re(r) + \Re(m) - \Re(z) - \Re(v) > 0, \quad \Re(x) > -1.$$

In the above sum we replace $z - k$ by k we get

$$\begin{aligned} & J_v(q^m, q^2) J_{x-v}(q^z, q^2) \\ &= \sum_{k \in \mathbb{Z}} J_{z-k}(q^{m+x-v-k}, q^2) J_{v+z-k}(q^{m-k}, q^2) J_x(q^k, q^2), \end{aligned}$$

The sum in the second member exists for

$$\forall z \in \mathbb{Z}, \quad \forall m, v, x \in \mathbb{C}, \quad \Re(x) > -1.$$

In fact there exist an infinity complex number $r \in \mathbb{C}$ for which

$$1 + \Re(r) + \Re(m) - \Re(z) - \Re(v) > 0.$$

Now using the definition of the normalized q -Bessel function

$$J_x(q^k, q^2) = \frac{(q^{2x+2}, q^2)_\infty}{(q^2, q^2)_\infty} q^{vk} j_x(q^k, q^2) = (1 - q) c_{q,x} q^{xk} j_x(q^k, q^2),$$

we obtain

$$\begin{aligned} & q^{mv+z(x-v)}(1-q)^2 c_{q,v} c_{q,x-v} j_v(q^m, q^2) j_{x-v}(q^z, q^2) \\ &= (1-q) c_{q,x} \sum_{k \in \mathbb{Z}} q^{xk} J_{z-k}(q^{x-v+m-k}, q^2) J_{v+z-k}(q^{m-k}, q^2) j_x(q^k, q^2). \end{aligned}$$

Let

$$\lambda = q^n, \quad n \in \mathbb{Z},$$

and replace

$$k \rightarrow k+n, m \rightarrow m+n, z \rightarrow z+n.$$

This implies

$$\begin{aligned} & q^{mv+z(x-v)}(1-q)^2 c_{q,v} c_{q,x-v} j_v(q^m \lambda, q^2) j_{x-v}(q^z \lambda, q^2) \\ &= (1-q) c_{q,x} \sum_{k \in \mathbb{Z}} q^{xk} J_{z-k}(q^{x-v+m-k}, q^2) J_{v+z-k}(q^{m-k}, q^2) j_x(q^k \lambda, q^2) \\ &= (1-q) c_{q,x} \sum_{k \in \mathbb{Z}} q^{2k(x+1)} q^{-k(x+2)} J_{z-k}(q^{x-v+m-k}, q^2) J_{v+z-k}(q^{m-k}, q^2) j_x(q^k \lambda, q^2). \end{aligned}$$

We put

$$E_{v,x}(q^m, q^z, q^k) = \frac{1}{(1-q)^2 c_{q,v} c_{q,x-v}} q^{-k(x+2)-mv-z(x-v)} J_{z-k}(q^{x-v+m-k}, q^2) J_{v+z-k}(q^{m-k}, q^2).$$

The Proposition 1 shows that the function:

$$\lambda \mapsto j_v(q^m \lambda, q^2) j_{x-v}(q^z \lambda, q^2), \quad \forall m, z \in \mathbb{Z},$$

belongs to the space $\mathcal{L}_{q,1,v}$. Theorem 1, part 1 leads to the following statement:

Proposition 2 For $z, m \in \mathbb{Z}$ and $x, v \in \mathbb{C}$ satisfying

$$\Re(x-v) > -1, \quad \Re(v) > -1$$

we have

$$j_v(q^m \lambda, q^2) j_{x-v}(q^z \lambda, q^2) = c_{q,x} \int_0^\infty E_{v,x}(q^m, q^z, t) j_x(\lambda t, q^2) t^{2x+1} d_q t,$$

and

$$E_{v,x}(q^m, q^z, q^k) = c_{q,x} \int_0^\infty j_v(q^m \lambda, q^2) j_{x-v}(q^z \lambda, q^2) j_x(q^k \lambda, q^2) \lambda^{2x+1} d_q \lambda.$$

3 Positivity of the q -translation operator

This section is a direct application of the previous one but before any things we recall that the q -translation operator possess the q -integral representation (see [2])

Proposition 3 *Let $f \in \mathcal{L}_{q,1,v}$ then*

$$T_{q,x}^v f(y) = \int_0^\infty f(z) D_v(x, y, z) z^{2v+1} d_q z,$$

where

$$D_v(x, y, z) = c_{q,v}^2 \int_0^\infty j_v(xt, q^2) j_v(yt, q^2) j_v(zt, q^2) t^{2v+1} d_q t.$$

When q tends to 1^- we obtain at least formally the classical one and the q -kernel $D_v(x, y, z)$ tends to the classical one ([1],[9]) which involves the area of some triangle. Hence the positivity of $T_{q,x}^v$ is subject of those of $D_v(x, y, z)$. A direct consequence of the above result is the fact that if the operator $T_{q,x}^v$ is positive and $f \in \mathcal{L}_{q,1,v}$ then $T_{q,x}^v f \in \mathcal{L}_{q,1,v}$ (in general it is not true without this hypothesis). Indeed

$$\begin{aligned} & \int_0^\infty |T_{q,x}^v f(y)| y^{2v+1} d_q t \\ & \leq \int_0^\infty T_{q,x}^v |f|(y) y^{2v+1} d_q t = \int_0^\infty |f(z)| \left[\int_0^\infty D_v(x, y, z) y^{2v+1} d_q y \right] z^{2v+1} d_q z. \end{aligned}$$

Putting

$$\phi : t \mapsto j_v(xt, q^2) j_v(zt, q^2),$$

then we can write

$$D_v(x, y, z) = c_{q,v} \mathcal{F}_{q,v} \phi(y).$$

This gives by the of the inversion formula in Theorem 1 the important result as the classical; one

$$\int_0^\infty D_v(x, y, z) y^{2v+1} d_q y = \mathcal{F}_{q,v}^2 \phi(0) = \phi(0) = 1.$$

Then we have

$$\int_0^\infty |T_{q,x}^v f(y)| y^{2v+1} d_q t \leq \|f\|_{q,v,1},$$

and we obtain

$$f \in \mathcal{L}_{q,1,v} \Rightarrow T_{q,x}^v f \in \mathcal{L}_{q,1,v}.$$

In [2] and as a first preamble of this theme the authors proved that

$$\begin{aligned} D_{-\frac{1}{2}}(q^m, q^r, q^k) &= \frac{q^{2(r-m)(k-m)-m}}{(1-q)(q; q)_I} (q^{2(r-m)+1}; q)_{\infty} {}_1\phi_1(0, q^{2(r-m)+1}, q; q^{2(k-m)+1}) \\ &= \frac{1}{1-q} q^{-m} J_{2(r-m)}(q^{k-m}, q), \end{aligned}$$

which implies that the correspondent domain of $T_{q,x}^{-\frac{1}{2}}$ is given by

$$Q_{-\frac{1}{2}} =]0, q_0],$$

where q_0 is the first zero of the q -hypergeometric function:

$$q \mapsto {}_1\phi_1(0, q, q, q);$$

a second statement is easily given by the Proposition 2 with $x = v = 0$ which gives

$$\begin{aligned} E_{0,0}(q^m, q^z, q^k) &= c_{q,0} \int_0^{\infty} j_0(q^m \lambda, q^2) j_0(q^z \lambda, q^2) j_0(q^k \lambda, q^2) \lambda d_q \lambda \\ &= \frac{1}{c_{q,0}} D_0(q^m, q^z, q^k), \end{aligned}$$

and then

$$\begin{aligned} D_0(q^m, q^z, q^k) &= c_{q,0} E_{0,0}(q^m, q^z, q^k) \\ &= \frac{1}{(1-q)} q^{-2k} \left[J_{z-k}(q^{m-k}, q^2) \right]^2, \end{aligned}$$

Hence we have

$$Q_0 =]0, 1[.$$

Now we explicit the kernel in the production formula in terms of D_v .

Proposition 4 For $n, m, k \in \mathbb{Z}$ and $-1 < v$ we have

$$E_{v,v}(q^m, q^n, q^k) = (1-q) \sum_{i=0}^{\infty} \frac{(q^{-2v}, q^2)_i}{(q^2, q^2)_i} q^{i(2+2v)} D_v(q^m, q^{i+n}, q^k).$$

Proof. From the following formula (see[7], §5) “which can be proved in a straightforward way by substitution of the defining series for the q -Bessel

functions on both sides, by interchanging summations, and by evaluating the q-binomial series which occurs”

$$J_{x-v}(\lambda, q^2) = \lambda^{-v} \sum_{i=0}^{\infty} \frac{(q^{-2v}, q^2)_i}{(q^2, q^2)_i} q^{i(2+x)} J_x(\lambda q^i, q^2),$$

where $\Re(x-v) > -1$ and $\Re(x) > -1$ we obtain

$$(1-q)c_{q,x-v} j_{x-v}(\lambda, q^2) = (1-q)c_{q,x} \sum_{i=0}^{\infty} \frac{(q^{-2v}, q^2)_i}{(q^2, q^2)_i} q^{i(2+2x)} j_x(\lambda q^i, q^2).$$

Put $x = v$ and change λ by $q^n \lambda$ we obtain

$$j_0(q^n \lambda, q^2) = (1-q)c_{q,v} \sum_{i=0}^{\infty} \frac{(q^{-2v}, q^2)_i}{(q^2, q^2)_i} q^{i(2+2v)} j_v(\lambda q^{i+n}, q^2),$$

which gives from Proposition 2

$$\begin{aligned} E_{v,v}(q^m, q^n, q^k) &= c_{q,v} \int_0^{\infty} j_v(q^m \lambda, q^2) j_0(q^n \lambda, q^2) j_v(q^k \lambda, q^2) \lambda^{2v+1} d_q \lambda \\ &= (1-q) \sum_{i=0}^{\infty} \frac{(q^{-2v}, q^2)_i}{(q^2, q^2)_i} q^{i(2+2v)} \\ &\quad \times \left[c_{q,v}^2 \int_0^{\infty} j_v(q^m \lambda, q^2) j_v(q^{i+n} \lambda, q^2) j_v(q^k \lambda, q^2) \lambda^{2v+1} d_q \lambda \right] \\ &= (1-q) \sum_{i=0}^{\infty} \frac{(q^{-2v}, q^2)_i}{(q^2, q^2)_i} q^{i(2+2v)} D_v(q^m, q^{i+n}, q^k). \end{aligned}$$

We justify the exchange of the signs sum and integral by

$$\begin{aligned} &\sum_{i=0}^{\infty} \frac{|(q^{-2v}, q^2)_i|}{(q^2, q^2)_i} q^{i(2+2v)} \int_0^{\infty} \left| j_v(q^m \lambda, q^2) j_v(q^{i+n} \lambda, q^2) j_v(q^k \lambda, q^2) \lambda^{2v+1} \right| d_q \lambda \\ &\leq \|j_v(\cdot, q^2)\|_{q,\infty}^2 \|j_v(\cdot, q^2)\|_{q,1,v} q^{-2(v+1)m} \sum_{i=0}^{\infty} \frac{|(q^{-2v}, q^2)_i|}{(q^2, q^2)_i} q^{i(2+2v)} < \infty. \end{aligned}$$

So we obtain the result. ■

Proposition 5 *Let $-1 < v < 0$ then*

$$Q_v \subsetneq]0, 1[.$$

Proof. For $m, n, k \in \mathbb{Z}$ and $-1 < v < 0$ we have

$$\begin{aligned} E_{v,v}(q^m, q^n, q^k) &= \frac{1}{(1-q)c_{q,v}} q^{-k(v+2)-mv} J_{n-k}(q^{m-k}, q^2) J_{v+n-k}(q^{m-k}, q^2) \\ &= (1-q) \sum_{i=0}^{\infty} \frac{(q^{-2v}, q^2)_i}{(q^2, q^2)_i} q^{i(2+2v)} D_v(q^m, q^{i+n}, q^k). \end{aligned}$$

In particular for $m = n = k = 0$


$$E_{v,v}(1, 1, 1) = \frac{1}{(1-q)c_{q,v}} J_0(1, q^2) J_v(1, q^2).$$

We introduce the following function

$$\phi_v : q \mapsto (q^2, q^2)_{\infty}^2 J_v(1, q^2).$$

Let $q_1 \simeq 0.658$ the first zero of ϕ_0 and consider the graph of the function

$$v \mapsto \phi_v(q_1), \quad v \in [0, 1].$$



We conclude that for $-1 < v < 0$ we have $\phi_v(q_1) < 0$. Then there exist a very small $\varepsilon > 0$ such that $\phi_v(q_1 - \varepsilon) < 0$ and $\phi_0(q_1 - \varepsilon) > 0$. Hence, this function

$$q \mapsto J_0(1, q^2) J_v(1, q^2) < 0,$$

takes some negative values in the interval $]0, 1[$, then for some entire $i \in \mathbb{N}$

$$\frac{(q^{-2v}, q^2)_i}{(q^2, q^2)_i} q^{i(2+2v)} D_v(1, q^i, 1) < 0.$$

As

$$\frac{(q^{-2v}, q^2)_i}{(q^2, q^2)_i} q^{i(2+2v)} > 0,$$

then $D_v(1, q^i, 1) < 0$ for some entire $i \in \mathbb{N}$. Which prove that

$$Q_v \not\subseteq]0, 1[.$$

This finish the proof. ■

Lemma 1 For $x \in \mathbb{R}_q^+$ and $\Re(v+t) > -1$ we have

$$(1-q)c_{q,v} \sum_{n \in \mathbb{Z}} q^{(1+v+t)n} j_v(q^n x, q^2) = \frac{(q^{1+v-t}; q^2)_\infty}{(q^{1+v+t}; q^2)_\infty} x^{-(1+v+t)}.$$

Proof. In [6] the following result was proved

$$\sum_{n \in \mathbb{Z}} q^{(1+t)n} J_v(q^n, q^2) = \frac{(q^{1+v-t}; q^2)_\infty}{(q^{1+v+t}; q^2)_\infty}, \quad \Re(v+t) > -1.$$

Then we get

$$(1-q)c_{q,v} \sum_{n \in \mathbb{Z}} q^{(1+v+t)n} j_v(q^n, q^2) = \frac{(q^{1+v-t}; q^2)_\infty}{(q^{1+v+t}; q^2)_\infty}.$$

Hence for $x = q^k \in \mathbb{R}_q^+$ we have

$$(1-q)c_{q,v} \sum_{n \in \mathbb{Z}} q^{(1+v+t)(n+k)} j_v(q^{n+k}, q^2) = \frac{(q^{1+v-t}; q^2)_\infty}{(q^{1+v+t}; q^2)_\infty}.$$

This finish the proof. ■

Theorem 2 Let $v \geq x > -1$ then

$$Q_x \subset Q_v.$$

As consequence

- If $0 \leq v$ then $Q_v =]0, 1[$.
- If $-\frac{1}{2} \leq v < 0$ then $]0, q_0] \subset Q_v \subsetneq]0, 1[$.
- If $-1 < v \leq -\frac{1}{2}$ then $Q_v \subset]0, q_0]$.

Proof. Let $v > -1$ and $0 < \mu < 1$. We will prove that

$$Q_v \subset Q_{v+\mu},$$

which suffices to show that $Q_x \subset Q_v$ if $x \leq v$. In the following we tack $q \in Q_v$. We have

$$c_{q,v+\mu} j_{v+\mu}(t, q^2) = c_{q,v} \sum_{i=0}^{\infty} \frac{(q^{2\mu}, q^2)_i}{(q^2, q^2)_i} q^{2(v+1)i} j_v(tq^i, q^2),$$

and then

$$\begin{aligned}
& c_{q,v} c_{q,v+\mu}^2 j_v(tx, q^2) j_{v+\mu}(ty, q^2) j_v(tz, q^2) \\
&= c_{q,v}^3 \sum_{i,j,s=0}^{\infty} \frac{(q^{2\mu}, q^2)_i}{(q^2, q^2)_i} \frac{(q^{2\mu}, q^2)_j}{(q^2, q^2)_j} q^{2(1+v)(i+j)} \\
&\quad \times j_v(tx, q^2) j_v(tq^i y, q^2) j_v(tq^j z, q^2),
\end{aligned}$$

which implies

$$\begin{aligned}
& c_{q,v} T_{v,v+\mu,v}(x, y, z) \\
&= c_{q,v} \sum_{i,j=0}^{\infty} \frac{(q^{2\mu}, q^2)_i}{(q^2, q^2)_i} \frac{(q^{2\mu}, q^2)_j}{(q^2, q^2)_j} q^{2(1+v)(i+j)} D_v(x, q^i y, q^j z) \geq 0,
\end{aligned}$$

where

$$T_{v,w,\alpha}(x, y, z) = c_{q,w}^2 \int_0^\infty j_v(tx, q^2) j_w(ty, q^2) j_w(tz, q^2) t^{2\alpha+1} d_q t.$$

Note that

$$T_{v,v,v}(x, y, z) = D_v(x, y, z).$$

The exchange of the signs sum and integral is valid since:

$$\begin{aligned}
& \sum_{i,j=0}^{\infty} q^{2(1+v)(i+j)} \int_0^\infty |j_v(q^m t, q^2)| |j_v(q^{n+i} t, q^2)| |j_v(q^{k+j} t, q^2)| t^{2v+1} d_q t \\
&= \sum_{i,j=0}^{\infty} q^{2(1+v)(i+j)} \int_0^1 |j_v(q^m t, q^2)| |j_v(q^{n+i} t, q^2)| |j_v(q^{k+j} t, q^2)| t^{2v+1} d_q t \\
&+ \sum_{i,j=0}^{\infty} q^{2(1+v)(i+j)} \int_1^\infty |j_v(q^m t, q^2)| |j_v(q^{n+i} t, q^2)| |j_v(q^{k+j} t, q^2)| t^{2v+1} d_q t.
\end{aligned}$$

Note that

$$\begin{aligned}
& \sum_{i,j=0}^{\infty} q^{2(1+v)(i+j)} \int_0^1 |j_v(q^m t, q^2)| |j_v(q^{n+i} t, q^2)| |j_v(q^{k+j} t, q^2)| t^{2v+1} d_q t \\
&\leq \|j_v(\cdot, q^2)\|_{q,\infty}^3 \left[\sum_{i,j=0}^{\infty} q^{2(1+v)(i+j)} \right] \int_0^1 t^{2v+1} d_q t < \infty,
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{i,j=0}^{\infty} q^{2(1+v)(i+j)} \int_1^{\infty} |j_v(q^m t, q^2)| |j_v(q^{n+i} t, q^2)| |j_v(q^{k+j} t, q^2)| t^{2v+1} d_q t \\
&= (1-q) \sum_{i,j,r'=0}^{\infty} q^{2(1+v)(i+j-r)} |j_v(q^{m-r}, q^2)| |j_v(q^{n+i-r}, q^2)| |j_v(q^{k+j-r}, q^2)| \\
&= \sum_{j,r'=0}^{\infty} q^{2(1+v)j} |j_v(q^{m-r}, q^2)| |j_v(q^{k+j-r}, q^2)| \times \left[(1-q) \sum_{i=0}^{\infty} q^{2(1+v)(i-r)} |j_v(q^{n+i-r}, q^2)| \right],
\end{aligned}$$

where $r' = r - 1$. We write

$$\begin{aligned}
\sum_{i=0}^{\infty} q^{2(1+v)(i-r)} |j_v(q^{n+i-r}, q^2)| &= q^{-2(1+v)n} (1-q) \sum_{i=0}^{\infty} q^{2(n+i-r)} |j_v(q^{n+i-r}, q^2)| \\
&= q^{-2(1+v)n} (1-q) \sum_{i=m-r}^{\infty} q^{2(1+v)i} |j_v(q^i, q^2)| \\
&< q^{-2(1+v)n} \|j_v(\cdot, q^2)\|_{q,1,v}.
\end{aligned}$$

Then

$$\begin{aligned}
& \sum_{j,r'=0}^{\infty} q^{2(1+v)j} |j_v(q^{m-r}, q^2)| |j_v(q^{k+j-r}, q^2)| \\
&= \sum_{s,r'=0}^{\infty} |j_v(q^{m-r}, q^2)| \left[\sum_{j=0}^{\infty} q^{2(1+v)j} |j_v(q^{k+j-r}, q^2)| \right] \\
&= \sum_{r'=0}^{\infty} |j_v(q^{m-r}, q^2)| \left[q^{2(1+v)(r-k)} \sum_{j=0}^{\infty} q^{2(k+j-r)} |j_0(q^{k+j-r}, q^2)| \right] \\
&= \sum_{s,r'=0}^{\infty} q^{2s} |j_v(q^{m-r}, q^2)| \left[q^{2(v+1)(r-k)} \sum_{j=k-r}^{\infty} q^{2(1+v)j} |j_0(q^j, q^2)| \right] \\
&\leq \frac{q^{-2(v+1)k}}{1-q} \|j_v(\cdot, q^2)\|_{q,1,v} \|j_v(\cdot, q^2)\|_{q,\infty} \sum_{r'=0}^{\infty} q^{2(v+1)r} < \infty.
\end{aligned}$$

Let $0 < \alpha < \mu$. We introduce the function $A_{\alpha,\mu,v}(x)$ as follows:

$$A_{\alpha,\mu,v}(x) = c_{q,v} \int_0^{\infty} t^{2\alpha} j_{v+\mu}(t, q^2) j_v(xt, q^2) t^{2v+1} d_q t.$$

From the inversion formula in Theorem 1 ($t \mapsto t^{2\alpha} j_{v+\mu}(t, q^2) \in \mathcal{L}_{q,1,v}$) we get

$$c_{q,v} \int_0^\infty A_{\alpha,\mu,v}(x) j_v(xt, q^2) x^{2v+1} d_q x = t^{2\alpha} j_{v+\mu}(t, q^2).$$

Let $x \leq 1$. From Lemma 1 we obtain

$$\begin{aligned} & A_{\alpha,\mu,v}(x) \\ &= c_{q,v}(1-q) \sum_{n \in \mathbb{Z}} q^{2(\alpha+v+1)n} j_v(xq^n, q^2) j_{v+\mu}(q^n, q^2) \\ &= c_{q,v}(1-q) \sum_{n \in \mathbb{Z}} \left[\sum_{i=0}^\infty (-1)^i \frac{q^{i(i+1)}}{(q^{2+2v}, q^2)_i (q^2, q^2)_i} \right] q^{2(\alpha+v+1+i)n} j_{v+\mu}(q^n, q^2) \\ &= c_{q,v}(1-q) \sum_{i=0}^\infty (-1)^i \frac{q^{i(i+1)}}{(q^{2+2v}, q^2)_i (q^2, q^2)_i} x^{2i} \left[\sum_{n \in \mathbb{Z}} q^{2(\alpha+v+1+i)n} j_{v+\mu}(q^n, q^2) \right] \\ &= \frac{c_{q,v}}{c_{q,v+\mu}} \sum_{i=0}^\infty (-1)^i \frac{q^{i(i+1)}}{(q^{2+2v}, q^2)_i (q^2, q^2)_i} x^{2i} \frac{(q^{2(1+v+\mu)-2(\alpha+v+1+i)}, q^2)_\infty}{(q^{2(\alpha+v+1+i)}, q^2)_\infty} \\ &= \frac{c_{q,v}}{c_{q,v+\mu}} \sum_{i=0}^\infty (-1)^i \frac{q^{i(i+1)}}{(q^{2+2v}, q^2)_i (q^2, q^2)_i} x^{2i} \frac{(q^{2(\mu-\alpha)} q^{-2i}, q^2)_\infty}{(q^{2(\alpha+v+1+i)}, q^2)_\infty} \geq 0. \end{aligned}$$

Note that

$$0 < \mu - \alpha < 1.$$

The exchange of the signs sum is valid since. Indeed, let $q^k = x$ then we have

$$\begin{aligned} & \sum_{i=0}^\infty q^{i(i+1)} q^{2ik} \left[\sum_{n \in \mathbb{Z}} q^{2(\alpha+v+1+i)n} |j_{v+\mu}(q^n, q^2)| \right] \\ &= \sum_{i=0}^\infty q^{i(i+1)} q^{2ik} \left[\sum_{n=0}^\infty q^{2(\alpha+v+1+i)n} |j_{v+\mu}(q^n, q^2)| \right] \\ &+ \sum_{i=0}^\infty q^{i(i+1)} q^{2ik} \left[\sum_{n=1}^\infty q^{-2(\alpha+v+1+i)n} |j_{v+\mu}(q^{-n}, q^2)| \right] \\ &\leq \sum_{i=0}^\infty q^{i(i+1)} q^{2ik} \left[\sum_{n=0}^\infty q^{2(\alpha+v+1)n} \right] \\ &+ \sum_{i=0}^\infty q^{i(i+1)} q^{2ik} \left[\sum_{n=1}^\infty q^{-2(\alpha+v+1+i)n+n^2+(2v+2\mu+1)n} \right]. \end{aligned}$$

One has to observe that the first sum exist. The second sum also:

$$\begin{aligned}
& \sum_{i=0}^{\infty} q^{i(i+1)} q^{2ik} \left[\sum_{n=1}^{\infty} q^{-2(\alpha+v+1+i)n+n^2+(2v+2\mu+1)n} \right] \\
&= \sum_{n=1}^{\infty} q^{-2(\alpha+v+1+i)n+n^2+(2v+2\mu+1)n} \left[\sum_{i=0}^{\infty} q^{i(i+1)} q^{2ik} \right] \\
&= \sum_{n=1}^{\infty} q^{-2(\alpha+v+1)n+(2v+2\mu+1)n+(2k+1)n} \left[\sum_{i=0}^{\infty} q^{(i-n)^2} q^{(2k+1)(i-n)} \right] \\
&= \sum_{n=1}^{\infty} q^{2(\mu-\alpha)n+(2k+1)n} \left[\sum_{i=-n}^{\infty} q^{i^2} q^{(2k+1)i} \right] \leq \sum_{n=1}^{\infty} q^{2(\mu-\alpha)n+(2k+1)n} \left[\sum_{i=-\infty}^{\infty} q^{i^2} q^{(2k+1)i} \right] < \infty.
\end{aligned}$$

If $x > 1$ then we obtain

$$\begin{aligned}
& A_{\alpha,\mu,v}(x) \\
&= c_{q,v}(1-q) \sum_{n \in \mathbb{Z}} q^{2(\alpha+v+1)n} j_v(xq^n, q^2) j_{v+\mu}(q^n, q^2) \\
&= c_{q,v}(1-q) \sum_{n \in \mathbb{Z}} \left[\sum_{i=0}^{\infty} (-1)^i \frac{q^{i(i+1)}}{(q^{2+2v}, q^2)_i (q^2, q^2)_i} \right] q^{2(\alpha+v+1+i)n} j_v(q^n x, q^2) \\
&= c_{q,v}(1-q) \sum_{i=0}^{\infty} (-1)^i \frac{q^{i(i+1)}}{(q^{2+2v+2\mu}, q^2)_i (q^2, q^2)_i} \left[\sum_{n \in \mathbb{Z}} q^{2(\alpha+v+1+i)n} j_v(q^n x, q^2) \right] \\
&= x^{-2(\alpha+v+1)} \sum_{i=0}^{\infty} (-1)^i \frac{q^{i(i+1)}}{(q^{2+2v+2\mu}, q^2)_i (q^2, q^2)_i} x^{-2i} \frac{(q^{2(1+v)-2(\alpha+v+1+i)}, q^2)_{\infty}}{(q^{2(\alpha+v+1+i)}, q^2)_{\infty}} \\
&= x^{-2(\alpha+v+1)} \sum_{i=0}^{\infty} (-1)^i \frac{q^{i(i+1)}}{(q^{2+2v+2\mu}, q^2)_i (q^2, q^2)_i} x^{-2i} \frac{(q^{-2\alpha} q^{-2i}, q^2)_{\infty}}{(q^{2(\alpha+v+1+i)}, q^2)_{\infty}} < 0.
\end{aligned}$$

Now, we write

$$\begin{aligned}
& c_{q,v} \int_0^{\infty} A_{\alpha,\mu,v}(x) T_{v,v+\mu,v}(x, y, z) x^{2v+1} d_q x \\
&= c_{q,v} \int_0^{\infty} A_{\alpha,\mu,v}(x) \left[c_{q,v}^2 \int_0^{\infty} j_w(xt, q^2) j_v(yt, q^2) j_v(zt, q^2) t^{2v+1} d_q t \right] x^{2v+1} d_q x \\
&= c_{q,v+\mu}^2 \int_0^{\infty} \left[c_{q,v} \int_0^{\infty} A_{\alpha,\mu,v}(x) j_v(xt, q^2) x^{2v+1} d_q x \right] j_{v+\mu}(yt, q^2) j_{v+\mu}(zt, q^2) t^{2v+1} d_q t \\
&= c_{q,v+\mu}^2 \int_0^{\infty} j_{v+\mu}(t, q^2) j_{v+\mu}(yt, q^2) j_{v+\mu}(zt, q^2) t^{2(v+\alpha)+1} d_q t = T_{v+\mu,v+\mu,v+\alpha}(1, y, z).
\end{aligned}$$

To justify the exchange of the signs integrals we write

$$\begin{aligned} & \int_0^\infty \left[\int_0^\infty |A_{\alpha,\mu,v}(x)| |j_v(xt, q^2)| x^{2v+1} d_q x \right] |j_v(yt, q^2)| |j_v(zt, q^2)| t d_q t \\ & \leq \left[\|j_v(\cdot, q^2)\|_{q,\infty}^2 \|j_v(\cdot, q^2)\|_{q,1,v} \|A_{\alpha,\mu,v}\|_{q,1,v} \right] \frac{1}{z}. \end{aligned}$$

Not that $x \mapsto A_{\alpha,\mu,v}(x)$ is continued at 0 and

$$|A_{\alpha,\mu,v}(x)| \leq \left[\|j_{v+\mu}(\cdot, q^2)\|_{q,\infty} \|j_v(\cdot, q^2)\|_{q,1,\alpha+v} \right] x^{-2(\alpha+v+1)}, \quad \text{as } x \rightarrow \infty.$$

On the other hand

$$\int_0^\infty A_{\alpha,\mu,v}(x) x^{2v+1} d_q x = 0,$$

then we obtain for all $\delta > 0$

$$T_{v+\mu,v+\mu,v+\alpha}(1, y, z) = \int_0^\infty A_{\alpha,\mu,v}(x) \left[T_{v,v+\mu,v}(x, y, z) - \delta \right] x^{2v+1} d_q x.$$

In the following we assume that $0 < y, z \leq 1$. Let

$$\delta_0 = \inf_{0 \leq y, z \leq 1} T_{v,v+\mu,v}(1, y, z)$$

Not that δ_0 exist and strictly positive, indeed the following function

$$(y, z) \mapsto T_{v,v+\mu,v}(1, y, z)$$

is continuous on the compact $[0, 1] \times [0, 1]$. Hence, there exist

$$(y_0, z_0) \in [0, 1] \times [0, 1]$$

such that

$$\delta_0 = T_{v,v+\mu,v}(1, y_0, z_0) = \inf_{0 \leq y, z \leq 1} T_{v,v+\mu,v}(1, y, z) \geq 0.$$

If we assume that $T_{v,v+\mu,v}(1, y_0, z_0) = 0$ then

$$\sum_{i,j=0}^{\infty} \frac{(q^{2\mu}, q^2)_i (q^{2\mu}, q^2)_j}{(q^2, q^2)_i (q^2, q^2)_j} q^{2(1+v)(i+j)} D_v(1, q^i y_0, q^j z_0) = 0,$$

which implies that

$$c_{q,v} \mathcal{F}_{q,v} \left[t \mapsto j_v(t, q^2) j_v(z_0 t, q^2) \right] (q^i y_0) = D_v(1, q^i y_0, z_0) = 0, \quad \forall i \in \mathbb{N}.$$

From Proposition 1 and the fact that there exist $\sigma_0 > 0$ such that

$$|j_v(z, q^2)| \leq \sigma_0 e^{|z|}, \quad \forall z \in \mathbb{C}$$

we see that this function

$$z \mapsto \mathcal{F}_{q,v} \left[t \mapsto j_v(t, q^2) j_v(z_0 t, q^2) \right] (z)$$

is analytic. Then for all $x \in \mathbb{R}_q$

$$\mathcal{F}_{q,v} \left[t \mapsto j_v(t, q^2) j_v(z_0 t, q^2) \right] (x) = 0 \Rightarrow j_v(t, q^2) j_v(z_0 t, q^2) = 0, \quad \forall t \in \mathbb{R}_q,$$

but this is absurd. Then $\delta_0 > 0$. Now we have

$$\delta_0 \leq T_{v,v+\mu,v}(x, y, z), \quad \forall 0 < x \leq 1.$$

If

$$\delta_0 > T_{v,v+\mu,v}(x, y, z), \quad \forall x > 1.$$

then

$$\begin{aligned} T_{v+\mu,v+\mu,v+\alpha}(1, y, z) &= \int_0^1 A_{\alpha,\mu,v}(x) \left[T_{v,v+\mu,v}(x, y, z) - \delta_0 \right] x^{2v+1} d_q x \\ &\quad + \int_1^\infty A_{\alpha,\mu,v}(x) \left[T_{v,v+\mu,v}(x, y, z) - \delta_0 \right] x^{2v+1} d_q x \geq 0. \end{aligned}$$

Otherwise, there exist $s > 0$ such that

$$\delta_0 > T_{v,v+\mu,v}(x, y, z), \quad \forall x > s.$$

because

$$|T_{v,v+\mu,v}(x, y, z)| < c_{q,v+\mu}^2 \|j_{v+\mu}(\cdot, q^2)\|_{q,\infty}^2 \times \|j_v(\cdot, q^2)\|_{q,v,1} x^{-2(v+1)} \rightarrow 0, \quad \text{as } x \rightarrow \infty.$$

For $\delta \geq \delta_0$ we obtain

$$\begin{aligned} T_{v+\mu,v+\mu,v+\alpha}(1, y, z) &= \int_0^1 A_{\alpha,\mu,v}(x) \left[T_{v,v+\mu,v}(x, y, z) - \delta \right] x^{2v+1} d_q x \\ &\quad + \int_1^s A_{\alpha,\mu,v}(x) \left[T_{v,v+\mu,v}(x, y, z) - \delta \right] x^{2v+1} d_q x \\ &\quad + \int_s^\infty A_{\alpha,\mu,v}(x) \left[T_{v,v+\mu,v}(x, y, z) - \delta \right] x^{2v+1} d_q x \\ &= I_1(\delta) + I_2(\delta) + I_3(\delta). \end{aligned}$$

Note that

$$\begin{cases} \delta \mapsto I_1(\delta) \text{ is a decreasing function tends towards } -\infty \text{ and } I_1(\delta_0) > 0 \\ \delta \mapsto I_2(\delta) \text{ is an increasing function tends towards } +\infty \text{ and } I_2(\delta_0) < 0 \\ \delta \mapsto I_3(\delta) \text{ is an increasing function tends towards } +\infty \text{ and } I_3(\delta_0) > 0 \end{cases}$$

then there exist $\delta > \delta_0$ such that

$$I_1(\delta) + I_2(\delta) = 0$$

which implies

$$T_{v+\mu, v+\mu, v+\alpha}(1, y, z) = I_3(\delta) > 0.$$

In the end

$$\begin{aligned} & \lim_{\alpha \rightarrow \mu} [T_{v+\mu, v+\mu, v+\alpha}(1, y, z)] \\ &= \lim_{\alpha \rightarrow \mu} \left[c_{q, v+\mu}^2 \int_0^\infty j_{v+\mu}(t, q^2) j_{v+\mu}(yt, q^2) j_{v+\mu}(zt, q^2) t^{2(v+\alpha)+1} d_q t \right] \\ &= c_{q, v+\mu}^2 \int_0^\infty j_{v+\mu}(t, q^2) j_{v+\mu}(yt, q^2) j_{v+\mu}(zt, q^2) \left(\lim_{\alpha \rightarrow \mu} t^{2(v+\alpha)+1} \right) d_q t \\ &= T_{v+\mu, v+\mu, v+\alpha}(1, y, z) = D_{v+\mu}(1, y, z) \geq 0. \end{aligned}$$

It is not hard to justify the exchange of the signs integral and limit, in fact

$$\begin{aligned} & \left| j_{v+\mu}(t, q^2) j_{v+\mu}(yt, q^2) j_{v+\mu}(zt, q^2) t^{2(v+\alpha)+1} \right| \\ & \leq \left\| t \mapsto j_{v+\mu}(yt, q^2) j_{v+\mu}(zt, q^2) t^{2(v+\alpha)+1} \right\|_{q, \infty} \times |j_{v+\mu}(t, q^2)|. \end{aligned}$$

From the following identity

$$D_{v+\mu}(x, y, z) = x^{-2(v+\mu+1)} D_{v+\mu} \left(1, \frac{y}{x}, \frac{z}{x} \right),$$

we deduce that

$$D_{v+\mu}(x, y, z) \geq 0, \quad \forall x, y, z \in \mathbb{R}_q^+.$$

The fact that

$$Q_{-\frac{1}{2}} =]0, q_0] \quad \text{and} \quad Q_0 =]0, 1[$$

leads to the result. ■

References

- [1] F.M.Cholewinski and D.T.Haimo, The Weierstrass Hankel convolution transform, *J. Analyse Math.*17, 1966p.1-58.
- [2] L. Dhaouadi, A. Fitouhi and J. El Kamel, Inequalities in q -Fourier Analysis, *Journal of Inequalities in Pure and Applied Mathematics*, Volume 7, Issue 5, Article 171, 2006.
- [3] A. Fitouhi, M.Hamza and F. Bouzeffour, The $q - j_\alpha$ Bessel function *J. Appr. Theory.*115,144-166(2002).
- [4] G. Gasper and M. Rahman, Basic hypergeometric series, *Encyclopedia of mathematics and its applications* 35, Cambridge university press, 1990.
- [5] F. H. Jackson, On a q -Definite Integrals, *Quarterly Journal of Pure and Application Mathematics* 41, 1910, 193-203.
- [6] T. H. Koornwinder and R. F. Swarttouw, On q -Analogues of the Hankel and Fourier Transform, *Trans. A. M. S.* 1992, 333, 445-461.
- [7] H. T. Koelink and R. F. Swarttouw, A q -Analogue of Graf's Addition Formula for the Hahn-Exton q -Bessel Function, *Journal of Approximation theory* 81. 260-273(1995).
- [8] R. F. Swarttouw, The Hahn-Exton q -Bessel functions PhD Thesis The Technical University of Delft ((1992).
- [9] G. N. Watson, A treatise of the theory of Bessel functions . Second edition, Cambridge University Press, London, New York, 1966.