



HAL
open science

Functions of q -positive type

Lazhar Dhaouadi

► **To cite this version:**

Lazhar Dhaouadi. Functions of q -positive type. Positivity, 2012, 16, pp.107-129. 10.1007/s11117-011-0115-y . hal-00162012

HAL Id: hal-00162012

<https://hal.science/hal-00162012>

Submitted on 12 Jul 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Functions of q -positive type

Lazhar Dhaouadi *

Abstract

In this paper we characterize the subspace of $\mathcal{L}_{q,1,v}$ of function which are the q -Bessel Fourier transform of positive functions in $\mathcal{L}_{q,1,v}$. As application we give a q -version of the Bochner's theorem.

1 Introduction and Preliminaries

Given a positive finite Borel measure μ on the real line \mathbb{R} , the Fourier transform Q of μ is the continuous function

$$Q(x) = \int_{\mathbb{R}} e^{-itx} d\mu(t).$$

The function Q is a positive definite function, i.e for any finite list of complex numbers z_1, \dots, z_n and real numbers x_1, \dots, x_n

$$\sum_{r=1}^n \sum_{l=1}^n z_r \bar{z}_l Q(x_r - x_l) \geq 0.$$

Bochner's theorem says the converse is true, i.e. every positive definite function Q is the Fourier transform of a positive finite Borel measure. In q -Fourier analysis, semelar phenomenon will appear. It is the subject of our article.

In the following we consider $0 < q < 1$ and we adopt the standard conventional notations of [2]. We put

$$\mathbb{R}_q^+ = \{q^n, \quad n \in \mathbb{Z}\},$$

*Institut Préparatoire aux Etudes d'Ingénieur de Bizerte (Université du 7 novembre Carthage). Route Menzel Abderrahmene Bizerte, 7021 Zarzouna, Tunisia. E-mail lazhardhaouadi@yahoo.fr

the set of q -real numbers and for complex a

$$(a; q)_0 = 1, \quad (a; q)_n = \prod_{i=0}^{n-1} (1 - aq^i), \quad n = 1 \dots \infty.$$

Jackson's q -integral (see [3]) in the interval $[0, \infty[$ is defined by

$$\int_0^\infty f(x) d_q x = (1 - q) \sum_{n=-\infty}^\infty q^n f(q^n).$$

Let $\mathcal{C}_{q,0}$ and $\mathcal{C}_{q,b}$ denote the spaces of functions defined on \mathbb{R}_q^+ continued at 0, which are respectively vanishing at infinity and bounded. These spaces are equipped with the topology of uniform convergence, and by $\mathcal{L}_{q,p,v}$ the space of functions f defined on \mathbb{R}_q^+ such that

$$\|f\|_{q,p,v} = \left[\int_0^\infty |f(x)|^p x^{2v+1} d_q x \right]^{1/p} < \infty.$$

The q -exponential function is defined by

$$e(z, q) = \sum_{n=0}^\infty \frac{z^n}{(q, q)_n} = \frac{1}{(z; q)_\infty}, \quad |z| < 1.$$

The normalized Hahn-Exton q -Bessel function of order $v > -1$ (see [5]) is defined by

$$j_v(z, q) = \sum_{n=0}^\infty (-1)^n \frac{q^{\frac{n(n-1)}{2}}}{(q, q)_n (q^{v+1}, q)_n} z^n.$$

The q -Bessel Fourier transform $\mathcal{F}_{q,v}$ introduced in [1,4] as follow

$$\mathcal{F}_{q,v} f(x) = c_{q,v} \int_0^\infty f(t) j_v(xt, q^2) t^{2v+1} d_q t,$$

where

$$c_{q,v} = \frac{1}{1 - q} \frac{(q^{2v+2}, q^2)_\infty}{(q^2, q^2)_\infty}.$$

Define the q -Bessel translation operator as follows:

$$T_{q,x}^v f(y) = c_{q,v} \int_0^\infty \mathcal{F}_{q,v}(f)(t) j_v(xt, q^2) j_v(yt, q^2) t^{2v+1} d_q t, \quad \forall x, y \in \mathbb{R}_q^+, \forall f \in \mathcal{L}_{q,v,1}.$$

Recall that $T_{q,x}^v$ is said positive if $T_{q,x}^v f \geq 0$ for $f \geq 0$. In the following we tack $q \in Q_v$ where

$$Q_v = \{q \in]0, 1[, \quad T_{q,x}^v \text{ is positive for all } x \in \mathbb{R}_q^+\}.$$

The q -convolution product of both functions $f, g \in \mathcal{L}_{q,1,v}$ is defined by

$$f *_q g(x) = c_{q,v} \int_0^\infty T_{q,x}^v f(y) g(y) y^{2v+1} d_q y.$$

In the end we denote by $\mathcal{A}_{q,v}$ the q -Wiener algebra

$$\mathcal{A}_{q,v} = \{f \in \mathcal{L}_{q,1,v}, \quad \mathcal{F}_{q,v}(f) \in \mathcal{L}_{q,1,v}\}.$$

The followings results in this sections was proved in [1].

Proposition 1 *Let $n, m \in \mathbb{Z}$ and $n \neq m$, then we have*

$$c_{q,v}^2 \int_0^\infty j_v(q^n x, q^2) j_v(q^m x, q^2) x^{2v+1} d_q x = \frac{q^{-2n(v+1)}}{1-q} \delta_{nm}.$$

Proposition 2

$$|j_v(q^n, q^2)| \leq \frac{(-q^2; q^2)_\infty (-q^{2v+2}; q^2)_\infty}{(q^{2v+2}; q^2)_\infty} \begin{cases} 1 & \text{if } n \geq 0 \\ q^{n^2+(2v+1)n} & \text{if } n < 0 \end{cases}.$$

Proposition 3 *The q -Bessel Fourier transform*

$$\mathcal{F}_{q,v} : \mathcal{L}_{q,1,v} \rightarrow \mathcal{C}_{q,0},$$

satisfying

$$\|\mathcal{F}_{q,v}(f)\|_{\mathcal{C}_{q,0}} \leq B_{q,v} \|f\|_{q,1,v},$$

where

$$B_{q,v} = \frac{1}{1-q} \frac{(-q^2; q^2)_\infty (-q^{2v+2}; q^2)_\infty}{(q^2; q^2)_\infty}.$$

Theorem 1 *Given $f \in \mathcal{L}_{q,1,v}$ then we have*

$$\mathcal{F}_{q,v}^2(f)(x) = f(x), \quad \forall x \in \mathbb{R}_q^+.$$

If $f \in \mathcal{L}_{q,1,v}$ and $\mathcal{F}_{q,v}(f) \in \mathcal{L}_{q,1,v}$ then

$$\|\mathcal{F}_{q,v}(f)\|_{q,2,v} = \|f\|_{q,2,v}.$$

Proposition 4 Let $f \in \mathcal{L}_{q,1,v}$ then

$$T_{q,x}^v f(y) = \int_0^\infty f(z) D_v(x, y, z) z^{2v+1} d_q z,$$

where

$$D_v(x, y, z) = c_{q,v}^2 \int_0^\infty j_v(xt, q^2) j_v(yt, q^2) j_v(zt, q^2) t^{2v+1} d_q t.$$

Proposition 5 Given two functions $f, g \in \mathcal{L}_{q,v,1}$ then

$$f *_q g \in \mathcal{L}_{q,v,1},$$

and

$$\mathcal{F}_{q,v}(f *_q g) = \mathcal{F}_{q,v}(f) \times \mathcal{F}_{q,v}(g).$$

Proposition 6 The q -Gauss kernel

$$G^v(x, t, q^2) = \frac{(-q^{2v+2}t, -q^{-2v}/t; q^2)_\infty}{(-t, -q^2/t; q^2)_\infty} e\left(-\frac{q^{-2v}}{t}x^2, q^2\right),$$

satisfying

$$\mathcal{F}_{q,v} \{e(-ty^2, q^2)\} (x) = G^v(x, t, q^2),$$

and for all function $f \in \mathcal{C}_{q,b}$

$$\lim_{a \rightarrow 0} c_{q,v} \int_0^\infty f(x) G^v(x, a^2, q^2) x^{2v+1} d_q x = f(0).$$

Theorem 2 Given $1 < p, p', r \leq 2$ and

$$\frac{1}{p} + \frac{1}{p'} - 1 = \frac{1}{r}.$$

If $f \in \mathcal{L}_{q,p,v}$ and $g \in \mathcal{L}_{q,p',v}$ then

$$f *_q g \in \mathcal{L}_{q,r,v}.$$

2 Functions of q -positive type

Definition 1 A function ϕ is of q -positive type if

$$\phi \in \mathcal{C}_{q,b} \cap \mathcal{L}_{q,1,v}$$

and for any finite list of complex numbers z_1, \dots, z_n and q -real numbers x_1, \dots, x_n

$$\sum_{r=1}^n \sum_{l=1}^n z_r \bar{z}_l T_{q,x_r}^v \phi(x_l) \geq 0. \quad (1)$$

Proposition 7 Let $\phi \in \mathcal{A}_{q,v}$ of q -positive type then $\mathcal{F}_{q,v}\phi$ is of q -positive type.

Proof. From Proposition 3 and the definition of the q -Wiener algebra

$$\mathcal{F}_{q,v}(\phi) \in \mathcal{C}_{q,b} \cap \mathcal{L}_{q,1,v}.$$

On the other hand, with the inversion formula in Theorem 1 we get

$$T_{q,x}^v \mathcal{F}_{q,v}(\xi)(y) = \int_0^\infty j_v(tx, q^2) j_v(ty, q^2) t^{2v+1} \phi(t) d_q t,$$

then

$$\begin{aligned} \sum_{r=1}^n \sum_{l=1}^n z_r \bar{z}_l T_{q,x_r}^v \mathcal{F}_{q,v} \xi(x_l) &= c_{q,v} \int_0^\infty \left[\sum_{r=1}^n \sum_{l=1}^n z_r \bar{z}_l j_v(x_r t, q^2) j_v(x_l t, q^2) \right] t^{2v+1} \phi(t) d_q t \\ &= c_{q,v} \int_0^\infty \left[\sum_{r=1}^n z_r j_v(x_r t, q^2) \right] \overline{\left[\sum_{l=1}^n z_l j_v(x_l t, q^2) \right]} t^{2v+1} \phi(t) d_q t \\ &= c_{q,v} \int_0^\infty \left| \sum_{r=1}^n z_r j_v(x_r t, q^2) \right|^2 t^{2v+1} \phi(t) d_q t \geq 0. \end{aligned}$$

This finish the proof. ■

Proposition 8 If ϕ is of q -positive type and $f \in \mathcal{L}_{q,2,v}$ then

$$\phi *_q f \in \mathcal{L}_{q,2,v},$$

and

$$\langle \phi *_q f, f \rangle \geq 0.$$

Proof. From Theorem 2 we see that $\phi *_q f \in \mathcal{L}_{q,2,v}$. On the other hand

$$\begin{aligned} \langle \phi *_q f, f \rangle &= c_{q,v}^2 \int_0^\infty \left[\int_0^\infty T_{q,x}^v \phi(y) f(y) y^{2v+1} d_q y \right] f(x) x^{2v+1} d_q x \\ &= (1-q)^2 c_{q,v}^2 \sum_{r=1}^\infty \sum_{l=1}^\infty q^{(2v+2)r} f(q^r) q^{(2v+2)l} f(q^l) T_{q,q^r}^v \phi(q^l) \geq 0. \end{aligned}$$

This finish the proof. ■

Corollary 1 *If ϕ is of q -positive type then*

$$\mathcal{F}_{q,v} \phi(x) \geq 0, \quad \forall x \in \mathbb{R}_q^+.$$

Proof. Given $x \in \mathbb{R}_q^+$ and let

$$f_x : t \mapsto c_{q,v} j_v(xt, q^2),$$

then with Proposition 2 we see that $f_x \in \mathcal{L}_{q,2,v}$ and by Proposition 1

$$\mathcal{F}_{q,v} f_x(y) = \delta_{q,v}(x, y),$$

which implies (see[1])

$$\langle \mathcal{F}_{q,v} \phi \times \mathcal{F}_{q,v} f_x, \mathcal{F}_{q,v} f_x \rangle = \mathcal{F}_{q,v} \phi(x) \delta_{q,v}(x, x) = \frac{1}{(1-q)x^{2v+2}} \mathcal{F}_{q,v} \phi(x).$$

From Proposition 8

$$\langle \mathcal{F}_{q,v} \phi \times \mathcal{F}_{q,v} f, \mathcal{F}_{q,v} f \rangle = \langle \phi *_q f, f \rangle \geq 0,$$

this leads to the result. ■

Proposition 9 *If ϕ is of q -positive type then $\mathcal{F}_{q,v} \phi \in \mathcal{L}_{q,1,v}$.*

Proof. From Proposition 6

$$\lim_{a \rightarrow 0} \int_0^\infty e(-a^2 x, q^2) \mathcal{F}_{q,v} \phi(x) x^{2v+1} d_q x = \lim_{a \rightarrow 0} c_{q,v} \int_0^\infty G^v(x, a^2, q^2) \phi(x) x^{2v+1} d_q x = \phi(0).$$

By the monotone convergence theorem and the preview corollary we see that

$$\int_0^\infty |\mathcal{F}_{q,v} \phi(x)| x^{2v+1} d_q x = \int_0^\infty \mathcal{F}_{q,v} \phi(x) x^{2v+1} d_q x = \phi(0).$$

This finish the proof. ■

Corollary 2 *If ϕ is of q -positive type then there exist a positive function $\xi \in \mathcal{A}_{q,v}$ such that*

$$\phi(x) = \mathcal{F}_{q,v}\xi(x), \quad \forall x \in \mathbb{R}_q^+.$$

Proof. From the inversion formula in theorem 1

$$\phi(x) = \mathcal{F}_{q,v}^2\phi(x), \quad \forall x \in \mathbb{R}_q^+.$$

Define the function ξ as follows

$$\xi(x) = \mathcal{F}_{q,v}\phi(x).$$

By the use of Corollary 1 and Proposition 9 we see that ξ is a positive function of $\mathcal{A}_{q,v}$. ■

Proposition 10 *Suppose ϕ is of q -positive type. If $f \in \mathcal{L}_{q,1,v}$ is positive function then the product $\phi\mathcal{F}_{q,v}f$ is of q -positive type.*

Proof. Proposition 5 and Proposition 9 give

$$\mathcal{F}_{q,v}(\phi\mathcal{F}_{q,v}f)(t) = \mathcal{F}_{q,v}\phi *_q f(t), \quad \forall t \in \mathbb{R}_q^+,$$

then

$$\begin{aligned} & \sum_{r=1}^n \sum_{l=1}^n z_r \overline{z_l} T_{q,x_r}^v (\phi\mathcal{F}_{q,v}f)(x_l) \\ &= c_{q,v} \int_0^\infty \left[\sum_{r=1}^n \sum_{l=1}^n z_r j_v(x_r t, q^2) \overline{z_l j_v(x_l t, q^2)} \right] \mathcal{F}_{q,v}(\phi\mathcal{F}_{q,v}f)(t) t^{2v+1} d_q t \\ &= c_{q,v} \int_0^\infty \left[\sum_{r=1}^n \sum_{l=1}^n z_r j_v(x_r t, q^2) \overline{z_l j_v(x_l t, q^2)} \right] \mathcal{F}_{q,v}\phi *_q f(t) t^{2v+1} d_q t \\ &= c_{q,v} \int_0^\infty \left| \sum_{r=1}^n z_r j_v(x_r t, q^2) \right|^2 \mathcal{F}_{q,v}\phi *_q f(t) t^{2v+1} d_q t. \end{aligned}$$

From the definition of the q -convolution product we write

$$\mathcal{F}_{q,v}\phi *_q f(t) = c_{q,v} \int_0^\infty \mathcal{F}_{q,v}\phi(z) T_{q,t} f(z) z^{2v+1} d_q z.$$

Proposition 4 give

$$T_{q,t}f(z) = c_{q,v} \int_0^\infty D_v(t, z, s) f(s) s^{2v+1} d_q s \geq 0.$$

This implies with Corollary 1

$$\mathcal{F}_{q,v} \phi *_q f(t) \geq 0,$$

which leads to the result. ■

Corollary 3 *Given two functions ϕ_1, ϕ_2 which are of q -positive type then the product $\phi_1 \times \phi_2$ is also of q -positive type.*

Proof. Let $\xi = \mathcal{F}_{q,v} \phi_2$ then with the inversion formula in theorem 1 we see that $\mathcal{F}_{q,v} \xi = \phi_2$. Proposition 9 give

$$\xi \in \mathcal{L}_{q,1,v},$$

and by Proposition 10 we achieved the proof. ■

3 q -Bochner's Theorem

We consider the set \mathcal{M}_q^+ of positives and bonded measures on \mathbb{R}_q^+ . The q -Bessel Fourier transform of $\xi \in \mathcal{M}_q^+$ is defined by

$$\mathcal{F}_{q,v}(\xi)(x) = \int_0^\infty j_v(tx, q^2) t^{2v+1} d_q \xi(t).$$

The q -convolution product of two measures $\xi, \rho \in \mathcal{M}_q^+$ is given by

$$\xi *_q \rho(f) = \int_0^\infty T_{q,x}^v f(t) t^{2v+1} d_q \xi(x) d_q \rho(t),$$

and we have

$$\mathcal{F}_{q,v}(\xi *_q \rho) = \mathcal{F}_{q,v}(\xi) \mathcal{F}_{q,v}(\rho).$$

The following Theorem (see[1]) is crucial for the proof of our main result.

Theorem 3 Let $(\xi_n)_{n \geq 0}$ be a sequences of probability measures of \mathcal{M}_q^+ such that

$$\lim_{n \rightarrow \infty} \mathcal{F}_{q,v}(\xi_n)(x) = \psi(x),$$

then there exists $\xi \in \mathcal{M}_q^+$ such that the sequence ξ_n converge strongly toward ξ and

$$\mathcal{F}_{q,v}(\xi) = \psi.$$

In the following we consider the function ψ defined by

$$\psi(x) = \begin{cases} 1 - x & \text{if } x < 1 \\ 0 & \text{otherwise} \end{cases}.$$

Now we are in a position to state and prove the q-analogue of the Bochner's theorem

Theorem 4 Let ϕ be a function defined on \mathbb{R}_q^+ continued at 0. Assume that the following function

$$\phi_n : x \mapsto \phi(x)\psi(q^n x),$$

satisfy (1) for all $n \in \mathbb{N}$ then there exist $\xi \in \mathcal{M}_q^+$ such that

$$\mathcal{F}_{q,v}(\xi) = \phi.$$

Proof. The function ϕ_n is of q -positive type. From Corollary 2 there exist ϱ_n a positive function of $\mathcal{A}_{q,v}$ such that

$$\mathcal{F}_{q,v}(\varrho_n) = \phi_n.$$

The measure ξ_n defined by

$$d_q \xi_n(x) = \varrho_n(x) d_q x,$$

belong to \mathcal{M}_q^+ and

$$\int_0^\infty x^{2v+1} d_q \xi_n(x) = \mathcal{F}_{q,v}(\varrho_n)(0) = \phi_n(0) = \phi(0).$$

Assume that $\phi(0) = 1$. On the other hand

$$\lim_{n \rightarrow \infty} \mathcal{F}_{q,v}(\xi_n)(x) = \lim_{n \rightarrow \infty} \phi_n(x) = \phi(x).$$

From Theorem 3 there exists $\xi \in \mathcal{M}_q^+$ such that the sequence ξ_n converge strongly toward ξ , and

$$\mathcal{F}_{q,v}(\xi) = \phi,$$

which leads to the result. ■

References

- [1] L. Dhaouadi, A. Fitouhi and J. El Kamel, Inequalities in q -Fourier Analysis, *Journal of Inequalities in Pure and Applied Mathematics*, Volume 7, Issue 5, Article 171, 2006.
- [2] G. Gasper and M. Rahman, Basic hypergeometric series, *Encyclopedia of mathematics and its applications* 35, Cambridge university press, 1990.
- [3] F. H. Jackson, On a q -Definite Integrals, *Quarterly Journal of Pure and Application Mathematics* 41, 1910, 193-203.
- [4] T. H. Koornwinder and R. F. Swarttouw, On q -Analogues of the Hankel and Fourier Transform, *Trans. A. M. S.* 1992, 333, 445-461.
- [5] R. F. Swarttouw, The Hahn-Exton q -Bessel functions PhD Thesis The Technical University of Delft (1992).