



HAL
open science

**Late Famennian correlation by miospores between the
Refrath 1 borehole (Bergisch Gladbach-Paffrath
Syncline, Germany) and the reference section of
Chanxhe (Dinant Syncline, Belgium)**

Maurice Strel, Christoph Hartkopf-Fröder

► **To cite this version:**

Maurice Strel, Christoph Hartkopf-Fröder. Late Famennian correlation by miospores between the Refrath 1 borehole (Bergisch Gladbach-Paffrath Syncline, Germany) and the reference section of Chanxhe (Dinant Syncline, Belgium). *Carnets de Géologie / Notebooks on Geology*, 2005, CG2005 (M02/10), pp.57-60. hal-00167981

HAL Id: hal-00167981

<https://hal.science/hal-00167981>

Submitted on 24 Aug 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Late Famennian correlation by miospores between the Refrath 1 borehole (Bergisch Gladbach-Paffrath Syncline, Germany) and the reference section of Chanxhe (Dinant Syncline, Belgium).

[Corrélations par miospores au Famennien supérieur entre le sondage de Refrath 1 (Synclinal Bergisch Gladbach-Paffrath, Allemagne) et la section de référence de Chanxhe (Synclinorium de Dinant, Belgique)]

Maurice STREEL¹

Christoph HARTKOPF-FRÖDER²

Key Words: Famennian; miospores; Germany; Belgium; biostratigraphy

STREEL M. & HARTKOPF-FRÖDER C. (2005).- Late Famennian correlation by miospores between the Refrath 1 borehole (Bergisch Gladbach-Paffrath Syncline, Germany) and the reference section of Chanxhe (Dinant Syncline, Belgium). *In*: STEEMANS P. & JAVAUX E. (eds.), Pre-Cambrian to Palaeozoic Palaeopalynology and Palaeobotany.- [Carnets de Géologie / Notebooks on Geology](#), Brest, Memoir 2005/02, Abstract 10 (CG2005_M02/10)

Mots-Clefs : Famennien ; miospores ; Allemagne ; Belgique ; biostratigraphie

The Bergisch Gladbach-Paffrath Syncline and the Dinant Syncline are both part of the Ardennes-Rhenish Massif but are separated by some 100 km (Fig. 1).

In the Upper Devonian succession of the Bergisch Gladbach-Paffrath Syncline (The Bergisches Land), the Knoppenbissen Formation is assigned the "Dasbergian" (HARTKOPF-FRÖDER *et alii*, 2004). In this syncline the thermal alteration index (TAI of 2 to 2+) is remarkably low compared to that of other Upper Devonian sediments in the Ardenne-Rhenisch area. Due to this low thermal maturity the preservation of miospores is very good to excellent (HARTKOPF-

FRÖDER & STREEL, 1994; HARTKOPF-FRÖDER, 2004). The sedimentary rocks of the Refrath 1 borehole are mainly dark grey, partly calcareous mudstones; rare marlstone intercalations are considered to be fine-grained turbidites. Both contain many fossils. Aside from the very well preserved miospores of the *Retispora lepidophyta - Knoxisporites literatus* (LL) miospore Zone, numerous other microfossils are known *i.e.* conodonts (PIECHA, 2004) and entomozocean ostracodes (GROOS-UFFENORDE, 2004). Some characteristic ammonoid species are present in the lowermost part of the borehole (KORN, 2004).



Figure 1: The Ardennes-Rhenish Massif and the location of the reference section and borehole, respectively in a neritic (Chanxhe) and a pelagic (Refrath 1) facies.

¹ Département de Géologie, Université de Liège, Allée du 6 août, B18, Sart-Tilman, 4000 Liège (Belgium)

² Geologisches Landesamt Nordrhein-Westfalen, De-Greif-Str. 195, Postfach 1080, D-47710 Krefeld (Germany)

In the Dinant Syncline, the Comblain-au-Pont Formation is assigned the "Strunian". Thermal alteration is rather high; therefore miospores in the formation are dark brown. Farther east, the Comblain-au-Pont Formation, the time-equivalent in Belgium of the French Etroeungt Formation, is made up of some dm- to several m-thick beds of green shale and siltstone interbedded with micaceous, argillaceous or calcareous sandstones. From bottom to top three lithologic units may be distinguished: a shale dominant unit, a sandstone dominant unit and a carbonate dominant unit (CONIL, 1964). The latter is the equivalent of the Etroeungt Limestone s.s. (BULTYNCK & DEJONGHE, 2001). The Chanxhe road section (Ourthe valley, eastern Belgium) displays the three lithologic units of the Comblain-au-Pont Formation. It is a site where miospores, foraminifers and conodonts have been described and occur in a sequence permitting intercorrelation of faunal components. Above the base of the *Retispora lepidophyta*-*Knoxisporites literatus* (LL)

miospore Zone, between the Df3δ and Df3ε foraminifer faunas, a late *expansa* conodont fauna is present. And above the Df3ε foraminifer fauna the succeeding *Retispora lepidophyta*-*Indotriradites explanatus* (LE) miospore Zone occurs. Benthic ostracodes belong to the "Eifelian Mega-Assemblage" (CASIER *et alii*, 2005) and allow ecological interpretation. In the Chanxhe section a decrease upward in the diameter of *Retispora lepidophyta* has been examined in great detail by MAZIANE *et alii* (2002). They reports a significant shift of mean size (more than or less than 60 μm) near the base of the carbonate dominant unit (around bed 115 on Fig. 1). This shift allows the recognition of two varieties of *Retispora lepidophyta*, i.e. *R. l. var. lepidophyta* dominant below and *R. l. var. minor* dominant above. Comparison with the range of sizes in other miospores supports the conclusion that this decrease is not the result of a process of sorting in the carbonate facies but rather a consequence of some evolutionary process.




Figure 2: Late and latest Famennian biostratigraphic correlation between the Comblain-au-Pont Formation at Chanxhe and the Knoppenbissen Formation in the Refrath borehole.




Figure 3: Location of the Refrath 1 borehole in the basinal facies of the Bergisch Gladbach-Paffrath Syncline.

The Chanxhe road section is a candidate reference section, in a neritic facies, for the transition between late Famennian and latest Famennian. Although an international agreement has yet not been reached to define lower limits of substages within the Famennian Stage, a subdivision into 4 substages has been adopted internationally. A proposal for a conodont-based definition of the lower limit of the uppermost Famennian at the base of the late *expansa* Zone (STREEL, 2005) is still under consideration by the International Subcommittee on Devonian Stratigraphy and, in this context, a search for a good reference section in a neritic facies is more important than ever.

The base of the late *expansa* Zone is absent in the Chanxhe section but new data from the Refrath 1 borehole allow this gap in our knowledge to be filled. In the Refrath 1 borehole, the lower part of the LL miospore Zone (which includes the presumed ancestral *Retispora macroreticulata* and large specimens of *R. lepidophyta*) is the equivalent of the medial portion of the *expansa* conodont Zone and the early part of the *hemisphaerica - dichotoma* entomozocean Zone. The conclusion is that the base of the late *expansa* conodont Zone and the base of the *Q. kob. kobeitusana* (Df3ε) foraminifer Zone are obviously much closer together than was believed previously (Fig. 2).

The few characteristic ammonoid species in the lowermost part of Refrath 1 suggest an age slightly older than the age indicated by the 3 groups of abundant microfossils (GROOS-UFFENORDE, 2004; HARTKOPF-FRÖDER, 2004; PIECHA 2004). This can be explained only if we accept that during their transport down the slope (Fig. 3) the turbiditic flows removed by erosion the upper levels of some limestone shoals or else that at present only a portion of the ammonoid fauna of the Knoppenbissen Formation is known.

The facies relationships shown here indicate that the turbiditic flows on the slope started when the transgression on the shelf (MAZIANE *et alii*, 2002, p. 222) had reached its maximum and was succeeded by cyclic regressive phases.

Bibliographic references

- BULTYNCK P. & DEJONGHE L. (2001).– Devonian lithostratigraphic units (Belgium).– *Geologica Belgica*, Liège, vol. 4/1-2, p. 39-69.
- CASIER J-G, LEBON A., MAMET B. & PRÉAT A. (2005).– Ostracods and lithofacies close to the Devonian-Carboniferous boundary in the Chanxhe and Rivage sections, northeastern part of the Dinant Basin, Belgium.– *Bulletin de l'Institut Royal des Sciences naturelles de Belgique*, Bruxelles, (Sciences de la Terre), vol. 75, p. 95-126.

- CONIL R. (1964).- Localités et coupes types pour l'étude du Tournaisien inférieur (Révision des limites sous l'aspect micropaléontologique).- *Mémoires de l'Académie Royale de Belgique*, Bruxelles, Classe des Sciences, n° 15, 4, 87 p.
- GROOS-UFFENORDE H. (2004).- Late Famennian ostracodes from the Refrath 1 borehole (Bergisch Gladbach-Paffrath Syncline; Ardennes-Rhenish Massif, Germany).- *Courier Forschungsinstitut Senckenberg*, Frankfurt am Main, 251, p. 185-235.
- HARTKOPF-FRÖDER C. & STREEL M. (1994).- Late Famennian miospore assemblages from the Bergish Gladbach- Paffrath Syncline, Rhenish Slate Mountains, Germany.- *Annales de la Société Géologique de Belgique*, Liège, 116, p. 333-357.
- HARTKOPF-FRÖDER C. (2004).- Palynostratigraphy of upper Famennian sediments from the Refrath 1 borehole (Bergisch Gladbach-Paffrath Syncline; Ardennes-Rhenish Massif, Germany).- *Courier Forschungsinstitut Senckenberg*, Frankfurt am Main, 251, p. 77-87.
- HARTKOPF-FRÖDER C., JUX U., KNAPP G. & PIECHA M. (2004).- The Late Devonian of the Bergisch Gladbach-Paffrath Syncline (Ardennes-Rhenish Massif, Germany): an overview.- *Courier Forschungsinstitut Senckenberg*, Frankfurt am Main, 251, p. 7-18.
- KORN D. (2004).- Late Famennian cephalopods from the Refrath 1 borehole (Bergisch Gladbach-Paffrath Syncline; Ardennes-Rhenish Massif, Germany).- *Courier Forschungsinstitut Senckenberg*, Frankfurt am Main, p. 175-183.
- MAZIANE N., HIGGS K.T. & STREEL M. (2002).- Biometry and paleoenvironment of *Retispora lepidophyta* (KEDO) PLAYFORD 1976 and associated miospores in the latest Famennian nearshore marine facies, eastern Ardenne (Belgium).- *Review of Palaeobotany and Palynology*, Amsterdam, vol. 118, n° 1-4, p. 211-226.
- PIECHA M. (2004).- Late Famennian conodonts from the Refrath 1 borehole (Bergisch Gladbach-Paffrath Syncline; Ardennes-Rhenish Massif, Germany).- *Courier Forschungsinstitut Senckenberg*, Frankfurt am Main, 251, p. 253-265.
- STREEL M. (2005).- Subdivision of the Famennian Stage into four Substages and correlation with the neritic and continental miospore zonation (SDS Business Meeting Florence, August 23, 2004).- SDS Newsletter 21 (online version): p. 14 & 16-17.