



**HAL**  
open science

## A remark on utility streams

Karl Schlechta

► **To cite this version:**

| Karl Schlechta. A remark on utility streams. 2007. hal-00169177

**HAL Id: hal-00169177**

**<https://hal.science/hal-00169177>**

Preprint submitted on 1 Sep 2007

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A REMARK ON UTILITY STREAMS

Karl Schlechta \*

Laboratoire d'Informatique Fondamentale de Marseille †

September 1, 2007

We write down in this very short comment some ideas which occurred to the author during an email discussion with Kaushik Basu on the paper [BM05].

The author suggested to consider only finite sequences (which solves the embedding problem, as the resulting set is countable), and to compare sequences of unequal length by repeating them until they have the same length, e.g. a sequence of length 2 will be repeated 3 times, and a sequence of length 3 2 times, and the results will then be compared.

Note that the author discussed somewhat related problems in Section 2.2.7 of [Sch04]. Considering sums, and not only orders, to evaluate sequences is generally difficult, in the sense that often no finite characterizations are possible - see again [Sch04].

We will write down now a few axioms, which seem reasonable, without discussion.

We have a domain  $X$ , and consider finite, non-empty sequences, noted  $\sigma$  etc., with values in  $X$ , the set of these sequences will be denoted  $\Sigma$ .  $X$  has an order  $<$ ,  $\equiv$  will express equivalence wrt. this order, and we put restrictions on a resulting order  $\prec$  on  $\Sigma$ , with equivalence  $\approx$ .  $\leq$  and  $\preceq$  etc. are defined in the obvious way.

## Notation 1.1

For  $\sigma$  and  $\sigma'$  of equal length, we write

$\sigma \leq \sigma'$  iff all  $\sigma_i \leq \sigma'_i$ ,

$\sigma < \sigma'$  iff  $\sigma \leq \sigma'$  and for one  $i$   $\sigma_i < \sigma'_i$ , and finally

$\sigma \ll \sigma'$  iff all  $\sigma_i < \sigma'_i$ .

The double use of  $\leq$  and  $<$  will not pose any problem.

---

\*ks@cmi.univ-mrs.fr, karl.schlechta@web.de, <http://www.cmi.univ-mrs.fr/~ks>

†UMR 6166, CNRS and Université de Provence, Address: CMI, 39, rue Joliot-Curie, F-13453 Marseille Cedex 13, France

$\{x\}$  is the sequence of length 1.

Concatenation is noted  $\circ$ . For singletons, we may use simple juxtaposition.

$\sigma^n$  is  $\sigma$  repeated  $n$  times.

### Axiom 1.1

(1) Singletons:

$$(1.1) x < x' \rightarrow \{x\} \prec \{x'\},$$

$$(1.2) x \equiv x' \rightarrow \{x\} \approx \{x'\}$$

(essentially Pareto).

(2) concatenation:

$$(2.1) \sigma \circ \sigma \approx \sigma \text{ (this expresses essentially that the mean value is interesting),}$$

$$(2.2) \sigma' \prec \sigma'' \rightarrow \sigma \circ \sigma' \prec \sigma \circ \sigma'',$$

$$(2.3) \sigma' \approx \sigma'' \rightarrow \sigma \circ \sigma' \approx \sigma \circ \sigma''.$$

(3) permutation:

$$\sigma \circ \sigma' \approx \sigma' \circ \sigma$$

(essentially Anonymity).

### Fact 1.1

These axioms allow to deduce:

$$(4) \sigma \approx \sigma' \rightarrow \sigma \approx \sigma \circ \sigma'$$

$$(5) \sigma \prec \sigma' \rightarrow \sigma \prec \sigma \circ \sigma'$$

(6) if there are  $i, j \leq \text{length}(\sigma) = \text{length}(\sigma')$ ,  $\sigma_i = \sigma'_j$ ,  $\sigma_j = \sigma'_i$ , and  $\sigma_k = \sigma'_k$  for all other  $k$ , then  $\sigma \approx \sigma'$  (real Anonymity)

(7) Weak Pareto:

$$(7.1) \sigma \leq \sigma' \rightarrow \sigma \preceq \sigma',$$

$$(7.2) \sigma < \sigma' \rightarrow \sigma \prec \sigma',$$

$$(7.3) \sigma \ll \sigma' \rightarrow \sigma \prec \sigma'.$$

and

(8) to compare sequences of different lengths, in the following sense: When  $\prec$  and  $\approx$  are defined between  $\sigma$ 's of equal length, and Axioms 1-3 hold, then the relation  $\prec$  (and  $\approx$ ) is determined for arbitrary sequences.

**Proof:**

Elementary.

(4) by (2.1) and (2.3).

(5) by (2.1) and (2.2).

(6) Let e.g.  $\sigma$  be  $\sigma_0 \circ a \circ \sigma_1 \circ b \circ \sigma_3$ , then  $ab \approx ba$  by (3), so  $\sigma_0 \circ ab \approx \sigma_0 \circ ba$  by (2.3) and (3), so  $b \circ \sigma_0 \circ a \approx a \circ \sigma_0 \circ b$  by (3), so  $b \circ \sigma_0 \circ a \circ \sigma_1 \approx a \circ \sigma_0 \circ b \circ \sigma_1$  by (2.3), etc.

(7) This follows from (1) and repeated use of (2.2) and (2.3).

(8) Let  $m := \text{length}(\sigma)$ ,  $n := \text{length}(\sigma')$ , then we obtain by using (2.1) once, and (2.3) repeatedly, that  $\sigma^n \approx \sigma$  and  $\sigma'^m \approx \sigma'$ , but  $\sigma^n$  and  $\sigma'^m$  have the same length.

□

## References

- [BM05] K.Basu, T.Mitra: "Possibility theorems for aggregating infinite utility streams equitably", in J.Roemer, K.Suzumura: "Intergenerational equity and sustainability", Palgrave, forthcoming
- [Sch04] K.Schlechta: "Coherent Systems", Elsevier, Amsterdam, 2004