



HAL
open science

FACTORS CONTRIBUTING TO YOLK RETENTION IN POULTRY: A REVIEW

Kashif Aziz Khan, Shakil Akhtar Khan, Asim Aslam, Masood Rabbani,
Muhammad Yasin Tipu

► **To cite this version:**

Kashif Aziz Khan, Shakil Akhtar Khan, Asim Aslam, Masood Rabbani, Muhammad Yasin Tipu. **FACTORS CONTRIBUTING TO YOLK RETENTION IN POULTRY: A REVIEW**. Pakistan Veterinary Journal, 2004, 24 (1), pp.46-51. hal-00179202v2

HAL Id: hal-00179202

<https://hal.science/hal-00179202v2>

Submitted on 29 Jul 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FACTORS CONTRIBUTING TO YOLK RETENTION IN POULTRY: A REVIEW

K. A. Khan, S. A. Khan, A. Aslam, M. Rabbani¹ and M.Y. Tipu
*Department of Pathology and ¹Department of Microbiology,
University of Veterinary and Animal Sciences, Lahore, Pakistan*

ABSTRACT

Yolk retention and yolk sac infection is considered as an important cause of death in chicken as well as in guinea fowl, duck, turkey, quail and goose. The factors which slow down the rate of yolk absorption and may in turn, lead to yolk retention are discussed. Yolk sac infection of bacterial origin is most important among these factors. Other factors which may contribute include posthatch starvation, type of initial feed, brooding temperature, prolonged exposure to hatcher environment and size of birds.

Key words: Yolk retention, poultry, yolk sac infection.

INTRODUCTION

During incubation, extraembryonic membranes encircle the yolk substance and constitute the yolk sac, which is attached to gut of the chick by yolk stalk. Just before hatching, the yolk sac is pulled from the egg cavity to abdomen of chick as an extension of intestine. Residual yolk comprises 20-25% of body weight at hatch but within the first week of life it becomes negligible in size (Ramnoff, 1960).

There are certain factors that affect yolk absorption and in turn may lead to its retention. Once bacteria get entry to yolk, other factors favour rapid bacterial growth and these include the fact that the yolk contains a lot of fat and water, favoured nutrients for bacteria. In addition, the yolk sac is maintained at the temperature of the hatcher and then at the chick's body temperature, which are the ideal temperatures for multiplication of certain bacteria (Anonymous, 2000). So yolk retention due to any cause may lead to yolk sac infection.

YOLK RETENTION

a) Incidence

Incidence of yolk retention and yolk sac infection is widely reported in literature. Anjum (1997) stated that it was the commonest cause of early chick mortality in Pakistan. Jordan (1990) and Singh *et al.* (1993) also reported it as the most frequent cause of death in chicks. Incidence of yolk retention and yolk sac infection was reported as 7.5, 10.5, 9.9, 5.1 and 8.9% in chickens by Schonhofen and Garcia (1981), Shrivastava (1982), Rathore *et al.* (1985), Suresh *et al.* (1988) and Bhattacharjee *et al.* (1996), respectively. Incidence was reported to be 15.20 and 20.71% in two

different strains of White Leg Horn by Viswanath *et al.* (1985). Ghodasara *et al.* (1992) found 31.45% mortality in chicks due to yolk sac infection.

Yolk retention is not only the cause of death in chicken but also in other species of poultry including guinea fowl, duck, turkey, quail and goose. It was reported as most frequent cause of death in indigenous guinea fowl by Rudy (1991). Sharma and Kaushik (1986b) and Roy and Misra (1989) found it as an important disease of ducks. Sharma and Kaushik (1986a) found that incidence of yolk retention was 20% in turkey while Thyagarajan *et al.* (1987) found it as 20.8%. Unabsorbed yolk was observed as principal lesion in quails died upto one week of age by Suneja *et al.* (1983) and incidence was reported as 16% by Sharma and Kaushik (1986c). Boado and Rojas (1990) found 7.3% incidence of omphalitis in goose.

Suneja *et al.* (1983) observed that incidence of omphalitis was most frequent in January. Similarly, Sainsbury (1992) reported it's higher incidence towards the end of winter or in the early spring.

b) Effects on host

Putrefactive and offensive odour was observed as characteristic clinical sign of yolk sac infection by Sainsbury (1992) and Anjum (1997). Abdomen of chick felt soft and distended with thickened, inflamed and moist umbilicus. Unabsorbed yolk sac was present in the abdomen and therefore it was named yolk retention. Yolk sac contents changed from viscid yellow green to watery yellow brown due to denaturation of yolk by bacteria (Jordan, 1990; Sainsbury, 1992; Anjum, 1997). Deeming (1995) reported that infected yolk sacs were, in general, larger in mass than uninfected sacs from poult of same age. Yolk sac and subcutaneous blood

vessels were dilated and engorged with blood (Jordan, 1990; Anjum, 1997). Chicks surviving more than four days might have pericarditis as well as infected yolk indicating systemic spread (Barnes and Gross, 1997). Haemorrhagic serous peritonitis was also observed by Jordan (1990) and Anjum (1997). Maximum deaths occurred upto 3 days of age (Sainsbury, 1992; Anjum 1997). In some cases, there might be no mortality with retained infected yolk as only manifestation (Barnes and Gross, 1997).

FACTORS CONTRIBUTING TOWARDS YOLK RETENTION

A. YOLK SAC INFECTION

A 1: Natural infection

a) Causative organisms

Different types of bacterial agents are attributed for causation of yolk sac infection/omphalitis. *Escherichia coli* was frequently the main one involved. Its isolation from yolk sac infection was reported by Zahdeh *et al.* (1984), Sarma *et al.* (1985), Linzitto *et al.* (1988), Jordan (1990), Utomo *et al.* (1990), Sainsbury (1992), Ali (1993), Choudhury *et al.* (1993), Deeming (1995), Rehman *et al.* (1996), Anjum (1997), Sharada *et al.* (1999) and Anonymous (2000). Next frequently found bacteria was genus *Salmonella* which was reported by Zahdeh *et al.* (1984), O'Brien (1988), Mutalib and Hanson (1989), Ali (1993), Choudhury *et al.* (1993), Rehman *et al.* (1996), Anjum (1997), Anonymous (2000) and Shivaprasad (2000).

Staphylococcus species were found to be involved by Zahdeh *et al.* (1984), Utomo *et al.* (1990), Choudhury *et al.* (1993), Deeming (1995), Rehman *et al.* (1996), Anjum (1997) and Anonymous (2000) and streptococcus species by Utomo *et al.* (1990), Ali (1993) and Anonymous (2000). Involvement of proteus species was observed by Zahdeh *et al.* (1984), Sarma *et al.* (1985), Utomo *et al.* (1990), Choudhury *et al.* (1993), Anjum (1997) and Anonymous (2000) and of bacillus species by Utomo *et al.* (1990), Ali (1993), Choudhury *et al.* (1993), Deeming (1995), Anjum (1997) and Anonymous (2000).

Other bacterial genera found to be involved include *Pseudomonas* (Zahdeh *et al.*, 1984; Sarma *et al.* 1985; Utomo *et al.*, 1990; Choudhury *et al.*, 1993; Anjum, 1997; Anonymous, 2000), *Klebsiella* (Zahdeh *et al.*, 1984; Choudhury *et al.*, 1993; Anonymous, 2000), *Clostridium* (Anjum, 1997; Anonymous, 2000), *Micrococcus* (Utomo *et al.*, 1990), *Yersinia* (Ali, 1993), *Enterobacter* (Utomo *et al.*, 1990; Ali, 1993)

Aerobacter (Bhatia *et al.*, 1971), *Citrobacter* (Utomo *et al.*, 1990), *Achromobacter* (Deeming, 1995), *Enterococci* (Zahdeh *et al.*, 1984; Anjum, 1997) and *Alcaligenes* (Utomo *et al.*, 1990). Involvement of *Aspergillus fumigatus* in yolk sac infection was also reported by Schonhofen and Garcia (1991).

b) Source of infection

Farm faecal contamination of shell was reported as source of infection by Anjum (1997). Poor hatchery hygiene condition was considered as another important source (Sainsbury, 1992; Anjum, 1997). William (1975) reported that the source of infection was the waste in the hatchery or contaminated poult boxes or poult box pads. Other source of infection include breeder, feed, environment, feathers, human skin, floor and dirty equipment (Anonymous, 2000).

c) Route of infection

Transmission of bacteria through unhealed naval was revealed by Jordan (1990) and Anjum (1997). Infection through blood stream and contamination of yolk before it is internalized into the chick were reported as other routes of infection by Anonymous (2000).

A 2: Experimental infection

Fuller and Jayne-Williams (1968) demonstrated sub-clinical yolk sac infection after oral administration of pure cultures of bacteria. It was concluded that the infection arose through translocation of bacteria across the gut wall. Seigo *et al.* (1970) inoculated *Bacillus cereus* through intrayolk, intraperitoneal, subcutaneous and oral routes. Omphalitis was reproduced only when inoculated into the yolk sac. Singh *et al.* (1997) studied the pathogenicity of *Escherichia coli* by intraperitoneal injection into 2-day-old chicks. Unabsorbed yolk sac was among the main lesions. Sander *et al.* (1998) observed retained yolk in chicks which received *Enterococcus faecalis* broth inoculation into yolk sac. Omphalitis was also observed as gross lesion in experimental infection with *Salmonella harder* and *Salmonella enteritidis* by Desmidt *et al.* (1998) and Dhillon *et al.* (2001), respectively. Khan *et al.* (2002) inoculated *E. coli* broth into yolk sac of day-old chicks and observed high yolk sac weight in these chicks as compared with the chicks inoculated with sterile broth. Seigo *et al.*, (1970) produced the disease by inoculating *Bacillus cereus* inside the egg shell of pipped eggs. Zahdeh (1987) observed severe oedematous swelling around navel orifice, severe omphalitis and incomplete withdrawal of yolk sac in chicks hatched from embryonated eggs that were dipped in 24-hours bacterial broth culture on 18th day.

B. POSTHATCH STARVATION

Slow absorption of yolk due to fasting has been reported by many workers. Moran and Reinhart (1980) observed that fasting led to a reduced uptake of yolk as compared to fully nourished birds. It was reported further that fasting favoured removal of moisture and lipid to a greater extent than protein while the converse was true if access to feed and water was permitted. Observations of Noy *et al.* (1996) also showed that yolk utilization was more rapid in fed than in fasted chicks, suggesting that the transport of yolk through the intestine could be increased by the greater intestinal activity found in fed chicks. Similar findings were observed by Santos and Silversides (1996) that starving chicks were unable to use the yolk sac nutrients, suggesting that yolk sac utilization seems to be correlated with activation of the digestive system.

In contrast, Murakami *et al.* (1992) found that posthatch starvation in chicks decreased carcass lipid content but did not modify the disappearance rate of yolk in the abdomen. Similarly, Chamblee *et al.* (1992) observed that availability of feed and water did not affect body weight or yolk sac absorption during first 24 hours. Al-Rawashdeh *et al.* (1995) observed non-significant difference in yolk weight of chicks after subjecting one-day-old chicks to five days starvation, with drinking water available freely. Baiao *et al.* (1998) delayed the housing of chicks for 24, 48 and 72 hours after hatching and observed that yolk sac absorption was not affected by the period of feed withdrawal between hatching and housing.

Efficient absorption of yolk due to fasting was reported by Pisarsaki *et al.* (1998b). The author delayed first feeding and watering of chicks for 24, 48 and 72 hours in two subsequent experiments and observed that yolk absorption/body weight ratio was higher in chicks fed 24-72 hours later than the control ones.

C. TYPE OF INITIAL FEED

Pisarsaki *et al.* (1998a) substituted commercial starter ration with ground corn in the first 12, 24 and 36 hours of life and concluded that corn feeding in early life led to slow absorption of yolk sac contents. The levels of lysine and methionine also have some effect on absorption of yolk sac in chicks (Wang *et al.*, 1994).

D. BROODING TEMPERATURE

Leeson *et al.* (1978) reported that incidence of unabsorbed yolk was increased by the fluctuating environmental temperature, with little difference for the cold and hot environments, compared with a control situation. Thaxton *et al.* (1974) found that yolk sac weight was not changed by lowered brooding temperature, while Yousaf (1985) observed that yolk

sac weight was decreased by cold temperature brooding but not until 72 hours.

E. MISCELLANEOUS FACTORS

Martin (1996) studied the effect of prolonged exposure to hatcher environment on yolk sac size and observed enlarged yolk sac in broiler chicks. In contrast, Chamblee *et al.* (1992) reported that chicks kept in hatcher for 24 hours and chicks kept in the hatcher for 12 hours and then on litter for 12 hours exhibited non-significant differences in body weight and yolk weight. Knizetova *et al.* (1989) observed rapid resorption of yolk sac contents in large duckling and goslings than in small ones.

CONCLUSION

It can be inferred from the above given findings that yolk retention and yolk sac infection is widely recorded in different species of birds and assumed as an important factor for early chick mortality. Some workers have related its incidence with winter.

The cases of yolk retention and yolk sac infection were recorded upto 10 days of age but high rate of mortality was observed upto 3 days of age. Putrefactive odour from the birds can be assumed as first signal for the farmers. The flabbiness and distention of abdomen, moist umbilicus and change in size, consistency and appearance of yolk can be thought as indicative of yolk sac infection. The affected birds also exhibit systemic manifestations such as pericarditis and peritonitis.

Diverse species of bacteria and *Aspergillus fumigatus* cause natural cases of yolk sac infection. Out of more than a dozen genera of bacteria, mainly the members of family Enterobacteriaceae are predominant. It is difficult to establish, which species of bacteria acted as primary factor and which one acted as secondary opportunist. On the basis of information relating to bacterial origin of yolk sac infection, it is tempting to speculate that *Escherichia coli* and *Salmonella* spp, recorded in natural cases, are the primary pathogens. The main route of infection is through unhealed navel but in some instances transmission is possible by blood stream and by contamination of yolk before it is inverted into the chick. Main sources of infection are faecal contamination of hatching eggs, contaminated hatchery equipments, poor hatcher environment and unhygienic poult boxes.

Experimental yolk sac infection can be produced in chicks by different routes i.e. intrayolk, intraperitoneal,

navel swabbing, oral and subcutaneous, but intrayolk and intraperitoneal are the best. Infection can also be produced by inoculation inside the egg shell of piped eggs or by exposing embryonating eggs to bacterial growth in broth. Different types of bacteria were used by different workers.

Reports regarding posthatch starvation are controversial. According to many workers yolk absorption is delayed due to posthatch starvation while other reported that it did not affect yolk absorption. Even efficient absorption of yolk due to fasting is also reported.

REFERENCES

- Ali, M.I., 1993. Studies on isolation and antibiotic sensitivity of bacteria causally associated with omphalitis in broiler chicks. M.Sc. (Hons.) Thesis, Dept. Vet. Microbiol. Univ. Agri. Faisalabad.
- Al-Rawashdeh, O.F., S.Q. Lafi, N.Q. Hailat, T.A. Abdul-Aziz, K.T. Freifij and A.Y.M. Nour, 1995. Effect of feed deprivation on the blood levels of flucose, beta-hydroxybutyrate, triglycerides and cholesterol and on body weight and yolk sac weight in one-day-old broiler chicks. *Acta Vet. Reograd*, 45(4): 175-186.
- Anjum, A.D., 1997. Poultry Diseases, VetAg Publications, Faisalabad-Pakistan, pp: 178-180.
- Anonymous, 2000. Controlling yolk sac infection. *Int. Hatchery Pract.*, 14(6): 27-28.
- Baiao, N.C., S.V. Cancado and C.G. Lucio, 1998. Effect of period of feed withdrawal between hatching and housing of the broilers on absorption of the yolk sac. *Arquivo Brasileiro de Medicina Veterinaria e Zootecnia*, 50 (4): 473-474.
- Barnes, H.J. and W.B. Gross, 1997. Colibacillosis. In: *Diseases of Poultry*, 10th Ed., Iowa State Univ. Press, Ames, Iowa, U.S.A. pp: 131-142.
- Bhatia, K.C., U.K. Sharma and N. Singh, 1971. Studies on persistent yolk sac condition in chicks. Incidence and pathology. *Indian J. Anim. Hlth.*, 10(1): 95-98.
- Bhattacharjee, P.S., R.L. Kundu, R.K. Biswas, J.U. Mazumder, E. Hossain and A.H. Miah, 1996. A retrospective analysis of chicken diseases diagnosed at the Central Disease Investigation Laboratory, Dhaka. *Bangladesh Vet. J.*, 30(3-4): 105-113.
- Boado, E. and G. Rojas, 1990. Diagnosis and pathological study of goose diseases. *Revista-cubana-de-ciencia-Avicola*, 17(1): 19-25.
- Chamblee, T.N., J.D. Brake, C.D. Schultz and J.P. Thaxton, 1992. Yolk sac absorption and initiation of growth in broilers. *Poult. Sci.*, 71: 1811-1816.
- Choudhury, B., A. Chanda, P. Dasgupta, R.K. Dutta, L. Saha, S. Bhui, L. Saha and S. Bhui, 1993. Studies on yolk sac infection in poultry, antibiogram of isolates and correlation between *in-vitro* and *in-vivo* drug action. *Indian J. Anim. Hlth.*, 32(1): 21-23.
- Deeming, D.C., 1995. Possible effect of microbial infection on yolk utilization in ostrich chicks. *Vet. Rec.*, 136(11): 270-271.
- Desmidt, M., R. Ducatelle and F. Haesebrouck, 1998. Serological and bacteriological observations on experimental infection with *Salmonella harder* in chickens. *Vet. Microbiol.*, 60(2-4): 259-269.
- Dhillon, A.S., H.L. Shivaprasad, T.P. Roy, B. Alisantosa, D. Schaberg, D. Bandli and S. Johnson, 2001. Pathogenicity of environmental origin Salmonellas in specific pathogen-free chicken. *Poult. Sci.*, 80(9): 1323-1328.
- Fuller, R. and D.J. Jayne-Williams, 1968. The origin of bacteria recovered from the peritoneum and yolk sac of healthy chickens. *British Poult. Sci.*, 9: 159-163.
- Ghodasara, D.J., B.P. Joshi, P.B. Jani, R.M. Gangopadhyay and K.S. Prajapati, 1992. Pattern of mortality in chicken. *Indian Vet. J.*, 69(10): 888-890.
- Jordan, F.T.W., 1990. *Poultry Diseases*, 3rd Ed., ELBS and Bailliere Tindall, London, U.K., pp:40.
- Khan, K. A., S.A. Khan, S. Hamid, A. Aslam and M. Rabbani, 2002. A study on the pathogenesis of yolk retention in broiler chicks. *Pakistan Vet. J.*, 22(4): 175-180.
- Knizetova, H., B. Knizi and J. Cerveny, 1989. Size of yolk sac in waterfowl and changes during 24 hours after hatching. *Proceedings of International Symposium on Waterfowl Production, The Satellite Conference for the XVIII World's Poult. Congress, September 11-18, Beijing, China*, pp: 242-245.
- Leeson, S., J.P. Walker and J.D. Summers, 1978. Environmental temperature and the incidence of unabsorbed yolks in sexed broiler chickens. *Poult. Sci.*, 57(1): 316-318.
- Linzitto, O.R., N.A. Mendenez and H.D. Aberio, 1988. *Escherichia coli* isolates from fowls: Pathological and bacteriological studies. *Therios*, 12: 327-330, 332-334.
- Martin, P.W., 1996. Enlarged yolk sac in broiler chicks. Effects of prolonged exposure to hatcher environment on yolk sac size in three-day old broilers. *Poult. Digest*, 55(6): 20, 22-25.

- Moran, E.T. and B.S. Reinhart, 1980. Poult yolk sac amount and composition upon placement. Effect of breeder age, egg weight, sex and subsequent change with feeding or fasting. *Poult. Sci.*, 59: 1521-1528.
- Murakami, H., Y. Akiba and M. Horiguchi, 1992. Growth and utilization of nutrients in newly hatched chick with or without removal of residual yolk. *Growth, Development and Aging*, 56(2): 75-84.
- Mutalib, A.A. and J. Hanson, 1989. Avian Salmonellosis. *Canadian Vet. J.*, 30(2): 178.
- Noy, Y., Z. Uni and D. Sklan, 1996. Routes of yolk utilization in the newly hatched chick. *British Poult. Sci.*, 37(5): 987-995.
- O'Brien, J.D.P., 1988. *Salmonella enteritidis* infection in broiler chickens. *Vet. Rec.*, 122(9): 214.
- Pisarski, R.K., H. Malec and L. Malec, 1998a. Effect of initial feeding of broilers on the growth rate, yolk sac absorption and selected blood chemical parameters. *Medycyna Weterynaryjna*, 54(6): 409-412.
- Pisarski, R.K., L. Malec, B.W. Borzemska and H. Malec, 1998b. Effect of first feeding time of broiler chicks upon the body weight, yolk sac absorption and some constituents of blood plasma. *Medycyna Weterynaryjna*, 54(9): 607-611.
- Ramhoff, A.L., 1960. *The Avian Embryo*, Mac Millan Company, New York, pp: 1077.
- Rathore, B.S., R. Singh and S.S. Khera, 1985. Survey on causes of poultry mortality in India, based on postmortem examinations conducted at ten diagnostic centres. *Indian J. Poult. Sci.*, 20(2): 135-139.
- Rehman, R., M. Rabbani, S.A. Khan and C.M. Saleem, 1996. Pathological aspects of early chick mortality due to bacterial infections. *Pakistan. J. Sci. Res.*, 48(3-4): 101-107.
- Roy, S. and S.K. Misra, 1989. Common duck diseases and their control. *Poult. Adviser*, 22(5): 33-39.
- Rudy, A., 1991. The effect of microbial contamination of incubators on the health of broiler chicks in first days of life. *Zeszyty-Naukowe-Akademii-Rolniczej-we-Wroclawiu,-Weterynaria*, 49:19-26.
- Sainsbury, D., 1992. *Poultry Health and Management*, 3rd Ed., Blackwell Scientific Publications, London, U.K., pp: 116.
- Sander, J.E., E.M. Willingham, J.L. Wilson and S.G. Thayer, 1998. The effect of inoculating *Enterococcus faecalis* into the yolk sac on chick quality and maternal antibody absorption. *Avian Dis.*, 42(2): 359-363.
- Santos, G.A. and F.G. Silversides, 1996. Utilization of the sex-linked gene for imperfect albinism I. Effect of early weight loss on chick metabolism. *Poult. Sci.*, 75: 1321-1329.
- Sarma, D.R.L., N.L. Char, M.R.K. Rao, D.I. Khan and G. Narayana, 1985. A comprehensive study on bacterial flora isolated from yolk sac infection (omphalitis) in chicks. *Indian J. Poult. Sci.*, 20(4): 262-266.
- Schonhofen, C.A. and R.G.F. Garcia, 1981. Aspergillus omphalitis in chicks. *Arquivos de biologia-e-Tecnologia*, 24(4): 437-438.
- Seigo, S., H. Iwamori and K. Hirai, 1970. Studies on the organisms isolated from chicks with omphalitis. *Japanese Poult. Sci.*, 4: 57-61.
- Sharada, R., G. Kirshnappa, R. Raghavan, R.N.S. Gowda and H.A. Upendra, 1999. Isolation and serotyping of *Escherichia coli* from different pathological conditions in poultry. *Indian J. Poult. Sci.*, 34(3): 366-369.
- Sharma, N.K. and R.K. Kaushik, 1986a. Incidence of infectious and non-infectious diseases in turkeys. *Indian J. Poult. Sci.*, 21(1): 53-56.
- Sharma, N.K. and R.K. Kaushik, 1986b. Surveillance of diseases of ducks. *Indian J. Anim. Hlth.*, 25(1): 1-5.
- Sharma, N.K. and R.K. Kaushik, 1986c. Surveillance of diseases of Japanese quails. *Indian J. Vet. Medicine*, 6(1): 48-50.
- Shivaprasad, H.L., 2000. Fowl typhoid and pullorum disease. *Review Sci. Technology*, 19(2): 405-424.
- Shrivastava, P.K., 1982. Mortality in chicks: Incidence and etiopathomorphological studies. *Vet. Res. J.*, 5(2): 143-144.
- Singh, H., B.B. Dash, P.K. Dash, S. Kumar, H. Singh and S. Kumar, 1993. Mortality pattern in indigenous guinea fowl under confinement rearing. *Indian J. Poult. Sci.*, 28(1): 56-62.
- Singh, M.P., R.S. Gupta and S.P. Singh, 1997. Pathogenicity of *Escherichia coli* isolated from colibacillosis in chicks and their antibiotic spectra. *Indian J. Comparative Microbiol. Immunol. Infec. Dis.*, 18(2): 142-146.
- Suneja, S.C., J.R. Sadana and C.K. Aggarwal, 1983. Studies on mortality pattern in quails as affected by sex, age and month of the year. *Haryana Veterinarian*, 22(2): 82-86.
- Suresh, S., Y.M. Sanhya and M. Morton, 1988. Disease patterns in 1987. *Poult. Adviser*, 21(4): 49-52.
- Thaxton, P., R.D. Wyatt and P.B. Hamilton, 1974. Effect of environmental temperature on paratyphoid infection in the neonatal chicken. *Poult. Sci.*, 53(1): 88-94.

- Thyagarajan, D., J. Ramakrishna, N. Ramamurthy, V. Venkataramanujam and V. Sundararasu, 1987. Mortality pattern in turkeys. *Kerala J. Vet. Sci.*, 18(1): 112-116.
- Utomo, B.N., S. Poemomo and Iskander, 1990. Bacteria isolated from chicken yolk sac infection at the Research Institute for Veterinary Science. *Penyakit Hewan*, 22(40): 102-105.
- Viswanath, A., A. Ramakrishnan and A.K.K. Unni, 1985. Mortality pattern of two strains of White Leghorns under hot humid conditions. *Kerala J. Vet. Sci.*, 16(1): 81-84.
- Wang, H., Q.G. Huo and S.B. Li, 1994. Nutrient transfer in yolk sac of feeding chick. *Acta Veterinaria et Zootechnica Sinica*, 25(2): 97-104.
- William, F.K., 1975. Omphalitis in poult, *Poult. Digest*, 12: 490.
- Yousaf, M.K., 1985. *Stress Physiology in Livestock*, Vol. III, Poultry, CRC Press, Florida, U.S.A., pp: 118-119.
- Zahdeh, A.H., A.H. Basoni and M.H. Awaad, 1984. Studies on the prevalence of bacterial agents isolated from yolk sacs of chicks showing omphalitis. *Poultry disease in Near East. Fac. Agriculture Univ. Jordan, Amman*, pp: 359-371.
- Zahdeh, A.H. 1987. Studies on the problem of omphalitis in baby chicks: Role of *Escherichia coli* *J. Egypt Vet. Med. Assoc.*, 47(1/2): 511-519.