



**HAL**  
open science

## On gradient bounds for the heat kernel on the Heisenberg group

Dominique Bakry, Fabrice Baudoin, Michel Bonnefont, Djilil Chafai

► **To cite this version:**

Dominique Bakry, Fabrice Baudoin, Michel Bonnefont, Djilil Chafai. On gradient bounds for the heat kernel on the Heisenberg group. 2007. hal-00179865v2

**HAL Id: hal-00179865**

**<https://hal.science/hal-00179865v2>**

Preprint submitted on 20 Nov 2007 (v2), last revised 5 Sep 2008 (v5)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On gradient bounds for the heat kernel on the Heisenberg group

D. Bakry, F. Baudoin, M. Bonnefont, D. Chafaï

Institut de Mathématiques de Toulouse

Université de Toulouse

CNRS 5219

November 21, 2007

## Abstract

It is known that the couple formed by the two dimensional Brownian motion and its Lévy area leads to the heat kernel on the Heisenberg group, which is one of the simplest sub-Riemannian space. The associated diffusion operator is hypoelliptic but not elliptic, which makes difficult the derivation of functional inequalities for the heat kernel. However, Driver and Melcher and more recently H.-Q. Li have obtained useful gradient bounds for the heat kernel on the Heisenberg group. We provide in this paper simple proofs of these bounds, and explore their consequences in terms of functional inequalities, including Cheeger and Bobkov type isoperimetric inequalities for the heat kernel.

**Keywords:** Heat kernel ; Heisenberg group ; functional inequalities ; hypoelliptic diffusions

**AMS-MSC:** 22E30 ; 60J60

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Elementary properties of the Heisenberg group</b>	<b>4</b>
<b>3</b>	<b>Reverse Poincaré inequalities</b>	<b>7</b>
<b>4</b>	<b>A proof of the Driver-Melcher inequality</b>	<b>9</b>
<b>5</b>	<b>Two proofs of the H.-Q. Li inequality</b>	<b>11</b>
5.1	Via a Cheeger type inequality . . . . .	11
5.2	Via a Poincaré type inequality and a complex quasi-commutation . . . . .	13
<b>6</b>	<b>Functional inequalities for the heat kernel</b>	<b>20</b>
6.1	Gross-Poincaré type inequalities . . . . .	20
6.2	Cheeger type isoperimetric inequalities . . . . .	21
6.3	Bobkov type isoperimetric inequalities . . . . .	23
6.4	Multi-times inequalities . . . . .	25

## 1 Introduction

Gradient bounds had proved to be a very efficient tool for the control of the rate of convergence to equilibrium, quantitative estimates on the regularization properties of heat kernels, functional

inequalities such as Poincaré, logarithmic Sobolev, Gaussian isoperimetric inequalities for heat kernel measures. The reader may take a look for instance at [2, 36, 37, 27, 1] and references therein. When dealing with the simplest examples, such as linear parabolic evolution equations (or heat kernels), those gradient bounds often rely on the control of the intrinsic Ricci curvature associated to the generator of the heat kernel. Those methods basically require some form of ellipticity of the generator.

## The elliptic case

Let  $\mathcal{M}$  be a complete connected Riemannian manifold of dimension  $n$  and let  $L$  be the associated Laplace-Beltrami operator, written in a local system of coordinates as

$$L(f)(x) = \sum_{i,j=1}^n a_{i,j}(x) \partial_{x_i x_j}^2 f(x).$$

The coefficients  $x \mapsto a_{i,j}(x)$  are smooth and the symmetric matrix  $(a_{i,j}(x))_{1 \leq i,j \leq n}$  is positive definite for every  $x$ . The “length of the gradient”  $|\nabla f|$  of a smooth  $f : \mathcal{M} \rightarrow \mathbb{R}$  is given by

$$\Gamma(f, f) = |\nabla f|^2 = \frac{1}{2}(L(f^2) - 2fLf) = \sum_{i,j=1}^n a_{i,j}(x) \partial_{x_i} f \partial_{x_j} f.$$

Let  $(P_t)_{t \geq 0} = (e^{tL})_{t \geq 0}$  be the heat semigroup generated by  $L$ . For every smooth  $f : \mathcal{M} \rightarrow \mathbb{R}$ , the function  $(t, x) \mapsto P_t(f)(x)$  is the solution of the heat equation associated to  $L$

$$\partial_t P_t(f)(x) = LP_t(f)(x) \quad \text{and} \quad P_0(f)(x) = f(x).$$

For every real number  $\rho \in \mathbb{R}$ , the following three propositions are equivalent (see [2, 27, 35]).

1.  $\forall f \in C_c^\infty(\mathcal{M}), \text{Ricci}(\nabla f, \nabla f) \geq \rho |\nabla f|^2$
2.  $\forall f \in C_c^\infty(\mathcal{M}), \forall t \geq 0, |\nabla P_t f|^2 \leq e^{-2\rho t} P_t(|\nabla f|^2)$
3.  $\forall f \in C_c^\infty(\mathcal{M}), \forall t \geq 0, |\nabla P_t f| \leq e^{-\rho t} P_t(|\nabla f|)$

This is the case for some  $\rho \in \mathbb{R}$  when  $\mathcal{M}$  is compact. This is also the case with  $\rho = 0$  when  $\mathcal{M}$  is  $\mathbb{R}^n$  equipped with the usual metric since  $\text{Ricci} \equiv 0$ . In this last example,  $L$  is the usual Laplace operator  $\Delta$  and the explicit formula for the heat kernel gives  $\nabla P_t f = P_t \nabla f$  for the usual gradient  $\nabla$  and thus  $|\nabla P_t f| \leq P_t |\nabla f|$ . Back to the general case, and following [2], the gradient bounds 2. or 3. above are equivalent to their infinitesimal version at time  $t = 0$ , which reads

$$\mathbb{I}_2(f, f) \geq \rho \Gamma(f, f)$$

where

$$\mathbb{I}_2(f, f) = \frac{1}{2}(L\Gamma(f, f) - 2\Gamma(f, Lf)) = |\nabla \nabla f|^2 + \text{Ric}(\nabla f, \nabla f).$$

The bound  $\mathbb{I}_2 \geq \rho \Gamma$  had proved to be a very efficient criterion for the derivation of gradient bounds for more general Markov processes, including for instance processes generated by an operator  $L$  with a first order linear part (i.e. with a potential).

In the equivalence above, one may add several other inequalities, including local Poincaré inequalities, local logarithmic Sobolev inequalities, and local Bobkov isoperimetric inequalities, and their respective reverse forms, with a specific constant involving  $e^{-\rho t}$ , see [2, 27]. Here the

term local means that they concern the probability measure  $P_t(\cdot)(x)$  for any fixed  $t$  and  $x$ . One may also replace in these inequalities  $e^{-\rho t}$  by any function  $c(t)$  continuous and differentiable at  $t = 0$  with  $c(0) = 1$  and  $c'(0) = -\rho$ . In the present paper, we will focus on the Heisenberg group, a non elliptic situation where these equivalences do not hold, but where some gradient bounds are still available and provide local inequalities of various types.

## The Heisenberg group

In recent years, some focus had been set on some degenerate situations, where the methods used for the elliptic case do not apply. One of the simplest example of such a situation is the Heisenberg group (see section 2 for the group structure). Namely, we consider on  $\mathbb{H} = \mathbb{R}^3$  the vector fields

$$X = \partial_x - \frac{y}{2}\partial_z \quad \text{and} \quad Y = \partial_y + \frac{x}{2}\partial_z$$

and the operator

$$L = X^2 + Y^2 = \partial_x^2 + \partial_y^2 + \frac{1}{4}(x^2 + y^2)\partial_z^2 + x\partial_{y,z}^2 - y\partial_{x,z}^2. \quad (1)$$

This operator is self-adjoint for the Lebesgue measure on  $\mathbb{R}^3$ . The matrix of second order derivatives associated to  $L$  is degenerate and thus  $L$  is not elliptic. If  $[U, V] = UV - VU$  stands for the commutator of  $U$  and  $V$ , then

$$Z := [X, Y] = \partial_z \quad \text{and} \quad [X, Z] = [Y, Z] = 0.$$

In particular,  $L$  is hypoelliptic in the Hörmander sense (the Lie algebra described by  $\{X, Y, Z\}$  is the Lie algebra of the Heisenberg group, see section 2). As a consequence, the heat semigroup  $(P_t)_{t \geq 0} = (e^{tL})_{t \geq 0}$  obtained by solving the heat equation associated to  $L$  admits a smooth density with respect to the Lebesgue measure on  $\mathbb{R}^3$ . It is remarkable that the Markov process associated to this semigroup is the couple formed by a Brownian motion on  $\mathbb{R}^2$  and its Lévy area, and for every fixed  $t > 0$  and  $\mathbf{x} \in \mathbb{H}$ , the probability distribution  $P_t(\cdot)(\mathbf{x})$  is a sort of Gaussian on  $\mathbb{H}$ . We refer to [6] and [34] for such probabilistic aspects. For this operator  $L$  we have also

$$\Gamma(f, f) = X(f)^2 + Y(f)^2 \quad (2)$$

and

$$\mathbb{E}_2(f, f) = X^2(f)^2 + Y^2(f)^2 + \frac{1}{2}(XY + YX)(f)^2 + \frac{1}{2}(Zf)^2 + 2(XZ(f)Y(f) - YZ(f)X(f)).$$

The presence of  $YZ(f)$  and  $XZ(f)$  in the  $\mathbb{E}_2$  expression forbids the existence of a constant  $\rho \in \mathbb{R}$  such that  $\mathbb{E}_2 \geq \rho\Gamma$  as functional quadratic forms. Therefore the methods used in the elliptic case to prove gradient bounds could not work. In other words, the Ricci tensor is everywhere  $-\infty$ . In fact, a closer inspection of the Ricci tensor of the elliptic operator  $X^2 + Y^2 + \epsilon Z^2$  when  $\epsilon$  goes to 0 shows that this operator has everywhere a Ricci tensor which is

$$\frac{1}{2\epsilon} \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

In the limit, one may consider that the lower bound of the Ricci tensor for  $L$  is everywhere  $-\infty$ . Despite this singularity, B. Driver and T. Melcher proved in [15] the existence of a finite positive constant  $C_2$  such that

$$\forall f \in \mathcal{P}^\infty(\mathbb{H}), \forall t \geq 0, \quad |\nabla P_t f|^2 \leq C_2 P_t(|\nabla f|^2). \quad (3)$$

where  $\mathcal{P}^\infty(\mathbb{H})$  is the class of smooth function from  $\mathbb{H}$  to  $\mathbb{R}$  with all partial derivatives of polynomial growth. Here  $C_2$  is the best constant, i.e. the smallest possible. As in the elliptic case, the gradient bound (3) implies a Poincaré inequality for  $P_t$ , since

$$P_t(f^2) - (P_t f)^2 = 2 \int_0^t P_s(|\nabla P_{t-s} f|^2) ds \leq 2tC_2 P_t(|\nabla f|^2). \quad (4)$$

The gradient bound (3) gives also a reverse Poincaré inequality for  $P_t$ , since

$$P_t(f^2) - (P_t f)^2 = 2 \int_0^t P_s(|\nabla P_{t-s} f|^2) ds \geq \frac{2t}{C_2} |(\nabla P_t f)|^2. \quad (5)$$

From the point of view of regularization, (5) is the most important, while (4) is more concerned with estimates on the heat kernel and concentration properties. More recently, H.-Q. Li showed in [28] that there exists a finite positive constant  $C_1$  such that

$$\forall f \in \mathcal{P}^\infty(\mathbb{H}), \forall t \geq 0, |\nabla P_t f| \leq C_1 P_t(|\nabla f|). \quad (6)$$

It is shown in [15] that  $C_1 \geq \sqrt{2}$  and  $C_2 \geq 2$ . The Jensen or Cauchy-Schwarz inequality for  $P_t$  gives  $C_2 \leq C_1^2$ , however, the exact values of  $C_1$  and  $C_2$  are not known to the authors knowledge. The gradient bound (6) is far more useful than (3), and has for instance many consequences in terms of functional inequalities for  $P_t$ , including Poincaré inequalities, Gross logarithmic Sobolev inequalities, Cheeger type inequalities, and Bobkov type inequalities, as presented in section 6. As we shall see later, (6) is much harder to obtain than (3).

More generally, one may consider for any  $p \geq 1$  and  $t \geq 0$  the best constant  $C_p(t)$  in  $[0, \infty]$  (i.e. the smallest possible, possibly infinite) such that

$$\forall f \in \mathcal{P}^\infty(\mathbb{H}), |\nabla P_t f|^p \leq C_p(t) P_t(|\nabla f|^p).$$

It is immediate that  $C_p(0) = 1$ . According to [15] and [28], for every  $p \geq 1$  and  $t > 0$ , the quantity  $C_p(t)$  belongs to  $(1, \infty)$  and does not depend on  $t$ . In particular,  $C_p$  is discontinuous at  $t = 0$ , and this reflects the fact that the  $\mathbb{E}_2$  curvature of  $L$  is  $-\infty$ .

The aim of this paper is mainly to provide simpler proofs of the gradient bounds (3) and (6). We also give in section 6 a collection of consequences of (6) in terms of functional inequalities for the heat kernel of the Heisenberg group. Section 2 gathers some elementary properties of the Heisenberg group used elsewhere. Section 3 provides a direct simple proof of the reverse Poincaré inequality (5) without using (3) or (6). Sections 4 and 5 provide elementary proofs of (3) and (6) respectively.

## 2 Elementary properties of the Heisenberg group

We summarize in this section the main properties of the Heisenberg group that we use in the present paper. For more details on the geometric aspects, we refer to [19, 33, 24]. The link with the Brownian motion and its Lévy area is considered for instance in [6] and [34]. From now on, we shall use the notation  $\langle f \rangle$  to denote the integral of a function  $f$  with respect to the Lebesgue measure in  $\mathbb{R}^3$  and  $\langle f, g \rangle = \langle fg \rangle$  for the scalar product of two functions in  $L^2(\mathbb{R}^3, \mathbb{R})$ . The Heisenberg group  $\mathbb{H}$  is the set of matrices

$$M(x, y, z) = \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix}$$

equipped with the following non-commutative product

$$M(x, y, z)M(x', y', z') = M(x + x', y + y', z + z' + xy').$$

The inverse of  $M(x, y, z)$  is  $M(-x, -y, xy - z)$ . It is often more convenient to work with the Lie algebra of the group. For that, we define

$$X = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad Y = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad Z = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

and we consider

$$N(x, y, z) = \exp(xX + yY + zZ),$$

which gives

$$N(x, y, z) = M\left(x, y, z + \frac{xy}{2}\right).$$

We shall therefore identify a point  $\mathbf{x} = (x, y, z)$  in  $\mathbb{R}^3$  with the matrix  $N(x, y, z)$  and endow  $\mathbb{R}^3$  with this group structure that we denote  $\mathbf{x} \bullet \mathbf{y}$  which is

$$(x, y, z) \bullet (x', y', z') = (x + x', y + y', z + z' + \frac{1}{2}(xy' - yx')).$$

The left invariant vector fields which are given by

$$X(f) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{x} \bullet (\epsilon, 0, 0)) - f(\mathbf{x})}{\epsilon} = (\partial_x - \frac{y}{2}\partial_z)(f),$$

$$Y(f) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{x} \bullet (0, \epsilon, 0)) - f(\mathbf{x})}{\epsilon} = (\partial_y + \frac{x}{2}\partial_z)(f),$$

$$Z(f) = \lim_{\epsilon \rightarrow 0} \frac{f(\mathbf{x} \bullet (0, 0, \epsilon)) - f(\mathbf{x})}{\epsilon} = \partial_z(f),$$

while the right invariant vector fields are given by

$$\hat{X}(f) = \lim_{\epsilon \rightarrow 0} \frac{f((\epsilon, 0, 0) \bullet \mathbf{x}) - f(\mathbf{x})}{\epsilon} = (\partial_x + \frac{y}{2}\partial_z)(f),$$

$$\hat{Y}(f) = \lim_{\epsilon \rightarrow 0} \frac{f((0, \epsilon, 0) \bullet \mathbf{x}) - f(\mathbf{x})}{\epsilon} = (\partial_y - \frac{x}{2}\partial_z)(f).$$

There is no  $\hat{Z}$  since for those points, left and right multiplications coincide. The Lie algebra structure is described by the identities  $[X, Y] = Z$  and  $[X, Z] = [Y, Z] = 0$ . In what follows, we are mainly interested in the operator  $L = X^2 + Y^2$ , and the associated heat semigroup  $(P_t)_{t \geq 0} = (e^{tL})_{t \geq 0}$ . We shall make a strong use of symmetries in what follows. They are described by the Lie algebra of the vector fields which commute with  $L$ . This Lie algebra is 4-dimensional and is generated by the vector fields  $\hat{X}$ ,  $\hat{Y}$ ,  $Z$ , and  $\Theta = x\partial_y - y\partial_x$ . The first ones, which correspond to the right action, commute with  $X, Y, Z$  (as it is the case on any Lie group), the last one reflects the rotational invariance of  $L$ . There is also another vector field which plays an important role : the dilation operator  $D$ , described by

$$D = \frac{1}{2}(x\partial_x + y\partial_y) + z\partial_z.$$

This operator  $D$  satisfies

$$[L, D] = L. \quad (7)$$

Let  $(T_t)_{t \geq 0} = (e^{tD})_{t \geq 0}$  be the group of dilations generated by  $D$ , that is

$$T_t f(x, y, z) = f(e^{t/2}x, e^{t/2}y, e^t z).$$

From the commutation relations, one deduces

$$P_t T_s = T_s P_{e^s t}, \quad (8)$$

and

$$P_t D = D P_t + t P_t L. \quad (9)$$

Since  $P_t$  commutes with left translations, if  $l_{\mathbf{x}}(f)(\mathbf{y}) = f(\mathbf{x}\mathbf{y})$ , then

$$P_t(f)(\mathbf{x}) = l_{\mathbf{x}} P_t(f)(0) = P_t(l_{\mathbf{x}} f)(0).$$

Here we have just formalized the fact that  $(P_t)_{t \geq 0}$  is a heat semigroup on a group. Moreover, since 0 is a fixed point of the dilation group, one has from equation (8)

$$P_t(f)(0) = P_1(T_{\log t} f)(0).$$

This explains why  $P_1(f)(0)$  gives the whole  $(P_t f)_{t \geq 0}$ . It is well known that

$$P_1(f)(0) = \int_{\mathbb{R}^3} f(\mathbf{y}) h(\mathbf{y}) d\mathbf{y},$$

where  $d\mathbf{y}$  is the Lebesgue measure on  $\mathbb{R}^3$  and the function  $h$  has the following Fourier representation in the  $z$  variable

$$h(x, y, z) = \frac{1}{8\pi^2} \int_{-\infty}^{+\infty} e^{i\lambda z} \exp\left(-\frac{r^2}{4} \lambda \coth \lambda\right) \frac{\lambda}{\sinh \lambda} d\lambda. \quad (10)$$

where  $r^2 = x^2 + y^2$  if  $\mathbf{x} = (x, y, z)$ . This formula appeared independently in the works of Gaveau and Lévy. It is not easy to deduce from this formula any good estimates on  $h$ , and it is not even easy to see that  $h$  is positive. Nevertheless, there are quite precise bounds on this function and its derivatives, see for instance [17, 18] and [28, 29], which may be expressed in terms of the Carnot-Carathéodory distance. The Carnot-Carathéodory distance may be defined as

$$d(\mathbf{x}, \mathbf{y}) = \sup_{\{f \text{ such that } \Gamma(f, f) \leq 1\}} f(x) - f(y)$$

where  $\Gamma$  is as in (2). Here, the explicit form is not easy to write, but we shall only need to express the distance from 0 to  $\mathbf{x}$ . In order to do that, it is better to describe the constant speed geodesics starting from 0. First, straight lines in the  $(x, y)$  plane passing through  $(0, 0)$  are geodesics, and the other ones are helices, whose orthogonal projection on the horizontal plane  $\{z = 0\}$  is a circle containing the origin. If a point  $\mathbf{x}(s)$  is moving at constant speed (for the Carnot-Carathéodory distance) on the geodesic, its horizontal projection  $\mathbf{p}(s)$  moves with unit speed (for the Euclidean distance) on this circle.

Moreover, the height  $z(t)$  of the point  $\mathbf{x}(s)$  is the surface between the segment  $(0, \mathbf{p}(s))$  and the circle, which comes from the fact that for any curve in  $\mathbb{R}^3$  whose tangent vector is a linear combination of  $X$  and  $Y$ , one has

$$dz = \frac{1}{2}(x dy - y dx),$$

that is the area spanned by the point  $\mathbf{p}(s)$  in the plane. Notice that all the geodesics end on the  $z$  axis, which is therefore the cut-locus of the point 0. Those geodesics may be parameterized as follows, using a complex notation for the horizontal projection  $\mathbf{p}(s)$

$$\mathbf{p}(s) = u \left( 1 - \exp \left( \frac{is}{|u|} \right) \right) \quad \text{and} \quad z(s) = \frac{|u|^2}{2} \left( \frac{s}{|u|} - \sin \left( \frac{s}{|u|} \right) \right). \quad (11)$$

Here,  $u$  is the center of the circle which is the horizontal projection of the geodesic, and  $s$  is distance from 0 ( $s \in (0, 2\pi |u|)$ ). When  $|u|$  goes to infinity, we recover the straight lines.

If we call  $d(\mathbf{x})$  the Carnot-Carathéodory distance from 0 to  $\mathbf{x}$ , it is easy to see from that if  $\mathbf{x} = (x, y, z)$ , and  $s = d(\mathbf{x})$ , then

$$d \left( \frac{x}{s}, \frac{y}{s}, \frac{z}{s^2} \right) = 1.$$

This corresponds to a change of  $u$  into  $\frac{u}{s}$ . Now, the unit ball for the Heisenberg metric is between two balls for the usual Riemannian metric of  $\mathbb{R}^3$ , and therefore one concludes easily that the ratio

$$\frac{(x^2 + y^2)^2 + z^2}{d^4(x, y, z)}$$

is bounded above and below. Although, if  $R = ((x^2 + y^2)^2 + z^2)^{1/4}$ , the function  $\Gamma(R, R)$  is bounded above but not below.

Let  $h$  be the heat kernel density at time 1 and at the origin, given by (10). The main properties of  $h$  used in the present paper are the following. Here for real valued functions  $a$  and  $b$ , we use the notation  $a \simeq b$  when the ratio  $a/b$  is bounded above and below by some positive constants. First,

$$h(\mathbf{x}) \simeq \frac{\exp(-\frac{d^2(\mathbf{x})}{4})}{\sqrt{1 + \|\mathbf{x}\| d(\mathbf{x})}}, \quad (12)$$

where  $\|\mathbf{x}\|$  denotes the Euclidean norm of the projection of  $\mathbf{x}$  onto the plane  $\{z = 0\}$ . Then, for some constant  $C$

$$\Gamma(\log h)(\mathbf{x}) \leq Cd(\mathbf{x}) \quad (13)$$

and

$$|Z(\log h)| \leq C. \quad (14)$$

The last one is not completely explicit in [28] but follows easily from the estimation of  $W_1$  page 376 of this paper.

**Lemma 2.1.** *The Schwartz space  $\mathcal{S}$  of smooth rapidly decreasing functions on the Heisenberg group  $\mathbb{H}$  is left globally stable by  $L$  and by  $P_t$  for any  $t \geq 0$ .*

*Proof.* If  $R = (x^2 + y^2)^2 + z^2$ , then an elementary computation shows that for any positive integer  $q$ , there exists a real constant  $B_q > 0$  such that  $L((1 + R)^{-q}) \leq B_q(1 + R)^{-q}$ . As a consequence,  $P_t((1 + R)^{-q}) \leq e^{B_q t}(1 + R)^{-q}$ . We may see the class  $\mathcal{S}$  as the class of smooth functions such that for any non negative integers  $a, b, c$  and any positive integer  $q$ , the function  $\hat{X}^a \hat{Y}^b Z^c(f)$  is bounded above by  $(1 + R)^{-q}$ . From that and the above, it is clear that if  $f$  is in  $\mathcal{S}$ , such is  $P_t f$ . On the other hand, the stability of  $\mathcal{S}$  by  $L$  is straightforward.  $\square$

### 3 Reverse Poincaré inequalities

We show here how to deduce a reverse Poincaré inequality as (5). The method is simple and direct, and does not rely on a gradient bound like (3) or (6).



**Theorem 3.1 (Reverse local Poincaré inequality).** *For any  $t \geq 0$  and any  $f \in \mathcal{C}_c^\infty(\mathbb{H})$ ,*

$$t\Gamma(P_t f, P_t f) \leq P_t(f^2) - (P_t f)^2.$$

*Proof.* Since we work on a group, it is enough to prove this for  $\mathbf{x} = 0$ . Then, thanks to the dilation properties, it is enough to prove it for  $t = 1$ . Now, consider the vector field  $\hat{X}$ , which coincides with  $X$  at  $\mathbf{x} = 0$ , and let as before  $h$  be the density of the heat kernel for  $t = 1$  and  $\mathbf{x} = 0$  given by (10). We want to bound, for a smooth compactly supported function  $f$

$$(\hat{X}P_1 f)^2 = (P_1(\hat{X}f))^2 = \langle f, \hat{X}h \rangle^2$$

where the last identity comes from integration by parts. The first remark is that we may suppose that  $\langle fh \rangle = 0$  since we may always add any constant to  $f$ . We then use Cauchy-Schwartz inequality under the measure  $h(\mathbf{x})d\mathbf{x}$  to get

$$(\hat{X}P_1 f)^2 \leq P_1(f^2)\langle \hat{X}(\log h)^2 h \rangle.$$

Using the same method for  $Y$ , we get

$$\Gamma(P_1 f, P_1 f) \leq (P_1(f^2) - (P_1 f)^2)\langle \hat{\Gamma}(\log h)h \rangle$$

where

$$\hat{\Gamma}(u, u) = \hat{X}(u)^2 + \hat{Y}(u)^2.$$

Now, the rotational invariance of  $h$ , which comes from  $[\Theta, L] = 0$ , shows that  $\hat{\Gamma}(\log h) = \Gamma(\log h)$ , and gives a reverse local Poincaré inequality with the constant  $C = \langle \Gamma(\log h)h \rangle$ . It remains to compute this constant. For that, we use the dilation operator  $D$  and the formula (9). In 0, we have  $Df = 0$ , and therefore it reads for  $t = 1$ , for any  $f$ ,

$$P_1((L - D)f) = 0,$$

which means that  $h$  is the invariant measure for the operator  $L - D$ , or in other words that

$$(L + D + 2)h = 0$$

since the adjoint of  $D$  is  $-D - 2$ . Multiply then both sides by  $\log h$  and taking integrals gives

$$\langle \log h, Lh \rangle = -\langle \Gamma(\log h, \log h), h \rangle$$

by using an integration by parts. Moreover, we have

$$\langle \log h, (D + 2)h \rangle = -\langle h, D \log h \rangle = -\langle Dh \rangle = 2\langle h \rangle = 2$$

and therefore

$$\langle \Gamma(\log h, \log h), h \rangle = 2.$$

If we had done the same reasoning on  $\mathbb{R}^n$  with the usual Laplace operator, and the corresponding dilation operator, we would have found a reverse Poincaré inequality with constant  $n/2$  instead of  $\frac{1}{2}$ . The reason comes from symmetry properties and the same will allow us to divide the constant by 2 in the Heisenberg case. In fact, because of the rotational invariance of  $h$ , we have, for any vector  $(a, b) \in \mathbb{R}^2$  with  $a^2 + b^2 = 1$ ,

$$\langle (a\hat{X}(\log h) + b\hat{Y}(\log h))^2, h \rangle = \langle (\hat{X}(\log h))^2, h \rangle = \frac{1}{2}\langle \hat{\Gamma}(\log h, \log h), h \rangle = 1.$$

Now, we may write

$$\Gamma(P_1 f, P_1 f)(0) = \sup_{a^2+b^2=1} (a\hat{X}P_1(f) + b\hat{Y}P_1(f))^2$$

and use the same Cauchy-Schwarz inequality to improve the bound to

$$\Gamma(P_1 f, P_1 f) \leq P_1(f^2) - (P_1 f)^2.$$

□

**Remark 3.2 (Optimal constants).** *Equality is achieved in theorem (3.1) when  $f = \hat{X}(\log h)$  for instance. To see it, note that by symmetry,  $\langle (\hat{X} \log h)^2 h \rangle = \langle (\hat{Y} \log h)^2 h \rangle = 1$ . By the rotational invariance of  $h$ , we get more generally that  $\langle (a\hat{X} \log h + b\hat{Y} \log h)^2 h \rangle = 1$  for any  $a^2 + b^2 = 1$ . As a consequence,  $\langle \hat{X}(\log h), \hat{Y}(\log h), h \rangle = 0$ . For  $f = \hat{X}(\log h)$ , this gives  $X P_1 f(0) = \langle \hat{X} f, h \rangle = -\langle f, \hat{X}(\log h) h \rangle = -1$  and  $Y P_1 f = 0$ , which is the desired equality. Notice the difference with the elliptic case: for the heat semigroup  $(P_t)_{t \geq 0}$  in  $\mathbb{R}^n$  or for any manifold with non negative Ricci curvature, one has for every  $t \geq 0$  and any smooth  $f$ ,*

$$2t\Gamma(P_t f, P_t f) \leq P_t(f^2) - (P_t f)^2.$$

*Notice also, as we already mentioned in the introduction, that the H.-Q. Li gradient bound (6) provides simply by semigroup interpolation a result similar to theorem 3.1, with a constant  $2tC_1^{-2}$  instead of  $t$ . These two constants are equal if and only if  $C_1 = \sqrt{2}$ . At the level of reverse local Poincaré inequalities, a necessary and sufficient condition for the efficiency of the semigroup interpolation technique is that  $C_1 = \sqrt{2}$ . Similarly, by using the Drivier and Melcher gradient bound (3) we obtain the condition  $C_2 = 2$ . It is thus tempting to conjecture that  $C_1^2 = C_2 = 2$ .*

**Remark 3.3 (Carnot groups).** *In the class of nilpotent groups, there is an interesting subclass, which are the Carnot groups, that is the nilpotent groups with dilations, see [19]. Let  $(X_1, \dots, X_{n_0})$  the generators at the first level of such a Carnot group, and  $L = X_1^2 + \dots + X_{n_0}^2$ . In those groups again there is a dilation operator  $D$  such that  $[L, D] = L$  and  $D^* = -D - \frac{n}{2}Id$ . The parameter  $n$  is what is called the homogeneous dimension of the group. The same applies in this case, except that we no longer have always enough rotations to insure that  $\Gamma(h, h) = \hat{\Gamma}(h, h)$  where the hat corresponds to the “chiral” action. But we may replace this argument by the fact that  $\hat{P}_t(\cdot)(0) = P_t(\cdot)(0)$  where  $(\hat{P}_t)_{t \geq 0} = (\exp(t\hat{L}))_{t \geq 0}$  and we would get as a bound*

$$\Gamma(P_t f, P_t f) \leq \frac{n}{2n_0 t} (P_t(f^2) - (P_t f)^2)$$

*recovering at the same time the Heisenberg and Euclidean cases. For the  $(2p+1)$ -dimensional Heisenberg group  $\mathbb{H}_{2p+1}$  we have  $n_0 = 2p$  while the homogeneous dimension is  $2p+2$ , and therefore here the inequality writes*

$$\Gamma(P_t f, P_t f) \leq \frac{p+1}{2pt} (P_t(f^2) - (P_t f)^2)$$

*and the constant approaches the Euclidean one when  $p$  goes to infinity.*

## 4 A proof of the Driver-Melcher inequality

We give here an elementary proof of the Driver and Melcher gradient bound (3). The argument is simply an integration by parts followed by the upper bound on  $\Gamma(\log h, \log h)$  obtained in section

3. Indeed, from the inequalities (12) and (13), it is quite clear that the constant

$$A = \int \|\mathbf{x}\| \Gamma(\log h, \log h)(\mathbf{x}) h(\mathbf{x}) d\mathbf{x} \quad (15)$$

is finite, where  $\|\mathbf{x}\|$  denotes as usual the Euclidean norm of the horizontal projection of the point  $\mathbf{x}$ . Then, we have the following theorem.

**Theorem 4.1.** *With the constant  $A$  defined by (15), we have for every  $t \geq 0$  and  $f \in C_c^\infty(\mathbb{H})$ ,*

$$\Gamma(P_t f, P_t f) \leq 2(A + 4)P_t(\Gamma(f, f)).$$

*Proof.* We assume that  $\mathbf{x} = 0$  (by group action) and  $t = 1$  (by dilation). Then, we write

$$XP_1 f(0) = P_1(\hat{X}f)(0) = \langle (X + yZ)f, h \rangle.$$

An integration by parts for  $\langle yZ(f), h \rangle = \langle y(XY - YX)(f), h \rangle$  gives

$$\langle X(f), (yY(\log h) + 1)h \rangle - \langle Y(f), yX(\log h)h \rangle$$

and a similar formula holds for  $YP_1 f$ . Next, we take a vector  $(a, b) \in \mathbb{R}^2$  of unit norm and we use the Cauchy-Schwarz inequality to get

$$(aXP_1(f)(0) + bYP_1(f)(0))^2 \leq P_1(X(f)^2)A_1 + P_1(Y(f)^2)A_2$$

where

$$A_1 = P_1[((yY(\log h) + 2)a - xY(\log h)b)^2]$$

and

$$A_2 = P_1[((xX(\log h) + 2)b - yY(\log h)a)^2].$$

The desired inequality comes then from the upper bound

$$\max(A_1, A_2) \leq A_1 + A_2 \leq 2(A + 4).$$

Note that the obtained constant  $2(A + 4)$  is certainly not the optimal one.  $\square$

**Remark 4.2 (Counter example).** *Unlike the elliptic case, the reverse local Poincaré (5) and the local Poincaré (4) inequalities are not in general equivalent. A simple example is provided on  $\mathbb{R}^2$  with the operator*

$$L = \partial_x^2 + x\partial_y$$

for which the corresponding diffusion process starting from  $(x, y)$  is up to some constant

$$U_t = \left( x + B_t, y + tx + \int_0^t B_s ds \right)$$

where  $(B_t)_{t \geq 0}$  is a Brownian motion on  $\mathbb{R}$ . In this example, the heat kernel is Gaussian and the semigroup  $(P_t)_{t \geq 0}$  is quite easy to compute, while  $\Gamma(f, f) = (\partial_x f)^2$ . In this situation, it is easy to see, using Cauchy-Schwarz inequality, that

$$tC\Gamma(P_t f) \leq P_t(f^2) - (P_t f)^2$$

for some constant  $C < 2$ , while the inequality

$$P_t f^2 - (P_t f)^2 \leq C(t)P_t \Gamma(f, f)$$

does not hold for any constant  $C(t)$ , as one may see with a function  $f$  depending on  $y$  only. For example, with  $f(x, y) = y$ , one has  $P_t f = y + tx$ , which depends on the variable  $x$ . This kind of hypoelliptic situation differs strongly from the case of the Heisenberg group, since here  $\Gamma(f, f) = 0$  does not imply that  $f$  is constant.

## 5 Two proofs of the H.-Q. Li inequality

In this section, we propose two alternate and independent proofs of the H.-Q. Li inequality (6). The first proof uses some basic symmetry considerations and a particular case of the Cheeger inequality of theorem 6.3 that we have to show by hands. The second proof relies on an explicit commutation between the complex gradient and the heat semigroup. Both mainly rely on the previous sharp estimates on the heat kernel that were obtained in [18].

### 5.1 Via a Cheeger type inequality

**Lemma 5.1.** *There exists a positive finite constant  $C$  such that for any smooth  $f : \mathbb{H} \rightarrow \mathbb{R}$  which vanishes on the unit ball centered at 0 for the Carnot-Carathéodory distance, we have*

$$\left| \int f h d\mathbf{x} \right| \leq C \int |\nabla f| h d\mathbf{x}$$

where  $h$  is as before the density of  $P_1(0, d\mathbf{x})$ .

*Proof.* Let us use the polar coordinates described in the formula 11, that is that we parameterize the exterior of the unit ball by  $(u, s)$ , with  $u \in \mathbb{C}^2$ ,  $|u| \geq \frac{1}{2\pi}$  and  $s \in (1, 2\pi|u|)$ , with

$$(x + iy, z) = \left( u \left( 1 - \exp\left(\frac{is}{|u|}\right) \right), \frac{|u|^2}{2} \left( \frac{s}{|u|} - \sin\left(\frac{s}{|u|}\right) \right) \right). \quad (16)$$

The unit ball is the set  $\{s = 1\}$ , and since  $f$  is supported outside the unit ball, we write

$$|f(u, s)| = \left| \int_1^s \nabla f(u, t) \cdot e_t dt \right| \leq \int_1^s |\nabla f|(u, t) dt$$

where  $e_t$  is the unit vector along the geodesic. Let us write  $A(u, t) du dt$  the Lebesgue measure on  $\mathbb{R}^3$  in those coordinates (we shall see the precise formula below). We write

$$\int |f(u, s)| h(u, s) du ds \leq \int |\nabla f|(u, t) \left( \int_t^{2\pi|u|} A(u, s) h(u, s) ds \right) du dt$$

and we shall have proved our inequality when we have proved that

$$\int_t^{2\pi|u|} A(u, s) h(u, s) ds \leq CA(u, t) h(u, t),$$

for any  $(u, t)$  such that  $|u| \geq \frac{1}{2\pi}$  and  $t \geq 1$ . In this computation, we forget the points in the  $(x, y)$  plane and the  $z$ -axis, but this is irrelevant since they have 0-measure. The computation of the Jacobian gives

$$A(u, s) = 16 |u| \sin\left(\frac{s}{2|u|}\right) \left( \frac{s}{2|u|} - \sin\left(\frac{s}{2|u|}\right) \right)$$

and the estimate (12) shows that we may replace  $h(u, s)$  by

$$\frac{\exp\left(-\frac{s^2}{4}\right)}{\sqrt{1 + 2s|u| \sin\left(\frac{s}{2|u|}\right)}}$$

since the Euclidean norm of the horizontal projection of the point whose coordinates are  $(u, s)$  is  $2|u|\sin(\frac{s}{2|u|})$ . Setting  $\tau = \frac{s}{2|u|}$  and  $r = |u|$ , the question is therefore to check that, for some constant  $C$ , for any  $r \geq \frac{1}{2\pi}$ , for any  $\tau_0 \geq \frac{1}{2r}$ , one has

$$r \int_{\tau_0}^{\pi} \frac{\sin \tau (\tau - \sin \tau)}{\sqrt{1 + 4r^2 \tau \sin \tau}} e^{-\tau^2 r^2} d\tau \leq C \frac{\sin \tau_0 (\tau_0 - \sin \tau_0)}{\sqrt{1 + 4r^2 \tau_0 \sin \tau_0}} e^{-\tau_0^2 r^2}.$$

Up to some constant, we may replace  $\sin \tau (\tau - \sin \tau)$  by  $\tau^4$  on  $(0, \frac{\pi}{2})$  and by  $\pi - \tau$  on  $(\frac{\pi}{2}, \pi)$ . In the same way, we may replace  $\sqrt{1 + 4r^2 \tau \sin \tau}$  by  $r\tau$  when  $\tau < \frac{\pi}{2}$  (since  $r\tau \geq \frac{1}{2}$ ) and by  $1 + r\sqrt{\pi - \tau}$  when  $\tau \in (\frac{\pi}{2}, \pi)$ .

We first consider the case where  $\tau_0 < \frac{\pi}{2}$ , and divide the integral into  $\int_{\tau_0}^{\pi/2}$  and  $\int_{\pi/2}^{\pi}$ . Using the above estimates, these integral can be bounded by the correct term by using the fact that

$$\int_A^{\infty} s^p \exp(-s^2) ds \leq C_p A^{p-1} \exp(-A^2).$$

When  $\tau_0 > \frac{\pi}{2}$ , one uses the same estimates, bounding above  $\pi - \tau$  by  $\pi - \tau_0$  and  $(1 + r\sqrt{\pi - \tau})^{-1}$  by 1 in the integral, and using the fact that  $r$  is bounded below on our domain.

Observe that the same reasoning on a ball of radius  $\epsilon$  would provide a constant which goes to infinity when  $\epsilon$  goes to 0, as for the usual heat kernel on  $\mathbb{R}^d$ .  $\square$

We are now in position to prove the H.-Q. Li inequality (6).

*Proof of (6).* With the help of the previous lemma, we may reduce the study of the H.-Q. Li inequality to functions which are

- either supported in a ball of radius 1 for the Carathéodory metric;
- either supported in a cylinder of radius 1 around the  $z$  axis;
- either supported outside a cylinder around the  $z$ -axis.

Indeed, let see how one may reduce first to the case of a function supported either in a ball or outside a ball. If  $f$  is any smooth function and  $\phi$  a cutoff function with values 1 on a ball of radius  $< 1$  and vanishing outside a ball of radius 1, we write  $f = f\phi + f(1 - \phi) = f_1 + f_2$ . Assuming that we know the inequality for  $f_1$  and  $f_2$ , then we bound

$$\langle \hat{X}(f), h \rangle \leq C(\langle |\nabla f_1| + |\nabla f_2|, h \rangle)$$

and since

$$|\nabla f_1| + |\nabla f_2| \leq |\nabla f| + 2|f| |\nabla \phi|$$

one uses lemma 5.1 to bound

$$\langle |f|, |\nabla \phi| h \rangle \leq C \|\nabla \phi\|_{\infty} \langle |\nabla f|, h \rangle$$

since  $|\nabla \phi|$  is supported outside the unit ball. We repeat the same operation with a cutoff function for the neighborhood of the  $z$ -axis.

Now, when  $f$  is supported inside the ball, we may use the method that we used in the proof of theorem 4.1, and the fact that  $|\nabla \log h|(\mathbf{x}) \leq Cd(\mathbf{x})$ , which is bounded on the unit ball.

If  $f$  is supported inside the cylinder around the  $z$ -axis, we write, with section 2 notations,

$$\langle \hat{X}(f), h \rangle = \langle X(f), h \rangle + \langle f, \frac{y}{2} Z(\log h) h \rangle$$

and then we use the fact that  $\frac{y}{2}Z(\log h)$  is bounded on the cylinder. It remains to bound

$$\langle |f|, h \rangle \leq C \langle |\nabla f|, h \rangle$$

thanks to lemma 5.1.

It remains to deal with a function which is supported outside a cylinder around the  $z$ -axis. We shall chose another integration by parts. For that, let us use a complex notation and write

$$\nabla(f) = X(f) + iY(f) \quad \text{and} \quad \hat{\nabla}(f) = \hat{X}(f) - i\hat{Y}(f).$$

Notice the change of sign in front of  $i$  in the second expression. We want to bound

$$\langle \hat{\nabla}(f), h \rangle = -\langle f, \hat{\nabla}h \rangle.$$

Now, since  $h$  is radial, we have

$$\hat{\nabla}h = \frac{x - iy}{x + iy} \nabla h$$

which comes from the fact that  $x\partial_y h = y\partial_x h$ . Let us call  $\Psi(x, y) = \exp(-2i\theta)$  the function  $\frac{x-iy}{x+iy}$ , where  $\theta$  is the angle in the plane  $(x, y)$ . Then, we integrate again by parts and get

$$\langle \hat{\nabla}f, h \rangle = -\langle f, \Psi(x, y)\nabla h \rangle = \langle \nabla f, \Psi(x, y)h \rangle + \langle f, \nabla(\Psi)h \rangle.$$

We then conclude observing that  $\Psi$  is bounded and  $|\nabla\Psi|$  is bounded outside the cylinder around the  $z$  axis. We therefore have

$$\left| \langle \hat{\nabla}(f), h \rangle \right| \leq \langle |\nabla f|, h \rangle + C \langle |f|, h \rangle$$

and we use again lemma 5.1 to conclude the proof.  $\square$

## 5.2 Via a Poincaré type inequality and a complex quasi-commutation

In  $\mathbb{R}^n$ , it is known that the gradient  $\nabla$  commute with the Laplace operator. This commutation leads to the commutation between  $\nabla$  and the heat semigroup  $P_t = e^{t\Delta}$  and therefore to the inequality:

$$|\nabla P_t f| = |P_t \nabla f| \leq P_t |\nabla f|.$$

In the Heisenberg group, the Lie algebra structure:

$$[X, Y] = Z, [X, Z] = [Y, Z] = 0$$

leads to the commutation:

$$(X + iY)L = (L - 2iZ)(X + iY),$$

where  $L = X^2 + Y^2$ . This leads to the *formal* commutation:

$$(X + iY)P_t = e^{t(L-2iZ)}(X + iY).$$

We shall use this commutation and give an analytical sense to the previous commutation of semigroups. It will give a new proof of H.Q. Li's inequality by studying the possible kernels for the integral operator  $e^{t(L-2iZ)}$  against complex gradients.

In order to exploit the rotational invariance, we shall use the cylindric coordinates  $x = r \cos \theta$ ,  $y = r \sin \theta$  in which the vector fields  $X$  and  $Y$  read

$$\begin{aligned} X &= \cos \theta \partial_r - \frac{\sin \theta}{r} \partial_\theta - \frac{1}{2} r \sin \theta \partial_z \\ Y &= \sin \theta \partial_r + \frac{\cos \theta}{r} \partial_\theta + \frac{1}{2} r \cos \theta \partial_z \\ Z &= \partial_z. \end{aligned}$$

The heat kernel associated to  $(P_t)_{t \geq 0}$  writes here in cylindric coordinates

$$p_t(r, z) = \frac{1}{8\pi^2} \int_{-\infty}^{+\infty} e^{i\lambda z} \frac{\lambda}{\sinh \lambda t} e^{-\frac{r^2}{4} \lambda \cotanh \lambda t} d\lambda. \quad (17)$$

We begin by the analytic properties of  $p_t(r, z)$  in the variable  $z$ .

**Lemma 5.2.** *Let  $t > 0$  and  $r \geq 0$ . The function*

$$z \rightarrow p_t(r, z) - \frac{1}{4\pi^2 (t + iz + \frac{r^2}{4})^2} - \frac{1}{4\pi^2 (t - iz + \frac{r^2}{4})^2}$$

*admits an analytic continuation on  $\{z \in \mathbb{C}, |\mathbf{Im} z| < \frac{r^2}{4} + 3t\}$ . The function*

$$z \rightarrow p_t(r, z)$$

*admits therefore a meromorphic continuation on  $\{z \in \mathbb{C}, |\mathbf{Im} z| < \frac{r^2}{4} + 3t\}$  with double poles at  $-i(t + \frac{r^2}{4})$  and  $i(t + \frac{r^2}{4})$ .*

*Proof.* Let  $t > 0$  and  $r \geq 0$ . By using the expression (17) for  $p_t(r, z)$ , and

$$\begin{aligned} \frac{1}{(t + iz + \frac{r^2}{4})^2} &= \int_0^{+\infty} e^{-i\lambda z} e^{-\lambda t} e^{-\lambda \frac{r^2}{4}} \lambda d\lambda, \\ \frac{1}{(t - iz + \frac{r^2}{4})^2} &= \int_0^{+\infty} e^{i\lambda z} e^{-\lambda t} e^{-\lambda \frac{r^2}{4}} \lambda d\lambda, \end{aligned}$$

we obtain

$$\begin{aligned} p_t(r, z) - \frac{1}{4\pi^2 (t + iz + \frac{r^2}{4})^2} - \frac{1}{4\pi^2 (t - iz + \frac{r^2}{4})^2} \\ = \frac{1}{8\pi^2} \int_{-\infty}^{+\infty} e^{i\lambda z} \left( \frac{e^{-\frac{r^2}{4} |\lambda| \cotanh |\lambda| t}}{\sinh |\lambda| t} - 2e^{-\frac{1}{4} |\lambda| r^2 - |\lambda| t} \right) |\lambda| d\lambda \end{aligned}$$

and the desired result follows easily.  $\square$

For any  $t > 0$ ,  $r \geq 0$ , and  $z \in \mathbb{C} - \{-i(t + \frac{1}{4}r^2)\}$  such that  $|\mathbf{Im} z| < \frac{r^2}{4} + 3t$ , let us denote

$$p_t^*(r, z) = p_t(r, z) - \frac{1}{4\pi^2 (t + iz + \frac{r^2}{4})^2}.$$

We have the following commutation property.

**Proposition 5.3.** *If  $f : \mathbb{H} \rightarrow \mathbb{R}$  is a smooth function with compact support, then*

$$(X + iY)\mathbf{P}_t f(0) = \int_{\mathbb{H}} p_t^*(r, z + 2it)(X + iY)f(r, \theta, z)rdrd\theta dz, \quad t > 0.$$

*Proof.* Due to the identities  $[X, Y] = Z$  and  $[X, Z] = [Y, Z] = 0$ , we have

$$(X + iY)L = (L - 2iZ)(X + iY).$$

If  $f(r, \theta, z) = e^{i\lambda z}g(\theta, z)$  we deduce from the previous commutation,

$$(X + iY)P_t f(0) = e^{2\lambda t}(P_t(X + iY)f)(0) = e^{2\lambda t} \int_{\mathbb{H}} p_t(r, z)(X + iY)f(r, \theta, z)rdrd\theta dz.$$

Let us now observe that for  $t > 0$ ,

$$(X + iY)\frac{1}{(t + iz + \frac{r^2}{4})^2} = 0$$

and thus

$$(X + iY)p_t^* = (X + iY)p_t,$$

consequently,

$$(X + iY)P_t f(0) = e^{2\lambda t} \int_{\mathbb{H}} p_t^*(r, z)(X + iY)f(r, \theta, z)rdrd\theta dz,$$

and the result follows by integrating by parts with respect to the variable  $z$ .  $\square$

As a first consequence, we deduce that there exists a finite constant  $C > 0$  such that for every smooth function compactly supported inside a Carnot-Carathéodory ball  $\mathbf{B}_R$  of radius  $R > 0$ ,

$$|\nabla P_1 f|(0) \leq CP_1(|\nabla f|)(0).$$

But of course, here, the constant  $C = \sup_{\mathbf{B}_R} \frac{|p_1^*(r, z + 2i)|}{p_1(r, z)}$  depends on  $R$  and we shall see below that it blows up when  $R \rightarrow +\infty$ .

Now, if  $f$  is a smooth function with compact support that vanishes in a ball with radius  $R > 0$ , where  $R$  is big enough to avoid the pole ( $r = 2, z = 0$ ) of  $z \rightarrow p_1(r, z + 2i)$ , we have the commutation:

$$(X + iY)\mathbf{P}_1 f(0) = \int_{\mathbb{H}} p_1(r, z + 2i)(X + iY)f(r, \theta, z)rdrd\theta dz, \quad t > 0.$$

**Proposition 5.4.** *There exists  $R > 0$  such that*

$$\sup_{r, z \notin \mathbf{B}_R} \frac{|p_1(r, z + 2i)|}{p_1(r, z)} < +\infty.$$

*Proof. Step 1.* We show that for any  $\eta > 0$ ,

$$\sup_{r \geq 3, r^2 \geq \eta|z|} \frac{\left| p_1^*(r, z + 2i) + \frac{1}{4\pi^2(-1 + iz + \frac{r^2}{4})^2} \right|}{p_1(r, z)} < +\infty.$$



For convenience, we may assume  $z > 0$ . Let us first observe that on our domain:

$$p_1^*(r, z + 2i) + \frac{1}{4\pi^2 \left(-1 + iz + \frac{r^2}{4}\right)^2} = \frac{1}{8\pi^2} \int_{-\infty}^{+\infty} e^{-2\lambda} e^{i\lambda z} \frac{\lambda}{\sinh \lambda} e^{-\frac{r^2}{4} \lambda \cotanh \lambda} d\lambda \quad (18)$$

From [18], it is known that for fixed  $r, z$ , the function

$$g : \lambda \rightarrow -i\lambda z + \frac{r^2}{4} \lambda \cotanh \lambda,$$

has a unique critical point in the strip  $\{|\operatorname{Im} \lambda| < \frac{\pi}{2}\}$ . This critical point is  $i\theta(r, z)$ , where  $\theta(r, z)$  the unique solution in  $(0, \frac{\pi}{2})$  of the equation

$$\mu\left(\frac{1}{2}\theta(r, z)\right)r^2 = 4z,$$

with  $\mu(\theta) = \frac{\theta}{\sin^2 \theta} - \cotan \theta$ . At this critical point, we have

$$g(i\theta(r, z)) = \frac{d^2(r, z)}{4},$$

where  $d(r, z)$  is the Carnot-Carathéodory distance to the point  $(r, z)$  (it does not depend on the angular variable  $\theta$ ). In fact, our function  $g$  corresponds to  $g(r, z, \lambda) = f\left(\frac{r}{\sqrt{2}}, \frac{z}{2}, 2\lambda\right)$  where  $f$  is the function studied in [18].

Moreover the function  $s \rightarrow \operatorname{Re}g(s + i\theta(r, z))$ , grows with  $|s|$ , and has a global minimum at  $s = 0$ , indeed a tedious computation shows that

$$\begin{aligned} \operatorname{Re}(g(s + i\theta(r, z)) - g(i\theta(r, z))) &= \frac{\sinh^2 2s}{\sinh^2 2s + \sin^2 2\theta(r, z)} (2s \cotanh 2s - 2\theta(r, z) \cotan 2\theta(r, z)) r^2 \\ &\geq \frac{\sinh^2 2s}{\sinh^2 2s + 1} (2s \cotanh 2s - 1) r^2 \\ &\geq 0. \end{aligned}$$

Let us finally observe that the previous computation also shows that there exists  $\delta > 0$  such that for  $s \in [-1, 1]$

$$\operatorname{Re}g(s + i\theta(r, z)) \geq \frac{d^2(r, z)}{4} + \delta r^2 s^2.$$

With all this in hands, we can now turn to our proof. We first start by changing the contour of integration in (18):

$$\begin{aligned} \int_{-\infty}^{+\infty} e^{-2\lambda} e^{i\lambda z} \frac{\lambda}{\sinh \lambda} e^{-\frac{r^2}{4} \lambda \cotanh \lambda} d\lambda &= \int_{\operatorname{Im} \lambda = \theta(\sqrt{r^2 - 8}, z)} e^{-2\lambda} e^{i\lambda z} \frac{\lambda}{\sinh \lambda} e^{-\frac{r^2}{4} \lambda \cotanh \lambda} d\lambda \\ &= \int_{\operatorname{Im} \lambda = \theta(\sqrt{r^2 - 8}, z)} e^{i\lambda z} \frac{\lambda}{\sinh \lambda} e^{-\left(\frac{r^2}{4} - 2\right) \lambda \cotanh \lambda} e^{2\lambda - 2\lambda \cotanh \lambda} d\lambda \end{aligned}$$

Therefore, by denoting

$$U(\lambda) = e^{2\lambda - 2\lambda \cotanh \lambda} \frac{\lambda}{\sinh \lambda}$$

we get

$$\begin{aligned} \left| \int_{-\infty}^{+\infty} e^{-2\lambda} e^{i\lambda z} \frac{\lambda}{\sinh \lambda} e^{-\frac{r^2}{4}\lambda \cotanh \lambda} d\lambda \right| &\leq e^{-\frac{d(\sqrt{r^2-8}, z)^2}{4}} \int_{|s|\leq 1} e^{-(r^2-8)\delta^2 s^2} \left| U(s + i\theta(\sqrt{r^2-8}, z)) \right| ds \\ &\quad + e^{-\frac{d(\sqrt{r^2-8}, z)^2}{4}} \int_{|s|\geq 1} e^{-(r^2-8)\delta^2} \left| U(s + i\theta(\sqrt{r^2-8}, z)) \right| ds \\ &\leq C_1 \frac{e^{-\frac{d(r, z)^2}{4}}}{r}, \end{aligned}$$

where we used the fact that on the domain on which we work, the difference  $d(\sqrt{r^2-8}, z) - d(r, z)$  is uniformly bounded. Finally, from the lower estimate of [29], on the considered domain,

$$p_t(r, z) \geq C_2 \frac{e^{-\frac{d(r, z)^2}{4}}}{r}.$$

**Step 2.** We show that there exists  $\eta > 0$  such that

$$\sup_{|z|\geq 1, r^2 \leq \eta|z|} \frac{\left| p_1^*(r, z + 2i) + \frac{1}{4\pi^2(-1 + iz + \frac{r^2}{4})^2} \right|}{p_1(r, z)} < +\infty.$$

We first start by giving an analytic representation of

$$p_1^*(r, z + 2i) + \frac{1}{4\pi^2(-1 + iz + \frac{r^2}{4})^2}$$

that is valid on the domain on which we work. As in the previous proof, we assume  $z > 0$ . Due to the Cauchy theorem, we can change the contour of integration in the representation (17), to get with  $0 < \varepsilon < \pi$ ,

$$\begin{aligned} p_1(r, z) &= \frac{1}{8\pi^2} \sum_{k=1}^{+\infty} \int_{|\lambda - ik\pi| = \varepsilon} e^{i\lambda z} \frac{\lambda}{\sinh \lambda} e^{-\frac{r^2}{4}\lambda \cotanh \lambda} d\lambda \\ &= \frac{-i}{8\pi^2} \sum_{k=1}^{+\infty} \int_{|\lambda| = \varepsilon} e^{i(-i\lambda + ik\pi)z} \frac{(-i\lambda + ik\pi)}{\sinh(-i\lambda + ik\pi)} e^{-\frac{r^2}{4}(-i\lambda + ik\pi)\cotanh(-i\lambda + ik\pi)} d\lambda \\ &= \frac{-i}{8\pi^2} \int_{|\lambda| = \varepsilon} \frac{e^{-(\pi-\lambda)(z - \frac{r^2}{4} \cotan \lambda)}}{1 + e^{-\pi(z - \frac{r^2}{4} \cotan \lambda)}} \left( \frac{\pi}{1 + e^{-\pi(z - \frac{r^2}{4} \cotan \lambda)}} - \lambda \right) \frac{d\lambda}{\sin \lambda} \end{aligned}$$

Therefore, for  $z > 0$ ,

$$\begin{aligned} &p_1^*(r, z + 2i) + \frac{1}{4\pi^2(-1 + iz + \frac{r^2}{4})^2} \\ &= \frac{-i}{8\pi^2} \int_{|\lambda| = \varepsilon} e^{2i\lambda} \frac{e^{-(\pi-\lambda)(z - \frac{r^2}{4} \cotan \lambda)}}{1 + e^{-\pi(z - \frac{r^2}{4} \cotan \lambda)}} \left( \frac{\pi}{1 + e^{-\pi(z - \frac{r^2}{4} \cotan \lambda)}} - \lambda \right) \frac{d\lambda}{\sin \lambda} \end{aligned}$$

On our domain, if  $\eta$  is small enough, when  $r, z \rightarrow +\infty$ ,  $\mathbf{Re}(z - \frac{r^2}{4} \cotan \lambda)$  goes uniformly on the circle  $|\lambda| = \varepsilon$  to  $+\infty$ . Consequently, on our domain

$$\left| p_1^*(r, z + 2i) + \frac{1}{4\pi^2(-1 + iz + \frac{r^2}{4})^2} \right| \leq c_1 \left| \int_{|\lambda| = \varepsilon} e^{2i\lambda} e^{-(\pi-\lambda)(z - \frac{r^2}{4} \cotan \lambda)} (\pi - \lambda) \frac{d\lambda}{\sin \lambda} \right|$$

for some finite positive constant  $c_1$ . By choosing  $\varepsilon = \pi - 2\theta(r, z)$ , we have

$$\begin{aligned} \int_{|\lambda|=\varepsilon} e^{2i\lambda} e^{-(\pi-\lambda)\left(z - \frac{r^2}{4} \cotan \lambda\right)} (\pi - \lambda) \frac{d\lambda}{\sin \lambda} \\ = \int_{|\lambda|=\pi-2\theta(r,z)} e^{2i\lambda} e^{-(\pi-\lambda)\left(z - \frac{r^2}{4} \cotan \lambda\right)} (\pi - \lambda) \frac{d\lambda}{\sin \lambda}, \end{aligned}$$

where the function  $\theta(r, z)$  has been introduced above. At this stage, we can follow step by step the proof of Theorem 2.17 in [18] (the only difference is in the function  $V$  which we take equal to  $V(\lambda) = e^{2i\lambda} \frac{\pi-\lambda}{\sin \lambda}$ ) to get an estimate on our domain :

$$\left| \int_{|\lambda|=\pi-2\theta(r,z)} e^{2i\lambda} e^{-(\pi-\lambda)\left(z - \frac{r^2}{4} \cotan \lambda\right)} (\pi - \lambda) \frac{d\lambda}{\sin \lambda} \right| \leq c_2 \frac{e^{-\frac{d(r,z)^2}{4}}}{\sqrt{rd(r,z)}}$$

for some finite positive constant  $c_2$ . Finally, the lower estimate of [29] leads to the conclusion.  $\square$

We can now reprove Li's inequality by using a partition of the unity (which is here simpler than in the previous subsection) and the Poincare inequality on balls:

Let  $f : \mathbb{H} \rightarrow \mathbb{R}$  be a smooth positive function compactly supported and let  $0 \leq \phi \leq 1$  be a smooth function that takes the value 1 on a ball  $\mathbf{B}_{R_1}$  and the value 0 outside the ball  $\mathbf{B}_{R_2}$  where  $R_1 < R_2$ , with  $R_1$  big enough to include the pole in  $\mathbf{B}_{R_1}$ .

We have

$$\begin{aligned} (X + iY)P_1 f(0) &= (X + iY)P_1 \phi f(0) + (X + iY)P_1 (1 - \phi) f(0) \\ &= \int_{\mathbb{H}} p_1^*(r, z + 2i)(X + iY)(f\phi)(r, \theta, z) r dr d\theta dz \\ &\quad + \int_{\mathbb{H}} p_1(r, z + 2i)(X + iY)(f(1 - \phi))(r, \theta, z) r dr d\theta dz \\ &= \int_{\mathbb{H}} \phi(r, \theta, z) p_1^*(r, z + 2i)(X + iY) f(r, \theta, z) r dr d\theta dz \\ &\quad + \int_{\mathbb{H}} (1 - \phi(r, \theta, z)) p_1(r, z + 2i)(X + iY) f(r, \theta, z) r dr d\theta dz \\ &\quad + \frac{1}{4\pi^2} \int_{\mathbb{H}} f(r, \theta, z) \frac{(X + iY)\phi(r, \theta, z)}{\left(-1 + iz + \frac{r^2}{4}\right)^2} r dr d\theta dz. \end{aligned}$$

Therefore

$$|\nabla P_1 f(0)| \leq C P_1 |\nabla f|(0) + \left| \frac{1}{4\pi^2} \int_{\mathbb{H}} f(r, \theta, z) \frac{(X + iY)\phi(r, \theta, z)}{\left(-1 + iz + \frac{r^2}{4}\right)^2} r dr d\theta dz \right|,$$

Now, we estimate  $\left| \int_{\mathbb{H}} f(r, \theta, z) \frac{(X+iY)\phi(r, \theta, z)}{(-1+iz+\frac{r^2}{4})^2} r dr d\theta dz \right|$  thanks to the Poincaré inequality (see [31]):

$$\begin{aligned}
& \left| \int_{\mathbb{H}} f(r, \theta, z) \frac{(X+iY)\phi(r, \theta, z)}{(-1+iz+\frac{r^2}{4})^2} r dr d\theta dz \right| \\
&= \left| \int_{\mathbb{H}} (f(r, \theta, z) - m) \frac{(X+iY)\phi(r, \theta, z)}{(-1+iz+\frac{r^2}{4})^2} r dr d\theta dz \right| \quad (m \text{ is the mean of } f \text{ on } \mathbf{B}_{R_2}) \\
&\leq C_1 \int_{\mathbf{B}_{R_2}} |f(r, \theta, z) - m| r dr d\theta dz \\
&\leq C_2 \int_{\mathbf{B}_{R_2}} |\nabla f|(r, \theta, z) r dr d\theta dz \\
&\leq C_3 P_1 |\nabla f|(0).
\end{aligned}$$

This completes the proof of H.Q. Li's inequality.

Interestingly, it is not possible to find a function  $\Phi$  on  $\mathbb{H}$  such that:

- $(X+iY)\phi = 0$ ;
- The ratio  $\frac{|p_1^*(r, z+2i) - \Phi(r, \theta, z)|}{p_1(r, z)}$  is bounded.

Indeed, the first point implies that  $\Phi$  can be written:

$$\Phi(r, \theta, z) = H \left( \frac{r^2}{4} + iz, re^{i\theta} \right),$$

where  $H : \{z_1 \in \mathbb{C}, \mathbf{Re}(z_1) \geq 0\} \times \mathbb{C} \rightarrow \mathbb{C}$  is analytic in  $z_1$  and  $z_2$ . Now, due to the estimate of Proposition 5.4 and the estimate on  $p_1$ , it would imply that for  $r$  and  $z$ , such that  $r^2 + |z|$  is big enough:

$$\left| H \left( \frac{r^2}{4} + iz, re^{i\theta} \right) + \frac{1}{4\pi^2 (-1+iz+\frac{r^2}{4})^2} \right| \leq Ae^{-B(r^2+|z|)}$$

where  $A$  and  $B$  are strictly positive constants. Now, we have the following lemma that prevents the the existence of such  $H$ :

**Lemma 5.5.** *Let  $f : \{z_1 \in \mathbb{C}, \mathbf{Re}(z_1) \geq 0\} \times \mathbb{C} \rightarrow \mathbb{C}$  be analytic in  $z_1$  and  $z_2$ . If there exist strictly positive constants  $A$  and  $B$  such that*

$$\forall r \geq 0, \forall z \in \mathbb{R}, \forall \theta \in [0, 2\pi], \quad |f(r^2 + iz, re^{i\theta})| \leq Ae^{-B(r^2+|z|)}$$

then  $f = 0$ .

*Proof.* Let  $r \geq 0, z \in \mathbb{R}$ . The function  $z_2 \rightarrow f(r^2 + iz, z_2)$  is analytic, therefore from the maximum principle we have

$$|f(r^2 + iz, z_2)| \leq Ae^{-B(|z_2|^2+|z|)},$$

for  $|z_2| \leq r$ . Consequently, on the set  $\mathbf{Re}(z_1) \geq |z_2|^2$  we have

$$|f(z_1, z_2)| \leq Ae^{-B(|z_2|^2+|\mathbf{Im}(z_1)|)}.$$

By using the analytic function  $z_1 \rightarrow f(z_1, z_2)$ , a translation, and a multiplication by  $e^{-z_1}$  we would therefore obtain a function  $g$  analytic on the set  $\mathbf{Re}(z) > 0$  such that

$$|g(z)| \leq \alpha e^{-\beta|z|}$$

with  $\alpha, \beta > 0$ , and such function has clearly to be 0 (Use for instance the conformal equivalence between the set  $\mathbf{Re}(z) > 0$  and the open unit disc to get a function  $h$  analytic on the disc that satisfy the estimate  $|g(z)| \leq \alpha' e^{-\frac{\beta'}{|z|}}$ ).

□

## 6 Functional inequalities for the heat kernel

Most of the consequences of the classical gradient bounds under a  $\mathbb{I}_2$  curvature assumption remain true under an H.-Q. Li gradient bound. In the sequel, we derive, by interpolation from the gradient bound (6), several local functional inequalities of Gross-Poincaré-Cheeger-Bobkov type for the heat kernel on the Heisenberg group. The term *local* means that these inequalities concern the probability measure  $P_t(\cdot)(\mathbf{x})$  at fixed  $t$  and  $\mathbf{x}$ , in contrast to inequalities for the invariant measure. These local inequalities can be seen as global inequalities for Gaussian measures on the Heisenberg group. In the literature, these inequalities and interpolations were mainly developed in Riemannian settings under a  $\mathbb{I}_2$  curvature assumption. Rigorously, the semigroup interpolations used in the sequel rely on the existence of an algebra of functions  $\mathcal{A}$  from  $\mathbb{H}$  to  $\mathbb{R}$  stable by the action of the heat kernel. Thanks to lemma 2.1, the Schwartz class  $\mathcal{S}$  of smooth and rapidly decreasing functions in  $\mathbb{R}^3$  may play this role in the case of the Heisenberg group  $\mathbb{H}$ .

### 6.1 Gross-Poincaré type inequalities

One of the first consequence of the gradient bound (6) is Gross-Poincaré type local inequalities, also called  $\varphi$ -Sobolev inequalities in [11, 22]. Namely, let  $\varphi : I \rightarrow \mathbb{R}$  be a smooth convex function defined on an open interval  $I \subset \mathbb{R}$  such that  $\varphi'' > 0$  on  $I$  and  $-1/\varphi''$  is convex on  $I$ .

**Theorem 6.1 (Local Gross-Poincaré inequalities).** *By using the notations of (6), for every  $t \geq 0$ , every  $\mathbf{x} \in \mathbb{H}$ , and every  $f \in C_c^\infty(\mathbb{H}, I)$ ,*

$$P_t(\varphi(f)) - \varphi(P_t f) \leq C_1^2 t P_t(\varphi''(f)|\nabla f|^2). \quad (19)$$

*Proof.* One can assume that the support of  $f$  is strictly included in  $I$ . Since  $L$  is a diffusion operator,  $L(\alpha(f)) = \alpha'(f)Lf + \alpha''(f)\Gamma f$  for any  $f \in C_c^\infty(\mathbb{H}, \mathbb{R})$  and any smooth  $\alpha : \mathbb{R} \rightarrow \mathbb{R}$ . By the semigroup and the diffusion properties,

$$P_t(\varphi(f)) - \varphi(P_t f) = \int_0^t \partial_s P_s(\varphi(P_{t-s}f)) ds = \int_0^t P_s(\varphi''(P_{t-s}f)|\nabla P_{t-s}f|^2) ds.$$

Now, (6) gives  $|\nabla P_{t-s}f|^2 \leq C_1^2 (P_{t-s}(|\nabla f|))^2$ . Next, by the Cauchy-Schwarz inequality or alternatively by the Jensen inequality for the bivariate convex function  $(u, v) \mapsto \varphi''(u)v^2$ , we get  $\varphi''(P_{t-s}f)(P_{t-s}(|\nabla f|))^2 \leq P_{t-s}(\varphi''(f)|\nabla f|^2)$ , which gives the desired result. □

- for  $\varphi(u) = u \log(u)$  on  $I = (0, \infty)$ , we get a Gross logarithmic Sobolev inequality, mentioned for instance in [28] (see also [20, 21]),

$$P_t(f \log(f)) - P_t(f) \log(P_t(f)) \leq C_1^2 t P_t(f^{-1} |\nabla f|^2); \quad (20)$$

- for  $\varphi(u) = u^p$  on  $I = (0, \infty)$  with  $1 < p \leq 2$ , we get a Beckner-Latała-Oleszkiewicz type inequality (see [7, 25])

$$\frac{P_t(f^p) - (P_t(f))^p}{p-1} \leq p C_1^2 t P_t(f^{p-2} |\nabla f|^2); \quad (21)$$

- for  $\varphi(u) = u^2$  on  $I = \mathbb{R}$ , we get a Poincaré inequality, mentioned in [15],

$$P_t(f^2) - (P_t(f))^2 \leq 2 C_1^2 t P_t(|\nabla f|^2). \quad (22)$$

We have seen in the introduction that a local Poincaré inequality like (22) can be also obtained from the Driver and Melcher gradient bound (3), with a constant  $2C_2$  instead of  $2C_1^2$ . However, the inequalities (20) and (21) need the stronger gradient bound (6) of H.-Q. Li. They also imply the local Poincaré inequality (22) by linearization. It is shown in [12, Theorem 4.4] that the convexity of the bivariate function  $(u, v) \mapsto \varphi''(u)v^2$  is equivalent to the convexity of the  $\varphi$ -entropy functional and also to the tensorization property of the  $\varphi$ -entropy functional. This fact is related to the infinite dimensional nature of (19). The inequality (21) interpolates between (20) (let  $p \rightarrow 1^+$ ) and (22) (take  $p = 2$ ). The linearity with respect to  $t$  of the constant in front of the right hand side of (19) is related to the fact that  $(P_t)_{t \geq 0}$  is a convolution semigroup, namely  $P_t(\cdot)(\mathbf{x})$  can be obtained from  $P_1(\cdot)(0)$  by  $\mathbf{x}$ -translation and  $\sqrt{t}$ -dilation in  $\mathbb{H}$ .

## 6.2 Cheeger type isoperimetric inequalities

As mentioned in the introduction, it is possible to deduce a reverse local Poincaré inequality from the gradient bounds (3) of Driver and Melcher or (6) of H.-Q. Li. However, the constants are not known precisely. A better constant is provided by theorem 3.1, which implies immediately that for every  $t \geq 0$  and every  $f \in \mathcal{C}_c^\infty(\mathbb{H}, \mathbb{R})$ ,

$$\|\nabla P_t f\|_\infty \leq \frac{1}{\sqrt{t}} \|f\|_\infty. \quad (23)$$

Cheeger derived in [13] a lower bound for the spectral gap of the Laplacian on a Riemannian manifold. This bound can be related to a sort of  $L^1$  Poincaré inequality, which has an isoperimetric content, see [14] and references therein. Here we derive such an inequality for the heat kernel by only using the gradient bound (6), by mixing arguments borrowed from [4] and [26].

**Theorem 6.2 (Local Cheeger type inequality).** *With the notations of (6), for every  $t \geq 0$ , every  $\mathbf{x} \in \mathbb{H}$ , and every  $f \in \mathcal{C}_c^\infty(\mathbb{H}, \mathbb{R})$ ,*

$$P_t(|f - P_t(f)(\mathbf{x})|)(\mathbf{x}) \leq 4C_1 \sqrt{t} P_t(|\nabla f|)(\mathbf{x}). \quad (24)$$

*Proof.* We adapt the method used in [26, p. 953] for the invariant measure in Riemannian

settings. For any  $g \in \mathcal{C}_c^\infty(\mathbb{H}, \mathbb{R})$  with  $\|g\|_\infty \leq 1$ , any  $t \geq 0$ , and any  $\mathbf{x} \in \mathbb{H}$ ,

$$\begin{aligned}
P_t((f - P_t(f)(\mathbf{x}))g)(\mathbf{x}) &= P_t(fg)(\mathbf{x}) - P_t(f)(\mathbf{x})P_t(g)(\mathbf{x}) \\
&= \int_0^t \partial_s P_s((P_{t-s}f)(P_{t-s}g))(\mathbf{x}) ds \\
&= 2 \int_0^t P_s(\Gamma(P_{t-s}f, P_{t-s}g))(\mathbf{x}) ds \\
&\leq 2 \int_0^t P_s(|\nabla P_{t-s}f| |\nabla P_{t-s}g|)(\mathbf{x}) ds \\
&\leq 2C_1 P_t(|\nabla f|)(\mathbf{x}) \int_0^t \frac{\|g\|_\infty}{\sqrt{(t-s)}} ds \\
&\leq 4C_1 \sqrt{t} P_t(|\nabla f|)(\mathbf{x}).
\end{aligned}$$

where we used the gradient bound (6) for  $f$  and the gradient bound (23) for  $g$ . The desired result follows then by  $L^1 - L^\infty$  duality by taking the supremum over  $g$ .  $\square$

Similarly, we get also the following correlation bound for every  $t \geq 0$  and  $f, g \in \mathcal{C}_c^\infty(\mathbb{H}, \mathbb{R})$ ,

$$|P_t(fg) - P_t(f)P_t(g)| \leq 2C_1^2 t \sqrt{P_t(|\nabla f|^2)} \sqrt{P_t(|\nabla g|^2)}. \quad (25)$$

When  $f = g$ , we recover the Poincaré inequality (22).

**Theorem 6.3 (Yet another local Cheeger type inequality).** *With the notations of (6), for every  $t \geq 0$ , every  $\mathbf{x} \in \mathbb{H}$ , and every ball  $B$  of  $\mathbb{H}$  for the Carnot-Carathéodory metric, there exists a real constant  $C_{B,t,\mathbf{x}} > 1$  such that for every function  $f \in \mathcal{C}_c^\infty(\mathbb{H}, \mathbb{R})$  which vanishes on  $B$ ,*

$$|P_t(f)(\mathbf{x})| \leq C_{B,t,\mathbf{x}} P_t(|\nabla f|)(\mathbf{x}). \quad (26)$$

*Proof.* Let  $g \in \mathcal{C}^\infty(\mathbb{H}, \mathbb{R})$  be such that  $\|g\|_\infty < \infty$  and  $g \equiv 1$  on  $B^c$ . Since  $fg = f$ , the computation made in the proof of theorem 6.2 provides

$$P_t(f)(\mathbf{x}) - P_t(f)(\mathbf{x})P_t(g)(\mathbf{x}) \leq 4C_1 \sqrt{t} \|g\|_\infty P_t(|\nabla f|)(\mathbf{x}).$$

For any arbitrary real number  $r \geq 1$ , the class of functions

$$\mathcal{C}_{B,r} = \{g \in \mathcal{C}^\infty(\mathbb{H}, \mathbb{R}) \text{ with } \|g\|_\infty \leq r \text{ and } g \equiv 1 \text{ on } B^c\}.$$

is not empty since it contains the constant function  $\equiv 1$ . Furthermore, since  $P_t(\cdot)(\mathbf{x})$  is a probability measure with non vanishing density, the following extrema

$$\alpha_-(B, r, t, \mathbf{x}) = \inf_{g \in \mathcal{C}_{B,r}} P_t(g)(\mathbf{x}) \quad \text{and} \quad \alpha_+(B, r, t, \mathbf{x}) = \sup_{g \in \mathcal{C}_{B,r}} P_t(g)(\mathbf{x})$$

are finite and non zero. Moreover, an elementary local perturbative argument on any element of the class  $\mathcal{C}_{B,r}$  shows that  $\alpha_-(B, r, t, \mathbf{x}) \alpha_+(B, r, t, \mathbf{x}) < 0$  as soon as  $r$  is large enough, say  $r \geq r_{B,t,\mathbf{x}}$ . Thus,  $P_t(f)(\mathbf{x})P_t(g)(\mathbf{x}) \leq 0$  for some  $g \in \mathcal{C}_{B,r}$ . The desired result follows then with  $C_{B,t,\mathbf{x}} = 4C_1 \sqrt{t} r_{B,t,\mathbf{x}}$ , since one can replace  $f$  by  $-f$  in the obtained inequality. Notice that  $C_{B,t,\mathbf{x}}$  blows up when  $\text{vol}(B) \searrow 0$ . Actually, this proof does not use the nature of the Heisenberg group  $\mathbb{H}$ , and relies roughly only on the diffusion property, the smoothness of the heat kernel and the gradient bound. However, on the Heisenberg group  $\mathbb{H}$ , the usage of translations and dilations and of the convolution semigroup nature of  $(P_t)_{t \geq 0}$  allows to precise the dependency of  $C_{B,t,\mathbf{x}}$  over  $t$  and  $\mathbf{x}$  by using  $\mathbf{x}$ -translation and  $\sqrt{t}$ -dilation.  $\square$

The isoperimetric content of (24) can be extracted by approximating an indicator with a smooth  $f$ , see for instance [4]. Namely, for any Borel set  $A \subset \mathbb{H}$  with smooth boundary, any  $t \geq 0$ , and any  $\mathbf{x} \in \mathbb{H}$ , we get by denoting  $\mu_{t,\mathbf{x}} = P_t(\cdot)(\mathbf{x})$ ,

$$\mu_{t,\mathbf{x}}(A)(1 - \mu_{t,\mathbf{x}}(A)) \leq 2C_1 \sqrt{t} \mu_{t,\mathbf{x}}^{\text{surface}}(\partial A) \quad (27)$$

where  $\mu_{t,\mathbf{x}}^{\text{surface}}$  is the surface measure of the boundary defined by

$$\mu^{\text{surface}}(\partial A) = \liminf_{r \rightarrow 0} \frac{\mu(A_r) - \mu(A)}{r} \quad \text{where } A_r = \{\mathbf{x} \in \mathbb{H}; d(\mathbf{x}, A) < r\}.$$

Here  $d$  is the Carnot-Carathéodory metric on  $\mathbb{H}$  considered in section 2. From (26), we get similarly for any ball  $B$  in  $\mathbb{H}$  and any Borel set  $A \subset B^c$  with smooth boundary,

$$\mu_{t,\mathbf{x}}(A) \leq C_{B,t,\mathbf{x}} \mu_{t,\mathbf{x}}^{\text{surface}}(\partial A). \quad (28)$$

### 6.3 Bobkov type isoperimetric inequalities

Let  $F_\gamma : \mathbb{R} \rightarrow [0, 1]$  be the cumulative probability function of the standard Gaussian distribution  $\gamma$  on the real line  $\mathbb{R}$ , given for every  $t \in \mathbb{R}$  by

$$F_\gamma(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t e^{-\frac{1}{2}u^2} du.$$

The Gaussian isoperimetric function  $\mathcal{I} : [0, 1] \rightarrow [0, (2\pi)^{-1/2}]$  is defined by  $\mathcal{I} = (F_\gamma)' \circ (F_\gamma)^{-1}$ . The function  $\mathcal{I}$  is concave, continuous on  $[0, 1]$ , smooth on  $(0, 1)$ , symmetric with respect to the vertical axis of equation  $u = 1/2$ , and satisfies to the differential equation

$$\mathcal{I}(u)\mathcal{I}''(u) = -1 \quad \text{for any } u \in [0, 1] \quad (29)$$

with  $\mathcal{I}(0) = \mathcal{I}(1) = 0$  and  $\mathcal{I}'(0) = -\mathcal{I}'(1) = \infty$ . Notice that  $\mathcal{I}(u) \geq u(1-u)$  for any real  $u \in [0, 1]$ , and that  $\mathcal{I}(u) \leq \min(u, 1-u)$  when  $u$  belongs to a neighborhood of  $1/2$ .

**Lemma 6.4 (Yet another uniform gradient bound).** *With the notations of (6), for every  $t \geq 0$  and  $f \in C_c^\infty(\mathbb{H}, (0, 1))$ ,*

$$\mathcal{I}(P_t f) - P_t(\mathcal{I}(f)) \leq C_1^2 \sqrt{2t} P_t(|\nabla f|). \quad (30)$$

*Proof.* The inequality (30) was obtained by Bobkov in [8] for the standard Gaussian measure on  $\mathbb{R}$ . Later, it was generalized in [4], by using semigroup techniques, to Riemannian settings under a  $\mathbb{I}_2$  curvature assumption. We give here a proof by adapting the argument given in [4, p. 261-263] from invariant measure settings to local settings. One may assume that  $\varepsilon \leq f \leq 1 - \varepsilon$  for some  $\varepsilon > 0$ . By the diffusion property and (29)

$$\begin{aligned} [\mathcal{I}(P_t f)]^2 - [P_t(\mathcal{I}(f))]^2 &= - \int_0^t \partial_s [P_s(\mathcal{I}(P_{t-s} f))]^2 ds \\ &= -2 \int_0^t P_s(\mathcal{I}(P_{t-s} f)) P_s \left( \mathcal{I}''(P_{t-s} f) |\nabla P_{t-s} f|^2 \right) ds \\ &= +2 \int_0^t P_s(\mathcal{I}(P_{t-s} f)) P_s \left( \frac{|\nabla P_{t-s} f|^2}{\mathcal{I}(P_{t-s} f)} \right) ds. \end{aligned}$$



Next, the Cauchy-Schwarz inequality or alternatively the Jensen inequality for the bivariate convex function  $(u, v) \mapsto u^2/\mathcal{I}(v) = -\mathcal{I}''(v)u^2$  gives

$$[\mathcal{I}(P_t f)]^2 - [P_t(\mathcal{I}(f))]^2 \geq 2 \int_0^t [P_s(|\nabla P_{t-s} f|)]^2 ds.$$

Now by using the gradient bound (6) we have

$$C_1 P_s(|\nabla P_{t-s} f|) \geq |\nabla P_s(P_{t-s} f)| = |\nabla P_t f|$$

and thus

$$[\mathcal{I}(P_t f)]^2 - [P_t(\mathcal{I}(f))]^2 \geq \frac{2t}{C_1^2} |\nabla P_t f|^2.$$

In particular, we obtain the following uniform gradient bound

$$\|\mathcal{I}''(P_t f)|\nabla P_t f\|_\infty = \left\| \frac{|\nabla P_t f|}{\mathcal{I}(P_t f)} \right\|_\infty \leq \frac{C_1}{\sqrt{2t}}.$$

We are now able to prove (30). By the diffusion property

$$\mathcal{I}(P_t f) - P_t(\mathcal{I}(f)) = - \int_0^t \partial_s P_s(\mathcal{I}(P_{t-s} f)) ds = - \int_0^t P_s(\mathcal{I}''(P_{t-s} f) |\nabla P_{t-s} f|^2) ds.$$

By (6) we get  $|\nabla P_{t-s} f|^2 \leq C_1 |\nabla P_{t-s} f| P_{t-s}(|\nabla f|)$  and thus

$$\mathcal{I}(P_t f) - P_t(\mathcal{I}(f)) \leq C_1 \left( \int_0^t \frac{C_1}{\sqrt{2(t-s)}} ds \right) P_t(|\nabla f|) = C_1^2 \sqrt{2t} P_t(|\nabla f|).$$

□

The isoperimetric content of (30) can be extracted by approximating an indicator with a smooth  $f$ , see [4]. Namely, for any Borel set  $A \subset \mathbb{H}$  with smooth boundary, any  $t \geq 0$ , and any  $\mathbf{x} \in \mathbb{H}$ , we get by denoting  $\mu_{t,\mathbf{x}} = P_t(\cdot)(\mathbf{x})$ ,

$$\mathcal{I}(\mu_{t,\mathbf{x}}(A)) \leq C_1^2 \sqrt{2t} \mu_{t,\mathbf{x}}^{\text{surface}}(\partial A). \quad (31)$$

**Corollary 6.5 (Yet another local Bobkov Gaussian isoperimetric inequality).** *With the notations of (6), for every  $t \geq 0$  and  $f \in C_c^\infty(\mathbb{H}, (0, 1))$ ,*

$$\mathcal{I}(P_t f) \leq P_t \left( \sqrt{(\mathcal{I}(f))^2 + 2C^4 t |\nabla f|^2} \right). \quad (32)$$

*Proof.* The desired result follows from the transportation-rearrangement argument given in [5, prop. 5 p. 427], which is inspired from [4, p. 273]. The method is not specific to the heat semigroup on the Heisenberg group. It is based in particular on a similar inequality for the standard Gaussian measure on  $\mathbb{R}$  obtained by Bobkov in [9]. □

One of the most important aspect of (32) is its stability by tensor product, in contrast with (30), while maintaining the same isoperimetric content. Moreover, one may recover from (32) the Gross logarithmic Sobolev inequality (20) by using the fact that  $\mathcal{I}'(u) \sim \sqrt{-2 \log(u)}$  and  $\mathcal{I}(u) \sim u \sqrt{-2 \log(u)}$  at  $u = 0$ . We ignore if (32) can be obtained directly by semigroup interpolation, as for the elliptic case in [4]. The proof given in [4] for the elliptic case is based directly on a curvature bound at the level of the infinitesimal generator, which is not implied by the gradient bound (6) on  $\mathbb{H}$ . We ignore also if one can adapt on the Heisenberg group the two points space approach used in [9] or the martingale representation approach used in [5, 10, 23, 27]. There is a lack of a direct proof of (32) on the Heisenberg group, despite the fact that (32) and (30) are equivalent, according to the argument of Barthe and Maurey in [5, prop. 5 p. 427].

**Remark 6.6 (Abstract Markov settings).** *In fact, up to specific constants, most of the proofs given above have nothing to do with the group structure of the space or with the convolution semigroup nature of  $(P_t)_{t \geq 0}$ . They remain actually valid in very general settings provided that the computations make sense. The key points are a  $\sqrt{\Gamma} - P_t$  sub-commutation and the semigroup and diffusion properties. Formally, let  $L$  be a diffusion operator on a smooth complete connected differential manifold  $\mathcal{M}$ , generating a Markov semigroup  $(P_t)_{t \geq 0} = (e^{tL})_{t \geq 0}$  with smooth density with respect to some reference Borel measure on  $\mathcal{M}$ . Let  $2\Gamma f = L(f^2) - 2fLf$  and suppose that there exists  $C : (0, \infty) \rightarrow (0, \infty)$  such that*

$$\sqrt{\Gamma P_t f} \leq C(t) P_t(\sqrt{\Gamma f}) \quad (33)$$

pointwise for every  $t \geq 0$  and every smooth  $f : \mathcal{M} \rightarrow \mathbb{R}$ . Let us define  $R(t)$  by

$$R(t) = \int_0^t C(s) \left( \int_0^s \frac{2}{C(u)^2} du \right)^{-\frac{1}{2}} ds.$$

Then for every  $t \geq 0$ , every  $x \in \mathcal{M}$ , and every smooth  $f : \mathcal{M} \rightarrow \mathbb{R}$ ,

$$P_t(|f - P_t(f)(x)|)(x) \leq 2R(t) P_t(\sqrt{\Gamma f})(x). \quad (34)$$

Moreover, for every  $t \geq 0$  and every smooth  $f : \mathcal{M} \rightarrow (0, 1)$ ,

$$\mathcal{I}(P_t(f)) - P_t(\mathcal{I}(f)) \leq R(t) P_t(\sqrt{\Gamma f}), \quad (35)$$

and

$$\mathcal{I}(P_t(f)) \leq P_t \left( \sqrt{(\mathcal{I}(f))^2 + R(t)^2 \Gamma f} \right), \quad (36)$$

where  $\mathcal{I}$  stands for the Gaussian isoperimetric function as in (29). Furthermore, if  $I$  is an open interval of  $\mathbb{R}$  and  $\varphi : I \rightarrow \mathbb{R}$  is a smooth convex function such that  $\varphi'' > 0$  on  $I$  and  $-1/\varphi''$  is convex on  $I$ , then for every  $t \geq 0$ , every  $x \in \mathcal{M}$ , and every smooth  $f : \mathcal{M} \rightarrow I$ ,

$$P_t(\varphi(f)) - \varphi(P_t f) \leq \left( \int_0^t C(u)^2 du \right) P_t(\varphi''(f) \Gamma f). \quad (37)$$

Finally, if  $P_t(\cdot)(x) \rightarrow \mu$  weakly as  $t \rightarrow \infty$  for some  $x \in \mathcal{M}$  and some probability measure  $\mu$  on  $\mathcal{M}$  then the four inequalities (34-37) above hold for  $\mu$  instead of  $P_t(\cdot)(x)$ . Here the constant in (34) is obtained partly by using a reverse local Poincaré inequality deduced from (33). On the Heisenberg group, we used an alternative constant for the reverse local Poincaré inequality, which was not deduced from (33).

## 6.4 Multi-times inequalities

Let  $\varphi : \mathcal{I} \rightarrow \mathbb{R}$  be fixed and as in (19). The  $\varphi$ -entropy functional

$$\mathbf{Ent}_\mu : f \mapsto \mathbf{Ent}_\mu(f) = \mathbb{E}_\mu(\varphi(f)) - \varphi(\mathbb{E}(f))$$

has the tensor product property. Namely, if  $\mu = \mu_1 \otimes \dots \otimes \mu_n$  is a probability measure on a product space  $E = E_1 \times \dots \times E_n$  then for every  $f : E \rightarrow \mathcal{I}$  in the domain of  $\mathbf{Ent}_\mu$ ,

$$\mathbf{Ent}_\mu(f) \leq \sum_{i=1}^n \mathbb{E}_\mu(\mathbf{Ent}_{\mu_i}(f))$$

where the expectations in  $\mathbf{Ent}_{\mu_i}(f)$  act only on the  $i^{\text{th}}$  coordinate. The details are given in [11]. Below, we use the notation  $\mathbf{Ent}(U) = \mathbb{E}(\varphi(U)) - \varphi(\mathbb{E}(U))$  for any real random variable  $U$  taking its values in  $\mathcal{I}$ . Now, let  $(X_t)_{t \geq 0}$  be the diffusion process on  $\mathbb{H}$  generated by  $L$ , with  $X_0 = 0$ . Let also  $F : \mathbb{H}^n \rightarrow \mathcal{I}$  be some fixed smooth function. Here  $\mathbb{H}^n$  stands for the  $n$ -product space  $\mathbb{H} \times \cdots \times \mathbb{H}$ . Since  $(X_t)_{t \geq 0}$  has independent stationary increments, i.e. is a Lévy process on  $\mathbb{H}$  associated to a convolution semigroup, we have, for any finite increasing sequence  $0 < t_1 < \cdots < t_n$  of fixed times,

$$\mathbf{Ent}(F(X_{t_1}, \dots, X_{t_n})) = \mathbf{Ent}_{\mathcal{L}(Q_1, \dots, Q_n)}(F \circ \pi)$$

where  $\pi : \mathbb{H}^n \rightarrow \mathbb{H}^n$  is defined by

$$\pi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n) = (\mathbf{x}_1, \mathbf{x}_1 \bullet \mathbf{x}_2, \dots, \mathbf{x}_1 \bullet \cdots \bullet \mathbf{x}_n)$$

for every  $(\mathbf{x}_1, \dots, \mathbf{x}_n) \in \mathbb{H}^n$ , and where  $Q_1, \dots, Q_n$  are independent random variables on the Heisenberg group  $\mathbb{H}$  with  $\mathcal{L}(Q_i) = \mathcal{L}((X_{t_{i-1}})^{-1} X_{t_i}) = \mathcal{L}(X_{t_i - t_{i-1}})$  for every  $i \in \{1, \dots, n\}$ , with  $t_0 = 0$ . The tensor product property of the entropy given above together with (19) gives

$$\mathbf{Ent}(F(X_{t_1}, \dots, X_{t_n})) \leq C_1^2 \mathbb{E}_{\mathcal{L}(X_{t_1}, \dots, X_{t_n})}(\varphi''(F) \mathcal{D}_{t_1, \dots, t_n}^2 F)$$

where  $C$  is as in (6) and (19), and where

$$\mathcal{D}_{t_1, \dots, t_n}^2 F = \sum_{i=1}^n (t_i - t_{i-1}) |\nabla_i(F \circ \pi)|^2 \circ \pi^{-1}$$

where  $\nabla_i$  denote the left invariant gradient  $\nabla$  of  $\mathbb{H}$  acting on the  $i^{\text{th}}$  coordinate  $\mathbf{x}_i$ . Only the distribution of  $\varphi''(F) \mathcal{D}_{t_1, \dots, t_n}^2 F$  under  $\mathcal{L}(X_{t_1}, \dots, X_{t_n})$  is of interest. Similarly, by using an argument of Bobkov detailed for instance in [5, p. 429-430], we get from (32), for any smooth function  $F : \mathbb{H}^n \rightarrow (0, 1)$ , by denoting  $\nu = \mathcal{L}(X_{t_1}, \dots, X_{t_n})$ ,

$$\mathcal{I}(\mathbb{E}_\nu(F)) \leq \mathbb{E}_\nu \left( \sqrt{(\mathcal{I}(F))^2 + 2C_2^4 \mathcal{D}_{t_1, \dots, t_n}^2 F} \right).$$

We ignore if such a cylindrical approach leads to functional inequalities for the paths space on  $\mathbb{H}$ , i.e. for the hypoelliptic Wiener measure, by letting  $n \rightarrow \infty$ .

## References

- [1] C. ANÉ, S. BLACHÈRE, D. CHAFAÏ, P. FOGÈRES, I. GENTIL, F. MALRIEU, C. ROBERTO, G. SCHEFFER, *Sur les inégalités de Sobolev logarithmiques*. Panoramas et Synthèses, 10. Société Mathématique de France, Paris, 2000. xvi+217 pp.
- [2] D. BAKRY, *On Sobolev and logarithmic Sobolev inequalities for Markov semigroups*. Taniguchi symposium. New trends in stochastic analysis (Charingworth, 1994), 43–75, World Sci. Publ., River Edge, NJ, **1997**.
- [3] D. BAKRY, *L'hypercontractivité et son utilisation en théorie des semigroupes*. Lectures on probability theory (Saint-Flour, **1992**), 1–114, Lecture Notes in Math., 1581, Springer, Berlin, **1994**.
- [4] D. BAKRY and M. LEDOUX, *Lévy-Gromov's isoperimetric inequality for an infinite-dimensional diffusion generator*. Invent. Math. 123 (**1996**), no. 2, 259–281.
- [5] F. BARTHE and B. MAUREY, *Some remarks on isoperimetry of Gaussian type*. Ann. Inst. H. Poincaré Probab. Statist. 36 (**2000**), no. 4, 419–434.
- [6] F. BAUDOIN, *An introduction to the geometry of stochastic flows*. Imperial College Press, London, **2004**. x+140 pp.
- [7] W. BECKNER, *A generalized Poincaré inequality for Gaussian measures*. Proc. Amer. Math. Soc. 105 (**1989**), no. 2, 397–400.

- [8] S.G. BOBKOV, *A functional form of the isoperimetric inequality for the Gaussian measure*. J. Funct. Anal. (1996) 135 39–49.
- [9] S.G. BOBKOV, *An isoperimetric inequality on the discrete cube, and an elementary proof of the isoperimetric inequality in Gauss space*. Ann. Probab. 25 (1997), no. 1, 206–214.
- [10] M. CAPITAINE and E.P. HSU and M. LEDOUX, *Martingale representation and a simple proof of logarithmic Sobolev inequalities on path spaces*. Electron. Comm. Probab. 2 (1997), 71–81.
- [11] D. CHAFAÏ, *Entropies, convexity, and functional inequalities: on  $\Phi$ -entropies and  $\Phi$ -Sobolev inequalities*. J. Math. Kyoto Univ. 44 (2004), no. 2, 325–363.
- [12] D. CHAFAÏ, *Binomial-Poisson entropic inequalities and the  $M/M/\infty$  queue*. ESAIM Probab. Stat. 10 (2006), 317–339.
- [13] J. CHEEGER, *A lower bound for the smallest eigenvalue of the Laplacian*. Problems in analysis (Papers dedicated to Salomon Bochner, 1969), pp. 195–199. Princeton Univ. Press, Princeton, N. J., 1970.
- [14] Th. COULHON and L. SALOFF-COSTE, *Isopérimétrie pour les groupes et les variétés*. Rev. Mat. Iberoamericana 9 (1993), no. 2, 293–314.
- [15] B.K. DRIVER and T. MELCHER, *Hypoelliptic heat kernel inequalities on the Heisenberg group*. J. Funct. Anal. 221 (2005), no. 2, 340–365.
- [16] P. FRIZ, H. OBERHAUSER, *Isoperimetry and Rough Path Regularity*, preprint [arXiv:0711.0163v1](https://arxiv.org/abs/0711.0163v1) [math.PR].
- [17] B. GAVEAU, *Principe de moindre action, propagation de la chaleur et estimées sous elliptiques sur certains groupes nilpotents*. Acta Math. 139 (1977), no. 1-2, 95–153.
- [18] R. BEALS, B. GAVEAU, P. GREINER, *Hamilton-Jacobi theory and the heat kernel on Heisenberg groups*, J. Math. Pures Appl. 79, 7 (2000) 633–689
- [19] M. GROMOV, *Metric structures for Riemannian and non-Riemannian spaces*. Based on the 1981 French original. With appendices by M. Katz, P. Pansu and S. Semmes. Progress in Mathematics, 152. Birkhuser Boston, Inc., Boston, MA, 1999. xx+585 pp.
- [20] L. GROSS, *Logarithmic Sobolev inequalities*. Amer. J. Math. 97 (1975), no. 4, 1061–1083.
- [21] L. GROSS, *Logarithmic Sobolev inequalities and contractivity properties of semi-groups*. Varenna (1992). Lecture Notes in Math. 1563 (1993) 54–88. Springer, Berlin.
- [22] Y.-Z. HU, *A unified approach to several inequalities for Gaussian and diffusion measures*, Séminaire de Probabilités, XXXIV, Springer, Berlin, 2000, 329–335.
- [23] E.P. HSU, *Stochastic analysis on manifolds*. Graduate Studies in Mathematics, 38. American Mathematical Society, Providence, RI, 2002. xiv+281 pp.
- [24] JUILLET, N. *Geometric inequalities and generalized Ricci bounds on the Heisenberg group*, preprint (2006).
- [25] R. LATAŁA and K. OLESZKIEWICZ, *Between Sobolev and Poincaré*. Geometric aspects of functional analysis, 147–168, Lecture Notes in Math., 1745, Springer, Berlin, 2000.
- [26] M. LEDOUX, *A simple analytic proof of an inequality by P. Buser*. Proc. Amer. Math. Soc. 121 (1994), no. 3, 951–959.
- [27] M. LEDOUX, *The geometry of Markov diffusion generators*. Probability theory. Ann. Fac. Sci. Toulouse Math. (6) 9 (2000), no. 2, 305–366.
- [28] H.-Q. LI, *Estimation optimale du gradient du semi-groupe de la chaleur sur le groupe de Heisenberg*. J. Funct. Anal. 236 (2006), no. 2, 369–394.
- [29] H.-Q. LI, *Estimations asymptotiques du noyau de la chaleur sur les groupes de Heisenberg*, CRAS Ser. I, 497-502, (2007).
- [30] H.-Q. LI, *Estimations optimales du noyau de la chaleur sur les groupes de type Heisenberg*, preprint, personal communication, (2007).
- [31] P. MAHEUX, L. SALOFF-COSTE, *Analyse sur les boules d'un opérateur sous-elliptique*, Math. Ann., 303, (1995), 713–740.
- [32] T. MELCHER, *Hypoelliptic heat kernel inequalities on Lie groups*, PhD thesis, University of California, San Diego (2004).
- [33] R. MONTGOMERY, *A tour of subriemannian geometries, their geodesics and applications*. Mathematical Surveys and Monographs, 91. American Mathematical Society, Providence, RI, 2002. xx+259 pp.
- [34] D. NEUENSCHWANDER, *Probabilities on the Heisenberg group. Limit theorems and Brownian motion*. Lecture Notes in Mathematics, 1630. Springer-Verlag, Berlin, 1996. viii+139 pp.

- [35] M.-K. VON RENESSE and K.-Th. STURM, *Transport inequalities, gradient estimates, entropy, and Ricci curvature*. Comm. Pure Appl. Math. 58 (2005), no. 7, 923–940.
- [36] N. Th. VAROPOULOS, *Small time Gaussian estimates of heat diffusion kernels. I. The semigroup technique*. Bull. Sci. Math. 113 (1989), no. 3, 253–277.
- [37] N. Th. VAROPOULOS and L. SALOFF-COSTE and Th. COULHON, *Analysis and Geometry on Groups*, Cambridge Tracts in Math., vol. 100, Cambridge Univ. Press, Cambridge, 1992.

Dominique BAKRY bakry[[@](mailto:bakry@math.ups-tlse.fr)]math.ups-tlse.fr  
Fabrice BAUDOIN fbaudoin[[@](mailto:fbaudoin@math.ups-tlse.fr)]math.ups-tlse.fr  
Michel BONNEFONT bonnefon[[@](mailto:bonnefon@math.ups-tlse.fr)]math.ups-tlse.fr  
Djalil CHAFAÏ chafai[[@](mailto:chafai@math.ups-tlse.fr)]math.ups-tlse.fr

INSTITUT DE MATHÉMATIQUES DE TOULOUSE (CNRS 5219)  
UNIVERSITÉ PAUL SABATIER  
118 ROUTE DE NARBONNE, F-31062 TOULOUSE, FRANCE.