



HAL
open science

VLTI-AMBER observations of Eta Carinae with high spatial resolution and spectral resolutions of 1,500 and 10,000

Gerd Weigelt, Romain G. Petrov, Olivier Chesneau, Kris Davidson, Armando Domiciano de Souza, Thomas Driebe, Renaud Foy, Didier Fraix-Burnet, Ted Gull, John D. Hillier, et al.

► To cite this version:

Gerd Weigelt, Romain G. Petrov, Olivier Chesneau, Kris Davidson, Armando Domiciano de Souza, et al.. VLTI-AMBER observations of Eta Carinae with high spatial resolution and spectral resolutions of 1,500 and 10,000. SPIE Astronomical Telescopes + Instrumentation, May 2006, Orlando, United States. 10.1117/12.671582 . hal-00199782

HAL Id: hal-00199782

<https://hal.science/hal-00199782>

Submitted on 7 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

VLTI-AMBER observations of Eta Carinae with high spatial resolution and spectral resolutions of 1,500 and 10,000 *

Gerd Weigelt^a, Romain G. Petrov^b, Olivier Chesneau^c, Kris Davidson^d, Armando Domiciano de Souza^{c,b}, Thomas Driebe^a, Renaud Foy^e, Didier Fraix-Burnet^f, Ted Gull^g, John D. Hillier^h, Karl-Heinz Hofmann^a, Stefan Kraus^a, Fabien Malbet^f, Alessandro Marconiⁱ, Philippe Mathias^c, Jean-Louis Monin^f, Florentin Millour^{f,b}, Keiichi Ohnaka^a, Frederik Rantakyro^k, Andrea Richichi^l, Dieter Schertl^a, Markus Schöller^k, Philippe Stee^c, Leonardo Testiⁱ, Markus Wittkowski^k

^aMax-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany; ^bLaboratoire Universitaire d'Astrophysique de Nice, UMR 6525 Université de Nice/CNRS, Parc Valrose, F-06108 Nice cedex 2, France; ^cLaboratoire Gemini, UMR 6203 Observatoire de la Côte d'Azur/CNRS, Avenue Copernic, 06130 Grasse, France; ^dSchool of Physics and Astronomy, University of Minnesota, 116 Church Street SE, Minneapolis, MN 55455, USA; ^eCentre de Recherche Astronomique de Lyon, UMR 5574 Université Claude Bernard/CNRS, 9 avenue Charles André, F-69561 Saint Genis Laval cedex, France; ^fLaboratoire d'Astrophysique de Grenoble, UMR 5571 Université Joseph Fourier/CNRS, BP 53, F-38041 Grenoble Cedex 9, France; ^gLaboratory for Extraterrestrial Planets and Stellar Astrophysics, Goddard Space Flight Center, 20771 Greenbelt, Maryland, USA; ^hDepartment of Physics and Astronomy, University of Pittsburgh, 3941 O'Hara Street, Pittsburgh, PA 15260, USA; ⁱINAF-Osservatorio Astrofisico di Arcetri, Istituto Nazionale di Astrofisica, Largo E. Fermi 5, I-50125 Firenze, Italy; ^kEuropean Southern Observatory, Casilla 19001, Santiago 19, Chile; ^lEuropean Southern Observatory, Karl Schwarzschild Strasse 2, D-85748 Garching, Germany;

ABSTRACT

We present the first interferometric NIR observations of the LBV η Carinae with high spectral resolution. The observations were carried out with three 8.2 m VLTI Unit Telescopes in the K -band. The raw data are spectrally dispersed interferograms obtained with spectral resolutions of 1,500 (MR-K mode) and 12,000 (HR-K mode). The observations were performed in the wavelength range around both the He I 2.059 μm and the Br γ 2.166 μm emission lines. The spectrally dispersed AMBER interferograms allow the investigation of the *wavelength dependence* of the visibility, differential phase, and closure phase of η Car. In the K -band continuum, a diameter of 4.0 ± 0.2 mas (Gaussian FWHM) was measured for η Car's optically thick wind region, whereas the Br γ and He I emission line regions are larger. If we fit Hillier et al.¹ model visibilities to the observed AMBER visibilities, we obtain 50% encircled-energy diameters of 4.3, 6.5 and 9.6 mas in the 2.17 μm continuum, the He I, and the Br γ emission lines, respectively. In the continuum near the Br γ line, an elongation along a position angle of $128^\circ \pm 15^\circ$ was found, consistent with previous VLTI/VINCI measurements.² We find good agreement between the measured visibilities and the predictions of the radiative transfer model of Hillier et al.¹ For the interpretation of the non-zero differential and closure phases measured within the Br γ line, we present a simple geometric model of an inclined, latitude-dependent wind zone. Our observations support theoretical models of anisotropic winds from fast-rotating, luminous hot stars with enhanced high-velocity mass loss near the polar regions.

Keywords: Stars: individual: η Carinae – Stars: mass-loss, emission-line, circumstellar matter, winds, outflows – Infrared: stars – Techniques: interferometric, high angular resolution, spectroscopic

* Based on observations obtained with the VLTI interferometer of the European Southern Observatory (ESO), Chile
Send offprint requests to: G. Weigelt; email: weigelt@mpifr-bonn.mpg.de

1. INTRODUCTION

η Car is one of the most luminous ($L \sim 4 \times 10^6 L_\odot$) and most massive ($M \sim 100 M_\odot$) unstable Luminous Blue Variables suffering from an extremely high mass loss at a rate of $\sim 10^{-3} M_\odot \text{yr}^{-1}$.³ It is surrounded by the expanding bipolar Homunculus nebula ejected during the Great Eruption in 1843. The inclination of the polar axis of the Homunculus nebula with the line-of-sight is $\sim 41^\circ$ with the southern pole pointing towards us.⁴ The first measurements of structures in the innermost sub-arcsecond region of the Homunculus were obtained by speckle-interferometric observations.^{5,6} These observations revealed a central object (component A) plus three compact and unusually bright objects (components B, C, and D) at distances ranging from approximately $0.1''$ to $0.2''$. Recent observations of η Car by Chesneau et al.⁷ using NACO and VLTI/MIDI revealed a butterfly-shaped dust environment at 3.74 and $4.05 \mu\text{m}$ and resolved the dusty emission from the individual speckle objects with unprecedented angular resolution in the NIR.

Spectroscopic studies of the Homunculus nebula^{8,9} showed that the wind of η Car is latitude-dependent. Van Boeckel et al.² resolved the optically thick, aspheric wind region with NIR interferometry using the VLTI/VINCI instrument. They measured a size of 5 mas (50% encircled energy diameter), an axis ratio of 1.25 ± 0.05 , and a position angle (PA) of the major axis of $134^\circ \pm 7^\circ$, and derived a mass-loss rate of $1.6 \times 10^{-3} M_\odot/\text{yr}^{-1}$. This aspheric wind can be explained by models for line-driven winds from luminous hot stars rotating near their critical speed.^{10,11} The models predict a higher wind speed and density along the polar axis than in the equatorial plane. Van Boeckel et al.² showed that the K broad-band observations obtained with VINCI are in agreement with the predictions from the detailed spectroscopic model by Hillier et al.¹ The Hillier et al. model was developed to explain STIS HST spectra.

A variety of observations suggest that the central source of η Car is a binary. Daminelli et al.¹² first noticed the 5.5-year periodicity in the spectroscopic changes of this object, which has been confirmed by several other studies since then.^{13–17} On the other hand, to date, the binary nature of the central object in η Car is still a matter of debate.^{18–21} HST STIS observations show that the He I lines are strongly variable and blue-shifted throughout most of the 5.54 year variability period. These observations cannot be explained in the context of a spherical wind model. It now appears likely that the He I emission lines originate in the bow shock and ionization zone, associated with the wind-wind interaction zone in a binary system.^{22,23} Very recent FUSE observations of η Car seem to have revealed a hot star component.²⁴

2. AMBER OBSERVATIONS AND DATA PROCESSING

AMBER^{26–28} is the near-infrared (J , H , K band) beam-combiner instrument of ESO's *Very Large Telescope Interferometer*. AMBER allows the measurement of visibilities, differential visibilities, differential phases, and closure phases.²⁹ AMBER offers three spectroscopic modes: low (LR mode; $R=\lambda/\Delta\lambda=75$), medium (MR mode; $R=1,500$), and high (HR mode; $R=12,000$) spectral resolutions. The fibers in AMBER limit the field-of-view to the diameter of the fibers on the sky ($\sim 60 \text{ mas}$). In AMBER the light is spectrally dispersed using a prism or grating. The AMBER detector is a Hawaii array detector with 512×512 pixels.

Our AMBER η Car observations were carried out on 2004 December 26, 2005 February 25, and 2005 February 26 with the three 8.2m *Unit Telescopes* UT2, UT3, and UT4.²⁵ Figure 2 (left) shows two AMBER raw interferograms taken in the wavelength range around the $\text{Br}\gamma$ line in HR (top) and MR (bottom) mode. In the MR data sets, the Doppler-broadened $\text{Br}\gamma$ line covers ~ 8 spectral channels, whereas in HR mode, the line is resolved by ~ 50 spectral channels. With projected baseline lengths up to 89 m , an angular resolution of $\sim 5 \text{ mas}$ was achieved in the K band. The MR-K observations were performed in the wavelength range around both the He I $2.059 \mu\text{m}$ and the $\text{Br}\gamma$ $2.166 \mu\text{m}$ emission lines. The HR-K observations were only performed in a wavelength range around the $\text{Br}\gamma$ line. The widths of the wavelength windows of the obtained MR-K and HR-K observations are approximately $0.05 \mu\text{m}$ and $0.02 \mu\text{m}$, respectively.

For the reduction of the AMBER data, we used version 2.4 of the *amdlib*[†] software package. This software uses the P2VM (*pixel-to-visibilities matrix*) algorithm³⁰ in order to extract complex visibilities for each baseline

[†]This software package is available from
<http://amber.obs.ujf-grenoble.fr>

and each spectral channel of an AMBER interferogram. From these three complex visibilities, the amplitude and the closure phase are derived.

3. RESULTS

Our AMBER observations of η Car were performed with the spectral resolutions of $R = 1,500$ and $R = 12,000$ in Dec. 2004 ($\phi = 0.268$) and Feb. 2005 ($\phi = 0.299$) in spectral windows around the He I and Br γ emission lines at $\lambda = 2.059$ and $2.166 \mu\text{m}$, respectively. From the measurements, we obtained spectra, visibilities, differential visibilities, differential phases, and closure phases. From the analysis of the data, we derived the following results:²⁵

In the K -band continuum, we resolved η Car's optically thick wind. From a Gaussian fit of the K -band continuum visibilities in the projected baseline range from 28–89 m, we obtained a FWHM diameter of 4.0 ± 0.2 mas.

When comparing the AMBER *continuum* visibilities with the NLTE radiative transfer model from Hillier et al.,¹ we find very good agreement between the model and observations. The best fit was obtained with a slightly rescaled version of the original Hillier et al.¹ model, corresponding to an observed FWHM diameter of 2.4 mas and a 50% encircled-energy diameter of 4.3 mas at $\lambda = 2.17 \mu\text{m}$. Taking the different FOVs into account, we found good agreement between the AMBER measurements and previous VLTI/VINCI observations of η Car presented by van Boeckel et al.²

If we fit Hillier et al.¹ model visibilities to the observed AMBER emission line visibilities, we obtain 50% encircled-energy diameters of 6.5 and 9.6 mas in He I and the Br γ emission lines, respectively.

In the continuum around the Br γ line, we found an elongation towards position angle $128^\circ \pm 15^\circ$ with a projected axis ratio of 1.21 ± 0.08 . This result confirms the earlier finding of van Boeckel et al.² using VLTI/VINCI and supports theoretical studies which predict an enhanced mass loss in polar direction for massive stars rotating close to their critical rotation rate.¹¹

For both the Br γ and the He I emission lines, we measured non-zero differential phases and non-zero closure phases within the emission lines, indicating a complex, asymmetric object structure. We developed a physically motivated model,²⁵ which shows that the asymmetries (DPs and CPs) measured within the wings of the Br γ line are consistent with the geometry expected for an aspherical, latitude-dependent stellar wind. Additional VLTI/AMBER measurements and radiative transfer modeling will be required to determine the precise parameters of such an inclined aspherical wind.

ACKNOWLEDGMENTS

This work is based on observations made with the European Southern Observatory telescopes. This research has also made use of the ASPRO observation preparation tool from the *Jean-Marie Mariotti Center* in France, the SIMBAD database at CDS, Strasbourg (France) and the Smithsonian/NASA Astrophysics Data System (ADS).

The data reduction software `amdlib` is freely available on the AMBER site <http://amber.obs.ujf-grenoble.fr>. It has been linked with the free software Yorick[‡] to provide the user-friendly interface `ammyorick`.

The NSO/Kitt Peak FTS data used here to identify the telluric lines in the AMBER data were produced by NSF/NOAO.

This project has benefitted from funding from the French Centre National de la Recherche Scientifique (CNRS) through the Institut National des Sciences de l'Univers (INSU) and its Programmes Nationaux (ASHRA, PNPS). The authors from the French laboratories would like to thank the successive directors of the INSU/CNRS directors. S. Kraus was supported for this research through a fellowship from the International Max Planck Research School (IMPRS) for Radio and Infrared Astronomy at the University of Bonn. C. Gil's work was partially supported by the Fundação para a Ciência e a Tecnologia through project POCTI/CTE-AST/55691/2004 from POCTI, with funds from the European program FEDER. K. Weis is supported by the state of North-Rhine-Westphalia (Lise-Meitner fellowship).

[‡]<ftp://ftp-icf.llnl.gov/pub/Yorick>




Figure 1. Left: Spectrally dispersed VLTI/AMBER Michelson interferograms of η Car. The two panels show the spectrally dispersed fringe signal (IF) as well as the photometric calibration signals from the three telescopes (P1-P3) in high (HR, upper panel) and medium spectral resolution mode (MR, lower panel). In both panels, the bright regions are the Doppler-broadened $\text{Br}\gamma$ emission line. Right: AMBER observables derived from our η Car data around the $\text{Br}\gamma$ line (MR mode, 2004 December 26; the HR mode and He I observations are described in Weigelt et al.²⁵). The top panel shows the spectrum as extracted from the interferometric channels, followed by the derived calibrated visibilities and the differential visibilities. In the two panels at the bottom, the differential phase and the closure phase are presented. The vertical grey line marks the rest-wavelength of $\text{Br}\gamma$ ($\lambda_{\text{vac}} = 2.1661 \mu\text{m}$). The left error bars correspond to the total error estimated for the continuum wavelength range, and the error bars towards the right visualizes the total error for the wavelength range within the line (see Weigelt et al.²⁵ for more details).

REFERENCES

1. D. J. Hillier, K. Davidson, K. Ishibashi, and T. Gull, “On the Nature of the Central Source in η Carinae,” *ApJ* **553**, pp. 837–860, June 2001.
2. R. van Boekel, P. Kervella, M. Schöller, T. Herbst, W. Brandner, A. de Koter, L. B. F. M. Waters, D. J. Hillier, F. Paresce, R. Lenzen, and A.-M. Lagrange, “Direct measurement of the size and shape of the present-day stellar wind of eta Carinae,” *A&A* **410**, pp. L37–L40, Oct. 2003.
3. K. Davidson and R. M. Humphreys, “Eta Carinae and Its Environment,” *A&A* **35**, pp. 1–32, 1997.
4. K. Davidson, N. Smith, T. R. Gull, K. Ishibashi, and D. J. Hillier, “The Shape and Orientation of the Homunculus Nebula Based on Spectroscopic Velocities,” *AJ* **121**, pp. 1569–1577, Mar. 2001.
5. G. Weigelt and J. Ebersberger, “Eta Carinae resolved by speckle interferometry,” *A&A* **163**, p. L5, July 1986.
6. K.-H. Hofmann and G. Weigelt, “Speckle masking observation of Eta Carinae,” *A&A* **203**, p. L21, Sept. 1988.
7. O. Chesneau, M. Min, T. Herbst, L. B. F. M. Waters, D. J. Hillier, C. Leinert, A. de Koter, I. Pascucci, W. Jaffe, R. Köhler, C. Alvarez, R. van Boekel, W. Brandner, U. Graser, A. M. Lagrange, R. Lenzen, S. Morel, and M. Schöller, “The sub-arcsecond dusty environment of Eta Carinae,” *A&A* **435**, pp. 1043–1061, June 2005.
8. N. Smith, “Dissecting the Homunculus nebula around Eta Carinae with spatially resolved near-infrared spectroscopy,” *MNRAS* **337**, pp. 1252–1268, Dec. 2002.
9. N. Smith, K. Davidson, T. R. Gull, K. Ishibashi, and D. J. Hillier, “Latitude-dependent Effects in the Stellar Wind of η Carinae,” *ApJ* **586**, pp. 432–450, Mar. 2003.
10. S. P. Owocki, S. R. Cranmer, and K. G. Gayley, “Inhibition FO Wind Compressed Disk Formation by Nonradial Line-Forces in Rotating Hot-Star Winds,” *ApJ* **472**, p. L115, Dec. 1996.
11. V. V. Dwarkadas and S. P. Owocki, “Radiatively Driven Winds and the Shaping of Bipolar Luminous Blue Variable Nebulae,” *ApJ* **581**, pp. 1337–1343, Dec. 2002.
12. A. Damineli, “The 5.52 Year Cycle of Eta Carinae,” *ApJ* **460**, p. L49, Mar. 1996.
13. A. Damineli, A. Kaufer, B. Wolf, O. Stahl, D. F. Lopes, and F. X. de Araújo, “ η Carinae: Binarity Confirmed,” *ApJ* **528**, pp. L101–L104, Jan. 2000.
14. K. Davidson, K. Ishibashi, T. R. Gull, R. M. Humphreys, and N. Smith, “ η Carinae: Testing a Binary Orbit Model with the Hubble Space Telescope/Space Telescope Imaging Spectrograph,” *ApJ* **530**, pp. L107–L110, Feb. 2000.
15. K. Davidson, J. Martin, R. M. Humphreys, K. Ishibashi, T. R. Gull, O. Stahl, K. Weis, D. J. Hillier, A. Damineli, M. Corcoran, and F. Hamann, “A Change in the Physical State of η Carinae?,” *AJ* **129**, pp. 900–906, Feb. 2005.
16. P. A. Whitelock, M. W. Feast, F. Marang, and E. Breedt, “The 2003 shell event in η Carinae,” *MNRAS* **352**, pp. 447–456, Aug. 2004.
17. K. Weis, O. Stahl, D. J. Bomans, K. Davidson, T. R. Gull, and R. M. Humphreys, “VLT UVES Observations of the Balmer Line Variations of η Carinae during the 2003 Spectroscopic Event,” *AJ* **129**, pp. 1694–1699, Mar. 2005.
18. K. Ishibashi, M. F. Corcoran, K. Davidson, J. H. Swank, R. Petre, S. A. Drake, A. Damineli, and S. White, “Recurrent X-Ray Emission Variations of η Carinae and the Binary Hypothesis,” *ApJ* **524**, pp. 983–987, Oct. 1999.
19. R. A. Duncan, S. M. White, J. E. Reynolds, and J. Lim, “Continuum and Spectral Observations of η Carinae at Wavelengths of 3 – 6 Centimeters,” in *ASP Conf. Ser. 179: Eta Carinae at The Millennium*, J. A. Morse, R. M. Humphreys, and A. Damineli, eds., p. 54, 1999.
20. M. Feast, P. Whitelock, and F. Marang, “Variability of η Carinae - III,” *MNRAS* **322**, pp. 741–748, Apr. 2001.
21. J. M. Pittard and M. F. Corcoran, “In hot pursuit of the hidden companion of eta Carinae: An X-ray determination of the wind parameters,” *A&A* **383**, pp. 636–647, Feb. 2002.
22. D. J. Hillier, T. Gull, K. Nielsen, G. Sonneborn, R. Iping, N. Smith, M. Corcoran, A. Damineli, F. W. Hamann, J. C. Martin, and K. Weis, “The UV Scattering Halo of the Central Source Associated with Eta Carinae,” *ApJ* **642**, p. 1098, May 2006.

23. K. E. Nielsen, M. F. Corcoran, T. R. Gull, D. J. Hillier, S. Ivarsson, K. Hamaguchi, and D. J. Lindler, “Eta Carinae across the 2003.5 minimum: Spectroscopic evidence for massive binary interaction,” *ApJ submitted*, 2006.
24. R. C. Iping, G. Sonneborn, T. R. Gull, D. L. Massa, and D. J. Hillier, “Detection of a Hot Binary Companion of η Carinae,” *ApJ* **633**, pp. L37–L40, Nov. 2005.
25. G. Weigelt and et al., “Near-Infrared interferometry of η Carinae with high spatial and spectral resolution using the VLTI and the AMBER instrument,” *A&A submitted*, 2006.
26. R. G. Petrov, F. Malbet, G. Weigelt, F. Lisi, P. Puget, P. Antonelli, U. Beckmann, S. Lagarde, E. Lecoarer, S. Robbe-Dubois, G. Duvert, S. Gennari, A. Chelli, M. Dugue, K. Rousselet-Perraut, M. Vannier, and D. Mourard, “Using the near infrared VLTI instrument AMBER,” in *Interferometry for Optical Astronomy II. Edited by Wesley A. Traub . Proceedings of the SPIE, Volume 4838, pp. 924-933 (2003).*, W. A. Traub, ed., pp. 924–933, Feb. 2003.
27. R. Petrov, F. Millour, O. Chesneau, G. Weigelt, D. Bonneau, P. Stee, S. Kraus, D. Mourard, and et al. , “First AMBER/VLTI observations of hot massive stars,” in *The power of optical/IR interferometry: recent scientific results and 2nd generation VLTI instrumentation*, A. Richichi, ed., 2006.
28. R. G. Petrov, F. Malbet, P. Antonelli, U. Beckmann, Y. Bresson, A. Chelli, D. M., G. Duvert, G. L., P. Kern, S. Lagarde, and et al., “AMBER, the near-infrared spectro-interferometric three-beam combiner of the VLTI,” *A&A submitted*, 2006.
29. F. Millour, M. Vannier, R. G. Petrov, O. Chesneau, L. Dessart, and P. Stee, “Differential interferometry with the AMBER/VLTI instrument: description, performances and illustration.,” in *EAS Publications Series*, **in press**, 2006.
30. E. Tatulli, F. Millour, A. Chelli, *et al.*, “Interferometric data reduction with amber/vlti. principle, estimators and illustration,” *A&A accepted*, 2006.