



HAL
open science

Conjugate and cut loci of a two-sphere of revolution with application to optimal control

Bernard Bonnard, Jean-Baptiste Caillau, Robert Sinclair, Minoru Tanaka

► **To cite this version:**

Bernard Bonnard, Jean-Baptiste Caillau, Robert Sinclair, Minoru Tanaka. Conjugate and cut loci of a two-sphere of revolution with application to optimal control. 2008. hal-00212075v1

HAL Id: hal-00212075

<https://hal.science/hal-00212075v1>

Preprint submitted on 22 Jan 2008 (v1), last revised 22 Feb 2008 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Conjugate and cut loci of a two-sphere of revolution with application to optimal control

Lieu conjugué et lieu de coupure d'une 2-sphère de révolution et application en contrôle optimal

Bernard Bonnard^a, Jean-Baptiste Caillau^b, Robert Sinclair^c, and Minoru Tanaka^d

^a*Institut de mathématiques de Bourgogne (UMR CNRS 5584), 9 avenue Savary, F-21078 Dijon*

^b*ENSEEIH-IRIT (UMR CNRS 5505), 2 rue Camichel, F-31071 Toulouse*

^c*Mathematical Biology Unit, Okinawa Institute of Science and Technology, Okinawa Industrial Technology Center Annex, 12-2 Suzaki, Uruma, Okinawa J-904-2234*

^d*Department of Mathematics, Tokai University, Hiratsuka City, Kanagawa Pref., J-259-1292*

Abstract

The objective of this article is to present a sharp result to determine when the cut locus for a class of metrics on a two-sphere of revolution is reduced to a single branch. This work is motivated by optimal control problems in space and quantum dynamics and give global optimal results in orbital transfer and for Redfield equations in quantum control.

Résumé

Le but de cet article est de présenter une condition suffisante permettant de garantir que le lieu de coupure d'une classe de métriques sur la 2-sphère de révolution est réduit à une branche simple. Ce travail est motivé par des problèmes de contrôle optimal en mécanique spatiale et mécanique quantique. Des résultats globaux d'optimalité sont obtenus en transfert orbital ainsi que dans le cas des équations de Redfield en contrôle quantique.

MSC: 53C20; 53C21; 49K15; 70Q05

Keywords: conjugate and cut loci; 2-spheres of revolution; optimal control; space and quantum mechanics

1. Introduction

The purpose of this article is to improve recent advanced results concerning the structure of the conjugate and cut loci on a two-surface of revolution [21,22] to analyse optimal control problems for both space and quantum control

Email addresses: Bernard.Bonnard@u-bourgogne.fr (Bernard Bonnard), caillau@n7.fr (Jean-Baptiste Caillau), sinclair@oist.jp (Robert Sinclair), m-tanaka@sm.u-tokai.ac.jp (Minoru Tanaka).

dynamics.

The determination of the cut and conjugate loci on a complete two-surface of revolution is a standard and difficult problem in Riemannian geometry. For a real analytic two-sphere, the cut locus at each point is a finite tree, whose extremities are conjugate points. This was stated by Poincaré [16] and proved by Myers [15].

The structure theorem of the cut locus of a point on a 2-dimensional Riemannian manifold was established by Hebda ([12]) and generalized by Shiohama and Tanaka ([18]) for the cut locus of a compact subset in an Alexandrov surface. By the structure theorem (see [18, Theorem A]), the cut locus is a local tree and is a union of countably many rectifiable Jordan arcs and the endpoints.

Still, the precise computations is a complicated problem and the complexity is to estimate the ramifying branches. Besides a construction due to Gluck and Singer [10] proves that there exists a smooth strictly convex surface of revolution, homeomorphic to \mathbb{S}^2 , whose cut locus is not stratifiable.

Even on an ellipsoid of revolution, the computation is not a standard exercise (in [3], the foreseen conjugate and cut loci are given as a conjecture). On an oblate ellipsoid the cut locus of a point different from the pole is a subarc of the antipodal parallel. For a prolate ellipsoid, the same holds replacing parallel by opposite meridian. In the first case the Gauss curvature is monotone increasing from the north pole to the equator and decreasing in the second case.

This result is a consequence of a general result in [22] : given a smooth metric on \mathbb{S}^2 of the form $dr^2 + m^2(r)d\theta^2$, where r is the angle along the meridian and θ the angle of revolution. Assume the following :

1. $m(\pi - r) = m(r)$ (reflective symmetry with respect to the equator)
2. The Gauss curvature is monotone non decreasing (resp. increasing) along a meridian from the north pole to the equator.

Then the cut locus of a point different from the pole is a simple branch located on the antipodal parallel (resp. opposite meridian).

This gives a nice computable criterion to decide whenever the cut locus is reduced to a simple branch. In parallel in recent research projects on geometric optimal control in orbital transfer or quantum control the optimality analysis can be reduced to optimal control problem on two-sphere of revolution for which the generalization of the previous result is crucial in several direction : first of all the monotonicity of the Gauss curvature is not satisfied, secondly the metric can have singularities. The key step is to relate the simple structure of the cut and conjugate loci to a tame property of the extremal flow.

The organization of this article is the following. In section 2, we present the systems from space and quantum dynamics motivating the analysis. In section 3, we give the sharp optimality result needed for the analysis in the Riemannian case, with application to our examples in section 4. The analysis is extended in section 5 to deal with almost Riemannian metrics on two-spheres of revolution encountered in our systems analysis and in parallel in extensions of Gauss-Bonnet theorem [1]. In a concluding section we discuss in details the contributions of this article as well as possible extensions.

It is also worth pointing that our analysis is related to homotopy methods in optimal control, deforming the round sphere \mathbb{S}^2 and keeping simple conjugate and cut loci to be compared to the opposite construction of [10] to generate complex such loci.

2. Motivating examples

2.1. Orbital transfer

The two-input coplanar transfer system [6] is modelled by a 2π -periodic system on a three-dimensional manifold M , of the form :

$$\frac{dq}{dl} = u_1 F_1(q, l) + u_2 F_2(q, l)$$

where the state q represents the geometric coordinates of the osculating ellipse, e.g. $q = (n, e, \theta)$, where n is the *mean motion*, e the *eccentricity* and θ the *argument of the pericenter*. The angular variable is the *longitude* $l \in \mathbb{S}^1$, while the trajectories are parameterized by the *cumulated* longitude $l \in \mathbb{R}$. When considering the energy minimization problem, the maximum principle tells us that the minimizers have to be selected among extremal curves solutions of the Hamiltonian

$$H(q, p, l) = \frac{1}{2}(H_1^2 + H_2^2)(q, p, l)$$

where the H_i 's are the Hamiltonian lifts $H_i(q, p, l) = \langle p, F_i(q, l) \rangle$, $i = 1, 2$. If low thrust is applied, we consider the long time behavior of the system which is approximated by the system with the *averaged* Hamiltonian

$$\bar{H}(q, p) = \frac{1}{2\pi} \int_0^{2\pi} H(q, p, l) dl.$$

Averaging generates Lie brackets of the initial vector fields, so that the averaged Hamiltonian turns to be a full rank quadratic form in the adjoint variable p , thus associated to a Riemannian metric. The computed expression is

$$\bar{g} = \frac{dn^2}{9n^{1/3}} + \frac{2n^{5/3}}{5(1-e^2)} de^2 + \frac{2n^{5/3}}{(5-4e^2)} e^2 d\theta^2.$$

Such a metric can be normalized with $n = (5\rho/2)^{6/5}$, $e = \sin r$ so that :

$$\bar{g} = d\rho^2 + (\rho^2/c^2)g$$

where $c = \sqrt{2/5}$ and $g = dr^2 + m^2(r)d\theta^2$ with

$$m^2(r) = \frac{\sin^2 r}{1 - (4/5)\sin^2 r}.$$

By homogeneity, we can restrict our optimality analysis to the Riemannian metric g with $r \in [0, \pi/2]$, where the pole $e = 0$ corresponds to circular orbits, while $e = 1$ corresponds to parabolic orbits. It can be extended to an *analytic metric* on a two-sphere of revolution, where (r, θ) are the spherical coordinates.

If we now come back to the original system modelling coplanar orbit transfer and fix the direction of the control, we obtain a single-input periodic system of the form :

$$\frac{dq}{dl} = uF(q, l)$$

and an interesting case motivated by cone constraint due to electro-ionic propulsion is the so-called *tangential case* when the control has to be directed in the velocity. By averaging, we obtain again a full rank averaged Hamiltonian. A remarkable feature is that the same geometric coordinates remain orthogonal for the new metric which writes [5]

$$\bar{g}_t = \frac{dn^2}{9n^{1/3}} + n^{5/3} \left[\frac{1 + \sqrt{1-e^2}}{4(1-e^2)^{3/2}} de^2 + \frac{1 + \sqrt{1-e^2}}{4(1-e^2)} e^2 d\theta^2 \right].$$

In the two-input case, the change of variables $e = \sin r$ only consisted in lifting the Poincaré disk on which (e, θ) are polar coordinates onto \mathbb{S}^2 , where (r, θ) are the standard angles. To normalize now, we set again $n = (5\rho/2)^{6/5}$ and slightly twist the previous lifting according to :

$$e = \sin r \sqrt{1 + \cos^2 r}$$

to obtain the normal form :

$$\bar{g}_t = d\rho^2 + (\rho^2/c_t)g_t$$

with $c_t = c^2 = 2/5$ and $g_t = dr^2 + m_t^2(r)d\theta^2$,

$$m_t^2 = \sin^2 r \left(\frac{1 - (1/2) \sin^2 r}{1 - \sin^2 r} \right)^2.$$

In both cases we can define a homotopy deforming the round metric on \mathbb{S}^2 introducing $g_\lambda = dr^2 + XR_\lambda(X)d\theta^2$, setting $X = \sin^2 r$, while $R_0 = 1$, $R_\lambda(X) = R(\lambda X)$ and we have :

- *bi-input case* $R(X) = \frac{1}{1-X}$ and we have for $\lambda \in [0, 1]$ a homotopy from the round metric $\lambda = 0$, to the orbital transfer for which $\lambda = 4/5$, while the limit case $\lambda = 1$ is *singular*, since R has a pole for $X = 1$.
- *tangential case* $R(X) = \left(\frac{1-X/2}{1-X} \right)^2$ and we have for $\lambda \in [0, 1]$ a homotopy from the round metric $\lambda = 0$ to the orbital transfer which is *singular* since R has a pole of order two for $X = 1$.

2.2. Quantum control

We consider a dissipative two-level quantum system whose dynamics is governed by the *Redfield form of the Lindblad equation*, see [20], [7] for the details :

$$\begin{aligned} \dot{q}_1 &= -\Gamma q_1 + u_2 q_3 \\ \dot{q}_2 &= -\Gamma q_2 - u_1 q_3 \\ \dot{q}_3 &= \gamma_- - \gamma_+ q_3 + (u_1 q_2 - u_2 q_1) \end{aligned}$$

where the state space $q = (q_1, q_2, q_3)$ is restricted to the *Bloch ball* : $q_1^2 + q_2^2 + q_3^2 \leq 1$, and the control is of the form $u = u_1 + iu_2$, u_1, u_2 being two real functions, $|u| \leq 1$ and the three parameters $\Gamma, \gamma_-, \gamma_+$ describing the interaction with the environment and satisfying constraints : $\Gamma \geq \frac{\gamma_+}{2} \geq 0, \gamma_+ \geq |\gamma_-|$.

To minimize the effect of dissipation, we consider the problem of minimizing time of transfer, but the energy minimization problem shares similar properties.

The system is written

$$\dot{q} = F_0(q) + u_1 F_1(q) + u_2 F_2(q)$$

and introducing the Hamiltonian lifts $H_i = \langle p, F_i(q) \rangle, i = 0, 1, 2$ outside the switching surface $H_i = 0, i = 1, 2$ the maximal principle tells us that time optimal control trajectories are extremals solutions of the Hamiltonian vector field :

$$H(q, p) = H_0(q, p) + \left(\sum_{i=1}^2 H_i^2(q, p) \right)^{1/2}.$$

Since $|u| \leq 1$, the problem is invariant by change of coordinates and feedback transformations of the form $u = \beta(q)v$, where $\beta(q)$ is an orthogonal matrix.

We consider only the case where $\gamma_- = 0$. Since the Bloch ball is invariant, we introduce the spherical coordinates $q_3 = r \cos \phi, q_1 = r \sin \phi \cos \theta, q_2 = r \sin \phi \sin \theta$ and using the relations :

$$\begin{aligned} r^2 &= q_1^2 + q_2^2 + q_3^2 \\ \rho &= \ln r \\ \theta &= \arctan q_2 / q_1 \\ \phi &= \arccos q_3 / r \end{aligned}$$

and the feedback transformation $v_1 = \cos \theta u_1 + u_2 \sin \theta, v_2 = -\sin \theta u_1 + u_2 \cos \theta$, the system takes the form :

$$\begin{aligned}\dot{\rho} &= -(\Gamma \sin^2 \phi + \gamma_+ \cos^2 \phi) \\ \dot{\theta} &= -(\cotan \phi)v_1 \\ \dot{\phi} &= v_2 + \frac{\sin 2\phi}{2}(\gamma_+ - \Gamma)\end{aligned}$$

If we consider the Hamiltonian systems describing the evolution of generic extremals we get

$$H = -p_\rho(\Gamma \sin^2 \phi + \gamma_+ \cos^2 \phi) + p_\phi \frac{\sin 2\phi}{2}(\gamma_+ - \Gamma) + R$$

where R is respectively $(H_1^2 + H_2^2)^{1/2}$ in the time minimal case and $\frac{1}{2}(H_1^2 + H_2^2)$ in the energy minimization case, with :

$$\begin{aligned}H_1 &= -p_\theta \cotan \phi \\ H_2 &= p_\phi.\end{aligned}$$

We observe the following.

- If $\gamma_+ = \Gamma$ both extremal flows are a suspension of the extremal flows associated to the metric on the two-sphere \mathbb{S}^2 with coordinates $(r = \phi, \theta)$ given by :

$$g = dr^2 + (\tan^2 r)d\theta^2$$

and corresponds to the limit case $\lambda = 1$ for the homotopy of the previous section, in the bi-input case.

- If $\gamma_+ \neq \Gamma$, the Hamiltonians admit ρ and θ as cyclic coordinates, hence p_ρ and p_θ are first integrals. A more deep analysis reveals that the optimality analysis is related to a geometric perturbation of the previous case for which :
- For fixed p_ρ , the reduced Hamiltonians are associated to a one parameter family of optimal control problems on the two-sphere of revolution.
- For each such problem, the extremal flow shares similar properties to the case $\gamma_+ = \Gamma$, which allows to make the computation of the conjugate and cut loci tractable.

3. Conjugate and cut loci on a two-sphere of revolution

The objective of this section is to characterize when the cut locus of a point different from the pole on a two-sphere of revolution is reduced to a single segment and the conjugate locus has the standard astroid shape. This is based mainly on the analysis in [22] but for extensions it is deduced from the properties of the extremal flow only. This is not restrictive since in complete Riemannian 2D-manifold the computation of the cut locus is obtained by evaluating the separating line of a point q_0 , $L(q_0)$ where minimizers starting from q_0 are intersecting while the conjugate locus is limit points of intersecting neighboring extremals or equivalently the envelope of such extremals.

A compact Riemannian manifold (M, g) homeomorphic to a 2-sphere is called a *2-sphere of revolution*, if M admits a point p such that for any two points q_1, q_2 on M with $d(p, q_1) = d(p, q_2)$, where $d(\cdot, \cdot)$ denotes the Riemannian distance function, there exists an isometry f on M satisfying $f(q_1) = q_2$ and $f(p) = p$. The point p is called a *pole* of M . It is proved in [22] that each pole of a 2-sphere of revolution has a unique cut point, which is also a pole of the 2-sphere. By fixing a pole p on a 2-sphere M of revolution, we introduce geodesic polar coordinate (r, θ) around the pole p . The Riemannian metric g is expressed as $g = dr^2 + m(r)^2 d\theta^2$ on $M \setminus \{p, q\}$, where

$$m(r(x)) := \sqrt{g\left(\left(\frac{\partial}{\partial \theta}\right)_x, \left(\frac{\partial}{\partial \theta}\right)_x\right)}, \quad (1)$$

and q denotes the unique cut point of p . The Gaussian curvature G at a point $x \in M \setminus \{p, q\}$ is equal to

$$G(x) = -\frac{m''(r(x))}{m(r(x))}. \quad (2)$$

Each unit speed geodesic $\mu : \mathbb{R} \rightarrow M$ passing through the pole p is called a *meridian*. Since q is the unique cut point of p , μ passes through q . It is easily checked that μ is periodic, i.e., $\mu(r + 4a) = \mu(r)$, where $2a := d(p, q)$. Each curve $r = c \in (0, 2a)$ is called a *parallel*.

Let $\gamma(s) = (r(s), \theta(s))$ be a unit speed geodesic on the manifold M . Then, there exists a constant ν such that

$$m(r(s))^2 \cdot \theta'(s) = m(r(s)) \cos \eta(s) = \nu \quad (3)$$

holds for any s , where $\eta(s)$ denotes the angle $\angle(\dot{\gamma}(s), (\partial/\partial\theta)_{\gamma(s)})$ made by $\dot{\gamma}(s) := d\gamma(\partial/\partial s)$ and $(\partial/\partial\theta)_{\gamma(s)}$. The relation (3) is called the *Clairaut relation*, and the constant ν is called the *Clairaut constant of γ* . Since γ is unit speed, it follows from (3) that

$$r'(s) = \varepsilon(r'(s)) \frac{\sqrt{m(r(s))^2 - \nu^2}}{m(r(s))}, \quad (4)$$

where $\varepsilon(r'(s))$ denotes the sign of $r'(s)$. In particular, $r'(s) = 0$ if and only if $m(r(s)) = |\nu|$. Hence, the geodesic γ stays in the closure of a connected component of $(m \circ r)^{-1}(|\nu|, \infty)$, and if $m(r(s)) = |\nu|$ at $s = s_0$, then γ is tangent to the parallel $r = r(s_0)$. It follows from (3) and (4) that

$$\theta(s_2) - \theta(s_1) \equiv \varepsilon(r'(s)) \int_{r(s_1)}^{r(s_2)} f(r, \nu) dr \quad \text{mod } 2\pi \quad (5)$$

holds, where

$$f(r, \nu) = \frac{\nu}{m(r) \sqrt{m(r)^2 - \nu^2}},$$

if $r'(s) \neq 0$ on (s_1, s_2) , and moreover the length $L(\gamma|_{[s_1, s_2]})$ of $\gamma|_{[s_1, s_2]}$ equals

$$L(\gamma|_{[s_1, s_2]}) = \varepsilon(r'(s)) \int_{r(s_1)}^{r(s_2)} \frac{m(r)}{\sqrt{m(r)^2 - \nu^2}} dr \quad (6)$$

if $r'(s) \neq 0$ on (s_1, s_2) . Hereafter, we assume that the function m satisfies

$$m(r) = m(2a - r) \quad (7)$$

for any $r \in (0, 2a)$, where $2a = d(p, q)$. The parallel $r = a$ is called the *equator of M* . By (7), M is reflectively symmetric with respect to the equator.

For technical reasons, we introduce the Riemannian universal covering manifold

$$\tilde{M} := ((0, 2a) \times \mathbb{R}, d\tilde{r}^2 + m(\tilde{r})^2 d\tilde{\theta}^2)$$

of $(M \setminus \{p, q\}, dr^2 + m(r)^2 d\theta^2)$. Note that the equations (3), (4), and (6) hold for geodesics on \tilde{M} . The equation (5) is replaced by

$$\tilde{\theta}(\tilde{\gamma}(s_2)) - \tilde{\theta}(\tilde{\gamma}(s_1)) = \varepsilon((\tilde{r} \circ \tilde{\gamma})'(s)) \int_{\tilde{r}(\tilde{\gamma}(s_1))}^{\tilde{r}(\tilde{\gamma}(s_2))} f(r, \nu) dr. \quad (8)$$

Here, we assume that $(\tilde{r} \circ \tilde{\gamma})'(s) \neq 0$ on (s_1, s_2) . For each $\nu \in [0, m(a)]$, let $\tilde{\gamma}_\nu$ denote a unit speed geodesic on \tilde{M} with the Clairaut constant ν emanating from a point on $\tilde{r}^{-1}(a)$. Since $\tilde{\gamma}_\nu$ satisfies the Clairaut relation,

$$\angle(\dot{\tilde{\gamma}}_\nu(0), (\partial/\partial\tilde{\theta})_{\tilde{\gamma}_\nu(0)}) = \arccos \frac{\nu}{m(a)}$$

holds.

Lemma 3.1 *If $m' \neq 0$ on $(0, a)$, then for each $\nu \in (0, m(a))$, the geodesic $\tilde{\gamma}_\nu$ intersects $\tilde{r} = a$ again at a point $\tilde{\gamma}_\nu(t_0(\nu))$, and the function, which is called the half period function of \tilde{M} ,*

$$\varphi(\nu) := 2 \int_{\xi(\nu)}^a \frac{\nu}{m(r) \sqrt{m(r)^2 - \nu^2}} dr \quad (9)$$

is well-defined and is equal to $\tilde{\theta}(\tilde{\gamma}_\nu(t_0(\nu))) - \tilde{\theta}(\tilde{\gamma}_\nu(0))$. Here, $\xi(\nu) := (m|_{[0, a]})^{-1}(\nu)$.

Proof. Choose any $v \in (0, m(a))$ and fix it. We may assume that $(\tilde{r} \circ \tilde{\gamma}_v)'(0) < 0$, since (7) holds. It is clear from (4) that $\tilde{\gamma}_v$ stays in $\tilde{r}^{-1}[\xi(v), 2a - \xi(v)]$. Since $m' \neq 0$ on $(0, a)$, it follows from [19, Lemma 7.1.7] that $(\tilde{r} \circ \tilde{\gamma}_v)'(t_1) = 0$ for some $t_1 > 0$, i.e., $\tilde{\gamma}_v$ is tangent to the parallel arc $\tilde{r} = \xi(v)$ at $\tilde{\gamma}_v(t_1)$. From (8), we get

$$\tilde{\theta}(\tilde{\gamma}_v(t_1)) - \tilde{\theta}(\tilde{\gamma}_v(0)) = \frac{1}{2}\varphi(v). \quad (10)$$

Since $(\tilde{r} \circ \tilde{\gamma}_v)'(t) > 0$ for any $t > t_1$ sufficiently close to t_1 and $(\tilde{r} \circ \tilde{\gamma}_v)'(t) \neq 0$ on (t_1, t) if $\tilde{r} \circ \tilde{\gamma}_v < 2a - \xi(v)$, there exists $t_0(v) (> t_1)$ such that $\tilde{r} \circ \tilde{\gamma}_v(t_0(v)) = a$ and $\tilde{r} \circ \tilde{\gamma}_v < a$ on $(t_1, t_0(v))$. Hence $\tilde{\gamma}_v$ intersects $\tilde{r} = a$ again at the point $\tilde{\gamma}_v(t_0(v))$. Since $\tilde{\theta}(\tilde{\gamma}_v(t_0(v))) - \tilde{\theta}(\tilde{\gamma}_v(t_1))$ is equal to $\varphi(v)/2$, the proof of our lemma is complete. \square

It is not difficult to calculate the parameter value $t_0(v)$ in Lemma 3.1. From (6) and (7), we get

$$t_0(v) = 2 \int_{\xi(v)}^a \frac{m(r)}{\sqrt{m(r)^2 - v^2}} dr. \quad (11)$$

Since

$$\frac{m}{\sqrt{m^2 - v^2}} = \frac{\sqrt{m^2 - v^2}}{m} + \frac{v^2}{m\sqrt{m^2 - v^2}},$$

we obtain

$$t_0(v) = 2 \int_{\xi(v)}^a \frac{\sqrt{m(r)^2 - v^2}}{m(r)} dr + v\varphi(v). \quad (12)$$

Lemma 3.2 *If the cut locus of a point q on the equator $r = a$ is a subset of the equator, then the function $\varphi(v)$ is well-defined and monotone non-increasing on $(0, m(a))$.*

Proof. If the cut locus C_q of q consists of a single point, then it is clear that $\varphi(v)$ is constant. Thus, we may assume that C_q is a subarc of the equator. Let q_0 be an endpoint of C_q . First we will prove that $q_0 := \gamma_{m(a)}(t_0)$ is conjugate to $q := \gamma_{m(a)}(0)$ along $\gamma_{m(a)}$, where $\gamma_{m(a)}$ denotes the subarc of the equator joining q to q_0 . Since $\gamma_{m(a)}$ does not contain a cut point of q in its interior, it is a minimal geodesic segment joining q to q_0 . If $\gamma_{m(a)}$ is the unique minimal geodesic segment joining q to q_0 , then it is clear that q_0 is conjugate to q along $\gamma_{m(a)}$. Hence we suppose that there exists a minimal geodesic segment $\alpha : [0, t_0] \rightarrow M$ joining q and q_0 which bounds a disc domain D together with $\gamma_{m(a)}|_{[0, t_0]}$. Here, we may assume that $(r \circ \alpha)'(0) < 0$ and $(r \circ \gamma_{m(a)})'(0) < 0$ by (7). Since D has no cut point of q , any geodesic segment β emanating from q with $(r \circ \beta)'(0) < 0$ must pass through the point q_0 , if $\angle(\dot{\beta}(0), \dot{\gamma}_{m(a)}(0)) < \angle(\dot{\alpha}(0), \dot{\gamma}_{m(a)}(0))$. Thus, we get a geodesic variation of $\gamma_{m(a)}|_{[0, t_0]}$, which is a family of geodesic segments joining q to q_0 . Hence, we have proved that q_0 is a conjugate point of q along $\gamma_{m(a)}$. This implies that the Gaussian curvature G is positive on the equator. Since $m''(a) = -G(q)m(a) < 0$ by (2) and $m'(a) = 0$, m' is positive on $(a - \delta, a)$ for some $\delta > 0$. Suppose that $m'(b) = 0$ for some $b \in (0, a)$. From [19, Lemma 7.1.4], the parallel $r = b$ is a geodesic. By choosing the maximal $b (< a)$ satisfying $m'(b) = 0$, we may assume that m' is positive on (b, a) . Thus, $m(r) > m(b)$ on $(b, a]$. Suppose that the geodesic $\gamma_{m(b)}$ is tangent to a parallel. Here, for each $v \in [0, m(a))$, γ_v denotes the unit speed geodesic emanating from q with the Clairaut constant v satisfying $(r \circ \gamma_v)'(0) < 0$. From (4), the possible parallel, to which $\gamma_{m(b)}$ is tangent, is the geodesic parallel $r = b$, but $\gamma_{m(b)}$ can not be tangent to another geodesic $r = b$. Hence, $(r \circ \gamma_{m(b)})'(0) \neq 0$, and in particular, $\gamma_{m(b)}$ does not intersect the equator again (see [19, Figure 7.1.2] on the behaviour of such a geodesic). Since M is compact, there exists a cut point of q along $\gamma_{m(b)}$. This contradicts the assumption that C_q is a subarc of the equator. Therefore, m' is non-zero on $(0, a)$ and the function $\varphi(v)$ is well-defined on $(0, m(a))$. It is now clear that γ_v intersects the equator again at $\gamma_v(t_0(v))$ and $\theta(\gamma_v(t)) \leq \pi$ for any $v \in (0, m(a))$ and any $t \in (0, t_0(v)]$. Here, for a technical reason, the geodesic polar coordinates (r, θ) are chosen so as to satisfy $\theta(\gamma_v(0)) = \theta(q) = 0$. In particular, $\varphi(v) \leq \pi$ for any $v \in (0, m(a))$.

Next, we will prove that φ is monotone non-increasing. Choose any two numbers $v_1 < v_2$ in $(0, m(a))$ and fix them. By the Clairaut relation and the inequality,

$$\arccos \frac{v_1}{m(a)} > \arccos \frac{v_2}{m(a)},$$

the geodesic $\gamma_{v_2}|_{(0, \delta)}$ lies in the domain D_{v_1} bounded by the equator and $\gamma_{v_1}|_{[0, t_0(v_1)]}$ for some $\delta > 0$. Since C_q is a subset of the equator, the geodesic segment γ_{v_2} lying in D_{v_1} does not pass through $\gamma_{v_1}|_{(0, t_0(v_1))}$, but passes through the equator and intersects at $\gamma_{v_2}(t_0(v_2))$. Thus, $\gamma_{v_2}(t_0(v_2))$ is a point on the subarc of the equator with endpoints $\gamma_{v_1}(0)$ and $\gamma_{v_1}(t_0(v_1))$, and in particular, $\varphi(v_2) \leq \varphi(v_1)$ holds. \square

Lemma 3.3 *If $m' \neq 0$ on $(0, a)$ and the function $\varphi : (0, m(a)) \rightarrow \mathbb{R}$ is monotone non-increasing, then the cut locus of each point on the equator $r = a$ is a subset of the equator.*

Proof. Let γ_v , $v \in [0, m(a)]$, denote the geodesic emanating from a point q on the equator that was defined in the proof of Lemma 3.2. It follows from Lemma 3.1 that for each $v \in [0, m(a))$, γ_v intersects the equator again at $\gamma_v(t_0(v))$. Choose any $v \in (0, m(a))$ and fix it. It follows from [19, Proposition 7.2.3] that $\gamma_v(t_1)$ is the first conjugate point of q along γ_v if and only if

$$\frac{\partial \theta}{\partial v}(r(\gamma_v(t_1)), v) = 0, \quad (13)$$

where

$$\theta(r, v) := \int_{\xi(v)}^a f(r, v) dr + \int_{\xi(v)}^r f(r, v) dr. \quad (14)$$

Notice that it follows from [19, Proposition 7.2.2] and [19, Corollary 7.2.1] that there is no conjugate point of q along $\gamma_v|_{[0, t]}$, if $(r \circ \gamma_v)' \neq 0$ on $[0, t]$. It is clear from (7) that

$$\theta(r, v) = \varphi(v) - \int_r^a f_v(r, v) dr \quad (15)$$

holds. Hence,

$$\frac{\partial \theta}{\partial v}(r \circ \gamma_v(t), v) = \varphi'(v) - \int_{r(\gamma_v(t))}^a f_v(r, v) dr < 0, \quad (16)$$

if $r \circ \gamma_v(t) < a$, since $\varphi'(v) \leq 0$ on $(0, m(a))$. Note that

$$f_v(r, v) = \frac{m(r)}{(m(r)^2 - v^2)^{\frac{3}{2}}} > 0.$$

Thus, there is no conjugate point of q along $\gamma_v|_{[0, t]}$, if $\gamma_v([0, t]) \subset r^{-1}(0, a)$. By taking the limit in (16), we may prove that there is no conjugate point of q along $\gamma_{m(0)}|_{[0, t]}$, if $\gamma_{m(0)}([0, t]) \subset r^{-1}[0, a)$. Therefore, we have proved that there is no conjugate point of q along $\gamma_v|_{[0, t]}$, if $\gamma_v([0, t]) \subset r^{-1}[0, a)$ and $v \in [0, m(a))$.

Suppose that there exists a cut point $x \notin r^{-1}(a)$ of q . From (7), we may assume that x is a point in $r^{-1}[0, a)$. Let $\gamma_{v_1} : [0, d(q, x)] \rightarrow M$ denote a minimal geodesic segment joining q to x . We may assume that x is conjugate to q along γ_{v_1} . Otherwise, there exists a minimal geodesic segment $\gamma_{v_2}|_{[0, d(q, x)]}$, $v_2 \in [0, m(a)) \setminus \{v_1\}$, joining q to x . Thus, both geodesic segments bound a disc domain D . Since the cut locus C_q in D has no circle, we may find an endpoint $y \in r^{-1}(0, a)$ in $C_q \cap D$. The endpoint y is conjugate to q along any minimal geodesic segment joining q to y . Therefore, by exchanging x and y , we may assume that x is conjugate to q along γ_{v_1} . This contradicts the fact that there is no conjugate point of q along $\gamma_v|_{[0, t]}$, if $\gamma_v([0, t]) \subset r^{-1}[0, a)$. Therefore, the cut locus of q is a subset of the equator. \square

Choose any point q in M , which is not a pole. We introduce geodesic polar coordinates (r, θ) around the pole p on M satisfying $\theta(q) = 0$. Put $u := r(q) \in (0, 2a)$. For each $v \in (0, m(u)]$, let $\alpha_v, \beta_v : [0, \infty) \rightarrow M$ denote the geodesics emanating from $q = \alpha_v(0) = \beta_v(0)$ with the Clairaut constant v satisfying

$$(r \circ \alpha_v)'(0) \leq 0 \leq (r \circ \beta_v)'(0).$$

Hence, from the Clairaut relation,

$$\angle(\dot{\alpha}_v(0), (\partial/\partial\theta)_q) = \angle(\dot{\beta}_v(0), (\partial/\partial\theta)_q) = \arccos \frac{v}{m(u)}.$$

From (7), the geodesics α_v and β_v intersect again at $(r, \theta)^{-1}(2a - u, \varphi(v)) = \alpha_v(t_0(v)) = \beta_v(t_0(v))$ (see [22], or [11]).

Lemma 3.4 *Assume that $m' \neq 0$ on $(0, a)$ and that φ is monotone non-increasing. Then, for each $v \in (0, m(u)]$, $\alpha_v|_{[0, t_0(v)]}$ is minimal. Furthermore, each point of $r^{-1}(2a - u) \cap \theta^{-1}[\varphi(m(u)), \pi]$ is a cut point of q .*

Proof. Since the geodesic $\gamma_{m(0)}$ meets the equator again at the antipodal point of $\gamma_{m(0)}(0)$, we have $\lim_{v \downarrow 0} \varphi(v) = \pi$. Thus, $0 < \varphi(v) \leq \pi$ for any $v \in (0, m(a))$, since φ is monotone non-increasing. If $\varphi(m(u)) = \pi$, then $\varphi(v) = \pi$ for any $v \in [0, m(u)]$. Hence the cut locus of q consists of a single point. It is now clear that both geodesics $\alpha|_{[0, t_0(v)]}$ and $\beta|_{[0, t_0(v)]}$ are minimal for any $v \in [0, m(u)]$. Hence we may assume that $\varphi(m(u)) < \pi$. Choose any point $x \in r^{-1}(2a - u) \cap \theta^{-1}[\varphi(m(u)), \pi]$. From (6), the length t_1 of α equals

$$\int_u^{2a-\xi(v_1)} \frac{m(r)}{\sqrt{m(r)^2 - v_1^2}} dr + \int_{2a-u}^{2a-\xi(v_1)} \frac{m(r)}{\sqrt{m(r)^2 - v_1^2}} dr \quad (17)$$

(if $\alpha = \beta_{v_1}|_{[0, t_1]}$), or to

$$\int_{\xi(v_1)}^u \frac{m(r)}{\sqrt{m(r)^2 - v_1^2}} dr + \int_{\xi(v_1)}^{2a-u} \frac{m(r)}{\sqrt{m(r)^2 - v_1^2}} dr \quad (18)$$

(if $\alpha = \alpha_{v_1}|_{[0, t_1]}$). By (7) and (11), both equations (17) and (18) equal

$$t_1 = 2 \int_{\xi(v_1)}^a \frac{m(r)}{\sqrt{m(r)^2 - v_1^2}} dr = t_0(v_1). \quad (19)$$

Therefore, $\alpha_{v_1}|_{[0, t_0(v_1)]}$ and $\beta_{v_1}|_{[0, t_0(v_1)]}$ are minimal geodesic segments joining q to x . Furthermore, for any $v \in \varphi^{-1}(\theta(x))$, $\alpha_v|_{[0, t_0(v)]}$ and $\beta_v|_{[0, t_0(v)]}$ are minimal, since $t_0'(v) = v\varphi'(v)$ by (12). Since x is arbitrarily taken, this implies that $\alpha_v|_{[0, t_0(v)]}$ and $\beta_v|_{[0, t_0(v)]}$ are minimal for any $v \in (0, m(u)]$ with $\varphi(v) < \pi$, hence for any $v \in (0, m(u)]$ from the limit argument. Therefore, $\alpha_v|_{[0, t_0(v)]}$ is minimal for all $v \in (0, m(u))$. The second claim is clear, since each point of $r^{-1}(2a - u) \cap \theta^{-1}(\varphi(m(u)), \pi]$ is joined by two minimal geodesic segments $\alpha_v|_{[0, t_0(v)]}$ and $\beta_v|_{[0, t_0(v)]}$ for some $v \in [0, m(u))$, and the cut locus is closed. \square

Theorem 3.5 *Let $(M, dr^2 + m(r)^2 d\theta^2)$ denote a 2-sphere of revolution, where $m : (0, 2a) \rightarrow (0, \infty)$ is a smooth function satisfying (7). If the cut locus of a point on $r = a$ is a subset of $r = a$, then, the cut locus of a point q with $r(q) \in (0, 2a) \setminus \{a\}$ is a subset of the antipodal parallel $r = 2a - r(q)$.*

Proof. Let $q \in r^{-1}((0, 2a) \setminus \{a\})$ be any point, and set $u := r(q)$. Since M is reflectively symmetric with respect to the meridian passing through q , the set $r^{-1}(2a - u) \cap \theta^{-1}[\varphi(m(u)), 2\pi - \varphi(m(u))]$ is a subset of C_q by Lemma 3.4. Choose any cut point x of q . Then, we have a minimal geodesic segment γ joining q to x . Since we may assume

that $0 < \theta(x) \leq \pi$, γ is equal to α_{v_1} , or β_{v_1} for some $v_1 \in [0, m(u)]$. Here v_1 denotes the Clairaut constant of γ . The point x is not an interior point of $\alpha_{v_1}|_{[0, t_0(v_1)]}$, or $\beta_{v_1}|_{[0, t_0(v_1)]}$, since both segments are minimal. Furthermore, γ is a subarc of $\alpha_{v_1}|_{[0, t_0(v_1)]}$, or $\beta_{v_1}|_{[0, t_0(v_1)]}$, since γ is minimal. Hence,

$$x = \alpha_{v_1}(t_0(v_1)) (= \beta_{v_1}(t_0(v_1))).$$

Since $r(\alpha_{v_1}(t_0(v_1))) = 2a - u$ and $\varphi(m(u)) \leq \varphi(v_1) = \theta(x) \leq \pi$, any cut point of q is a point of $r^{-1}(2a - u) \cap \theta^{-1}[\varphi(m(u)), 2\pi - \varphi(m(u))]$, which is a subarc of the antipodal parallel of q . \square

Theorem 3.6 *Let $(M, dr^2 + m(r)^2 d\theta^2)$ denote a 2-sphere of revolution, where $m : (0, 2a) \rightarrow (0, \infty)$ is a smooth function satisfying (7). Assume that the cut locus of a point on $r = a$ is a subset of $r = a$. If the function φ defined by (9) is such that $\varphi''(v) \geq 0$ on $(0, m(a))$ and $\varphi''(v) > 0$ whenever $\varphi'(v) = 0$, then the first conjugate locus of any point q which is not a pole of M has exactly four cusps. Recall that the function φ is defined in (9).*

Proof. Choose any point $q \in M$ which is not a pole. Set $u := r(q) \in (0, 2a)$. It follows from [19, Proposition 7.2.3] that $\alpha_v(t_c(v))$ is the first conjugate point of q along α_v if and only if

$$\frac{\partial \theta_\alpha}{\partial v}(r(\alpha_v(t_c(v))), v) = 0, \quad (20)$$

where

$$\theta_\alpha(r, v) = \int_{\xi(v)}^u f(r, v) dr + \int_{\xi(v)}^r f(r, v) dr. \quad (21)$$

From (7), it is clear that

$$\theta_\alpha(r, v) = \varphi(v) - \int_r^{2a-u} f(r, v) dr \quad (22)$$

holds. Hence,

$$\frac{\partial \theta_\alpha}{\partial v}(r, v) = \varphi'(v) - \int_r^{2a-u} f_v(r, v) dr, \quad (23)$$

where

$$f_v(r, v) = \frac{m(r)}{(m(r)^2 - v^2)^{\frac{3}{2}}}.$$

From (20) and (23), it follows that

$$\varphi'(v) = \int_{u_\alpha(v)}^{2a-u} f_v(r, v) dr, \quad (24)$$

where $u_\alpha(v) = r(\alpha_v(t_c(v)))$. Hence, the first conjugate point of q along α_v is given by

$$\theta = \theta_\alpha(u_\alpha(v), v), \quad r = u_\alpha(v). \quad (25)$$

Since we assume that $\varphi'(v) \leq 0$ for each $v \in (0, m(u))$,

$$2a - u \leq u_\alpha(v) \quad (26)$$

by (24). Furthermore, $\varphi'(v) = 0$ if and only if $2a - u = u_\alpha(v)$. By differentiating (24) with respect to v , we have

$$\varphi''(v) + \int_{2a-u}^{u_\alpha(v)} f_{vv}(r, v) dr = -f_v(u_\alpha(v), v)u'_\alpha(v). \quad (27)$$

Since $f_{vv}(r, v) = 3vm(r)(m(r)^2 - v^2)^{-5/2} > 0$ and $\varphi''(v) \geq 0$, $u'_\alpha(v) \leq 0$ on $(0, m(u))$. If $u'_\alpha(v) = 0$, then, by (27), $\varphi''(v) = 0$ and $2a - u = u_\alpha(v)$. This implies that $\varphi''(v) = \varphi'(v) = 0$. This contradicts the assumption of

our theorem. Therefore, $u'_\alpha(v) < 0$ on $(0, m(u))$. In particular, there is no cusp on the open arc defined by (25), $v \in (0, m(u))$. Let $v \in (0, m(u))$ be any fixed number. The velocity vector $v_\alpha(v)$ of the curve by (25) is given by

$$v_\alpha(v) = f(u_\alpha(v), v) u'_\alpha(v) \left(\frac{\partial}{\partial \theta} \right)_{\alpha_v(t_c(v))} + u'_\alpha(v) \left(\frac{\partial}{\partial r} \right)_{\alpha_v(t_c(v))}. \quad (28)$$

Hence, $v_\alpha(v)$ is parallel to

$$f(u_\alpha(v), v) \left(\frac{\partial}{\partial \theta} \right)_{\alpha_v(t_c(v))} + \left(\frac{\partial}{\partial r} \right)_{\alpha_v(t_c(v))}.$$

Since $\lim_{v \downarrow 0} f(u_\alpha(v), v) = 0$ and $\lim_{v \uparrow m(u)} f(u_\alpha(v), v) = \infty$,

$$\lim_{v \downarrow 0} \frac{1}{\|v_\alpha(v)\|} v_\alpha(v) = \left(\frac{\partial}{\partial r} \right)_{\alpha_0(t_c(0))} \quad (29)$$

and

$$\lim_{v \uparrow m(u)} \frac{1}{\|v_\alpha(v)\|} v_\alpha(v) = \frac{1}{m(\alpha_{m(u)}(t_c(m(u))))} \left(\frac{\partial}{\partial \theta} \right)_{\alpha_{m(u)}(t_c(m(u)))}, \quad (30)$$

where $\|v_\alpha(v)\| := \sqrt{g(v_\alpha(v), v_\alpha(v))}$. Hence the subarc of the conjugate locus given by (25) is tangent to the parallel $r = r(\alpha_{m(0)}(t_c(0)))$ at $\alpha_{m(0)}(t_c(0))$ and to the opposite meridian $\theta^{-1}(\pi)$ at $\alpha_{m(u)}(t_c(m(u)))$.

Next, we will argue about the first conjugate locus of q along β_v , $v \in (0, m(u))$. It follows from [19, Proposition 7.2.3] that $\beta_v(t_f(v))$ is the first conjugate point of q along β_v if and only if

$$\frac{\partial \theta_\beta}{\partial v} (r(\beta_v(t_f(v))), v) = 0, \quad (31)$$

where

$$\theta_\beta(r, v) = \int_u^{2a-\xi(v)} f(r, v) dr + \int_r^{2a-\xi(v)} f(r, v) dr. \quad (32)$$

From (7), it is clear that

$$\theta_\beta(r, v) = \varphi(v) + \int_r^{2a-u} f(r, v) dr \quad (33)$$

holds. Hence, we have

$$\varphi'(v) + \int_{u_\beta(v)}^{2a-u} f_v(r, v) dr = 0, \quad (34)$$

where $u_\beta(v) := r(\beta_v(t_f(v)))$. By using the same argument above, we may conclude that $u'_\beta(v) > 0$ on $(0, m(u))$. In particular, there is no cusp on the open arc defined by

$$\theta = \theta_\beta(u_\beta(v), v), \quad r = u_\beta(v), \quad (35)$$

on $v \in (0, m(u))$. It is easy to prove that

$$\lim_{v \downarrow 0} \frac{1}{\|v_\beta(v)\|} v_\beta(v) = \left(\frac{\partial}{\partial r} \right)_{\beta_0(t_f(0))} \quad (36)$$

and

$$\lim_{v \uparrow m(u)} \frac{1}{\|v_\beta(v)\|} v_\beta(v) = \frac{1}{m(\beta_{m(u)}(t_f(m(u))))} \left(\frac{\partial}{\partial \theta} \right)_{\beta_{m(u)}(t_f(m(u)))}, \quad (37)$$

where $v_\beta(v)$ denotes the velocity vector of the curve defined by (35). Since $\beta_{m(u)} = \alpha_{m(u)}$, by (30) and (37), we get

$$\lim_{v \uparrow m(u)} \frac{1}{\|v_\beta(v)\|} v_\beta(v) = \lim_{v \uparrow m(u)} \frac{1}{\|v_\alpha(v)\|} v_\alpha(v). \quad (38)$$

Therefore, the point $\alpha_{m(u)}(t_c(m(u))) = \beta_{m(u)}(t_f(m(u)))$ is a cusp of the first conjugate locus of q . Since M has a reflective symmetry with respect to the meridian passing through q , the points $\alpha_{m(0)}(t_c(m(u)))$ and $\beta_{m(0)}(t_c(m(u)))$ lying on the opposite half meridian to q are cusps of the conjugate locus. Therefore, the conjugate locus of q has exactly four cusps which consists of one pair lying on the parallel $r = 2a - r(q)$, the other lying on the opposite half meridian to q . \square

The following dual to Theorem 3.6 is also true, but we do not know examples satisfying the assumption in the theorem.

Theorem 3.7

Let $(M, dr^2 + m(r)^2 d\theta^2)$ denote a 2-sphere of revolution, where $m : (0, 2a) \rightarrow (0, \infty)$ is a smooth function satisfying (7). Assume that $\varphi : (0, m(a)) \rightarrow \mathbb{R}$ is well-defined, i.e., $m' \neq 0$ on $(0, a)$, and φ is monotone non-decreasing on $(0, m(a))$. If $\varphi''(v) \leq 0$ on $(0, m(a))$ and $\varphi''(v) < 0$ whenever $\varphi'(v) = 0$, then the first conjugate locus of any point q which is not a pole of M has exactly four cusps and the cut locus of q is a subarc of the opposite half meridian to q .

Proof. It is proved in Theorem 3.6 that the first conjugate point of q along $\alpha_v, \beta_v, v \in (0, m(u))$, is given by

$$\theta = \theta_\alpha(u_\alpha(v), v), \quad r = u_\alpha(v),$$

and

$$\theta = \theta_\beta(u_\beta(v), v), \quad r = u_\beta(v),$$

respectively. By making use of the assumption that $\varphi'(v) \geq 0$, we may prove that

$$u'_\alpha(v) > 0 \quad \text{and} \quad u'_\beta(v) < 0 \tag{39}$$

on $(0, m(u))$. The first claim is now clear from the argument in the proof of Theorem 3.6.

Since

$$\frac{\partial \theta_\alpha}{\partial v}(u_\alpha(v), v) = 0 \quad \text{and} \quad \frac{\partial \theta_\beta}{\partial v}(u_\beta(v), v) = 0,$$

we get

$$\frac{\partial}{\partial v} \theta_\alpha(u_\alpha(v), v) = u'_\alpha(v) f(u_\alpha(v), v), \quad \frac{\partial}{\partial v} \theta_\beta(u_\beta(v), v) = -u'_\beta(v) f(u_\beta(v), v). \tag{40}$$

Hence, by (39), both functions $\theta_\alpha(u_\alpha(v), v)$ and $\theta_\beta(u_\beta(v), v)$ are strictly monotone increasing. In particular,

$$\theta_\alpha(u_\alpha(v), v) > \lim_{v \downarrow 0} \theta_\alpha(u_\alpha(v), v), \quad \theta_\beta(u_\beta(v), v) > \lim_{v \downarrow 0} \theta_\beta(u_\beta(v), v) \tag{41}$$

for any $v \in (0, m(u)]$. Since the first conjugate points of $\alpha_{m(0)}$ and $\beta_{m(0)}$ lie in $\theta^{-1}(\pi)$ respectively,

$$\lim_{v \downarrow 0} \theta_\alpha(u_\alpha(v), v) = \lim_{v \downarrow 0} \theta_\beta(u_\beta(v), v) = \pi.$$

Here, we should recall that the geodesic polar coordinates (r, θ) is chosen so as to satisfy $\theta(q) = \pi$. Therefore, by (41), there is no conjugate point of q in $M \setminus \theta^{-1}(\pi)$. This implies that the cut locus of q is a subarc of the opposite half meridian $\theta^{-1}(\pi)$. \square

4. Applications

Next, we apply our results to the problems introduced in section 2.

Let g_λ be the family of analytic metrics on \mathbb{S}^2 defined by

$$g_\lambda = dr^2 + m_\lambda^2(r)d\theta^2$$

with

$$m_\lambda(r) = \sqrt{\lambda + 1} \sin r / \sqrt{1 + \lambda \cos^2 r}, \quad \lambda \geq 0. \quad (42)$$

It is clear that m_λ satisfies

$$m_\lambda(r) = m_\lambda(\pi - r),$$

so the Riemannian manifold $M_\lambda := (\mathbb{S}^2, g_\lambda)$ is a 2-sphere of revolution that is reflectively symmetric with respect to the equator $r = \pi/2$. This family of metrics contains the metric \bar{g} associated with the averaged controlled Kepler equation introduced in §2 (set $\lambda = 4$), and defines a path between the following two remarkable metrics.

The case $\lambda = 0$. It corresponds to the standard metric $g_0 = dr^2 + \sin^2 r d\theta^2$ on \mathbb{S}^2 with Gaussian curvature equal to one which is obtained by restricting the Euclidian metric on \mathbb{R}^3 to the sphere.

The case $\lambda = \infty$. In the limit case, the metric becomes

$$g_\infty = dr^2 + \tan^2 r d\theta^2$$

and is singular along the equator $r = \pi/2$. The Gauss curvature is $-2/\cos^2 r$ which is strictly negative on each hemisphere and tends to $-\infty$ when r tends to $\pi/2$.

Lemma 4.1 *The Gaussian curvature G_λ of M_λ is given by*

$$G_\lambda(q) = \frac{(\lambda + 1)(1 - 2\lambda \cos^2 r(q))}{(1 + \lambda \cos^2 r(q))^2} \quad (43)$$

at a point $q \in M_\lambda$. Furthermore, we have

$$\left(\frac{\partial}{\partial r} \right)_q G_\lambda = \frac{2\lambda(\lambda + 1) \sin 2r(q)}{(1 + \lambda \cos^2 r(q))^3} (2 - \lambda \cos^2 r(q)). \quad (44)$$

In particular, if $\lambda > 2$, then G_λ is not monotone along a meridian from a pole to the point on the equator.

Proof. By (42), we have

$$m'_\lambda(r) = \frac{(\lambda + 1) \cos r}{(1 + \lambda \cos^2 r) \sin r} m_\lambda(r). \quad (45)$$

Since

$$G_\lambda(q) = -\frac{m''_\lambda(r(q))}{m_\lambda(r(q))}$$

for $q \in r^{-1}(0, 2\pi)$, it follows from (45) that

$$G_\lambda(q) = \frac{(\lambda + 1)(1 - 2\lambda \cos^2 r(q))}{(1 + \lambda \cos^2 r(q))^2}. \quad (46)$$

By making use of (46), it is easy to show (44). \square

Lemma 4.2 *Let a, b, c be positive numbers satisfying $c > b$. Then,*

$$\begin{aligned} & \int \frac{1}{x(x+a)\sqrt{(x-b)(c-x)}} dx \\ &= \frac{2}{a} \left\{ \frac{1}{\sqrt{bc}} \arctan \left(\sqrt{\frac{c}{b}} t \right) - \frac{1}{\sqrt{(a+c)(a+b)}} \arctan \left(\sqrt{\frac{a+c}{a+b}} t \right) \right\} \end{aligned} \quad (47)$$

holds, where

$$t = \sqrt{\frac{x-b}{c-x}}.$$

In particular,

$$\int_b^c \frac{1}{x(x+a)\sqrt{(x-b)(c-x)}} dx = \frac{\pi}{a} \left\{ \frac{1}{\sqrt{bc}} - \frac{1}{\sqrt{(a+c)(a+b)}} \right\}$$

holds.

Proof. From direct computation, we get

$$\begin{aligned} & \frac{d}{dx} \left\{ \frac{1}{\sqrt{bc}} \arctan \left(\sqrt{\frac{c}{b}} t \right) - \frac{1}{\sqrt{(a+c)(a+b)}} \arctan \left(\sqrt{\frac{a+c}{a+b}} t \right) \right\} \\ &= \frac{a}{2} \cdot \frac{1}{x(x+a)\sqrt{(x-b)(c-x)}}. \end{aligned}$$

Hence, we obtain (47). □

Proposition 4.3 For the 2-sphere of revolution M_λ , we get

$$\varphi(v) = \pi - \frac{\lambda \pi v}{\sqrt{\lambda+1}\sqrt{\lambda+1+\lambda v^2}} \quad (48)$$

for each $v \in [0, m_\lambda(\pi/2)]$.

Proof. By putting $x = m_\lambda(r)^2$, we have, from (42) and (45),

$$dr = \frac{(1 + \lambda \cos^2 r) \tan r}{2(\lambda + 1)x} dx. \quad (49)$$

Since

$$x = m_\lambda(r)^2 = \frac{(\lambda + 1)(1 - \cos^2 r)}{1 + \lambda \cos^2 r}, \quad (50)$$

we obtain

$$\cos^2 r = \frac{\lambda + 1 - x}{\lambda x + \lambda + 1}. \quad (51)$$

Since

$$\tan^2 r = \frac{1}{\cos^2 r} - 1,$$

we have

$$\tan^2 r = \frac{(\lambda + 1)x}{\lambda + 1 - x}. \quad (52)$$

Combining (49), (51), and (52), we obtain

$$dr = \frac{(\lambda + 1)^{\frac{3}{2}}}{2(\lambda x + \lambda + 1)\sqrt{x(\lambda + 1 - x)}} dx. \quad (53)$$

Hence, we get

$$\varphi(v) = (\lambda + 1)^{\frac{3}{2}} v \int_{v^2}^{\lambda+1} \frac{1}{x(\lambda x + \lambda + 1)\sqrt{(x - v^2)(\lambda + 1 - x)}} dx. \quad (54)$$

It follows from Lemma 4.2 that

$$\varphi(v) = \pi - \frac{\lambda \pi v}{\sqrt{\lambda+1}\sqrt{\lambda+1+\lambda v^2}}.$$

□

Theorem 4.4 *If $\lambda > 0$, then, for each point q of M_λ distinct from a pole, the cut locus of q is a subarc of the antipodal parallel to q and the first conjugate locus of q has exactly four cusps.*

Proof. It is clear that

$$\varphi'(v) = \frac{-\lambda \pi \sqrt{\lambda + 1}}{(\lambda + 1 + \lambda v^2)^{\frac{3}{2}}}$$

and

$$\varphi''(v) = \frac{3\pi\lambda^2 v \sqrt{\lambda + 1}}{(\lambda + 1 + \lambda v^2)^{\frac{5}{2}}}.$$

In particular,

$$\varphi'(v) < 0 < \varphi''(v)$$

on $(0, m_\lambda(\pi/2))$, if $\lambda > 0$. The claims of Theorem 4.4 are now clear from Lemma 3.3, Theorem 3.5, and Theorem 3.6. \square

Remark 1 *If $\lambda > 2$, then the Gaussian curvature of M_λ is not monotone along a meridian from a pole to the point on the equator. Therefore, the family $M_\lambda, \lambda > 2$, is a new example which has the simple cut locus structure.*

On Fig. 1 we represent conjugate and cut loci in averaged orbital transfer ($\lambda = 4$). Since to loose optimality an extremal trajectory has to cross the equator, $e = 1$, we conclude that extremals are optimal in the physical elliptic domain. The conjugate and cut loci for $\lambda = \infty$ are given on Figs. 2 and 3. In this singular case, since the curvature outside the equator is strictly negative, geodesics starting from a point not on the equator have again to cross it to have conjugate points, the cut locus still being included in the antipodal parallel. For a point on the equator, the conjugate and cut loci accumulate near the the point itself, and the cut locus is the equator minus the point. The first two cusps of the conjugate locus disappear and are replaced by a contact of order two at the initial point [1].

5. The singular case

In this section, we outline the analysis of the case where the metric $g = dr^2 + m^2(r)d\theta^2$ on the 2-sphere of revolution is singular on the equator and we refer to [8] for more details. The singularity encountered coming from the Grusin model example was analysed by [1] and is described near the point identified to 0 by the local model :

$$g_s = dx^2 + \frac{dy^2}{x^2}$$

and the analysis can be extended to the case of order p :

$$g_s = dx^2 + p^2 \frac{dy^2}{x^{2p}}$$

called a *generalized Grusin singularity*.

While Riemannian metric and the Gauss curvature explode when approaching the y-axis, still the extremal curves are described by a smooth Hamiltonian system

$$H = \frac{1}{2} \left(p_x^2 + \frac{x^{2p}}{p^2} p_y^2 \right).$$

They are associated to SR-geometry in dimension 3.




FIG. 1. Conjugate and cut loci in averaged orbital transfer ($\lambda = 4$) for an initial condition different from the poles.

Indeed, the case $p = 1$ is deduced from the Heisenberg case with corresponding Hamiltonian

$$H = \frac{1}{2}[(p_x^2 + p_y^2) - 2p_z(xp_y - yp_x) + (x^2 + y^2)p_z^2]$$

and using cylindrical coordinates, we get :

$$H = \frac{1}{2}(p_r^2 + (p_\theta/r - rp_z)^2).$$

As θ is cyclic, p_θ is a first integral and for $p_\theta = 0$, the reduced Hamiltonian in the (r, z) space is with the desired singularity :

$$H = \frac{1}{2}(p_r^2 + r^2 p_z^2).$$

Similarly, the case of order 2 can be deduced by the so-called Martinet flat case, with Hamiltonian

$$H = \frac{1}{2}((p_x + \frac{y^2}{2} p_z)^2 + p_y^2).$$

As x is cyclic, p_x is a first integral and for $p_x = 0$, the reduced Hamiltonian is of the form

$$H = \frac{1}{2}(p_y^2 + \frac{y^4}{4} p_z^2).$$

In particular in both cases $p = 1, 2$ the extremal flow, the conjugate and cut loci can be deduced from the analysis of the SR-problem, see [9].




FIG. 2. Conjugate and cut loci in the singular case ($\lambda = \infty$) for an initial condition not on the equator and different from the poles.

case $p = 1$ The extremal trajectories with initial condition $x(0) = y(0) = 0$ and parameterized by arc-length are given, with $\lambda = p_y(0) \geq 0$, by :

- $\lambda = 0 : x(t) = \pm t, y(t) = 0$
- $\lambda > 0 : x(t) = \pm \frac{1}{\lambda} \sin(\lambda t), y(t) = \frac{t}{2\lambda} - \frac{1}{4\lambda^2} \sin(2\lambda t)$

while extremals for $\lambda < 0$ are by reflection with respect to the x -axis.

For $\lambda > 0$, the first conjugate time is at $t_{1c} = \tau/\lambda$, $\tau \simeq 4.5$, while due to symmetry, the optimality is lost at time π/λ , when crossing the y -axis.

case $p = 2$ The extremal trajectories with initial condition $x(0) = y(0) = 0$ and parameterized by arc-length, with $\lambda = p_y(0) \geq 0$ are :

- $\lambda = 0 : x(t) = t, y(t) = 0$
- $\lambda > 0 : x(t) = -\frac{2k}{\sqrt{\lambda}} \operatorname{cnu}, y(t) = \frac{2}{3\lambda^{3/2}} [(2k^2 - 1)(E(u) - E(K)) + k^2 t \sqrt{\lambda} + 2k^2 \sin u \operatorname{cnu} \operatorname{dnu}]$, where $u = K + t \sqrt{\lambda}$, $k^2 = k'^2 = 1/2$, and the curves deduced from the previous ones using the reflections with respect to the x and y -axis.

For $\lambda > 0$, the first conjugate time is at time $t_{1c} \simeq \frac{3K}{\sqrt{\lambda}}$, while due to symmetries the optimality is lost at time $\frac{2K}{\sqrt{\lambda}}$, when crossing the y -axis.

Hence, for both cases, we have the same geometric situation, optimality is lost due to the symmetry with respect to the y -axis and the conjugate and cut loci are disjoint, because the first conjugate point occurs after the crossing of the y -axis. It can be generalized to any order using quasi-homogeneity.

Proposition 5.1 Consider a metric of the form $g_s = dx^2 + \frac{dy^2}{x^{2p}}$. Then at the origin the conjugate and cut loci are disjoint, the cut locus is the y -axis minus 0, while the conjugate locus is the set $y = \pm c_p x^{p+1}$ minus 0.




FIG. 3. Conjugate and cut loci in the singular case ($\lambda = \infty$) for an initial condition on the equator.

Hence, the model gives locally the cut and conjugate loci of a point of the equator observed on Figure 3, which corresponds to a singularity of order 1.

Having resolved the singularity at the equator, we can extend the result of the regular case.

Definition 5.2 Consider a metric on a 2-sphere of revolution of the form $dr^2 + m^2(r)d\theta^2$, with m' nonzero on $]0, \pi/2[$ and $m(\pi - r) = m(r)$, smooth except at the equator for which we have a Grusin singularity. The extremal flow is called tame if the half-period function $\varphi(v)$ in section 3 is monotone non-increasing for $v \in]0, m(\pi/2) = +\infty[$.

Since the extremal flow remains smooth the analysis of the regular case can be extended.

Proposition 5.3 In the tame case, we have

1. The cut locus of a point different from the pole and not on the equator is a subset of the antipodal parallel.

2. The cut locus of a point on the equator is a subset of the equator accumulating at the point.

6. Conclusion

The contribution of this article is two-fold. First of all we give a simple and computable criterion to decide when the cut and conjugate loci on a surface of revolution are the simplest possible. It is based on two case studies in optimal control, in the framework of Hamiltonian dynamics. Such specialized results are important because even on a surface of revolution the computations of such objects are in general untractable. This provides a bridge between Riemannian geometry and optimal control, with promising extensions to the singular case and Zermelo navigation problem on Riemannian manifolds [2]. Secondly, we give a neat proof of the structure of the conjugate and cut loci in coplanar orbital transfer, which were previously computed in [4] using the explicit parameterization of the extremal flow for each initial point while it is sufficient to consider the half period mapping. In the tangential case they were obtained using a normal form and numerical simulations. Applications and generalizations of these results, using Hamiltonian formalism, will provide very significant improvements in the understanding of Redfield equations describing the interaction of a two-level quantum system controlled by a laser with an environment. Based on the theoretical concept and results of our analysis in the Riemannian case, this will be analyzed in a forthcoming article. Roughly speaking, if the interaction is weak, it defines a Zermelo navigation problem on a two-sphere of revolution for which conjugate and cut loci have a similar simple structure.

References

- [1] A. Agrachev, U. Boscain and M. Sigalotti, A Gauss-Bonnet like formula on two-dimensional almost Riemannian manifolds, *Preprint SISSA 55/2006/M* (2006).
- [2] D. Bao, C. Robles and Z. Shen, Zermelo navigation on Riemannian manifolds, *J. Differential Geom.* **66** (2004), no. 3, 377–435.
- [3] M. Berger, A panoramic view of Riemannian geometry, *Springer-Verlag, Berlin*, 2003.
- [4] B. Bonnard, J.-B. Caillau, Optimality results in orbit transfer, *C. R. Acad. Sci. Paris, Ser. I* **345** (2007), 319–324.
- [5] B. Bonnard, J.-B. Caillau and R. Dujol, Energy minimization of single-input orbit transfer by averaging and continuation, *Bull. Sci. Math.* **130** (2006), no. 8, 707–719.
- [6] B. Bonnard and J.-B. Caillau, Riemannian metric of the averaged energy minimization problem in orbital transfer with low thrust, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **24** (2007), no. 3, 395–411.
- [7] B. Bonnard and D. Sugny, Optimal control with applications in space and quantum dynamics, *Preprint Institut math. Bourgogne* (2007).
- [8] B. Bonnard, J. B. Caillau, R. Sinclair and M. Tanaka, One-parameter family of Clairaut-Liouville metrics with application to optimal control, *HAL preprint* No 00177686 (2007), url: hal.archives-ouvertes.fr/hal-00177686.
- [9] B. Bonnard and M. Chyba, Singular trajectories and their role in control theory, *Springer-Verlag, Berlin*, 2003.
- [10] H. Gluck and D. Singer, Scattering of geodesic fields I and II, *Ann. of Math.* **108** (1978), 347–372, and **110** (1979), 205–225.
- [11] J. Gravesen, S. Markvorsen, R. Sinclair and M. Tanaka, The cut locus of a torus of revolution, *Asian J. Math.* **9** (2005), 103–120.
- [12] J. J. Hebda, Metric structure of cut loci in surfaces and Ambrose’s problem, *J. Differential Geom.* **40** (1994), 621–642.
- [13] J. Itoh and K. Kiyohara, The cut loci and the conjugate loci on ellipsoids, *Manuscripta Math.* **114** (2004), 247–264.
- [14] K. Fleischmann, Die geodätischen Linien auf Rotationsflächen, *Liegnitz, Dissertation, Seyffarth*, 1915. Persistent URL: <http://www-gdz.sub.uni-goettingen.de/cgi-bin/digbib.cgi?PPN317487884>.
- [15] S. B. Myers, Connections between Differential Geometry and Topology: I. Simply connected surfaces and II. Closed surfaces, *Duke Math. J.* **1** (1935), 376–391, and **2** (1936), 95–102.
- [16] H. Poincaré, Sur les lignes géodésiques des surfaces convexes, *Trans. Amer. Math. Soc.* **6** (1905), 237–274.
- [17] T. Sakai, Riemannian geometry, *Transl. Math. Monogr.* **149**, American Mathematical Society, Providence, R.I., 1996.

- [18] K. Shiohama and M. Tanaka, Cut loci and distance spheres on Alexandrov surfaces, *Actes de la table ronde de géométrie différentielle en l'honneur de Marcel Berger*, 531–560, Séminaires & Congrès, Collection Soc. Math. France **1**, 1996.
- [19] K. Shiohama, T. Shioya and M. Tanaka, *The Geometry of Total Curvature on Complete Open Surfaces*, Cambridge tracts in mathematics **159**, 2003.
- [20] D. Sugny, C. Kontz and H. Jauslin, Time-optimal control of a two-level dissipative quantum system, *Phys. Rev. A* **76** 023419 (2007).
- [21] R. Sinclair and M. Tanaka, A bound on the number of endpoints of the cut locus, *LMS J. Comput. Math.* **9** (2006), 21–39.
- [22] R. Sinclair and M. Tanaka, The cut locus of a two-sphere of revolution and Toponogov's comparison theorem, *Tohoku Math. J.* **59** (2007), no. 2, 379–399.
- [23] M. Tanaka, On a characterization of a surface of revolution with many poles, *Mem. Fac. Sci. Kyushu Univ. Ser. A* **46** (1992), 251–268.