



Large deviations for the Boussinesq Equations under Random Influences

Jinqiao Duan, Annie Millet

► To cite this version:

Jinqiao Duan, Annie Millet. Large deviations for the Boussinesq Equations under Random Influences. Stochastic Processes and their Applications, 2009, 119 (6), pp.2052-2081. 10.1016/j.spa.2008.10.004 . hal-00250132v3

HAL Id: hal-00250132

<https://hal.science/hal-00250132v3>

Submitted on 7 Nov 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LARGE DEVIATIONS FOR THE BOUSSINESQ EQUATIONS UNDER RANDOM INFLUENCES

JINQIAO DUAN AND ANNIE MILLET

ABSTRACT. A Boussinesq model for the Bénard convection under random influences is considered as a system of stochastic partial differential equations. This is a coupled system of stochastic Navier-Stokes equations and the transport equation for temperature. Large deviations are proved, using a weak convergence approach based on a variational representation of functionals of infinite-dimensional Brownian motion.

1. INTRODUCTION

The need to take stochastic effects into account for modeling complex systems has now become widely recognized. Stochastic partial differential equations (SPDEs) arise naturally as mathematical models for nonlinear macroscopic dynamics under random influences. It is thus desirable to understand the impact of such random influences on the system evolution [24, 8, 20].

The Navier-Stokes equations are often coupled with other equations, especially, with the scalar transport equations for fluid density, salinity, or temperature. These coupled equations (often with the Boussinesq approximation) model a variety of phenomena in environmental, geophysical, and climate systems [9, 10, 17]. We consider the Boussinesq equations in which the scalar quantity is temperature, under different boundary conditions for the temperature at different parts (top and bottom) of the boundary. This is a Bénard convection problem. With other boundary conditions, the Boussinesq equations model various phenomena in weather and climate dynamics, for example. We take random forcings into account and formulate the Bénard convection problem as a system of stochastic partial differential equations (SPDEs). This is a coupled system of the stochastic Navier-Stokes equations and the stochastic transport equation for temperature.

In various papers about large deviation principle (LDP) for solutions u^ε to SPDEs or to evolution equations in a semi-linear framework [3, 5, 4, 6, 14, 15, 18, 21, 26], the strategy used is similar to the classical one for diffusion processes. A very general version of Schilder's theorem yields the LDP for the Gaussian noise $\sqrt{\varepsilon}W$ driving

2000 *Mathematics Subject Classification*. Primary 60H15, 60F10; Secondary 35R60, 76D05, 76R05 .

Key words and phrases. Boussinesq equations, Bénard convection, large deviations, stochastic PDEs, stochastic Navier-Stokes equations, impact of noise on system evolution, multiplicative noise.

This research was partly supported by the NSF Grant 0620539 (J. Duan) and by the research project BMF2003-01345 (A. Millet).

the stochastic forcing term, with a good rate function $\tilde{I}$ written in terms of its reproducing kernel Hilbert space (RKHS). However, since the noise is not additive, the process u^ε is not a continuous function of the noise, which creates technical difficulties. As if the contraction principle were true, one defines deterministic controlled equations u_h which are similar to the stochastic one, replacing the stochastic integral with respect to the noise $\sqrt{\varepsilon}W$ by deterministic integrals in terms of elements h of its RKHS. Once well-posedness of this controlled equation is achieved, one proves that solution u^ε to the stochastic evolution equation satisfies a LDP with a rate function I defined in terms of $\tilde{I}$ and of u_h by means of an energy minimization problem. In order to transfer the LDP from the noise to the process, there are two classical proofs, each of which contains two main steps. One way consists in proving a continuity property of the map $h \mapsto u_h$ on level sets of the rate function $\tilde{I}$ and then some Freidlin-Wentzell inequality, which states continuity of the process with respect to the noise except on an exponentially small set. Another classical method in proving LDP for evolution equations is to establish both some exponential tightness and exponentially good approximations for some approximating sequence where the diffusion coefficient is stepwise constant. These methods require some time Hölder regularity that one can obtain when the diffusion coefficient is controlled in term of the L^2 -norm of the solution, but not in the framework we will use here, where the bilinear term creates technical problems. An alternative approach [11] for large deviations is based on nonlinear semi-group theory and infinite-dimensional Hamilton-Jacobi equations, and it also requires establishing exponential tightness.

The method used in the present paper is related to the Laplace principle. One proves directly that the level sets of the rate function I are compact and then establishes weak convergence of solutions to stochastic controlled equations written in terms of the noise $\sqrt{\varepsilon}W$ shifted by a random element h_ε of its RKHS. This is again some kind of continuity property written in terms of the distributions. Unlike [22], well-posedness and a priori estimates are proved directly for very general stochastic controlled equations with a forcing term including a stochastic integral and a deterministic integral with respect to a random element h_ε of the RKHS of the noise, and for diffusion coefficients which may depend on the gradient. Indeed, if the well-posedness for the stochastic controlled equation can be deduced from that of the stochastic equation by means of a Girsanov transformation, the a priori estimates uniform in $\varepsilon > 0$, which are a key ingredient of the proof of the weak convergence result, cannot be deduced from the corresponding ones for the stochastic Bénard equation since as $\varepsilon \rightarrow 0$, the $p > 1$ moments of the Girsanov density go to infinity exponentially fast. Well-posedness has been proved in [12] for the stochastic Boussinesq equation only in the particular case of an additive noise on the velocity component. This weak convergence approach has been introduced in [1, 2]. This method has been recently applied to SPDEs [22, 25] or SDEs in infinite dimensions [19]. Finally note that the proofs of the weak convergence and compactness property require more assumptions on the diffusion coefficient σ which may not depend on the gradient. Indeed, in order to prove convergence of integrals defined in terms of

elements h_ε of the RKHS of the noise only using weak convergence of h_ε , we also need to deal with localized integral estimates of time increments. With additional assumptions on the diffusion coefficient we are able to provide complete details of the proof of this statement which was missing in [22].

This paper is organized as follows. The mathematical formulation for the stochastic Bénard model is in §2. Then the well-posedness and general a priori estimates for the model are proved in §3. Finally, a large deviation principle is shown in §4.

2. MATHEMATICAL FORMULATION

Let $D = (0, l) \times (0, 1)$ be a rectangular domain in the vertical plane. Denote by $x = (x_1, x_2)$ the spatial variable, $u = (u_1, u_2)$ the velocity field, p the pressure field, θ the temperature field, and (e_1, e_2) the standard basis in $\mathbb{R}^2$.

We consider the following stochastic coupled Navier-Stokes and heat transport equations for the Bénard convection problem [13]:

$$\frac{\partial}{\partial t} u^\varepsilon + u^\varepsilon \cdot \nabla u^\varepsilon - \nu \Delta u^\varepsilon + \nabla p^\varepsilon = \theta^\varepsilon e_2 + \sqrt{\varepsilon} n_1(t), \quad \nabla \cdot u^\varepsilon = 0, \quad (2.1)$$

$$\frac{\partial}{\partial t} \theta^\varepsilon + u^\varepsilon \cdot \nabla \theta^\varepsilon - \kappa \Delta \theta^\varepsilon = \sqrt{\varepsilon} n_2(t), \quad (2.2)$$

with boundary conditions

$$u^\varepsilon = 0 \text{ \& \; } \theta^\varepsilon = 0 \text{ on } x_2 = 0 \text{ and } x_2 = 1, \quad (2.3)$$

$$u^\varepsilon, p^\varepsilon, \theta^\varepsilon, u_{x_1}^\varepsilon, \theta_{x_1}^\varepsilon \text{ are periodic in } x_1 \text{ with period } l, \quad (2.4)$$

where n_1, n_2 are noise forcing terms and $\varepsilon > 0$ is a small parameter.

We consider the abstract functional setting for this system as in [13, 12]; see also [7, 23]. Let $L^2(D)$ be endowed with the usual scalar product and the induced norm. Consider another Hilbert space of vector-valued functions:

$$\dot{\mathbf{L}}^2(D) = \{u \in L^2(D)^2, \nabla \cdot u = 0, u|_{x_2=0} = u|_{x_2=1} = 0, u \text{ is periodic in } x_1 \text{ with period } l\}$$

$$\dot{L}^2(D) = \{\theta \in L^2(D), \theta|_{x_2=0} = \theta|_{x_2=1} = 0, \theta \text{ is periodic in } x_1 \text{ with period } l\}$$

Let $H = \dot{\mathbf{L}}^2(D) \times \dot{L}^2(D)$ be the product Hilbert space. We denote by the same notations, $(\cdot, \cdot)$ and $|\cdot|$, the scalar product and the induced norm, in $\dot{\mathbf{L}}^2(D)$, $\dot{L}^2(D)$ and H ,

$$(\phi, \psi) = \int_D \phi(x) \psi(x) dx, \quad |\phi| = \sqrt{(\phi, \phi)} = \sqrt{|\phi_1|^2 + |\phi_2|^2}.$$

Define $V = V_1 \times V_2$, where

$$V_1 = \{v \in H^1(D)^2 : \nabla \cdot v = 0, v|_{x_2=0} = v|_{x_2=1} = 0; v \text{ is periodic in } x_1 \text{ with period } l\},$$

$$V_2 = \{f \in H^1(D) : f|_{x_2=0} = f|_{x_2=1} = 0; f \text{ is periodic in } x_1 \text{ with period } l\}.$$

Then V is a product Hilbert space with the scalar product and the induced norm,

$$((\phi, \psi)) = \int_D \nabla \phi \cdot \nabla \psi dx, \quad \|\phi\| = \sqrt{((\phi, \phi))} = \sqrt{\|\phi_1\|^2 + \|\phi_2\|^2},$$

where, to ease the notation, the space variable x is omitted when writing integrals on D . Again, we also use the same notations for the scalar product and the induced norm in V_1 and V_2 . Let V' be the dual space of V . We have the dense and continuous embeddings $V \hookrightarrow H = H' \hookrightarrow V'$ and denote by $\langle \phi, \psi \rangle$ the duality between $\phi \in V$ (resp. V_i) and $\psi \in V'$ (resp. V'_i). Recall that there exists some positive constant c_1 such that for $u \in V_1$, $\theta \in V_2$,

$$|u|_{L^4(D)^2}^2 \leq c_1 |u| \|u\|, \quad \text{and} \quad |\theta|_{L^4(D)}^2 \leq c_1 |\theta| \|\theta\|. \quad (2.5)$$

Furthermore, the Poincaré inequality yields the existence of a positive constant c_2 such that

$$|\phi| \leq c_2 \|\phi\|, \quad \forall \phi \in V. \quad (2.6)$$

To lighten the notations, we will set for $\phi = (u, \theta)$, $u \in L^4$, $\theta \in L^4$ and $\phi \in L^4$ for vectors of dimensions 2, 1 and 3 whose components belong to $L^4(D)$ and denote the corresponding norms by $|\cdot|_{L^4}$.

Consider an unbounded linear operator $A = (\nu A_1, \kappa A_2) : H \rightarrow H$ with $D(A) = D(A_1) \times D(A_2)$ where $D(A_1) = V_1 \cap H^2(D)^2$, $D(A_2) = V_2 \cap H^2(D)$ and define

$$\langle A_1 u, v \rangle = ((u, v)), \quad \langle A_2 \theta, \eta \rangle = ((\theta, \eta)), \quad \forall u, v \in D(A_1), \quad \forall \theta, \eta \in D(A_2).$$

Both the Stokes operator A_1 and the Laplace operator A_2 are self-adjoint, positive, with compact self-adjoint inverses. They map V to V' . We also introduce the bilinear operators B_1 and B_2 as follows: for $u, v, w \in V_1$ and $\theta, \eta \in V_2$,

$$\begin{aligned} \langle B_1(u, v), w \rangle &= \int_D [u \cdot \nabla v] w dx := \sum_{i,j=1,2} \int_D u_i \partial_i v_j w_j dx, \\ \langle B_2(u, \theta), \eta \rangle &= \int_D [u \cdot \nabla \theta] \eta dx := \sum_{i=1,2} \int_D u_i \partial_i \theta \eta dx. \end{aligned}$$

With the notation $\phi^\varepsilon = (u^\varepsilon, \theta^\varepsilon)$ and under the above formulation, we assume that the noise terms n_1 and n_2 are respectively $\sigma_1(t, \phi) \frac{\partial}{\partial t} W^1(t)$, $\sigma_2(t, \phi) \frac{\partial}{\partial t} W^2(t)$, where $W^1(t), W^2(t)$ are independent Wiener processes defined on a filtered probability space $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P})$, taking values in $\dot{L}^2(D)$ and $\dot{L}^2(D)$, with linear symmetric positive covariant operators Q_1 and Q_2 , respectively. We denote $Q = (Q_1, Q_2)$. It is a linear symmetric positive covariant operator in the Hilbert space H . We assume that Q_1, Q_2 and thus Q are trace class (and hence compact [8]), i.e., $\text{tr}(Q) < \infty$.

As in [22], let $H_0 = Q^{\frac{1}{2}} H$. Then H_0 is a Hilbert space with the scalar product

$$(\phi, \psi)_0 = (Q^{-\frac{1}{2}} \phi, Q^{-\frac{1}{2}} \psi), \quad \forall \phi, \psi \in H_0$$

together with the induced norm $|\cdot|_0 = \sqrt{(\cdot, \cdot)_0}$. The embedding $i : H_0 \rightarrow H$ is Hilbert-Schmidt and hence compact, and moreover, $i i^* = Q$.

Let L_Q be the space of linear operators S such that $S Q^{\frac{1}{2}}$ is a Hilbert-Schmidt operator (and thus a compact operator [8]) from H to H . The norm in the space L_Q is defined by $|S|_{L_Q}^2 = \text{tr}(S Q S^*)$, where S^* is the adjoint operator of S .

Note that the above formulation is equivalent to projecting the first governing equation from $\dot{L}^2(D)^2$ into the “divergence-free” space and thus the pressure term is absent. With these notation, the above Boussinesq system (2.1)-(2.2) becomes

$$du^\varepsilon + [\nu A_1 u^\varepsilon + B_1(u^\varepsilon, u^\varepsilon) - \theta^\varepsilon e_2]dt = \sqrt{\varepsilon} \sigma_1(t, \phi^\varepsilon) dW^1(t), \quad (2.7)$$

$$d\theta^\varepsilon + [\kappa A_2 \theta^\varepsilon + B_2(u^\varepsilon, \theta^\varepsilon) - u_2^\varepsilon]dt = \sqrt{\varepsilon} \sigma_2(t, \phi^\varepsilon) dW^2(t). \quad (2.8)$$

Thus, we write this system for $\phi^\varepsilon = (u^\varepsilon, \theta^\varepsilon)$ as

$$d\phi^\varepsilon + [A\phi^\varepsilon + B(\phi^\varepsilon) + R\phi^\varepsilon]dt = \sqrt{\varepsilon} \sigma(t, \phi^\varepsilon) dW(t), \quad \phi^\varepsilon(0) = \xi := (u_0^\varepsilon, \theta_0^\varepsilon), \quad (2.9)$$

where $W(t) = (W^1(t), W^2(t))$ and

$$A\phi = (\nu A_1 u, \kappa A_2 \theta), \quad (2.10)$$

$$B(\phi) = (B_1(u, u), B_2(u, \theta)), \quad (2.11)$$

$$R\phi = (-\theta e_2, -u_2), \quad (2.12)$$

$$\sigma(t, \phi) = (\sigma_1(t, \phi), \sigma_2(t, \phi)). \quad (2.13)$$

The noise intensity $\sigma : [0, T] \times V \rightarrow L_Q(H_0, H)$ is assumed to satisfy the following:

Assumption A: There exist positive constants K and L such that

(A.1) $\sigma \in C([0, T] \times H; L_Q(H_0, H))$

(A.2) $|\sigma(t, \phi)|_{L_Q}^2 \leq K(1 + \|\phi\|^2), \quad \forall t \in [0, T], \forall \phi \in V.$

(A.3) $|\sigma(t, \phi) - \sigma(t, \psi)|_{L_Q}^2 \leq L\|\phi - \psi\|^2, \quad \forall t \in [0, T], \forall \phi, \psi \in V.$

In what follows, to ease the notation, we will suppose that $\sigma(t, \phi) = \sigma(\phi)$; however, all the results have a straightforward extension to time-dependent noise intensity under the assumption A. When no confusion arises, we set $L^p := L^p(D)$ for $1 \leq p < +\infty$ and denote by C a constant which may change from one line to the next one.

3. WELL-POSEDNESS

The goal for this paper is to show the large deviation principle for $(\phi^\varepsilon, \varepsilon > 0)$ as $\varepsilon \rightarrow 0$, where ϕ^ε denotes the solution to the stochastic Bénard equation (2.9).

Let $\mathcal{A}$ be the class of H_0 -valued $(\mathcal{F}_t)$ -predictable stochastic processes ϕ with the property $\int_0^T |\phi(s)|_0^2 ds < \infty$, a.s. Let

$$S_M = \left\{ h \in L^2(0, T; H_0) : \int_0^T |h(s)|_0^2 ds \leq M \right\}.$$

The set S_M endowed with the following weak topology is a Polish space (complete separable metric space) [2]: $d_1(h, k) = \sum_{i=1}^\infty \frac{1}{2^i} \left| \int_0^T (h(s) - k(s), \tilde{e}_i(s))_0 ds \right|$, where $\{\tilde{e}_i(s)\}_{i=1}^\infty$ is a complete orthonormal basis for $L^2(0, T; H_0)$. Define

$$\mathcal{A}_M = \{\phi \in \mathcal{A} : \phi(\omega) \in S_M, \text{ a.s.}\}. \quad (3.1)$$

As in [22], we prove existence and uniqueness of the solution to the Bénard equation. However, in what follows, we will need some precise bounds on the norm of the solution to a more general equation, which contains an extra forcing (or control) term driven by an element of $\mathcal{A}_M$. These required estimates cannot be deduced from

the corresponding ones by means of a Girsanov transformation. More precisely, let $h \in \mathcal{A}$, $\varepsilon \geq 0$ and consider the following generalized Bénard equation with initial condition $\phi_h^\varepsilon(0) = \xi$. For technical reasons, we need to add some control in the forcing term, with intensity $\tilde{\sigma} \in C([0, T] \times H; L_Q(H_0, H))$ satisfying similar stronger conditions:

Assumptions $\tilde{\mathbf{A}}$: There exist positive constants $\tilde{K}$ and $\tilde{L}$ such that:

$$(\tilde{\mathbf{A}}.1) \quad |\tilde{\sigma}(t, \phi)|_{L_Q}^2 \leq \tilde{K}(1 + |\phi|_{L^4}^2), \quad \forall t \in [0, T], \forall \phi \in L^4(D)^3.$$

$$(\tilde{\mathbf{A}}.2) \quad |\tilde{\sigma}(t, \phi) - \tilde{\sigma}(t, \psi)|_{L_Q}^2 \leq \tilde{L}|\phi - \psi|_{L^4}^2, \quad \forall t \in [0, T], \forall \phi, \psi \in L^4(D)^3.$$

Notice that since $V \subset L^4(D)^3$, the assumption $\tilde{\mathbf{A}}$ is stronger than $\mathbf{A}$. For σ , $\tilde{\sigma} \in C(H; L_Q(H_0, H))$ which satisfy Assumptions $\mathbf{A}$ and $\tilde{\mathbf{A}}$ respectively, set

$$d\phi_h^\varepsilon(t) + [A\phi_h^\varepsilon(t) + B(\phi_h^\varepsilon(t)) + R\phi_h^\varepsilon(t)]dt = \sqrt{\varepsilon}\sigma(\phi_h^\varepsilon(t))dW(t) + \tilde{\sigma}(\phi_h^\varepsilon(t))h(t)dt. \quad (3.2)$$

Recall that a stochastic process $\phi_h^\varepsilon(t, \omega)$ is called the weak solution for the generalized stochastic Bénard problem (3.2) on $[0, T]$ with initial condition ξ if ϕ_h^ε is in $C([0, T]; H) \cap L^2((0, T); V)$, a.s., and satisfies

$$\begin{aligned} & (\phi_h^\varepsilon(t), \psi) - (\xi, \psi) + \int_0^t [(\phi_h^\varepsilon(s), A\psi) + \langle B(\phi_h^\varepsilon(s)), \psi \rangle + (R\phi_h^\varepsilon(s), \psi)]ds \\ &= \sqrt{\varepsilon} \int_0^t (\sigma(\phi_h^\varepsilon(s))dW(s), \psi) + \int_0^t (\tilde{\sigma}(\phi_h^\varepsilon(s))h(s), \psi) ds, \quad a.s., \end{aligned} \quad (3.3)$$

for all $\psi \in D(A)$ and all $t \in [0, T]$. In most of the analysis here, we work in the Banach space $X := C([0, T]; H) \cap L^2((0, T); V)$ with the norm

$$\|\phi\|_X = \left\{ \sup_{0 \leq s \leq T} |\phi(s)|^2 + \int_0^T \|\phi(s)\|^2 ds \right\}^{\frac{1}{2}}. \quad (3.4)$$

Theorem 3.1. (*Well-Posedness and A priori bounds*)

Fix $M > 0$; then there exists $\varepsilon_0 := \varepsilon_0(\nu, \kappa, K, L, \tilde{K}, \tilde{L}, T, M) > 0$, such that the following existence and uniqueness result is true for $0 \leq \varepsilon \leq \varepsilon_0$. Let the initial datum satisfy $\mathbb{E}|\xi|^4 < \infty$, let $h \in \mathcal{A}_M$ and $\varepsilon \in [0, \varepsilon_0]$; then there exists a pathwise unique weak solution ϕ_h^ε of the generalized stochastic Bénard problem (3.2) with initial condition $\phi_h^\varepsilon(0) = \xi \in H$ and such that $\phi_h^\varepsilon \in X$ a.s. Furthermore, there exists a constant $C_1 := C_1(\nu, \kappa, K, L, T, M)$ such that for $\varepsilon \in [0, \varepsilon_0]$ and $h \in \mathcal{A}_M$,

$$E\|\phi_h^\varepsilon\|_X^2 \leq 1 + E\left(\sup_{0 \leq t \leq T} |\phi_h^\varepsilon(t)|^4 + \int_0^T \|\phi_h^\varepsilon(t)\|^2 dt\right) \leq C_1(1 + E|\xi|^4). \quad (3.5)$$

Remark 3.2. Note that if $\sigma = 0$, i.e., when the noise term is absent, we deduce the existence and uniqueness of the solution to the “deterministic” control equation defined in terms of an element $h \in L^2((0, T); H_0)$ and an initial condition $\xi \in H$

$$d\phi(t) + [A\phi(t) + B(\phi(t)) + R\phi(t)]dt = \tilde{\sigma}(\phi(t))h(t)dt, \quad \phi(0) = \xi. \quad (3.6)$$

If $h \in S_M$, the solution ϕ to (3.6) satisfies

$$\sup_{0 \leq s \leq T} |\phi(s)|^2 + \int_0^T \|\phi(s)\|^2 ds \leq \tilde{C}_1(\nu, \kappa, \tilde{K}, \tilde{L}, T, M, |\xi|). \quad (3.7)$$

Remark 3.3. Finally, note that when ϕ_h^ε is a solution to the stochastic Boussinesq equation (2.9), a similar argument shows that Theorem 3.1 holds for any $\varepsilon \geq 0$ if the coefficients σ (resp. $\tilde{\sigma}$) belong to $C([0, T] \times H; L_Q(H_0, H))$ and are such that in the upper estimates of the L_Q -norm appearing in the right-hand sides of conditions (A.2) and (A.3) (resp. $(\tilde{A}.1)$ and $(\tilde{A}.2)$), one replaces the V (resp. the L^4) norms of ϕ and $\phi - \psi$ by their H -norms.

Indeed, in that case, for any fixed $\varepsilon > 0$, the control of the V -norm of the solution, or of its finite-dimensional approximation, only comes from the operators A and B . Thus Lemmas 3.6 and 3.7 below prove that for α small enough, the V -norm can be dealt with.

The proof of this theorem will require several steps. The following lemmas gather some properties of B_1 and B_2 . We refer the reader to [7] or [23] for the results on B_1 which are classical and sketch some proofs of the corresponding results on B_2 .

Lemma 3.4. For $u, v, w \in V_1$ and $\theta, \eta \in V_2$,

$$\begin{aligned} \langle B_1(u, v), v \rangle &= 0, & \langle B_2(u, \theta), \theta \rangle &= 0, \\ \langle B_1(u, v), w \rangle &= -\langle B_1(u, w), v \rangle, & \langle B_2(u, \theta), \eta \rangle &= -\langle B_2(u, \eta), \theta \rangle. \end{aligned}$$

Let $u \in V_1$, $\theta \in V_2$ and $\phi = (u, \theta) \in V$; note that $|\phi|^2 = |u|^2 + |\theta|^2$ and $\|\phi\|^2 = \|u\|^2 + \|\theta\|^2$. The following lemma provides upper bound estimates of B_1 and B_2 .

Lemma 3.5. Let c_1 denote the constant in (2.5); then for any $u \in V_1$, $\theta, \eta \in V_2$ and $\phi = (u, \theta)$, one has

$$|B_1(u, u)|_{V_1'} \leq |u|_{L^4}^2 \leq c_1 |u| \|u\|, \quad (3.8)$$

$$|\langle B_2(u, \theta), \eta \rangle| \leq |u|_{L^4} |\theta|_{L^4} \|\eta\| \leq c_1 |\phi| \|\phi\| \|\eta\|. \quad (3.9)$$

Proof. We only check the properties on B_2 . For $\phi = (u, \theta) \in V$ and $\eta \in V_2$, Lemma 3.4, Hölder's inequality, and (2.5) imply

$$|\langle B_2(u, \theta), \eta \rangle| = |\langle B_2(u, \eta), \theta \rangle| \leq \|\eta\| |u|_{L^4} |\theta|_{L^4} \leq c_1 \|\eta\| |u|^{\frac{1}{2}} \|u\|^{\frac{1}{2}} |\theta|^{\frac{1}{2}} \|\theta\|^{\frac{1}{2}}.$$

This yields (3.9). $\square$

Lemma 3.6. Let $\phi = (u, \theta) \in V$, and let $v \in L^4(D)^2$ and $\eta \in L^4(D)$. For any constant $\alpha > 0$, the following estimates hold:

$$|\langle B_1(u, u), v \rangle| \leq \alpha \|u\|^2 + \frac{3^3 c_1^2}{4^4 \alpha^3} |u|^2 |v|_{L^4}^4, \quad (3.10)$$

$$|\langle B_2(\phi), \eta \rangle| \leq \alpha \|\phi\|^2 + \frac{3^3 c_1^2}{4^4 \alpha^3} |u|^2 |\eta|_{L^4}^4. \quad (3.11)$$

Proof. We only check (3.11). The first part of (3.9) and Young's inequality yield

$$\begin{aligned} |\langle B_2(\phi), \eta \rangle| &= |\langle B_2(u, \eta), \theta \rangle| \leq |\eta|_{L^4} |u|_{L^4} |\nabla \theta|_{L^2} \leq \sqrt{c_1} |\eta|_{L^4} |u|_{L^2}^{\frac{1}{2}} |\nabla u|_{L^2}^{\frac{1}{2}} |\nabla \theta|_{L^2} \\ &\leq \sqrt{c_1} |\eta|_{L^4} |u|_{L^2}^{\frac{1}{2}} \|\phi\|^{\frac{3}{2}} \leq \alpha \|\phi\|^2 + \frac{3^3 c_1^2}{4^4 \alpha^3} |u|^2 |\eta|_{L^4}^4. \end{aligned}$$

$\square$

The following lemma allows rewriting differences of B_i for $i = 1, 2$ and deducing estimates for the difference of B .

Lemma 3.7. *Let $\phi = (u, \theta)$ and $\psi = (v, \eta)$ belong to V . Then*

$$\begin{aligned}\langle B_1(u, u) - B_1(v, v), u - v \rangle &= -\langle B_1(u - v, u - v), v \rangle, \\ \langle B_2(\phi) - B_2(\psi), \theta - \eta \rangle &= -\langle B_2(\phi - \psi), \eta \rangle.\end{aligned}$$

Furthermore, for some constant $c > 0$ and for any constant $\alpha > 0$,

$$|\langle B(\phi) - B(\psi), \phi - \psi \rangle| \leq c \|\phi - \psi\| \|\phi - \psi\| \|\psi\| \quad (3.12)$$

$$\leq \alpha \|\phi - \psi\|^2 + \frac{3^3 c^2}{2^4 \alpha^3} \|\phi - \psi\|^2 \|\psi\|_{L^4}^4. \quad (3.13)$$

Proof. Integration by parts, the boundary conditions and $\operatorname{div}(u) = \nabla \cdot u = 0$ yield

$$\begin{aligned}\langle B_2(\phi) - B_2(\psi), \theta - \eta \rangle &= \int_D (u \cdot \nabla \theta)(\theta - \eta) dx - \int_D (v \cdot \nabla \eta)(\theta - \eta) dx \\ &= - \int_D (u \cdot \nabla(\theta - \eta)) \theta dx + \int_D (v \cdot \nabla(\theta - \eta)) \eta dx\end{aligned}$$

Since $\langle B_2(u, w), w \rangle = \int_D (u \cdot \nabla w) w dx = 0$ for any $w \in V_2$, we deduce that

$$\langle B_2(\phi) - B_2(\psi), \theta - \eta \rangle = - \int_D (u \cdot \nabla(\theta - \eta)) \eta dx + \int_D (v \cdot \nabla(\theta - \eta)) \eta dx,$$

which completes the proof of the second identity. The proof of the first one, which is similar and classical, is omitted. Finally, combining these identities with the upper estimates in Lemmas 3.5 and 3.6 concludes the proof. $\square$

For $\phi = (u, \theta) \in V$, define

$$F(\phi) = -A\phi - B(\phi) - R\phi. \quad (3.14)$$

We at first prove crucial monotonicity properties of F . Let $\nu \wedge \kappa := \min(\nu, \kappa)$.

Lemma 3.8. *Assume that $\phi = (u, \theta) \in V$ and $\psi = (v, \eta) \in V$; then for some constant $c > 0$ we have*

$$\langle F(\phi) - F(\psi), \phi - \psi \rangle + (\nu \wedge \kappa) \|\phi - \psi\|^2 \leq c \|\phi - \psi\| \|\phi - \psi\| \|\psi\| + \|\phi - \psi\|^2. \quad (3.15)$$

Proof. Set $U := u - v$, $\Theta := \theta - \eta$ and $\Phi = \phi - \psi = (U, \Theta)$. Integrating by parts we deduce from Lemma 3.7

$$\langle F(\phi) - F(\psi), \Phi \rangle = -\nu \|U\|^2 - \kappa \|\Theta\|^2 - \langle B_1(U, U), v \rangle - \langle B_2(\Phi), \eta \rangle + 2\langle U_2, \Theta \rangle.$$

Thus (3.12) yields (3.15). $\square$

The proof of Theorem 3.1 involves Galerkin approximations. Let $\{\varphi_n\}_{n \geq 1}$ be a complete orthonormal basis of the Hilbert space H such that $\varphi_n \in \operatorname{Dom}(A)$, domain of definition of the operator A . For any $n \geq 1$, let $H_n = \operatorname{span}(\varphi_1, \dots, \varphi_n) \subset \operatorname{Dom}(A)$ and $P_n : H \rightarrow H_n$ denote the orthogonal projection onto H_n . Note that P_n contracts the H and V norms and that its norm as a linear operator of $L^4(D)^3$

is bounded in n . Suppose that the H -valued Wiener process W with covariance operator Q is such that

$$P_n Q^{\frac{1}{2}} = Q^{\frac{1}{2}} P_n, \quad n \geq 1,$$

which is true if $Qh = \sum_{n \geq 1} \lambda_n \varphi_n$ with trace $\sum_{n \geq 1} \lambda_n < \infty$. Then for $H_0 = Q^{\frac{1}{2}} H$ and $(\phi, \psi)_0 = (Q^{-\frac{1}{2}} \phi, Q^{-\frac{1}{2}} \psi)$ given $\phi, \psi \in H_0$, we see that $P_n : H_0 \rightarrow H_0 \cap H_n$ is a contraction both of the H and H_0 norms. Let $W_n = P_n W$, $\sigma_n = P_n \sigma$ and $\tilde{\sigma}_n = P_n \tilde{\sigma}$.

For $h \in \mathcal{A}_M$, consider the following stochastic ordinary differential equation on the n -dimensional space H_n defined by

$$d(\phi_{n,h}^\varepsilon, \psi) = [\langle F(\phi_{n,h}^\varepsilon), \psi \rangle + (\tilde{\sigma}_n(\phi_{n,h}^\varepsilon)h, \psi)] dt + \sqrt{\varepsilon} (\sigma_n(\phi_{n,h}^\varepsilon) dW_n, \psi), \quad (3.16)$$

for $\psi = (v, \eta) \in H_n$ and $\phi_{n,h}^\varepsilon(0) = P_n \xi$.

Note that for $\psi = (v, \eta) \in V$, the map $\phi \in H_n \mapsto \langle (A + R)(\phi), \psi \rangle$ is globally Lipschitz, while using Lemma 3.5 the map $\phi = (u, \theta) \in H_n \mapsto \sum_{i,j=1,2} \int_D u_i v_j \partial_i u_j dx + \sum_{i=1,2} \int_D u_i \eta \partial_i \theta dx$ is locally Lipschitz. Furthermore, conditions **(A.3)** and **(A.2)** imply that the maps $\phi \in H_n \rightarrow \sigma_n(\phi)$ and $\phi \in H_n \rightarrow \tilde{\sigma}_n(\phi)$ are globally Lipschitz from H_n to $n \times n$ matrices. Hence by a well-posedness result for stochastic ordinary differential equations [16], there exists a maximal solution to (3.16), i.e., a stopping time $\tau_{n,h}^\varepsilon \leq T$ such that (3.16) holds for $t < \tau_{n,h}^\varepsilon$ and as $t \uparrow \tau_{n,h}^\varepsilon < T$, $|\phi_{n,h}^\varepsilon(t)| \rightarrow \infty$. For every $N > 0$, set

$$\tau_N = \inf\{t : |\phi_{n,h}^\varepsilon(t)| \geq N\} \wedge T. \quad (3.17)$$

Almost surely, $\phi_{n,h}^\varepsilon \in C([0, T], H_n)$ on $\{\tau_N = T\}$. The following proposition shows that $\tau_{n,h}^\varepsilon = T$ a.s. and gives estimates on $\phi_{n,h}^\varepsilon$ depending only on the physical constants ν and κ , K , $\tilde{K}$, T , M , $\mathbb{E}|\xi|^{2p}$ which are valid for all n and all $\varepsilon \in [0, \varepsilon_0]$ for some $\varepsilon_0 > 0$. Its proof depends on the following version of Gronwall's lemma.

Lemma 3.9. *Let X , Y and I be non-decreasing, non-negative processes, φ be a non-negative process and Z be a non-negative integrable random variable. Assume that $\int_0^T \varphi(s) ds \leq C$ almost surely and that there exist positive constants $\alpha, \beta \leq \frac{1}{2(1+Ce^C)}$, $\gamma \leq \frac{\alpha}{2(1+Ce^C)}$ and $\tilde{C} > 0$ such that for $0 \leq t \leq T$,*

$$X(t) + \alpha Y(t) \leq Z + \int_0^t \varphi(r) X(r) dr + I(t), \quad \text{a.s.} \quad (3.18)$$

$$\mathbb{E}(I(t)) \leq \beta \mathbb{E}(X(t)) + \gamma \mathbb{E}(Y(t)) + \tilde{C}. \quad (3.19)$$

Then if $X \in L^\infty([0, T] \times \Omega)$, we have for $t \in [0, T]$

$$\mathbb{E}[X(t) + \alpha Y(t)] \leq 2(1 + Ce^C)(\mathbb{E}(Z) + \tilde{C}). \quad (3.20)$$

Proof. Iterating inequality (3.18) and ignoring Y , an induction argument on n yields for $t \in [0, T]$, $n \geq 1$

$$X(t) \leq Z + \int_0^t \varphi(s_1) \left[Z + \int_0^{s_1} \varphi(s_2) X(s_2) ds_2 + I(s_1) \right] ds_1 + I(t)$$

$$\begin{aligned}
&\leq Z + I(t) + \sum_{1 \leq k \leq n} \int_0^t \varphi(s_1) \int_0^{s_1} \varphi(s_2) \cdots \int_0^{s_{k-1}} \varphi(s_k) [Z + I(s_k)] ds_k \cdots ds_1 \\
&\quad + \int_0^t \varphi(s_1) \int_0^{s_1} \varphi(s_2) \cdots \int_0^{s_n} \varphi(s_{n+1}) X(s_{n+1}) ds_{n+1} ds_n \cdots ds_1.
\end{aligned}$$

Recall that $X(s, \omega)$ is a.e. bounded and $\int_0^T \varphi(s) ds \leq C$; thus $X(t) \leq e^C [Z + I(t)]$. Using this inequality in (3.18) and the fact that I is non-decreasing, we deduce that $X(t) + \alpha Y(t) \leq [Z + I(t)] (1 + Ce^C)$. Taking expected values and using (3.19), we conclude the proof. $\square$

Proposition 3.10. *There exists $\varepsilon_{0,p} := \varepsilon_{0,p}(\nu, \kappa, K, \tilde{K}, T, M)$ such that for $0 \leq \varepsilon \leq \varepsilon_{0,p}$ the following result holds for an integer $p \geq 1$ (with the convention $x^0 = 1$). Let $h \in \mathcal{A}_M$ and $\xi \in L^{2p}(\Omega, H)$. Then $\tau_{n,h} = T$ a.s. and equation (3.16) has a unique solution with a modification $\phi_{n,h}^\varepsilon \in C([0, T], H_n)$ and satisfying*

$$\begin{aligned}
&\sup_n \mathbb{E} \left(\sup_{0 \leq t \leq T} |\phi_{n,h}^\varepsilon(t)|^{2p} + \int_0^T \|\phi_{n,h}^\varepsilon(s)\|^2 |\phi_{n,h}^\varepsilon(s)|^{2(p-1)} ds \right) \\
&\leq C(p, \nu, \kappa, K, \tilde{K}, T, M) (\mathbb{E}|\xi|^{2p} + 1).
\end{aligned} \tag{3.21}$$

Proof. Itô's formula yields that for $t \in [0, T]$ and τ_N defined by (3.17),

$$|\phi_{n,h}^\varepsilon(t \wedge \tau_N)|^2 = |P_n \xi|^2 + 2\sqrt{\varepsilon} \int_0^{t \wedge \tau_N} (\sigma_n(\phi_{n,h}^\varepsilon(s)) dW_n(s), \phi_{n,h}^\varepsilon(s)) \tag{3.22}$$

$$\begin{aligned}
&+ 2 \int_0^{t \wedge \tau_N} \langle F(\phi_{n,h}^\varepsilon(s)), \phi_{n,h}^\varepsilon(s) \rangle ds + 2 \int_0^{t \wedge \tau_N} (\tilde{\sigma}_n(\phi_{n,h}^\varepsilon(s)) h(s), \phi_{n,h}^\varepsilon(s)) ds \\
&+ \varepsilon \int_0^{t \wedge \tau_N} |\sigma_n(\phi_{n,h}^\varepsilon(s)) P_n|_{L_Q}^2 ds.
\end{aligned} \tag{3.23}$$

Apply again Itô's formula for x^p when $p \geq 2$ and then use Lemma 3.4. With the convention $p(p-1)x^{p-2} = 0$ for $p = 1$, this yields for $t \in [0, T]$,

$$\begin{aligned}
&|\phi_{n,h}^\varepsilon(t \wedge \tau_N)|^{2p} + 2p \int_0^{t \wedge \tau_N} |\phi_{n,h}^\varepsilon(r)|^{2(p-1)} [\nu \|u_{n,h}^\varepsilon(r)\|^2 + \kappa \|\theta_{n,h}^\varepsilon(r)\|^2] dr \\
&\leq |P_n \xi|^{2p} + \sum_{1 \leq j \leq 5} T_j(t),
\end{aligned} \tag{3.24}$$

where

$$\begin{aligned}
T_1(t) &= 4p \int_0^{t \wedge \tau_N} |(\theta_{n,h}^\varepsilon(r), u_{n,h,2}^\varepsilon(r))| |\phi_{n,h}^\varepsilon(r)|^{2(p-1)} dr, \\
T_2(t) &= 2p\sqrt{\varepsilon} \left| \int_0^{t \wedge \tau_N} (\sigma_n(\phi_{n,h}^\varepsilon(r)) dW_n(r), \phi_{n,h}^\varepsilon(r)) |\phi_{n,h}^\varepsilon(r)|^{2(p-1)} \right|, \\
T_3(t) &= 2p \int_0^{t \wedge \tau_N} |(\tilde{\sigma}_n(\phi_{n,h}^\varepsilon(r)) h(r), \phi_{n,h}^\varepsilon(r))| |\phi_{n,h}^\varepsilon(r)|^{2(p-1)} dr,
\end{aligned}$$

$$\begin{aligned}
T_4(t) &= p\varepsilon \int_0^{t \wedge \tau_N} |\sigma_n(\phi_{n,h}^\varepsilon(r)) P_n|_{L_Q}^2 |\phi_{n,h}^\varepsilon(r)|^{2(p-1)} dr, \\
T_5(t) &= 2p(p-1)\varepsilon \int_0^{t \wedge \tau_N} |\Pi_n \sigma_n^*(\phi_{n,h}^\varepsilon(r)) \phi_{n,h}^\varepsilon(r)|_{H_0}^2 |\phi_{n,h}^\varepsilon(r)|^{2(p-2)} dr.
\end{aligned}$$

The Cauchy-Schwarz inequality implies that $2|(\theta_{n,h}^\varepsilon(r), u_{n,h,2}^\varepsilon(r))| \leq |\phi_{n,h}^\varepsilon(r)|^2$. Hence

$$T_1(t) \leq 2p \int_0^{t \wedge \tau_N} |\phi_{n,h}^\varepsilon(r)|^{2p} dr. \quad (3.25)$$

Since $h \in \mathcal{A}_M$, the Cauchy-Schwarz inequality, (A.2), (2.5) and the Poincaré inequality (2.6) imply the existence of some positive constant c such that for every $\delta_1 > 0$,

$$\begin{aligned}
T_3(t) &\leq 2p \int_0^{t \wedge \tau_N} [\tilde{K}(1 + c \|\phi_{n,h}^\varepsilon(r)\|^2)]^{\frac{1}{2}} |h(r)|_0 |\phi_{n,h}^\varepsilon(r)|^{2p-1} dr \\
&\leq \delta_1 \int_0^{t \wedge \tau_N} \|\phi_{n,h}^\varepsilon(r)\|^2 |\phi_{n,h}^\varepsilon(r)|^{2(p-1)} dr + \frac{p^2 \tilde{K} c}{\delta_1} \int_0^{t \wedge \tau_N} |h(r)|_0^2 |\phi_{n,h}^\varepsilon(r)|^{2p} dr \\
&\quad + \delta_1 \int_0^{t \wedge \tau_N} |\phi_{n,h}^\varepsilon(r)|^{2(p-1)} dr.
\end{aligned} \quad (3.26)$$

Using (A.2), we deduce that

$$\begin{aligned}
T_4(t) + T_5(t) &\leq 2p^2 K \varepsilon \int_0^{t \wedge \tau_N} \|\phi_{n,h}^\varepsilon(r)\|^2 |\phi_{n,h}^\varepsilon(r)|^{2(p-1)} dr \\
&\quad + 2p^2 K \varepsilon \int_0^{t \wedge \tau_N} |\phi_{n,h}^\varepsilon(r)|^{2(p-1)} dr.
\end{aligned} \quad (3.27)$$

Finally, the Burkholder-Davies-Gundy inequality, (A.2) and Schwarz's inequality yield that for $t \in [0, T]$ and $\delta_2 > 0$,

$$\begin{aligned}
\mathbb{E} \left(\sup_{0 \leq s \leq t} |T_2(s)| \right) &\leq 6p\sqrt{\varepsilon} \mathbb{E} \left\{ \int_0^{t \wedge \tau_N} |\phi_{n,h}^\varepsilon(r)|^{2(2p-1)} |\sigma_{n,h}(\phi_{n,h}^\varepsilon(r)) P_n|_{L_Q}^2 dr \right\}^{\frac{1}{2}} \\
&\leq \delta_2 \mathbb{E} \left(\sup_{0 \leq s \leq t \wedge \tau_N} |\phi_{n,h}^\varepsilon(s)|^{2p} \right) + \frac{9p^2 K \varepsilon}{\delta_2} \mathbb{E} \int_0^{t \wedge \tau_N} |\phi_{n,h}^\varepsilon(r)|^{2(p-1)} dr \\
&\quad + \frac{9p^2 K \varepsilon}{\delta_2} \mathbb{E} \int_0^{t \wedge \tau_N} \|\phi_{n,h}^\varepsilon(r)\|^2 |\phi_{n,h}^\varepsilon(r)|^{2(p-1)} dr.
\end{aligned} \quad (3.28)$$

Consider the following property $I(i)$ for an integer $i \geq 0$:

I(i) There exists $\varepsilon_{0,i} := \varepsilon_{0,i}(\nu, \kappa, K, \tilde{K}, T, M) > 0$ such that for $0 \leq \varepsilon \leq \varepsilon_{0,i}$

$$\sup_n \mathbb{E} \int_0^{t \wedge \tau_N} |\phi_{n,h}^\varepsilon(r)|^{2i} dr \leq C(i) := C(i, \nu, \kappa, K, \tilde{K}, T, M) < +\infty.$$

The property $I(0)$ obviously holds with $\varepsilon_{0,0} = 1$ and $C(0) = T$. Assume that for some integer i with $1 \leq i \leq p$, the property I(i-1) holds; we prove that I(i) holds.

Set $\delta_1 = \frac{(\nu \wedge \kappa)^i}{2}$, $\varphi_i(r) = 2i + \frac{i^2 c \tilde{K}}{\delta_1} |h(r)|_0^2$, $Z = \delta_1 \int_0^{\tau_N} |\phi_{n,h}^\varepsilon(r)|^{2(i-1)} dr + |\xi|^{2i}$, $X(t) = \sup_{0 \leq s \leq t} |\phi_{n,h}^\varepsilon(s \wedge \tau_N)|^{2i}$, $Y(t) = \int_0^{t \wedge \tau_N} \|\phi_{n,h}^\varepsilon(s)\|^2 |\phi_{n,h}^\varepsilon(s)|^{2(i-1)} ds$ and $I(t) = \sup_{0 \leq s \leq t} 2i\sqrt{\varepsilon} \left| \int_0^{t \wedge \tau_N} (\sigma_n(\phi_{n,h}^\varepsilon(r)) dW_n(r), \phi_{n,h}^\varepsilon(r)) |\phi_{n,h}^\varepsilon(r)|^{2(i-1)} \right|$. Then $\int_0^T \varphi_i(s) ds \leq C_i(M) := 2iT + \frac{i^2 c \tilde{K}}{\delta_1} M$. Let $\alpha = i(\nu \wedge \kappa)$, $\beta = \delta_2 = \frac{1}{2[1+C_i(M)e^{C_i(M)}]}$ and $\tilde{C} = \frac{9i^2 K}{\delta_2} \mathbb{E} \int_0^{\tau_N} |\phi_{n,h}^\varepsilon(s)|^{2(i-1)} ds$. Let

$$\varepsilon_{0,i} = 1 \wedge \frac{\nu \wedge \kappa}{8iK} \wedge \frac{\nu \wedge \kappa}{144iK[1+C_i(M)e^{C_i(M)}]^2} \wedge \varepsilon_{0,i-1}.$$

Then for $0 \leq \varepsilon \leq \varepsilon_{0,i}$ inequalities (3.24)-(3.28) show that the assumptions of Lemma 3.9 hold with $\gamma = \frac{9i^2 K \varepsilon}{\delta_2} \leq \alpha\beta$, which yields I(i).

An induction argument shows that $I(p-1)$ holds, and hence the previous computations with $i = p$ and Lemma 3.9 yield that for $t = T$ and $0 \leq \varepsilon \leq \varepsilon_{0,p}$,

$$\sup_n \mathbb{E} \left(\sup_{0 \leq s \leq \tau_N} |\phi_{n,h}^\varepsilon(s)|^{2p} + \int_0^{\tau_N} \|\phi_{n,h}^\varepsilon(s)\|^2 |\phi_{n,h}^\varepsilon(s)|^{2(p-1)} ds \right) \leq C(p, \nu, \kappa, K, \tilde{K}, T, M).$$

As $N \rightarrow \infty$, $\tau_N \uparrow \tau_{n,h}$ and on $\{\tau_{n,h} < T\}$, $\sup_{0 \leq s \leq t \wedge \tau_N} |\phi_{n,h}(s)| \rightarrow \infty$. Hence $\mathbb{P}(\tau_{n,h} < T) = 0$ and for almost all ω , for $N(\omega)$ large enough, $\tau_{N(\omega)}(\omega) = T$ and $\phi_{n,h}(\cdot)(\omega) \in C([0, T], H_n)$. By the Lebesgue monotone convergence theorem, we complete the proof of the proposition. $\square$

We now have the following bound in $L^4(D)^3$.

Proposition 3.11. *Let $h \in \mathcal{A}_M$ and $\xi \in L^4(\Omega, H)$. Let $\varepsilon_{0,2}$ be defined as in Proposition 3.10 with $p = 2$. Then there exists a constant $C_2 := C_2(\nu, \kappa, K, \tilde{K}, T, M)$ such that for $0 \leq \varepsilon \leq \varepsilon_{0,2}$,*

$$\sup_n \mathbb{E} \int_0^T |\phi_{n,h}^\varepsilon(s)|_{L^4}^4 ds \leq C_2(1 + \mathbb{E}|\xi|^4). \quad (3.29)$$

Proof. Let $f_{n,h}(t) = u_{n,h,i}(t)$ or $\theta_{n,h}^\varepsilon(t)$, with $i = 1, 2$. Then (3.21) with $p = 2$ implies that

$$\sup_n \mathbb{E} \int_0^T \|f_{n,h}(s)\|^2 |f_{n,h}(s)|^2 ds \leq C_2(\nu, \kappa, K, \tilde{K}, T, M)(1 + \mathbb{E}|\xi|^4).$$

Hence by the second part of (3.8), we finish the proof of (3.29). $\square$

The following result is a consequence of Itô's formula; it will be used in what follows for various choices of coefficients.

Lemma 3.12. *Let $\xi \in L^4(\Omega, H)$ be $\mathcal{F}_0$ -measurable, $\rho' : [0, T] \times \Omega \rightarrow [0 + \infty[$ be adapted such that for almost every ω the map $t \rightarrow \rho'(t, \omega) \in L^1([0, T])$ and for $t \in [0, T]$, set $\rho(t) = \int_0^t \rho'(s) ds$. For $i = 1, 2$, let σ_i satisfy assumption (A.1), $\bar{\sigma}_i \in C([0, T] \times H, L_Q^2)$ and let $\bar{\sigma}$ satisfy Assumption $\tilde{\mathbf{A}}$. Let F satisfy condition (3.15),*

$h_\varepsilon \in \mathcal{A}_M$ and $\phi_i \in L^2([0, T], V) \cap L^\infty([0, T], H)$ a.s. and be such that $\phi_i(0) = \xi$ and satisfy the equation

$$d\phi_i(t) = F(\phi_i(t))dt + \sqrt{\varepsilon}\sigma_i(t, \phi_i(t))dW(t) + (\bar{\sigma}(t, \phi_i(t))h_\varepsilon(t) + \bar{\sigma}_i(t))dt. \quad (3.30)$$

Let $\Phi = \phi_1 - \phi_2$ and c_1 and c_2 denote the constants in (2.5) and (2.6) respectively. Then for every $t \in [0, T]$,

$$\begin{aligned} e^{-\rho(t)} |\Phi(t)|^2 &\leq \int_0^t e^{-\rho(s)} \left\{ -(\nu \wedge \kappa) \|\Phi(s)\|^2 + \varepsilon |\sigma_1(s, \phi_1(s)) - \sigma_2(s, \phi_2(s))|_{L_Q^2}^2 \right. \\ &\quad \left. + |\Phi(s)|^2 \left[-\rho'(s) + 2 + \frac{8c_1^2}{\nu \wedge \kappa} \|\phi_2(s)\|^2 + \frac{2\tilde{L}c_1c_2}{\nu \wedge \kappa} |h_\varepsilon(s)|_0^2 \right] \right\} ds \\ &\quad + 2 \int_0^t e^{-\rho(s)} (\bar{\sigma}_1(s) - \bar{\sigma}_2(s), \Phi(s)) ds + I(t), \end{aligned} \quad (3.31)$$

where $I(t) = 2\sqrt{\varepsilon} \int_0^t e^{-\rho(s)} ([\sigma_1(s, \phi_1(s)) - \sigma_2(s, \phi_2(s))] dW(s), \Phi(s))$.

Proof. Itô's formula, (3.15) and condition ($\tilde{\mathbf{A}}.2$) imply that for $t \in [0, T]$,

$$\begin{aligned} e^{-\rho(t)} |\Phi(t)|^2 &= \int_0^t e^{-\rho(s)} \left\{ -\rho'(s) |\Phi(s)|^2 + \varepsilon |\sigma_1(s, \phi_1(s)) - \sigma_2(s, \phi_2(s))|_{L_Q^2}^2 \right. \\ &\quad \left. + 2 \langle F(\phi_1(s)) - F(\phi_2(s)), \Phi(s) \rangle + 2([\bar{\sigma}(s, \phi_1(s)) - \bar{\sigma}(s, \phi_2(s))] h_\varepsilon(s), \Phi(s)) \right\} ds \\ &\quad + \int_0^t e^{-\rho(s)} 2(\bar{\sigma}_1(s) - \bar{\sigma}_2(s), \Phi(s)) ds + I(t) \\ &\leq \int_0^t e^{-\rho(s)} \left\{ -\rho'(s) |\Phi(s)|^2 + \varepsilon |\sigma_1(s, \phi_1(s)) - \sigma_2(s, \phi_2(s))|_{L_Q^2}^2 - 2(\nu \wedge \kappa) \|\Phi(s)\|^2 \right. \\ &\quad \left. + 4c_1 |\Phi(s)| \|\Phi(s)\| \|\phi_2(s)\| + 2|\Phi(s)|^2 + 2\sqrt{\tilde{L}c_1c_2} \|\Phi(s)\| |h_\varepsilon(s)|_0 |\Phi(s)| \right\} ds \\ &\quad + \int_0^t e^{-\rho(s)} 2(\bar{\sigma}_1(s) - \bar{\sigma}_2(s), \Phi(s)) ds + I(t). \end{aligned}$$

The inequalities $4c_1 |\Phi(s)| \|\Phi(s)\| \|\phi_2(s)\| \leq \frac{(\nu \wedge \kappa)}{2} \|\Phi(s)\|^2 + \frac{8c_1^2}{\nu \wedge \kappa} \|\phi_2(s)\|^2 |\Phi(s)|^2$ and $2\sqrt{\tilde{L}c_1c_2} \|\Phi(s)\| |h_\varepsilon(s)|_0 |\Phi(s)| \leq \frac{(\nu \wedge \kappa)}{2} \|\Phi(s)\|^2 + \frac{2\tilde{L}c_1c_2}{\nu \wedge \kappa} |h_\varepsilon(s)|_0^2 |\Phi(s)|^2$ conclude the proof of (3.31). $\square$

We are now ready to prove the main result of this section.

Proof of Theorem 3.1:

Let $\Omega_T = [0, T] \times \Omega$ be endowed with the product measure $ds \otimes d\mathbb{P}$ on $\mathcal{B}([0, T]) \otimes \mathcal{F}$. Let $\varepsilon_{0,2}$ be defined by Proposition 3.10 with $p = 2$ and set $\varepsilon_0 := \varepsilon_{0,2} \wedge \frac{\nu \wedge \kappa}{2L}$. The proof consists of several steps.

Step 1: Inequalities (3.21) and (3.29) imply the existence of a subsequence of $\{\phi_{n,h}^\varepsilon\}_{n \geq 0}$ (still denoted by the same notation), of processes $\phi_h^\varepsilon \in L^2(\Omega_T, V) \cap L^4(\Omega_T, L^4(D)^3) \cap L^4(\Omega, L^\infty([0, T], H))$, $F_h^\varepsilon \in L^2(\Omega_T, V')$, $S_h^\varepsilon, \tilde{S}_h^\varepsilon \in L^2(\Omega_T, L_Q)$, and of random variables $\check{\phi}_h^\varepsilon(T) \in L^2(\Omega, H)$, for which the following properties hold:

- (i) $\phi_{n,h}^\varepsilon \rightarrow \phi_h^\varepsilon$ weakly in $L^2(\Omega_T, V)$,
- (ii) $\phi_{n,h}^\varepsilon \rightarrow \phi_h^\varepsilon$ weakly in $L^4(\Omega_T, L^4(D)^3)$,
- (iii) $\phi_{n,h}^\varepsilon$ is weak star converging to ϕ_h^ε in $L^4(\Omega, L^\infty([0, T], H))$,
- (iv) $\phi_{n,h}^\varepsilon(T) \rightarrow \tilde{\phi}_h^\varepsilon(T)$ weakly in $L^2(\Omega, H)$,
- (v) $F(\phi_{n,h}^\varepsilon) \rightarrow F_h^\varepsilon$ weakly in $L^2(\Omega_T, V')$,
- (vi) $\sigma_n(\phi_{n,h}^\varepsilon)P_n \rightarrow S_h^\varepsilon$ weakly in $L^2(\Omega_T, L_Q)$,
- (vii) $\tilde{\sigma}_n(\phi_{n,h}^\varepsilon)h \rightarrow \tilde{S}_h^\varepsilon$ weakly in $L^{\frac{4}{3}}(\Omega_T, H)$.

Indeed, (i)-(iv) are straightforward consequences of Propositions 3.10 and 3.11, and of uniqueness of the limit of $\mathbb{E} \int_0^T \phi_{n,h}^\varepsilon(t) \psi(t) dt$ for appropriate ψ .

Furthermore, given $\psi = (v, \eta) \in L^2(\Omega_T, V)$, we have

$$\begin{aligned}
& \mathbb{E} \int_0^T \left[\nu \langle A_1(u_{n,h}^\varepsilon(t), v(t)) \rangle + \kappa \langle A_2(\theta_{n,h}^\varepsilon(t)), \eta(t) \rangle \right] dt \\
&= -\nu \mathbb{E} \int_0^T (\nabla u_{n,h}^\varepsilon(t), \nabla v(t)) dt - \kappa \mathbb{E} \int_0^T (\nabla \theta_{n,h}^\varepsilon(t), \nabla \eta(t)) dt \\
&\rightarrow -\nu \mathbb{E} \int_0^T (\nabla u_h^\varepsilon(t), \nabla v(t)) dt - \kappa \mathbb{E} \int_0^T (\nabla \theta_h^\varepsilon(t), \nabla \eta(t)) dt. \tag{3.32}
\end{aligned}$$

Using (3.21) with $p = 2$, (3.8), (3.9), the Cauchy-Schwarz and Poincaré inequalities, we deduce

$$\begin{aligned}
& \sup_n \mathbb{E} \int_0^T \left| \langle B_1(u_{n,h}^\varepsilon(t), u_{n,h}^\varepsilon(t)), v(t) \rangle + \langle B_2(\phi_{n,h}^\varepsilon(t)), \eta(t) \rangle + \langle R\phi_{n,h}^\varepsilon(t), \psi(t) \rangle \right| dt \\
&\leq C \sup_n \mathbb{E} \int_0^T \left\{ \|u_{n,h}^\varepsilon(t)\| |u_{n,h}^\varepsilon(t)| \|v(t)\| + \|\phi_{n,h}^\varepsilon(t)\| |\phi_{n,h}^\varepsilon(t)| \|\eta(t)\| \right. \\
&\quad \left. + |\theta_{n,h}^\varepsilon(t)| |v_2(t)| + |u_{n,h,2}^\varepsilon(t)| |\eta(t)| \right\} dt \\
&\leq C_3(\nu, \kappa, K, T, M)(1 + E|\xi|^4) + \mathbb{E} \int_0^T \|\psi(t)\|^2 dt.
\end{aligned}$$

Hence $\{B(\phi_{n,h}^\varepsilon(t)) + R\phi_{n,h}^\varepsilon(t), n \geq 1\}$ has a subsequence converging weakly in $L^2(\Omega_T, V')$. This convergence and (3.32) prove (v).

Since P_n contracts the $|\cdot|_0$ and $|\cdot|$ norms, (A.2) and (3.21) imply that

$$\sup_n \mathbb{E} \int_0^T |\sigma_n(\phi_{n,h}^\varepsilon(t))P_n|_{L_Q}^2 dt \leq K \sup_n \mathbb{E} \int_0^T (1 + \|\phi_{n,h}^\varepsilon(t)\|^2) dt < \infty,$$

which proves (vi). Finally, using Assumption (A.1), Hölder's inequality and (3.29), we deduce that for $h \in \mathcal{A}_M$, for any $n \geq 1$,

$$\begin{aligned}
\mathbb{E} \int_0^T |\tilde{\sigma}_n(\phi_{n,h}^\varepsilon(s)) h(s)|_H^{\frac{4}{3}} ds &\leq \mathbb{E} \int_0^T [\tilde{K}(1 + |\phi_{n,h}^\varepsilon(s)|_{L^4}^2)]^{\frac{2}{3}} |h(s)|_0^{\frac{4}{3}} ds \\
&\leq \tilde{K}^{\frac{4}{3}} \left(\mathbb{E} \int_0^T |h(s)|_0^2 ds \right)^{\frac{2}{3}} \left(\mathbb{E} \int_0^T [1 + |\phi_{n,h}^\varepsilon(s)|_{L^4}^2] ds \right)^{\frac{1}{3}}
\end{aligned}$$

$$\leq C(M, T, K, \tilde{K}, \nu, \kappa).$$

This completes the proof of (vii).

Step 2: For $\delta > 0$, let $f \in H^1(-\delta, T + \delta)$ be such that $\|f\|_\infty = 1$, $f(0) = 1$ and for any integer $j \geq 1$ set $g_j(t) = f(t)\varphi_j$, where $\{\varphi_j\}_{j \geq 1}$ is the previously chosen orthonormal basis for H . Itô's formula implies that for any $j \geq 1$, and for $0 \leq t \leq T$,

$$(\phi_{n,h}^\varepsilon(T), g_j(T)) = (\phi_{n,h}^\varepsilon(0), g_j(0)) + \sum_{i=1}^4 I_{n,k}^i, \quad (3.33)$$

where

$$\begin{aligned} I_{n,k}^1 &= \int_0^T (\phi_{n,h}^\varepsilon(s), \varphi_j) f'(s) ds, \\ I_{n,k}^2 &= \sqrt{\varepsilon} \int_0^T (\sigma_n(\phi_{n,h}^\varepsilon(s)) P_n dW_n(s), g_j(s)), \\ I_{n,k}^3 &= \int_0^T \langle F(\phi_{n,h}^\varepsilon(s)), g_j(s) \rangle ds, \\ I_{n,k}^4 &= \int_0^T (\tilde{\sigma}_n(\phi_{n,h}^\varepsilon(s)) h(s), g_j(s)) ds. \end{aligned}$$

Since $f' \in L^2([0, T])$ and for every $X \in L^2(\Omega)$, $(t, \omega) \mapsto \varphi_j X(\omega) f'(t) \in L^2(\Omega, H)$, (i) above implies that as $n \rightarrow \infty$, $I_{n,k}^1 \rightarrow \int_0^T (\phi_h^\varepsilon(s), \varphi_j) f'(s) ds$ weakly in $L^2(\Omega)$. Similarly, (v) implies that as $n \rightarrow \infty$, $I_{n,k}^3 \rightarrow \int_0^T \langle F_h^\varepsilon(s), g_j(s) \rangle ds$ weakly in $L^2(\Omega)$, while (vii) implies that $I_{n,k}^4 \rightarrow \int_0^T (\tilde{S}_h^\varepsilon(s), g_j(s)) ds$ weakly in $L^{\frac{4}{3}}(\Omega)$. To prove the convergence of $I_{n,k}^2$, as in [22], let $\mathcal{P}_T$ denote the class of predictable processes in $L^2(\Omega_T, L_Q(H_0, H))$ with the inner product

$$(G, J)_{\mathcal{P}_T} = \mathbb{E} \int_0^T (G(s), J(s))_{\mathcal{P}_T} ds = \mathbb{E} \int_0^T \text{trace}(G(s) Q J(s)^*) ds.$$

The map $\mathcal{T} : \mathcal{P}_T \rightarrow L^2(\Omega)$ defined by $\mathcal{T}(G)(t) = \int_0^t (G(s) dW(s), g_j(s))$ is linear and continuous because of the Itô isometry. Furthermore, (vi) shows that for every $G \in \mathcal{P}_T$, as $n \rightarrow \infty$, $(\sigma_n(\phi_{n,h}^\varepsilon) P_n, G)_{\mathcal{P}_T} \rightarrow (S_h^\varepsilon, G)_{\mathcal{P}_T}$ weakly in $L^2(\Omega)$.

Finally, as $n \rightarrow \infty$, $P_n \xi = \phi_{n,h}^\varepsilon(0) \rightarrow \xi$ in H and by (iv), $(\phi_{n,h}^\varepsilon(T), g_j(T)) \rightarrow (\tilde{\phi}_h^\varepsilon(T), g_j(T))$ weakly in $L^2(\Omega)$. Therefore, (3.33) leads to, as $n \rightarrow \infty$,

$$\begin{aligned} (\tilde{\phi}_h^\varepsilon(T), \varphi_j) f(T) &= (\xi, \varphi_j) + \int_0^T (\phi_h^\varepsilon(s), \varphi_j) f'(s) ds + \sqrt{\varepsilon} \int_0^T (S_h^\varepsilon(s) dW(s), g_j(s)) \\ &\quad + \int_0^T \langle F_h^\varepsilon(s), g_j(s) \rangle ds + \int_0^T (\tilde{S}_h^\varepsilon(s), g_j(s)) ds. \end{aligned} \quad (3.34)$$

For $\delta > 0$, $k > \frac{1}{\delta}$, $t \in [0, T]$, let $f_k \in H^1(-\delta, T + \delta)$ be such that $\|f_k\|_\infty = 1$, $f_k = 1$ on $(-\delta, t - \frac{1}{k})$ and $f_k = 0$ on $(t, T + \delta)$. Then $f_k \rightarrow 1_{(-\delta, t)}$ in L^2 , and $f'_k \rightarrow -\delta_t$ in

the sense of distributions. Hence as $k \rightarrow \infty$, (3.34) written with $f := f_k$ yields

$$\begin{aligned} 0 &= (\xi, \varphi_j) - (\phi_h^\varepsilon(t), \varphi_j) + \sqrt{\varepsilon} \left(\int_0^t S_h^\varepsilon(s) dW(s), \varphi_j \right) \\ &\quad + \left\langle \int_0^t F_h^\varepsilon(s) ds, \varphi_j \right\rangle + \left(\int_0^t \tilde{S}_h^\varepsilon(s) ds, \varphi_j \right). \end{aligned}$$

Note that j is arbitrary and $\mathbb{E} \int_0^T |S_h^\varepsilon(s)|_{L_Q}^2 ds < \infty$; we deduce that for $0 \leq t \leq T$,

$$\phi_h^\varepsilon(t) = \xi + \sqrt{\varepsilon} \int_0^t S_h^\varepsilon(s) dW(s) + \int_0^t F_h^\varepsilon(s) ds + \int_0^t \tilde{S}_h^\varepsilon(s) ds \in H. \quad (3.35)$$

Indeed, $\int_0^t F_h^\varepsilon(s) ds$, as a linear combination of H -valued terms, also belongs to H . Moreover, let $f = 1_{(-\delta, T+\delta)}$. Using (3.34) again, we obtain

$$\tilde{\phi}_h^\varepsilon(T) = \xi + \sqrt{\varepsilon} \int_0^T S_h^\varepsilon(s) dW(s) + \int_0^T F_h^\varepsilon(s) ds + \int_0^T \tilde{S}_h^\varepsilon(s) ds.$$

This equation and (3.35) yield that $\tilde{\phi}_h^\varepsilon(T) = \phi_h^\varepsilon(T)$ a.s.

Step 3: In (3.35) we still have to prove that $ds \otimes d\mathbb{P}$ a.s. on Ω_T , one has

$$S_h^\varepsilon(s) = \sigma(\phi_h^\varepsilon(s)), \quad F_h^\varepsilon(s) = F(\phi_h^\varepsilon(s)) \quad \text{and} \quad \tilde{S}_h^\varepsilon(s) = \tilde{\sigma}(\phi_h^\varepsilon(s)) h(s).$$

Let

$$\begin{aligned} \mathcal{X} &:= \{ \psi \in L^4(\Omega_T, L^4(D)^3) \cap L^4(\Omega, L^\infty([0, T], H)) \cap L^2(\Omega_T, V) : \\ &\quad \int_0^T (\|\psi(t)\|^2 + \|\phi_h^\varepsilon(t)\|^2) |\psi(t) - \phi_h^\varepsilon(t)|^2 dt < +\infty \text{ a.s.} \}. \end{aligned}$$

Then (i)-(iii) yield $\phi_h^\varepsilon \in \mathcal{X}$ and since $\|u\| \leq C(m)|u|$ for every $u \in H_m$, using (3.8) and the fact that $\phi_h^\varepsilon \in L^2(\Omega_T, V)$, we deduce that for any $m \geq 1$, $L^\infty(\Omega_T, H_m) \subset \mathcal{X}$. Let $\psi = (v, \eta) \in L^\infty(\Omega_T, H_m)$. For every $t \in [0, T]$, if $a \wedge b := \inf(a, b)$ and c_1 is the constant in (2.5), set

$$r(t) = \int_0^t \left[2 + \frac{8c_1^2}{\nu \wedge \kappa} \|\psi(s)\|^2 + \frac{2c_1 c_2 \tilde{L}}{\nu \wedge \kappa} |h(s)|_0^2 \right] ds. \quad (3.36)$$

Then $r(t) < \infty$ for all $t \in [0, T]$ and Fatou's lemma implies

$$\mathbb{E}(|\phi_h^\varepsilon(T)|^2 e^{-r(T)}) \leq \mathbb{E}(\liminf_n |\phi_{n,h}^\varepsilon(T)|^2 e^{-r(T)}) \leq \liminf_n \mathbb{E}(|\phi_{n,h}^\varepsilon(T)|^2 e^{-r(T)}).$$

Apply Itô's formula to (3.35) and (3.16), and for $\phi = \phi_h^\varepsilon$ or $\phi = \phi_{n,h}^\varepsilon$, let $\phi = \psi + (\phi - \psi)$. After simplification, this yields

$$\begin{aligned} \mathbb{E}|\xi|^2 + \mathbb{E} \int_0^T e^{-r(s)} \left[-r'(s) \{ |\phi_h^\varepsilon(s) - \psi(s)|^2 + 2(\phi_h^\varepsilon(s) - \psi(s), \psi(s)) \} + 2\langle F_h^\varepsilon(s), \phi_h^\varepsilon(s) \rangle \right. \\ \left. + \varepsilon |S_h^\varepsilon(s)|_{L_Q}^2 + 2(\tilde{S}_h^\varepsilon(s), \phi_h^\varepsilon(s)) \right] ds \leq \liminf_n (\mathbb{E}|P_n(\xi)|^2 + X_n), \end{aligned} \quad (3.37)$$

where

$$\begin{aligned} X_n = & \mathbb{E} \int_0^T e^{-r(s)} \left[-r'(s) \left\{ |\phi_{n,h}^\varepsilon(s) - \psi(s)|^2 + 2(\phi_{n,h}^\varepsilon(s) - \psi(s), \psi(s)) \right\} \right. \\ & \left. + 2\langle F(\phi_{n,h}^\varepsilon(s)), \phi_{n,h}^\varepsilon(s) \rangle + \varepsilon |\sigma_n(\phi_{n,h}^\varepsilon(s)) P_n|_{L_Q}^2 + 2(\tilde{\sigma}(\phi_{n,h}^\varepsilon(s)) h(s), \phi_{n,h}^\varepsilon(s)) \right] ds. \end{aligned}$$

Set $a \vee b := \max(a, b)$. Inequalities (3.15), (A.3), (A.2), (3.36), the Poincaré and Schwarz inequalities imply that for $0 \leq \varepsilon \leq \varepsilon_0 \leq \frac{\nu \wedge \kappa}{2L}$,

$$\begin{aligned} Y_n := & \mathbb{E} \int_0^T e^{-r(s)} \left[-r'(s) |\phi_{n,h}^\varepsilon(s) - \psi(s)|^2 \right. \\ & + \left[2\langle F(\phi_{n,h}^\varepsilon(s)) - F(\psi(s)), \phi_{n,h}^\varepsilon(s) - \psi(s) \rangle + \varepsilon |\sigma_n(\phi_{n,h}^\varepsilon(s)) P_n - \sigma_n(\psi(s)) P_n|_{L_Q}^2 \right. \\ & \left. + 2(\{\tilde{\sigma}_n(\phi_{n,h}^\varepsilon(s)) - \tilde{\sigma}_n(\psi(s))\} h(s), \phi_{n,h}^\varepsilon(s) - \psi(s)) \right] ds \\ \leq & \mathbb{E} \int_0^T e^{-r(s)} |\phi_{n,h}^\varepsilon(s) - \psi(s)|^2 \left\{ -r'(s) + 2 + \frac{8c_1^2}{\nu \wedge \kappa} \|\psi(s)\|^2 + \frac{2c_1 c_2 \tilde{L}}{\nu \wedge \kappa} |h(s)|_0^2 \right\} ds \\ \leq & 0. \end{aligned} \tag{3.38}$$

Furthermore, $X_n = Y_n + \sum_{i=1}^2 Z_n^i$, with

$$\begin{aligned} Z_n^1 = & \mathbb{E} \int_0^T e^{-r(s)} \left[-2r'(s) (\phi_{n,h}^\varepsilon(s) - \psi(s), \psi(s)) + 2\langle F(\phi_{n,h}^\varepsilon(s)), \psi(s) \rangle \right. \\ & + 2\langle F(\psi(s)), \phi_{n,h}^\varepsilon(s) \rangle - 2\langle F(\psi(s)), \psi(s) \rangle + 2\varepsilon (\sigma_n(\phi_{n,h}^\varepsilon(s)) P_n, \sigma(\psi(s)))_{L_Q} \\ & \left. + 2(\tilde{\sigma}_n(\phi_{n,h}^\varepsilon(s)) h(s), \psi(s)) + 2(\tilde{\sigma}(\psi(s)) h(s), \phi_{n,h}^\varepsilon(s)) - 2(P_n \tilde{\sigma}(\psi(s)) h(s), \psi(s)) \right] ds, \\ Z_n^2 = & \mathbb{E} \int_0^T e^{-r(s)} \left[2\varepsilon (\sigma_n(\phi_{n,h}^\varepsilon(s)) P_n, [\sigma(\psi(s)) P_n - \sigma(\psi(s))])_{L_Q} - \varepsilon |P_n \sigma(\psi(s)) P_n|_{L_Q}^2 \right] ds. \end{aligned}$$

The weak convergence properties (i)-(vii) imply that, as $n \rightarrow \infty$, $Z_n^1 \rightarrow Z^1$ where

$$\begin{aligned} Z^1 = & \mathbb{E} \int_0^T e^{-r(s)} \left[-2r'(s) (\phi_h^\varepsilon(s) - \psi(s), \psi(s)) + 2\langle F_h^\varepsilon(s), \psi(s) \rangle + 2\langle F(\psi(s)), \phi_h^\varepsilon(s) \rangle \right. \\ & - 2\langle F(\psi(s)), \psi(s) \rangle + 2\varepsilon (S_h^\varepsilon(s), \sigma(\psi(s)))_{L_Q} + 2(\tilde{S}_h^\varepsilon(s), \psi(s)) \\ & \left. + 2(\tilde{\sigma}(\psi(s)) h(s), \phi_h^\varepsilon(s)) - 2(\tilde{\sigma}(\psi(s)) h(s), \psi(s)) \right] ds. \end{aligned} \tag{3.39}$$

Now we study (Z_n^2) ; when $n \rightarrow \infty$, $|\sigma(\psi(s))(P_n - Id_{H_0})|_{L_Q} \rightarrow 0$ a.s., and by (A.2),

$$\mathbb{E} \int_0^T e^{-r(s)} \sup_n |\sigma(\psi(s))(P_n - Id_{H_0})|_{L_Q}^2 ds < \infty.$$

Hence the Lebesgue dominated convergence theorem implies that, as $n \rightarrow \infty$,

$$\mathbb{E} \int_0^T e^{-r(s)} |\sigma(\psi(s))(P_n - Id_{H_0})|_{L_Q}^2 ds \rightarrow 0.$$

Since $\sup_n \mathbb{E} \int_0^T e^{-r(s)} |\sigma_n(\phi_{n,h}^\varepsilon(s)) P_n|_{L_Q}^2 ds < \infty$, by the Cauchy-Schwarz inequality, we see that $Z_n^2 \rightarrow -\varepsilon \mathbb{E} \int_0^T e^{-r(s)} |\sigma(\psi(s))|_{L_Q}^2 ds$.

Thus, (3.37)-(3.39) imply that for any $m \geq 1$ and any $\psi \in L^\infty(\Omega_T, H_m)$,

$$\begin{aligned} & \mathbb{E} \int_0^T e^{-r(s)} \left\{ -r'(s) |\phi_h^\varepsilon(s) - \psi(s)|^2 + 2 \langle F_h^\varepsilon(s) - F(\psi(s)), \phi_h^\varepsilon(s) - \psi(s) \rangle \right. \\ & \left. + \varepsilon |S_h^\varepsilon(s) - \sigma(\psi(s))|_{L_Q}^2 + 2 \left(\tilde{S}_h^\varepsilon(s) - \tilde{\sigma}(\psi(s)) h(s), \phi_h^\varepsilon(s) - \psi(s) \right) \right\} ds \leq 0. \end{aligned} \quad (3.40)$$

By a density argument, this inequality extends to all $\psi \in \mathcal{X}$. Taking $\psi = \phi_h^\varepsilon \in \mathcal{X}$, we conclude that $S_h^\varepsilon(s) = \sigma(\phi_h^\varepsilon(s)) ds \otimes d\mathbb{P}$ a.e. For a real number λ , $\tilde{\psi} = (v, \eta) \in L^\infty(\Omega_T, H_m)$ for some m , set $\psi_\lambda = \phi_h^\varepsilon - \lambda \tilde{\psi} \in \mathcal{X}$. Thus applying (3.40) to ψ_λ and neglecting $\varepsilon |\sigma(\phi_h^\varepsilon(s)) - \sigma(\psi_\lambda(s))|_{L_Q}^2$, we obtain

$$\begin{aligned} & \mathbb{E} \int_0^T e^{-r(s)} \left[-\lambda^2 r'(s) |\tilde{\psi}(s)|^2 + 2\lambda \left\{ \langle F_h^\varepsilon(s) - F(\psi_\lambda(s)), \tilde{\psi}(s) \rangle \right. \right. \\ & \left. \left. + \left(\tilde{S}_h^\varepsilon(s) - \tilde{\sigma}(\psi_\lambda(s)) h(s), \tilde{\psi}(s) \right) \right\} \right] ds \leq 0. \end{aligned} \quad (3.41)$$

Using (A.2), (2.5) and (2.6), we have for almost every $(s, \omega) \in \Omega_T$ as $\lambda \rightarrow 0$,

$$|([\tilde{\sigma}(\psi_\lambda(s)) - \tilde{\sigma}(\phi_h^\varepsilon(s))] h(s), \tilde{\psi}(s))| \leq \sqrt{L c_1 c_2} \lambda \|\tilde{\psi}(s)\| |h(s)|_0 |\tilde{\psi}(s)| \rightarrow 0.$$

Furthermore, (A.1) (2.5) and (2.6) imply that for some constant $c > 0$,

$$\begin{aligned} & \mathbb{E} \int_0^T \sup_{|\lambda| \leq 1} \left| \left(\tilde{\sigma}(\psi_\lambda(s)) h(s), \tilde{\psi}(s) \right) \right| ds \\ & \leq \sqrt{\tilde{K} c} \mathbb{E} \int_0^T (1 + 2\|\phi_h^\varepsilon(s)\|^2 + 2\|\tilde{\psi}(s)\|^2)^{\frac{1}{2}} |h(s)|_0 |\tilde{\psi}(s)| ds \\ & \leq c \tilde{K} M + \mathbb{E} \int_0^T \left[\{1 + 2\|\phi_h^\varepsilon(s)\|^2 + 2\|\tilde{\psi}(s)\|^2\} |\tilde{\psi}(s)|^2 \right] ds < \infty. \end{aligned}$$

Hence, the Lebesgue dominated convergence theorem yields, as $\lambda \rightarrow 0$,

$$\mathbb{E} \int_0^T \left(\{ \tilde{S}_h^\varepsilon(s) - \tilde{\sigma}(\psi_\lambda(s)) \} h(s), \tilde{\psi}(s) \right) ds \rightarrow \mathbb{E} \int_0^T \left(\{ \tilde{S}_h^\varepsilon(s) - \tilde{\sigma}(\phi_h(s)) \} h(s), \tilde{\psi}(s) \right) ds.$$

Furthermore, (3.15) yields for $\lambda \neq 0$

$$|\langle F(\psi_\lambda(s)) - F(\phi_h^\varepsilon(s)), \tilde{\psi}(s) \rangle| \leq \lambda^2 \left[(\nu \wedge \kappa) \|\tilde{\psi}(s)\|^2 + 2c_1 \|\tilde{\psi}(s)\|^2 |\tilde{\psi}(s)| + |\tilde{\psi}(s)|^2 \right].$$

Using again the dominated convergence theorem, we deduce as $\lambda \rightarrow 0$,

$$\mathbb{E} \int_0^T \langle F_h^\varepsilon(s) - F(\psi_\lambda(s)), \tilde{\psi}(s) \rangle ds \rightarrow \mathbb{E} \int_0^T \langle F_h^\varepsilon(s) - F(\phi_h^\varepsilon(s)), \tilde{\psi}(s) \rangle ds.$$

Thus, dividing (3.41) by $\lambda > 0$ and letting $\lambda \rightarrow 0$ we obtain that for every m and $\tilde{\psi} \in L^\infty(\Omega_T, H_m)$,

$$\mathbb{E} \int_0^T \left[\langle F_h^\varepsilon(s) - F(\phi_h^\varepsilon(s)), \tilde{\psi}(s) \rangle + (\{\tilde{S}_h^\varepsilon(s) - \tilde{\sigma}(\phi_h^\varepsilon(s))\} h(s), \tilde{\psi}(s)) \right] ds \leq 0,$$

while a similar calculation for $\lambda < 0$ yields the opposite inequality. Therefore for almost every $(s, \omega) \in \Omega_T$, for every $\tilde{\psi}$ in a dense subset of $L^2(\Omega_T, V)$,

$$\mathbb{E} \int_0^T \left[\langle F_h^\varepsilon(s) - F(\phi_h^\varepsilon(s)), \tilde{\psi}(s) \rangle + (\{\tilde{S}_h^\varepsilon(s) - \tilde{\sigma}(\phi_h^\varepsilon(s))\} h(s), \tilde{\psi}(s)) \right] ds = 0. \quad (3.42)$$

Hence a.e. for $t \in [0, T]$, (3.35) can be rewritten as

$$\phi_h^\varepsilon(t) = \xi + \sqrt{\varepsilon} \int_0^t \sigma(\phi_h^\varepsilon(s)) dW_s + \int_0^t [F(\phi_h^\varepsilon(s)) + \tilde{\sigma}(\phi_h^\varepsilon(s)) h(s)] ds. \quad (3.43)$$

Furthermore, (i), (iv) and (3.21) for $p = 2$ imply that

$$E \left(\int_0^T \|\phi_h^\varepsilon(t)\|^2 dt \right) \leq \sup_n \mathbb{E} \int_0^T \|\phi_{n,h}^\varepsilon(t)\|^2 dt \leq C(1 + E|\xi|^4), \quad (3.44)$$

$$E \left(\sup_{0 \leq t \leq T} |\phi_h^\varepsilon(t)|^4 \right) \leq \sup_n \mathbb{E} \left(\sup_{0 \leq t \leq T} |\phi_{n,h}^\varepsilon(t)|^4 \right) \leq C(1 + E|\xi|^4). \quad (3.45)$$

Since $|x|^2 \leq 1 \vee |x|^4$ for any $x \in \mathbb{R}$, this completes the proof of (3.5).

Step 4: To complete the proof of Theorem 3.1, we show that ϕ_h^ε has a $C([0, T], H)$ -valued modification and that the solution to (3.43) is unique in $X := C([0, T], H) \cap L^2([0, T], V)$. Note that (3.5) implies that if $\tilde{\tau}_N = \inf\{t \geq 0 : |\phi_h^\varepsilon(t)| \geq N\} \wedge T$ for $N > 0$, $\mathbb{P}(\tilde{\tau}_N < T) \leq CN^{-2}$. The Borel-Cantelli lemma yields $\tilde{\tau}_N \rightarrow T$ a.s. when $N \rightarrow \infty$.

We at first prove uniqueness. Let $\psi = (v, \eta) \in X$ be another solution to (3.43). Then if $\bar{\tau}_N = \inf\{t \geq 0 : |\psi(t)| \geq N\} \wedge T$ for $N > 0$, since $|\psi(\cdot)|$ is a.s. bounded on $[0, T]$, as $N \rightarrow \infty$, we have $\bar{\tau}_N \rightarrow T$ a.s. and hence $\tau_N = \tilde{\tau}_N \wedge \bar{\tau}_N \rightarrow T$, a.s.

Let $\phi_h^\varepsilon = (u_h^\varepsilon, \theta_h^\varepsilon)$, $\Phi = \phi_h^\varepsilon - \psi$, and $a = \frac{8c_1^2}{\nu \wedge \kappa}$, where c_1 is the constant defined in (2.5). Set $\rho'(t) := a\|\psi(t)\|^2$, $h_\varepsilon := h$, $\sigma_1 = \sigma_2 = \sigma$, $\bar{\sigma} = \tilde{\sigma}$, $\bar{\sigma}_1 = \bar{\sigma}_2 = 0$. Then $\phi_1 = \phi_h^\varepsilon$ and $\phi_2 = \psi$ satisfy (3.30). Set

$$\mathcal{I}(t) = \sup_{\tau \leq t} 2\sqrt{\varepsilon} \int_0^\tau e^{-a \int_0^s \|\psi(r)\|^2 dr} \left([\sigma(\phi_h^\varepsilon(s)) - \sigma(\psi(s))] dW(s), \Phi(s) \right),$$

Then using Lemma 3.12 and condition **(A.3)** yields for $0 \leq \varepsilon \leq \varepsilon_0 \leq \frac{\nu \wedge \kappa}{2L}$

$$\begin{aligned} \zeta(t) : &= e^{-\rho(t \wedge \tau_N)} |\Phi(t \wedge \tau_N)|^2 \\ &\leq \mathcal{I}(t \wedge \tau_N) + \int_0^{t \wedge \tau_N} e^{-\rho(s)} \left\{ [\varepsilon L - \nu \wedge \kappa] \|\Phi(s)\|^2 \right. \\ &\quad \left. + |\Phi(s)|^2 [-a\|\psi(s)\|^2 + 2 + \frac{8c_1^2}{\nu \wedge \kappa} \|\psi(s)\|^2 + \frac{2\tilde{L}c_1c_2}{\nu \wedge \kappa} |h(s)|_0^2] \right\} ds. \end{aligned}$$

Thus

$$\zeta(t) + \frac{\nu \wedge \kappa}{2} Y(t) \leq \int_0^t \left(\frac{2\tilde{L}c_1c_2}{\nu \wedge \kappa} |h(s \wedge \tau_n)|_0^2 + 2 \right) \zeta(s) ds + \mathcal{I}(t \wedge \tau_n),$$

where $Y(t) = \int_0^{t \wedge \tau_n} e^{-\rho(s)} \|\Phi(s)\|^2 ds$. Burkholder's inequality and Assumption **(A.3)** imply that for all $\beta > 0$ and $\varepsilon \in [0, \varepsilon_0]$,

$$\mathbb{E} \mathcal{I}(t \wedge \tau_n) \leq 6\sqrt{\varepsilon_0} \mathbb{E} \left(\int_0^{t \wedge \tau_N} e^{-2\rho(s)} L \|\Phi(s)\|^2 |\Phi(s)|^2 ds \right)^{\frac{1}{2}} \leq \beta \mathbb{E} \sup_{0 \leq s \leq t} \zeta(s) + \frac{9L\varepsilon_0}{\beta} \mathbb{E} Y(t).$$

Since $\int_0^T \left(\frac{2\tilde{L}c_1c_2}{\nu \wedge \kappa} |h(s \wedge \tau_N)|_0^2 + 2 \right) ds \leq \frac{2M\tilde{L}c_1c_2}{\nu \wedge \kappa} + 2T := C$, Lemma 3.9 implies that for $\beta = (2[1 + Ce^C])^{-1}$ and $\varepsilon_0 L$ small enough to have $\frac{9\varepsilon_0 L}{\beta} \leq \frac{\nu \wedge \kappa}{2} \beta$, one has

$$\mathbb{E} \sup_{0 \leq s \leq T} e^{-a \int_0^{s \wedge \tau_N} \|\psi(r)\|^2 dr} |\Phi(s \wedge \tau_N)|^2 = 0. \quad (3.46)$$

Since $\lim_{N \rightarrow \infty} \tau_N = T$ a.s., we thus deduce $|\Phi(s, \omega)| = 0$ a.s. on Ω_T . Thus if ϕ_h^ε is in $C([0, T], H)$, we conclude that $\phi_h^\varepsilon(t) = \psi(t)$, a.s., for every $t \in [0, T]$.

Finally, set

$$\tilde{\rho}'(t) = \frac{8c_1^2}{\nu \wedge \kappa} \|\phi_h^\varepsilon(s)\|^2 + 2 + \frac{2\tilde{L}c_1c_2}{\nu \wedge \kappa} |h(s)|_0^2, \quad (3.47)$$

let $h_\varepsilon := h$, $\sigma_1 = P_n \sigma P_n$, $\sigma_2 = \sigma$, $\bar{\sigma}_1 = 0$, $\bar{\sigma}_2(s) = [\tilde{\sigma}(\phi_h^\varepsilon(s)) - P_n \tilde{\sigma}(\phi_h^\varepsilon(s))] h(s)$ and $\bar{\sigma} = P_n \tilde{\sigma}$. Then $\tilde{\rho}(t) = \int_0^t \tilde{\rho}'(s) ds < +\infty$ a.s. Then $\phi_1 = \phi_{n,h}^\varepsilon$ and $\phi_2 = \phi_h^\varepsilon$ satisfy (3.30). Set $\Phi_{n,h}^\varepsilon = \phi_{n,h}^\varepsilon - \phi_h^\varepsilon$ and let $0 \leq \varepsilon \leq \varepsilon_0 \leq \frac{\nu \wedge \kappa}{4L}$. By Lemma 3.12 and condition **(A.3)**, we deduce that for every $t \in [0, T]$,

$$\begin{aligned} \mathbb{E}(e^{-\tilde{\rho}(t)} |\Phi_{n,h}^\varepsilon(t)|^2) &\leq \mathbb{E} \int_0^t e^{-\tilde{\rho}(s)} \left\{ [2\varepsilon L - (\nu \wedge \kappa)] \|\Phi_{n,h}^\varepsilon\|^2 + 2\varepsilon |P_n \sigma(\phi_h^\varepsilon(s)) P_n - \sigma(\phi_h^\varepsilon(s))|_{L_Q}^2 \right. \\ &\quad \left. + |\Phi_{n,h}^\varepsilon(s)|^2 [-\tilde{\rho}'(s) + 2 + \frac{8c_1^2}{\nu \wedge \kappa} \|\phi_h^\varepsilon(s)\|^2 + \frac{2\tilde{L}c_1c_2}{\nu \wedge \kappa} |h(s)|_0^2] \right\} ds \\ &\quad + \mathbb{E} \int_0^t e^{-\tilde{\rho}(s)} 2 |\Phi_{n,h}^\varepsilon(s)| |P_n \tilde{\sigma}(\phi_h^\varepsilon(s)) - \tilde{\sigma}(\phi_h^\varepsilon(s))|_{L_Q} |h(s)|_0 ds \\ &\leq \mathcal{R}(t, n) - \frac{\nu \wedge \kappa}{2} \mathbb{E} \int_0^t e^{-\tilde{\rho}(s)} \|\Phi_{n,h}^\varepsilon(s)\|^2 ds, \end{aligned}$$

where

$$\mathcal{R}(t, n) = \mathbb{E} \int_0^t [2\varepsilon |P_n \sigma(\phi_h^\varepsilon(s)) P_n - \sigma(\phi_h^\varepsilon(s))|_{L_Q}^2 + |P_n \tilde{\sigma}(\phi_h^\varepsilon(s)) - \tilde{\sigma}(\phi_h^\varepsilon(s))|_{L_Q}^2] ds,$$

and the last inequality follows from Schwarz's inequality and the definition of $\tilde{\rho}$.

Furthermore, for almost every (s, ω) , one has $|P_n \sigma(\phi_h^\varepsilon(s)) P_n - \sigma(\phi_h^\varepsilon(s))|_{L_Q} \rightarrow 0$ and $|P_n \tilde{\sigma}(\phi_h^\varepsilon(s)) - \tilde{\sigma}(\phi_h^\varepsilon(s))|_{L_Q} \rightarrow 0$ as $n \rightarrow \infty$. Thus the dominated convergence theorem shows that $\lim_n \sup_t \mathcal{R}(t, n) \rightarrow 0$, and thus that $\lim_{n \rightarrow \infty} I(n) = 0$, where

$$I(n) = \sup_{0 \leq t \leq T} \mathbb{E}(e^{-\tilde{\rho}(t)} |\Phi_{n,h}^\varepsilon(t)|^2) + \mathbb{E} \int_0^T e^{-\tilde{\rho}(s)} \|\Phi_{n,h}^\varepsilon(s)\|^2 ds.$$

Using again Lemma 3.12 and the Burkholder-Davies-Gundy inequality, a similar computation yields that for $0 \leq \varepsilon \leq \varepsilon_0 \leq \frac{\nu \wedge \kappa}{4L}$:

$$\begin{aligned} \mathbb{E} \left(\sup_{0 \leq t \leq T} e^{-\tilde{\rho}(t)} |\Phi_{n,h}^\varepsilon(t)|^2 \right) &\leq \frac{1}{2} \mathbb{E} \left(\sup_{0 \leq t \leq T} e^{-\tilde{\rho}(t)} |\Phi_{n,h}^\varepsilon(t)|^2 \right) \\ &\quad + 18\varepsilon \mathbb{E} \int_0^T e^{-\tilde{\rho}(s)} |\sigma_n(\phi_{n,h}^\varepsilon(s)) P_n - \sigma(\phi_h^\varepsilon(s))|_{L_Q}^2 ds \\ &\quad + \mathbb{E} \int_0^T \left[2\varepsilon |P_n \sigma(\phi_h^\varepsilon(s)) P_n - \sigma(\phi_h^\varepsilon(s))|_{L_Q}^2 + |P_n \tilde{\sigma}(\phi_h^\varepsilon(s)) - \tilde{\sigma}(\phi_h^\varepsilon(s))|_{L_Q}^2 \right] ds \\ &\leq C[I(n) + \mathcal{R}(T, n)]. \end{aligned}$$

Therefore, $\phi_{n,h}^\varepsilon$ has a subsequence converging a.s. uniformly to ϕ_h^ε in H . Since $\phi_{n,h}^\varepsilon \in C([0, T], H)$, we conclude that ϕ_h^ε has a modification in $C([0, T], H)$. $\square$

4. LARGE DEVIATIONS

We consider large deviations via a weak convergence approach [1, 2], based on variational representations of infinite-dimensional Wiener processes. The solution to the stochastic Bénard model (2.9) is denoted as $\phi^\varepsilon = \mathcal{G}^\varepsilon(\sqrt{\varepsilon}W)$ for a Borel measurable function $\mathcal{G}^\varepsilon : C([0, T], H) \rightarrow X$. The space $X = C([0, T]; H) \cap L^2((0, T); V)$ endowed with the metric associated with the norm defined in (3.4) is Polish. Let $\mathcal{B}(X)$ denote its Borel σ -field. We recall some classical definitions.

Definition 4.1. *The random family $\{\phi^\varepsilon\}$ is said to satisfy a large deviation principle on X with the good rate function I if the following conditions hold:*

***I is a good rate function.** The function $I : X \rightarrow [0, \infty]$ is such that for each $M \in [0, \infty[$ the level set $\{\phi \in X : I(\phi) \leq M\}$ is a compact subset of X . For $A \in \mathcal{B}(X)$, set $I(A) = \inf_{\phi \in A} I(\phi)$.*

***Large deviation upper bound.** For each closed subset F of X :*

$$\limsup_{\varepsilon \rightarrow 0} \varepsilon \log \mathbb{P}(\phi^\varepsilon \in F) \leq -I(F).$$

***Large deviation lower bound.** For each open subset G of X :*

$$\liminf_{\varepsilon \rightarrow 0} \varepsilon \log \mathbb{P}(\phi^\varepsilon \in G) \geq -I(G).$$

To establish the large deviation principle, we need to strengthen the hypothesis on the growth condition and Lipschitz property of σ (and $\tilde{\sigma}$) as follows:

Assumption A Bis There exist positive constants K and L such that

$$(A.4) \quad |\sigma(t, \phi)|_{L_Q}^2 \leq K(1 + |\phi|^2), \quad \forall t \in [0, T], \quad \forall \phi \in V.$$

$$(A.5) \quad |\sigma(t, \phi) - \sigma(t, \psi)|_{L_Q}^2 \leq L|\phi - \psi|^2, \quad \forall t \in [0, T], \quad \forall \phi, \psi \in V.$$

Note that due to the continuous embedding $V \hookrightarrow H$, assumptions (A.4-A.5) imply (A.2-A.3) as well as ($\tilde{A}$.1- $\tilde{A}$.2). Thus the conclusions of Theorem 3.1 hold if $\tilde{\sigma} = \sigma$ satisfy assumptions (A.4-A.5).

The proof of the large deviation principle will use the following technical lemma which studies time increments of the solution to the stochastic control equation. For any integer $k = 0, \dots, 2^n - 1$, and $s \in [kT2^{-n}, (k+1)T2^{-n}[$, set $\underline{s}_n = kT2^{-n}$ and $\bar{s}_n = (k+1)T2^{-n}$. Given $N > 0$, $h \in \mathcal{A}_M$, $\varepsilon \geq 0$ small enough, let ϕ_h^ε denote the solution to (3.2) given by Theorem 3.1, and for $t \in [0, T]$, let

$$G_N(t) = \left\{ \omega : \left(\sup_{0 \leq s \leq t} |\phi_h^\varepsilon(s)(\omega)|^2 \right) \vee \left(\int_0^t \|\phi_h^\varepsilon(s)(\omega)\|^2 ds \right) \leq N \right\}.$$

Lemma 4.2. *Let $M, N > 0$, σ and $\tilde{\sigma}$ satisfy Assumptions (A.1), (A.4) and (A.5), $\xi \in L^4(\Omega, H)$ be $\mathcal{F}_0$ -measurable and ϕ^ε be a solution to (3.2). Then there exists a positive constant $C := C(\nu, \kappa, K, L, T, M, N, \varepsilon_0)$ such that for any $h \in \mathcal{A}_M$, $\varepsilon \in [0, \varepsilon_0]$,*

$$I_n(h, \varepsilon) := \mathbb{E} \left[1_{G_N(T)} \int_0^T |\phi_h^\varepsilon(s) - \phi_h^\varepsilon(\bar{s}_n)|^2 ds \right] \leq C 2^{-\frac{n}{2}}. \quad (4.1)$$

Proof. Let $h \in \mathcal{A}_M$, $\varepsilon \geq 0$; Itô's formula yields $I_n(h, \varepsilon) = \sum_{1 \leq i \leq 6} I_{n,i}$, where

$$\begin{aligned} I_{n,1} &= 2\sqrt{\varepsilon} \mathbb{E} \left(1_{G_N(T)} \int_0^T ds \int_s^{\bar{s}_n} (\sigma(\phi_h^\varepsilon(r)) dW_r, \phi_h^\varepsilon(r) - \phi_h^\varepsilon(s)) \right), \\ I_{n,2} &= \varepsilon \mathbb{E} \left(1_{G_N(T)} \int_0^T ds \int_s^{\bar{s}_n} |\sigma(\phi_h^\varepsilon(r))|_{L_Q}^2 dr \right), \\ I_{n,3} &= 2\mathbb{E} \left(1_{G_N(T)} \int_0^T ds \int_s^{\bar{s}_n} (\tilde{\sigma}(\phi_h^\varepsilon(r)) h(r), \phi_h^\varepsilon(r) - \phi_h^\varepsilon(s)) dr \right), \\ I_{n,4} &= -2\mathbb{E} \left(1_{G_N(T)} \int_0^T ds \int_s^{\bar{s}_n} [\nu(A_1 u_h^\varepsilon(r), u_h^\varepsilon(r) - u_h^\varepsilon(s)) + \kappa(A_2 \theta_h^\varepsilon(r), \theta_h^\varepsilon(r) - \theta_h^\varepsilon(s))] dr \right), \\ I_{n,5} &= -2\mathbb{E} \left(1_{G_N(T)} \int_0^T ds \int_s^{\bar{s}_n} (B(\phi_h^\varepsilon(r)), \phi_h^\varepsilon(r) - \phi_h^\varepsilon(s)) dr \right), \\ I_{n,6} &= 2\mathbb{E} \left(1_{G_N(T)} \int_0^T ds \int_s^{\bar{s}_n} [(u_{h,2}^\varepsilon(r), \theta_h^\varepsilon(r) - \theta_h^\varepsilon(s)) + (\theta_h^\varepsilon(r), u_{h,2}^\varepsilon(r) - u_{h,2}^\varepsilon(s))] dr \right). \end{aligned}$$

Clearly $G_N(T) \subset G_N(r)$ for $r \in [0, T]$. The Burkholder-Davis-Gundy inequality, (A.4) and the definition of $G_N(r)$ yield for $0 \leq \varepsilon \leq \varepsilon_0$

$$\begin{aligned} |I_{n,1}| &\leq 2\sqrt{\varepsilon} \mathbb{E} \int_0^T ds \left| \int_s^{\bar{s}_n} (\sigma(\phi_h^\varepsilon(r)) dW_r, \phi_h^\varepsilon(r) - \phi_h^\varepsilon(s)) 1_{G_N(r)} \right| \\ &\leq 6\sqrt{\varepsilon} \int_0^T ds \mathbb{E} \left(\int_s^{\bar{s}_n} |\sigma(\phi_h^\varepsilon(r))|_{L_Q}^2 |\phi_h^\varepsilon(r) - \phi_h^\varepsilon(s)|^2 1_{G_N(r)} dr \right)^{\frac{1}{2}} \\ &\leq 12\sqrt{\varepsilon} \sqrt{KN(1+N)} \int_0^T ds (T2^{-n})^{\frac{1}{2}} \leq C(\varepsilon_0, K, N, T) 2^{-\frac{n}{2}}. \quad (4.2) \end{aligned}$$

Property (A.4) implies that for $\varepsilon \leq \varepsilon_0$,

$$|I_{n,2}| \leq \varepsilon K \mathbb{E} \left(1_{G_N(T)} \int_0^T ds \int_s^{\bar{s}_n} (1 + |\phi_h^\varepsilon(r)|^2) dr \right) \leq \varepsilon_0 K (1+N) T^2 2^{-n}. \quad (4.3)$$

Schwarz's inequality, Fubini's theorem, (A.4) and the definition of $\mathcal{A}_M$ yield

$$\begin{aligned} |I_{n,3}| &\leq 2\sqrt{K} \mathbb{E} \left(1_{G_N(T)} \int_0^T ds \int_s^{\bar{s}_n} (1 + |\phi_h^\varepsilon(r)|^2)^{\frac{1}{2}} |h(r)|_0 |\phi_h^\varepsilon(r) - \phi_h^\varepsilon(s)| dr \right) \\ &\leq 4\sqrt{KN(1+N)} \mathbb{E} \int_0^T |h(r)|_0 dr \int_{\underline{\tau}_n}^r ds \leq C(K, N, M, T) 2^{-n}. \end{aligned} \quad (4.4)$$

Schwarz's inequality and (3.5) imply that for some constant $\tilde{C} := C(\varepsilon_0, \nu, \kappa, K, T)$

$$\begin{aligned} I_{n,4} &\leq \mathbb{E} \left(1_{G_N(T)} \int_0^T ds \int_s^{\bar{s}_n} dr \left[-\nu \|u_h^\varepsilon(r)\|^2 - \kappa \|\theta_h^\varepsilon(r)\|^2 + \nu \|u_h^\varepsilon(r)\| \|u_h^\varepsilon(s)\| \right. \right. \\ &\quad \left. \left. + \kappa \|\theta_h^\varepsilon(r)\| \|\theta_h^\varepsilon(s)\| \right] \right) \\ &\leq \frac{\nu + \kappa}{2} \mathbb{E} \left(1_{G_N(T)} \int_0^T ds \|\phi_h^\varepsilon(s)\|^2 \int_s^{\bar{s}_n} dr \right) \leq \tilde{C} 2^{-n}. \end{aligned} \quad (4.5)$$

Inequalities (3.5), (3.8) and (3.9), Schwarz's inequality and Fubini's theorem imply that for some constant $\tilde{C} := C(\varepsilon_0, \nu, \kappa, K, T)$,

$$\begin{aligned} |I_{n,5}| &\leq 2c_1 \mathbb{E} \left(1_{G_N(T)} \int_0^T ds \int_s^{\bar{s}_n} dr \left[|u_h^\varepsilon(r)| \|u_h^\varepsilon(r)\| (\|u_h^\varepsilon(r)\| + \|u_h^\varepsilon(s)\|) \right. \right. \\ &\quad \left. \left. + |\phi_h^\varepsilon(r)| \|\phi_h^\varepsilon(r)\| (\|\theta_h^\varepsilon(r)\| + \|\theta_h^\varepsilon(s)\|) \right] \right) \\ &\leq 3c_1 \sqrt{N} \mathbb{E} \int_0^T dr (\|u_h^\varepsilon(r)\|^2 + \|\phi_h^\varepsilon(r)\|^2) \int_{\underline{\tau}_n}^r ds \\ &\quad + c_1 \sqrt{N} \mathbb{E} \int_0^T ds (\|u_h^\varepsilon(s)\|^2 + \|\phi_h^\varepsilon(s)\|^2) \int_s^{\bar{s}_n} dr \leq \sqrt{N} \tilde{C} 2^{-n}. \end{aligned} \quad (4.6)$$

Finally, Schwarz's inequality implies that

$$|I_{n,6}| \leq 4\mathbb{E} \left(1_{G_N(T)} \int_0^T ds \int_s^{\bar{s}_n} (|u_h^\varepsilon(r)| + |u_h^\varepsilon(s)|) (|\theta_h^\varepsilon(r)| + |\theta_h^\varepsilon(s)|) dr \right) \leq \frac{16T^2N}{2^n}. \quad (4.7)$$

Collecting the upper estimates from (4.2)-(4.7), we conclude the proof of (4.1). $\square$

Let ε_0 be defined as in Theorem 3.1 and $(h_\varepsilon, 0 < \varepsilon \leq \varepsilon_0)$ be a family of random elements taking values in $\mathcal{A}_M$. Let $\phi_{h_\varepsilon}^\varepsilon$ be the solution of the corresponding stochastic control equation with initial condition $\phi_{h_\varepsilon}^\varepsilon(0) = \xi \in H$:

$$d\phi_{h_\varepsilon}^\varepsilon + [A\phi_{h_\varepsilon}^\varepsilon + B(\phi_{h_\varepsilon}^\varepsilon) + R\phi_{h_\varepsilon}^\varepsilon]dt = \sigma(\phi_{h_\varepsilon}^\varepsilon)h_\varepsilon dt + \sqrt{\varepsilon} \sigma(\phi_{h_\varepsilon}^\varepsilon)dW(t). \quad (4.8)$$

Note that $\phi_{h_\varepsilon}^\varepsilon = \mathcal{G}^\varepsilon \left(\sqrt{\varepsilon} (W + \frac{1}{\sqrt{\varepsilon}} \int_0^\cdot h_\varepsilon(s)ds) \right)$ due to the uniqueness of the solution.

For all ω and $h \in L^2([0, T], H_0)$, let ϕ_h be the solution of the corresponding control equation (3.6) with initial condition $\phi_h(0) = \xi(\omega)$:

$$d\phi_h + [A\phi_h + B(\phi_h) + R\phi_h]dt = \sigma(\phi_h)h dt. \quad (4.9)$$

Note that here we may assume that h and ξ are random, but ϕ_h may be defined pointwise by (3.6).

Let $\mathcal{C}_0 = \{\int_0^\cdot h(s)ds : h \in L^2([0, T], H_0)\} \subset C([0, T], H_0)$. For every $\omega \in \Omega$, define $\mathcal{G}^0 : C([0, T], H_0) \rightarrow X$ by $\mathcal{G}^0(g)(\omega) = \phi_h(\omega)$ for $g = \int_0^\cdot h(s)ds \in \mathcal{C}_0$ and $\mathcal{G}^0(g) = 0$ otherwise.

Proposition 4.3. (*Weak convergence*)

Suppose that σ does not depend on time and satisfies Assumptions (A.1), (A.4) and (A.5). Let $\xi \in H$, be $\mathcal{F}_0$ -measurable such that $E|\xi|_H^4 < +\infty$, and let h_ε converge to h in distribution as random elements taking values in $\mathcal{A}_M$. (Note that here $\mathcal{A}_M$ is endowed with the weak topology induced by the norm defined in (3.4)). Then as $\varepsilon \rightarrow 0$, $\phi_{h_\varepsilon}^\varepsilon$ converges in distribution to ϕ_h in $X = C([0, T]; H) \cap L^2((0, T); V)$ endowed with norm (3.4). That is, $\mathcal{G}^\varepsilon\left(\sqrt{\varepsilon}(W + \frac{1}{\sqrt{\varepsilon}} \int_0^\cdot h_\varepsilon(s)ds)\right)$ converges in distribution to $\mathcal{G}^0(\int_0^\cdot h(s)ds)$ in X , as $\varepsilon \rightarrow 0$.

Proof. Since $\mathcal{A}_M$ is a Polish space (complete separable metric space), by the Skorokhod representation theorem, we can construct processes $(\tilde{h}_\varepsilon, \tilde{h}, \tilde{W})$ such that the joint distribution of $(\tilde{h}_\varepsilon, \tilde{W})$ is the same as that of (h_ε, W) , the distribution of $\tilde{h}$ coincides with that of h , and $\tilde{h}_\varepsilon \rightarrow \tilde{h}$, a.s., in the (weak) topology of S_M . Hence a.s. for every $t \in [0, T]$, $\int_0^t \tilde{h}_\varepsilon(s)ds - \int_0^t \tilde{h}(s)ds \rightarrow 0$ weakly in H_0 . Let $\Phi_\varepsilon = \phi_{h_\varepsilon}^\varepsilon - \phi_h$, or in component form $\Phi_\varepsilon = (U_\varepsilon, \Theta_\varepsilon) = (u_{h_\varepsilon}^\varepsilon - u_h, \theta_{h_\varepsilon}^\varepsilon - \theta_h)$; then

$$\begin{aligned} d\Phi_\varepsilon + [A\Phi_\varepsilon + B(\phi_{h_\varepsilon}^\varepsilon) - B(\phi_h) + R\Phi_\varepsilon]dt \\ = [\sigma(\phi_{h_\varepsilon}^\varepsilon)h_\varepsilon - \sigma(\phi_h)h]dt + \sqrt{\varepsilon}\sigma(\phi_{h_\varepsilon}^\varepsilon)dW(t), \quad \Phi_\varepsilon(0) = 0. \end{aligned} \quad (4.10)$$

Let ε_0 be defined as in Theorem 3.1. Set $\sigma_1 = \sigma$, $\sigma_2 = 0$, $\bar{\sigma} = \sigma$, $\bar{\sigma}_1 = 0$, $\bar{\sigma}_2(s) = \sigma(\phi_h(s))(h_\varepsilon(s) - h(s))$ and $\rho = 0$. Then $\phi_1 = \phi_{h_\varepsilon}^\varepsilon$ and $\phi_2 = \phi_h$ satisfy (3.30). Thus, Lemma 3.12, (A.4) and (A.5) yield for $0 \leq \varepsilon \leq \varepsilon_0 \wedge \frac{\nu \wedge \kappa}{4L}$:

$$\begin{aligned} |\Phi_\varepsilon(t)|^2 + (\nu \wedge \kappa) \int_0^t \|\Phi_\varepsilon(s)\|^2 ds &\leq \sum_{i=1}^3 T_i(t, \varepsilon) \\ &+ \int_0^t |\Phi_\varepsilon(s)|^2 \left[2 + \frac{8c_1^2}{\nu \wedge \kappa} \|\phi_h(s)\|^2 + \frac{2Lc_1c_2}{\nu \wedge \kappa} |h(s)|_0^2 \right] ds, \end{aligned} \quad (4.11)$$

where

$$\begin{aligned} T_1(t, \varepsilon) &= 2\sqrt{\varepsilon} \int_0^t (\Phi_\varepsilon(s), \sigma(\phi_{h_\varepsilon}^\varepsilon(s)) dW(s)) \\ T_2(t, \varepsilon) &= \varepsilon K \int_0^t (1 + |\phi_{h_\varepsilon}^\varepsilon(s)|^2) ds, \\ T_3(t, \varepsilon) &= 2 \int_0^t (\sigma(\phi_h(s)) (h_\varepsilon(s) - h(s)), \Phi_\varepsilon(s)) ds. \end{aligned}$$

Our goal here is to show that as $\varepsilon \rightarrow 0$, $\sup_{0 \leq t \leq T} |\Phi_\varepsilon(t)|^2 + \int_0^T \|\Phi_\varepsilon(s)\|^2 ds \rightarrow 0$ in probability, which implies that $\phi_{h_\varepsilon} \rightarrow \phi_h$ in distribution in $X := C([0, T]; H) \cap$

$L^2((0, T); V)$. Fix $N > 0$ and for $t \in [0, T]$ let

$$\begin{aligned} G_N(t) &= \left\{ \sup_{0 \leq s \leq t} |\phi_h(s)|^2 \leq N \right\} \cap \left\{ \int_0^t \|\phi_h(s)\|^2 ds \leq N \right\}, \\ G_{N,\varepsilon}(t) &= G_N(t) \cap \left\{ \sup_{0 \leq s \leq t} |\phi_{h_\varepsilon}^\varepsilon(s)|^2 \leq N \right\} \cap \left\{ \int_0^t \|\phi_{h_\varepsilon}^\varepsilon(s)\|^2 ds \leq N \right\}. \end{aligned}$$

Claim 1. For any $\varepsilon_0 > 0$, $\sup_{0 < \varepsilon \leq \varepsilon_0} \sup_{h, h_\varepsilon \in \mathcal{A}_M} \mathbb{P}(G_{N,\varepsilon}(T)^c) \rightarrow 0$ as $N \rightarrow \infty$.

Indeed, for $\varepsilon > 0$, $h, h_\varepsilon \in \mathcal{A}_M$, the Markov inequality and estimate (3.5) imply

$$\begin{aligned} \mathbb{P}(G_{N,\varepsilon}(T)^c) &\leq \mathbb{P}\left(\sup_{0 \leq s \leq T} |\phi_h(s)|^2 > N\right) + \mathbb{P}\left(\sup_{0 \leq s \leq T} |\phi_{h_\varepsilon}^\varepsilon(s)|^2 > N\right) \\ &\quad + \mathbb{P}\left(\int_0^T (\|\phi_h(s)\|^2 ds > N)\right) + \mathbb{P}\left(\int_0^T \|\phi_{h_\varepsilon}^\varepsilon(s)\|^2 ds > N\right) \\ &\leq \frac{1}{N} \sup_{h, h_\varepsilon \in \mathcal{A}_M} \mathbb{E}\left(\sup_{0 \leq s \leq T} |\phi_h(s)|^2 + \sup_{0 \leq s \leq T} |\phi_{h_\varepsilon}^\varepsilon(s)|^2 + \int_0^T (\|\phi_h(s)\|^2 + \|\phi_{h_\varepsilon}^\varepsilon(s)\|^2) ds\right) \\ &\leq C_1(\nu, \kappa, K, L, T, M) (1 + E|\xi|^4) N^{-1}. \end{aligned}$$

Claim 2. For fixed $N > 0$, $h, h_\varepsilon \in \mathcal{A}_M$ such that as $\varepsilon \rightarrow 0$, $h_\varepsilon \rightarrow h$ a.s. in the weak topology of $L^2([0, T], H_0)$, one has as $\varepsilon \rightarrow 0$

$$\mathbb{E}\left[1_{G_{N,\varepsilon}(T)} \left(\sup_{0 \leq t \leq T} |\Phi_\varepsilon(t)|^2 + \int_0^T \|\Phi_\varepsilon(t)\|^2 dt\right)\right] \rightarrow 0. \quad (4.12)$$

Indeed, (4.11) and Gronwall's lemma imply that on $G_{N,\varepsilon}(T)$,

$$\sup_{0 \leq t \leq T} |\Phi_\varepsilon(t)|^2 \leq \left[\sup_{0 \leq t \leq T} (T_1(t, \varepsilon) + T_3(t, \varepsilon)) + \varepsilon K T (1 + N) \right] e^{2T + \frac{8c_1^2 N}{\nu \wedge \kappa} + \frac{2Lc_1 c_2 M}{\nu \wedge \kappa}}.$$

Thus, using again (4.11) we deduce that for some constant $\tilde{C} = C(\nu, \kappa, K, L, T, M, N)$, one has for every $\varepsilon > 0$:

$$\mathbb{E}(1_{G_{N,\varepsilon}(T)} |\Phi_\varepsilon|_X^2) \leq \tilde{C} \left(\varepsilon K T (1 + N) + \mathbb{E}\left[1_{G_{N,\varepsilon}(T)} \sup_{0 \leq t \leq T} (T_1(t, \varepsilon) + T_3(t, \varepsilon))\right] \right). \quad (4.13)$$

Since the sets $G_{N,\varepsilon}(\cdot)$ decrease, $\mathbb{E}(1_{G_{N,\varepsilon}(T)} \sup_{0 \leq t \leq T} |T_1(t, \varepsilon)|) \leq \mathbb{E}(\lambda_\varepsilon)$, where

$$\lambda_\varepsilon := 2\sqrt{\varepsilon} \sup_{0 \leq t \leq T} \left| \int_0^t 1_{G_{N,\varepsilon}(s)} (\Phi_\varepsilon(s), \sigma(\phi_{h_\varepsilon}^\varepsilon(s)) dW(s)) \right|.$$

The scalar-valued random variables λ_ε converge to 0 in L^1 as $\varepsilon \rightarrow 0$. Indeed, by the Burkholder-Davis-Gundy inequality, (A.4) and the definition of $G_{N,\varepsilon}(s)$, we have

$$\begin{aligned} \mathbb{E}(\lambda_\varepsilon) &\leq 6\sqrt{\varepsilon} \mathbb{E}\left\{ \int_0^T 1_{G_{N,\varepsilon}(s)} |\Phi_\varepsilon(s)|^2 |\sigma(\phi_{h_\varepsilon}^\varepsilon(s))|_{L_Q}^2 ds \right\}^{\frac{1}{2}} \\ &\leq 6\sqrt{\varepsilon} \mathbb{E}\left[\left\{ 4N \int_0^T 1_{G_{N,\varepsilon}(s)} K (1 + |\phi_{h_\varepsilon}^\varepsilon(s)|^2) ds \right\}^{\frac{1}{2}} \right] \end{aligned}$$

$$\leq 12\sqrt{\varepsilon}\sqrt{KT}(1+N). \quad (4.14)$$

For $k = 0, \dots, 2^n$ set $t_k = kT2^{-n}$; for $s \in]t_k, t_{k+1}]$, set $\bar{s}_n = t_{k+1}$ and $\underline{s}_n = t_k$. Then for any $n \geq 1$,

$$\mathbb{E}\left(1_{G_{N,\varepsilon}(T)} \sup_{0 \leq t \leq T} |T_3(t, \varepsilon)|\right) \leq 2 \sum_{i=1}^3 \tilde{T}_i(N, n, \varepsilon) + 2 \mathbb{E}(\bar{T}_4(N, n, \varepsilon, \omega)),$$

where

$$\begin{aligned} \tilde{T}_1(N, n, \varepsilon) &= \mathbb{E}\left[1_{G_{N,\varepsilon}(T)} \sup_{0 \leq t \leq T} \left| \int_0^t \left(\sigma(\phi_h(s))(h_\varepsilon(s) - h(s)), [\Phi_\varepsilon(s) - \Phi_\varepsilon(\bar{s}_n)] \right) ds \right| \right], \\ \tilde{T}_2(N, n, \varepsilon) &= \mathbb{E}\left[1_{G_{N,\varepsilon}(T)} \sup_{0 \leq t \leq T} \left| \int_0^t \left([\sigma(\phi_h(s)) - \sigma(\phi_h(\bar{s}_n))](h_\varepsilon(s) - h(s)), \Phi_\varepsilon(\bar{s}_n) \right) ds \right| \right], \\ \tilde{T}_3(N, n, \varepsilon) &= \mathbb{E}\left[1_{G_{N,\varepsilon}(T)} \sup_{1 \leq k \leq 2^n} \sup_{t_{k-1} \leq t \leq t_k} \left| \left(\sigma(\phi_h(t_k)) \int_{t_{k-1}}^t (h_\varepsilon(s) - h(s)) ds, \Phi_\varepsilon(t_k) \right) \right| \right] \\ \bar{T}_4(N, n, \varepsilon) &= 1_{G_{N,\varepsilon}(T)} \sum_{k=1}^{2^n} \left| \left(\sigma(\phi_h(t_k)) \int_{t_{k-1}}^{t_k} (h_\varepsilon(s) - h(s)) ds, \Phi_\varepsilon(t_k) \right) \right|. \end{aligned}$$

Using Schwarz's inequality, (A.4) and Lemma 4.2, we deduce that for some constant $\bar{C}_1 := C(\nu, \kappa, K, T, M, N)$ and any $\varepsilon \in]0, \varepsilon_0]$,

$$\begin{aligned} \tilde{T}_1(N, n, \varepsilon) &\leq \sqrt{K} \mathbb{E}\left[1_{G_{N,\varepsilon}(T)} \int_0^T (1 + |\phi_h(s)|^2)^{\frac{1}{2}} |h_\varepsilon(s) - h(s)|_0 |\Phi_\varepsilon(s) - \Phi_\varepsilon(\bar{s}_n)| ds\right] \\ &\leq \sqrt{2K(1+N)} \left(\mathbb{E} \int_0^T |h_\varepsilon(s) - h(s)|_0^2 ds \right)^{\frac{1}{2}} \\ &\quad \times \left(\mathbb{E}\left[1_{G_{N,\varepsilon}(T)} \int_0^T \{|\phi_{h_\varepsilon}^\varepsilon(s) - \phi_{h_\varepsilon}^\varepsilon(\bar{s}_n)|^2 + |\phi_h(s) - \phi_h(\bar{s}_n)|^2\} ds\right] \right)^{\frac{1}{2}} \\ &\leq \bar{C}_1 2^{-\frac{n}{4}}. \end{aligned} \quad (4.15)$$

A similar computation based on (A.5) and Lemma 4.2 yields for some constant $\bar{C}_2 := C(\nu, \kappa, K, L, T, M, N)$ and any $\varepsilon \in]0, \varepsilon_0]$

$$\begin{aligned} \tilde{T}_2(N, n, \varepsilon) &\leq \sqrt{L} \left(\mathbb{E}\left[1_{G_{N,\varepsilon}(T)} \int_0^T |\phi_h(s) - \phi_h(\bar{s}_n)|^2 ds\right] \right)^{\frac{1}{2}} \left(\mathbb{E} \int_0^T |h_\varepsilon(s) - h(s)|_0^2 4N ds \right)^{\frac{1}{2}} \\ &\leq \bar{C}_2 2^{-\frac{n}{4}}. \end{aligned} \quad (4.16)$$

Using Schwarz's inequality and (A.4) we deduce for $\bar{C}_3 = C(K, N, M)$ and any $\varepsilon \in]0, \varepsilon_0]$

$$\begin{aligned} \tilde{T}_3(N, n, \varepsilon) &\leq \sqrt{K} \mathbb{E}\left[1_{G_{N,\varepsilon}(T)} \sup_{1 \leq k \leq 2^n} (1 + |\phi_h(t_k)|^2)^{\frac{1}{2}} \int_{t_{k-1}}^{t_k} |h_\varepsilon(s) - h(s)|_0 ds |\Phi_\varepsilon(t_k)|\right] \\ &\leq 2\sqrt{KN(1+N)} \mathbb{E}\left(\sup_{1 \leq k \leq 2^n} \int_{t_{k-1}}^{t_k} |h_\varepsilon(s) - h(s)|_0 ds\right) \end{aligned}$$

$$\leq 8\sqrt{KN(1+N)}\sqrt{M}2^{-\frac{n}{2}} = \bar{C}_3 2^{-\frac{n}{2}}. \quad (4.17)$$

Finally, note that the weak convergence of h_ε to h implies that for any $a, b \in [0, T]$, $a < b$, as $\varepsilon \rightarrow 0$, the integral $\int_a^b h_\varepsilon(s)ds \rightarrow \int_a^b h(s)ds$ in the weak topology of H_0 . Therefore, since for $\phi \in H$ the operator $\sigma(\phi)$ is compact from H_0 to H , we deduce that $\left| \sigma(\phi) \left(\int_a^b h_\varepsilon(s)ds - \int_a^b h(s)ds \right) \right|_H \rightarrow 0$ as $\varepsilon \rightarrow 0$. Hence a.s. for fixed n as $\varepsilon \rightarrow 0$, $\bar{T}_4(N, n, \varepsilon, \omega) \rightarrow 0$. Furthermore, $\bar{T}_4(N, n, \varepsilon, \omega) \leq \sqrt{K}\sqrt{1+N}\sqrt{4N}\sqrt{M}$ and hence the dominated convergence theorem proves that for any fixed n , $\mathbb{E}(\bar{T}_4(N, n, \varepsilon)) \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Thus, given $\alpha > 0$, we may choose n_0 large enough to have $(\bar{C}_1 + \bar{C}_2)2^{-\frac{n}{4}} + \bar{C}_3 2^{-\frac{n}{2}} \leq \alpha$ for $n \geq n_0$. Then for fixed $n \geq n_0$, let $\varepsilon_1 \in]0, \varepsilon_0]$ be such that for $0 < \varepsilon \leq \varepsilon_1$, $\mathbb{E}[\bar{T}_4(N, n, \varepsilon)] \leq \alpha$. Using (4.15)-(4.17), we deduce that for $\varepsilon \in]0, \varepsilon_1]$,

$$\mathbb{E} \left[1_{G_{N,\varepsilon}(T)} \sup_{0 \leq t \leq T} |T_3(t, \varepsilon)| \right] \leq 2\alpha. \quad (4.18)$$

Claim 2 is a straightforward consequence of inequalities (4.13), (4.14) and (4.18).

To conclude the proof of Proposition 4.3, let $\delta > 0$ and $\alpha > 0$ and set

$$\Lambda_\varepsilon := |\Phi_\varepsilon|_X^2 = \sup_{0 \leq t \leq T} |\Phi_\varepsilon(t)|^2 + \int_0^T \|\Phi_\varepsilon(s)\|^2 ds.$$

Then the Markov inequality implies that

$$\mathbb{P}(\Lambda_\varepsilon > \delta) = \mathbb{P}(G_{N,\varepsilon}(T)^c) + \frac{1}{\delta} \mathbb{E} \left(1_{G_{N,\varepsilon}(T)} |\Phi_\varepsilon|_X^2 \right)$$

Using *Claim 1*, one can choose N large enough to make sure that $\mathbb{P}(G_{N,\varepsilon}(T)^c) < \alpha$ for every $\varepsilon \leq \varepsilon_0$. Fix N ; *Claim 2* shows that for ε small enough, $\mathbb{E} \left(1_{G_{N,\varepsilon}(T)} |\Phi_\varepsilon|_X^2 \right) < \delta\alpha$. This concludes the proof of the proposition. $\square$

The following compactness result will show that the rate function of the LDP satisfied by the solution to (4.8) is a good rate function. The proof is similar to that of Proposition 4.3 and easier.

Proposition 4.4. (*Compactness*)

Let M be any fixed finite positive number and let $\xi \in H$ be deterministic. Define

$$K_M = \{\phi_h \in C([0, T]; H) \cap L^2((0, T); V) : h \in S_M\},$$

where ϕ_h is the unique solution of the deterministic control equation:

$$d\phi_h(t) + [A\phi_h(t) + B(\phi_h(t)) + R\phi_h(t)]dt = \sigma(\phi_h(t))h(t)dt, \quad \phi_h(0) = \xi, \quad (4.19)$$

and σ does not depend on time and satisfies (A.1), (A.4) and (A.5). Then K_M is a compact subset of X .

Proof. Let (ϕ_n) be a sequence in K_M , corresponding to solutions of (4.19) with controls (h_n) in S_M :

$$d\phi_n(t) + [A\phi_n(t) + B(\phi_n(t)) + R\phi_n(t)]dt = \sigma(\phi_n(t))h_n(t)dt, \quad \phi_n(0) = \xi. \quad (4.20)$$

Since S_M is a bounded closed subset in the Hilbert space $L^2((0, T); H_0)$, it is weakly compact. So there exists a subsequence of (h_n) , still denoted as (h_n) , which converges weakly to a limit h in $L^2((0, T); H_0)$. Note that in fact $h \in S_M$ as S_M is closed. We now show that the corresponding subsequence of solutions, still denoted as (ϕ_n) , converges in X to ϕ which is the solution of the following “limit” equation

$$d\phi(t) + [A\phi(t) + B(\phi(t)) + R\phi(t)]dt = \sigma(\phi(t))h(t)dt, \quad \phi(0) = \xi. \quad (4.21)$$

This will complete the proof of the compactness of K_M . To ease notation we will often drop the time parameters $s, t, \dots$ in the equations and integrals.

Let $\Phi_n = \phi_n - \phi$, or in component form $\Phi_n = (U_n, \Theta_n) = (u_n - u, \theta_n - \theta)$; then

$$d\Phi_n + [A\Phi_n + B(\phi_n) - B(\phi) + R\Phi_n]dt = [\sigma(\phi_n)h_n - \sigma(\phi)h]dt, \quad \Phi_n(0) = 0. \quad (4.22)$$

Set $\sigma_1 = \sigma_2 = 0$, $\bar{\sigma} = \sigma$, $\bar{\sigma}_1 = 0$, $\bar{\sigma}_2(s) = \sigma(\phi(s)) [h(s) - h_n(s)]$, $h_\varepsilon = h_n$, $\rho = 0$. Then $\phi_1 := \phi_n$ and $\phi_2 := \phi$ satisfy (3.30).

Thus Lemma 3.12 yields the following integral inequality

$$\begin{aligned} |\Phi_n(t)|^2 + (\nu \wedge \kappa) \int_0^t \|\Phi_n(s)\|^2 ds &\leq 2 \int_0^t (\sigma(\phi(s)) [h(s) - h_n(s)], \Phi_n(s)) ds \\ &+ \int_0^t \left\{ 2 + \frac{8c_1^2}{\nu \wedge \kappa} \|\phi(s)\|^2 + \frac{2Lc_1c_2}{\nu \wedge \kappa} |h_n(s)|_0^2 \right\} |\Phi_n(s)|^2 ds. \end{aligned} \quad (4.23)$$

For $N \geq 1$ and $k = 0, \dots, 2^N$, set $t_k = k2^{-N}$. For $s \in]t_{k-1}, t_k]$, $1 \leq k \leq 2^N$, let $\bar{s}_N = t_k$. Inequality (3.7) implies that there exists a constant $\bar{C} > 0$ such that

$$\sup_n \left[\sup_{0 \leq t \leq T} (|\phi(t)|^2 + |\phi_n(t)|^2) + \int_0^T (\|\phi(s)\|^2 + \|\phi_n(s)\|^2) ds \right] = \bar{C} < +\infty.$$

Thus Gronwall's inequality implies

$$\sup_{t \leq T} |\Phi_n(t)|^2 \leq \exp \left(2T + \frac{8c_1^2 \bar{C}}{\nu \wedge \kappa} + \frac{2Lc_1c_1M}{\nu \wedge \kappa} \right) \sum_{i=1}^4 I_{n,N}^i, \quad (4.24)$$

where

$$\begin{aligned} I_{n,N}^1 &= \int_0^T |(\sigma(\phi(s)) [h_n(s) - h(s)], \Phi_n(s) - \Phi_n(\bar{s}_N))| ds, \\ I_{n,N}^2 &= \int_0^T \left| \left([\sigma(\phi(s)) - \sigma(\phi(\bar{s}_N))] [h_n(s) - h(s)], \Phi_n(\bar{s}_N) \right) \right| ds, \\ I_{n,N}^3 &= \sup_{1 \leq k \leq 2^N} \sup_{t_{k-1} \leq t \leq t_k} \left| \left(\sigma(\phi(t_k)) \int_{t_{k-1}}^t (h_n(s) - h(s)) ds, \Phi_n(t_k) \right) \right|, \\ I_{n,N}^4 &= \left| \sum_{k=1}^{2^N} \left(\sigma(\phi(t_k)) \int_{t_{k-1}}^{t_k} [h_n(s) - h(s)] ds, \Phi_n(t_k) \right) \right|. \end{aligned}$$

Schwarz's inequality, (A.4), (A.5) and Lemma 4.2 imply that for some constant C which does not depend on n and N ,

$$\begin{aligned} I_{n,N}^1 &\leq \left(\int_0^T K(1 + \bar{C}) |h_n(s) - h(s)|_0^2 ds \right)^{\frac{1}{2}} \\ &\quad \times \left(2 \int_0^T (|\phi_n(s) - \phi_n(\bar{s}_N)|^2 + |\phi(s) - \phi(\bar{s}_N)|^2) ds \right)^{\frac{1}{2}} \\ &\leq C 2^{-\frac{N}{4}}, \end{aligned} \quad (4.25)$$

$$I_{n,N}^2 \leq \left(L \int_0^T |\phi(s) - \phi(\bar{s}_N)|^2 ds \right)^{\frac{1}{2}} \left(\bar{C} \int_0^T |h_n(s) - h(s)|_0^2 ds \right)^{\frac{1}{2}} \leq C 2^{-\frac{N}{4}}, \quad (4.26)$$

$$I_{n,N}^3 \leq K \left(1 + \sup_t |\phi(t)| \right) \sup_t (|\phi(t)| + \phi_n(t)) 2^{-\frac{N}{2}} 2M \leq C 2^{-\frac{N}{2}}. \quad (4.27)$$

Thus, given $\alpha > 0$, one may choose N large enough to have $\sup_n \sum_{i=1}^3 I_{n,N}^i \leq \alpha$. Then, for fixed N and $k = 1, \dots, 2^N$, as $n \rightarrow \infty$, the weak convergence of h_n to h implies that of $\int_{t_{k-1}}^{t_k} (h_n(s) - h(s)) ds$ to 0 weakly in H_0 . Since $\sigma(\phi(t_k))$ is a compact operator, we deduce that for fixed k the sequence $\sigma(\phi(t_k)) \int_{t_{k-1}}^{t_k} (h_n(s) - h(s)) ds$ converges to 0 strongly in H as $n \rightarrow \infty$. Since $\sup_n \sup_k |\Phi_n(t_k)| \leq 2\tilde{C}$, we have $\lim_n I_{n,N}^4 = 0$. Thus as $n \rightarrow \infty$, $\sup_{0 \leq t \leq T} |\Phi_n(t)|^2 \rightarrow 0$. Using this convergence and (4.24), we deduce that $\|\Phi_n\|_X \rightarrow 0$ as $n \rightarrow \infty$. This shows that every sequence in K_M has a convergent subsequence. Hence K_M is a compact subset of X . $\square$

With the above results, we have the following large deviation theorem.

Theorem 4.5. *Suppose that σ does not depend on time and satisfies (A.1), (A.4) and (A.5), let ϕ^ε be the solution of the stochastic Bénard problem (2.9). Then $\{\phi^\varepsilon\}$ satisfies the large deviation principle in $C([0, T]; H) \cap L^2((0, T); V)$, with the good rate function*

$$I_\xi(\psi) = \inf_{\{h \in L^2(0, T; H_0) : \psi = \mathcal{G}^0(\int_0^\cdot h(s) ds)\}} \left\{ \frac{1}{2} \int_0^T |h(s)|_0^2 ds \right\}. \quad (4.28)$$

Here the infimum of an empty set is taken as infinity.

Proof. Propositions 4.4 and 4.3 imply that $\{\phi^\varepsilon\}$ satisfies the Laplace principle which is equivalent to the large deviation principle in $X = C([0, T], H) \cap L^2((0, T), V)$ with the above-mentioned rate function; see Theorem 4.4 in [1] or Theorem 5 in [2]. $\square$

Acknowledgements. This work was partially done while J. Duan was a Professor Invité at the Université Paris 1, France. J. Duan would like to thank this University for financial support and the colleagues at Samos-Matisse, Centre d'Économie de la Sorbonne, for their hospitality. It was completed while A. Millet was visiting the Mittag Leffler Institute, Sweden, which provided financial support. She would like to thank the center for excellent working conditions and a very friendly atmosphere.

The authors would also like to thank Padma Sundar for helpful discussions and an anonymous referee for his careful reading and helpful comments.

REFERENCES

- [1] A. Budhiraja & P. Dupuis, A variational representation for positive functionals of infinite dimensional Brownian motion, *Prob. and Math. Stat.* **20** (2000), 39-61.
- [2] A. Budhiraja, P. Dupuis & V. Maroulas, Large deviations for infinite dimensional stochastic dynamical systems, *Ann. Prob.* **36** (4) (2008), 1390-1420.
- [3] S. Cerrai and M. Rockner, Large deviations for stochastic reaction-diffusion systems with multiplicative noise and non-Lipschitz reaction term, *Ann. Prob.* **32** (2004), 1100-1139.
- [4] F. Chenal and A. Millet, Uniform large deviations for parabolic SPDEs and applications. *Stochastic Process. Appl.* **72** (1997), no. 2, 161-186.
- [5] M. H. Chang, Large deviations for the Navier-Stokes equations with small stochastic perturbations. *Appl. Math. Comput.* **76** (1996), 65-93.
- [6] P. L. Chow, Large deviation problem for some parabolic Ito equations. *Comm. Pure Appl. Math.* **45** (1992), 97-120.
- [7] P. Constantin and C. Foias, *Navier-Stokes Equations*, U. of Chicago Press, Chicago, 1988.
- [8] G. Da Prato & J. Zabczyk, *Stochastic Equations in Infinite Dimensions*, Cambridge University Press, 1992.
- [9] H. A. Dijkstra, *Nonlinear Physical Oceanography*, Kluwer Academic Publishers, Boston, 2000.
- [10] J. Duan, H. Gao and B. Schmalfuss, Stochastic Dynamics of a Coupled Atmosphere-Ocean Model, *Stochastics and Dynamics* **2** (2002), 357-380.
- [11] J. Feng and T. G. Kurtz, *Large Deviations for Stochastic Processes*. AMS, 2007.
- [12] B. Ferrario, The Bénard Problem with random perturbations: Dissipativity and invariant measures. *Nonlinear Differential Equations and Applications (NoDEA)* **4** (1997), 101-121.
- [13] C. Foias, O. Manley & R. Temam, Attractors for the Bénard Problem: Existence and physical bounds on their fractal dimension. *Nonlinear Analysis* **11** (1987), 939-967.
- [14] M. I. Freidlin & A. D. Wentzell, Reaction-diffusion equation with randomly perturbed boundary condition, *The Annals. of Prob.* **20**(2) (1992), 963-986.
- [15] G. Kallianpur and J. Xiong, Large deviations for a class of stochastic partial differential equations. *Ann. Prob.* **24** (1996), 320-345.
- [16] H. Kunita, *Stochastic flows and stochastic differential equations*. Cambridge ; New York : Cambridge University Press, 1990.
- [17] T. Ozgokmen, T. Iliescu, P. Fischer, A. Srinivasan and J. Duan. Large eddy simulation of stratified mixing in two-dimensional dam-break problem in a rectangular enclosed domain. *Ocean Modeling* **16** (2007), 106-140.
- [18] S. Peszat, Large deviation estimates for stochastic evolution equations. *Prob. Theory Rel. Fields.* **98** (1994), 113-136.
- [19] J. Ren and X. Zhang, Freidlin-Wentzell's large deviations for homeomorphism flows of non-Lipschitz SDEs. *Bull. Sci. Math.* **129** (2005), 643-655.
- [20] B. L. Rozovskii, *Stochastic Evolution Equations*. Kluwer Academic Publishers, Boston, 1990.
- [21] R. Sowers, Large deviations for a reaction-diffusion system with non-Gaussian perturbations. *Ann. Prob.* **20** (1992), 504-537.
- [22] S. S. Sritharan & P. Sundar, Large deviations for the two-dimensional Navier-Stokes equations with multiplicative noise, *Stoch. Proc. and Appl.* **116** (2006), 1636-1659.
- [23] R. Temam, *Navier-Stokes Equations and Nonlinear Functional Analysis*, 2nd Edition, SIAM, Philadelphia, 1995.
- [24] E. Waymire & J. Duan (Eds.), *Probability and Partial Differential Equations in Modern Applied Mathematics*. Springer-Verlag, New York, 2005.

- [25] W. Wang & J. Duan, Reductions and deviations for stochastic partial differential equations under fast dynamical boundary conditions. *Stochastic Analysis and Applications*, accepted, 2008.
- [26] J. Zabczyk, On large deviations for stochastic evolution equations. Stochastic Systems and Optimization. *Lecture Notes on Control and Inform. Sci.*, Springer, Berlin, 1988.

(J. Duan) DEPARTMENT OF APPLIED MATHEMATICS, ILLINOIS INSTITUTE OF TECHNOLOGY, CHICAGO, IL 60616, USA

E-mail address, J. Duan: `duan@iit.edu`

(A. Millet) SAMOS-MATISSE, CENTRE D'ÉCONOMIE DE LA SORBONNE (UMR 8174), UNIVERSITÉ PARIS 1, CENTRE PIERRE MENDÈS FRANCE, 90 RUE DE TOLBIAC, F- 75634 PARIS CEDEX 13, FRANCE *and* LABORATOIRE DE PROBABILITÉS ET MODÈLES ALÉATOIRES (UMR 7599), UNIVERSITÉS PARIS 6-PARIS 7, BOÎTE COURRIER 188, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE

E-mail address, A. Millet: `amillet@univ-paris1.fr` *and* `annie.millet@upmc.fr`