



**HAL**  
open science

# Existence of travelling-wave solutions and local well-posedness of the Fowler equation

Borys Alvarez-Samaniego, Pascal Azerad

► **To cite this version:**

Borys Alvarez-Samaniego, Pascal Azerad. Existence of travelling-wave solutions and local well-posedness of the Fowler equation. 2008. hal-00257423v1

**HAL Id: hal-00257423**

**<https://hal.science/hal-00257423v1>**

Preprint submitted on 19 Feb 2008 (v1), last revised 12 Mar 2009 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# EXISTENCE OF TRAVELLING-WAVE SOLUTIONS AND LOCAL WELL-POSEDNESS OF THE FOWLER EQUATION

BORYS ALVAREZ-SAMANIEGO AND PASCAL AZERAD

ABSTRACT. We study the existence of travelling-waves and local well-posedness in a subspace of  $C_b^1(\mathbb{R})$  for a nonlinear evolution equation recently proposed by Andrew C. Fowler to study the dynamics of dunes.

## 1. INTRODUCTION

1.1. **General setting.** Dunes are land formations of sand which are subject to different forms and sizes based on their interaction with the wind or water or some other mobile medium. In the case of dunes in the desert their shapes depend mainly on the amount of sand available and on the change of the direction of the wind with time (see Herrmann and Sauermann [6]). Some examples of dune patterns are longitudinal, transverse, star and Barchan dunes, however, there are more than 100 categories of dunes.

An interesting topic is to try to understand if the shape of a dune is maintained when it moves. With regard to Barchan dunes, for example, Herrmann and Sauermann [6] have given some arguments, mainly related with the fact that Barchan dunes constantly lose sand at the two horns and tend to disappear if these dunes are not supplied with new sand, against the hypothesis that Barchan dunes are solitary waves. Recently, Durán, Schwämmle and Herrmann [2] considered a minimal model for dunes consisting of three coupled equations of motion to study, from a numerical point of view, the mechanisms of dune interactions for the case when a small Barchan dune collides with a bigger one; four different cases were observed, depending only on the relative sizes of the two dunes, namely, coalescence, breeding, budding, and solitary wave behavior.

In this paper, we are concerned with the following evolution equation proposed by Fowler (see [3], [4] and [5] for more details) to study nonlinear dune formation:

$$\frac{\partial u}{\partial t}(x, t) + \frac{\partial}{\partial x} \left[ \frac{u^2}{2}(x, t) - \frac{\partial u}{\partial x}(x, t) + \int_0^{+\infty} \xi^{-1/3} \frac{\partial u}{\partial x}(x - \xi, t) d\xi \right] = 0, \quad (1.1)$$

where  $u = u(x, t)$  represents the dune amplitude,  $x \in \mathbb{R}$ , and  $t \geq 0$ . The second and fourth terms of equation (1.1) correspond to the nonlinear and nonlocal terms respectively, while the third term is the dissipative term. To the authors' knowledge, ours is the first study to report a rigorous mathematical proof of the existence of travelling-waves for dune morphodynamics. The authors hope that these results could be of interest for geographers, geologists, oceanographers and others.

---

2000 *Mathematics Subject Classification.* 47J35; 35G25; 76B25.

*Key words and phrases.* Nonlocal evolution equation; travelling-wave.

**1.2. Organization of the paper.** In Section 2 we study the existence of travelling-wave solutions to equation (1.1). The main result of this section is Theorem 2.1 which implies that there exists a travelling-wave solution  $u(x, t) = \phi(x - ct)$  to equation (2.5), for  $\eta$  in a neighborhood of zero, where  $\phi \in C_b^1(\mathbb{R})$ , and  $c > 0$ ; the idea of its proof is to use the implicit function theorem on suitable Banach spaces. Then, by a scaling argument we extend this for any  $\eta \in \mathbb{R}$ .

Section 3 is devoted to proving local well-posedness (LWP) for the integral equation associated to the initial value problem (IVP) for equation (1.1). Inspired by the regularity of the travelling-wave obtained in Section 2, we consider a suitable subspace of  $C_b^1(\mathbb{R})$ . Our main result of this section is stated in Theorem 3.1; it gives local-in-time existence of the solution of the integral equation associated to the IVP (3.1), with initial data  $u(\cdot, 0) = u_0(\cdot)$  belonging to the subspace  $X$  of  $C_b^1(\mathbb{R})$ , where

$$X := \{f \in C_b^1(\mathbb{R}); f' \text{ is uniformly continuous}\}.$$

**1.3. Notations.** - We denote by  $\mathbb{R}$  and  $\mathbb{C}$  the sets of all real and complex numbers respectively.  $\mathbb{N}$  denotes the set of all natural numbers.

- We denote by  $C(c_1, c_2, \dots)$  a constant which depends on the parameters  $c_1, c_2, \dots$ .  $C$  is assumed to be a non-decreasing function of its arguments.

- The norm of a measurable function  $f \in L^p(\Omega)$ , for  $\Omega$  a subset of  $\mathbb{R}$ , is written  $\|f\|_{L^p(\Omega)}^p = \int_{\Omega} |f|^p dx$  for  $1 \leq p < +\infty$ , and  $\|f\|_{L^\infty(\Omega)} = \text{ess sup}_{\Omega} |f|$ . The inner product of two functions  $f, g \in L^2(\Omega)$  is written as  $(f, g) = \int_{\Omega} f \bar{g} dx$ . We will often omit set  $\Omega$  when context is clear.

- We denote by  $\hat{f} = \mathcal{F}f$  the Fourier transform of  $f$  ( $\mathcal{F}^{-1}$  and  $\check{\cdot}$  are used to denote the inverse of the Fourier transform), where  $\hat{f}(\xi) := \frac{1}{\sqrt{2\pi}} \int e^{-i\xi x} f(x) dx$  for  $f \in L^1(\mathbb{R})$  (it follows that  $\widehat{f * g} = \sqrt{2\pi} \hat{f} \hat{g}$  for  $f, g \in L^1(\mathbb{R})$ ).

- The Schwartz space of rapidly decreasing functions on  $\mathbb{R}$  is denoted  $\mathcal{S}(\mathbb{R})$ .

- We denote  $\Lambda := (1 - \partial_x^2)^{1/2}$  and  $H^s(\mathbb{R})$  ( $s \in \mathbb{R}$ ) the usual Sobolev space  $H^s(\mathbb{R}) = \{u \in \mathcal{S}'(\mathbb{R}), \|u\|_{H^s} < \infty\}$ , where  $\|u\|_{H^s} = \|\Lambda^s u\|_{L^2}$ .

- Let  $\Omega \subset \mathbb{R}$ .  $C^0(\Omega) = C(\Omega)$  is used to denote the space of all continuous complex-valued functions on  $\Omega$ . Moreover,  $C^k(\Omega) = \{u : \Omega \mapsto \mathbb{C} ; u, u', \dots, u^{(k)} \in C^0(\Omega)\}$ , for  $k \in \mathbb{N}$ . We write  $C^\infty(\Omega)$  to denote the set of infinitely differentiable complex-valued functions on  $\Omega$ . Similarly, we use the notations  $C^0(\Omega; Y) = C(\Omega; Y)$ ,  $C^k(\Omega; Y)$ ,  $C^\infty(\Omega; Y)$  when functions take values in the Banach space  $Y$ .

- We write  $C_\infty(\mathbb{R})$  to denote the space of all continuous complex-valued functions defined on  $\mathbb{R}$  which tend to zero at infinity.

- We denote by  $C_b(\mathbb{R}) = C_b^0(\mathbb{R})$  the space of all bounded continuous real-valued functions on  $\mathbb{R}$  with the norm  $\|\cdot\|_{L^\infty}$ . Moreover, for every  $k \in \mathbb{N}$ , we write

$$C_b^k(\mathbb{R}) := \{f \in C^k(\mathbb{R}) ; f, f', \dots, f^{(k)} \in C_b(\mathbb{R})\},$$

where  $\|f\|_{C_b^k} := \sum_{i=0}^k \|f^{(i)}\|_{L^\infty}$ , for all  $f \in C_b^k(\mathbb{R})$ .

- If  $X$  and  $Y$  are two Banach spaces, we denote by  $\mathfrak{L}(X, Y)$  the set of all continuous linear mappings defined on  $X$  with values in  $Y$ ; if  $X = Y$ , we denote by  $\mathfrak{L}(X)$ .

## 2. EXISTENCE OF TRAVELLING-WAVE SOLUTIONS OF THE FOWLER EQUATION

We begin this section with some notations and preliminary results. We define

$$\psi(x) := \chi_{(0, \infty)}(x) \cdot x^{-1/3}, \quad \text{for all } x \in \mathbb{R}, \quad (2.1)$$

where  $\chi_A$  is used to denote the characteristic function of the set  $A$ . We also define

$$g[u] := \psi * \partial_x u. \quad (2.2)$$

We note that, since  $\psi \in \mathcal{S}'(\mathbb{R})$ , it follows that for  $\phi \in \mathcal{S}(\mathbb{R})$ , one has that  $\psi * \phi \in C^\infty(\mathbb{R}) \cap \mathcal{S}'(\mathbb{R})$  and  $\widehat{\psi * \phi} = \sqrt{2\pi} \hat{\psi} \hat{\phi}$  (see Rudin [8]). Then, for  $\varphi \in \mathcal{S}(\mathbb{R})$ ,  $g[\varphi](\cdot) = \psi * \partial_x \varphi(\cdot) = \int_0^{+\infty} \xi^{-1/3} \partial_x \varphi(\cdot - \xi) d\xi$ . Next lemma gives the Fourier transform of function  $\psi$ .

LEMMA 2.1. *For the function  $\psi$  defined by (2.1) we have*

$$\hat{\psi}(\xi) = \frac{1}{\sqrt{2\pi}} \Gamma\left(\frac{2}{3}\right) \left(\frac{1}{2} - i \frac{\sqrt{3}}{2} \operatorname{sgn}(\xi)\right) |\xi|^{-2/3}, \quad (2.3)$$

where

$$\operatorname{sgn}(\xi) = \begin{cases} -1, & \xi < 0, \\ 1, & \xi > 0, \end{cases}$$

and  $\Gamma$  is the gamma function.

*Proof.* We define the function  $\psi_n(x) := \chi_{(0,n)}(x) x^{-1/3}$ , for all  $x \in \mathbb{R}$ , and  $n \in \mathbb{N}$ . It is not difficult to see that  $\psi_n \rightarrow \psi$  in  $\mathcal{S}'(\mathbb{R})$  as  $n$  goes to infinity. Let  $\varphi \in \mathcal{S}(\mathbb{R})$ . Then

$$\begin{aligned} \langle \hat{\psi}_n, \varphi \rangle &= \frac{1}{\sqrt{2\pi}} \int \left[ \int_0^n \frac{\cos(\xi x)}{\xi^{1/3}} d\xi - i \int_0^n \frac{\sin(\xi x)}{\xi^{1/3}} d\xi \right] \varphi(x) dx \\ &= \frac{1}{\sqrt{2\pi}} \int x^{-2/3} \left[ \int_0^{nx} \frac{\cos(u)}{u^{1/3}} du - i \int_0^{nx} \frac{\sin(u)}{u^{1/3}} du \right] \varphi(x) dx. \end{aligned}$$

Since

$$\int_0^{+\infty} \frac{\cos x}{x^{1/3}} dx = \frac{1}{2} \Gamma\left(\frac{2}{3}\right), \quad \text{and} \quad \int_0^{+\infty} \frac{\sin x}{x^{1/3}} dx = \frac{\sqrt{3}}{2} \Gamma\left(\frac{2}{3}\right),$$

it follows that

$$\left| x^{-2/3} \int_0^{nx} \frac{e^{-iu}}{u^{1/3}} du \varphi(x) \right| \leq C |x|^{-2/3} |\varphi(x)|,$$

for all  $n \in \mathbb{N}$ , and  $x \in \mathbb{R}$ . Therefore, the dominated convergence theorem implies that

$$\lim_{n \rightarrow \infty} \langle \hat{\psi}_n, \varphi \rangle = \frac{1}{\sqrt{2\pi}} \int x^{-2/3} \Gamma\left(\frac{2}{3}\right) \left(\frac{1}{2} - i \frac{\sqrt{3}}{2} \operatorname{sgn}(x)\right) \varphi(x) dx.$$

This completes the proof of the lemma.  $\square$

REMARK 2.1. *Let  $s \in \mathbb{R}$ . If  $u \in H^s(\mathbb{R})$ , one can define  $g[u]$  through its Fourier transform by*

$$\widehat{g[u]}(\xi) := \Gamma\left(\frac{2}{3}\right) \left(\frac{\sqrt{3}}{2} \operatorname{sgn}(\xi) + \frac{i}{2}\right) \xi^{1/3} \hat{u}(\xi), \quad (2.4)$$

for almost every  $\xi \in \mathbb{R}$ . Thus, if  $u \in H^s(\mathbb{R})$ , it follows that  $g[u] \in H^{s-1/3}(\mathbb{R})$  and  $\|g[u]\|_{H^{s-1/3}} \leq \Gamma\left(\frac{2}{3}\right) \|u\|_{H^s}$ .

In this section we consider the following, more general, version of equation (1.1):

$$\partial_t u(x, t) + \partial_x \left( \frac{u^2}{2} - \partial_x u + \eta g[u] \right)(x, t) = 0, \quad (2.5)$$

where  $\eta \in \mathbb{R}$ . We will show existence of travelling-wave solutions to equation (2.5), for any  $\eta \in \mathbb{R}$ . First, we consider the case  $\eta = 0$ . For any  $d \in \mathbb{R}$  (see Johnson [7]),

$$u_d(x, t) = \frac{d}{2} \left[ 1 - \tanh \left( \frac{d}{4} \left( x - \frac{d}{2} t \right) \right) \right] \quad (2.6)$$

is a solution to equation (2.5) with  $\eta = 0$ .

REMARK 2.2. *Let  $\lambda > 0$ . We define*

$$u_\lambda(x, t) := \frac{1}{\lambda} u \left( \frac{x}{\lambda}, \frac{t}{\lambda^2} \right), \text{ for } x \in \mathbb{R}, \text{ and } t \geq 0. \quad (2.7)$$

*It is straightforward to check that if  $u$  is a solution to the equation*

$$\partial_t u(x, t) + \partial_x \left( \frac{u^2}{2} - \partial_x u + \lambda^{2/3} \eta g[u] \right) (x, t) = 0, \quad (2.8)$$

*then  $u_\lambda$  satisfies equation (2.5). Hence, if  $\phi$  is a travelling-wave solution of equation (2.8) with speed  $c$ , then  $\phi_\lambda(\cdot) = \frac{1}{\lambda} \phi(\frac{\cdot}{\lambda})$  is a travelling-wave solution of equation (2.5) with speed  $c/\lambda$ .*

We define, for  $c \in \mathbb{R}$ , the functions

$$g_c(x) := c \left( 1 - \tanh \left( \frac{c}{2} x \right) \right), \text{ and } h_c(x) := g'_c(x) = -\frac{c^2}{2} \operatorname{sech}^2 \left( \frac{c}{2} x \right). \quad (2.9)$$

REMARK 2.3. *Let  $c \in \mathbb{R}$ . We see that  $g[g_c] = I_1 + I_2$ , where  $I_j := \psi_j * h_c$  for  $j = 1, 2$ , with  $\psi_1 := \psi \cdot \chi_{(0,1)}$ , and  $\psi_2 := \psi \cdot \chi_{(0,+\infty)}$ . Now, we state some immediate properties of the function  $g[g_c]$ .*

**a.)** *Let  $p > 3$ . Since  $\psi_1 \in L^1(\mathbb{R})$ , and  $\psi_2 \in L^p(\mathbb{R})$ , it follows from the Young inequality for convolution that  $g[g_c] \in L^p(\mathbb{R})$ .*

**b.)** *Furthermore,  $g[g_c] \in C_\infty(\mathbb{R})$ . In fact, it follows from the dominated convergence theorem that  $I_1$  is continuous and  $I_1(x) \rightarrow 0$  as  $|x| \rightarrow \infty$ . Moreover, Hölder's inequality and the dominated convergence theorem imply that*

$$I_2(x) \leq C(c) \left( \int_1^{+\infty} \frac{\operatorname{sech}^4 \left( \frac{c}{2} (x - \xi) \right)}{\xi^{4/3}} d\xi \right)^{1/4} \rightarrow 0, \text{ as } |x| \rightarrow +\infty.$$

*The continuity of  $I_2$  is shown similarly to the continuity of  $I_1$ .*

Let  $c \in \mathbb{R}$ . In the sequel, we will consider the following spaces:

$$\begin{aligned} \mathfrak{X} = \mathfrak{X}_c &:= \left\{ \varphi \in C_b^1(\mathbb{R}) ; \int \varphi' h'_c dx = 0 \right\}, \\ \tilde{\mathfrak{X}} = \tilde{\mathfrak{X}}_c &:= \left\{ g_c + \varphi ; \varphi \in \mathfrak{X} \right\}, \end{aligned}$$

where  $\| \cdot \|_{\mathfrak{X}} = \| \cdot \|_{C_b^1}$ . One sees that  $(\mathfrak{X}, \| \cdot \|_{\mathfrak{X}})$  is a Banach space.

REMARK 2.4. *Assume that  $\varphi \in C_b^1(\mathbb{R})$ . By integration by parts one has that*

$$g[\varphi](x) = \int_0^1 \frac{1}{\xi^{1/3}} \varphi'(x - \xi) d\xi + \varphi(x - 1) - \frac{1}{3} \int_1^{+\infty} \frac{1}{\xi^{4/3}} \varphi(x - \xi) d\xi. \quad (2.10)$$

*Then  $g[\varphi] \in C_b(\mathbb{R})$ . Moreover,*

$$\|g[\varphi]\|_{L^\infty} \leq C \| \varphi \|_{C_b^1}. \quad (2.11)$$

*Hence, if  $\phi \in \tilde{\mathfrak{X}}$ , it follows from Remark 2.3-b.) above that  $g[\phi] \in C_b(\mathbb{R})$ .*

Suppose now that  $u(x, t) = \phi(x - ct)$  is a solution to equation (2.5), where  $\phi \in \tilde{\mathfrak{X}}$ . Then  $-c\phi' + \frac{d}{dx}(\frac{\phi^2}{2} - \phi' + \eta g[\phi]) = 0$ . Thus, a sufficient condition to guarantee that  $\phi$  satisfies the last equation is

$$F(\eta, \phi) = F_c(\eta, \phi) := c\phi - \frac{\phi^2}{2} + \phi' - \eta g[\phi] = 0. \quad (2.12)$$

We denote by  $\tau_c$  the function given by  $\tau_c(x) := c - g_c(x) = c \tanh(\frac{c}{2}x)$ , for  $x \in \mathbb{R}$ . We now define the function  $G = G_c$ , which is well defined on  $\mathbb{R} \times \mathfrak{X}$  by Remarks 2.3 and 2.4 above, as

$$\begin{aligned} G : \mathbb{R} \times \mathfrak{X} &\mapsto C_b(\mathbb{R}) \\ (\eta, \varphi) &\mapsto G(\eta, \varphi) = \tau_c \varphi - \frac{\varphi^2}{2} + \varphi' - \eta g[\varphi] - \eta g[g_c]. \end{aligned} \quad (2.13)$$

Assume that  $\phi = \varphi + g_c \in \tilde{\mathfrak{X}}$ . Since  $F(0, g_c) = 0$ , it follows that  $F(\eta, \phi) = G(\eta, \varphi)$ . Hence,  $\phi$  satisfies equation (2.12) if and only if  $\varphi$  verifies the equation  $G(\eta, \varphi) = 0$ .

The following theorem implies the existence of a travelling-wave solution,  $u(x, t) = \phi(x - ct)$  with  $c > 0$  and  $\phi \in \tilde{\mathfrak{X}}$ , to equation (2.5) for  $\eta$  in a neighborhood of zero; its proof uses the implicit function theorem.

**THEOREM 2.1.** *Suppose  $c > 0$ . Then there exist  $\delta, \delta_0 > 0$  such that for every  $\eta \in (-\delta, \delta)$ , there is exactly one  $\varphi_\eta = \varphi_{\eta, c} \in \mathfrak{X}$  for which  $\|\varphi_\eta\|_{\mathfrak{X}} \leq \delta_0$  and  $G(\eta, \varphi_\eta) = 0$ . Moreover, the mapping  $\eta \mapsto \varphi_\eta$  is a  $C^\infty$ -map on a neighborhood of 0.*

*Proof.* Let  $c > 0$ . The mapping  $G = G_c$  is defined on the Banach space  $\mathbb{R} \times \mathfrak{X}$  taking values in the Banach space  $(C_b(\mathbb{R}), \|\cdot\|_{L^\infty})$ , and satisfies  $G(0, 0) = 0$ .

We now claim that  $\partial_1 G$  and  $\partial_2 G$  exist as partial F-derivatives (Fréchet derivative) on  $\mathbb{R} \times \mathfrak{X}$  and that the partial F-derivative  $\partial_2 G(0, 0) : \mathfrak{X} \mapsto C_b(\mathbb{R})$  is bijective. In fact, let us take  $(\eta, \varphi) \in \mathbb{R} \times \mathfrak{X}$ . One can see that

$$\partial_1 G(\eta, \varphi) \cdot = -(g[\varphi] + g[g_c]) \cdot$$

and

$$\partial_2 G(\eta, \varphi) \cdot = (\tau_c - \varphi) \cdot + \partial_x \cdot - \eta g[\cdot]. \quad (2.14)$$

Then  $\partial_1 G(\eta, \varphi) \in \mathfrak{L}(\mathbb{R}, C_b(\mathbb{R}))$ , and  $\partial_2 G(\eta, \varphi) \in \mathfrak{L}(\mathfrak{X}, C_b(\mathbb{R}))$ . Moreover, we obtain that  $\|\partial_1 G(\eta, \varphi)\|_{\mathfrak{L}(\mathbb{R}, C_b(\mathbb{R}))} \leq C \cdot (\|\varphi\|_{C_b^1} + \|g[g_c]\|_{L^\infty})$ , and  $\|\partial_2 G(\eta, \varphi)\|_{\mathfrak{L}(\mathfrak{X}, C_b(\mathbb{R}))} \leq C \cdot (1 + |\eta| + \|\tau_c - \varphi\|_{L^\infty})$ , where we have used inequality (2.11). Hence,  $\partial_1 G, \partial_2 G$  exist as partial F-derivatives on  $\mathbb{R} \times \mathfrak{X}$ .

We will now show that the partial F-derivative  $\partial_2 G(0, 0) = \tau_c + \partial_x : \mathfrak{X} \mapsto C_b(\mathbb{R})$  is bijective. We begin with the injectivity; we emphasize here that the definition of the space  $\mathfrak{X} \subset C_b^1(\mathbb{R})$  was chosen to ensure the injectivity of the mapping  $\partial_2 G(0, 0)$ . Let  $f$  be an element of  $\mathfrak{X}$  such that  $\tau_c f + f' = 0$ . By solving the last ordinary differential equation, one gets

$$f(x) = f(0) \cdot e^{-\int_0^x \tau_c(s) ds} = f(0) \cdot \operatorname{sech}^2\left(\frac{c}{2}x\right).$$

Since  $f \in \mathfrak{X}$ , it follows that

$$\int f'(x) h'_c(x) dx = -f(0) \frac{2}{c^2} \int (h'_c)^2(x) dx = 0.$$

Then  $f(0) = 0$ , and therefore  $f = 0$ .

We will now show that the mapping  $\partial_2 G(0, 0)$  is onto. Let  $y$  be an element of

$C_b(\mathbb{R})$ . By the method of variation of parameters, we obtain that the function

$$g(x) := \lambda l_c(x) + l_c(x) \int_0^x \frac{y(s)}{l_c(s)} ds \quad (2.15)$$

is a solution to the equation  $\tau_c g + g' = y$ , for any  $\lambda \in \mathbb{R}$ , where  $l_c := -\frac{2}{c^2} h_c = \operatorname{sech}^2(\frac{c}{2}x)$ . We will prove that  $g \in \mathfrak{X}$  for a suitably chosen real number  $\lambda$ . First, we remark that there exists a unique  $\lambda = \lambda_{y,c} \in \mathbb{R}$  such that  $\int g' h'_c dx = 0$ . In fact take

$$\lambda := \frac{c^2}{2 \int (h'_c)^2(x) dx} \int \left[ (h'_c)^2(x) \int_0^x \frac{y(s)}{h_c(s)} ds + y(x) h'_c(x) \right] dx, \quad (2.16)$$

where we note that

$$0 < \int (h'_c)^2(x) dx \leq \frac{c^6}{4} \int \operatorname{sech}^4\left(\frac{c}{2}x\right) dx = C(c), \quad \int |h'_c(x)| dx = c^2,$$

and

$$\begin{aligned} \int (h'_c)^2(x) \left| \int_0^x \frac{y(s)}{h_c(s)} ds \right| dx &\leq \frac{c^4}{2} \|y\|_{L^\infty} \int \frac{\sinh^2(\frac{c}{2}x)}{\cosh^6(\frac{c}{2}x)} \left| \int_0^x \frac{1 + \cosh(cs)}{2} ds \right| dx \\ &\leq \frac{c^3}{4} \|y\|_{L^\infty} \int \left[ c|x| \operatorname{sech}^4\left(\frac{c}{2}x\right) + 2 \operatorname{sech}^2\left(\frac{c}{2}x\right) \right] dx \leq C(c) \|y\|_{L^\infty}. \end{aligned}$$

It remains to show that  $g$  given by (2.15) and (2.16) belongs to  $C_b^1(\mathbb{R})$ . It is immediate to see that  $g \in C(\mathbb{R})$ , we need to show that  $g$  is bounded. We have that

$$\begin{aligned} \operatorname{sech}^2\left(\frac{c}{2}x\right) \left| \int_0^x \frac{y(s)}{\operatorname{sech}^2(\frac{c}{2}s)} ds \right| &\leq \frac{\|y\|_{L^\infty}}{1 + \cosh(cx)} \left| \int_0^x (1 + \cosh(cs)) ds \right| \\ &\leq \|y\|_{L^\infty} \left( \frac{|x|}{1 + \cosh(cx)} + \frac{1}{c} |\tanh(cx)| \right) \leq C(c) \|y\|_{L^\infty}. \end{aligned}$$

Then  $g \in C_b(\mathbb{R})$ . Moreover, since  $g$  satisfies the equation  $\tau_c g + g' = y$ , it follows that  $g \in C_b^1(\mathbb{R})$ . Hence,  $g \in \mathfrak{X}$ . Therefore,  $\partial_2 G(0, 0)$  is a surjective mapping.

It is not difficult to see, by using inequality (2.11), that  $G$ ,  $\partial_1 G$  and  $\partial_2 G$  are continuous on  $\mathbb{R} \times \mathfrak{X}$ . Then, the implicit function theorem implies the first part of the theorem. Furthermore, from (2.13) one can see that function  $G$  is quadratic in  $\varphi$  and linear in  $\eta$ , therefore it is not difficult to verify that  $\partial_{i,j}^2 G(\eta, \varphi)$  is independent of  $(\eta, \varphi) \in \mathbb{R} \times \mathfrak{X}$ , for all  $i, j \in \{1, 2\}$ . Hence,  $\partial_{i_1, \dots, i_k}^k G(\eta, \varphi) = 0$  for all  $k \geq 3$ , where  $i_1, \dots, i_k \in \{1, 2\}$ , and  $(\eta, \varphi) \in \mathbb{R} \times \mathfrak{X}$ . Finally, the second part of the theorem is then a consequence of the fact that the mapping  $G$  is a  $C^\infty$ -map on  $\mathbb{R} \times \mathfrak{X}$ .  $\square$

**COROLLARY 2.1.** *Let  $\eta \in \mathbb{R}$  and  $c > 0$ . Then there exists  $\lambda_0 = \lambda_0(\eta, c) > 0$  such that for every  $d \in (\frac{c}{\lambda_0}, +\infty)$ , there is a travelling-wave solution  $\tilde{\phi} \in C_b^1(\mathbb{R})$  of equation (2.5) with speed  $d$ .*

*Proof.* By Theorem 2.1 there exists  $\lambda_0 > 0$  such that for every  $\lambda \in (0, \lambda_0)$ , there is a  $\phi = \phi_{\lambda, \eta, c} \in C_b^1(\mathbb{R})$  such that  $u(x, t) = \phi(x - ct)$  is a solution to equation (2.8). Now we can see, by using Remark 2.2, that  $\tilde{\phi}(\cdot) = \frac{1}{\lambda} \phi(\frac{\cdot}{\lambda})$  is a travelling-wave solution of equation (2.5) with speed  $c/\lambda$ .  $\square$

### 3. LOCAL THEORY IN A SUBSPACE OF $C_b^1(\mathbb{R})$

In Section 2, we proved the existence of a travelling-wave solution  $u(x, t) = \phi(x - ct)$  to equation (2.5) for any  $\eta \in \mathbb{R}$ , where  $c$  is an appropriate positive number and  $\phi \in C_b^1(\mathbb{R})$ . Motivated by this last result, we will consider in this section the local well-posedness theory for the following initial value problem (IVP)

$$\begin{cases} \partial_t u(x, t) + \partial_x \left( \frac{1}{2} u^2 - \partial_x u + g[u] \right)(x, t) = 0, \\ u(0) = u_0, \end{cases} \quad (3.1)$$

where  $g[u]$  is given by (2.2), and  $u_0$  belongs to a suitable subspace of  $C_b^1(\mathbb{R})$ . The Cauchy problem associated to the IVP (3.1) for initial data  $u_0 \in L^2(\mathbb{R})$  was recently studied by Alibaud, Azerad and Isèbe [1].

**3.1. The Linear Equation.** First, we consider the linear part associated to the IVP (3.1), namely

$$\begin{cases} \partial_t u(x, t) - \partial_x^2 u(x, t) + \partial_x g[u](x, t) = 0, \\ u(0) = u_0. \end{cases} \quad (3.2)$$

By formally taking the Fourier transform of the last expression, we get

$$\hat{u}(\xi, t) = \hat{K}(\xi, t) \hat{u}_0(\xi), \quad (3.3)$$

where

$$\hat{K}(\xi, t) = e^{-t[\xi^2 - \xi^{4/3}(a + ib \operatorname{sgn}(\xi))]}, \quad (3.4)$$

for  $\xi \in \mathbb{R}$  and  $t \geq 0$ , with  $a := \frac{1}{2}\Gamma(\frac{2}{3})$  and  $b := -\frac{\sqrt{3}}{2}\Gamma(\frac{2}{3})$ . For  $\xi \in \mathbb{R}$ , we define

$$\Phi(\xi) := (a + ib \operatorname{sgn}(\xi)). \quad (3.5)$$

We note that  $|\Phi(\xi)| = \Gamma(\frac{2}{3})$ , for all  $\xi \in \mathbb{R}$ .

**REMARK 3.1.** *For every  $t > 0$ , the kernel  $K(\cdot, t)$  is not a nonnegative function. More precisely, for every  $t > 0$ , there exists  $x \in \mathbb{R}$  such that  $K(x, t) < 0$ . This fact implies, in particular, that the IVP (3.1) is non-monotone (see [1] for more details).*

For  $t \geq 0$ , we define the operator  $E(t)$  by

$$\begin{cases} E(t)\phi(x) = \frac{1}{\sqrt{2\pi}}(K(\cdot, t) * \phi)(x), & \text{for } t > 0 \text{ and } x \in \mathbb{R}, \\ E(0)\phi = \phi, \end{cases} \quad (3.6)$$

where  $\phi \in C_b(\mathbb{R})$  (see Lemma 3.11 below). Now, we define the following spaces

$$Y := \{g \in C_b(\mathbb{R}); g \text{ is uniformly continuous}\}; \quad (3.7)$$

$$X := \{f \in C_b^1(\mathbb{R}); f' \text{ is uniformly continuous}\}. \quad (3.8)$$

One can see that  $(Y, \|\cdot\|_{C_b(\mathbb{R})})$ , and  $(X, \|\cdot\|_{C_b^1(\mathbb{R})})$  are Banach spaces and that  $X \hookrightarrow Y$ . In Sub-section 3.2 we will show local-in-time well-posedness of the IVP (3.1), with initial data  $u_0 \in X$ .

The following lemma contains a calculus result.

**LEMMA 3.1.** *Let  $h : \mathbb{R} \mapsto \mathbb{C}$  be a function which satisfies the following conditions:*

- i.)**  $h \in L^1(\mathbb{R}) \cap C_\infty(\mathbb{R}) \cap C^2(\mathbb{R} \setminus \{0\})$ ;
- ii.)**  $h' \in L^1(\mathbb{R})$ ,  $|h'(x)| \rightarrow 0$  as  $|x| \rightarrow +\infty$ . Moreover, there exist  $\lim_{x \downarrow 0} h'(x) = h'(0^+)$ , and  $\lim_{x \uparrow 0} h'(x) = h'(0^-)$ ;

iii.)  $h'' \in L^1(\mathbb{R})$ .

Then  $\hat{h} \in L^1(\mathbb{R}) \cap C_\infty(\mathbb{R})$ , and

$$\|\hat{h}\|_{L^1} \leq \sqrt{\frac{2}{\pi}} \left[ \|h\|_{L^1} + |h'(0^+) - h'(0^-)| + \|h''\|_{L^1} \right]. \quad (3.9)$$

*Proof.* Since  $h \in L^1(\mathbb{R})$ , it follows from the Riemann-Lebesgue lemma that  $\hat{h} \in C_\infty(\mathbb{R})$ . After using integration by parts twice, we see that

$$\hat{h}(\xi) = \frac{1}{\sqrt{2\pi}} \left[ \int_{-\infty}^{+\infty} h''(x) \frac{e^{-i\xi x}}{(i\xi)^2} dx + \frac{h'(0^+) - h'(0^-)}{(i\xi^2)} \right], \quad \text{for } \xi \neq 0.$$

Expression (3.9) follows from the last equation and from the fact that  $\|\hat{h}\|_{L^\infty} \leq \frac{1}{\sqrt{2\pi}} \|h\|_{L^1}$ .  $\square$

REMARK 3.2. *It is well-known that  $W^{1,1}(\mathbb{R}) \subset C_\infty(\mathbb{R}) \cap AC(\mathbb{R})$ , where  $AC(\mathbb{R})$  denotes the space of all complex-valued functions, which are absolutely continuous on  $\mathbb{R}$ . Therefore, it follows from Lemma 3.1 above that if  $f \in W^{2,1}(\mathbb{R})$ , then  $\hat{f} \in L^1(\mathbb{R}) \cap C_\infty(\mathbb{R})$  and*

$$\|\hat{f}\|_{L^1} \leq \sqrt{\frac{2}{\pi}} \left[ \|f\|_{L^1} + \|f''\|_{L^1} \right]. \quad (3.10)$$

REMARK 3.3. *Suppose now that  $t \in (0, 1)$ . Since*

$$\begin{aligned} K(x, t) &= \frac{1}{\sqrt{2\pi}} \int e^{ix\xi} e^{-t[\xi^2 - \xi^{4/3}\Phi(\xi)]} d\xi \\ &= \frac{t^{-1/2}}{\sqrt{2\pi}} \int e^{i(t^{-1/2}x)\xi} e^{-[\xi^2 - \xi^{4/3}\Phi(\xi)]} e^{-(1-t^{1/3})\xi^{4/3}\Phi(\xi)} d\xi, \end{aligned}$$

*it follows that*

$$K(x, t) = t^{-1/2} (K(\cdot, 1) * G(\cdot, 1 - t^{1/3}))(t^{-1/2}x), \quad \text{for } x \in \mathbb{R}, \quad (3.11)$$

*where*

$$G(\cdot, 1 - t^{1/3}) = \frac{1}{\sqrt{2\pi}} \mathcal{F}^{-1}(e^{-(1-t^{1/3})\xi^{4/3}(a+ib \operatorname{sgn}(\xi))})(\cdot). \quad (3.12)$$

The next three lemmas are elementary calculus results which will be used in the sequel.

LEMMA 3.2. *Suppose that  $\alpha > -1$ , and  $\beta > 0$ . Then*

$$I(\alpha, \beta) := \int |\xi|^\alpha e^{-\beta|\xi|^{4/3}} d\xi = C(\alpha) \beta^{-\frac{3}{4}(\alpha+1)}.$$

*Proof.* The assertion of the lemma follows from the fact that

$$I(\alpha, \beta) = \beta^{-\frac{3}{4}(\alpha+1)} \int |\tau|^\alpha e^{-|\tau|^{4/3}} d\tau.$$

$\square$

LEMMA 3.3. *Suppose that  $\alpha > -1$ ,  $\beta > 0$ , and  $t > 0$ . Then*

$$I(\alpha, \beta, t) := \int |\xi|^\alpha e^{-t[\xi^2 - \beta|\xi|^{4/3}]} d\xi \leq C(\alpha, \beta) \left[ e^{\frac{4}{27}\beta^3 t} + t^{-\frac{\alpha+1}{2}} \right].$$

*Proof.* It is elementary to check that  $\xi^2 - \beta\xi^{4/3} \geq -\frac{4}{27}\beta^3$  for all  $\xi \in \mathbb{R}$ , and  $\xi^2 - \beta\xi^{4/3} \geq \xi^2/2$  for  $\xi \geq (2\beta)^{3/2}$ . Then

$$\begin{aligned} I(\alpha, \beta, t) &\leq 2 \left[ \int_0^{(2\beta)^{3/2}} \xi^\alpha e^{\frac{4}{27}\beta^3 t} d\xi + \int_{(2\beta)^{3/2}}^{+\infty} \xi^\alpha e^{-\frac{1}{2}\xi^2} d\xi \right] \\ &\leq C(\alpha, \beta) \left[ e^{\frac{4}{27}\beta^3 t} + \int_0^{+\infty} \left(\frac{2}{t}\right)^{\frac{\alpha}{2}} u^\alpha e^{-u^2} \sqrt{\frac{2}{t}} du \right], \end{aligned}$$

where in the last inequality we have used the fact that  $\alpha > -1$ . The result now follows.  $\square$

LEMMA 3.4. *Suppose that  $g \in W^{1,1}(\mathbb{R})$ , and  $l \in L^\infty(\mathbb{R})$ . If  $f := g * l$ , then  $f \in C^1(\mathbb{R}) \cap W^{1,\infty}(\mathbb{R})$ , and  $f'(x) = (g' * l)(x)$  for all  $x \in \mathbb{R}$ .*

*Proof.* Since  $g \in L^1(\mathbb{R})$  and  $l \in L^\infty(\mathbb{R})$ , it follows from Young's inequality that  $f \in L^\infty(\mathbb{R})$ . Moreover, since  $|f(x+h) - f(x)| \leq \|g(\cdot+h) - g(\cdot)\|_{L^1} \|l\|_{L^\infty}$  for all  $x \in \mathbb{R}$ , and  $\|g(\cdot+h) - g(\cdot)\|_{L^1} \rightarrow 0$  as  $h$  tends to zero, it follows that  $f \in C(\mathbb{R})$ . Let  $x \in \mathbb{R}$ . Since  $W^{1,1}(\mathbb{R}) \subset AC(\mathbb{R})$ , we see that

$$\begin{aligned} &\left| \frac{f(x+h) - f(x)}{h} - \int g'(x-y)l(y)dy \right| = \\ &\left| \int_0^1 (g'(x-y+th) - g'(x-y))l(y)tdtdy \right| \\ &\leq \|l\|_{L^\infty} \int_0^1 \|g'(\cdot+th) - g'(\cdot)\|_{L^1} dt \rightarrow 0 \quad \text{as } h \rightarrow 0, \end{aligned}$$

where the last expression is a consequence of the dominated convergence theorem.  $\square$

The following five lemmas provide more explicit estimates than the corresponding results mentioned in [1]. The next lemma gives an upper bound, which goes to infinity as  $t$  tends to 1, for  $\|G(\cdot, 1 - t^{1/3})\|_{L^1}$  when  $t \in [0, 1)$ .

LEMMA 3.5. *Let  $t_0 \in (0, 1)$ . Then, for all  $t \in [0, t_0]$ , the function  $G(\cdot, 1 - t^{1/3})$  given by (3.12) belongs to  $L^1(\mathbb{R}) \cap C_\infty(\mathbb{R})$ . Moreover,*

$$\|G(\cdot, 1 - t^{1/3})\|_{L^1} \leq C[(1 - t^{1/3})^{3/4} + (1 - t^{1/3})^{-3/4}] \leq C \cdot (1 - t_0^{1/3})^{-3/4}, \quad (3.13)$$

for all  $t \in [0, t_0]$ , where  $C$  is a positive constant independent of  $t$ .

*Proof.* Let  $t \in [0, 1)$ . We define  $g(\xi, t) := e^{-(1-t^{1/3})\xi^{4/3}\Phi(\xi)}$  for  $\xi \in \mathbb{R}$ . It follows that  $g(\cdot, t)$  is continuous. Furthermore,

$$\partial_\xi g(\xi, t) = -\frac{4}{3}(1 - t^{1/3})\xi^{1/3}\Phi(\xi)e^{-(1-t^{1/3})\xi^{4/3}\Phi(\xi)}, \quad \text{for } \xi \neq 0. \quad (3.14)$$

Then  $|\partial_\xi g(\xi, t)| \rightarrow 0$  as  $|\xi| \rightarrow +\infty$ , and  $\partial_\xi g(0^+, t) = 0 = \partial_\xi g(0^-, t)$ . Moreover,

$$\begin{aligned} \partial_\xi^2 g(\xi, t) &= \left[ -\frac{4}{9}(1 - t^{1/3})\xi^{-2/3}\Phi(\xi) + \left(\frac{4}{3}(1 - t^{1/3})\xi^{1/3}\Phi(\xi)\right)^2 \right] \\ &\quad \times e^{-(1-t^{1/3})\xi^{4/3}\Phi(\xi)}, \quad \text{for } \xi \neq 0. \end{aligned}$$

We see that  $g(\cdot, t) \in C_\infty(\mathbb{R}) \cap C^2(\mathbb{R} \setminus \{0\})$ . In addition,  $\|g(\cdot, t)\|_{L^1} = C \cdot (1 - t^{1/3})^{-3/4}$ , and  $\|\partial_\xi g(\cdot, t)\|_{L^1} = 4$ . Furthermore,

$$\begin{aligned} \|\partial_\xi^2 g(\cdot, t)\|_{L^1} &\leq C \cdot (1 - t^{1/3}) \int |\xi|^{-2/3} e^{-(1-t^{1/3})a\xi^{4/3}} d\xi \\ &\quad + C \cdot (1 - t^{1/3})^2 \int |\xi|^{2/3} e^{-(1-t^{1/3})a\xi^{4/3}} d\xi \\ &\leq C \cdot (1 - t^{1/3})^{3/4}, \end{aligned}$$

where the last inequality is a consequence of Lemma 3.2 above. The result now follows from Lemma 3.1.  $\square$

Lemmas 3.6 and 3.9 below provide estimates for  $\|K(\cdot, t)\|_{L^1}$  and  $\|\partial_x K(\cdot, t)\|_{L^1}$ , for any  $t > 0$ .

LEMMA 3.6. *Suppose that  $t > 0$ . Then the function  $K(\cdot, t) \in L^1(\mathbb{R}) \cap C_\infty(\mathbb{R})$ , and*

$$\|K(\cdot, t)\|_{L^1} \leq C \cdot (1 + t^2 e^{\frac{4}{27}a^3 t}), \quad (3.15)$$

where  $C$  is a positive constant independent of  $t$ .

*Proof.* Let  $t > 0$ . It follows from (3.4) that

$$\partial_\xi \hat{K}(\xi, t) = -t \left[ 2\xi - \frac{4}{3} \xi^{1/3} \Phi(\xi) \right] \hat{K}(\xi, t), \quad \text{for } \xi \neq 0,$$

and

$$\partial_\xi^2 \hat{K}(\xi, t) = \left\{ -t \left[ 2 - \frac{4}{9} \xi^{-2/3} \Phi(\xi) \right] + t^2 \left[ 2\xi - \frac{4}{3} \xi^{1/3} \Phi(\xi) \right]^2 \right\} \hat{K}(\xi, t), \quad \text{for } \xi \neq 0.$$

Then  $\hat{K}(\cdot, t) \in C_\infty(\mathbb{R}) \cap C^2(\mathbb{R} \setminus \{0\})$ . Moreover,  $|\partial_\xi \hat{K}(\xi, t)| \rightarrow 0$  as  $|\xi| \rightarrow +\infty$ , and  $\partial_\xi \hat{K}(0^+, t) = 0 = \partial_\xi \hat{K}(0^-, t)$ . Furthermore,

$$\|\hat{K}(\cdot, t)\|_{L^1} = 2 \int_0^{+\infty} e^{-t[\xi^2 - a\xi^{4/3}]} d\xi \leq C \left[ e^{\frac{4}{27}a^3 t} + \frac{1}{\sqrt{t}} \right],$$

where the last inequality is a consequence of Lemma 3.3. Again using Lemma 3.3, we see that

$$\begin{aligned} \|\partial_\xi \hat{K}(\cdot, t)\|_{L^1} &\leq C [1 + t^{1/3} + t e^{\frac{4}{27}a^3 t}], \quad \text{and} \\ \|\partial_\xi^2 \hat{K}(\cdot, t)\|_{L^1} &\leq C [\sqrt{t} + t^{5/6} + t^{7/6} + (t + t^2) e^{\frac{4}{27}a^3 t}] \leq C [\sqrt{t} + t^2 e^{\frac{4}{27}a^3 t}]. \end{aligned}$$

Lemma 3.1, applied to  $h(\cdot) = \hat{K}(\cdot, t)$ , implies that

$$\|K(\cdot, t)\|_{L^1} \leq C \left[ \frac{1}{\sqrt{t}} + t^2 e^{\frac{4}{27}a^3 t} \right], \quad \text{for all } t > 0. \quad (3.16)$$

Suppose now that  $t \in (0, 1)$ . Then

$$\begin{aligned} \int |K(x, t)| dx &= \frac{1}{\sqrt{t}} \int |K(\cdot, 1) * G(\cdot, 1 - t^{1/3})|(x/\sqrt{t}) dx \\ &= \int (1 + y^2)^{1/2} \frac{|K(\cdot, 1) * G(\cdot, 1 - t^{1/3})|(y)}{(1 + y^2)^{1/2}} dy \\ &\leq C \|\hat{K}(\cdot, 1) \hat{G}(\cdot, 1 - t^{1/3})\|_{H^1}, \end{aligned} \quad (3.17)$$

where the first equality above comes from (3.11). From the fact that  $e^{-(1-h^{1/3})a\xi^{4/3}} \leq 1$ , for all  $h \in [0, 1)$ ,  $\xi \in \mathbb{R}$ , and equation (3.14) we have that  $|\hat{G}(\xi, 1 - h^{1/3})| \leq C$ , and  $|\partial_\xi \hat{G}(\xi, 1 - h^{1/3})| \leq C|\xi|^{1/3}$  for all  $\xi \in \mathbb{R}$ . Now, from Lemma 3.3, we obtain

$$\begin{aligned} \|\hat{K}(\cdot, 1)\hat{G}(\cdot, 1 - h^{1/3})\|_{H^1} &\leq \|\hat{K}(\cdot, 1)\hat{G}(\cdot, 1 - h^{1/3})\|_{L^2} \\ &+ \|\partial_\xi \hat{K}(\cdot, 1)\hat{G}(\cdot, 1 - h^{1/3})\|_{L^2} + \|\hat{K}(\cdot, 1)\partial_\xi \hat{G}(\cdot, 1 - h^{1/3})\|_{L^2} \leq C, \end{aligned} \quad (3.18)$$

for all  $h \in [0, 1)$ . The assertion of the lemma now follows from (3.16)-(3.18).  $\square$

The following result gives an upper bound for  $\|\partial_x K(\cdot, t)\|_{L^1}$  when  $t \in (0, 1)$ .

LEMMA 3.7. *Let  $t \in (0, 1)$ . Then,  $K(\cdot, t) \in L^1(\mathbb{R}) \cap C^1(\mathbb{R}) \cap W^{1,\infty}(\mathbb{R})$ . In addition,  $\partial_x K(\cdot, t)(x) = t^{-1}(\partial_x K(\cdot, 1) * G(\cdot, 1 - t^{1/3}))(t^{-1/2}x)$  for all  $x \in \mathbb{R}$ ,  $\partial_x K(\cdot, t) \in L^1(\mathbb{R}) \cap C_\infty(\mathbb{R})$ , and*

$$\|\partial_x K(\cdot, t)\|_{L^1} \leq \frac{C}{\sqrt{t}} [(1 - t^{1/3})^{3/4} + (1 - t^{1/3})^{-3/4}], \quad (3.19)$$

where  $C$  is a positive constant independent of  $t$ .

*Proof.* Let  $f$  denote the function given by  $f(\xi) := \xi \hat{K}(\xi, 1)$  for all  $\xi \in \mathbb{R}$ . Then  $f \in C_\infty(\mathbb{R}) \cap C^2(\mathbb{R})$ . Furthermore,

$$f'(\xi) = \left[1 - \xi \left(2\xi - \frac{4}{3}\xi^{1/3}\Phi(\xi)\right)\right] \hat{K}(\xi, 1), \quad \text{for } \xi \neq 0,$$

and

$$f''(\xi) = \left\{ \left[-4\xi + \frac{16}{9}\xi^{1/3}\Phi(\xi)\right] - \left[1 - \xi \left(2\xi - \frac{4}{3}\xi^{1/3}\Phi(\xi)\right)\right] \left[2\xi - \frac{4}{3}\xi^{1/3}\Phi(\xi)\right] \right\} \hat{K}(\xi, 1),$$

for  $\xi \neq 0$ . By using Lemma 3.3, it follows that  $\|f\|_{L^1} \leq C$ ,  $\|f'\|_{L^1} \leq C$ , and  $\|f''\|_{L^1} \leq C$ . Moreover,  $|f'(\xi)| \rightarrow 0$  as  $|\xi| \rightarrow +\infty$ , and  $f'(0^+) = 1 = f'(0^-)$ . Thus, Lemma 3.1 implies that  $\partial_x K(\cdot, 1) \in C_\infty(\mathbb{R}) \cap L^1(\mathbb{R})$ . Therefore, using Lemma 3.6, we have that  $K(\cdot, 1) \in W^{1,1}(\mathbb{R})$ . Let  $t \in (0, 1)$ . Lemma 3.2 implies that

$$\|G(\cdot, 1 - t^{1/3})\|_{L^\infty} \leq C \cdot (1 - t^{1/3})^{-3/4}. \quad (3.20)$$

Thus, applying Lemma 3.4 to equation (3.11), taking into account (3.20), one sees that  $K(\cdot, t) \in C^1(\mathbb{R}) \cap W^{1,\infty}(\mathbb{R})$ , and  $\partial_x K(\cdot, t)(x) = t^{-1}(\partial_x K(\cdot, 1) * G(\cdot, 1 - t^{1/3}))(t^{-1/2}x)$  for all  $x \in \mathbb{R}$ . Furthermore,

$$\begin{aligned} \|\partial_x K(\cdot, t)\|_{L^1} &= t^{-1/2} \int |\partial_x K(\cdot, 1) * G(\cdot, 1 - t^{1/3})|(y) dy \\ &\leq C t^{-1/2} [(1 - t^{1/3})^{3/4} + (1 - t^{1/3})^{-3/4}], \end{aligned}$$

where in the last step we have used Young's inequality and Lemma 3.5.  $\square$

Next lemma will be useful to study  $\|\partial_x K(\cdot, t)\|_{L^1}$  for  $t \geq t_0$ , where  $t_0 > 0$ .

LEMMA 3.8. *Suppose that  $t > 0$ . Then,  $\partial_x K(\cdot, t) \in L^1(\mathbb{R}) \cap C_\infty(\mathbb{R})$ , and*

$$\|\partial_x K(\cdot, t)\|_{L^1} \leq C \left[ \frac{1}{t} + t^2 e^{\frac{4}{27}a^3 t} \right], \quad (3.21)$$

where  $C$  is a positive constant independent of  $t$ .

*Proof.* Let  $t > 0$ . For  $\xi \in \mathbb{R}$ , we define  $h(\xi, t) := \xi \hat{K}(\xi, t)$ . Then

$$\partial_\xi h(\xi, t) = \left[ 1 - t \left( 2\xi^2 - \frac{4}{3} \xi^{\frac{4}{3}} \Phi(\xi) \right) \right] \hat{K}(\xi, t), \quad \text{for } \xi \neq 0,$$

and

$$\partial_\xi^2 h(\xi, t) = -t \left\{ \left[ 4\xi - \frac{16}{9} \xi^{\frac{1}{3}} \Phi(\xi) \right] + \left[ 1 - t \left( 2\xi^2 - \frac{4}{3} \xi^{\frac{4}{3}} \Phi(\xi) \right) \right] \left[ 2\xi - \frac{4}{3} \xi^{\frac{1}{3}} \Phi(\xi) \right] \right\} \hat{K}(\xi, t),$$

for  $\xi \neq 0$ . We see that  $h(\cdot, t) \in C_\infty(\mathbb{R}) \cap C^2(\mathbb{R})$ ,  $|\partial_\xi h(\xi, t)| \rightarrow 0$  as  $|\xi| \rightarrow +\infty$ , and  $\partial_\xi h(0^+, t) = 1 = \partial_\xi h(0^-, t)$ . Moreover, by Lemma 3.3, we have that

$$\begin{aligned} \|h(\cdot, t)\|_{L^1} &\leq C \left[ \frac{1}{t} + e^{\frac{4}{27} a^3 t} \right], \\ \|\partial_\xi h(\cdot, t)\|_{L^1} &\leq C \left[ \frac{1}{t^{1/2}} + \frac{1}{t^{1/6}} + (1+t) e^{\frac{4}{27} a^3 t} \right], \quad \text{and} \\ \|\partial_\xi^2 h(\cdot, t)\|_{L^1} &\leq C \left[ 1 + t^{1/3} + t^{2/3} + (t+t^2) e^{\frac{4}{27} a^3 t} \right]. \end{aligned}$$

The proof of the lemma is now completed by applying Lemma 3.1.  $\square$

The next result provides a unified upper bound for  $\|\partial_x K(\cdot, t)\|_{L^1}$  for any  $t > 0$ , which takes the best of the corresponding bounds obtained in Lemmas 3.7 and 3.8.

**LEMMA 3.9.** *Suppose that  $t > 0$ . Then the function  $\partial_x K(\cdot, t) \in L^1(\mathbb{R}) \cap C_\infty(\mathbb{R})$ . Moreover,*

$$\|\partial_x K(\cdot, t)\|_{L^1} \leq C \left[ \frac{1}{\sqrt{t}} + t^2 e^{\frac{4}{27} a^3 t} \right], \quad (3.22)$$

where  $C$  is a positive constant independent of  $t$ .

The following lemma will be used in the proof of Lemma 3.11 below.

**LEMMA 3.10.**

$$\lim_{A \rightarrow +\infty} \int_{|y| > A} |K(\cdot, 1) * G(\cdot, 1 - h^{1/3})|(y) dy = 0, \quad \text{uniformly in } h \in [0, 1]. \quad (3.23)$$

*Proof.* We recall that

$$\hat{K}(\xi, 1) = e^{-[\xi^2 - \xi^{4/3} \Phi(\xi)]}, \quad \text{and} \quad \hat{G}(\xi, 1 - h^{1/3}) = \frac{1}{\sqrt{2\pi}} e^{-(1-h^{1/3})\xi^{4/3} \Phi(\xi)}.$$

From (3.18) we see that

$$\begin{aligned} &\int_{|y| > A} |K(\cdot, 1) * G(\cdot, 1 - h^{1/3})|(y) dy \\ &\leq \left( \int_{|y| > A} \frac{dy}{1+y^2} \right)^{\frac{1}{2}} \left( \int (1+y^2) |K(\cdot, 1) * G(\cdot, 1 - h^{1/3})(y)|^2 dy \right)^{\frac{1}{2}} \\ &\leq \frac{C}{\sqrt{A}} \|\hat{K}(\cdot, 1) \hat{G}(\cdot, 1 - h^{1/3})\|_{H^1} \leq \frac{C'}{\sqrt{A}}. \end{aligned}$$

This concludes the proof.  $\square$

The next lemma shows that  $(E(t))_{t \geq 0}$  is a  $C^0$ -semigroup on the Banach space  $Y$  and also on the Banach space  $X$ .

LEMMA 3.11. **i.)** If  $u_0 \in C_b(\mathbb{R})$ , then  $u(t) := E(t)u_0 \in C_b(\mathbb{R})$  for every  $t \geq 0$ . In addition,

$$\|E(t)\|_{\mathcal{L}(C_b(\mathbb{R}))} \leq C \cdot (1 + t^2 e^{\frac{4}{27}a^3 t}), \quad \text{for all } t > 0. \quad (3.24)$$

Moreover,  $(E(t))_{t \geq 0}$  is a  $C^0$ -semigroup on  $Y$ .

**ii.)**  $(E(t))_{t \geq 0}$  is a  $C^0$ -semigroup on  $X$ . Furthermore, (3.24) remains true if the space  $C_b(\mathbb{R})$  is replaced by  $C_b^1(\mathbb{R})$ .

*Proof.* **i.)** Let  $u_0 \in C_b(\mathbb{R})$ , and  $t > 0$ . Since  $u(x, t) = E(t)u_0(x) = \frac{1}{\sqrt{2\pi}} \int K(x - y, t)u_0(y)dy$ , it follows that

$$\|u(t)\|_{L^\infty} \leq \frac{1}{\sqrt{2\pi}} \|u_0\|_{L^\infty} \|K(\cdot, t)\|_{L^1} \leq C \cdot (1 + t^2 e^{\frac{4}{27}a^3 t}) \|u_0\|_{L^\infty},$$

where the last inequality is a consequence of Lemma 3.6. Moreover,

$$|u(x + h, t) - u(x, t)| \leq \frac{\|u_0\|_{L^\infty}}{\sqrt{2\pi}} \|K(\cdot + h, t) - K(\cdot, t)\|_{L^1} \rightarrow 0 \text{ as } h \rightarrow 0.$$

Thus, we have proved that if  $u_0 \in C_b(\mathbb{R})$ , then  $u(t) \in C_b(\mathbb{R})$  for all  $t \geq 0$ . In addition, one can see that  $E(t + s)\phi = E(t)E(s)\phi$ , for all  $t, s \geq 0$ , and  $\phi \in C_b(\mathbb{R})$ .

Suppose now that  $t = 0$ ,  $u_0 \in Y \setminus \{0\}$ , and  $h \in (0, 1)$ . Since  $\hat{K}(0, h) = \frac{1}{\sqrt{2\pi}} \int K(z, h)dz = 1$ , and using (3.11) we have that

$$\begin{aligned} |u(x, h) - u_0(x)| &= \frac{1}{\sqrt{2\pi}} \left| \int K(z, h)(u_0(x - z) - u_0(x))dz \right| \\ &= \frac{1}{\sqrt{2\pi}} \left| \int h^{-1/2} (K(\cdot, 1) * G(\cdot, 1 - h^{1/3}))(h^{-1/2}z)(u_0(x - z) - u_0(x))dz \right| \\ &= \frac{1}{\sqrt{2\pi}} \left| \int (K(\cdot, 1) * G(\cdot, 1 - h^{1/3}))(y)(u_0(x - y\sqrt{h}) - u_0(x))dy \right|. \end{aligned} \quad (3.25)$$

Let  $\epsilon > 0$ . By Lemma 3.10, there exists  $A > 0$ , such that for every  $h \in [0, 1)$ ,

$$\int_{|y| > A} |K(\cdot, 1) * G(\cdot, 1 - h^{1/3})|(y)dy < \frac{\sqrt{2\pi}\epsilon}{4\|u_0\|_{L^\infty}}. \quad (3.26)$$

Since  $u_0$  is uniformly continuous, there exists  $\delta > 0$  such that for all  $z, w \in \mathbb{R}$ ,

$$\text{if } |z - w| < \delta, \text{ then } |u_0(z) - u_0(w)| < \sqrt{2\pi}\epsilon / (2C\|K(\cdot, 1)\|_{L^1}), \quad (3.27)$$

where  $C$  is a positive constant such that  $\|G(\cdot, 1 - h^{1/3})\|_{L^1} \leq C$  for all  $h \in [0, 1/2)$  (see Lemma 3.5). Let  $h \in (0, \min\{\frac{1}{2}, \frac{\delta^2}{A^2}\})$ . Using (3.25)-(3.27), we get

$$\begin{aligned} |u(x, h) - u_0(x)| &\leq \frac{2\|u_0\|_{L^\infty}}{\sqrt{2\pi}} \int_{|y| > A} |K(\cdot, 1) * G(\cdot, 1 - h^{1/3})(y)|dy \\ &\quad + \frac{1}{\sqrt{2\pi}} \int_{|y| \leq A} |K(\cdot, 1) * G(\cdot, 1 - h^{1/3})(y)| |u_0(x - y\sqrt{h}) - u_0(x)|dy \\ &\leq \frac{\epsilon}{2} + \frac{\epsilon}{2C\|K(\cdot, 1)\|_{L^1}} \|K(\cdot, 1) * G(\cdot, 1 - h^{1/3})\|_{L^1} \leq \epsilon, \end{aligned}$$

for all  $x \in \mathbb{R}$ , where the last inequality is a consequence of Young's inequality. Therefore,

$$\lim_{h \rightarrow 0} \|u(h) - u_0\|_{L^\infty} = 0. \quad (3.28)$$

We notice that if  $u_0 \in Y$ , then  $u(t) = E(t)u_0 \in Y$  for all  $t \geq 0$ . In fact, assume  $t > 0$  and let  $\epsilon > 0$  be given. Since  $u_0$  is uniformly continuous, there exists  $\delta > 0$

such that if  $|h| < \delta$ , then  $|u_0(x+h) - u_0(x)| < \epsilon\sqrt{2\pi}/\|K(\cdot, t)\|_{L^1}$ , for any  $x \in \mathbb{R}$ . Suppose  $|h| < \delta$ , then

$$|u(x+h, t) - u(x, t)| \leq \frac{1}{\sqrt{2\pi}} \int |K(y, t)| |u_0(x-y+h) - u_0(x-y)| dy \leq \epsilon, \quad \text{for all } x \in \mathbb{R}.$$

Hence,  $u(t)$  is uniformly continuous, for all  $t > 0$ .

Assume now that  $t > 0$ , and  $u_0 \in Y$ . It follows from (3.28) and the semigroup property that

$$\lim_{h \downarrow 0} \|E(t+h)u_0 - E(t)u_0\|_{L^\infty} = 0.$$

On the other hand, for  $h > 0$ , one can see that

$$\begin{aligned} |u(x, t-h) - u(x, t)| &= \frac{1}{\sqrt{2\pi}} \left| \int K(x-y, t-h) (u_0(y) - u(y, h)) dy \right| \\ &\leq \frac{1}{\sqrt{2\pi}} \|K(\cdot, t-h)\|_{L^1} \|u(h) - u_0\|_{L^\infty} \\ &\leq C \cdot (1 + (t-h)^2 e^{\frac{4}{27}a^3(t-h)}) \|u(h) - u_0\|_{L^\infty}, \end{aligned}$$

where the last inequality is a consequence of Lemma 3.6. Equation (3.28) and the last inequality imply that

$$\lim_{h \downarrow 0} \|E(t-h)u_0 - E(t)u_0\|_{L^\infty} = 0.$$

**ii.)** Let  $u_0 \in X$ . By item **i.)** above, we already know that  $u \in C([0, +\infty); Y)$ , where  $u(t) = E(t)u_0$  for all  $t \geq 0$ . We will now prove that  $\partial_x u(t) \in C_b(\mathbb{R})$ ,  $\partial_x u(t)$  is uniformly continuous, and  $\lim_{h \rightarrow 0} \|\partial_x u(t+h) - \partial_x u(t)\|_{L^\infty} = 0$ , for all  $t \geq 0$ . Suppose first that  $t > 0$ . Then

$$\begin{aligned} &\left| \frac{u(x+h, t) - u(x, t)}{h} - \frac{1}{\sqrt{2\pi}} K(\cdot, t) * u'_0(x) \right| \\ &= \frac{1}{\sqrt{2\pi}} \left| \left( K(\cdot, t) * \frac{u_0(\cdot+h) - u_0(\cdot)}{h} \right) (x) - K(\cdot, t) * u'_0(x) \right| \\ &\leq \frac{1}{\sqrt{2\pi}} \int |K(x-y, t)| \left| \frac{u_0(y+h) - u_0(y)}{h} - u'_0(y) \right| dy. \end{aligned}$$

The dominated convergence theorem and Lemma 3.6 imply that the last expression tends to zero as  $h$  goes to zero. Then there exists

$$\partial_x u(x, t) = \frac{1}{\sqrt{2\pi}} K(\cdot, t) * u'_0(x), \quad \text{for all } x \in \mathbb{R}, \text{ and } t > 0. \quad (3.29)$$

It is easy to see that the last expression is also valid if we only require that  $u_0 \in C_b^1(\mathbb{R})$ . It follows from (3.29) and Lemma 3.6 that

$$\|\partial_x u(\cdot, t)\|_{L^\infty} \leq C \cdot (1 + t^2 e^{\frac{4}{27}a^3 t}) \|u'_0\|_{L^\infty}, \quad \text{for all } t > 0.$$

Using the fact that  $u'_0$  is uniformly continuous, (3.29), and Lemma 3.6, it follows that  $\partial_x u(\cdot, t)$  is uniformly continuous, for all  $t > 0$ .

Finally, since  $(E(t))_{t \geq 0}$  is a  $C^0$ -semigroup on the space  $Y$  and using (3.29), we see that  $\lim_{h \rightarrow 0} \|\partial_x u(t+h) - \partial_x u(t)\|_{L^\infty} = 0$ , for all  $t \geq 0$ .  $\square$

**3.2. Local Theory in the Space  $X$ .** In this Sub-section we will use the Banach fixed-point theorem on an appropriate complete metric space to find a local-in-time solution to the integral equation associated to the IVP (3.1). The following lemma will be helpful during the proof of Theorem 3.1 below.

LEMMA 3.12. *Suppose that  $u \in C([0, T]; X)$ . We define*

$$D(\cdot, t) := \int_0^t K(\cdot, t-s) * \frac{1}{2} \partial_x u^2(\cdot, s) ds, \quad \text{for } t \in [0, T]. \quad (3.30)$$

Then  $D \in C([0, T]; X)$ .

*Proof. i.)* Let  $t \in (0, T]$ . Now we first prove that  $D(t) \in X$ . In fact,

$$\begin{aligned} \|D(\cdot, t)\|_{L^\infty} &\leq \sup_{s \in [0, T]} \|u(\cdot, s)\|_{C_b^1}^2 \int_0^t \|K(\cdot, t-s)\|_{L^1} ds \\ &\leq C \|u\|_{C([0, T]; X)}^2 \int_0^t (1 + (t-s)^2 e^{\frac{4}{27} a^3 (t-s)}) ds, \end{aligned}$$

where the last inequality is a consequence of Lemma 3.6. Then

$$\|D(\cdot, t)\|_{L^\infty} \leq C' \|u\|_{C([0, T]; X)}^2 \nu(t), \quad (3.31)$$

where

$$\nu(r) := r + r^2 e^{\frac{4}{27} a^3 r}, \quad \text{for all } r \geq 0. \quad (3.32)$$

Moreover,

$$\|D(\cdot + h, t) - D(\cdot, t)\|_{L^\infty} \leq \int_0^t \|K(\cdot, t-s)\|_{L^1} \left\| \frac{1}{2} \partial_x u^2(\cdot + h, s) - \frac{1}{2} \partial_x u^2(\cdot, s) \right\|_{L^\infty} ds.$$

Using the fact that  $\partial_x u(\cdot, s)u(\cdot, s)$  is uniformly continuous on  $\mathbb{R}$ , for all  $s \in [0, T]$ , Lemma 3.6, and the dominated convergence theorem, it follows from the last inequality that  $D(t)$  is uniformly continuous on  $\mathbb{R}$ . Now we claim that there exists  $\frac{\partial D}{\partial x}(x, t)$  (in the classical sense), for all  $x \in \mathbb{R}$ , and

$$\begin{aligned} \frac{\partial D}{\partial x}(x, t) &= \int_0^t \partial_x (K(\cdot, t-s) * \frac{1}{2} \partial_x u^2(\cdot, s))(x) ds \\ &= \int_0^t \partial_x K(\cdot, t-s) * \frac{1}{2} \partial_x u^2(\cdot, s)(x) ds, \quad \text{for all } x \in \mathbb{R}. \end{aligned} \quad (3.33)$$

We now establish the last claim. It follows from Lemmas 3.6, 3.9, and 3.4 that  $K(\cdot, t-s) * \frac{1}{2} \partial_x u^2(\cdot, s) \in C^1(\mathbb{R}) \cap W^{1, \infty}(\mathbb{R})$ , and

$$\partial_x (K(\cdot, t-s) * \frac{1}{2} \partial_x u^2(\cdot, s))(x) = (\partial_x K(\cdot, t-s) * \frac{1}{2} \partial_x u^2(\cdot, s))(x), \quad (3.34)$$

for all  $x \in \mathbb{R}$  and  $s \in [0, t)$ . Moreover,

$$\begin{aligned} &\left| \frac{D(x+h, t) - D(x, t)}{h} - \int_0^t \partial_x K(\cdot, t-s) * \frac{1}{2} \partial_x u^2(\cdot, s)(x) ds \right| \\ &= \left| \int_0^t \left( \frac{K(\cdot+h, t-s) - K(\cdot, t-s)}{h} - \partial_x K(\cdot, t-s) \right) * \frac{1}{2} \partial_x u^2(\cdot, s)(x) ds \right|. \end{aligned} \quad (3.35)$$

In addition,

$$\begin{aligned}
& \left\| \left( \frac{K(\cdot + h, t - s) - K(\cdot, t - s)}{h} - \partial_x K(\cdot, t - s) \right) * \frac{1}{2} \partial_x u^2(\cdot, s) \right\|_{L^\infty} \\
& \leq \|u\|_{C([0, T]; X)}^2 \left\| \frac{K(\cdot + h, t - s) - K(\cdot, t - s)}{h} - \partial_x K(\cdot, t - s) \right\|_{L^1} \\
& \leq \|u\|_{C([0, T]; X)}^2 \left( \frac{1}{|h|} \left| \int_0^h \|\partial_x K(\cdot + y, t - s)\|_{L^1} dy \right| + \|\partial_x K(\cdot, t - s)\|_{L^1} \right) \\
& \leq C \|u\|_{C([0, T]; X)}^2 \left( \frac{1}{\sqrt{t - s}} + (t - s)^2 e^{\frac{4}{27} a^3 (t - s)} \right) \in L^1((0, t), ds), \quad (3.36)
\end{aligned}$$

where the last inequality is a consequence of Lemma 3.9. The claim now follows from (3.34)-(3.36) and the dominated convergence theorem.

It follows directly from (3.33) and Lemma 3.9 that

$$\|\partial_x D(\cdot, t)\|_{L^\infty} \leq C \|u\|_{C([0, T]; X)}^2 \mu(t), \quad (3.37)$$

where

$$\mu(r) := \sqrt{r} + r^2 e^{\frac{4}{27} a^3 r}, \quad \text{for all } r \geq 0. \quad (3.38)$$

The fact that  $\partial_x D(\cdot, t)$  is uniformly continuous on  $\mathbb{R}$  can be shown similarly to the analogous result for  $D(\cdot, t)$ , using Lemma 3.9 instead of Lemma 3.6.

**ii.)** We will now prove that  $D \in C([0, T]; X)$ . Let  $t \in [0, T]$ . We first assume that  $h > 0$ . Then

$$\|D(\cdot, t + h) - D(\cdot, t)\|_{L^\infty} \leq I_1(t, h) + I_2(t, h),$$

where

$$\begin{aligned}
I_1(t, h) & := \int_0^t \left\| (K(\cdot, t + h - s) - K(\cdot, t - s)) * \frac{1}{2} \partial_x u^2(\cdot, s) \right\|_{L^\infty} ds, \quad \text{and} \\
I_2(t, h) & := \int_t^{t+h} \left\| K(\cdot, t + h - s) * \frac{1}{2} \partial_x u^2(\cdot, s) \right\|_{L^\infty} ds.
\end{aligned}$$

We see that

$$\begin{aligned}
& \left\| (K(\cdot, t + h - s) - K(\cdot, t - s)) * \frac{1}{2} \partial_x u^2(\cdot, s) \right\|_{L^\infty} \\
& \leq \|u\|_{C([0, T]; X)}^2 \|K(\cdot, t + h - s) - K(\cdot, t - s)\|_{L^1} \\
& \leq C \|u\|_{C([0, T]; X)}^2 (1 + T^2 e^{\frac{8}{27} a^3 T}) \in L^1((0, t), ds),
\end{aligned}$$

where the last inequality follows from Lemma 3.6 and the fact that  $h \in (0, T)$ . Thus, using Lemma 3.11 and the dominated convergence theorem we have that

$$I_1(t, h) = \sqrt{2\pi} \int_0^t \|(E(h) - 1)E(t - s) \frac{1}{2} \partial_x u^2(\cdot, s)\|_{L^\infty} ds \rightarrow 0, \quad \text{as } h \downarrow 0.$$

Moreover, using Lemma 3.6 we get like in (3.31) that

$$I_2(t, h) \leq C \|u\|_{C([0, T]; X)}^2 \nu(h) \rightarrow 0, \quad \text{as } h \downarrow 0,$$

where  $\nu(\cdot)$  is given by (3.32). Hence,

$$\lim_{h \downarrow 0} \|D(\cdot, t + h) - D(\cdot, t)\|_{L^\infty} = 0. \quad (3.39)$$

On the other hand, it follows from (3.33) that

$$\|\partial_x D(\cdot, t + h) - \partial_x D(\cdot, t)\|_{L^\infty} \leq J_1(t, h) + J_2(t, h),$$

where

$$\begin{aligned} J_1(t, h) &:= \int_0^t \|(\partial_x K(\cdot, t+h-s) - \partial_x K(\cdot, t-s)) * \frac{1}{2} \partial_x u^2(\cdot, s)\|_{L^\infty} ds, \quad \text{and} \\ J_2(t, h) &:= \int_t^{t+h} \|\partial_x K(\cdot, t+h-s) * \frac{1}{2} \partial_x u^2(\cdot, s)\|_{L^\infty} ds. \end{aligned}$$

It follows directly from Lemma 3.9 that

$$\begin{aligned} J_2(t, h) &\leq \|u\|_{C([0, T]; X)}^2 \int_0^h \|\partial_x K(\cdot, \tau)\|_{L^1} d\tau \\ &\leq C \|u\|_{C([0, T]; X)}^2 \mu(h) \rightarrow 0, \quad \text{as } h \downarrow 0, \end{aligned}$$

where  $\mu(\cdot)$  is given by (3.38). To estimate  $J_1(t, h)$  we first extend  $\partial_x K$  for all times in the following way:

$$H(\cdot, s) := \begin{cases} \partial_x K(\cdot, s), & \text{if } s \in [0, T], \\ 0, & \text{if } s \in \mathbb{R} \setminus [0, T]. \end{cases}$$

We note that  $H \in L^1(\mathbb{R}^2)$ . In fact, by Lemma 3.9 we get

$$\int \int |H(x, s)| dx ds = \int_0^T \|\partial_x K(\cdot, s)\|_{L^1} ds \leq C \mu(T).$$

Then

$$\begin{aligned} J_1(t, h) &\leq \|u\|_{C([0, T]; X)}^2 \int_0^t \|\partial_x K(\cdot, \tau+h) - \partial_x K(\cdot, \tau)\|_{L^1} d\tau \\ &\leq \|u\|_{C([0, T]; X)}^2 \int \int |H(x, \tau+h) - H(x, \tau)| dx d\tau \rightarrow 0, \quad \text{as } h \downarrow 0, \end{aligned}$$

where the last assertion follows from the continuity of translations in  $L^1(\mathbb{R}^2)$ . Hence,

$$\lim_{h \downarrow 0} \|\partial_x D(\cdot, t+h) - \partial_x D(\cdot, t)\|_{L^\infty} = 0. \quad (3.40)$$

It follows from (3.39) and (3.40) that  $\lim_{h \downarrow 0} \|D(\cdot, t+h) - D(\cdot, t)\|_{C_b^1} = 0$ .

The case when  $t \in (0, T]$  and  $h < 0$  can be shown similarly to the previous case. This finishes the proof of the lemma.  $\square$

The next theorem is the main result of this section, it states local-in-time existence of the solution of the integral equation associated to the IVP (3.1).

**THEOREM 3.1.** *Suppose  $u_0 \in X$ . Then there exist  $T = T(\|u_0\|_{C_b^1}) > 0$  and a unique function  $u \in C([0, T]; X)$  satisfying the integral equation*

$$u(\cdot, t) = E(t)u_0(\cdot) - \frac{1}{2} \int_0^t E(t-s) \partial_x u^2(\cdot, s) ds, \quad (3.41)$$

where  $E(t)$  is defined by (3.6).

*Proof.* Let  $M := 1 + 2\|u_0\|_{C_b^1}$ . Let  $T > 0$  be fixed.  $T$  will be suitably chosen later. We now consider the nonlinear operator  $A$  given by

$$(Af)(\cdot, t) := E(t)u_0(\cdot) - \frac{1}{2} \int_0^t E(t-s) \partial_x f^2(\cdot, s) ds,$$

defined on the complete metric space

$$\Theta_T^M := \left\{ f \in C([0, T]; X); \sup_{t \in [0, T]} \|f(\cdot, t)\|_{C_b^1} \leq M \right\}.$$

Let  $f \in \Theta_T^M$ . It follows from Lemmas 3.11 and 3.12 that  $Af \in C([0, T]; X)$ .

We will now prove that we can choose  $T = \tilde{T} > 0$  small enough such that  $A(\Theta_T^M) \subset \Theta_T^M$ . Suppose  $f \in \Theta_T^M$ . By Lemma 3.11 we know that  $\lim_{h \downarrow 0} \|(E(h) - 1)u_0\|_{C_b^1} = 0$ . Then there exists  $\delta = \delta(\|u_0\|_{C_b^1}) > 0$  such that if  $0 \leq h \leq \delta$ , then  $\|E(h)u_0\|_{C_b^1} \leq \frac{1}{2}(1 + 3\|u_0\|_{C_b^1})$ . If  $T \leq \delta$ , using Lemmas 3.6 and 3.9, and (3.33), we get

$$\begin{aligned} \|(Af)(\cdot, t)\|_{C_b^1} &\leq \frac{1}{2}(1 + 3\|u_0\|_{C_b^1}) \\ &+ \frac{1}{2\sqrt{2\pi}} \int_0^t (\|K(\cdot, t-t')\|_{L^1} + \|\partial_x K(\cdot, t-t')\|_{L^1}) \|f\|_{C([0, T]; X)}^2 dt' \\ &\leq \frac{1}{2}(1 + 3\|u_0\|_{C_b^1}) + M^2 C \int_0^t \left[ \frac{1}{\sqrt{\tau}} + \tau^2 e^{\frac{4}{27}a^3\tau} \right] d\tau \\ &\leq \frac{1}{2}(1 + 3\|u_0\|_{C_b^1}) + M^2 C \mu(T), \end{aligned}$$

for all  $t \in [0, T]$ , where  $\mu(\cdot)$  is given by (3.38). Take  $T^\dagger > 0$  such that  $M^2 C \mu(T^\dagger) \leq \frac{1}{2}(1 + \|u_0\|_{C_b^1})$ . Thus, if  $\tilde{T} \in (0, \min\{\delta, T^\dagger\}]$ , then  $\|(Af)(\cdot, t)\|_{C_b^1} \leq M$  for all  $t \in [0, \tilde{T}]$ .

Finally, we will prove that there exists  $T' \in (0, \tilde{T}]$  such that  $A$  is contractive on  $\Theta_{T'}^M$ . Suppose that  $f, g \in \Theta_{T'}^M$ . Let  $t \in [0, \tilde{T}]$ . Then

$$\begin{aligned} &\|(Af)(\cdot, t) - (Ag)(\cdot, t)\|_{C_b^1} \\ &\leq C \int_0^t (\|K(\cdot, t-t')\|_{L^1} + \|\partial_x K(\cdot, t-t')\|_{L^1}) \|\partial_x f^2(\cdot, t') - \partial_x g^2(\cdot, t')\|_{L^\infty} dt' \\ &\leq C \int_0^t (\|K(\cdot, t-t')\|_{L^1} + \|\partial_x K(\cdot, t-t')\|_{L^1}) \\ &\quad \times [\|f(\cdot, t')\|_{L^\infty} \|\partial_x(f(\cdot, t') - g(\cdot, t'))\|_{L^\infty} + \|f(\cdot, t') - g(\cdot, t')\|_{L^\infty} \|\partial_x g(\cdot, t')\|_{L^\infty}] dt' \\ &\leq CM \|f - g\|_{C([0, \tilde{T}]; X)} \mu(t). \end{aligned}$$

Taking  $T' \in (0, \tilde{T}]$  such that  $CM \mu(T') < 1$ , it follows that  $A$  is a contraction on  $\Theta_{T'}^M$ . Therefore, the mapping  $A$  has a unique fixed point  $u \in \Theta_{T'}^M$ , which satisfies equation (3.41) with  $T' = T'(\|u_0\|_{C_b^1}) > 0$ . The unicity of the solution of equation (3.41) in the class  $C([0, T']; X)$  is a consequence of Proposition 3.1 below.  $\square$

The next proposition shows the continuous dependance of the solutions of equation (3.41) on the initial data.

**PROPOSITION 3.1.** *Suppose that  $u, v \in C([0, T]; X)$  are solutions of equation (3.41) with initial data  $u_0, v_0 \in X$  respectively. Then for all  $t \in [0, T]$  we have*

$$\|u(\cdot, t) - v(\cdot, t)\|_{C_b^1} \leq C(T) e^{\alpha t} \|u_0 - v_0\|_{C_b^1}, \quad (3.42)$$

where  $\alpha := \tilde{C}^2 B(\frac{1}{2}, \frac{1}{2})$  and  $C(T) := C'(T)(1 + 2\tilde{C}\sqrt{T})$ , where  $C'(T)$  is given by (3.45) and  $\tilde{C} = C(T, \|u\|_{C([0, T]; X)}, \|v\|_{C([0, T]; X)})$  is given by (3.47). Here  $B(\cdot, \cdot)$

denotes the beta function defined by

$$B(x, y) := \int_0^1 t^{x-1}(1-t)^{y-1} dt,$$

for  $\Re(x), \Re(y) > 0$ .

*Proof.* Let  $t \in [0, T]$ . We write  $w(\cdot, t) := u(\cdot, t) - v(\cdot, t)$ . Then

$$\|w(\cdot, t)\|_{C_b^1} \leq \|E(t)(u_0 - v_0)\|_{C_b^1} + \frac{1}{2} \left\| \int_0^t E(t-t')(\partial_x u^2(\cdot, t') - \partial_x v^2(\cdot, t')) dt' \right\|_{C_b^1}. \quad (3.43)$$

It follows from Lemma 3.11 that

$$\|E(t)(u_0 - v_0)\|_{C_b^1} \leq C'(T)\|u_0 - v_0\|_{C_b^1}, \quad (3.44)$$

where

$$C'(T) := C \cdot (1 + T^2 e^{\frac{4}{27} a^3 T}). \quad (3.45)$$

Moreover, by Lemmas 3.6 and 3.9, we get

$$\begin{aligned} & \frac{1}{2} \left\| \int_0^t E(t-t')(\partial_x u^2(\cdot, t') - \partial_x v^2(\cdot, t')) dt' \right\|_{C_b^1} \\ & \leq \frac{\|u\|_{C([0, T]; X)} + \|v\|_{C([0, T]; X)}}{\sqrt{2\pi}} \\ & \quad \times \int_0^t (\|K(\cdot, t-t')\|_{L^1} + \|\partial_x K(\cdot, t-t')\|_{L^1}) \|w(\cdot, t')\|_{C_b^1} dt' \\ & \leq C \cdot (\|u\|_{C([0, T]; X)} + \|v\|_{C([0, T]; X)}) \\ & \quad \times \int_0^t \left[ 1 + (t-t')^2 e^{\frac{4}{27} a^3 (t-t')} + \frac{1}{\sqrt{t-t'}} \right] \|w(\cdot, t')\|_{C_b^1} dt' \\ & \leq \tilde{C} \int_0^t \frac{\|w(\cdot, t')\|_{C_b^1}}{\sqrt{t-t'}} dt', \end{aligned} \quad (3.46)$$

where

$$\tilde{C} := C \cdot (1 + T^{5/2} e^{\frac{4}{27} a^3 T}) (\|u\|_{C([0, T]; X)} + \|v\|_{C([0, T]; X)}). \quad (3.47)$$

Thus, it follows from (3.43), (3.44) and (3.46) that

$$\|w(\cdot, t)\|_{C_b^1} \leq C'(T)\|u_0 - v_0\|_{C_b^1} + \tilde{C} \int_0^t \frac{\|w(\cdot, t')\|_{C_b^1}}{\sqrt{t-t'}} dt'.$$

Then

$$\begin{aligned} & \|w(\cdot, t)\|_{C_b^1} \leq C'(T)\|u_0 - v_0\|_{C_b^1} \\ & + \tilde{C} \int_0^t \frac{1}{\sqrt{t-t'}} \left[ C'(T)\|u_0 - v_0\|_{C_b^1} + \tilde{C} \int_0^{t'} \frac{\|w(\cdot, r)\|_{C_b^1}}{\sqrt{t'-r}} dr \right] dt' \\ & \leq C'(T)(1 + 2\tilde{C}\sqrt{T})\|u_0 - v_0\|_{C_b^1} + \tilde{C}^2 \int_0^t \int_r^t \frac{\|w(\cdot, r)\|_{C_b^1}}{\sqrt{t-t'}\sqrt{t'-r}} dt' dr \\ & = C'(T)(1 + 2\tilde{C}\sqrt{T})\|u_0 - v_0\|_{C_b^1} + \tilde{C}^2 B\left(\frac{1}{2}, \frac{1}{2}\right) \int_0^t \|w(\cdot, r)\|_{C_b^1} dr. \end{aligned}$$

The proposition now follows by applying Gronwall's inequality to the last expression.  $\square$

**3.3. Future Work.** Some interesting problems remain, though: the study of the global well-posedness for the IVP (3.1) with initial data belonging to the space  $X$ , and the nonlinear stability theory of the travelling-wave solution of equation (1.1). These two problems will be addressed elsewhere.

**Acknowledgements:** The authors were supported by the ANR-France project COPTER (Conception, optimisation et prototypage d'ouvrages de lutte contre l'érosion en domaine littoral) under grant No. NT05-2\_42253. The authors wish to thank Rémi Carles (I3M-Université Montpellier 2) and Nathaël Alibaud (Département de Mathématiques de Besançon) for fruitful discussions.

#### REFERENCES

- [1] N. Alibaud, P. Azerad and D. Isèbe. *A non-monotone nonlocal conservation law for dune morphodynamics*. Preprint.
- [2] O. Durán, V. Schwämmle and H.J. Herrmann. *Breeding and solitary wave behavior of dunes*. Phys. Rev. E 72, 021308 (2005).
- [3] A.C. Fowler. *Dunes and drumlins*. In: Geomorphological fluid mechanics, N.J. Balmforth and A. Provenzale (eds.), Springer-Verlag, Berlin, pp. 430-454, 2001.
- [4] A.C. Fowler. *Evolution equations for dunes and drumlins*. Mathematics and environment (Spanish) (Paris, 2002). RACSAM Rev. R. Acad. Cienc. Exactas Fs. Nat. Ser. A Mat. 96 (2002), no. 3, 377-387.
- [5] A.C. Fowler. *Mathematics and the environment*. October 27, 2006. Preprint.
- [6] H.J. Herrmann and G. Sauer mann. *The shape of dunes*. Physica A 283 (2000), 24-30.
- [7] R.S. Johnson. *A modern introduction to the mathematical theory of water waves*. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 2004.
- [8] W. Rudin. *Functional Analysis*. Second edition. International Series in Pure and Applied Mathematics. McGraw-Hill, Inc., New York, 1991.

INSTITUT DE MATHÉMATIQUES ET MODÉLISATION DE MONTPELLIER (I3M)-UMR 5149 CNRS,  
UNIVERSITÉ MONTPELLIER 2, CC 051, PLACE EUGÈNE BATAILLON, 34095 MONTPELLIER CEDEX  
5, FRANCE

*E-mail address:* balvarez@math.uic.edu

*E-mail address:* azerad@math.univ-montp2.fr