

# A new test procedure of independence in copula models via chi-square-divergence

Salim Bouzebda, Amor Keziou

#### ▶ To cite this version:

Salim Bouzebda, Amor Keziou. A new test procedure of independence in copula models via chi-square-divergence. 2008. hal-00262077v1

# HAL Id: hal-00262077 https://hal.science/hal-00262077v1

Preprint submitted on 10 Mar 2008 (v1), last revised 19 Jun 2011 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A new test procedure of independence in Copula models via $\chi^2$ -divergence

Salim BOUZEBDA\* & Amor KEZIOU\*\*

\*LSTA-Université Paris 6, 175, rue du Chevaleret–Bt.A, Boîte 158, 75013 Paris,

bouzebda@ccr.jussieu.fr

\*\*Laboratoire de Mathématiques (UMR 6056) CNRS, Université de Reims Champagne-Ardenne and LSTA-Université Paris 6. UFR Sciences, Moulin de la Housse, B.P. 1039, 51687 Reims, France.

amor.keziou@univ-reims.fr

Key Words: Dependence function; Gumbel copula; pseudo-likelihood; Asymptotic theory; Multivariate rank statistic; Semiparametric estimation.

Abstract. We introduce a new test procedure of independence in the framework of parametric copulas with unknown marginals. The method is based essentially on the dual representation of  $\chi^2$ -divergence on signed finite measures. The asymptotic properties of the proposed estimate and the test statistic are studied under null and alternative hypothesis, with simple and standard limit distributions both when the parameter is an interior point or not.

### 1 Introduction and motivations

Parametric models for copulas have been intensively investigated during the last decades. *Copula* have become popular in applied statistics, because of the fact that they constitute a flexible and robust way to model dependence between the margins of random vectors. In this framework, semiparametric inference methods, based on *pseudo-likelihood*, have been applied to copulas by a number of authors (see, e.g., Shih and Louis (1995), Wang and Ding (2000) Tsukahara (2005) and the references therein). Throughout the available literature, investigations on the asymptotic properties of parametric estimators, as well as the relevant test statistics, have privileged the case where the parameter is an interior point of the admissible domain. However, for most parametric copula models of interest, the boundaries of the admissible parameter spaces include some important parameter values, typically among which, that corresponding to the independence of margins. This paper concentrates on this specific problem. We aim, namely, to investigate parametric inference procedures, in the case where the parameter belongs to the boundary of the admissible domain. In particular, the usual limit laws both for parametric copula estimators and test statistics become invalid under these limiting cases, and, in particular, under marginal independence. Motivated by this observation, we will introduce a new semiparametric inference procedure based on  $\chi^2$ -divergence and duality techniques. We will show that the proposed estimators remain asymptotically normal, even under the marginal independence assumption. This will allow us to introduce test statistics of independence, whose study will be made, both under the null and alternative hypotheses.

It is well known since the work of Sklar (1959) that the joint behavior of a bivariate vector  $(X_1, X_2)$  with d.f.  $F(x_1, x_2) := P(X_1 \le x_1, X_2 \le x_2)$ , and continuous marginal d.f.'s  $F_i(x_i) := P(X_i \le x_i), i = 1, 2$ , is characterized by the copula (or dependence function) Cassociated with  $F(\cdot, \cdot)$ . The copula function is defined, for all  $(u_1, u_2) \in (0, 1)^2$ , through the identity

$$C(u_1, u_2) := P \{F_1(X_1) \le u_1, F_2(X_2) \le u_2\}.$$

Many useful multivariate models for dependence between  $X_1$  and  $X_2$  turn out to be generated by *parametric* families of copulas of the form  $\{C_{\theta}; \theta \in \Theta\}$ , typically indexed by a vector valued parameter  $\theta \in \Theta \subseteq \mathbb{R}^p$  (see, e.g., Kimeldorf and Sampson (1975a), Kimeldorf and Sampson (1975b), and Joe (1993)). The nonparametric approach to copula estimation has been initiated by Deheuvels (1979b), who introduced and investigated the *empirical copula process*. In addition, Deheuvels (1980, 1981) described the limiting behavior of this empirical process (see, also Fermanian *et al.* (2004a) and the references therein). In this paper, we consider semiparametric copula models with unknown general margins.

In order to estimate the unknown *true* value of the parameter  $\theta \in \Theta$ , which we denote, throughout the sequel, by  $\theta_T \in \Theta$ , some semiparametric estimation procedures, based on the maximization, on the parameter space  $\Theta$ , of properly chosen *pseudo-likelihood* criterion, have been proposed by Oakes (1994), and studied by Genest et al. (1995), Shih and Louis (1995) and Wang and Ding (2000), Tsukahara (2005) among others. In each of these papers, some asymptotic normality properties are established for  $\sqrt{n}(\tilde{\theta} - \theta_T)$ , where  $\tilde{\theta} = \tilde{\theta}_n$  denotes a properly chosen estimator of  $\theta_T$ . This is achieved, provided that  $\theta_T$  lies in the *interior*, denoted by  $\Theta$ , of the parameter space  $\Theta \subseteq \mathbb{R}^p$ . On the other hand, the case where  $\theta_T \in \partial \Theta :=$  $\overline{\Theta} - \mathring{\Theta}$  is a *boundary value* of  $\Theta$ , has not been studied in a systematical way until present. Moreover, it turns out that, for the above-mentioned estimators, the asymptotic normality of  $\sqrt{n}(\theta - \theta_T)$ , may fail to hold for  $\theta_T \in \partial\Theta$ ; indeed, under some regularity conditions, when  $\theta$  is univariate, we can prove that the limit law is the distribution of  $Z1_{(Z\geq 0)}$  where Z is a centred normal variable, and that the limit law of the generalized pseudo-likelihood ratio statistic is a mixture of  $\chi^2_1$  law with one degree of freedom and Dirac measure at zero; see Bouzebda and Keziou (2007). Furthermore, when the parameter is multivariate, the derivation of the limit distributions under the null hypothesis of independence, becomes much more complex; see Self and Liang (1987). Also, the limit distributions are not standard which yields formidable numerical difficulties to calculate the critical value of the test. We cite below some examples of parametric copulas, for which marginal independence is verified for some specific values of the parameter  $\theta$ , on the boundary  $\partial \Theta$  of the admissible parameter set  $\Theta$ . We start with examples for which  $\theta$  varies within subsets of  $\mathbb{R}$ . Such is the case for the Gumbel (1960) family of copulas, which is one of the most popular model used to model bivariate extreme values. This family of copulas is defined, for  $0 < u_1, u_2 < 1$ , by

$$C_{\theta}(u_1, u_2) := \exp\left\{-\left[(-\log u_1)^{\theta} + (-\log u_2)^{\theta}\right]^{1/\theta}\right\}, \text{ for } \theta \in [1, \infty).$$
(1.1)

A useful family of copulas, due to Joe (1993), is given, for  $0 < u_1, u_2 < 1$ , by

$$C_{\theta}(u_1, u_2) := 1 - \left[ (1 - u_1)^{\theta} + (1 - u_2)^{\theta} - (1 - u_1)^{\theta} (1 - u_2)^{\theta} \right]^{1/\theta}, \text{ for } \theta \in [1, \infty).$$
(1.2)

The Galambos (1975) family of copulas is given, for  $0 < u_1, u_2 < 1$ , by

$$C_{\theta}(u_1, u_2) := u_1 u_2 \exp\left\{\left[(-\log u_1)^{-\theta} + (-\log u_2)^{-\theta}\right]^{-1/\theta}\right\}, \text{ for } \theta \in [0, \infty).$$
(1.3)

The Hüsler and Reiss (1989) family of copulas is defined, for  $0 < u_1, u_2 < 1$ , by

$$C_{\theta}(u_1, u_2) := u_2^{\Phi\left(\theta^{-1} + \frac{\theta}{2}\log\left[(-\log u_2)/(-\log u_1)\right]\right)} u_1^{\Phi\left(\theta^{-1} + \frac{\theta}{2}\log\left[(-\log u_1)/(-\log u_2)\right]\right)}, \qquad (1.4)$$

for  $\theta \in [0, \infty)$ , with  $\Phi(\cdot)$  denoting the standard normal N(0, 1) distribution function.

The Gumbel-Barnett copulas are given, for  $0 < u_1, u_2 < 1$ , by

$$C_{\theta}(u_1, u_2) := u_1 u_2 \exp\left\{-(1-\theta)(\log u_1)(\log u_2)\right\}, \quad \text{for} \quad \theta \in [0, 1].$$
(1.5)

The Clayton copulas are such that, for  $0 < u_1, u_2 < 1$ ,

$$C_{\theta}(u_1, u_2) = \left(u_1^{-\theta} - u_2^{-\theta} - 1\right)^{-1/\theta}, \quad \text{for} \quad \theta \in [0, \infty).$$
(1.6)

Parametric families of copulas with parameter  $\theta$  varying in  $\mathbb{R}^p$ , for some  $p \ge 2$ , include the following classical examples. Below, we set  $\theta = (\theta_1, \theta_2)^\top \in \mathbb{R}^2$ .

$$C_{\theta}(u_{1}, u_{2}) := \left\{ 1 + \left[ (u_{1}^{-\theta_{1}} - 1)^{\theta_{2}} + (u_{2}^{-\theta_{1}} - 1)^{\theta_{2}} \right]^{1/\theta_{2}} \right\}^{-1/\theta_{1}}, \ \theta \in ]0, \infty[\times[1, \infty[; (1.7)]]$$
$$C_{\theta}(u_{1}, u_{2}) := \exp\left\{ - \left[ \theta_{2}^{-1} \log\left( \exp\left( -\theta_{2} (\log u_{1})^{\theta_{1}} \right) + \exp\left( -\theta_{2} (\log u_{2})^{\theta_{1}} \right) - 1 \right) \right]^{1/\theta_{1}} \right\}, \ \theta \in [1, \infty[\times]0, \infty[.$$

For other examples of the kind, we refer to Joe (1997).

For each of the above examples, the independence case  $C_{\theta_T}(u_1, u_2) = u_1 u_2$  occurs at the boundary of the parameter space  $\Theta$ , i.e., when  $\theta_T = 1$  for the models (1.1), (1.2) and (1.5),  $\theta_T = 0$  for the models (1.3), (1.4) and (1.6),  $\theta_T = (0, 1)$  for the bivariate parameter model (1.7), and  $\theta_T = (1, 0)$  for the bivariate parameter model (1.8). In the sequel, we will denote by  $\theta_0$  the value of the parameter (when it exists), corresponding to the independence of the margins, i.e., the value of the parameter for which we have

$$C_{\theta_0}(u_1, u_2) := u_1 u_2$$
, for all  $(u_1, u_2) \in (0, 1)^2$ .

Hence,  $\theta_0 = 1$  for the models (1.1), (1.2) and (1.5),  $\theta_0 = 0$  for the models (1.3), (1.4) and (1.6),  $\theta_0 = (0, 1)$  for the model (1.7), and  $\theta_0 = (1, 0)$  for the model (1.8).

In contrast with the preceding examples, where  $\theta_0 \in \partial \Theta$  is a boundary value of  $\Theta$ , the case where  $\theta_0$  is an interior point of  $\Theta$  may, at times, occur, but is more seldom. An example where  $\theta_0 \in \mathring{\Theta}$  is given by the Farlie-Gumbel-Morgenstern (FGM) copula, defined by

$$C_{\theta}(u_1, u_2) := u_1 u_2 + \theta u_1 u_2 (1 - u_1)(1 - u_2), \ \theta \in \Theta := [-1, 1], \tag{1.9}$$

and for which  $\theta_0 = 0 \in \mathring{\Theta} = ]-1, 1[.$ 

In the present article, we will treat, in turn, parametric estimation of  $\theta_T$ , and tests of the independence assumption  $\theta_T = \theta_0$ . We consider both the case where  $\theta_0 \in \mathring{\Theta}$  is an interior point of  $\Theta$ , and the case where  $\theta_0 \in \partial \Theta$  is a boundary value of  $\Theta$ . To treat this case, we propose a new inference procedure, based on an estimation of  $\chi^2$ -divergence by duality technique. This method may be applied independently of the dimension of the parameter space. Also the limit law of the estimate of the parameter is normal and the limiting distribution of the proposed test statistic is  $\chi^2$  under independence, either when  $\theta_0$  is an interior point, or when  $\theta_0$  is a boundary point of  $\Theta$ , independently of the dimension of the parameter space. The idea is to include the parameter domain  $\Theta$  into an enlarged space, say  $\Theta_e$ , in order to render  $\theta_0$  an interior point of the new parameter space,  $\Theta_e$ . The conclusion is then obtained through an application of  $\chi^2$ -divergence and duality technique. Our methods rely on the fact that, under appropriate assumptions, the definition of the density  $c_{\theta}(\cdot, \cdot) :=$  $\frac{\partial^2}{\partial u_1 \partial u_2} C_{\theta}(\cdot, \cdot)$  of  $C_{\theta}(\cdot, \cdot)$ , pertaining to the models we consider, may be extended beyond the "standard" domain of variation  $\Theta$  of  $\theta$ . On the other hand, the definition of  $c_{\theta}(\cdot, \cdot)$  which corresponds to these extensions, is then, in general, not any more a density, and may, at times, become negative. For example, such is the case for the parametric models (1.1), (1.2), (1.3) and (1.4), for which  $c_{\theta}(\cdot, \cdot)$  is meaningful for some  $\theta \notin \Theta$ , but then, becomes negative over some non-negligible (with respect to Lebesgue's measure) subsets of  $(0, 1)^2$ . This implies that the log-likelihood of the data is not properly defined on the whole space  $\Theta_e$ . For this reason, we will use the  $\chi^2$ -divergence between signed finite measures. We will discuss this problem in more details, below, in section 2.

The remainder of the present paper is organized as follows. In section 2, we present our semiparametric inference procedure, based upon an optimization of the  $\chi^2$ -divergence between the model ( $C_{\theta}, \theta \in \Theta_e$ ) and the empirical copula associated to the data, and by using the *dual representation* of  $\chi^2$ -divergence. We then derive the asymptotic limiting distribution of the proposed estimator. It will become clear later on from our results, that the asymptotic normality of the estimate holds, even under the independence assumption, when, either,  $\theta_0$ is an interior, or a boundary point of  $\Theta$ , independently of the dimension of the parameter space. The proposed test statistic of independence is also studied, under the null hypothesis  $\mathcal{H}_0$  of independence, as well as under the alternative hypothesis. The limiting asymptotic distribution of the test statistic under the alternative hypothesis is used to derive an approximation to the power function. An application of the forthcoming results will allow us to evaluate the sample size necessary to guarantee a pre-assigned power level, with respect to a specified alternative. Finally, section 4 reports a short simulation results. The proofs of these results will be postponed to the appendix.

# 2 A semi-parametric estimation procedure through $\chi^2$ divergence

As mentioned earlier, the problem of estimating  $\theta$ , when  $\theta \in \partial \Theta$ , has not been systematically considered in the scientific literature; and the classical asymptotic normality property of the estimators is no more satisfied. To overcome this difficulty, in what follows, we enlarge the parameter space  $\Theta$  into a wider space  $\Theta_e \supset \Theta$ . This is tailored to let  $\theta_0$  become an interior point of  $\Theta_e$ . Naturally, we assume that the definition of the function  $c_{\theta}$  may be extended to  $\Theta_e$ . The difficulty associated with this construction is that, subject to a proper definition, the densities  $c_{\theta}$  of  $C_{\theta}$  with respect to the Lebesgue measure, may become negative on some non negligible subsets of  $I := (0, 1)^2$  (in this case  $c_{\theta}$  becomes the density of a signed measure, see remark 3.2). Note that just as Deheuvels empirical copula is not a copula, is not necessarily a copula density and fail to integrate to 1. When such is the case, a semiparametric estimation of  $\theta_T$  via log-likelihood cannot be used. To overcome this difficulty, we introduce a new inference procedure, based on  $\chi^2$ -divergence methods, and duality technique. Recall that the  $\chi^2$ -divergence between a bounded signed measure Q, and a probability P on  $\mathcal{D}$ , when  $Q \ll P$  is absolutely continuous with respect to P, is defined by

$$\chi^2(Q,P) := \int_{\mathcal{D}} \varphi\left(\frac{dQ}{dP}\right) \, dP, \text{ where } \varphi \ : \ x \in \mathbb{R} \mapsto \varphi(x) := \frac{1}{2} \left(x-1\right)^2. \tag{2.1}$$

In the sequel, we denote by  $\chi^2(\theta_0, \theta_T)$  the  $\chi^2$ -divergence between  $C_{\theta_0}$  and  $C_{\theta_T}$ . Applying the dual representation of  $\phi$ -divergence obtained by Broniatowski and Keziou (2006) Theorem 4.4, we readily obtain that  $\chi^2(\theta_0, \theta_T)$  can be rewritten into

$$\chi^2(\theta_0, \theta_T) := \sup_{f \in \mathcal{F}} \left\{ \int_I f \ dC_{\theta_0} - \int_I \varphi^*(f) \ dC_{\theta_T} \right\},$$
(2.2)

where  $\varphi^*$  is used to denote the convex conjugate of  $\varphi$ , namely, the function defined by

$$\varphi^*$$
:  $t \in \mathbb{R} \mapsto \varphi^*(t) := \sup_{x \in \mathbb{R}} \{tx - \varphi(x)\} = \frac{t^2}{2} + t,$ 

and  $\mathcal{F}$  is an arbitrary class of measurable functions, fulfilling the following condition  $\forall f \in \mathcal{F}; \quad \int |f| \ dC_{\theta_0}$  is finite and  $\varphi'(dC_{\theta_0}/dC_{\theta_T}) = \varphi'(1/c_{\theta_T}) \in \mathcal{F}.$  Furthermore, the sup in (2.2) is unique and achieved at  $f = \varphi'(1/c_{\theta_T}).$ 

Define the new parameter space  $\Theta_e$  of  $\theta$  as follows. Set

$$\Theta_e := \left\{ \theta \in \mathbb{R}^d \text{ such that } \int |\varphi'(1/c_\theta(u_1, u_2))| \ du_1 du_2 < \infty \right\}.$$
(2.3)

By choosing the class of functions  $\mathcal{F}$ , via

$$\mathcal{F} := \{ (u_1, u_2) \in I \mapsto \varphi'(1/c_\theta(u_1, u_2)) - 1 ; \theta \in \Theta_e \}$$

we infer from (2.2) the relation

$$\chi^{2}(\theta_{0},\theta_{T}) := \sup_{\theta \in \Theta_{e}} \left\{ \int_{I} \left( \frac{1}{c_{\theta}(u_{1},u_{2})} - 1 \right) du_{1} du_{2} - \int_{I} \left( \frac{1}{2} \frac{1}{c_{\theta}(u_{1},u_{2})^{2}} - \frac{1}{2} \right) dC_{\theta_{T}}(u_{1},u_{2}) \right\} (2.4)$$

It turns out that the supremum in (2.4) is reached iff  $\theta = \theta_T$ . Moreover, in general,  $\theta_T$  is an interior point of the new parameter space  $\Theta_e$ , especially under the null hypothesis of independence, namely, when  $\theta_T = \theta_0$ . Set

$$m(\theta, u_1, u_2) := \int_I \left(\frac{1}{c_{\theta}(u_1, u_2)} - 1\right) du_1 du_2 - \left\{\frac{1}{2}\frac{1}{c_{\theta}(u_1, u_2)^2} - \frac{1}{2}\right\}.$$
 (2.5)

In what follows, we propose to estimate the  $\chi^2$ -divergence  $\chi^2(\theta_0, \theta_T)$  between  $C_{\theta_0}$  and  $C_{\theta_T}$ , by

$$\widehat{\chi}^2(\theta_0, \theta_T) := \sup_{\theta \in \Theta_e} \int_I m(\theta, u_1, u_2) \ dC_n(u_1, u_2), \tag{2.6}$$

and to estimate the parameter  $\theta_T$  by

$$\widehat{\theta}_n := \arg \sup_{\theta \in \Theta_e} \left\{ \int_I m(\theta, u_1, u_2) \ dC_n(u_1, u_2) \right\},\tag{2.7}$$

where  $C_n$  is the modified empirical copula, defined by

$$C_n(u_1, u_2) := \frac{1}{n} \sum_{k=1}^n \mathbb{1}_{\{F_{1n}(X_{1k}) \le u_1\}} \mathbb{1}_{\{F_{2n}(X_{2k}) \le u_2\}}, \ (u_1, u_2) \in I,$$
(2.8)

and  $F_{jn}(t) := \frac{1}{n} \sum_{k=1}^{n} \mathbb{1}_{]-\infty,t]}(X_{jk}), \ j = 1, 2.$ 

**Remark 2.1** The set  $\Theta_e$  defined above is generally with non empty interior  $\mathring{\Theta}_e$ . In particular, we may check that  $\theta_0$  (the value corresponding to independence) belongs to  $\mathring{\Theta}_e$ , since the integral in (2.3) is finite; it is equal to zero when  $\theta = \theta_0$ , for any copula density  $c_{\theta}$ . For example, in the case of FGM copulas (1.9), it is easy to show that  $\Theta_e = \mathbb{R}$ . However, the explicit calculation of the integral in (2.3) may be complicated for some copulas, in such cases we use the Monte Carlo method to compute this integral for all  $\theta \in \Theta_e$ .

Statistic of the form

$$\Psi_n := \int_I \psi(u_1, u_2) \ dC_n(u_1, u_2),$$

belong to the general class of *multivariate rank statistics*. Their asymptotic properties have been investigated at length by a number of authors, among whom we may cite Ruymgaart *et al.* (1972), Ruymgaart (1974) and Rüschendorf (1976). In particular, the previous authors have provided regularity conditions, imposed on  $\psi(\cdot, \cdot)$ , which imply the asymptotic normality of  $\Psi_n$ . The corresponding arguments have been modified by Genest *et al.* (1995), Tsukahara (2005) and Fermanian *et al.* (2004b), as to establish almost sure convergence of the estimators they consider. In the same spirit, the limiting behavior, as  $n \to \infty$ , of our estimator and test statistic  $\mathbf{T}_n$ , will make an instrumental use of the general theory of multivariate rank statistics. Using some similar arguments as in Qin and Lawless (1994), the existence and consistency of our estimator and test statistic will be established through an application of the law of the iterated logarithm for empirical copula processes, in combination with general arguments from multivariate rank statistics theory. In the sequel, without loss of generality, we will limit ourselves to the case where the parameter is univariate. The extensions of our result in a multivariate framework may be achieved, at the price of additional technicalities, and under similar assumptions. We will make use of the following definitions.

- **Definition 2.1** (i) Let Q be the set of continuous functions q on [0,1] which are positive on (0,1), symmetric about 1/2, increasing on [0,1/2] and satisfy  $\int_0^1 \{q(t)\}^{-2} dt < \infty$ .
  - (ii) A function  $r : (0,1) \to (0,\infty)$  is called u-shaped if it is symmetric about 1/2 and increasing on (0,1/2].
- (iii) For  $0 < \beta < 1$  and u-shaped function r, we define

$$r_{\beta}(t) := \begin{cases} r(\beta t) & if \quad 0 < t \le 1/2; \\ r\{1 - \beta(1 - t)\} & if \quad 1/2 < t \le 1. \end{cases}$$

If for  $\beta > 0$  in a neighborhood of 0, there exists a constant  $M_{\beta}$ , such that  $r_{\beta} \leq M_{\beta}r$ on (0,1), then r is called a reproducing u-shaped function. We denote by  $\mathcal{R}$  the set of reproducing u-shaped functions.

Typical examples of elements in  $\mathcal{Q}$  and  $\mathcal{R}$  are given by

$$q(t) = [t(1-t)]^{\varsigma}, \ 0 < \zeta < 1/2, \quad r(t) = \varrho [t(1-t)]^{-\varsigma}, \ \varsigma \ge 0, \ \varrho \ge 0.$$

We will describe the asymptotic properties of the proposed estimate  $\widehat{\theta}_n$  under the following conditions.

(C.1) There exist functions  $r_{1,k}, r_{2,k} \in \mathcal{R}$  such that

$$\left|\frac{\partial^k}{\partial\theta^k} m(\theta_T, u_1, u_2)\right| \le r_{1,k}(u_1) r_{2,k}(u_2), \text{ for } k = 1, 2,$$

and

$$\left|\frac{\partial^3}{\partial \theta^3} m(\theta, u_1, u_2)\right| \le r_{1,3}(u_1) r_{2,3}(u_2) \text{ on a neighborhood } N(\theta_T) \text{ of } \theta_T,$$
  
where  $\int_I \{r_{1,k}(u_1) r_{2,k}(u_2)\}^2 dC_{\theta_T}(u_1, u_2) < \infty \text{ for } k = 1, 2, 3;$ 

- (C.2) The function  $(u_1, u_2) \in I \mapsto \frac{\partial}{\partial \theta} m(\theta_T, u_1, u_2)$  is of bounded variation on I;
- (C.3) For each  $\theta$ , the function  $\frac{\partial}{\partial \theta} m(\theta, u_1, u_2) : I \to \mathbb{R}$  is continuously differentiable, and there exist functions  $r_i \in \mathcal{R}$ ,  $\tilde{r}_i \in \mathcal{R}$  and  $q_i \in \mathcal{Q}$  for i = 1, 2, such that

$$\left|\frac{\partial^2}{\partial\theta\partial u_i}m(\theta, u_1, u_2)\right| \le \widetilde{r}_i(u_i)r_j(u_j), i, j = 1, 2 \text{ and } i \ne j,$$

and

$$\int_{I} \{q_i(u_i)\tilde{r}_i(u_i)r_j(u_j)\} \, dC_{\theta_T}(u_1, u_2) < \infty, \ i, j = 1, 2 \text{ and } i \neq j.$$

 $\operatorname{Set}$ 

$$\Xi := -E\left[\frac{\partial^2}{\partial \theta^2} m\left(\theta_T, F_1(X_{1k}), F_2(X_{2k})\right)\right],\tag{2.9}$$

and

$$\Sigma^2 := \operatorname{Var}\left[\frac{\partial}{\partial \theta}m\left(\theta_T, F_1(X_1), F_2(X_2)\right) + W_1(\theta_T, X_1) + W_2(\theta_T, X_2)\right], \qquad (2.10)$$

where

$$W_{i}(\theta_{T}, X_{i}) := \int_{I} \mathbb{1}_{\{F_{i}(X_{i}) \leq u_{i}\}} \frac{\partial^{2}}{\partial \theta \partial u_{i}} m(\theta_{T}, u_{1}, u_{2}) c_{\theta_{T}}(u_{1}, u_{2}) du_{1} du_{2}, \ i = 1, 2.$$

We can see that  $\Xi$  and  $W_i(X_i)$  can be defined, respectively, by

$$\Xi = E\left[\frac{\partial}{\partial \theta}m^2\left(\theta_T, F_1(X_{1k}), F_2(X_{2k})\right)\right]$$

and, for i = 1, 2,

$$W_i(\theta_T, X_i) = -\int_I \mathbb{1}_{\{F_i(X_i) \le u_i\}} \frac{\partial}{\partial \theta} m\left(\theta_T, u_1, u_2\right) \frac{\partial}{\partial u_i} m\left(\theta_T, u_1, u_2\right) c_{\theta_T}(u_1, u_2) du_1 du_2.$$

**Theorem 2.1** Assume that the conditions (C.1-2-3) hold.

(a) Let  $B(\theta_T, n^{-1/3}) := \{\theta \in \Theta_e, |\theta - \theta_T| \le n^{-1/3}\}$ . Then, as  $n \to \infty$ , with probability one, the function  $\theta \mapsto \int m(\theta, u_1, u_2) \ dC_n(u_1, u_2)$  reaches its maximum value at some point  $\widehat{\theta}_n$  in the interior of the interval  $B(\theta_T, n^{-1/3})$ . As a consequence, the estimate  $\widehat{\theta}_n$  of  $\theta$ is  $n^{1/3}$ -consistent, and satisfies

$$\int_{I} \frac{\partial}{\partial \theta} m\left(\widehat{\theta}_{n}, u_{1}, u_{2}\right) \ dC_{n}(u_{1}, u_{2}) = 0.$$

(b) As  $n \to \infty$ ,

$$\sqrt{n}(\widehat{\theta}_n - \theta_T) \xrightarrow{d} N(0, \Sigma^2 / \Xi^2).$$

The proof of Theorem 2.1 is postponed to section 6.

## **3** A test based on " $\chi^2$ -divergence"

One of the motivations of the present work is to build a statistical test of independence, based on  $\chi^2$ -divergence. In the framework of the parametric copula model, the null hypothesis, namely, the independence case  $C_{\theta}(u_1, u_2) = u_1 u_2$  corresponds to the condition that  $\mathcal{H}_0$  :  $\theta_T = \theta_0$ . We consider the alternative composite hypothesis  $\mathcal{H}_1$  :  $\theta_T \neq \theta_0$ . The corresponding generalized pseudo-likelihood ratio statistic is then given by

$$\mathbf{S}_{n}(\theta_{0},\tilde{\theta}) = 2\log \frac{\sup_{\theta \in \Theta} \prod_{k=1}^{n} c_{\theta}(\widehat{F}_{1n}(X_{1k}), \widehat{F}_{2n}(X_{2k}))}{\prod_{k=1}^{n} c_{\theta_{0}}(\widehat{F}_{1n}(X_{1k}), \widehat{F}_{2n}(X_{2k}))},$$

where, for j = 1, 2,  $\hat{F}_{jn}$  stands for n/(n+1) times the marginal empirical distribution function of the *j*-th variable  $X_j$ . The rescaling by the factor n/(n+1), avoids difficulties arising from potential unboundedness of  $\log c_{\theta}(u_1, u_2)$ , when either  $u_1$  or  $u_2$  tends to 1. Since,  $\theta_0$  is a boundary value of the parameter space  $\Theta$ , we can see, that the convergence in distribution of  $\mathbf{S}_n$  to a  $\chi^2$  random variable is not likely to hold. In order to bring a solution to this problem, we introduce the statistic

$$\mathbf{T}_n(\theta_0, \widehat{\theta}_n) := 2n \sup_{\theta \in \Theta_e} \int_I m(\theta, u_1, u_2) \ dC_n(u_1, u_2).$$
(3.1)

Below, we will show that, under the null hypothesis  $\mathcal{H}_0$ , the just-given statistic  $\mathbf{T}_n(\theta_0, \widehat{\theta}_n)$ converges in distribution to a  $\chi^2$  random variable. This property allows us to build a test of  $\mathcal{H}_0$  against  $\mathcal{H}_1$ , asymptotically of level  $\alpha$ . The limit law of  $\mathbf{T}_n(\theta_0, \widehat{\theta}_n)$  will also be given under the alternative hypothesis  $\mathcal{H}_1$ . The following additional conditions will be needed for the statement of our results.

(C.4) We have

$$\lim_{\theta \to \theta_0} \frac{\partial}{\partial u_i} m(\theta, u_1, u_2) = 0,$$

and there exist  $M_1 > 0$  and  $\delta_1 > 0$  such that, for all  $\theta$  in some neighborhood of  $\theta_0$ , one has, for i = 1, 2,

$$\left|\frac{\partial}{\partial \theta} m(\theta, u_1, u_2) \frac{\partial}{\partial u_i} m(\theta, u_1, u_2) c_{\theta}(u_1, u_2)\right| < M_1 r(u_i)^{-1.5+\delta_1} r(u_{3-i})^{0.5+\delta_1},$$
  
where  $r(u) := u(1-u)$  for  $u \in (0, 1);$ 

(C.5) There exist a neighborhood  $N(\theta_T)$  of  $\theta_T$ , and functions  $r_i \in \mathcal{R}$  such that for all  $\theta \in N(\theta_T)$ , we have

$$\left|\frac{\partial}{\partial\theta}m(\theta, u_1, u_2)\right| \le r_1(u_1)r_2(u_2) \text{ with } \int_I \{r_1(u_1)r_2(u_2)\}^2 dC_{\theta_T}(u_1, u_2) < \infty;$$

(C.6) There exist functions  $r_i, \tilde{r}_i \in \mathcal{R}, q_i \in \mathcal{Q} \ i = 1, 2$  such that

$$\left| \begin{array}{l} m(\theta_T, u_1, u_2) \right| \leq r_1(u_1) r_2(u_2), \\ \left| \frac{\partial}{\partial u_i} m(\theta, u_1, u_2) \right| \leq \widetilde{r}_i(u_i) r_j(u_j), i, j = 1, 2 \text{ and } i \neq j \end{array} \right|$$

with

$$\int_{I} \left\{ r_1(u_1) r_2(u_2) \right\}^2 dC_{\theta_T}(u_1, u_2) < \infty,$$

and

$$\int_{I} \{q_i(u_i)\widetilde{r}_i(u_i)r_j(u_j)\} \, dC_{\theta_T}(u_1, u_2) < \infty, \ i, j = 1, 2 \text{ and } i \neq j.$$

**Theorem 3.1** Assume that conditions (C.1-2-3-4) hold. Then, under the null hypothesis  $\mathcal{H}_0$ , the statistic  $\mathbf{T}_n$  converges in distribution to a  $\chi_1^2$  random variable (with 1 degree of freedom).

**Theorem 3.2** Assume that conditions (C.1-2-3) and (C.5-6) hold. Then, under the alternative hypothesis  $\mathcal{H}_1$ , we have

$$\sqrt{n}\left(\frac{\mathbf{T}_n}{2n} - \chi^2(\theta_0, \theta_T)\right)$$

converges to a centered normal random variable with variance

$$\sigma_{\chi^2}^2 := \operatorname{Var}\left[m\left(\theta_T, F_1(X_{1k}), F_2(X_{2k})\right) + Y_1(\theta_T, X_1) + Y_2(\theta_T, X_2)\right],$$
(3.2)

where

$$Y_i(\theta_T, X_i) := \int_I \mathbb{1}_{\{F_i(X_i) \le u_i\}} \frac{\partial}{\partial u_i} m\left(\theta_T, u_1, u_2\right) c_{\theta_T}(u_1, u_2) \ du_1 du_2.$$

The proof of Theorem 3.1 is postponed until section 6.

**Remark 3.1** An application of Theorem 3.1, leads to reject the null hypothesis  $\mathcal{H}_0: \theta_T = \theta_0$ , whenever the value of the statistic  $\mathbf{T}_n$  exceeds  $q_{1-\alpha}$ , namely, the  $(1-\alpha)$ -quantile of the law of the random variable  $\chi_1^2$ . The test corresponding to this rejection rule is then, asymptotically of level  $\alpha$ , when  $n \to \infty$ . Accordingly, the critical region is given by

$$CR := \{\mathbf{T}_n > q_{1-\alpha}\}. \tag{3.3}$$

The fact that this test is consistent follows from Theorem 2.3. Further, this theorem can be used to give an approximation to the power function  $\theta_T \mapsto \beta(\theta_T) := P_{\theta_T} \{CR\}$  in a similar way to Keziou and Leoni-Aubin (2005) and Keziou and Leoni-Aubin (2007). We so obtain that

$$\beta(\theta_T) \approx 1 - \Phi\left(\frac{\sqrt{n}}{\sigma_{\chi^2}^2} \left(\frac{q_{1-\alpha}}{2n} - \chi^2(\theta_0, \theta_T)\right)\right), \qquad (3.4)$$

where  $\Phi$  denotes, as usual, the distribution function of N(0,1) standard normal random variable. A useful consequence of (3.4) is the possibility of computing an approximate value of the sample size ensuring a specified power  $\beta(\theta_T)$ , with respect to some pre-assigned alternative  $\theta_T \neq \theta_0$ . Let  $n_0$  be the positive root of the equation

$$\beta = 1 - \Phi\left(\frac{\sqrt{n}}{\sigma_{\chi^2}^2} \left(\frac{q_{1-\alpha}}{2n} - \chi^2(\theta_0, \theta_T)\right)\right),\tag{3.5}$$

which can be rewritten into

$$n_0 = \frac{(a+b) - \sqrt{a(a+2b)}}{2\chi^2(\theta_0, \theta_T)},\tag{3.6}$$

where  $a := \sigma_{\chi^2}^2 (\Phi^{-1}(1-\beta))^2$  and  $b := q_{1-\alpha}\chi^2(\theta_0, \theta_T)$ . The sought-after approximate value of the sample size is then given

$$n^* := \lfloor n_0 \rfloor + 1,$$

where  $\lfloor u \rfloor$  denote the integer part of u.

**Remark 3.2** If the parameter  $\theta$  does not belong to  $\Theta$ , i.e.,  $\theta \in \Theta_e - \Theta$ , the densities  $c_{\theta}$ , with respect to the Lebesgue measure, associated to  $C_{\theta}$ , may become negative on some non negligible subsets of I. So, in this case

$$C_{\theta}(u_1, u_2) := \int_{(0, u_1) \times (0, u_2)} c_{\theta}(u_1, u_2) \ d\lambda(u_1, u_2); \ \forall u \in I,$$
(3.7)

may be a signed measure (and not a copula) for some  $\theta \in \Theta_e - \Theta$ . Hence, the formula (2.7) may lead to a value of the estimator  $\hat{\theta}_n$  belonging to  $\Theta_e - \Theta$ , and the corresponding  $C_{\hat{\theta}_n}$  is not necessarily a copula. So, for point estimation, the pseudo-maximum likelihood estimator (restricted to vary in the admissible domain) should be used instead of (2.7). The latter estimator may not be meaningful, and is likely to have a larger mean square error. The main advantage of our formula (2.6), where the supremum is taken on the extended space  $\Theta_e$ , instead of the admissible domain  $\Theta$ , is that it permits easily to build a test of independence even when the dimension of the parameter space  $\Theta$  is larger than 1. We can show that the proposed test statistic, based on the formula (2.6), has a  $\chi^2(p)$  limit law with  $p = \dim(\Theta)$ degrees of freedom.

**Remark 3.3** The asymptotic variances (2.10) and (3.2) may be consistently estimated respectively by the sample variances of

$$\frac{\partial}{\partial \theta} m\left(\widehat{\theta}_n, F_{1n}(X_{1,k}), F_{2n}(X_{2,k})\right) + W_1(\widehat{\theta}_n, X_{1,k}) + W_2(\widehat{\theta}_n, X_{2,k}), \quad k = 1, \dots, n,$$
(3.8)

$$\left(\widehat{\theta}_{n}, F_{1n}(X_{1,k}), F_{2n}(X_{2,k})\right) + Y_{1}(\widehat{\theta}_{n}, X_{1,k}) + Y_{2}(\widehat{\theta}_{n}, X_{2,k}), \quad k = 1, \dots, n,$$
(3.9)

as was done in Genest et al. (1995). Similarly, the parameter  $\Xi$  in (2.9) may be consistently estimated by the sample mean of

$$\left[\frac{\partial}{\partial \theta} m\left(\widehat{\theta}_n, F_{1n}(X_{1,k}), F_{2n}(X_{2,k})\right)\right]^2, \quad k = 1, \dots, n.$$
(3.10)

#### 4 Simulation results

In this section, we present some simulation results aiming to illustrate the theoretical results of 3.1. These results relate to the accuracy of the approximations of the laws of the test statistics  $\mathbf{T}_n$ , by the limit law given in Theorem 3.1. As an example, we consider the Clayton family of copulas

$$C_{\theta}(u_1, u_2) := \left(u_1^{-\theta} - u_2^{-\theta} - 1\right)^{-1/\theta}; \quad \text{for} \quad \theta \in [0, \infty).$$
(4.1)

The case of the test statistic  $\mathbf{T}_n$  based on the  $\chi^2$ -divergence. Under the null hypothesis of independence of margins (here  $\theta_T = \theta_0 := 0$ ), Theorem 3.1 shows that the asymptotic law of the proposed test statistic  $\mathbf{T}_n$  is a  $\chi_1^2$  law with one degree of freedom; see, e.g., Theorem 3.1. We have illustrated this convergence by simulation: in Figure 1 and Figure 2 are plotted the distribution function of the limit law  $\chi_1^2$ , and the distribution functions of the statistic  $\mathbf{T}_n$ obtained from 500 independent runs for sample sizes n = 25 and n = 100. We observe that the approximation is also good even for moderate sample sizes (see Figure 1 and Figure 2). The integral in the expression of  $m(\theta, \cdot, \cdot)$  is calculated by the Monte Carlo method, and the supremum in (3.1) is considered on the extended space  $\Theta_e$ ; the supremum in (3.1) has been computed on a neighborhood of  $\theta_0 := 0$  by the Newton-Raphson algorithm taking  $\theta_0 := 0$ as an initial point.

#### 5 Conclusion

We have presented a new approach based on  $\chi^2$ -divergence and duality technique in the framework of parametric copulas. It seems that the procedure introduced here is particularly well adapted to the boundary problem.



Figure 1: Simulated distribution function of the statistic  $\mathbf{T}_n$  and its limit law  $\chi_1^2$ . (Sample size 25, 500 replications.)



Figure 2: Simulated distribution function of the statistic  $\mathbf{T}_n$  and its limit law  $\chi_1^2$ . (Sample size 100, 500 replications.)

### 6 Appendix

First we give a technical Lemma which we will use to prove our results.

**Lemma 6.1** Let F have a continuous margins and let C have continuous partial derivatives. Assume that j is a continuous function, with bounded variation on I. Then

$$\int_{I} j(u_1, u_2) \ d\left(C_n(u_1, u_2) - C(u_1, u_2)\right) = O\left(n^{-1/2} (\log \log n)^{1/2}\right) \quad (a.s.).$$
(6.1)

**Proof of Lemma 6.1.** Recall that the *modified empirical copula*  $C_n$ , is slightly different from the *empirical copula*  $\mathbb{C}_n(u_1, u_2)$ , introduced by Deheuvels (1979a), and defined by

$$\mathbb{C}_n(u_1, u_2) = F_n\left(F_{1n}^{-1}(u_1), F_{2n}^{-1}(u_2)\right) \quad \text{for} \quad (u_1, u_2) \in (0, 1)^2, \tag{6.2}$$

where  $F_{1n}^{-1}(\cdot)$  and  $F_{2n}^{-1}(\cdot)$  denote the empirical quantile functions, associated with  $F_{1n}(x_1) = F_n(x_1, \infty)$  and  $F_{2n}(x_2) = F_n(\infty, x_2)$ , respectively, and defined by

$$F_{jn}^{-1}(t) := \inf\{x \in \mathbb{R} \mid F_{jn}(x) \ge t\}, \quad j = 1, 2.$$
(6.3)

Here,  $F_n(\cdot, \cdot)$  denotes the joint empirical distribution function, associated with the sample  $\{(X_{1k}, X_{2k}); k = 1, ..., n\}$ , defined by

$$F_n(x_1, x_2) = \frac{1}{n} \sum_{k=1}^n \mathbb{1}_{\{X_{1k} \le x_1\}} \mathbb{1}_{\{X_{2k} \le x_2\}}, \quad -\infty < x_1, x_2 < \infty.$$
(6.4)

We know that  $\mathbb{C}_n$  and  $C_n$  coincide on the grid  $\{(i/n, j/n), 1 \leq i \leq j \leq n\}$ . The subtle difference lies in the fact that  $\mathbb{C}_n$  is left-continuous with right-hand limits, whereas  $C_n$  on the other hand is right continuous with left-hand limits. The difference between  $\mathbb{C}_n$  and  $C_n$ , however, is small

$$\sup_{\mathbf{u}\in I} |\mathbb{C}_n(\mathbf{u}) - C_n(\mathbf{u})| \le \max_{1\le i,j\le n} \left| \mathbb{C}_n\left(\frac{i}{n},\frac{j}{n}\right) - \mathbb{C}_n\left(\frac{i-1}{n},\frac{j-1}{n}\right) \right| \le \frac{2}{n}.$$
 (6.5)

Using integration by parts, as in Fermanian et al. (2004a), we see that

$$\begin{split} \sqrt{n} \int_{I} \jmath(u_{1}, u_{2}) \ d(C_{n} - C)(u_{1}, u_{2}) &= \int_{I} \sqrt{n}(C_{n} - C)(u_{1}, u_{2}) \ d\jmath(u_{1}, u_{2}) \\ &- \int_{I} \sqrt{n}(C_{n} - C)(u_{1}, 1) \ d\jmath(u_{1}, u_{2}) - \int_{I} \sqrt{n}(C_{n} - C)(1, u_{2}) \ d\jmath(u_{1}, u_{2}) \\ &- \int_{[0,1]} \sqrt{n}(C_{n}(u_{1}, 1) - u_{1}) \ d\jmath(u_{1}, 1) - \int_{[0,1]} \sqrt{n}(C_{n}(1, u_{2}) - u_{2}) \ d\jmath(1, u_{2}). \end{split}$$

Hence,

$$\left|\sqrt{n} \int_{I} j(u_1, u_2) \ d(C_n - C)(u_1, u_2)\right| \le 5\sqrt{n} \sup_{\mathbf{u} \in I} |(C_n - C)(\mathbf{u})| \int_{I} d|j(\mathbf{u})|.$$

From this and (6.5), applying Theorem 3.1 in Deheuvels (1979a), we obtain the following result

$$\int_{I} j(u_1, u_2) \ d(C_n - C)(u_1, u_2) = O\left(n^{-1/2} (\log \log n)^{1/2}\right) \quad (a.s.).$$

**Proof of Theorem 2.1.** (a) Under the assumption (C.1), a straightforward calculus yields

$$\int_{I} \frac{\partial}{\partial \theta} m(\theta_T, u_1, u_2) \ dC_{\theta_T}(u_1, u_2) = 0.$$
(6.6)

Under the assumptions (C.1), and by applying Proposition 2.2 in Genest *et al.* (1995), we can see that, as  $n \to \infty$ ,

$$\int_{I} \frac{\partial}{\partial \theta} m(\theta_T, u_1, u_2) \ dC_n(u_1, u_2) \to \int_{I} \frac{\partial}{\partial \theta} m(\theta_T, u_1, u_2) \ dC_{\theta_T}(u_1, u_2) = 0, \tag{6.7}$$

and

$$\int_{I} \frac{\partial^2}{\partial \theta^2} m(\theta_T, u_1, u_2) \ dC_n(u_1, u_2) \to \int_{I} \frac{\partial^2}{\partial \theta^2} m(\theta_T, u_1, u_2) \ dC_{\theta_T}(u_1, u_2) := -\Xi < 0, \tag{6.8}$$

almost surely. Now, for any  $\theta = \theta_T + v n^{-1/3}$ , with  $|v| \leq 1$ , consider a Taylor expansion of  $\int m(\theta, u_1, u_2) \, dC_n(u_1, u_2)$  in  $\theta$  in a neighborhood of  $\theta_T$ , using (C.1) part 2, one finds

$$n \int_{I} m(\theta, u_{1}, u_{2}) \ dC_{n}(u_{1}, u_{2}) - n \int_{I} m(\theta_{T}, u_{1}, u_{2}) \ dC_{n}(u_{1}, u_{2})$$

$$= n^{2/3} v \int_{I} \frac{\partial}{\partial \theta} m(\theta_{T}, u_{1}, u_{2}) \ dC_{n}(u_{1}, u_{2}) + n^{1/3} \frac{v^{2}}{2} \int_{I} \frac{\partial^{2}}{\partial \theta^{2}} m(\theta_{T}, u_{1}, u_{2}) \ dC_{n}(u_{1}, u_{2}) + O(1) \quad (a.s.),$$
(6.9)

uniformly in v with  $|v| \leq 1$ . On the other hand, under condition (C.2), by Lemma 6.1, we have

$$\int_{I} \frac{\partial}{\partial \theta} m(\theta_T, u_1, u_2) dC_n(u_1, u_2) = O(n^{-1/2} (\log \log n)^{1/2}) \quad (a.s.)$$

Therefore, using (6.8) and (6.9), we obtain for any  $\theta = \theta_T + v n^{-1/3}$  with |v| = 1,

$$n \int_{I} m(\theta, u_1, u_2) \ dC_n(u_1, u_2) - n \int_{I} m(\theta_T, u_1, u_2) \ dC_n(u_1, u_2)$$
  
=  $O(n^{1/6} (\log \log n)^{1/2}) - 2^{-1} \Xi n^{1/3} + O(1) \ (a.s.).$  (6.10)

Observe that the right-hand side vanishes when  $\theta = \theta_T$ , and that the left-hand side, by (6.8), becomes negative for all *n* sufficiently large. Thus, by the continuity of  $\theta \mapsto \int m(\theta, u_1, u_2) \, dC_n(u_1, u_2)$ , it holds that as  $n \to \infty$ , with probability one,

$$\theta \mapsto \int m(\theta, u_1, u_2) \ dC_n(u_1, u_2)$$

reaches its maximum value at some point  $\hat{\theta}_n$  in the interior of the interval *B*. Therefore, the estimate  $\hat{\theta}_n$  satisfies

$$\int_{I} \frac{\partial}{\partial \theta} m(\widehat{\theta}_n, u_1, u_2) \ dC_n(u_1, u_2) = 0 \text{ and } | \widehat{\theta}_n - \theta_T | = O(n^{-1/3}).$$
(6.11)

(b) Making use of the first part of Theorem 2.1, and once more, by a Taylor expansion of

$$\int_{I} \frac{\partial}{\partial \theta} m(\widehat{\theta}_n, u_1, u_2) \ dC_n(u_1, u_2),$$

with respect to  $\hat{\theta}_n$ , in the neighborhood of  $\theta_T$ , we obtain that

$$0 = \int_{I} \frac{\partial}{\partial \theta} m(\widehat{\theta}_{n}, u_{1}, u_{2}) \ dC_{n}(u_{1}, u_{2}) = \int_{I} \frac{\partial}{\partial \theta} m(\theta_{T}, u_{1}, u_{2}) \ dC_{n}(u_{1}, u_{2})$$
$$+ (\widehat{\theta}_{n} - \theta_{T}) \int_{I} \frac{\partial^{2}}{\partial \theta^{2}} m(\theta_{T}, u_{1}, u_{2}) \ dC_{n}(u_{1}, u_{2}) + o(n^{-1/2}).$$

Hence,

$$\sqrt{n}(\widehat{\theta}_n - \theta_T) = \left(-\int_I \frac{\partial^2}{\partial \theta^2} m(\theta_T, u_1, u_2) \ dC_n(u_1, u_2)\right)^{-1} \sqrt{n} W_n(\theta_T) + o(1), \tag{6.12}$$

where

$$W_n(\theta_T) := \int_I \frac{\partial}{\partial \theta} m(\theta_T, u_1, u_2) \ dC_n(u_1, u_2).$$

Applying Proposition 3 page 362 in Tsukahara (2005), under assumptions (C.1) part 1 and (C.3), we have

$$\sqrt{n}W_n(\theta_T) \xrightarrow{d} N(0, \Sigma^2).$$
 (6.13)

Finally, by combining (6.13) and (6.8) in connection with Slutsky's Theorem, we conclude that

$$\sqrt{n}(\widehat{\theta}_n - \theta_T) \to N(0, \Sigma^2 / \Xi^2).$$
 (6.14)

This completes the proof.

**Proof of Theorem 3.1.** Assume that  $\theta_T = \theta_0$ . From (6.12), using (6.8) and (6.13), we obtain

$$\sqrt{n}\left(\widehat{\theta}_n - \theta_T\right) = -\frac{1}{\Sigma}\sqrt{n}W_n(\theta_T) + o_P(1).$$
(6.15)

Expanding in Taylor series  $\mathbf{T}_n(\theta_0, \widehat{\theta}_n)$  in  $\widehat{\theta}_n$  around  $\theta_T$ , we get

$$\mathbf{T}_{n}(\theta_{0},\widehat{\theta}_{n}) = 2nW_{n}(\theta_{T})(\widehat{\theta}_{n}-\theta_{T}) - \Sigma n(\widehat{\theta}_{n}-\theta_{T})^{2} + o_{P}(1).$$
(6.16)

Now, use (6.15) combined with (6.16) to obtain

$$\mathbf{T}_n(\theta_0, \widehat{\theta}_n) = \frac{1}{\Sigma} n W_n(\theta_T)^2 + o_P(1).$$
(6.17)

By (6.13), we have

$$\sqrt{n}W_n(\theta_T) \to N\left(0, \Sigma^2\right) \tag{6.18}$$

in distribution. When  $\theta_T = \theta_0$ , under Assumption (C.4), we can see that  $\Sigma^2$  in (6.18) is equal to  $1/\Xi$ ; see Proposition 2.2 in Genest *et al.* (1995). Combining this with (6.17) to conclude that

$$\mathbf{T}_n \xrightarrow{d} \chi_1^2,$$

under the null hypothesis  $\mathcal{H}_0$ .

**Proof of Theorem 3.2.** Rewriting  $\frac{\mathbf{T}_n}{2n}$  as

$$\frac{\mathbf{T}_n(\theta_0, \widehat{\theta}_n)}{2n} = \sup_{\theta \in \Theta_e} \int_I m(\theta, u_1, u_2) \ dC_n(u_1, u_2), \tag{6.19}$$

and making use of a Taylor expansion of  $\frac{\mathbf{T}_n(\theta_0, \hat{\theta}_n)}{2n}$ , with respect to  $\hat{\theta}_n$ , in a neighborhood of  $\theta_T$ , under (C.5) to obtain

$$\frac{\mathbf{T}_n(\theta_0, \widehat{\theta}_n)}{2n} = \int_I m(\theta_T, u_1, u_2) \ dC_n(u_1, u_2) + o_P(n^{-1/2}).$$

Hence, one finds

$$\sqrt{n} \left( \frac{\mathbf{T}_n}{2n} - \chi^2(\theta_0, \theta_T) \right) = \sqrt{n} \left( \int_I m(\theta_T, u_1, u_2) \ dC_n - \int_I m(\theta_T, u_1, u_2) \ dC_{\theta_T} \right) \\
+ o_P(1).$$

Finally, under condition (C.6), application once more of Proposition 3 page 362 in Tsukahara (2005), concludes the proof.

#### References

- Bouzebda, S. and Keziou, A. (2007). A test of independence in some copula models. *Submitted*.
- Broniatowski, M. and Keziou, A. (2006). Minimization of φ-divergences on sets of signed measures. Studia Sci. Math. Hungar., 43(4), 403–442.
- Deheuvels, P. (1979a). La fonction de dépendance empirique et ses propriétés. Un test non paramétrique d'indépendance. Acad. Roy. Belg. Bull. Cl. Sci. (5), 65(6), 274–292.
- Deheuvels, P. (1979b). Propriétés d'existence et propriétés topologiques des fonctions de dépendance avec applications à la convergence des types pour des lois multivariées. C. R. Acad. Sci. Paris Sér. A-B, 288(2), A145–A148.
- Deheuvels, P. (1980). Nonparametric test of independence. In Nonparametric asymptotic statistics (Proc. Conf., Rouen, 1979) (French), volume 821 of Lecture Notes in Math., pages 95–107. Springer, Berlin.
- Deheuvels, P. (1981). A Kolmogorov-Smirnov type test for independence and multivariate samples. *Rev. Roumaine Math. Pures Appl.*, 26(2), 213–226.
- Fermanian, J.-D., Radulović, D., and Wegkamp, M. (2004a). Weak convergence of empirical copula processes. *Bernoulli*, **10**(5), 847–860.
- Fermanian, J.-D., Radulović, D., and Wegkamp, M. (2004b). Weak convergence of empirical copula processes. http://www.crest.fr/pageperso/fermanian/fermanian.htm.
- Galambos, J. (1975). Order statistics of samples from multivariate distributions. J. Amer. Statist. Assoc., 70(351, part 1), 674–680.
- Genest, C., Ghoudi, K., and Rivest, L.-P. (1995). A semiparametric estimation procedure of dependence parameters in multivariate families of distributions. *Biometrika*, 82(3), 543–552.

- Gumbel, E. J. (1960). Bivariate exponential distributions. J. Amer. Statist. Assoc., 55, 698–707.
- Hüsler, J. and Reiss, R.-D. (1989). Maxima of normal random vectors: between independence and complete dependence. *Statist. Probab. Lett.*, 7(4), 283–286.
- Joe, H. (1993). Parametric families of multivariate distributions with given margins. J. Multivariate Anal., 46(2), 262–282.
- Joe, H. (1997). Multivariate models and dependence concepts, volume 73 of Monographs on Statistics and Applied Probability. Chapman & Hall, London.
- Keziou, A. and Leoni-Aubin, S. (2005). Test of homogeneity in semiparametric two-sample density ratio models. C. R. Math. Acad. Sci. Paris, 340(12), 905–910.
- Keziou, A. and Leoni-Aubin, S. (2007). On empirical likelihood for semiparametric twosample density ratio models. *Journal of Statistical Planning and Inference; In press.*
- Kimeldorf, G. and Sampson, A. (1975a). One-parameter families of bivariate distributions with fixed marginals. *Comm. Statist.*, 4, 293–301.
- Kimeldorf, G. and Sampson, A. (1975b). Uniform representations of bivariate distributions. Comm. Statist., 4(7), 617–627.
- Oakes, D. (1994). Multivariate survival distributions. J. Nonparametr. Statist., **3**(3-4), 343–354.
- Qin, J. and Lawless, J. (1994). Empirical likelihood and general estimating equations. Ann. Statist., 22(1), 300–325.
- Rüschendorf, L. (1976). Asymptotic distributions of multivariate rank order statistics. Ann. Statist., 4(5), 912–923.
- Ruymgaart, F. H. (1974). Asymptotic normality of nonparametric tests for independence. Ann. Statist., 2, 892–910.

- Ruymgaart, F. H., Shorack, G. R., and van Zwet, W. R. (1972). Asymptotic normality of nonparametric tests for independence. Ann. Math. Statist., 43, 1122–1135.
- Self, S. G. and Liang, K.-Y. (1987). Asymptotic properties of maximum likelihood estimators and likelihood ratio tests under nonstandard conditions. J. Amer. Statist. Assoc., 82(398), 605–610.
- Shih, J. H. and Louis, T. A. (1995). Inferences on the association parameter in copula models for bivariate survival data. *Biometrics*, 51(4), 1384–1399.
- Sklar, M. (1959). Fonctions de répartition à n dimensions et leurs marges. Publ. Inst. Statist. Univ. Paris, 8, 229–231.
- Tsukahara, H. (2005). Semiparametric estimation in copula models. *Canad. J. Statist.*, **33**(3), 357–375.
- Wang, W. and Ding, A. A. (2000). On assessing the association for bivariate current status data. *Biometrika*, 87(4), 879–893.