



HAL
open science

$^{13}\text{C}/^{12}\text{C}$ composition, a novel parameter to study the downward migration of paper sludge in soils

Eric Lichtfouse, Karyne Rogers, Cécile Payet, Jean-Christophe Renat

► To cite this version:

Eric Lichtfouse, Karyne Rogers, Cécile Payet, Jean-Christophe Renat. $^{13}\text{C}/^{12}\text{C}$ composition, a novel parameter to study the downward migration of paper sludge in soils. *Geochemical Transactions*, 2002, 3, pp.48-50. 10.1186/1467-4866-3-48 . hal-00263862

HAL Id: hal-00263862

<https://hal.science/hal-00263862>

Submitted on 13 Mar 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

$^{13}\text{C}/^{12}\text{C}$ composition, a novel parameter to study the downward migration of paper sludge in soils

Eric Lichtfouse,^{*a} Karyne Rogers,^b Cécile Payet^c and Jean-Christophe Renat^c

^a *GéoSol-CST, Earth Sciences, University of Burgundy, 6, Bd Gabriel, 21000 Dijon, France;*

E-mail: Eric.Lichtfouse@u-bourgogne.fr

^b *Stable Isotope Laboratory, Institute of Geological and Nuclear Sciences, PO Box 31-312, Lower Hutt, New Zealand; E-mail: K.Rogers@gns.cri.nz*

^c *Traitement Valorisation Décontamination Ltd, 3, rue du Coteau, F-54180 Heillecourt, France.*

Abstract. $\delta^{13}\text{C}$ values of crop and forest soils were measured 8 years after disposal of paper sewage sludge. The carbon transfer from paper sludge downward to the first humic layer is evidenced by a ^{13}C -enrichment of up to + 5.6‰ due to the input of ^{13}C -enriched sludge carbonates. $^{13}\text{C}/^{12}\text{C}$ composition is thus a novel, sensitive parameter to follow to downward transfer of paper sludge carbon.

Introduction

Large amounts of agricultural, industrial and municipal wastes are produced daily by human activities¹⁻⁵. In 1980, France's annual waste production reached about 5.5 million dry tons (m.d.t.) of municipal waste, 1.8 m.d.t. of urban and industrial sewage sludge, 78 m.d.t. from the agriculture and agro-industry, and 6.75 m.d.t. from forestry⁶. Disposal of organic wastes onto agricultural and forestry lands has several potential benefits such as long-term fertilisation, improving soil water-holding capacity and improvement of aggregate stability. However, land-based waste disposal must be carefully controlled because of potential hazards associated with application of wastes, include pathogens, heavy metals and toxic organic by-products, as reviewed by Wilson et al.². So far, the long-term changes of soil properties induced by organic waste disposal such as paper sludge are not well understood, notably due to the lack of analytical approaches to follow the fate of waste matter into the soil profile. Nonetheless, several recent reports show that stable carbon isotopes can be used to study environmental issues⁷⁻⁹. More specifically, since the main biochemical components of plants are isotopically distinguished, e.g. cellulose being ^{13}C -enriched versus lignin and lipids¹⁰⁻¹², we hypothesised that paper sludge might have a distinct $\delta^{13}\text{C}$ ratio which could be used to study their long-term fate in soils. Moreover, since paper sludges

contain carbonates, which are ^{13}C -enriched, it could be feasible to isotopically distinguish soil carbon from sludge carbon. Here, we wish to report an isotopic investigation of crop and forest soils treated with paper sludge in 1992.




Figure 1 Sampling of soil layers from the wood plot in August 2000. A layer of paper sludge has been disposed of to this soil in 1992.

Table 1 $\delta^{13}\text{C}$ values and %carbon of non-demineralised soil samples cored in August 2000 from experimental control sites and from sites treated with paper sludge in 1992. Light-blue chunks of solidified paper sludge found on the soil surface yielded a %C value of 17.36% and $\delta^{13}\text{C}$ value of -16.84‰. Sample deviation: $\pm 0.05\%$ and $\pm 0.05\text{‰}$ (3 repeats).

HORIZON, thickness	WOOD SOILS				CROP SOILS			
	C (%)		$\delta^{13}\text{C}$ (‰)		C (%)		$\delta^{13}\text{C}$ (‰)	
	Control site	Paper site	Control site	Paper site	Control site	Paper site	Control site	Paper site
Litter ~ 3 cm, grasses ¹	40.24	44.94	-27.63	-27.24	44.25	43.01	-28.20	-27.58
Blue sludge ² , 10 cm		18.28		-20.97		5.82		-22.81
Black humic ³ , 10 cm	4.80	4.34	-27.66	-25.72	2.50	5.20	-27.48	-21.84
Dark-brown ³ , 20 cm	2.26	2.30	-27.79	-27.63	1.04	0.56	-28.16	-28.70
Light-brown ³ , 20 cm	1.03	1.03	-27.35	-27.34	0.55	0.31	-27.72	-27.81

¹Woodland sites: litter, mostly fern debris. Crop site: living grasses. ²Horizon visibly composed of about half black soil and half 2 mm-blue chunks.

³Horizons without visible blue chunks.

Soil sampling and isotope analysis

In August 2000, solid chunks of blue paper from the surface, organic litter from woodland sites, grasses from crop sites, and soils of increasing depth were sampled from ~ 1 m³ holes cored at the four sites, as shown on Figure 1 for the woodland site. Samples were air-dried at 20°C, 2 mm-sieved (soils), then finely ground using a ball mortar. Total %C and $\delta^{13}\text{C}$ values (‰) were measured by continuous flow isotopic ratio mass spectrometry as described elsewhere¹³⁻¹⁵. Note that %C and $\delta^{13}\text{C}$ values refer to total soil carbon, including organic and inorganic (carbonates) carbon. TOC and $\delta^{13}\text{C}_{(\text{org})}$ values of all demineralised samples were also measured and show that the total %C and $\delta^{13}\text{C}$ have notable contribution of carbonates, as detailed at the end of this article. Demineralisation was performed by overnight treatment of the sample in 1N HCl, followed by washing with distilled water and centrifuging the samples until neutral. The samples were air-dried and treated in the same manner as above.

the total C content of litter in the woodland sludge-treated site (45%) is higher than the woodland control site (40%), the reverse is observed for the crop sites: 43 versus 44%. Similarly, the black humic horizon underlying the sludge give higher total C values in the crop sludge-treated site, 5.2 versus 2.5%, but lower C values in the woodland sludge-treated site amounting to 4.3 versus 4.8%. Therefore, although one should expect the upward and downward transfer of some carbon from the blue sludge layer 8 years after disposal, carbon values do not show unambiguous differences. Nonetheless, the next section will show that the migration of sludge carbon can be clearly evidenced by isotope analysis.

Results and discussion

Paper sludge disposal

In 1992, crop and woodland sites from the Lorraine region, France were treated with 186-306 tons of paper sewage sludge in order to study the effects of waste recycling. Precautions were taken to minimise potential environmental hazards, e.g. input of heavy metals. From 1992 to 1997, comparison of plants grown on both the treated and control sites showed the absence of visual toxic effects. Plants developed well with roots growing through the blue sludge layer. An investigation of the blue sludge layer from 1992 to 1997 showed a decrease of calcium content, from about 23 to 10%, and of organic matter content, from about 35 to 20%. In 2000, the blue sludge layer is still clearly apparent under a fern litter layer, as shown for the woodland site on Figure 1. Here, we analysed samples of litter, grasses, sludge layer, and soil layers of increasing depth cored in July 2000, in order to study the downward carbon transfer from the paper sludge.

Carbon content

Total carbon content and $\delta^{13}\text{C}$ values of samples from woodland and crop sites treated with paper sludge are reported on Table 1. All sites show a decrease of total C content with depth from ~ 42% for litter and grasses, to 0.5-1% at the bottom of the core. Although the blue sludge layers yielded high carbon contents, 18.2% for the woodland site and 5.8% for the crop site respectively, carbon contents do not clearly show the transfer of paper sludge-derived carbon to other layers. Specifically, while

¹³C/¹²C isotopic composition

$\delta^{13}\text{C}$ values of samples from sites treated with paper sludge and from control sites are drawn on Figure 2. We observe a notable ¹³C-enrichment in the blue sludge layer for both sites, yielding $\delta^{13}\text{C}$ values of -20.97‰ for the woodland site and -22.81‰ for the crop site, and in the underlying black humic layer (-25.72, -21.84‰ respectively), relative to the average $\delta^{13}\text{C}$ values from control plots amounting to $-27.6 \pm 0.2\text{‰}$ for the woodland site and to $-27.9 \pm 0.3\text{‰}$ from the crop site. Moreover, the soil $\delta^{13}\text{C}$ values of sludge-treated sites increase toward the original isotopic value of the paper sludge (-16.84‰) with decreasing depth. These findings have several implications. First, the blue

$$\delta_{\text{layer}} = x \cdot \delta_{\text{sludge}} + (1-x) \cdot \delta_{\text{control}}$$

sludge layer is composed of a mixture of carbon derived from the $\delta^{13}\text{C}$ -enriched paper sludge and from the soil, in agreement with visual observation of both blue and dark particles in the blue sludge layer. Second, the notable ¹³C-enrichment of the underlying black humic horizons shows clearly the downward migration of sludge-derived carbon where other data such as %C contents and visual observation do not yield clear trends. Third, the fraction x of paper sludge-derived carbon can be calculated by isotope balance according to the following equation:

where δ_{layer} refer to the soil layer, δ_{sludge} to solid chunks of pure paper sludge (-16.84‰), and δ_{control} to average $\delta^{13}\text{C}$ values of control plots. In the woodland sites, the percentage x.100 of sludge-derived carbon amounts to 76% in the blue sludge layer and to 21% in the underlying black humic layer, thus showing a notable downward carbon transfer. In crop plots, values amount respectively to 56 and 67% as the result of a downward carbon transfer, which could be explained by the lesser initial stratification of crop soils.

Sludge carbonates

$\delta^{13}\text{C}_{(\text{org})}$ analysis of demineralised samples show that the ¹³C-enrichment of the non-demineralised samples is due to the presence of carbonates. Specifically, demineralised blue paper chunks yield $\delta^{13}\text{C}_{(\text{org})}$ values of -25.41‰ versus -16.84‰ for the non-demineralised sample. The blue sludge layers give $\delta^{13}\text{C}_{(\text{org})}$ values of -27.22‰ for the woodland site and -26.28‰ for the crop site, versus respectively -20.97‰ and -22.81‰ for the non-demineralised samples. Similarly, the black humic layers give $\delta^{13}\text{C}_{(\text{org})}$ of -26.97‰ for the woodland site and -26.27‰ for the crop site, versus -25.72‰ and -21.84‰ respectively for the non-demineralised samples. Since the demineralised values are

similar to control values (Table 1), the observed ¹³C-increases of non-demineralised layers can be explained by the total carbonate contribution from the sludge.

Conclusion

The downward transfer of paper sludge 8 years after disposal of to crop and woodland soils has been assessed using ¹³C isotope analyses. The observed isotopic shifts are due to the presence of enriched carbon, derived from carbonates in the paper sludge.

References

- 1 S. C. Wilson, V. Burnett, K. S. Waterhouse and K. C. Jones. Volatile organic compounds in digested United Kingdom sewage sludges. *Environmental Science and Technology*, 1994, **28**, 259.
- 2 S. C. Wilson, R. Duarte-Davidson and K. C. Jones. Screening the environmental fate of organic contaminants in sewage sludges applied to agricultural soils : 1. The potential for downward movement to groundwaters. *The Science of the Total Environment*, 1996, **185**, 45.
- 3 G. A. O'Connor. Organic compounds in sludge-amended soils and their potential for uptake by crop plants. *The Science of the Total Environment*, 1996, **185**, 71.
- 4 R. L. Chaney, J. A. Ryan and G. A. O'Connor. Organic contaminants in municipal biosolids: risk assessment, quantitative pathways analysis, and current research priorities. *The Science of the Total Environment*, 1996, **185**, 187.
- 5 C. Payet, C. Bryselbout, J. L. Morel and E. Lichtfouse. Fossil fuel biomarkers in sewage sludges: environmental significance. *Naturwissenschaften*, 1999, **86**, 484.
- 6 M. Mustin. *Le Compost*. 1987, F. Dubusc Publisher. Paris. 954 pages (in French).
- 7 B. Sherwood Lollar and T. A. Abrajano (Eds.). Compound-specific isotope analysis: tracing organic contaminant sources and processes in geochemical systems. *Organic Geochemistry*, 1999, **30**, 721.
- 8 W. Amelung, R. Bol and C. Friedrich. Natural ¹³C abundance: a tool to trace the incorporation of dung-derived carbon into soil particle-size fractions. *Rapid Communications in Mass Spectrometry*, 1999, **13**, 1291.
- 9 E. Lichtfouse. Compound-specific isotope analysis (CSIA). Application to archaeology, biomedical sciences, biosynthesis, environment, extraterrestrial chemistry, food science, forensic science, humic substances, microbiology, organic geochemistry, soil science and sport. *Rapid Communications in Mass Spectrometry*, 2000, **14**, 1337.
- 10 R. Park and S. Epstein. Metabolic fractionation of ¹³C and ¹²C in plants. *Plant Physiology*, 1961, **36**, 133.
- 11 R. Benner, M. L. Fogel, E. K. Sprague and R. E. Hodson. Depletion of ¹³C in lignin and its implications for stable isotope studies. *Nature*, 1987, **329**, 708.
- 12 M. O'Leary. Carbon isotope fractionation in plants. *Phytochemistry*,




Figure 2 ¹³C/¹²C isotopic composition measured in 2000 of woodland and crop soils treated with paper sludge in 1992, versus non-treated soils (control). Note the ¹³C-enrichment of the black humic layer is due to the downward carbon transfer from the paper sludge.

- 1981, **20**, 553.
- 13 E. Lichtfouse, S. Dou, C. Girardin, M. Grably, J. Balesdent, F. Behar, M. Vandenbroucke. Unexpected ¹³C-enrichment of organic components from wheat crop soils : evidence for the in situ origin of soil organic matter. *Organic Geochemistry*, 1995, **23**, 865.
 - 14 K. M. Rogers. Effects of sewage contamination on macro-algae and shellfish at Moa Point, New Zealand using stable carbon and nitrogen isotopes. *New Zealand Journal of Marine and Freshwater Research*, 1999, **33**, 181.
 - 15 K. M. Rogers, H. E. G. Morgans, G. S. Wilson. Identification of a Waipawa Formation equivalent in the Te Uri Member of the Whangai Formation – implications for depositional history and age. *New Zealand Journal of Geology and Geophysics*, 2001, **44**, 345.

