



**HAL**  
open science

# Geometric Interpretation of Second Elliptic Integrable System

Idrisse Khemar

► **To cite this version:**

| Idrisse Khemar. Geometric Interpretation of Second Elliptic Integrable System. 2008. hal-00266503v1

**HAL Id: hal-00266503**

**<https://hal.science/hal-00266503v1>**

Preprint submitted on 23 Mar 2008 (v1), last revised 8 Apr 2009 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Geometric Interpretation of Second Elliptic Integrable System

Idrisse Khemar

(joint work with Francis Burstall)

## Abstract.

In this paper we give a geometrical interpretation of all the second elliptic integrable systems associated to 4-symmetric spaces. We first show that a 4-symmetric space  $G/G_0$  can be embedded into the twistor space of the corresponding symmetric space  $G/H$ . Then we prove that the second elliptic system is equivalent to the vertical harmonicity of an admissible twistor lift  $J$  taking values in  $G/G_0 \hookrightarrow \Sigma(G/H)$ . We begin the paper by an example:  $G/H = \mathbb{R}^4$ . We study also the structure of 4-symmetric bundles over Riemannian symmetric space.

## Introduction

The first example of second elliptic integrable system associated to a 4-symmetric space was given in [12]: the authors showed that the Hamiltonian stationary Lagrangian surfaces in  $\mathbb{C}^2$  are solution of one such integrable system, and after they generalized their result to complex two-dimensional Hermitian symmetric space, [14]. Then we gave in [17] a new class of geometrical problems for surfaces in the Euclidean space of dimension 8 by using the identification  $\mathbb{R}^8 = \mathbb{O}$ , and proved that they are solution of a second elliptic integrable system. Using the left multiplication in  $\mathbb{O}$  by the vectors of the canonical basis of  $\text{Im } \mathbb{O}$  we defined a family  $\{\omega_i, 1 \leq i \leq 7\}$  of canonical symplectic forms in  $\mathbb{O}$ . This allowed us to define the notion of  $\omega_I$ -isotropic surfaces, for  $I \subsetneq \{1, \dots, 7\}$ . Using the cross-product in  $\mathbb{O}$  we defined a map  $\rho: Gr_2(\mathbb{O}) \rightarrow S^6$  from the Grassmannian of plan of  $\mathbb{O}$  to  $S^6$ . This allowed us to associate to each surface  $\Sigma$  of  $\mathbb{O}$  a function  $\rho_\Sigma: \Sigma \rightarrow S^6$ . In the case of  $\omega_I$ -isotropic surfaces,  $\rho_\Sigma$  takes values into a subsphere  $S^I = S(\oplus_{i \notin I, i > 0} \mathbb{R}e_i) \simeq S^{6-|I|}$ . Then we showed that the surfaces in  $\mathbb{O}$  such that  $\rho_\Sigma$  is harmonic ( $\rho$ -harmonic surfaces) are solutions of a completely integrable system  $\mathcal{S}$ . More generally we showed that the  $\omega_I$ -isotropic  $\rho$ -harmonic surfaces are solutions of a completely integrable system  $\mathcal{S}_I$ . Hence we built a family  $(\mathcal{S}_I)$  indexed by  $I$ , of set of surfaces solutions of a integrable system, all included in  $\mathcal{S} = \mathcal{S}_\emptyset$ , such that  $I \subset J$  implies  $\mathcal{S}_J \subset \mathcal{S}_I$ . Each  $\mathcal{S}_I$  is a second elliptic integrable system (in the sense of C.L. Terng). This means that

the equations of this system are equivalent to the free curvature equation :

$$d\alpha_\lambda + \frac{1}{2}[\alpha_\lambda \wedge \alpha_\lambda] = 0,$$

for all  $\lambda \in \mathbb{C}$ , and where  $\alpha_\lambda = \lambda^{-2}\alpha'_2 + \lambda^{-1}\alpha_{-1} + \alpha_0 + \lambda\alpha_1 + \lambda^2\alpha''_2$ .

By restriction to  $\mathbb{H} \subset \mathbb{O}$  of our theory we obtain a new class of surfaces: the  $\omega_I$ -isotropic  $\rho$ -harmonic surfaces of  $\mathbb{H}$ . Then  $\rho(Gr_2(\mathbb{H})) = S^2$  and  $|I| = 0, 1$  or  $2$ . For  $|I| = 1$  we obtain the Hamiltonian Stationary Lagrangian surfaces in  $\mathbb{R}^4$  and for  $|I| = 2$ , the special Lagrangian surfaces. By restriction to  $\text{Im } \mathbb{H}$ , we obtain the CMC surfaces of  $\mathbb{R}^3$ .

Besides in [18], we found a supersymmetric interpretation of all the second elliptic integrable system associated to a 4-symmetric space in terms of super harmonic maps into a symmetric space. This leads us to conjecture that this system has a geometrical interpretation in terms of surfaces with values in a symmetric space, such that a certain associated map is harmonic as this is the case for Hamiltonian stationary Lagrangian surfaces in Hermitian symmetric spaces or for  $\rho$ -harmonic surfaces of  $\mathbb{O}$ .

In this paper we give the answer to this conjecture. That is to say, we give a geometrical interpretation – in terms of vertical harmonic twistor lift – of all the second elliptic integrable systems associated to a 4-symmetric space. Indeed given a 4-symmetric space  $G/G_0$ , and its order four automorphism  $\sigma: G \rightarrow G$ , then the involution  $\tau = \sigma^2$  gives rise to the symmetric space  $G/H$ , with  $H = G^\tau$ . Then we prove that the second elliptic integrable system associated to the 4-symmetric space  $G/G_0$  is the equation of vertical harmonicity for admissible twistor lift in  $G/H$ . More precisely, given a 4-symmetric space  $G/G_0$ , and its associated symmetric space  $G/H$ , then  $G/G_0$  is a subbundle of the twistor space  $\Sigma(G/H)$ . We prove that the second elliptic integrable systems associated to  $G/G_0$ , is the system of equations for maps  $J: \mathbb{C} \rightarrow G/G_0 \subset \Sigma(G/H)$  such that  $J$  is compatible with the Gauss map of  $X: \mathbb{C} \rightarrow G/H$ , the projection of  $J$  into  $G/H$ , i.e.  $X$  is  $J$ -holomorphic (admissible twistor lift), and such that  $J$  is vertically harmonic.

We begin the paper by an example:  $\mathbb{R}^4$ . This case was just mentioned quickly in the end of [17] as a restriction of the difficult problem in  $\mathbb{O}$ . So here we study it independently and in details. However, here we give also a formulation of this problem in terms of twistor lift which is the right formulation. Besides, in dimension 4 we have unicity of the twistor lift (in  $\Sigma^+(G/H)$  and  $\Sigma^-(G/H)$  respectively) so we are in this case in the presence of a theory of surface (and not as in the general case, a theory of twistor lift) and so we can speak about  $\rho$ -harmonic surfaces in this dimension (which are exactly the solution of the second elliptic integrable system). In our work we are led to prove some theorems on the structure of 4-symmetric bundles. Indeed we want to answer the following questions: given a Riemannian symmetric spaces, do there exist 4-symmetric bundle over it? In other words in its twistor bundle do there exist 4-symmetric subbundle, and if yes how can we characterize these 4-symmetric components? are they isomorphic? How are they distributed in the twistor space? Do they form a partition of the twistor space?. etc.. The 4-symmetric

space have been classified (at least in the compact, see [16, 21]). However, our point of view is different: we want to have an intrinsic point of view as long as possible so we deal with the Riemannian symmetric space and its (locally) 4-symmetric bundle defined over it, and we try to forget as much as possible the four automorphism of the Lie algebra. Our aim is to give a formulation of our problem as general and intrinsic as possible. For example, our definition of vertical harmonicity holds for any Riemannian manifold, moreover we give the following characterization: define a (locally) 4-symmetric bundle over  $M$  is equivalent to give oneself  $J_0 \in \Sigma(T_{p_0}M)$  which leaves invariant the curvature. We obtain the following image: the twistor bundle in the disjoint union of all the maximal (locally) 4-symmetric subbundle, which are orbits (defined by some subgroups of  $\text{Is}(M)$ ). Each isomorphism class of orbits defined a different second elliptic integrable system.

Our paper is organized as follows. In Section 1 we deals with the  $\rho$ -harmonic surfaces in  $\mathbb{R}^4$ . Section 2 contains our main result: the interpretation of the second elliptic integrable systems associated to a 4-symmetric space in terms of the vertical harmonicity of an admissible twistor lift. Then Section 3 and 4 are devoted to the study of the structure of 4-symmetric bundles over symmetric spaces. The last Section gives some example of 4-symmetric bundles.

## 1 $\rho$ -harmonic surfaces in $\mathbb{H}$

### 1.1 Cross product, complex structure and Grassmannian of plan in $\mathbb{H}$

We consider the space  $\mathbb{R}^4 = \mathbb{H}$  with its canonical basis  $(1, i, j, k)$  (which we denote also by  $(e_i)_{0 \leq i \leq 3}$ ). Let  $P = q \wedge q'$  be a oriented plan of  $\mathbb{H}$  (itself oriented by its canonical basis) then there exists an unique positive complex structure  $I_P \in \Sigma^+(P)$  on the plan  $P$ . It is defined by  $I_P(q) = q'$ ,  $I_P(q') = -q$  if  $(q, q')$  is orthogonal. Next, we can extend it in a unique way to a positive (resp. negative) complex structure of  $\mathbb{H} = P \oplus P^\perp$ ,  $J_P^+$  (resp.  $J_P^-$ ) given by

$$\begin{aligned} J_P^+ &= I_P \oplus I_{P^\perp} \\ J_P^- &= I_P \oplus -I_{P^\perp} \end{aligned} \tag{1}$$

( $P^\perp$  is oriented so that  $P \oplus P^\perp$  is positively oriented). Hence we obtain a surjective map:

$$\begin{aligned} J^+ : Gr_2(\mathbb{H}) &\rightarrow \Sigma^+(\mathbb{H}) \\ q \wedge q' &\mapsto J_{q \wedge q'}^+ \end{aligned} \tag{2}$$

and in the same way a surjective map  $J^- : Gr_2(\mathbb{H}) \rightarrow \Sigma^-(\mathbb{H})$ .

Besides, we have

$$J_{q \wedge q'}^+ = L_{q \times_L q'} = \frac{1}{2}(L_{q'}L_{\bar{q}} - L_qL_{\bar{q}'}),$$

where  $q \times_L q' = -\text{Im}(q \cdot \bar{q}') = \text{Im}(q' \cdot \bar{q})$  is the left cross product (it is a bilinear skew map from  $\mathbb{H} \times \mathbb{H}$  to  $\text{Im } \mathbb{H}$ ). Indeed, if  $(q, q')$  is orthonormal then  $q \times_L q' =$

$-q \cdot \bar{q}' \in S(\text{Im } \mathbb{H})$  so  $L_{q \times_L q'}$  is a complex structure of  $\mathbb{H}$  and it is positive (because  $\{L_u, u \in S^2\}$  is connected and  $L_i \in \Sigma^+(\mathbb{H})$  because  $(1, L_i(1), j, L_i(j))$  is positively oriented). Moreover if  $(q, q')$  is orthonormal then  $L_{q \times_L q'}(q) = (q' \bar{q})q = q'$ . Hence  $L_{q \times_L q'} = J_{q \wedge q'}^+$ . Thus we obtain a diffeomorphism:

$$\begin{array}{ccc} \Sigma^+(\mathbb{H}) & \xrightarrow{\sim} & S^2 \\ J & \mapsto & J(1) \end{array} \quad (3)$$

Under this identification, the map (2) become

$$\begin{array}{ccc} \rho_+ : Gr_2(\mathbb{H}) & \rightarrow & S^2 \\ q \wedge q' & \mapsto & q \times_L q' \end{array}$$

We can do the same for  $\Sigma^-(\mathbb{H})$ . We obtain that  $J_{q \wedge q'}^- = \overline{R_{q \times_R q'}} = -R_{q \times_R q'} = \frac{1}{2}(R_{q'} R_{\bar{q}} - R_q R_{\bar{q}'})$ , where  $q \times_R q' = -\text{Im}(\bar{q} \cdot q') = \text{Im}(\bar{q}' \cdot q)$  is the right cross product (it is a bilinear skew map from  $\mathbb{H} \times \mathbb{H}$  to  $\text{Im } \mathbb{H}$ ). Then we have the same identification between  $\Sigma^-(\mathbb{H})$  and  $S^2$ , as in (3). Under this identification  $J^-$  become

$$\begin{array}{ccc} \rho_- : Gr_2(\mathbb{H}) & \rightarrow & S^2 \\ q \wedge q' & \mapsto & q \times_R q' \end{array}$$

## 1.2 Action of $SO(4)$

Recall the following 2-sheeted covering of  $SO(4)$ :

$$\begin{array}{ccc} \chi : S^3 \times S^3 & \rightarrow & SO(4) \\ (a, b) & \mapsto & L_a R_{\bar{b}} \end{array}$$

and set  $Spin(3)_+ = L_a, a \in S^3, Spin(3)_- = R_{\bar{b}}, b \in S^3$ , then  $SO(4) = Spin(3)_+ Spin(3)_- = Spin(3)_- Spin(3)_+$ . We have the two following representation of  $Spin(3)_\varepsilon$ :

$$\chi^+ : L_a \mapsto \text{int}_a = L_a R_{\bar{a}} \in SO(\text{Im } \mathbb{H}), \quad \chi^- : R_{\bar{b}} \mapsto \text{int}_{\bar{b}} = L_b R_{\bar{b}} \in SO(\text{Im } \mathbb{H}).$$

Then the map  $\rho_\varepsilon$  is  $Spin(3)$ -equivariant: for all  $q, q' \in \mathbb{H}, g = L_a R_{\bar{b}} \in SO(4)$ ,

$$\begin{aligned} (gq) \times_L (gq') &= a(q \times_L q') \bar{a} = \text{int}_a(q \times_L q') \\ (gq) \times_R (gq') &= b(q \times_R q') \bar{b} = \text{int}_b(q \times_R q'). \end{aligned}$$

Hence we have  $\forall g \in SO(4)$ ,

$$\rho_\varepsilon(g(q \wedge q')) = \chi_g^\varepsilon(\rho_\varepsilon(q \wedge q'))$$

(where we have extended  $\chi^\varepsilon$  to  $SO(4)$  in an obvious way:  $\chi^+(L_a R_{\bar{b}}) = \chi^+(L_a)$ ,  $\chi^-(L_a R_{\bar{b}}) = \chi^-(R_{\bar{b}})$ ). Besides the map  $J^\varepsilon$  is also  $Spin(3)$ -equivariant, in other words the identification (3) is  $Spin(3)$ -equivariant:

$$\begin{aligned} \forall g \in SO(4), \\ g J_{q \wedge q'}^+ g^{-1} &= L_a R_{\bar{b}} L_{q \times_L q'} R_b L_{\bar{a}} = L_{a(q \times_L q')} a^{-1} = J_{g(q \wedge q')}^+ \end{aligned}$$

The action of  $Spin(3)_+ = SU(\mathbb{R}^4, R_e)$  (resp.  $Spin(3)_- = SU(\mathbb{R}^4, L_e)$ ) on  $\Sigma^-(\mathbb{H})$  (resp.  $\Sigma^+(\mathbb{H})$ ) is trivial. Hence  $SO(4)$  acts on  $\Sigma^\varepsilon(\mathbb{H})$  only by its component  $Spin(3)_\varepsilon$  (in the same way it acts on  $S_\varepsilon^2$  only by its component  $Spin(3)_\varepsilon$  via  $\chi^\varepsilon$ ). In fact, the equality  $gJ_{q \wedge q'}^+ g^{-1} = J_{g(q \wedge q')}^+$  results immediately from the definition of  $J_{q \wedge q'}^+$  and the fact that  $g$  is a positive isometrie. This natural equality which is equivalent to what we called the fundamental property in [17]:  $(gq) \times (gq') = \chi_g(q \times q')$ , is characteristic of dimension 4: in this case it is possible to associate in a natural way (which depends only on the metric and the orientation) to each plan a complex structure, which is not possible in greater dimension. In dimension 8, we must choose a octonionic structure in  $\mathbb{R}^8$  to do that. (see [17]).

### 1.3 The Grassmannian $Gr_2(\mathbb{H})$ is a product of sphere

**Theorem 1** *The map*

$$\begin{aligned} \rho_+ \times \rho_- : Gr_2(\mathbb{H}) &\rightarrow S^2 \times S^2 \\ q \wedge q' &\mapsto (q \times_L q', q \times_R q') \end{aligned}$$

*is a diffeomorphism.*

**Proof.**  $SO(3) \times SO(3)$  acts transitively on  $S^2 \times S^2$  so  $SO(4)$  acts transitively on  $S^2 \times S^2$  via  $\chi^+ \times \chi^-$ , thus  $\rho_+ \times \rho_-$  is surjective.

Let  $\rho \in S(\text{Im } \mathbb{H})$ ,  $g = L_a R_{\bar{b}}$ ,  $g' = L_{a'} R_{\bar{b}'}$   $\in SO(4)$  then we have

$$\begin{aligned} \rho_+ \times \rho_-(g(1 \wedge e)) = \rho_+ \times \rho_-(g'(1 \wedge e)) &\iff (aea^{-1}, beb^{-1}) = (a'ea'^{-1}, b'eb'^{-1}) \\ &\iff a'^{-1}a, b'^{-1}b \in S^1(e) \\ &\implies (L_{a'} R_{\bar{b}'})^{-1} (L_a R_{\bar{b}})(1 \wedge e) = 1 \wedge e \\ &\implies g(1 \wedge e) = g'(1 \wedge e). \end{aligned}$$

Hence, since  $SO(4)$  acts transitively on  $Gr_2(\mathbb{H})$ , we have proved that  $\rho_+ \times \rho_-$  is injective and that

$$\rho_+ \times \rho_-(g(1 \wedge e)) = \rho_+ \times \rho_-(g'(1 \wedge e)) \iff (a'^{-1}a, b'^{-1}b) \in S^1(e) \times S^1(e)$$

(in the previous sequence of implications, the last propositions implies the first one so all the propositions are equivalent). This completes the proof.  $\blacksquare$

As it is the case in [17], it is useful here to introduce a function  $\tilde{\rho}_\varepsilon$  on  $Spin(3)_\varepsilon$  corresponding to  $\rho_\varepsilon$ : we define  $\tilde{\rho}_{\varepsilon e} : Spin(3)_\varepsilon \rightarrow S^2$  by  $\tilde{\rho}_{\varepsilon e}(g) = \chi_g^\varepsilon(e)$  (where  $e \in S(\text{Im } \mathbb{H}) = S^2$ ), i.e. under the identification  $Spin(3)_\varepsilon = S^3$  we have  $\tilde{\rho}_{\varepsilon e}(a) = \text{int}_a(e) = aea^{-1}$ , which is nothing but the Hopf fibration  $S^3 \rightarrow S^3/S^1(e)$ . If  $\rho_\varepsilon(e_1 \wedge e_2) = e$  then  $\tilde{\rho}_{\varepsilon e}(g) = \rho_\varepsilon(g(e_1 \wedge e_2))$ . In the following, we will forget the index  $e$ . Hence, if we take  $e_1 \wedge e_2$  such that  $\rho_\varepsilon(e_1 \wedge e_2) = e$  for  $\varepsilon = \pm 1$  (i.e.  $e_1 \wedge e_2 = (1 \wedge e)^\perp$  which means also that  $(e, e_1, e_2)$  is a direct orthonormal basis

of  $\text{Im } \mathbb{H}$ ) then we have the following commutative diagram:

$$\begin{array}{ccc} S^3 \times S^3 & \xrightarrow{\chi} & SO(4) \\ \tilde{\rho}_+ \times \tilde{\rho}_- \downarrow & & \downarrow \begin{array}{c} g \\ g(e_1 \wedge e_2) \end{array} \\ S^2 \times S^2 & \xleftarrow[\rho_+ \times \rho_-]{\cong} & Gr_2(\mathbb{H}) \end{array}$$

Let us now consider the restriction to  $\text{Im } \mathbb{H} = \mathbb{R}^3$  of this diagram. First the universal covering  $Spin(3) \rightarrow SO(3)$  is obtained by restriction to  $\Delta_3 = \{(a, a), a \in S^3\} \simeq S^3$  of  $\chi: S^3 \times S^3 \rightarrow SO(4)$ , which gives the covering  $(a, a) \mapsto \text{int}_a$ . Then supposing in addition that  $e_1, e_2 \in \text{Im } \mathbb{H}$ , the restriction to  $SO(3)$  of  $SO(4) \rightarrow Gr_2(\mathbb{H})$  is only the surjective map  $g \in SO(3) \mapsto g(e_1 \wedge e_2) \in Gr_3(\mathbb{R}^3)$ . And the restriction to  $Gr_2(\mathbb{R}^3)$  of  $\rho_+ \times \rho_-$  gives the diffeomorphism  $\rho: u \wedge v \in Gr_2(\mathbb{R}^3) \rightarrow u \times v \in S^2$ . Finally the restriction to  $\Delta_3$  of  $\tilde{\rho}_+ \times \tilde{\rho}_-$  gives the Hopf fibration  $\tilde{\rho}: a \in S^3 \mapsto aea^{-1} \in S^2$ . So by restriction to  $\mathbb{R}^3$ , we obtain the classical commutative diagram:

$$\begin{array}{ccc} S^3 & \xrightarrow{x_3} & SO(3) \\ \text{Hopf} \downarrow & & \downarrow \\ S^2 & \xleftarrow{\cong} & Gr_2(\mathbb{R}^3) \end{array}$$

**Remark 1** Besides if we use  $\Sigma^\varepsilon(\mathbb{H})$  instead of the sphere  $S^2$  the Hopf fibration  $\tilde{\rho}_\varepsilon$  become  $SU(2, J_{1 \wedge e}^{-\varepsilon}) \rightarrow \Sigma^\varepsilon(\mathbb{H}) = SU(2, J_{1 \wedge e}^{-\varepsilon})/U(1)_\varepsilon = SO(4)/U(2, J_{1 \wedge e}^\varepsilon)$  where  $U(1)_+ = R_{S^1(e)} = \exp(\mathbb{R}.R_e)$ ,  $U(1)_- = L_{S^1(e)} = \exp(\mathbb{R}.L_e)$ .

#### 1.4 The $\rho$ -harmonic $\omega_I$ -isotropic surfaces

We recall here in the particular case of  $\mathbb{H} = \mathbb{R}^4$  our result obtained in [17] about  $\rho$ -harmonic surfaces. To do that, we need to introduce some notations and definitions. We have

$$\rho_\varepsilon(q \wedge q') = -\varepsilon \sum_{i=1}^3 \omega_i^\varepsilon(q, q') e_i$$

where  $(e_i)_{1 \leq i \leq 3} = (i, j, k)$  and  $\omega_i^\varepsilon = \langle \cdot, J_{1 \wedge e_i}^\varepsilon \cdot \rangle$  (i.e.  $\omega_i^+ = \langle \cdot, L_{e_i} \cdot \rangle$ ,  $\omega_i^- = \langle \cdot, R_{e_i} \cdot \rangle$ ). Let us set, for  $I \subsetneq \{1, 2, 3\}$ ,

$$Q_I^\varepsilon = \{P \in Gr_2(\mathbb{H}) \mid \omega_i^\varepsilon(P) = 0, i \in I\},$$

then  $Q_\emptyset = Gr_2(\mathbb{H})$ ,  $Q_{\{k\}} = \{P \in Gr_2(\mathbb{H})$ , Lagrangian for  $\omega_k^\varepsilon\}$ , and  $Q_{\{k, l\}}^\varepsilon$  is the set of special Lagrangian plan (more precisely the  $\omega_k^\varepsilon$ -Lagrangian plan  $P$  such that  $\det_{\mathbb{C}^2}(P) = \pm i$  under the identification:  $x \in \mathbb{R}^4 \mapsto (x_0 + ix_k, x_l + i\varepsilon x_{k \wedge l}) \in \mathbb{C}^2$ , with  $(k, l, k \wedge l)$  cyclic permutation of  $(1, 2, 3)$ ; for example, if  $(k, l) = (1, 2)$ , it is the identification  $(z_1, z_2) \in \mathbb{C}^2 \mapsto z_1 + z_2 j \in \mathbb{H}$  for  $\varepsilon = 1$  and  $(z_1, z_2) \mapsto z_1 + j z_2$

for  $\varepsilon = -1$ ). We have also  $\rho_\varepsilon(Q_I) = S^I = S(\bigoplus_{i \notin I} \mathbb{R}e_i) = S^2, S^1, \{\pm e_k\}$  for  $|I| = 0, 1, 2$  respectively. Besides we have for  $I = \{i\} \subset \{1, 2, 3\}$ , that  $J^+(Q_I) = L_{S^I} = S^1(\mathbb{R}L_{e_j} \oplus \mathbb{R}L_{e_k})$  is the circle of positive complex structure which anticommute with  $L_{e_i}$ ; and for  $I = \{i, j\} \subset \{1, 2, 3\}$ ,  $J^+(Q_I) = L_{S^I} = \{\pm L_{e_k}\}$ .

We denote by  $G_I^\varepsilon$  the subgroup of  $Spin(3)_\varepsilon$  which conserves  $\omega_i^\varepsilon$ ,  $i \in I$ ; this is the subgroup of  $Spin(3)_\varepsilon$  which commutes with  $L_{e_i}$ ,  $i \in I$ . Then  $G_I^\varepsilon = S^3, S^1, \{\pm 1\}$  for  $|I| = 0, 1, 2$  respectively. We can also consider instead of  $Spin(3)_\varepsilon$  the group  $SO(4)$  (which is equivalent to add the component  $Spin(3)_{-\varepsilon}$  which is useless), then we have  $G_I^\varepsilon = SO(4), U(2, J_{1 \wedge e_i}^\varepsilon), SU(2, J_{1 \wedge e_i}^\varepsilon)$  for  $|I| = 0, 1, 2$  respectively. Let  $e \in S(\bigoplus_{i \notin I} \mathbb{R}e_i)$ . The inner automorphism  $\text{int}_{J_{1 \wedge e}^\varepsilon}$ , define on  $G_I^\varepsilon$  an involution which gives rise to the symmetric space  $S^I = G_I^\varepsilon / G_{I \cap \{k\}}^\varepsilon$  and in the Lie algebra of  $G_I^\varepsilon$ ,  $\mathfrak{g}_I^\varepsilon$ , to the eigenspace decomposition of  $\text{Ad}J_{1 \wedge e}^\varepsilon$ :

$$g_I^\varepsilon = \mathfrak{g}_0^\varepsilon(I) \oplus \mathfrak{g}_2^\varepsilon(I)$$

with  $\mathfrak{g}_0^\varepsilon(I) = \ker(\text{Ad}J_{1 \wedge e}^\varepsilon - \text{Id})$ ,  $\mathfrak{g}_2^\varepsilon(I) = \ker(\text{Ad}J_{1 \wedge e}^\varepsilon + \text{Id})$ .

Let us introduce  $\mathcal{G}_I^\varepsilon = G_I^\varepsilon \times \mathbb{R}^4$  the group of affine isometries of which the linear part is in  $G_I^\varepsilon$ , and its Lie algebra:  $\mathfrak{g}^\varepsilon(I) = \mathfrak{g}_I^\varepsilon \oplus \mathbb{R}^4$ . Consider the automorphism of the group  $\mathcal{G}_I^\varepsilon$ :  $\tau_e^\varepsilon = \text{int}(-\varepsilon J_{1 \wedge e}^\varepsilon, 0)$  with  $e \in S(\bigoplus_{i \notin I} \mathbb{R}e_i)$ . This is an order four automorphism which gives us an eigenspace decomposition of  $\mathfrak{g}^\varepsilon(I)^\mathbb{C}$ :

$$\mathfrak{g}^\varepsilon(I)^\mathbb{C} = \bigoplus_{k \in \mathbb{Z}_4} \tilde{\mathfrak{g}}_k^\varepsilon(I)$$

with  $\tilde{\mathfrak{g}}_{\pm 1}^\varepsilon(I) = \mathfrak{g}_{\pm 1}^\varepsilon = \ker(J_{1 \wedge e}^\varepsilon \pm i\text{Id})$ ,  $\tilde{\mathfrak{g}}_0^\varepsilon(I) = \mathfrak{g}_0^\varepsilon(I)^\mathbb{C}$ ,  $\tilde{\mathfrak{g}}_2^\varepsilon(I) = \mathfrak{g}_2^\varepsilon(I)^\mathbb{C}$ . Moreover we have  $[\tilde{\mathfrak{g}}_k^\varepsilon(I), \tilde{\mathfrak{g}}_l^\varepsilon(I)] \subset \tilde{\mathfrak{g}}_{k+l}^\varepsilon(I)$ .

We fix a value of  $\varepsilon = \pm 1$ . Then let us define as in [17]:

**Definition 1** *Let  $\Sigma$  be a immersed surface in  $\mathbb{H}$ , then a map  $\rho_\Sigma: \Sigma \rightarrow S^2$  is associated to it, defined by  $\rho_\Sigma(z) = \rho_\varepsilon(T_z \Sigma)$  i.e. if  $X: \Sigma \rightarrow \mathbb{H}$  is the immersion then  $\rho_\Sigma = X^* \rho_\varepsilon$ . We will say that  $\Sigma$  is  $\rho$ -harmonic if  $\rho_\Sigma$  is harmonic. Let  $I \subsetneq \{1, 2, 3\}$ , we will say that  $\Sigma$  is  $\omega_I$ -isotropic if  $\forall z \in \Sigma, T_z \Sigma \in Q_I^\varepsilon$  (i.e.  $\Sigma$  is  $\omega_i^\varepsilon$ -isotropic for all  $i \in I$ ) which is equivalent to:  $\rho_\Sigma$  takes values in  $S^I = S(\bigoplus_{i \notin I} \mathbb{R}e_i) \subset S^2$ . Hence for  $|I| = 1$ , the  $\rho$ -harmonic  $\omega_I^\varepsilon$ -isotropic surfaces are the Hamiltonian stationary Lagrangian surfaces in  $\mathbb{C}^2$ , and for  $|I| = 2$ , these are the special Lagrangian surfaces in  $\mathbb{C}^2$  (see above for the identification  $\mathbb{R}^4 \simeq \mathbb{C}^2$ ). If it could be some ambiguity as concerned the value of  $\varepsilon = \pm 1$ , we will use the qualificatifs "left" and "right" respectively to design these two values. A lifted conformal left (resp. right)  $\omega_I$ -isotropic immersion -  $LC\omega_I$  - (if  $I = \emptyset$  we will say a lifted conformal immersion or simply a lift) is a map  $U = (F, X): \Sigma \rightarrow \mathcal{G}_I^\varepsilon$  such that  $X$  is a conformal  $\omega_I^\varepsilon$ -isotropic immersion and  $\tilde{\rho}_\varepsilon \circ F = \rho_\Sigma$*

We have obtained the following result in [17]:

**Theorem 2** *Let  $\Omega$  be a simply connected open domain, and  $\alpha$  a 1-form on  $\Omega$  with values in  $\mathfrak{g}(I)$ , then*



- $\alpha$  is the Maurer-Cartan form of a  $LC\omega_I$  if and only if

$$d\alpha + \alpha \wedge \alpha = 0, \quad \alpha''_{-1} = 0 \quad \text{and} \quad \alpha'_{-1} \text{ does not vanish}$$

- furthermore,  $\alpha$  corresponds to some  $\rho$ -harmonic  $\omega_I$ -isotropic conformal immersion if and only if the extended Maurer-Cartan form  $\alpha_\lambda = \lambda^{-2}\alpha'_2 + \lambda^{-1}\alpha_{-1} + \alpha_0 + \lambda\alpha_1 + \lambda^2\alpha''_2$  satisfies

$$d\alpha_\lambda + \alpha_\lambda \wedge \alpha_\lambda = 0, \quad \forall \lambda \in \mathbb{C}^*.$$

Let us recall the proof given in [17].

**Proof.** To fix ideas, we take  $\varepsilon = 1$ .  $\alpha$  is a Maurer-Cartan form if and only if it satisfies the Maurer-Cartan equation. In this case, it can be integrated by  $U = (F, X): \Sigma \rightarrow \mathcal{G}_I$ ,  $\alpha = U^{-1}.dU$ ,  $U(z_0) = 1$ . Hence  $\alpha = U^{-1}.dU = (F^{-1}.dF, F^{-1}.dX)$ . Moreover,  $F^{-1}.dX = \alpha_{-1} + \alpha_1$  is real and  $\mathfrak{g}_{\pm 1} = \{V \pm iL_e V, V \in \mathbb{H}\}$  so  $\alpha_{-1} = \overline{\alpha_1}$ . Hence  $\alpha''_{-1} = 0 \iff \alpha''_1 = \alpha'_1 = 0 \iff \alpha_{-1} = (F^{-1} \frac{\partial X}{\partial z}) dz \iff F^{-1} \frac{\partial X}{\partial y} = L_e(F^{-1} \frac{\partial X}{\partial x}) \iff F^{-1}.dX = h(q_0 du + q'_0 dv)$  with  $h \in C^\infty(\omega, \mathbb{R})$ ,  $q_0, q'_0 \in C^\infty(\Omega, S^3)$ ,  $\langle q_0, q'_0 \rangle = 0$  and  $\rho(q_0, q'_0) = e$ . Thus we have  $\alpha''_{-1} = 0$  and  $\alpha'_{-1} \neq 0 \iff dX = e^f(q du + q' dv)$  with  $f \in C^\infty(\omega, \mathbb{R})$ ,  $(q, q')$  orthonormal and  $\rho(q, q') = \tilde{\rho}_e(F)$  i.e.  $\rho_X = \tilde{\rho}_e(F)$ . This proves the first point. Hence we have the decomposition  $\alpha = \alpha_2 + \alpha_{-1} + \alpha_0 + \alpha_1 = \alpha'_2 + \alpha'_{-1} + \alpha_0 + \alpha'_1 + \alpha''_2$ . Furthermore, using the commutation relations  $[\tilde{\mathfrak{g}}_k(I), \tilde{\mathfrak{g}}_l(I)] \subset \tilde{\mathfrak{g}}_{k+l}(I)$ ,  $[\mathfrak{g}_{\pm 1}, \mathfrak{g}_{\pm 1}] = \{0\}$ , we obtain

$$\begin{aligned} d\alpha_\lambda + \alpha_\lambda \wedge \alpha_\lambda &= \lambda^{-2}(d\alpha'_2 + [\alpha_0 \wedge \alpha'_2]) \\ &\quad + \lambda^{-1}(d\alpha'_{-1} + [\alpha'_{-1} \wedge \alpha_0] + [\alpha''_1 \wedge \alpha'_2]) \\ &\quad + (d\alpha_0 + \frac{1}{2}[\alpha_0 \wedge \alpha_0] + \frac{1}{2}[\alpha'_2 \wedge \alpha''_2]) \\ &\quad + \lambda(d\alpha''_1 + [\alpha''_1 \wedge \alpha_0] + [\alpha'_{-1} \wedge \alpha''_2]) \\ &\quad + \lambda^2(d\alpha''_2 + [\alpha_0 \wedge \alpha''_2]), \end{aligned}$$

the coefficient of  $\lambda^{-1}, \lambda^0, \lambda$  are respectively the projection of  $d\alpha + \alpha \wedge \alpha$  on  $\mathfrak{g}_{-1}, \mathfrak{g}_0, \mathfrak{g}_1$  respectively so they vanish and hence

$$d\alpha_\lambda + \alpha_\lambda \wedge \alpha_\lambda = d\beta_{\lambda^2} + \beta_{\lambda^2} \wedge \beta_{\lambda^2}$$

where  $\beta_\lambda = \lambda^{-1}\alpha'_2 + \alpha_0 + \lambda\alpha''_2$  is the extended Maurer-Cartan form of  $\beta = F^{-1}.dF$ , the Maurer-Cartan form of the lift  $F \in G_I$  of  $\rho_X \in S^I$ . According to [6], we know that  $\rho_X$  is harmonic if and only if  $d\beta_\lambda + \beta_\lambda \wedge \beta_\lambda = 0, \forall \lambda \in \mathbb{C}^*$ . This proves the second point and completes the proof.  $\blacksquare$

**Remark 2** We have  $\rho_-(x, y) = -\text{Im}(\bar{x}.y) = \rho_+(\bar{x}, \bar{y})$ . Hence  $X: \Omega \rightarrow \mathbb{H}$  is  $\rho_-$ -harmonic if and only if  $\bar{X}$  is  $\rho_+$ -harmonic, and  $X$  is  $\omega_I^-$ -isotropic if and only if  $\bar{X}$  is  $\omega_I^+$ -isotropic. Besides if  $U = (F, X): \Omega \rightarrow G_I \times \mathbb{H}$  is a left  $LC\omega_I$  then we have  $F = L_a$  and  $aea^{-1} = \rho_X = \rho_+(q, q')$  with  $dX = e^\omega(q du + q' dv)$ ,  $(q, q')$  orthonormal. Thus,  $\rho_-(\bar{q}, \bar{q}') = aea^{-1}$  and hence  $\underline{U} = (R_{\bar{a}}, \bar{X})$  is a right  $LC\omega_I$ .

**Remark 3** The restriction to  $\text{Im } \mathbb{H} = \mathbb{R}^3$  of the left (or right) cross product gives us the usual cross product of  $\mathbb{R}^3$ . Hence a surface in  $\text{Im } \mathbb{H}$  is left (resp. right)  $\rho$ -harmonic if and only if it is a constant mean curvature surface. In the same way, it is easy to see that a surface in  $S^3$  is left (resp. right)  $\rho$ -harmonic if and only if it is a constant mean curvature surface.

**Remark 4** We can apply now the Dorfmeister-Pedit-Wu method to obtain a Weierstrass representation of  $\rho$ -harmonic surfaces (see [6, 12, 13, 14, 17, 18]).

## 2 Second Elliptic Integrable System

### 2.1 4-symmetric spaces and twistor space

**Definition 2** Let  $M$  be a Riemannian symmetric space. We will say that a Lie group  $G$  acts symmetrically on  $M$  or that  $M$  is a  $G$ -symmetric space if  $G$  acts transitively and isometrically on  $M$  and if there exists an involutive automorphism of  $G$ ,  $\sigma$ , such that  $H$  the isotropy subgroup at a fixed point  $p_0 \in M$ , satisfies  $(G^\sigma)^0 \subset H \subset G^\sigma$ . We will say also that  $G/H$  is a symmetric realisation of  $M$ . We will say that a  $G$ -homogeneous space  $N = G/G_0$  is a 4-symmetric bundle over the  $G$ -symmetric space  $M$  if there exists an order four automorphism  $\tau$  of  $G$ , such that  $(G^\tau)^0 \subset G_0 \subset G^\tau$ , and  $(G, \tau)$  gives rise to the symmetric space  $M$ , i.e.  $\sigma = \tau^2$  and  $G_0 \subset H$ .

A  $G$ -homogeneous space  $N = G/G_0$  is a locally 4-symmetric space if there exists a order four automorphism of the Lie algebra  $\mathfrak{g} = \text{Lie } G$ ,  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  such that  $\mathfrak{g}^\tau = \text{Lie } G_0$ . We will say that  $G/G_0$  is a locally 4-symmetric bundle over the  $G$ -symmetric space  $M$  if  $\tau^2 = T_1\sigma$  (and  $G_0 \subset H$ ).

Let us consider  $M$  a  $G$ -symmetric space with  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  an order four automorphism such that  $\tau^2 = T_1\sigma$ . The automorphism  $\tau$  gives us an eigenspace decomposition of  $\mathfrak{g}^{\mathbb{C}}$ :

$$\mathfrak{g}^{\mathbb{C}} = \bigoplus_{k \in \mathbb{Z}_4} \tilde{\mathfrak{g}}_k$$

where  $\tilde{\mathfrak{g}}_k$  is the  $e^{ik\pi/2}$ -eigenspace of  $\tau$ . We have clearly  $\tilde{\mathfrak{g}}_0 = \mathfrak{g}_0^{\mathbb{C}}$ ,  $\overline{\tilde{\mathfrak{g}}_k} = \tilde{\mathfrak{g}}_{-k}$  and  $[\tilde{\mathfrak{g}}_k, \tilde{\mathfrak{g}}_l] \subset \tilde{\mathfrak{g}}_{k+l}$ . We define  $\mathfrak{g}_2$ ,  $\mathfrak{m}$  and  $\underline{\mathfrak{g}}_1$  by

$$\tilde{\mathfrak{g}}_2 = \mathfrak{g}_2^{\mathbb{C}}, \quad \mathfrak{m}^{\mathbb{C}} = \tilde{\mathfrak{g}}_{-1} \oplus \tilde{\mathfrak{g}}_1 \quad \text{and} \quad \underline{\mathfrak{g}}_1^{\mathbb{C}} = \bigoplus_{k \in \mathbb{Z}_4 \setminus \{0\}} \tilde{\mathfrak{g}}_k,$$

it is possible because  $\overline{\tilde{\mathfrak{g}}_2} = \tilde{\mathfrak{g}}_2$  and  $\overline{\tilde{\mathfrak{g}}_{-1}} = \tilde{\mathfrak{g}}_1$ . Let us set  $\mathfrak{g}_{-1} = \tilde{\mathfrak{g}}_{-1}$ ,  $\mathfrak{g}_1 = \tilde{\mathfrak{g}}_1$ ,  $\mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_2$ . Then

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$$

is the eigenspace decomposition of the involutive automorphism  $\sigma$ ,  $\mathfrak{h}$  is the Lie algebra of  $H$ , the isotropy subgroup of  $G$  at a reference point  $p_0$ , and  $\mathfrak{m}$  is identified to the tangent space  $T_{p_0}M$ . Besides we remark that  $\tau|_{\mathfrak{m}} \in \Sigma(\mathfrak{m})$  (since  $\tau|_{\mathfrak{m}^{\mathbb{C}}} = -i\text{Id}_{\mathfrak{g}_{-1}} \oplus i\text{Id}_{\mathfrak{g}_1}$ )<sup>1</sup>, which gives us the following theorem (proved

<sup>1</sup>We choose a metric in  $\mathfrak{m}$  stabilized by  $\tau|_{\mathfrak{m}}$  (and of course by  $\text{Ad}H$ ), see section 3.1

in section 3.2).

**Theorem 3** *Let us consider  $M$  a Riemannian  $G$ -symmetric space and  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  an order four automorphism such that  $\tau^2 = T_1\sigma$ . Let us make  $G$  acting on  $\Sigma(M): g \cdot J = gJg^{-1}$ . Let  $J_0 \in \Sigma(T_{p_0}M)$  be the complex structure corresponding to  $\tau|_{\mathfrak{m}} \in \Sigma(\mathfrak{m})$  (resp. to  $-\tau|_{\mathfrak{m}} = \tau|_{\mathfrak{m}}^{-1} \in \Sigma(\mathfrak{m})$ ), under the identification  $T_{p_0}M = \mathfrak{m}$ . Then the orbit of  $J_0$  under the action of  $G$  is an immersed submanifold of  $\Sigma(M)$ . Denoting by  $G_0$  the stabilizer of  $J_0$ , then  $\text{Lie } G_0 = \mathfrak{g}^\tau$  and thus  $G/G_0$  is a locally 4-symmetric bundle over  $M$ , and the natural map*

$$\begin{aligned} i: G/G_0 &\longrightarrow \Sigma(M) \\ g \cdot G_0 &\longmapsto gJ_0g^{-1}. \end{aligned}$$

*is an injective immersion and a morphism of bundle. Moreover, if the image of  $G$  in  $\text{Is}(M)$  (the group of isometry of  $M$ ) is closed, then  $i$  is an embedding.*

## 2.2 The second elliptic integrable system associated to a 4-symmetric space

We give ourself  $M$  a Riemannian  $G$ -symmetric space with  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  an order four automorphism such that  $\tau^2 = T_1\sigma$ , and  $N = G/G_0$  the associated locally 4-symmetric space given by theorem 3. We use the same notations as in Section 2.1. Then let us recall what is a second elliptic system according to C.L. Terng (see [20]).

**Definition 3** *The second  $(\mathfrak{g}, \tau)$ -system is the equation for  $(u_0, u_1, u_2): \mathbb{C} \rightarrow \oplus_{j=0}^2 \tilde{\mathfrak{g}}_{-j}$ ,*

$$\begin{cases} \partial_{\bar{z}}u_2 + [\bar{u}_0, u_2] = 0 & (a) \\ \partial_{\bar{z}}u_1 + [\bar{u}_0, u_1] + [\bar{u}_1, u_2] = 0 & (b) \\ -\partial_{\bar{z}}u_0 + \partial_z\bar{u}_0 + [u_0, \bar{u}_0] + [u_1, \bar{u}_1] + [u_2, \bar{u}_2] = 0. & (c) \end{cases} \quad (4)$$

*It is equivalent to say that the 1-form*

$$\alpha_\lambda = \sum_{i=0}^2 \lambda^{-i} u_i dz + \lambda^i \bar{u}_i d\bar{z} = \lambda^{-2} \alpha'_2 + \lambda^{-1} \alpha'_1 + \alpha_0 + \lambda \alpha''_1 + \lambda^2 \alpha''_2 \quad (5)$$

*satisfies the zero curvature equation:*

$$d\alpha_\lambda + \frac{1}{2}[\alpha_\lambda \wedge \alpha_\lambda] = 0, \quad (6)$$

*for  $\lambda \in \mathbb{C}^*$ . We will speak about the  $(G, \tau)$ -system ( $\tau$  is an automorphism of  $\text{Lie } G = \mathfrak{g}$ ) when we look at solutions of the  $(\mathfrak{g}, \tau)$ -system in  $G$ , i.e. maps  $U: \mathbb{C} \rightarrow G$  such that their Maurer-Cartan form is solution of the  $(\mathfrak{g}, \tau)$ -system, in other words when we integrate the zero curvature equation (6) in  $G$ . We will call (geometric) solution of the second elliptic integrable system associated to the locally 4-symmetric space  $G/G_0$  a map  $J: \mathbb{C} \rightarrow G/G_0$  which can be lifted into a solution  $U: \mathbb{C} \rightarrow G$  of (4).*

**Remark 5** In (4),  $\{\operatorname{Re}(a), (b), (c)\}$  is equivalent to  $d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0$ . Hence the additional condition added to the Maurer-Cartan equation by the zero curvature equation (6) is  $\operatorname{Im}(\partial_{\bar{z}}\alpha'_2 + [\alpha''_0, \alpha'_2]) = 0$  or equivalently

$$d(\star\alpha_2) + [\alpha_0 \wedge (\star\alpha_2)] = 0.$$

The first example of second elliptic system was given by F. Hélein and P. Romon (see [12, 14]): they showed that the equations for Hamiltonian stationary Lagrangian surfaces in 4-dimension Hermitian symmetric spaces are the second elliptic system associated to certain 4-symmetric spaces. Then in [17], we found an other example in  $\mathbb{O}$ : the  $\rho$ -harmonic surfaces in  $\mathbb{O}$ , which by restriction to  $\mathbb{H}$  gives us the  $\rho$ -harmonic surfaces in  $\mathbb{H}$  (studied in section 1) which generalize the Hamiltonian stationary Lagrangian surfaces in  $\mathbb{C}^2$ .

**Definition 4** Let  $M$  be a Riemannian manifold and  $\nabla$  its Levi-Civita connection which induces a connection on  $\operatorname{End}(TM)$ . Let us define for each  $(p, J_p) \in \Sigma(M)$  the orthogonal projection

$$\operatorname{pr}^\perp(p, J_p): \operatorname{End}(T_p M) \rightarrow T_{J_p}(\Sigma(T_p M))$$

( $T_p M$  is an Euclidean vector space so  $\Sigma(T_p M)$  is a submanifold of the Euclidean space  $\operatorname{End}(T_p M)$  and so  $T_{J_p}\Sigma(T_p M)$  is a vector subspace of  $\operatorname{End}(T_p M)$  and we can consider the orthogonal projection on this subspace). Given  $J: \mathbb{C} \rightarrow \Sigma(M)$  we set

$$\Delta J(z) = 4\operatorname{pr}^\perp(J(z)).\nabla_{\frac{\partial}{\partial \bar{z}}}(\nabla_{\frac{\partial}{\partial z}} J).$$

(in fact we endow the bundle  $\Sigma(M)$  with the connection defined by the horizontal distribution  $(\mathcal{H}_J \oplus (T_J\Sigma(T_p M))^\perp) \cap T_J\Sigma(M)$ , where  $\mathcal{H}_J$  is the connection of  $\operatorname{End}(TM): T\operatorname{End}(TM) = \mathcal{H} \oplus \operatorname{End}(TM)$ ). We will say that  $J$  is vertically harmonic if  $\Delta J = 0$ .

**Definition 5** Let  $(L, j)$  be a Riemann surface,  $M$  a oriented manifold and  $X: L \rightarrow M$  a immersion. Let  $J: L \rightarrow X^*(\Sigma(M))$  be a almost complex structure on the vector bundle  $X^*(TM)$ . Then we will say that  $J$  is an admissible twistor lift of  $X$  if one of the following equivalent statements holds:

- (i)  $X$  is  $J$ -holomorphic:  $\star dX := dX \circ j = J.dX$
- (ii)  $J$  is an extension of the complex structure on the oriented tangent plan  $P = X_*(TL)$  induced by  $j$ , the complex structure of  $L$ , or equivalently  $J$  induces the complex structure  $j$  in  $L$ .
- (iii)  $X$  is a conformal immersion and  $J$  stabilizes the tangent plan  $X_*(TL)$ , i.e. for all  $z \in L$ ,  $J_z$  stabilizes  $X_*(T_z L)$  and induces on it the same orientation, which we will denote by  $J \circ X_*(TL)$
- (iv)  $X$  is a conformal immersion and  $J$  is an extension of the unique positive complex structure  $I_P$  of the tangent plan  $P = X_*(TL)$ .

Finally, we will say that a map  $J: L \rightarrow \Sigma(M)$  is an admissible twistor lift if its projection  $X = \text{pr}_M \circ J: L \rightarrow M$  is an immersion and  $J$  is an admissible twistor lift of it.

**Theorem 4** *Let  $L$  be a simply connected Riemann surface and  $(G, \tau)$  a locally 4-symmetric bundle over a symmetric space  $M = G/H$ . Let  $J_0 \in \Sigma(T_{p_0}M)$  be the complex structure corresponding to  $\tau|_{\mathfrak{m}}^{-1}$  (see Section 2.1). Let be  $J_X: L \rightarrow i(G/G_0) \subset \Sigma(G/H)$ . Then the two following statements are equivalent:*

- $J_X$  is an admissible twistor lift.
- Any lift  $F: L \rightarrow G$  of  $J_X$  ( $FJ_0F^{-1} = J_X$ ) gives rise to a Maurer-Cartan form  $\alpha = F^{-1}.dF$  which satisfies:  $\alpha''_{-1} = \alpha'_1 = 0$  and  $\alpha'_{-1}$  does not vanish.

Furthermore, under these statements,  $J_X: L \rightarrow \Sigma(G/H)$  is vertically harmonic if and only if  $J_X: L \rightarrow G/G_0$  is solution of the second elliptic integrable system associated to the locally 4-symmetric space  $(G, \tau)$ , i.e.

$$d\alpha_\lambda + \frac{1}{2}[\alpha_\lambda \wedge \alpha_\lambda] = 0, \quad \forall \lambda \in \mathbb{C}^*,$$

where  $\alpha_\lambda = \lambda^{-2}\alpha'_2 + \lambda^{-1}\alpha'_{-1} + \alpha_0 + \lambda\alpha'_1 + \lambda^2\alpha''_2$  is the extended Maurer-Cartan form of  $\alpha$ .

**Proof.** For the first point, let us make  $F^{-1}$  acting on the equation  $dX \circ j = J_X.dX$ , we obtain  $\alpha_{\mathfrak{m}} \circ j = \tau|_{\mathfrak{m}}(\alpha_{\mathfrak{m}})$  which is equivalent to  $\alpha''_{-1} = \alpha'_1 = 0$ . Thus  $\alpha_{-1}(\frac{\partial}{\partial z}) = \alpha_{\mathfrak{m}}(\frac{\partial}{\partial z}) = F^{-1}.\frac{\partial X}{\partial z}$ , and  $X$  is an immersion if and only if  $\alpha'_{-1}$  does not vanish.

For the second point, let us recall that  $\text{End}(T_pM) = \text{sym}(T_pM) \oplus^{\perp} \mathfrak{so}(T_pM)$  and given  $J \in \Sigma(T_pM)$ , we have  $T_J\Sigma(T_pM) = \text{Ant}(J) = \{A \in \mathfrak{so}(T_pM) | AJ + JA = 0\}$  and  $(T_J\Sigma(T_pM))^{\perp} \cap \mathfrak{so}(T_pM) = \text{Com}(J) = \{A \in \mathfrak{so}(T_pM) | [A, J] = 0\}$ .

Now, let us compute the connection  $X^*\nabla$  on  $X^*(\text{End}(TM))$ , in terms of the Lie algebra setting. Let  $A$  be a section of  $X^*(\text{End}(TM))$  and  $Y$  a section of  $X^*(TM)$ . Let  $A_0 \in C^\infty(L, \text{End}(T_{p_0}M))$  be defined by  $A_{F.p_0} = FA_0F^{-1}$  and  $A_{\mathfrak{m}} \in C^\infty(L, \text{End}(\mathfrak{m}))$  its image under the identification  $T_{p_0}M = \mathfrak{m}$ . Then  $A_{F.p_0}$  corresponds to  $\text{Ad}F \circ A_{\mathfrak{m}} \circ \text{Ad}F^{-1}$  (under the identification  $TM = [\mathfrak{m}]$ ). In particular  $(J_X)_{\mathfrak{m}} = \tau|_{\mathfrak{m}}$  (we suppose  $F(p_0) = 1$ ). We set also  $Y = \text{Ad}F(\xi).p_0$ ,  $\xi \in C^\infty(L, \mathfrak{m})$ . From now, we do the identification  $TM = [\mathfrak{m}]$  without precising it. We have

$$\begin{aligned} (\nabla A)(Y) &= \nabla(A Y) - A(\nabla Y) \\ &= \text{Ad}F([d(A_{\mathfrak{m}}\xi) + [\alpha, A_{\mathfrak{m}}.\xi]]_{\mathfrak{m}} - A_{\mathfrak{m}}(d\xi + [\alpha, \xi]_{\mathfrak{m}})) \\ &= \text{Ad}F((dA_{\mathfrak{m}}\xi + (\text{ad}\alpha_{\mathfrak{h}} \circ A_{\mathfrak{m}} - A_{\mathfrak{m}} \circ \text{ad}\alpha_{\mathfrak{h}})\xi)). \end{aligned}$$

Hence

$$\nabla A = \text{Ad}F(dA_{\mathfrak{m}} + [\text{ad}_{\mathfrak{m}}\alpha_{\mathfrak{h}}, A_{\mathfrak{m}}]).$$

In particular,<sup>2</sup>

$$\nabla_{\frac{\partial}{\partial \bar{z}}} J_X = 2\text{Ad}F(\text{ad}_m \alpha'_2 \circ \tau_{|m})$$

(because  $\text{ad}_m \mathfrak{g}_0$  commutes with  $\tau_{|m}$  whereas  $\text{ad}_m \mathfrak{g}_2$  anticommutes with it) and thus

$$\begin{aligned} \nabla_{\frac{\partial}{\partial \bar{z}}}(\nabla_{\frac{\partial}{\partial \bar{z}}} J_X) &= 2\text{Ad}F(\text{ad}_m(\partial_{\bar{z}} \alpha'_2) \circ \tau_{|m} + [\text{ad}_m(\alpha''_0), \text{ad}_m(\alpha'_2) \circ \tau_{|m}]) \\ &= 2\text{Ad}F(\text{ad}_m(\partial_{\bar{z}} \alpha'_2) \circ \tau_{|m} + \text{ad}_m([\alpha''_0, \alpha'_2]) \circ \tau_{|m} \\ &\quad + [\text{ad}_m \alpha''_2, \text{ad}_m(\alpha'_2) \circ \tau_{|m}]) \\ &= 2\text{Ad}F(\text{ad}_m(\partial_{\bar{z}} \alpha'_2 + [\alpha''_0, \alpha'_2]) \circ \tau_{|m} + [\text{ad}_m \alpha''_2, \text{ad}_m(\alpha'_2) \circ \tau_{|m}]) \end{aligned}$$

but  $\text{Ad}F([\text{ad}_m \alpha''_2, \text{ad}_m(\alpha'_2) \circ \tau_{|m}])$  commutes with  $\text{Ad}F(\tau_{|m}) = J_X$  so it is orthogonal to  $T_J \Sigma(T_p M)$  thus

$$\text{pr}^\perp(J_X) \cdot \nabla_{\frac{\partial}{\partial \bar{z}}}(\nabla_{\frac{\partial}{\partial \bar{z}}} J_X) = 2\text{Ad}F(\text{ad}_m(\partial_{\bar{z}} \alpha'_2 + [\alpha''_0, \alpha'_2]) \circ \tau_{|m}).$$

Hence, since  $\text{ad}_m$  is injective,

$$\Delta J_X = 0 \iff \partial_{\bar{z}} \alpha'_2 + [\alpha''_0, \alpha'_2] = 0. \quad (7)$$

This completes the proof.  $\blacksquare$

**Remark 6** The equivalence (7) holds for any map  $J_X: L \rightarrow i(G/G_0)$ . Indeed, we have not used the fact that  $J_X$  is an admissible twistor lift to prove this equivalence.

**Theorem 5** *Let  $J_X: L \rightarrow G/G_0 \hookrightarrow \Sigma(G/H)$  be an admissible twistor lift. Then  $J_X: L \rightarrow G/G_0$  is harmonic<sup>3</sup> if and only if  $X: L \rightarrow G/H$  is harmonic and  $J_X$  is vertically harmonic.*

**Proof.**  $J_X: L \rightarrow G/G_0$  is harmonic if and only if the Maurer cartan form  $\alpha = F^{-1} \cdot dF$  of the lift  $F: L \rightarrow G$  of  $J_X$  ( $F J_0 F^{-1} = J_X$ ) satisfies (see [3])

$$\partial_{\bar{z}} \underline{\alpha}'_1 + [\alpha''_0, \underline{\alpha}'_1] + [\underline{\alpha}''_1, \underline{\alpha}'_1]_{\mathfrak{g}_1} = 0$$

(where  $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$  is the reductive decomposition corresponding to the homogeneous space  $G/G_0$ , see Section 2.1) which splits into

$$\begin{cases} \partial_{\bar{z}} \alpha'_2 + [\alpha''_0, \alpha'_2] + [\alpha''_1, \alpha'_1] + [\alpha''_{-1}, \alpha'_{-1}] = 0 \\ \partial_{\bar{z}} \alpha'_{-1} + [\alpha''_0, \alpha'_{-1}] + [\alpha''_2, \alpha'_1] + [\alpha''_1, \alpha'_2] = 0 \\ \partial_{\bar{z}} \alpha'_1 + [\alpha''_0, \alpha'_1] + [\alpha''_2, \alpha'_{-1}] + [\alpha''_{-1}, \alpha'_2] = 0. \end{cases} \quad (8)$$

then, using  $\alpha''_{-1} = \alpha'_1 = 0$ , we obtain

$$\begin{cases} \partial_{\bar{z}} \alpha'_2 + [\alpha''_0, \alpha'_2] = 0 \\ \partial_{\bar{z}} \alpha'_{-1} + [\alpha''_0, \alpha'_{-1}] = 0 \\ [\alpha''_2, \alpha'_{-1}] = 0 \end{cases}$$

<sup>2</sup>In all the proof, we will merge  $\alpha'_k$  (resp.  $\alpha''_k$ ) with  $\alpha'_k(\frac{\partial}{\partial \bar{z}})$  (resp.  $\alpha''_k(\frac{\partial}{\partial \bar{z}})$ ), and in particular write  $[\alpha''_k, \alpha'_l]$  instead of  $[\alpha''_k(\frac{\partial}{\partial \bar{z}}), \alpha'_l(\frac{\partial}{\partial \bar{z}})]$ .

<sup>3</sup>with respect to the killing form.

(in the second equation, we have used  $[\alpha_1'', \alpha_2'] = -\overline{[\alpha_2'', \alpha_{-1}']} = 0$ ). Besides  $X: L \rightarrow G/H$  is harmonic if and only if we have

$$\partial_{\bar{z}}\alpha'_m + [\alpha''_h, \alpha'_m] = 0$$

which splits into

$$\begin{cases} \partial_{\bar{z}}\alpha'_{-1} + [\alpha''_0, \alpha'_{-1}] + [\alpha''_2, \alpha'_1] = 0 \\ \partial_{\bar{z}}\alpha'_1 + [\alpha''_0, \alpha'_1] + [\alpha''_2, \alpha'_{-1}] = 0. \end{cases} \quad (9)$$

and using  $\alpha''_{-1} = \alpha'_1 = 0$ , we obtain

$$\begin{cases} \partial_{\bar{z}}\alpha'_{-1} + [\alpha''_0, \alpha'_{-1}] = 0 \\ [\alpha''_2, \alpha'_{-1}] = 0. \end{cases}$$

This completes the proof. ■

### 3 Structure of 4-symmetric bundles over symmetric spaces

#### 3.1 4-symmetric spaces

Let  $G$  be a Lie group with Lie algebra  $\mathfrak{g}$ ,  $\tau: G \rightarrow G$  an order four automorphism with the fixed point subgroup  $G^\tau$ , and the corresponding Lie algebra  $\mathfrak{g}_0 = \mathfrak{g}^\tau$ . Let  $G_0$  be a subgroup of  $G$  such that  $(G^\tau)^0 \subset G_0 \subset G^\tau$ , then  $\text{Lie } G_0 = \mathfrak{g}_0$  and  $G/G_0$  is a 4-symmetric space. The automorphism  $\tau$  gives us an eigenspace decomposition of  $\mathfrak{g}^{\mathbb{C}}$  for which we use the notation of section 2.1. Then  $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$  is the eigenspace decomposition of the involutive automorphism  $\sigma = \tau^2$ . Let  $H$  be a subgroup of  $G$  such that  $(G^\sigma)^0 \subset H \subset G^\sigma$  then  $\text{Lie } H = \mathfrak{h}$  and  $G/H$  is a symmetric space. We will often suppose that  $G_0$  and  $H$  are chosen such that  $G_0 = G^\tau \cap H$ . With this condition,  $G_0 \subset H$  so that  $G/G_0$  is a bundle over  $G/H$ . Recall that the tangent bundle  $TM$  is canonically isomorphic to the subbundle  $[\mathfrak{m}]$  of the trivial bundle  $M \times \mathfrak{g}$ , with fiber  $\text{Ad}g(\mathfrak{m})$  over the point  $x = g.H \in M$ . Under this identification the canonical  $G$ -invariant connection of  $M$  is just the flat differentiation in  $M \times \mathfrak{g}$  followed by the projection on  $[\mathfrak{m}]$  along  $[\mathfrak{h}]$  (which is defined in the same way as  $\mathfrak{m}$ ) (see [2] or [4]). For the homogeneous space  $N = G/G_0$  we have the following reductive decomposition

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \underline{\mathfrak{g}}_1 \quad (10)$$

( $\underline{\mathfrak{g}}_1$  can be written  $\underline{\mathfrak{g}}_1 = \mathfrak{m} \oplus \mathfrak{g}_2$ ) with  $[\mathfrak{g}_0, \underline{\mathfrak{g}}_1] \subset \underline{\mathfrak{g}}_1$ . As for the symmetric space  $G/H$ , we can identify the tangent bundle  $TN$  with the subbundle  $[\underline{\mathfrak{g}}_1]$  of the trivial bundle  $N \times \mathfrak{g}$ , with fiber  $\text{Ad}g(\underline{\mathfrak{g}}_1)$  over the point  $x = g.G_0 \in N$ .

The symmetric space  $M = G/H$  (resp. the homogeneous space  $N = G/G_0$ ) is Riemannian if it admits an  $G$ -invariant metric, which is equivalent to say that  $\mathfrak{m}$  (resp.  $\underline{\mathfrak{g}}_1$ ) admit an  $\text{Ad}(H)$ -invariant (resp.  $\text{Ad}(G_0)$ -invariant) inner product or equivalently, that  $\text{Ad}_{\mathfrak{m}}(H)$  (resp.  $\text{Ad}_{\underline{\mathfrak{g}}_1}(G_0)$ ) be relatively compact<sup>4</sup>. We

<sup>4</sup>In the literature, it is often supposed that  $\text{Ad}_{\mathfrak{m}}(H)$  is compact. We will see that these two hypothesis are in fact equivalent.

remark that the Levi-Civita connection coincide with the previous canonical  $G$ -invariant connection and in particular is independent of the  $G$ -invariant metric chosen. We will always suppose that the symmetric spaces  $M$  which we consider are Riemannian. We will in addition to that suppose that the  $\text{Ad}(H)$ -invariant inner product in  $\mathfrak{m}$  is also invariant by  $\tau|_{\mathfrak{m}}$  (such a inner product always exists when  $\text{Ad}_{\mathfrak{m}}(H)$  is compact, see the appendix). We will also suppose that  $M$  is connected, then  $G^0$  acts transitively on  $M$  and so we can suppose that  $G$  is connected.

We want to study the Riemannian symmetric spaces  $M$  such that there exists a 4-symmetric space  $(G, \tau)$  which gives rise to  $M$  in the same way as above. For this let us recall the following theorem:

**Theorem 6** [2, 15] *Let  $M$  be a Riemannian manifold.*

- (a) *The group  $\text{Is}(M)$  of all the isometries of  $M$  is a Lie group and acts differentiably on  $M$ .*
- (b) *Let  $p_0 \in M$ , then an isometry  $f$  of  $M$  is determined by the image  $f(p_0)$  of the point  $p_0$  and the corresponding tangent map  $T_{p_0}f$  (i.e. if  $f(p_0) = g(p_0)$  and  $T_{p_0}f = T_{p_0}g$  then  $f = g$ ).*
- (c) *The isotropy subgroup  $\text{Is}_{p_0}(M) = \{f \in \text{Is}(M); f(p_0) = p_0\}$  is a closed subgroup of  $\text{Is}(M)$  and the linear isotropy representation  $\rho_{p_0}: f \in \text{Is}_{p_0}(M) \mapsto T_{p_0}f \in O(T_{p_0}M)$  is an isomorphism from  $\text{Is}_{p_0}(M)$  onto a closed subgroup of  $O(T_{p_0}M)$ . Hence  $\text{Is}_{p_0}(M)$  is a compact subgroup of  $\text{Is}(M)$ .*
- (d) *If  $M$  is a Riemannian homogeneous space,  $M = G/H$  with  $G = \text{Is}(M)$ ,  $H = \text{Is}_{p_0}(M)$  and  $\mathfrak{m}$  an  $\text{Ad}H$ -invariant space such that  $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$ , then the previous closed subgroup, image of  $H$  by the preceding isomorphism  $\rho_{p_0}$ , i.e. the linear isotropy subgroup  $H^*$  can be identified to  $\text{Ad}_{\mathfrak{m}}H$ . More precisely the linear isometry  $\xi \in \mathfrak{m} \mapsto \xi.p_0 \in T_{p_0}M$  gives rise to an isomorphism from  $O(\mathfrak{m})$  onto  $O(T_{p_0}M)$  which sends  $\text{Ad}_{\mathfrak{m}}H$  onto  $H^*$ . Hence the linear adjoint representation of  $H$  on  $\mathfrak{m}$ :  $g \in H \mapsto \text{Ad}_{\mathfrak{m}}g \in \text{Ad}_{\mathfrak{m}}H$  is an isomorphism (of Lie groups).  $H \cong H^* \cong \text{Ad}_{\mathfrak{m}}H$ .*

### 3.1.1 First convenient hypothesis.

There may be more than one Lie group  $G$  acting symmetrically on a Riemannian symmetric space  $M$ . Besides, we have a convenient way to work on Riemannian symmetric spaces: it is to consider that  $G$  is a subgroup of the group of isometries of  $M$ ,  $\text{Is}(M)$ , which is equivalent to suppose that  $G$  act effectively on  $M$ , i.e.  $H$ , the isotropy subgroup at a fixed point  $p_0$  does not contain non-trivial normal subgroup of  $G$  (see [2]). It is always possible because the kernel  $K$  of the natural morphism  $\phi_H: G \rightarrow \text{Is}(M)$  is the maximal normal subgroup of  $G$  contained in  $H$ <sup>5</sup>, and  $G' = G/K$  acts transitively and effectively on  $M = G/H$  with isotropy subgroup  $H' = H/K$ . Thus  $M = G'/H'$  and since  $K \subset H \subset G^\sigma$ , then  $\sigma$  gives

<sup>5</sup> $K = \ker \phi_H = \ker \rho_{p_0} = \ker \text{Ad}_{\mathfrak{m}}$



rise to an involutive morphism  $\sigma': G' \rightarrow G'$  such that  $(G'^{\sigma'})^0 \subset H' \subset G'^{\sigma'}$ . Now, let us suppose that there exists an order four automorphism  $\tau: G \rightarrow G$  such that  $\sigma = \tau^2$ . Then it gives rise to an isomorphism  $\tau': G/K \rightarrow G/\tau(K)$ . We would like that  $\tau(K) = K$ . It is the case if  $\tau(H) = H$ :  $K$  and  $\tau(K)$  are respectively the maximal normal subgroup of  $G$  contained in  $H$  and  $\tau(H)$  and so if  $\tau(H) = H$  then  $K = \tau(K)$ .

Let us suppose that  $\tau(K) = K$ , then  $\tau$  gives rise to an order four automorphism  $\tau': G/K \rightarrow G/K$  such that  $\sigma' = \tau'^2$ . With our convention we have  $G'_0 = G'^{\tau'} \cap H'$ , then we obtain a 4-symmetric bundle  $N'_{min} = G'/G'_0$  over  $M$ . Hence, when  $G'_0$  describes all the possible choice:  $(G'^{\tau'})^0 \subset G'_0 \subset G'^{\tau'} \cap H'$ , we obtain a family of 4-symmetric bundle  $N' = G'/G'_0$  over  $M$  which are discrete covering of  $N'_{min} = G'/(G'^{\tau'} \cap H')$  and of which  $N'_{max} = G'/(G'^{\tau'})^0$  is a discrete covering. For example, if we choose  $G'_0 = \pi_K(G_0K)$ , we obtain the 4-symmetric bundle over  $M$ ,  $N' = (G/K)/\pi_K(G_0K) = G/G_0K = N/K$ <sup>6</sup>.

Let us come back to the general case (i.e. we do not suppose that  $\tau(K) = K$ ). Since  $\tau(\mathfrak{h}) = \mathfrak{h}$ , we have  $\tau(H^0) = H^0$  and thus denoting by  $K_0$  the maximal normal subgroup of  $G$  contained in  $H^0$  (we have  $K^0 \subset K_0 \subset K \cap H^0$ ), then  $\tau(K_0) = K_0$  for the same reason as above (in particular, if  $K_0 = K$  i.e.  $K \subset H^0$ , then we are in the previous case:  $\tau(K) = K$ ). Hence  $\tau$  gives rise to an order four automorphism  $\tilde{\tau}: G/K_0 \rightarrow G/K_0$  and we are in the case considered above if we consider the symmetric space  $\tilde{M} = G/H^0$  (instead of  $M$ ). Let us precise this point. Indeed  $\tilde{M}$  is a  $(G/K_0)$ -symmetric space and  $\tilde{G} = G/K_0$  acts effectively on it (the isotropy subgroup  $\tilde{H} = H^0/K_0$  does not contain non trivial normal subgroup of  $G/K_0$ ): as above  $\sigma$  gives rise to an involutive automorphism  $\tilde{\sigma}$  of  $\tilde{G} = G/K_0$  such that  $\tilde{H} = (\tilde{G}^{\tilde{\sigma}})^0$  and  $\tilde{\tau}$  is an order four automorphism of  $\tilde{G}/K_0$  such that  $\tilde{\tau}^2 = \tilde{\sigma}$ . Finally, as above we obtain a family of 4-symmetric bundle  $\tilde{N} = \tilde{G}/\tilde{G}_0$  over  $\tilde{M}$  when  $\tilde{G}_0$  describe the set of all possible choice:  $(\tilde{G}^{\tilde{\tau}})^0 \subset \tilde{G}_0 \subset \tilde{G}^{\tilde{\tau}} \cap \tilde{H}$ .

Moreover, the involution  $\tilde{\sigma}$  of  $\tilde{G}/K_0$  gives rise also to the  $\tilde{G}/K_0$ -symmetric space  $\tilde{M}$  (i.e.  $(\tilde{G}^{\tilde{\sigma}})^0 \subset \tilde{H}/K_0 \subset \tilde{G}^{\tilde{\sigma}}$  or equivalently  $\tilde{M}$  belongs to the family of  $\tilde{G}/K_0$ -symmetric spaces defined by  $\tilde{\sigma}$  (of which  $\tilde{M}$  is a discrete covering)).

In the same way, we have  $\tau(G^\sigma) = G^\sigma$  and thus we can do the same as above for the symmetric space  $M_{min} = G/G^\sigma$ .

Nevertheless, in general, it is possible that  $\tau(K) \neq K$  and then  $\tau$  does not give rise to an order four automorphism of  $G' = G/K$  but only to the isomorphism  $\tau': G/K \rightarrow G/\tau(K)$ . However, the tangent map  $T_e\tau' = T_e\tilde{\tau}$  is an order four automorphism of the Lie algebra  $\text{Lie}(G/K) = \text{Lie}(G/\tau(K)) = \text{Lie}(G/K_0) = \mathfrak{g}/\mathfrak{k}$ , and we have  $(T_e\tau')^2 = T_e\sigma'$ , thus  $N/K = (G/K)/\pi_K(G_0K)$  is a locally 4-symmetric bundle over  $M$  ( $\text{Lie } \pi_K(G_0K) = \mathfrak{g}^{T_e\tau}$ ).

Hence we have two good settings to study the Riemannian symmetric spaces

---

<sup>6</sup>In the writing  $N' = N/K$ ,  $K$  does not act freely on  $N$  in general: it is  $K' = K/K \cap G_0$  which acts freely on  $N$  and we have  $N' = N/K = N/K'$ . In particular it is possible that  $N/K = N$  for a non trivial  $K$  (see section 5.3).

$M$  over which a 4-symmetric bundle can be defined, if we want to work only with subgroup of  $\text{Is}(M)$ .

**The first possibility** is to consider that we begin by giving ourself an order four automorphism  $\tau: G \rightarrow G$  and that we always choose the Riemannian symmetric space  $\tilde{M} = G/H$  with  $H = (G^{\tau^2})^0$  (respectively  $M_{min} = G/H$  with  $H = G^{\tau^2}$ ). In other words, in the family of  $G$ -symmetric space corresponding to  $\sigma = \tau^2$  (i.e.  $(G^\sigma)^0 \subset H \subset G^\sigma$ ), we choose the "maximal" one  $\tilde{M} = G/(G^\sigma)^0$ , which is a discrete covering of all the other (respectively the "minimal" one  $M_{min} = G/G^\sigma$ , of which all the other are discrete coverings). Then according to what precede, we can always suppose that  $G$  is a subgroup of  $\text{Is}(\tilde{M})$  (respectively of  $\text{Is}(M_{min})$ ).

**The second possibility** is to work with locally 4-symmetric spaces. In other words we begin by a Riemannian symmetric space over which there exists a locally 4-symmetric bundle. It means that we work with the following setting: a Riemannian symmetric spaces  $M$  with  $G$  a subgroup of  $\text{Is}(M)$  acting symmetrically on  $M$  and a order four automorphism  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$ , such that  $T_1\sigma = \tau^2$ . To define the locally 4-symmetric space  $N$  in this setting, we must tell how we define  $G_0$ . We will set

$$G_0 = \{g \in H \mid \text{Ad}_m g \circ \tau|_{\mathfrak{m}} \circ \text{Ad}_m g^{-1} = \tau|_{\mathfrak{m}}\}. \quad (11)$$

First, we have to verify that if  $\tau$  can be integrated by an automorphism of  $G$ , also denoted by  $\tau$ , then we have  $G_0 = G^\tau \cap H$ . Indeed, if  $g \in G^\tau \cap H$ , then  $\text{Ad}g \circ \tau \circ \text{Ad}g^{-1} = \text{Ad}(g.\tau(g)^{-1}) \circ \tau = \tau$  and since  $\text{Ad}H$  stabilises  $\mathfrak{m}$ , we have  $\text{Ad}_m g \circ \tau|_{\mathfrak{m}} \circ \text{Ad}_m g^{-1} = \tau|_{\mathfrak{m}}$  by taking the restriction to  $\mathfrak{m}$  of the preceding equation. Conversely, suppose that  $g \in H$  and  $\text{Ad}_m g \circ \tau|_{\mathfrak{m}} \circ \text{Ad}_m g^{-1} = \tau|_{\mathfrak{m}}$ , then  $\text{Ad}(g.\tau(g)^{-1}) \circ \tau|_{\mathfrak{m}} = \tau|_{\mathfrak{m}}$  so since  $\tau|_{\mathfrak{m}}$  is surjective,  $\text{Ad}(g.\tau(g)^{-1})|_{\mathfrak{m}} = \text{Id}_{\mathfrak{m}}$  and since the adjoint representation of  $H$  on  $\mathfrak{m}$  is injective (because we suppose that  $G$  is a subgroup of  $\text{Is}(M)$ , and thus  $H$  is a subgroup of  $\text{Is}_{p_0}(M)$ ) it follows that  $g.\tau(g)^{-1} = 1$ . Finally,  $g \in G^\tau \cap H$ . Thus our definition (11) is coherent with our convention which holds when  $\tau$  can be integrated by an automorphism of  $G$ .

Besides, it is easy to see that  $\text{Lie } G_0 = \{a \in \mathfrak{h} \mid \text{ada}|_{\mathfrak{m}} \circ \tau|_{\mathfrak{m}} = \tau|_{\mathfrak{m}} \circ \text{ada}|_{\mathfrak{m}}\} = \mathfrak{g}_0$ . (Indeed,  $\forall a \in \mathfrak{g}_0$ ,  $\text{ada} \circ \tau = \tau \circ \text{ada}$ , and  $\forall a \in \mathfrak{g}_2$ ,  $\text{ada} \circ \tau = -\tau \circ \text{ada}$ , moreover  $\tau|_{\mathfrak{m}} \circ \text{ada}|_{\mathfrak{m}} = 0 \Rightarrow \text{ada}|_{\mathfrak{m}} = 0 \Rightarrow a = 0$  because  $a \in \mathfrak{h} \mapsto \text{ada}|_{\mathfrak{m}}$  is the tangent map of  $h \in H \mapsto \text{Ad}h|_{\mathfrak{m}}$  which is an injective morphism). Hence  $N = G/G_0$  is a locally 4-symmetric bundle over  $M$ .

Further, let  $\pi: \tilde{G} \rightarrow G$  be the universal covering of  $G$ , and  $D = \ker \pi$ . Then  $\tau$  can be integrated by  $\tilde{\tau}: \tilde{G} \rightarrow G$ . Set  $\tilde{\sigma} = \tilde{\tau}^2$ , then  $\sigma \circ \pi = \pi \circ \tilde{\sigma}$  and  $T_1\sigma = T_1\tilde{\sigma} = (T_1\tilde{\tau})^2$ .  $\tilde{G}$  acts almost effectively on  $M$  with isotropy subgroup  $\tilde{H} = \pi^{-1}(H)$  and almost effectively on  $\tilde{M} = \tilde{G}/\tilde{H}^0$  which is the universal covering of  $M$  (see [15]). Besides, if  $\tilde{G}$  does not act effectively on  $\tilde{M}$ , then we take  $D_0$  the maximal normal subgroup of  $\tilde{G}$  included in  $\tilde{H}^0$ , and then we quotient by it, so that we obtain an effective action of  $\tilde{G}/D_0$  on  $\tilde{M}$  and  $\tilde{\tau}$  gives rise to an automorphism of  $\tilde{G}/D_0$ , according to above. Thus we are in the first possibility. Besies it is easy to see that  $\forall g \in \tilde{G}$ ,  $\text{Ad}g = \text{Ad}\pi(g)$  (more precisely

$T_1\pi \circ \text{Ad}g = \text{Ad}\pi(g) \circ T_1\pi$  and we identify  $\tilde{\mathfrak{g}}$  and  $\mathfrak{g}$  so that  $T_1\pi = \text{Id}$ ). Thus  $\tilde{G}_0 = \tilde{G}^{\tilde{\tau}} \cap \tilde{H}^0 = \{g \in \tilde{H}^0 \mid \text{Ad}g \circ \tau_{|\mathfrak{m}} \circ \text{Ad}g^{-1} = \tau_{|\mathfrak{m}}\} \subset \pi^{-1}(G_0)$ . Hence the 4-symmetric space  $\tilde{G}/\tilde{G}_0$  is a discrete covering of the locally 4-symmetric space  $G/G_0$  and we have the following commutative diagram:

$$\begin{array}{ccc} \tilde{G}/\tilde{G}_0 & \longrightarrow & G/G_0 \\ \downarrow & & \downarrow \\ \tilde{M} & \longrightarrow & M. \end{array} \quad (12)$$

**In conclusion**, the two possibilities are equivalent, but we will use the second one because it works with any symmetric space  $M$ , whereas the first needs that we choose a certain covering of  $M$  (for example its universal covering).

**Remark 7** We see that in the preceding reasoning (this using the universal covering  $\tilde{G}$ ) we need only the automorphism of Lie algebra  $\tau$  (and not the symmetric space  $M$ ). Hence, we can consider that we work in the Lie algebra setting and give ourself an order four automorphism  $\tau$  of  $\mathfrak{g}$ . Then we consider the family of associated pairs  $(G, H)$  where  $G$  is a connected Lie group with Lie algebra  $\mathfrak{g}$  and  $H$  a closed Lie subgroup with Lie algebra  $\mathfrak{h} = \mathfrak{g}^\sigma$ . To each such pair corresponds the locally symmetric space  $M = G/H$  and defining  $G_0$  by (11), the locally 4-symmetric bundle  $N = G/G_0$  over  $M$ . Let  $\tilde{G}$  be a simply connected Lie group with Lie algebra  $\mathfrak{g}$ , then  $\tau$  and  $\sigma$  integrates in  $\tilde{G}$  and thus for  $\tilde{H}$  the closed subgroup with Lie algebra  $\mathfrak{h}$ , we can take all subgroups such that  $(\tilde{G}^{\tilde{\sigma}})^0 \subset \tilde{H} \subset \tilde{G}^{\tilde{\sigma}}$  (which implies that  $\tilde{H}$  is closed). If we suppose  $\tilde{H}$  connected, i.e.  $\tilde{H} = (\tilde{G}^{\tilde{\sigma}})^0$ , then  $\tilde{M} = \tilde{G}/\tilde{H}$  is a symmetric space and is also the universal covering of all the locally symmetric spaces  $M = G/H$  when  $(G, H)$  describes all the associated pairs (see [15]), and we have the above commutative diagram between the 4-symmetric bundle  $\tilde{N} = \tilde{G}/\tilde{G}_0$  over  $\tilde{M}$  and the locally 4-symmetric bundle  $N = G/G_0$  over  $M$ . Moreover if  $\tilde{M}$  is Riemannian then all the symmetric spaces  $M = G/H$  when  $(G, H)$  describes all the symmetric associated pairs are Riemannian (see appendix, corollary 3).

**Remark 8** Let us consider  $M$  a  $G$ -symmetric space,  $G \subset \text{Is}(M)$ ,  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  an order four automorphism such that  $\tau^2 = T_1\sigma$ . Then we have  $\tau_{|\mathfrak{m}} \in \Sigma(\mathfrak{m})$  ( $\tau_{|\mathfrak{m}^c} = -i\text{Id}_{\mathfrak{g}_{-1}} \oplus i\text{Id}_{\mathfrak{g}_1}$ ) and it is easy to see that

$$\forall a \in \mathfrak{h}, \quad \tau_{|\mathfrak{h}}(a) = \text{ad}_{\mathfrak{m}}^{-1}(\tau_{|\mathfrak{m}} \circ \text{ad}_{a|\mathfrak{m}} \circ \tau_{|\mathfrak{m}}^{-1}).$$

In other words, under the identification  $\mathfrak{h} \simeq \text{ad}_{\mathfrak{m}}\mathfrak{h} \subset \mathfrak{so}(\mathfrak{m})$ ,  $\tau_{|\mathfrak{h}}$  is the restriction to  $\mathfrak{h}$  of  $\text{Ad}\tau_{|\mathfrak{m}}: \mathfrak{so}(\mathfrak{m}) \rightarrow \mathfrak{so}(\mathfrak{m})$ . Hence  $\tau$  is determined by  $\tau_{|\mathfrak{m}}$ . Besides  $\tau_{|\mathfrak{h}}$  is the tangent map of the isomorphism  $\tau_H$ :

$$\tau_H(g) = \text{Ad}_{\mathfrak{m}}^{-1}(\tau_{|\mathfrak{m}} \circ \text{Ad}_{\mathfrak{m}}g \circ \tau_{|\mathfrak{m}}^{-1}),$$

for  $g \in H^0$  (and more generally for  $g \in \text{Ad}_{\mathfrak{m}}^{-1} \circ (\text{Int}\tau_{|\mathfrak{m}})^{-1} \circ \text{Ad}_{\mathfrak{m}}(H)$ ). Under the identification  $H \simeq \text{Ad}_{\mathfrak{m}}H \subset O(\mathfrak{m})$  it is the restriction to  $H^0$  of the involution  $\text{Int}\tau_{|\mathfrak{m}}: O(\mathfrak{m}) \rightarrow O(\mathfrak{m})$ . According to the definition (11) of  $G_0$ , we have

$G_0 = H^{\tau_H}$ . Besides  $\tau_H(H^0) = H^0$ , thus  $H^0/G_0^0$  is an  $H^0$ -symmetric space. If  $\text{Int}\tau_{|\mathfrak{m}}(\text{Ad}_{\mathfrak{m}}H) = (\text{Ad}_{\mathfrak{m}}H)$ , then  $\tau_H$  is defined in  $H$  and  $\tau_H(H) = H$ , then  $H/G_0$  is an  $H$ -symmetric space (if  $\tau_H(H) \neq H$  it is only a locally symmetric space). Obviously, if  $\tau$  can be integrated in  $G$  then  $\tau_H = \tau_{|H}$ .

**Theorem 7** *Let  $M$  a  $G$ -symmetric space,  $G \subset \text{Is}(M)$ ,  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  an order four automorphism such that  $\tau^2 = T_1\sigma$ . Let  $\text{Aut}(\mathfrak{m})$  be the subgroup of  $O(\mathfrak{m})$  defined by:*

$$\text{Aut}(\mathfrak{m}) = \{F \in O(\mathfrak{m}) \mid F(\text{ad}_{\mathfrak{m}}[v, v'])F^{-1} = \text{ad}_{\mathfrak{m}}[Fv, Fv']\}$$

*it is the subgroup of  $O(\mathfrak{m})$  which leaves invariant  $\text{ad}_{\mathfrak{m}}([\cdot, \cdot]_{|\mathfrak{m} \times \mathfrak{m}}) \in (\Lambda^2 \mathfrak{m}^*) \otimes \mathfrak{so}(\mathfrak{m})$ . Its Lie Algebra*

$$\text{Der}(\mathfrak{m}) = \{A \in \mathfrak{so}(\mathfrak{m}) \mid [A, \text{ad}_{\mathfrak{m}}[v, v']] = \text{ad}_{\mathfrak{m}}[Av, v'] + \text{ad}_{\mathfrak{m}}[v, Av'], \forall v, v' \in \mathfrak{m}\}$$

*is the Lie subalgebra of  $\mathfrak{so}(\mathfrak{m})$  which (acting by derivation) leaves invariant  $\text{ad}_{\mathfrak{m}}([\cdot, \cdot]_{|\mathfrak{m} \times \mathfrak{m}}) \in (\Lambda^2 \mathfrak{m}^*) \otimes \mathfrak{so}(\mathfrak{m})$ .*

*Then  $\tau_{|\mathfrak{m}} \in \text{Aut}(\mathfrak{m})$  and  $\tau$  can be extended in an unique way to the Lie algebra  $\text{Der}(\mathfrak{m}) \oplus \mathfrak{m}$  endowed with the Lie bracket*

$$[(A, v), (A', v')] = ([A, A'] + \text{ad}_{\mathfrak{m}}[v, v'], A.v' - A'.v)$$

*and of which  $\mathfrak{g}$  is a Lie subalgebra, under the inclusion  $a + v \in \mathfrak{h} \oplus \mathfrak{m} \mapsto (\text{ad}_{\mathfrak{m}}a, v) \in \text{Der}(\mathfrak{m}) \oplus \mathfrak{m}$ , by setting*

$$\mathcal{I}_{|\mathfrak{m}} = \tau_{|\mathfrak{m}} \quad \text{and} \quad \mathcal{I}_{|\text{Der}(\mathfrak{m})} = \text{Int}\tau_{|\mathfrak{m}}. \quad (13)$$

*Conversely, given  $\tau_{\mathfrak{m}} \in O(\mathfrak{m})$ , the linear map  $\mathcal{I}$  defined by (13) is an automorphism of the Lie algebra  $\text{Der}(\mathfrak{m}) \oplus \mathfrak{m}$  if and only if  $\tau_{\mathfrak{m}} \in \text{Aut}(\mathfrak{m})$ . Besides it satisfies  $\mathcal{I}^2 = \text{Id}_{\text{Der}(\mathfrak{m})} \oplus -\text{Id}_{\mathfrak{m}}$  (and in particular is of order four) if and only if  $\tau_{\mathfrak{m}} \in \Sigma(\mathfrak{m})$ .*

*Hence, define a locally 4-symmetric bundle over the Riemannian symmetric space  $M$  (which the realisation  $M = G/H$ , i.e.  $\tau$  is an automorphism of  $\mathfrak{g}$  such that  $\tau^2 = T_1\sigma$ ) is equivalent to give ourself  $\tau_{\mathfrak{m}} \in \Sigma(\mathfrak{m}) \cap \text{Aut}(\mathfrak{m})$  such that the order four automorphism  $\mathcal{I}$  of  $\text{Der}(\mathfrak{m}) \oplus \mathfrak{m}$  stabilizes  $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$ , i.e. such that  $\tau_{\mathfrak{m}}(\text{ad}_{\mathfrak{m}}\mathfrak{h})\tau_{\mathfrak{m}}^{-1} = \text{ad}_{\mathfrak{m}}\mathfrak{h}$  (i.e.  $\text{ad}_{\mathfrak{m}}\mathfrak{h}$  is a subalgebra of  $\text{Der}(\mathfrak{m})$  stable by  $\text{Ad}\tau_{\mathfrak{m}}$ ). Then  $\tau = \mathcal{I}_{|\mathfrak{g}}$  is an order four automorphism of  $\mathfrak{g}$  such that  $\tau^2 = \text{Id}_{\mathfrak{h}} \oplus -\text{Id}_{\mathfrak{m}} = T_1\sigma$ .*

**Proof.** First  $\tau_{|\mathfrak{m}} \in \text{Aut}(\mathfrak{m})$ : that follows from the fact that  $\tau$  is an automorphism, so  $\tau \circ \text{ada} \circ \tau^{-1} = \text{ad}\tau(a), \forall a \in \mathfrak{g}$ .

Second,  $\text{Der}(\mathfrak{m}) \oplus \mathfrak{m}$  is a Lie subalgebra. We have to check the Jacobi identity is satisfied. It is a straightforward computation (see [15]). Then we have to check that  $\mathcal{I}$  is an automorphism if and only if  $\tau_{\mathfrak{m}} \in \text{Aut}(\mathfrak{m})$ .

If  $\tau_{\mathfrak{m}} \in \text{Aut}(\mathfrak{m})$  then

- if  $A, A' \in \text{Der}(\mathfrak{m})$ ,  $\mathcal{I}([A, A']) = [\mathcal{I}(A), \mathcal{I}(A')]$  because  $\mathcal{I}_{\text{Der}(\mathfrak{m})} = \text{Int}\tau_{|\mathfrak{m}}$  is an automorphism of  $\text{Der}(\mathfrak{m})$ .

- if  $A \in \text{Der}(\mathfrak{m})$ ,  $v \in \mathfrak{m}$ ,  $\underline{\tau}([A, v]) = \tau_{\mathfrak{m}}(A.v) = \tau_{\mathfrak{m}}A\tau_{\mathfrak{m}}^{-1}(\tau_{\mathfrak{m}}.v) = [\underline{\tau}(A), \underline{\tau}(v)]$
- if  $v, v' \in \mathfrak{m}$ ,  $\underline{\tau}([v, v']) = \text{Int}\tau_{\mathfrak{m}}(\text{ad}_{\mathfrak{m}}[v, v']) = \text{ad}_{\mathfrak{m}}([\tau_{\mathfrak{m}}v, \tau_{\mathfrak{m}}v']) = [\underline{\tau}(v), \underline{\tau}(v')]$   
because  $\tau_{\mathfrak{m}} \in \text{Aut}(\mathfrak{m})$ .

Finally  $\underline{\tau}$  is an automorphism and the unique extension of  $\tau$  (because it is determined by  $\tau|_{\mathfrak{m}}$ , see remark 8).

Conversely if  $\underline{\tau}$  is an automorphism of Lie algebra then

$$\tau_{\mathfrak{m}}\text{ad}_{\mathfrak{m}}([v, v'])\tau_{\mathfrak{m}}^{-1} = (\underline{\tau}\text{ad}_{\mathfrak{m}}([v, v'])\underline{\tau}^{-1})|_{\mathfrak{m}} = (\text{ad}\underline{\tau}([v, v']))|_{\mathfrak{m}} = \text{ad}_{\mathfrak{m}}([\underline{\tau}(v), \underline{\tau}(v')]) = \text{ad}_{\mathfrak{m}}([\tau_{\mathfrak{m}}v, \tau_{\mathfrak{m}}v']).$$

Thus  $\tau_{\mathfrak{m}} \in \text{Aut}(\mathfrak{m})$ .

The last assertion of the theorem follows from what precedes. This completes the proof.  $\blacksquare$

**Remark 9** Let  $\tau_{\mathfrak{m}} \in \Sigma(\mathfrak{m})$  then the condition  $\text{Ad}\tau_{\mathfrak{m}}(\text{ad}_{\mathfrak{m}}\mathfrak{h}) = \text{ad}_{\mathfrak{m}}\mathfrak{h}$  implies that there exists an automorphism  $\tau_{\mathfrak{h}}$  of  $\mathfrak{h}$  defined by  $\forall a \in \mathfrak{h}$ ,  $\text{Ad}\tau_{\mathfrak{m}}(\text{ad}_{\mathfrak{m}}a) = \text{ad}_{\mathfrak{m}}\tau_{\mathfrak{h}}(a)$ , i.e.  $\tau_{\mathfrak{h}} = \text{ad}_{\mathfrak{m}}^{-1} \circ \text{Ad}\tau_{\mathfrak{m}} \circ \text{ad}_{\mathfrak{m}}$ . Then the condition  $\tau_{\mathfrak{m}} \in \text{Aut}(\mathfrak{m})$  is equivalent to

$$\tau_{\mathfrak{h}}([v, v']) = [\tau_{\mathfrak{m}}v, \tau_{\mathfrak{m}}v'], \forall v, v' \in \mathfrak{m}.$$

And obviously, if these two conditions are satisfied then we have  $\tau_{\mathfrak{h}} = \tau|_{\mathfrak{h}}$  (where  $\tau = \underline{\tau}_{\mathfrak{g}}$  is given by the theorem 7).

**Remark 10** Let us consider the map

$$s: g \in \text{Is}_{p_0}(M) \mapsto \text{Ad}_{\mathfrak{m}}g \circ \tau_{\mathfrak{m}} \circ \text{Ad}_{\mathfrak{m}}g^{-1} \in \Sigma(\mathfrak{m})$$

and set  $\underline{G}_0 = \{g \in \text{Is}_{p_0}(M) | s(g) = \tau_{\mathfrak{m}}\}$ .  $\text{Is}_{p_0}(M)$  acts on  $\Sigma(\mathfrak{m})$  by  $g.J = \text{Ad}_{\mathfrak{m}}g \circ J \circ \text{Ad}_{\mathfrak{m}}g^{-1}$  and  $s(g) = g.\tau_{\mathfrak{m}}$ ,  $\underline{G}_0 = \text{Stab}_{\text{Is}_{p_0}(M)}(\tau_{\mathfrak{m}})$ . In the same way, the subgroup  $H = \text{Is}_{p_0}(M) \cap G$  acts on  $\Sigma(M)$  and  $G_0 = \text{Stab}_H(\tau_{\mathfrak{m}})$ . Then  $s(\text{Is}_{p_0}(M)) = \text{Is}_{p_0}(M)/\underline{G}_0$  is a compact submanifold of  $\Sigma(\mathfrak{m})$ , and  $s(H) = H/G_0$  is a relatively compact (immersed) submanifolds of  $\Sigma(\mathfrak{m})$ .

### 3.1.2 Second convenient hypothesis.

An other convenient hypothesis on  $G$  is to consider that it is a closed subgroup of  $\text{Is}(M)$  (and not only a immersed subgroup). It is always possible to work with this hypothesis. Let us make precise this point. Let  $\sigma_{p_0}$  be the symmetry of  $M$  around  $p_0$  (defined by  $\sigma$ ):  $\sigma_{p_0} \in \text{Is}(M)$ ,  $\sigma_{p_0}(p_0) = p_0$  and  $T_{p_0}\sigma_{p_0} = -\text{Id}$ . Then  $\sigma_{p_0}$  belongs to the isotropy subgroup  $\text{Is}_{p_0}(M) = \{f \in \text{Is}(M); f(p_0) = p_0\}$ , and we can define the involution of  $\text{Is}(M)$ :

$$\sigma_{\text{Is}(M)} = \text{Int}(\sigma_{p_0}): g \in \text{Is}(M) \mapsto \sigma_{p_0} \circ g \circ \sigma_{p_0}^{-1} \in \text{Is}(M).$$

It is easy to see that we have

$$(\text{Is}(M)^{\sigma_{\text{Is}(M)}})^0 \subset \text{Is}_{p_0}(M) \subset \text{Is}(M)^{\sigma_{\text{Is}(M)}} \quad (14)$$

(see [15, 2]). The result of this is that  $\sigma: G \rightarrow G$  is the restriction of  $\sigma_{\text{Is}(M)}$  to  $G \subset \text{Is}(M)$  (they induce  $\sigma_{p_0}$  on  $M = G/H$  and the identity on  $H$ , thus, since  $G$  is locally isomorphic to  $M \times H$ , they are identical, see also [15]). Moreover there exists a unique subgroup  $\bar{G}$  of  $\text{Diff}(M)$  such that for any  $G$ -invariant Riemannian metric  $b$  on  $M$ , the group  $\bar{G}$  is the closure of  $G$  in  $\text{Is}(M, b)$ :  $\text{Is}(M, b)$  is closed in  $\text{Diff}(M)$  and so the closure of  $G$  in  $\text{Is}(M, b)$  is its closure in  $\text{Diff}(M)$  and thus it does not depend on  $b$  (see [2, 15]). Then  $\sigma$  extends in a unique way to an involutive morphism  $\bar{\sigma}: \bar{G} \rightarrow \bar{G}$ , which is the restriction of  $\sigma_{\text{Is}(M)}$  to  $\bar{G}$ . Hence denoting by  $\hat{H}$  the isotropy subgroup of  $\bar{G}$  at  $p_0$ ,  $\hat{H} = \text{Is}_{p_0}(M) \cap \bar{G}$ , we have according to (14),  $(\bar{G}^{\bar{\sigma}})^0 \subset \hat{H} \subset \bar{G}^{\bar{\sigma}}$ . Besides  $\bar{\sigma}$  gives rise to the cartan decomposition  $\text{Lie } \bar{G} = \text{Lie } \hat{H} \oplus \mathfrak{m}$ .

In addition to that, we have  $\hat{H} = \bar{H}$ . Indeed, let  $\Phi: U \times \text{Is}_{p_0}(M) \rightarrow \text{Is}(M)$  be a local trivialisation of  $\text{Is}(M) \rightarrow M$ , such that  $\Phi(p_0, h) = h$ , and  $\Phi(U \times H) = \Phi(U \times \text{Is}_{p_0}(M)) \cap G$  (take  $\Phi(p, h) = \phi(p).h$ , with  $\phi: U \rightarrow G$  a local section such that  $\phi(p_0) = 1$ ). Further, if  $g \in \text{Is}_{p_0}(M) \cap \bar{G}$  and  $(g_n)$  is a sequence of  $G \cap \Phi(U \times \text{Is}_{p_0}(M))$  such that  $g_n \rightarrow g$ , then  $\Phi^{-1}(g_n) = (u_n, h_n) \in U \times H$  converges to  $\Phi^{-1}(g) = (p_0, g)$ , thus  $h_n \rightarrow g$  so  $g \in \bar{H}$ .

Moreover,  $\bar{H}$  is a closed subgroup of  $\text{Is}_{p_0}(M)$ , thus it is compact. Hence, we have the symmetric realisation  $M = \bar{G}/\bar{H}$  and  $\text{Ad}_{\mathfrak{m}}(\bar{H})$  is compact: we have showed that the hypothesis  $\text{Ad}_{\mathfrak{m}}(H)$  relatively compact and  $\text{Ad}_{\mathfrak{m}}(H)$  compact gives the same symmetric spaces. Moreover, by using the preceding reasoning (to prove  $\hat{H} = \bar{H}$ ) it is easy to see that if  $\text{Ad}_{\mathfrak{m}}(H)$  is compact then  $G$  is closed in  $\text{Is}(M)$  (see also [15]) so that the hypothesis " $\text{Ad}_{\mathfrak{m}}(H)$  is compact" and " $G$  is closed in  $\text{Is}(M)$ " are in fact equivalent.

Besides, the closure of  $G$  is the same in  $\text{Is}(M)$  and in  $\text{Is}(\tilde{M})$  with  $\tilde{M} = G/H^0$ : since  $M$  and  $\tilde{M}$  are complete (a Riemannian homogeneous space is complete) then  $\text{Is}(M)$  and  $\text{Is}(\tilde{M})$  are complete (see [15]), and thus the closure of  $G$  in one of this group is the completed of  $G$ .

Now, let us suppose that we have a locally 4-symmetric bundle over  $M$ .

**Theorem 8** *Let us consider  $M$  a  $G$ -symmetric space with  $G \subset \text{Is}(M)$  and  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  an order four automorphism such that  $\tau^2 = T_1\sigma$ . Then the extension  $\underline{\tau}$  of  $\tau$ , given by theorem 7 stabilizes the Lie algebra,  $\text{Lie } \bar{G}$ , of the closure of  $G$  in  $\text{Is}(M)$ :*

$$\underline{\tau}(\text{Lie } \bar{G}) = \text{Lie } \bar{G}.$$

*Then denoting by  $\bar{\tau} := \underline{\tau}|_{\text{Lie } \bar{G}}$  the extension of  $\tau$  to  $\text{Lie } \bar{G}$  (given by theorem 7), the subgroup fixed by  $\bar{\tau}$  (defined by (11)) is the closure of  $G_0$ :*

$$\hat{G}_0 := \{g \in \bar{H} \mid \tau|_{\mathfrak{m}} \circ \text{Ad}_{\mathfrak{m}}g \circ \tau|_{\mathfrak{m}}^{-1} = \text{Ad}_{\mathfrak{m}}g\} = \bar{G}_0.$$

*Finally the new locally 4-symmetric bundle over  $M$  defined by  $\bar{\tau}$  is  $\bar{G}/\bar{G}_0$ , and using the notation of remark 10 the fiber of  $\bar{G}/\bar{G}_0$ ,  $\hat{S}_0 := s(\bar{H}) = \bar{H}/\bar{G}_0$  is the closure of the fiber  $S_0$  of  $G/G_0$ :  $\bar{S}_0 = s(\bar{H}) = (\bar{H}/\bar{G}_0)$ , in the maximal fiber over  $M$ :  $\underline{S}_0 := s(\text{Is}_{p_0}(M)) = \text{Is}_{p_0}(M)/\underline{G}_0$ .*

**Theorem 9** *Let us consider  $M$  a Riemannian symmetric space and  $\tilde{M}$  its universal covering.*

- *We have  $R_{p_0}(\cdot, \cdot) = \text{ad}_{\mathfrak{m}}([\cdot, \cdot]_{\mathfrak{m} \times \mathfrak{m}})$  and thus <sup>7</sup>*

$$\begin{aligned} \text{Der}(\mathfrak{m}) &= \mathfrak{Is}_{p_0}(\tilde{M}) \supset \mathfrak{Is}_{p_0}(M) \supset \text{Lie Hol}(M) \\ \text{Aut}(\mathfrak{m}) &\supset \text{Is}_{p_0}(\tilde{M}) \supset \text{Is}_{p_0}(M) \supset \text{Hol}(M) \end{aligned} \quad (15)$$

*(using the identification  $T_{p_0}M = \mathfrak{m}$ ) and  $\text{Der}(\mathfrak{m}) \oplus \mathfrak{m} = \mathfrak{Is}(\tilde{M})$ .*

- *Moreover the following statements are equivalent:*

- $\mathfrak{Is}_{p_0}(\tilde{M}) = \mathfrak{Is}_{p_0}(M)$  (i.e.  $\mathfrak{Is}(\tilde{M}) = \mathfrak{Is}(M)$ )
- $M = M' \times M_0$ , with  $M'$  of the semisimple type (i.e.  $\text{Is}(M')$  is semisimple) and  $M_0$  is Euclidean.
- $h_0 = \mathfrak{so}(\mathfrak{m}_0)$ , where  $h_0$  and  $\mathfrak{m}_0$  are respectively the Euclidean part of  $\mathfrak{Is}_{p_0}(M)$  and  $\mathfrak{m}$  respectively, in the decomposition  $\mathfrak{Is}(M) = \mathfrak{g}' \oplus \mathfrak{g}_0$ , with  $\mathfrak{g}'$  semisimple and  $\mathfrak{g}_0$  of the Euclidean type.

- *Besides the following statements are also equivalent:*

- $\mathfrak{Is}_{p_0}(\tilde{M}) = \mathfrak{Is}_{p_0}(M) \oplus \mathfrak{so}(\mathfrak{m}_0)$
- $h_0 = 0$
- Let  $\tilde{M} = M' \times M_0$  be the decomposition of  $\tilde{M}$  into the semisimple and Euclidean type,  $\Gamma$  the group of deck transformations of the covering  $\pi: \tilde{M} \rightarrow M$ . Then the projection on the Euclidean factor (of  $\text{Is}(\tilde{M}) = \text{Is}(M') \times \text{Is}(M_0)$ ) of  $\Gamma$  satisfies  $\Gamma_0 \cong \mathbb{Z}^r$  with  $r = \dim M_0$  so that  $M_0/\Gamma_0 = \mathbb{T}^r$ .

*Further  $\text{Aut}(\mathfrak{m})$  stabilizes  $\mathfrak{Is}_{p_0}(M)$  if and only if one of the 6 previous statements holds i.e. if and only if  $\mathfrak{Is}(\tilde{M})/\mathfrak{Is}(M) = \{0\}$  or  $\mathfrak{so}(\mathfrak{m}_0)$ . Denoting by  $\text{Aut}(\mathfrak{m})^*$  the subgroup of  $\text{Aut}(\mathfrak{m})$  which stabilizes  $\mathfrak{Is}_{p_0}(M)$ , then the maximal subalgebra of  $\mathfrak{Is}_{p_0}(M)$  stable by  $\text{Aut}(\mathfrak{m})$  is  $\mathfrak{Is}_{p_0}(M)$  if  $\text{Aut}(\mathfrak{m}) = \text{Aut}(\mathfrak{m})^*$  and  $\mathfrak{h}' = \mathfrak{Is}_{p_0}(M')$  if not.*

**Theorem 10** *Let us consider  $M$  a  $G$ -symmetric space with  $G \subset \text{Is}(M)$  and  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  an order four automorphism such that  $\tau^2 = T_1\sigma$ .*

*Then the extension  $\underline{\tau}$  of  $\tau$ , given by theorem 7 define a maximal locally 4-symmetric bundle over  $M$ . Indeed let  $\underline{\mathfrak{g}}$  be the maximal subalgebra of  $\mathfrak{Is}(M)$  stabilized by  $\underline{\tau}$  and  $\underline{G}$  the subgroup of  $\text{Is}(M)$  generated by it. Then  $\underline{G}$  is a closed subgroup of  $\text{Is}(M)$  acting symmetrically on  $M$ , and  $\underline{\tau}|_{\underline{\mathfrak{g}}}$ , define a maximal locally 4-symmetric bundle over  $M$ , with the realisation  $M = \underline{G}/\underline{H}$ .*

*We can also define a minimal locally 4-symmetric bundle over  $M$ , by considering the subalgebra  $\mathfrak{g}' \oplus \mathfrak{m}_0$  (where  $\mathfrak{g}'$  is the semisimple part of  $\mathfrak{Is}(M)$  and  $\mathfrak{m}_0$  the Euclidean part of  $\mathfrak{m}$ ).*

*In conclusion, given any (even dimensional) Riemannian symmetric space  $M$ , define over it a locally 4-symmetric bundle is equivalent to give ourself  $\tau_{\mathfrak{m}} \in \Sigma(\mathfrak{m}) \cap \text{Aut}(\mathfrak{m}) \supset \Sigma(T_{p_0}M) \cap \text{Is}_{p_0}(M)$ . Then the order four automorphism of  $\mathfrak{Is}(\tilde{M})$ ,  $\underline{\tau}$ , defined by  $\tau_{\mathfrak{m}}$ , define the maximal locally 4-symmetric bundle over  $M$ ,  $\underline{N} = \underline{G}/\underline{G}_0$  with  $\underline{G}_0 = \{g \in \underline{H} \mid \tau|_{\mathfrak{m}} \circ \text{Ad}_{\mathfrak{m}g} \circ \tau|_{\mathfrak{m}}^{-1} = \text{Ad}_{\mathfrak{m}g}\}$ . Moreover, any locally*

<sup>7</sup> $\text{Hol}(M)$  is the holonomy group of  $M$

4-symmetric bundle over  $M$  is a subbundle of one such maximal bundle and define such a subbundle  $N$  is equivalent to give oneself a Lie subgroup  $G \subset \text{Is}(M)$  acting symmetrically on  $M$  such that  $\underline{\tau}(\mathfrak{g}) = \mathfrak{g}$  i.e.  $\text{Ad}_{\tau_{\mathfrak{m}}}(\mathfrak{h}) = \mathfrak{h}$ . In this case, the closure  $\bar{N}$  of  $N = G/G_0$  in the (unique) maximal locally 4-symmetric bundle over  $M$ , containing  $N$ ,  $\bar{N}$ , is also a locally 4-symmetric bundle over  $M$  and we have  $\bar{N} = \bar{G}/G_0$ ,  $M = \bar{G}/\bar{H}$ .

**Proof of theorem 8** We have to check that  $\underline{\tau}(\text{Lie } \bar{G}) = \text{Lie } \bar{G}$ , i.e. according to the theorem 7,  $\text{Int}_{\tau_{\mathfrak{m}}}(\text{Lie } \bar{H}) = \text{Lie } \bar{H}$ . We still have  $\text{Int}_{\tau_{\mathfrak{m}}}(H^0) = H^0$ , thus  $\text{Int}_{\tau_{\mathfrak{m}}}(\bar{H}^0) = \bar{H}^0$ . It remains to verify that  $(\bar{H})^0 = \bar{H}^0$ . But this is simply the results of the fact that  $\tilde{M} = \bar{G}/\bar{H}^0$  is a discrete covering of  $M = \bar{G}/\bar{H}$ . Indeed  $(\bar{H})^0$  is closed thus  $(\bar{H})^0 \supset \bar{H}^0$  and then we have

$$\tilde{M} = \bar{G}/\bar{H}^0 \xrightarrow{\text{fibration}} \bar{G}/(\bar{H})^0 \xrightarrow{\text{covering}} \bar{G}/\bar{H} = M$$

and  $\tilde{M} \xrightarrow{\text{covering}} M$ , hence  $(\bar{H})^0/\bar{H}^0$  is discret but the two groups are connected ( $(\bar{H})^0$  is enough) thus  $(\bar{H})^0 = \bar{H}^0$ . We have proved that  $\underline{\tau}(\text{Lie } \bar{G}) = \text{Lie } \bar{G}$ . Using the notation of remark 10, we have, since  $\bar{H}$  is compact,  $s(\bar{H}) = \overline{s(\bar{H})}$ , hence using the same method as for  $\hat{H}$ , we can easily show that  $\hat{G}_0 := s^{-1}(\tau_{\mathfrak{m}}) \cap \bar{H} = \bar{G}_0$  and thus  $s(\bar{H}) = \bar{H}/\bar{G}_0$ . Finally, the new locally 4-symmetric space is  $\bar{G}/\bar{G}_0$ . This completes the proof. ■

**Proof of theorem 9** For the first point see [15]. For the following points, see section 4.1 and 4.2. ■

**Proof of theorem 10** First  $\underline{G}$  is closed : it a immediate consequence of the maximality and theorem 8. Then we have to prove that  $\bar{G}/\bar{G}_0$  is the closure  $\bar{N}$  of  $N = G/G_0$  in  $\underline{G}/\underline{G}_0$ . Let  $\pi_{J_0}: \underline{G} \rightarrow \underline{G}/\underline{G}_0$  be the projection map, then we have  $\pi_{J_0}(G) = \underline{G}/\underline{G}_0 \cap G = G/G_0$  (according to definition (11)) and thus  $\pi_{J_0}(\bar{G}) \subset \overline{\pi_{J_0}(G)} = \bar{N}$  but  $\pi_{J_0}(\bar{G}) = \bar{G}/\underline{G}_0 \cap \bar{G} = \bar{G}/\bar{G}_0$  (according to definition (11) and  $\hat{G}_0 = \bar{G}_0$ ). Hence  $\bar{G}/\bar{G}_0 \subset \bar{N}$ . These are together subbundle (over  $M$ ) of  $\bar{N}$  and using a trivialisation of  $\bar{N} = \underline{G}/\underline{G}_0 \rightarrow M$  (same reasoning as for  $\hat{H}$ ) it is easy to see that the fiber of  $\bar{N}$  (over  $p_0$ ) is  $\bar{H}/\bar{G}_0$  which implies that  $\bar{G}/\bar{G}_0 = \bar{N}$ . This completes the proof. ■

**Remark 11** In particular, if we suppose that we have an order four automorphism of  $G$ ,  $\sigma = \tau^2$ . Then since  $\tau$  is uniformly continuous, it extends into an order four automorphism  $\bar{\tau}: \bar{G} \rightarrow \bar{G}$  (because  $\text{Is}(M)$  is complete) and obviously  $\bar{\sigma} = \bar{\tau}^2$ .

**Remark 12** According to the definition (11),  $\tau_{\mathfrak{m}}$  and  $-\tau_{\mathfrak{m}}$  gives rise to the same group  $G_0$ . Moreover  $\tau_{\mathfrak{m}} = (\tau^{-1})_{\mathfrak{m}}$  and in particular if  $\tau$  integrates in  $G$  then  $G^\tau = G^{\tau^{-1}}$ . Besides  $(\tau^{-1})^2 = (T_1\sigma)^{-1} = T_1\sigma$ , hence  $\tau^{-1}$  define the same locally 4-symmetric bundle over  $M$  as  $\tau$ . Moreover, given any  $\tau_{\mathfrak{m}} \in \Sigma(\mathfrak{m}) \cap \text{Aut}(\mathfrak{m})$ , then  $-\tau_{\mathfrak{m}} \in \Sigma(\mathfrak{m}) \cap \text{Aut}(\mathfrak{m})$  and gives rise (according to theorem 7) to the automorphism  $\underline{\tau}^{-1}$  which gives rise to the same maximal locally 4-symmetric bundle over  $M$  and the same family of 4-symmetric subbundle over  $M$ .



From now, we will always suppose that  $G$  is a closed subgroup of  $\text{Is}(M)^0$ . The result of this is that the isotropy subgroup of  $G$  at the point  $p_0$ ,  $H = \text{Stab}_G(p_0)$  is compact and can be identified (via the adjoint representation on  $\mathfrak{m}$ , resp. via the linear isotropy representation) to a closed subgroup of  $O(\mathfrak{m})$  (resp. of  $O(T_{p_0}M)$ ). Then according to theorem 10, to study the case of non closed subgroup of  $\text{Is}(M)^0$  (or equivalently the non closed locally 4-symmetric bundle over  $M$ ), we have just to consider the non closed subgroups of our closed group  $G$ , acting symmetrically on  $M$ , and whose Lie algebra is stable by  $\tau$ .

### 3.2 Twistor subbundle

We give ourself a locally 4-symmetric bundle  $N = G/G_0$  (defined by an order four automorphism  $\tau$  and by (11)) over a symmetric space  $M = G/H$ . We will show that  $G/G_0$  is a subbundle of the twistor bundle  $\Sigma(G/H)$ . Under the isomorphism between  $TM$  and  $[\mathfrak{m}] = \{(g.p_0, \text{Ad}g(\xi)), \xi \in \mathfrak{m}, g \in G\}$ ,  $T_{p_0}M$  is identified to  $\mathfrak{m}$ :  $\xi \in \mathfrak{m} \mapsto \xi.p_0 \in T_{p_0}M$  is an isomorphism of vector spaces. Then to  $\tau|_{\mathfrak{m}} \in \Sigma(\mathfrak{m})$  (resp. to  $-\tau|_{\mathfrak{m}} = \tau|_{\mathfrak{m}}^{-1} \in \Sigma(\mathfrak{m})$ ) corresponds  $J_0 \in \Sigma(T_{p_0}M)$ , and more generally to  $\text{Ad}g \circ \tau|_{\mathfrak{m}} \circ \text{Ad}g^{-1} \in \Sigma(\text{Ad}g(\mathfrak{m}))$  (resp.  $\text{Ad}g \circ \tau|_{\mathfrak{m}}^{-1} \circ \text{Ad}g^{-1}$ ) corresponds  $gJ_0g^{-1} \in \Sigma(T_{g.p_0}M)$ . Thus we have defined a map

$$\begin{aligned} \rho_{J_0}: \quad G &\longrightarrow \Sigma(M) \\ g &\longmapsto gJ_0g^{-1} \end{aligned}$$

which according to the definition (11) of  $G_0$  gives rise under quotient to the injective map:

$$\begin{aligned} i: \quad G/G_0 &\longrightarrow \Sigma(M) \\ g.G_0 &\longmapsto gJ_0g^{-1}. \end{aligned}$$

Moreover,  $i$  is an embedding. Indeed,  $G$  acts smoothly on the manifolds  $\Sigma(M)$  and so the map  $g \in G \mapsto gJ_0g^{-1} \in \Sigma(M)$  is a subimmersion of constant rank. Thus  $i: G/G_0 \rightarrow \Sigma(M)$  is an injective subimmersion and so it is an injective immersion. We can add that  $i: G/\text{Stab}_G(J_0) \rightarrow G.J_0$  is a homeomorphism if the orbit  $G.J_0$  is locally closed in  $\Sigma(M)$  (see [5]). We will show directly that  $i(G/G_0) = G.J_0$  is a subbundle of  $\Sigma(M)$ .

First, let us precise the fibration  $G/G_0 \rightarrow G/H$ . We have the isomorphism of bundle:  $G/G_0 \cong G \times_H H/G_0$ . In particular, the fiber type of  $G/G_0$  is  $H/G_0$ . Besides  $i$  is a morphism of bundle (over  $M$ ). Since  $i$  is also an injective immersion, we can identify the fibers of  $G/G_0$  and  $i(G/G_0)$  respectively over the point  $g.p_0 \in M$ . The fiber of  $i(G/G_0)$  over  $p = g.p_0$  is  $gS_0g^{-1}$  where  $S_0 = \text{Int}(H)(J_0) \subset \Sigma(T_{p_0}M)$  is the fiber over  $p_0$ .<sup>8</sup>

Now let us show that  $i(G/G_0)$  is a subbundle of  $\Sigma(M)$ . Let  $\sigma: U \subset G/H \rightarrow G$  be a local section of the fibration  $\pi_H: G \rightarrow G/H$ . Then we have the following trivialisation of  $\Sigma(U)$ :

$$\Phi: (p, J) \in U \times \Sigma(T_{p_0}M) \longmapsto (p, \sigma(p)J\sigma(p)^{-1}) \in \Sigma(U)$$

<sup>8</sup>we remark that  $H \subset O(T_{p_0}M)$ ,  $G_0 \subset U(T_{p_0}M, J_0)$  and  $S_0 = H/G_0$  is a compact submanifold of  $\Sigma(T_{p_0}M)$ .

and we have  $\Phi(U \times S_0) = \bigsqcup_{p \in U} \{p\} \times (\sigma(p)S_0\sigma(p)^{-1}) = i(G/G_0) \cap \Sigma(U)$ . Thus  $i(G/G_0)$  is a subbundle of  $\Sigma(M)$ , hence  $i$  is an embedding.

Let us recapitulate what precede:

**Theorem 11** *The map*

$$\begin{aligned} i: G/G_0 &\longrightarrow \Sigma(M) \\ g.G_0 &\longmapsto gJ_0g^{-1}. \end{aligned}$$

*is an embedding and a morphism of bundle from  $G/G_0$  into  $\Sigma(M)$ . Besides the fiber of  $i(G/G_0)$  over the point  $p = g.p_0$  is  $gS_0g^{-1}$ , with  $S_0 = \text{Int}(H)(J_0)$  and  $J_0 \in \Sigma(T_{p_0}M)$  corresponding to  $\tau_{\mathfrak{m}} \in \Sigma(\mathfrak{m})$  (resp. to  $\tau_{\mathfrak{m}}^{-1}$ ).*

**Remark 13** If we consider a locally 4-symmetric bundle  $N = G/G_0$  over  $M$ , with  $G_0$  not defined by (11), then  $i$  is not injective in general: to obtain a injective map  $i$ , we must consider the locally 4-symmetric space  $G/\pi_K^{-1}(G'_0) = (G/K)/G'_0$  where  $K = \ker \text{Ad}_{\mathfrak{m}}$  and  $G'_0$  is the subgroup of  $H' = H/K$  defined by (11). In particular, we see that in general a 4-symmetric space  $G/G_0$  is not a submanifold of a twistor space (see section 5). Moreover, we can see the aim of our definition (11) (and in particular of our convention  $G_0 = G^\tau \cap H$ ): it is to obtain a injective map  $i$ .

**Remark 14** For any covering  $\pi: \tilde{G} \rightarrow G$ ,  $\tilde{G}$  acting symmetrically on  $M$ , we have  $\tilde{\rho}_{J_0}(\tilde{G}) = \rho_{J_0}(G) = i_{J_0}(G/G_0)$ : the locally 4-symmetric subbundle of  $\Sigma(M)$ ,  $i_{J_0}(G/G_0)$  does not depend on the choice of the group  $G$  (we have chosen for  $G$ , the subgroup of  $\text{Is}(M)$  generated by  $\mathfrak{g}$ ).

Moreover,  $\rho_{J_0}(\underline{G}) = i_{J_0}(\underline{N})$  is a maximal locally 4-symmetric subbundle in  $\Sigma(M)$ . Now, suppose that we work with a non closed subgroup  $G' \subset \text{Is}(M)$ , then  $\rho_{J_0}(G') = i_{J_0}(G'/G'_0)$  is an immersed subbundle in  $\Sigma(M)$ :  $\Phi(U \times S'_0) = \bigsqcup_{p \in U} \{p\} \times (\sigma(p)S'_0\sigma(p)^{-1}) = i(G'/G'_0) \cap \Sigma(U)$  but the fiber  $S'_p = \sigma(p)S'_0\sigma(p)^{-1}$  is only a (non closed relatively compact) immersed submanifold in  $\Sigma(T_pM)$ . And since  $i_{J_0}$  is an embedding (from  $\underline{N}$  into  $\Sigma(M)$ ) we have  $i_{J_0}(\overline{G'}/\overline{G'_0}) = i_{J_0}(\overline{N'}) = i_{J_0}(\overline{N'})$ . In others words, taking the closure of  $G'$  in  $\text{Is}(M)$  is equivalent to take the closure of  $N'$  in  $\underline{N}$  according to theorem 10 which is equivalent to take the closure of  $i_{J_0}(N')$  in  $i_{J_0}(\underline{N})$ .

**Remark 15** The maximal locally 4-symmetric bundles  $\underline{N}$  are disjoint: these are orbits in  $\Sigma(M)$ . More precisely these are suborbits of  $\text{Is}(M)^0$ -orbits in the form  $\underline{G} \cdot J_0$  in  $\Sigma(M) \cap \text{Aut}(M)$  with  $\text{Aut}(M) = \bigsqcup_{p \in M} p \times \text{Aut}(T_pM)$  (see section 4.1 and 4.2). In particular,  $\Sigma(M) \cap \text{Aut}(M)$  is the disjoint union of all the maximal locally 4-symmetric bundles over  $M$ . Moreover, the set of maximal locally 4-symmetric bundles over  $M$ ,  $\mathcal{N}(M)$ , contains the subset  $\mathcal{N}^*(M)$  of elements which are  $\text{Is}(M)^0$ -orbits, i.e. elements  $\rho_{J_0}(\text{Is}(M)^0)$  with  $J_0 \in \Sigma(\mathfrak{m}) \cap \text{Aut}^*(\mathfrak{m})$ :  $\mathcal{N}^*(M) = \Sigma(M) \cap \text{Aut}^*(M)/\text{Is}(M)^0 \subset \Sigma(M)/\text{Is}(M)^0$ .

**Remark 16** The Riemannian manifold  $M = G/H$  is orientable if and only if  $\text{Ad}_m H \subset SO(\mathfrak{m})$  (or equivalently  $H^* \subset SO(T_{p_0} M)$ ). Besides  $\tau_m \in \Sigma^\varepsilon(\mathfrak{m})$ , and to fix ideas, let us suppose that  $\varepsilon = 1$ . Then, if  $M$  is orientable,  $i$  is an embedding from  $G/G_0$  into  $\Sigma^+(M)$ . Moreover, if we work with  $\tilde{M} = G/\tilde{H}^0$ , we are sure that  $H^0 \subset SO(T_{p_0} \tilde{M})$ . Hence, if we work with what we called the first possibility (see section 3.1.1),  $i$  takes values in  $\Sigma^+(\tilde{M})$ . In other words, given a locally 4-symmetric bundle over  $M$ , the corresponding 4-symmetric bundle over its universal covering  $\tilde{M}$  (see remark 7) is embedded in  $\Sigma^+(\tilde{M})$ .

Let us consider more generally any covering  $\pi: \tilde{M} \rightarrow M$  then it induces the covering  $\pi_\Sigma: \Sigma(\tilde{M}) \rightarrow \Sigma(M)$  which is also a morphism of bundle over  $\pi: \tilde{M} \rightarrow M$ . It is defined by

$$\pi_\Sigma: J_{\tilde{x}} \in \Sigma(T_{\tilde{x}} \tilde{M}) \mapsto T_{\tilde{x}} \pi \circ J_{\tilde{x}} \circ (T_{\tilde{x}} \pi)^{-1} \in \Sigma(T_x M).$$

Now, let us suppose that  $\pi$  comes from a covering  $\tilde{\pi}: \tilde{G} \rightarrow G$  and that we have  $M = G/H$ ,  $\tilde{M} = \tilde{G}/\tilde{H}^0$  (symmetric realisation) with  $\tilde{H} = \tilde{\pi}^{-1}(H)$  and  $G \subset \text{Is}(M)$ ,  $\tilde{G} \subset \text{Is}(\tilde{M})$  (see above). Then we have

$$T_{\tilde{x}} \pi \circ (\tilde{g} J_{\tilde{p}_0} \tilde{g}^{-1}) \circ (T_{\tilde{x}} \pi)^{-1} = g J_{p_0} g^{-1}.$$

with  $\tilde{x} = \tilde{g} \cdot \tilde{p}_0$ ,  $g = \tilde{\pi}(\tilde{g})$ . Hence the restriction<sup>9</sup> of  $\pi_\Sigma$  to  $\tilde{G}/\tilde{G}_0$  gives rise to the morphism of bundle (12). Moreover<sup>10</sup>

$$S_0 = \text{Int}(H)(J_0) = \bigcup_{h \in \tilde{H}/\tilde{H}^0} h \tilde{S}_0 h^{-1}$$

with, since  $\tilde{H}^0 \subset SO(T_{p_0} \tilde{M})$ ,  $\tilde{S}_0 \subset \Sigma^+(T_{p_0} \tilde{M})$ . Further if  $H \subset O(T_{p_0} M)$  is not included in  $SO(T_{p_0} M)$  (i.e.  $M$  is not orientable), then we have

$$\pi_\Sigma(\Sigma^+(\tilde{M})) = \Sigma(M).$$

**Remark 17** Let us see what happens when we change  $M$ , in theorem 4. Let  $\tilde{G}$  be a covering of  $G$ , acting symmetrically on a covering  $\tilde{M}$  of  $M$ ,  $\pi: \tilde{M} \rightarrow M$ , with  $\tilde{G} \subset \text{Is}(\tilde{M})$ . Then according to remark 16, we have  $\pi_\Sigma \circ i_{J_{\tilde{p}_0}} = i_{J_{p_0}} \circ \pi_0$ , with  $\pi_0: \tilde{G}/\tilde{G}_0 \rightarrow G/G_0$  the morphism of bundle (over  $\pi: \tilde{M} \rightarrow M$ ) given by (12). Then given any solution  $\alpha$  of the  $(\mathfrak{g}, \tau)$ -system (6), let us integrate it in  $\tilde{G}$  and  $G$  respectively,  $\tilde{U}: L \rightarrow \tilde{G}$ ,  $U: L \rightarrow G$  with  $\tilde{U}(0) = 1$ ,  $U(0) = 1$  (0 is a reference point in  $L$ ), we have  $\tilde{\pi} \circ \tilde{U} = U$ . Then let us project these lifts in  $\tilde{G}/\tilde{G}_0$  and  $G/G_0$  respectively: we obtain the geometric solutions  $\tilde{J}: L \rightarrow \tilde{G}/\tilde{G}_0$  and  $J: L \rightarrow G/G_0$  respectively and we have  $\pi_0 \circ \tilde{J} = J$ . Then let us embed these into the twistor spaces  $\Sigma(\tilde{M})$  and  $\Sigma(M)$  to obtain the admissible twistor lifts  $\tilde{J}_{\tilde{X}}: L \rightarrow i_{\tilde{j}_0}(\tilde{G}/\tilde{G}_0)$  and  $J_X: L \rightarrow i_{j_0}(G/G_0)$  respectively which are related by  $\pi_\Sigma \circ \tilde{J}_{\tilde{X}} = J_X$ , and in particular  $\pi \circ \tilde{X} = X$ .

<sup>9</sup> i.e.  $\pi_\Sigma \circ i_{J_{\tilde{p}_0}} = i_{J_{p_0}} \circ \pi_0$ ,  $\pi_0: \tilde{G}/\tilde{G}_0 \rightarrow G/G_0$  given by (12).

<sup>10</sup> In fact,  $h \tilde{S}_0 h^{-1}$  means obviously  $T_{h \cdot p_0} \pi \circ (h \tilde{S}_0 h^{-1}) \circ T_{h \cdot p_0} \pi^{-1}$ .  $\pi_\Sigma$  allows to consider the fibers  $\Sigma(T_{\tilde{x}} \tilde{M})$  as included in the fiber  $\Sigma(T_x M)$ , with  $x = \pi(\tilde{x})$ .

## 4 Splitting of $M$ into the 3 type of symmetric space

**Theorem 12** [15, 2] *Let  $M$  be a simply connected Riemannian symmetric space. Then  $M$  is a product*

$$M = M_0 \times M_- \times M_+$$

where  $M_0$  is an Euclidean space,  $M_-$  and  $M_+$  are Riemannian symmetric space of the compact and noncompact type respectively. In particular

$$M = M_0 \times M'$$

where  $M'$  has a group of isometries  $G = \text{Is}(M')$  semisimple and its isotropy subgroup at  $p_0 \in M'$ ,  $H$ , (which is connected because  $M'$  is simply connected) is equal to the holonomy group of  $M'$ . Hence a Riemannian symmetric space  $M$  of which the isometry group is semisimple (which is equivalent to say that its universal covering have not Euclidean factor, or equivalently the Lie algebra of  $G$  does not contain no trivial abelian ideal, i.e. its Killing form is no degenerate) has a unique symmetric writing  $G/H$ , with  $G$  acting effectively. In this unique realisation, we have necessarily  $G = \text{Is}(M)^{0 \ 11}$  and  $H = \text{Is}_{p_0}^0(M) = \text{Is}_{p_0}(M) \cap \text{Is}(M)^0 (\supset \text{Is}_{p_0}(M)^0)$ . Further the Lie algebra  $\mathfrak{Is}_{p_0}(M) = \text{Der}(\mathfrak{m}) = \mathfrak{Sol}(M)$  is spanned by  $[\mathfrak{m}, \mathfrak{m}] = \{R_{p_0}(X, Y), X, Y \in T_{p_0}(M)\}$ .

Moreover the covering of such a Riemannian symmetric space  $M$ , has a decomposition into a product of irreducible Riemannian symmetric spaces (i.e the linear isotropy representation is irreducible)

$$\tilde{M} = M_1 \times \cdots \times M_r.$$

**Theorem 13** *Let us consider the decomposition of  $(\mathfrak{g}, T_1\sigma)$  into the sum of orthogonal (for the Killing form) ideals of the compact, noncompact and Euclidean type respectively:*

$$\mathfrak{g} = \mathfrak{l}_0 \oplus \mathfrak{l}_- \oplus \mathfrak{l}_+$$

and let  $\mathfrak{l}_\alpha = \mathfrak{h}_\alpha \oplus \mathfrak{m}_\alpha$  be the eigenspace decomposition of the involution  $T_1\sigma|_{\mathfrak{l}_\alpha}$ . Suppose now that we have an order four automorphism  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  with  $\tau^2 = T_1\sigma$ . Then  $\tau(\mathfrak{l}_\alpha) = \mathfrak{l}_\alpha$ ,  $\tau(\mathfrak{h}_\alpha) = \mathfrak{h}_\alpha$ ,  $\tau(\mathfrak{m}_\alpha) = \mathfrak{m}_\alpha$  for  $\alpha = 0, -, +$ . Hence  $\tau_{\mathfrak{m}} = \bigoplus_{\alpha} \tau_{\mathfrak{m}_\alpha}$ , with  $\tau_{\mathfrak{m}_\alpha} \in \Sigma(\mathfrak{m}_\alpha)$ , and  $\tau|_{\mathfrak{l}_\alpha}$  is the automorphism of  $\mathfrak{l}_\alpha$  defined by  $\tau_{\mathfrak{m}_\alpha}$  according to theorem 7 and we have  $\tau_{\mathfrak{l}_\alpha}^2 = T_1\sigma|_{\mathfrak{l}_\alpha}$ . Moreover, we have  $\text{Aut}(\mathfrak{m}) = \prod_{\alpha} \text{Aut}(\mathfrak{m}_\alpha)$ .

**Corollary 1** *Let  $M$  be a  $G$ -symmetric space,  $G \subset \text{Is}(M)$  and  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  an order four automorphism with  $\tau^2 = T_1\sigma$ . Let  $\tilde{M}$  be its universal covering, which has a symmetric realisation  $\tilde{M} = \tilde{G}/\tilde{H}^0$ , with  $\pi: \tilde{G} \rightarrow G$  a covering of  $G$ ,  $\tilde{H} = \pi^{-1}(H)$  and  $\tilde{G} \subset \text{Is}(\tilde{M})$ , such that  $\tau$  integrates into  $\tilde{\tau}: \tilde{G} \rightarrow \tilde{G}$ . Then*

<sup>11</sup>as usual, we suppose that  $G$  is connected

the decomposition of  $\mathfrak{g}$  into 3 ideals of different type gives rise to the following decomposition of  $\tilde{G}$ :

$$\tilde{G} = L_0 \times L_- \times L_+$$

which induces the following decomposition of  $\tilde{H}^0$  and  $\tilde{G}_0 = \tilde{H}^0 \cap \tilde{G}^{\tilde{\tau}}$ , corresponding also to the decomposition  $\mathfrak{h} = \oplus_{\alpha} \mathfrak{h}_{\alpha}$  and  $\mathfrak{g}_0 = \oplus_{\alpha} (\mathfrak{g}_0)_{\alpha}$ :

$$\tilde{H}^0 = H_0 \times H_- \times H_+ \quad (16)$$

$$\tilde{G}_0 = (G_0)_0 \times (G_0)_- \times (G_0)_+. \quad (17)$$

Hence  $\tilde{M} = M_0 \times M_- \times M_+$  and  $\tilde{N} = N_0 \times N_- \times N_+$  with  $M_{\alpha} = L_{\alpha}/H_{\alpha}$ ,  $N_{\alpha} = L_{\alpha}/(G_0)_{\alpha}$ . Besides  $\tilde{\sigma}$  and  $\tilde{\tau}$  have the decompositions  $\tilde{\sigma} = \prod_{\alpha} \tilde{\sigma}_{\alpha}$  and  $\tilde{\tau} = \prod_{\alpha} \tilde{\tau}_{\alpha}$ , and  $H_{\alpha} = (\tilde{G}^{\tilde{\sigma}_{\alpha}})^0$ ,  $(G_0)_{\alpha} = H_{\alpha} \cap L_{\alpha}^{\tau_{\alpha}} = (G_{\alpha})_0$ .  $N_{\alpha}$  is a 4-symmetric bundle over  $M_{\alpha}$ .

**Theorem 14** Let us consider the decomposition of  $(\mathfrak{g}, T_1\sigma)$  into the sum of orthogonal (for the Killing form) ideals:

$$\mathfrak{g} = \oplus_{i=0}^r \mathfrak{g}_i \quad (18)$$

with  $\mathfrak{g}_0$  abelian and  $(\mathfrak{g}_i, T_1\sigma|_{\mathfrak{g}_i})$  irreducible, and let  $\mathfrak{g}_i = \mathfrak{h}_i \oplus \mathfrak{m}_i$  be the eigenspace decomposition of  $T_1\sigma|_{\mathfrak{g}_i}$ . Suppose now that we have an order four automorphism  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  such that  $\tau^2 = T_1\sigma$ .

There exists a unique decomposition of  $\mathfrak{g}$ :

$$\mathfrak{g} = \mathfrak{g}_0 \oplus (\oplus_{i=1}^{r'} \mathfrak{g}'_i) \quad (19)$$

where  $\mathfrak{g}'_i = \mathfrak{g}_i$  or  $\mathfrak{g}'_i = \mathfrak{g}_i \oplus \mathfrak{g}_j$  with  $(\mathfrak{g}_i, T_1\sigma|_{\mathfrak{g}_i})$  and  $(\mathfrak{g}_j, T_1\sigma|_{\mathfrak{g}_j})$  isomorphic, such that  $\tau(\mathfrak{g}'_i) = \mathfrak{g}'_i$ ,  $\tau(\mathfrak{h}'_i) = \mathfrak{h}'_i$ ,  $\tau(\mathfrak{m}'_i) = \mathfrak{m}'_i$ . Moreover if  $\mathfrak{g}'_i = \mathfrak{g}_i \oplus \mathfrak{g}_j$  then  $\tau(\mathfrak{g}_i) = \mathfrak{g}_j$ ,  $\tau(\mathfrak{h}_i) = \mathfrak{h}_j$ ,  $\tau(\mathfrak{m}_i) = \mathfrak{m}_j$ . Hence  $\tau_{\mathfrak{m}} = \oplus_{i=0}^{r'} \tau_{\mathfrak{m}'_i}$  with  $\tau_{\mathfrak{m}'_i} \in \Sigma(\tau_{\mathfrak{m}'_i})$ , and  $\tau|_{\mathfrak{g}'_i}$  is the automorphism of  $\mathfrak{g}'_i$  defined by  $\tau_{\mathfrak{m}'_i}$  according to theorem 7 and we have  $\tau|_{\mathfrak{g}'_i}^2 = T_1\sigma|_{\mathfrak{g}'_i}$ .

**Corollary 2** Let  $M$  be a  $G$ -symmetric space,  $G \subset \text{Is}(M)$  and  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  an order four automorphism with  $\tau^2 = T_1\sigma$ . Let  $\tilde{M}$  be its universal covering, which has a symmetric realisation  $\tilde{M} = \tilde{G}/\tilde{H}^0$ , with  $\pi: \tilde{G} \rightarrow G$  a covering of  $G$ ,  $\tilde{H} = \pi^{-1}(H)$  and  $\tilde{G} \subset \text{Is}(\tilde{M})$ , such that  $\tau$  integrates into  $\tilde{\tau}: \tilde{G} \rightarrow \tilde{G}$ . Then the decomposition of  $\mathfrak{g}$ , (18), gives rise to the following decomposition of  $\tilde{G}$ :

$$\tilde{G} = L_0 \times L_1 \times \cdots \times L_r$$

which induces the following decomposition of  $\tilde{H}^0$ , corresponding also to the decomposition  $\mathfrak{h} = \oplus_{i=0}^r \mathfrak{h}_i$ :

$$\tilde{H}^0 = H_0 \times H_1 \times \cdots \times H_r$$

$\tilde{\sigma}$  have the decomposition of  $\tilde{\sigma} = \prod_{i=0}^r \tilde{\sigma}_i$  with  $\tilde{\sigma}_i$  involution of  $L_i$  and  $H_i = (\tilde{G}^{\tilde{\sigma}_i})^0$ . Moreover there exists an unique decomposition of  $\tilde{G}$ :

$$\tilde{G} = L'_0 \times L_1 \times \cdots \times L'_r, \quad (20)$$

where  $L'_i = L_i$  or  $L'_i = L_i \times L_j$  with  $(L_i, \tilde{\sigma}_i)$  and  $(L_j, \tilde{\sigma}_j)$  isomorphic. Then  $\tau$  have the decomposition  $\tilde{\tau} = \prod_{i=0}^{r'} \tilde{\tau}'_i$  with  $\tilde{\tau}'_i$  order four automorphism of  $L'_i$ . Further, by identifying  $(L_i, \tilde{\sigma}_i)$  and  $(L_j, \tilde{\sigma}_j)$  (when  $L'_i = L_i \times L_j$ ), then in 20, we have either  $L'_i = L_i$  and then  $\tilde{\tau}'_i = \tilde{\tau}_i$  is an order four automorphism of  $L_i$  so that  $(L'_i)^{\tilde{\tau}'_i} = (L_i)^{\tilde{\tau}_i}$ , or  $L'_i = L_i \times L_i$  and then

$$\tilde{\tau}'_i: (a, b) \in L_i \times L_i \mapsto (\sigma_i(b), a) \in L_i \times L_i$$

so that  $(L'_i)^{\tilde{\tau}'_i} = \Delta(H_i) \subset H_i \times H_i$ . Hence  $\tilde{M} = M_0 \times M_1 \times \dots \times M_r$  with  $M_i = L_i/H_i$  and  $\tilde{N} = N'_0 \times N'_1 \times \dots \times N'_r$ , where either  $N'_i = N_i = L_i/(G_i)_0$  is a 4-symmetric bundle over  $M_i$ , or  $N'_i = G_i \times G_i/\Delta(H_i)$  is a 4-symmetric bundle over  $M_i \times M_i = G_i \times G_i/H_i \times H_i$  (and the fiber  $H_i \times H_i/\Delta(H_i) \simeq H_i$  is a group).

**Proof of theorems 13,14 and corollaries 1,2** Use the facts that  $\tau_m$  leaves invariant the metric in  $\mathfrak{m}$  and the restriction to  $\mathfrak{m}$  of the killing form. ■

#### 4.1 The semisimple case

**Definition 6** We will say that the Riemannian symmetric space  $M$  is of semisimple type if  $\text{Is}(M)$  is semisimple.

**Theorem 15** If  $M$  is of semisimple type then each (connected) locally 4-symmetric bundle over  $M$  is maximal and in the form  $\underline{N}^0 = \text{Is}(M)^0/\underline{G}_0$ , i.e. is an  $\text{Is}(M)^0$ -orbit in  $\Sigma(M) \cap \text{Aut}(M)$ . In other words the set of locally 4-symmetric bundles over  $M$  is  $\mathcal{N}(M) = \Sigma(M) \cap \text{Aut}(M)/\text{Is}(M)^0 \subset \Sigma(M)/\text{Is}(M)^0$ .

**Remark 18** The "size" of a maximal (locally) 4-symmetric bundle over  $M$  in the twistor bundle  $\Sigma(M)$  depends on the "size" of the isotropy subgroup  $\text{Is}_{p_0}(M)$  and on  $J_0 \in \Sigma(T_{p_0}M)$ . In other words, if we want a fiber  $S_0 \subset \Sigma(T_{p_0}M)$  of maximal dimension, we must find  $J_0 \in \Sigma(T_{p_0}M) \cap \text{Aut}(T_{p_0}M) \supset \Sigma(T_{p_0}M) \cap \text{Is}_{p_0}(M)$ <sup>12</sup> such that  $T_{J_0}\underline{S}_0 = \underline{\mathfrak{g}}_2(J_0) := \{A \in \mathfrak{Is}_{p_0}(M) \mid AJ_0 + J_0A = 0\}$  is of maximal dimension, or equivalently such that  $\underline{\mathfrak{g}}_0(J_0) = \{A \in \mathfrak{Is}_{p_0}(M) \mid AJ_0 - J_0A = 0\}$  is of minimal dimension.

**Remark 19** It is possible that there exists different non isomorphic locally 4-symmetric bundles over  $M$  (see section 5.3). And it is also possible that there does not exist locally 4-symmetric bundle over  $M$ . For example:  $M = S^1 \times S^3$ , then  $\text{Is}(M) = SO(2) \times SO(4)$  and  $\text{Is}_{p_0}(M) = SO(3)$ , and there does not exist  $J_0 \in \Sigma(\mathbb{R}^4)$  such that  $J_0SO(3)J_0^{-1} = SO(3)$ .

Moreover we have the following obvious theorem (see also [16]):

<sup>12</sup> $\text{Aut}(T_{p_0}M)$  (=  $\text{Aut}(\mathfrak{m})$  under the usual identification) is the group of automorphism of  $T_{p_0}M$  which stabilizes the metric  $g_{p_0}$  and the curvature  $R_{p_0}$ .

**Theorem 16** Let  $(\mathfrak{g}, \sigma)$  be an orthogonal symmetric Lie algebra. Then set  $\mathfrak{g}^* = \mathfrak{h} \oplus \mathfrak{im}$  and  $\sigma^* = \text{Id}_{\mathfrak{h}} \oplus \text{Id}_{\mathfrak{im}}$ . Then  $(\mathfrak{g}^*, \sigma^*)$  is an orthogonal symmetric Lie algebra. If  $(\mathfrak{g}, \sigma)$  is of the compact type then  $(\mathfrak{g}^*, \sigma^*)$  is of the non compact type and conversely. Now, for  $\tau_{\mathfrak{m}} \in \text{End}(\mathfrak{m})$ , set  $\tau_{\mathfrak{m}}^*: \mathfrak{im} \mapsto i\tau_{\mathfrak{m}}(v)$ . Then

$$\tau_{\mathfrak{m}} \in \text{Aut}(\mathfrak{m}) \iff \tau_{\mathfrak{m}}^* \in \text{Aut}(\mathfrak{im})$$

and  $\tau_{\mathfrak{m}} \in \Sigma(\mathfrak{m})$  if and only if  $\tau_{\mathfrak{m}}^* \in \Sigma(\mathfrak{im})$ . In this case  $(\tau_{\mathfrak{m}} \in \text{Aut}(\mathfrak{m}) \cap \Sigma(M))$  let  $\tau$  (resp.  $\tau^*$ ) be the automorphism of  $\mathfrak{g}$  (resp.  $\mathfrak{g}^*$ ) defined by  $\tau_{\mathfrak{m}}$  (resp.  $\tau_{\mathfrak{m}}^*$ ) and denoting by  $A^{\mathbb{C}} \in \text{End}(V^{\mathbb{C}})$  the extension to  $V^{\mathbb{C}}$  of  $A \in \text{End}(V)$  ( $V$  real vector space) then we have

$$\tau^{\mathbb{C}} = \tau^{*\mathbb{C}} \quad \text{i.e.} \quad \tau^* = \tau|_{\mathfrak{g}^*}^{\mathbb{C}}$$

**Theorem 17** Let  $M$  be an irreducible symmetric spaces of type II (compact type) or type IV (non compact type) then there does not exist any (non trivial) locally 4-symmetric bundle over  $M$ . Equivalently  $\text{Aut}(M) \cap \Sigma(M) = \emptyset$ , in other words, there does not exist any automorphism  $\tau$  of  $\mathfrak{Is}(M)$  such that  $\tau^2 = T_1\sigma$ .

**Proof.** By duality, it is enough to prove the assertion for the compact type. In this case let  $\tilde{M}$  be the universal covering of  $M$ , we have  $\tilde{M} = H \times H/\Delta(H)$  and  $\tilde{\sigma}: (a, b) \in G \times G \mapsto (b, a)$ . Then an automorphism  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  must send  $\mathfrak{g}_1 = \mathfrak{h} \oplus \{0\}$  either on  $\mathfrak{g}_1$  or on  $\mathfrak{g}_2 = \{0\} \oplus \mathfrak{h}$  and idem for  $\mathfrak{g}_2$ , and thus for any automorphism we have  $\tau^2(\mathfrak{g}_i) = \mathfrak{g}_i$  and hence we can not have  $\tau^2 = T_1\sigma$ . This completes the proof.  $\blacksquare$

## 4.2 The Euclidean case

**Theorem 18** Let  $M = \mathbb{R}^{2n}$  with its canonical inner product. Then  $\text{Is}(M) = O(2n) \times \mathbb{R}^{2n}$  the group of affine isometries in  $\mathbb{R}^{2n}$ . Then for any  $p_0 \in \mathbb{R}^{2n}$ , we have  $\text{Is}_{p_0}(M) = \{(F, \text{Id} - F)p_0, F \in O(2n)\} \simeq O(2n)$ . In particular for  $p_0 = 0$ ,  $\text{Is}_{p_0}(M) = O(2n)$ . We have  $\forall p_0 \in \mathbb{R}^{2n}$ ,  $\text{Is}(M) = \text{Is}_{p_0}(M) \times \mathbb{R}^{2n}$ . Further  $M = G/H$  is a symmetric realisation with  $G$  acting effectively if and only if  $G = H \times \mathbb{R}^{2n}$  with  $H \subset \text{Is}_{p_0}(\mathbb{R}^{2n})$  for some  $p_0 \in \mathbb{R}^{2n}$ . Then we  $G = H_0 \times \mathbb{R}^{2n}$  with  $H_0 = \text{pr}_{O(2n)}(H) \subset O(2n)$ . The cartan involution for this realisation is

$$\sigma = \text{Int}(-\text{Id}, 2p_0): (h, x) \in G \mapsto (h, 2(\text{Id} - h)p_0 - x)$$

giving rise to the symmetry around  $p_0$ :  $\sigma_0: x \in \mathbb{R}^{2n} \mapsto -(x - p_0) + p_0 \in \mathbb{R}^{2n}$ . Let us fix  $p_0 = 0$ , so that for any symmetric realisation  $H \subset \text{Is}_{p_0}(M) = O(2n)$  and  $\sigma = \text{Int}(-\text{Id}, 0)$ .

All (connected) locally 4-symmetric bundles over  $M$  are globally 4-symmetric bundle over  $M$ . The twistor bundle,  $\Sigma(\mathbb{R}^{2n}) \times \mathbb{R}^{2n}$ , is a globally 4-symmetric bundle over  $M$ . All the (connected) 4-symmetric bundles over  $\mathbb{R}^{2n}$  are in the form:  $S_0 \times \mathbb{R}^{2n}$  where  $S_0$  is a compact Riemannian symmetric space embedded<sup>13</sup>, in  $\Sigma^{\varepsilon}(\mathbb{R}^{2n})$ . Besides  $\text{Aut}(T_{p_0}M) = \text{Is}_{p_0}(M) = O(2n)$  so that any

<sup>13</sup>only immersed if  $H$  is not closed in  $O(2n)$

$J_0 \in \Sigma(\mathbb{R}^{2n})$  define the maximal 4-symmetric bundle  $\Sigma(\mathbb{R}^{2n}) \times \mathbb{R}^{2n} = O(2n) \times \mathbb{R}^{2n}/U(\mathbb{R}^{2n}, J_0)$ .

**Theorem 19** *Let  $M$  be an Euclidean Riemannian symmetric space (i.e. its universal covering is an Euclidean space  $\mathbb{R}^{2n}$ ). Then  $M = \mathbb{R}^{2n} \oplus \mathbb{T}^{2q}$ ,  $Is(M) = O(2p) \times (\mathfrak{S}_{2q} \otimes \{\pm 1\}) \times M$  ( $\mathfrak{S}_{2q}$  is the group of permutation) and denoting by  $\pi: \mathbb{R}^{2n} \rightarrow M$  the universal covering, and  $p_0 = \pi(0)$ , then  $Is_{p_0}(M) = O(2p) \times (\mathfrak{S}_{2q} \otimes \{\pm 1\})$ . Moreover  $Aut(\mathbb{R}^{2n}) = O(2n)$ , and  $J_0 \in \Sigma(\mathbb{R}^{2n})$  define the (connected) maximal 4-symmetric bundle over  $M: (\Sigma(E^{2l}) \times \{J_0|_{E^{2l+1}}\}) \times M$ , where  $E^{2l}$  is the (unique) maximal subspace in  $\mathbb{R}^{2p}$  stable by  $J_0$ . In particular,  $Aut^*(M) \cap \Sigma(M) = \Sigma(\mathbb{R}^{2p}) \times \Sigma(\mathbb{R}^{2q}) \times M$ .*

**Proof.** Let  $\tilde{\pi}: \tilde{G} \rightarrow G$  be a covering of  $G = Is(M)^0$  acting symmetrically and effectively on  $\tilde{M} = \mathbb{R}^{2n}$  and  $\tilde{\sigma}: \tilde{G} \rightarrow \tilde{G}$  the corresponding involution. Then setting  $\tilde{H} = (\tilde{G}^{\tilde{\sigma}})^0$ , we have according to the previous theorem  $\tilde{G} = \tilde{H} \ltimes \mathbb{R}^{2n}$  and  $\tilde{H} \subset SO(2n)$ . Then setting  $D = \ker \pi$ ,  $D$  is a discrete central subgroup of  $\tilde{G}$ . Besides it is easy to see that  $\text{Cent}(\tilde{G}) = \text{Cent}(\tilde{H} \times \mathbb{R}^{2n}) = \mathbb{R}^{2p}$  where  $\mathbb{R}^{2p}$  is the maximal subspace of  $\mathbb{R}^{2n}$  fixed by  $\tilde{H}$ , i.e.  $\tilde{H} \subset SO(2p) \times \{\text{Id}_{2q}\}$  ( $2p + 2q = 2n$ ). Hence  $D = \bigoplus_{i=1}^r \mathbb{Z}e_i$  with  $(e_i)_{1 \leq i \leq r}$   $\mathbb{R}$ -free so that  $G = Is(M)^0 = \tilde{G}/D = \tilde{H} \ltimes M'$  with  $M' = \mathbb{R}^{2p} \oplus \mathbb{R}^{2q-r} \oplus \mathbb{R}^r/\mathbb{Z}^r$ . Moreover we have  $\sigma: (h, x) \text{ in } \tilde{H} \times M' \rightarrow (h, -x)$  because  $\tilde{\sigma} = \text{Int}(-\text{Id}, 0)$  (see the previous theorem) and thus  $G^\sigma = \tilde{H}$  but the isotropy subgroup of  $G$  at  $p_0$  satisfies  $H \supset \tilde{\pi}(\tilde{H})$  (because  $\tilde{H}$  is connected), but  $\tilde{\pi}(\tilde{H}) = \tilde{H}$  ( $D \cap \tilde{H} = \{1\}$ ) and thus  $H = \tilde{H}$ . Thus  $M = G/H = M'$ . Now, we have to compute  $Is(M)$ , we know that  $Is(M)^0 = H \times M \subset SO(2p) \times M'$ . In the other hand, we know that  $Is(M) \subset O(2n)$  (an isometry is determined by its tangent map at  $p_0$ ) and thus  $g \in O(2n)$  is in  $Is(M)$  if and only if  $g(D) = D$  which is equivalent to  $g \in [O(2p + 2q - r) \times (GL_r(\mathbb{Z}) \cap O(\mathbb{R}^r))] \times M = [O(2p + 2q - r) \times (\mathfrak{S}_r \times \{\pm \text{Id}\})] \times M$ . Hence  $Is(M)^0 = SO(2p + 2q - r)$  and thus  $r = 2q$ . Finally  $M = \mathbb{R}^{2p} \oplus \mathbb{T}^{2q}$ ,  $Is(M) = O(2p) \times (\mathfrak{S}_r \times \{\pm \text{Id}\}) \times M$ ,  $Is_{p_0}(M) = O(2p) \times (\mathfrak{S}_r \times \{\pm \text{Id}\})$ , and  $Is_{p_0}(M)^0 = H = O(2p)$ . We conclude by remarking that  $J_0 \in \Sigma(\mathbb{R}^{2n})$  satisfies  $J_0 H J_0^{-1} = H$  for  $H \subset SO(2p)$  connected and maximal if and only if  $H = SO(E^{2l})$  and  $J_0 \in \Sigma(E^{2l}) \times \Sigma(E^{2l+1})$ . This completes the proof.  $\blacksquare$

**Remark 20** We can use the second elliptic integrable system in the Euclidean case to "modelize" this system in the general case. Indeed, let us consider  $M$  a Riemannian symmetric space of the semisimple type (then its isotropy subgroup  $H = Is_{p_0}(M)$  is essentially its holonomy group, i.e. they have the same identity component) with  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  an order four automorphism such that  $\tau^2 = T_1\sigma$ . Then we can associate to the corresponding locally 4-symmetric bundle  $N$  over  $M$ , the 4-symmetric bundle over  $M_0 = \mathfrak{m} = H \times \mathfrak{m}/H: N_0 = H \times \mathfrak{m}/G_0 = S_0 \times \mathfrak{m} \subset \Sigma(\mathfrak{m}) \times \mathfrak{m}$ , and to the second elliptic integrable system in  $N$ , its "linearized" in  $N_0$ . We conjecture that the "concrete" geometrical interpretation (i.e. in terms of the second fundamental form of the surface  $X$  etc...) is the same for the linearized and the initial system. This is what happens in dimension 4.



**Remark 21** The second elliptic integrable system can be viewed as "a coupling" between the harmonic map equation in  $S_0 = H/G_0$  and a kind of Dirac equation in  $\mathfrak{g}_{-1}$ :  $\partial_{\bar{z}}u_1 + [\bar{u}_0, u_1] + [\bar{u}_1, u_2] = 0$ . In the Euclidean case, the projection on the "group part",  $\mathfrak{g} = \mathfrak{h} \ltimes \mathfrak{m} \rightarrow \mathfrak{h}$ , of the second elliptic system is only the harmonic map equation in  $H/G_0$ . In other words, the second elliptic integrable system is only the harmonic map equation in  $H/G_0$  and a kind of Dirac equation in  $\mathbb{C}^n$  ( $\cong (\mathfrak{g}_{-1}, J_0)$ ). In particular, if we apply any method of integrable systems theory using loop groups (DPW, Dressing action etc..) or something else (spectral curves) to the second elliptic system in  $G/G_0$  and then project in the group part (pr:  $H \ltimes \mathfrak{m} \rightarrow H$ ), we obtain the same method applied to the first elliptic integrable system in  $H/G_0$  i.e. the harmonic map equation in  $H/G_0$ . For example, if we apply the DPW method: given  $\mu = (\mu_{\mathfrak{h}}, \mu_{\mathfrak{m}})$  a homomorphic potential, we have  $\text{pr}(\mathcal{W}_{G/G_0}(\mu)) = \mathcal{W}_{H/G_0}(\mu_{\mathfrak{h}})$  where  $\mathcal{W}_{G/G_0}, \mathcal{W}_{H/G_0}$  are the weierstrass representations for each elliptic system. So to solve the second elliptic system, we can first solve the harmonic map equation in  $H/G_0$ , by using any method of integrable systems theory which gives us a lift  $h$  in  $H$  of a harmonic map in  $H/G_0$ , and then we have to solve the Dirac equation with parameters  $u_0, u_2$  given by the lift :  $h^{-1}\partial_{\bar{z}}h = u_0 + u_2$  following  $\mathfrak{h} = \mathfrak{g}_0 \oplus \mathfrak{g}_2$  (see [17]). However, the Dirac equation is not intrinsic since it depends on the lift  $h$  of the harmonic map (see [17]).

In the particular case where  $S_0$  is a group and  $H = G_0 \times S_0$ , (for example  $S_0 = G_0 \times G_0/G_0$ ), then we have a canonical lift and then the Dirac equation become intrinsic (see [17]). It is in particular what happens for Hamiltonian stationary Lagrangian surfaces : in  $\mathbb{C}^2$  we have a intrinsic Dirac equation whereas in the others Hermitian symmetric spaces this equation does not exist (see [12]-[14]). It is also what happens in [17] when we take for  $S_0$  the subsphere  $S^3 \subset S^6$  ( $S^6$  embeds in  $\Sigma^+(\mathbb{R}^8)$  by the left multiplication in  $\mathbb{O}$ ).

## 5 Example of 4-symmetric bundles

We use the notations of section 3.1.

### 5.1 The sphere

Let us consider  $M = S^{2n} = SO(2n+1)/SO(2n)$  with  $G = SO(2n+1)$ ,  $H = SO(2n)$  and the involution  $\sigma = \text{Intdiag}(\text{Id}_{2n}, -1)$ . Then  $G^\sigma = SO(2n) \sqcup O^-(2n) \times \{-1\}$ . Hence  $H = (H^\sigma)^0$ ,  $M_{min} = \mathbb{R}\mathbb{P}^{2n}$  and  $M_{max} = S^{2n}$ .<sup>14</sup> We have also

$$\mathfrak{h} = \mathfrak{so}(2n), \quad \mathfrak{m} = \left\{ \begin{pmatrix} 0 & v \\ -v^t & 0 \end{pmatrix}, v \in \mathbb{R}^{2n} \right\} = \{i_{\mathfrak{m}}(v), v \in \mathbb{R}^{2n}\}$$

where  $i_{\mathfrak{m}}: \mathbb{R}^{2n} \rightarrow \mathfrak{m}$  is defined in an obvious way. Now, let us consider the action of  $H$  on  $\mathfrak{m}$ : for  $h \in SO(2n)$ ,  $\xi = i_{\mathfrak{m}}(v) \in \mathfrak{m}$ , we have

$$\text{Ad}_{\mathfrak{m}}h(\xi) = i_{\mathfrak{m}}(h.v)$$

---

<sup>14</sup> $M_{max}$  is simply connected and  $M_{min}$  is the adjoint space.

hence  $K = \ker \text{Ad}_{\mathfrak{m}} = \{\text{Id}\}$  and the action of  $G$  is effective (in fact  $SO(2n+1)$  is simple because  $2n+1$  is odd). Identifying  $\mathfrak{m}$  with  $\mathbb{R}^{2n}$  via  $i_{\mathfrak{m}}$  we have:  $\forall h \in SO(2n)$ ,  $\text{Ad}_{\mathfrak{m}} h = h$  i.e.  $\text{Ad}_{\mathfrak{m}} = \text{Id}$ . Moreover  $SO(2n+1)$  is the connected isometry group of  $S^{2n}$ . Now, according to theorem 7, define a locally 4-symmetric bundle over  $M = S^{2n}$  is equivalent to give oneself  $\tau_{\mathfrak{m}} \in \Sigma(\mathfrak{m}) \cap \text{Aut}(\mathfrak{m}) = \Sigma(\mathfrak{m})$ . Further, given  $J_0 \in \Sigma^\varepsilon(\mathbb{R}^{2n})$ , let us define the order four automorphism of  $G$ :  $\tau = \text{Int}(\text{diag}(-J_0, 1))$ . Then  $\tau^2 = \sigma$  and since  $\tau_H = \text{Int} J_0$  and  $\tau|_{\mathfrak{m}} = J_0$ , we obtain all the locally 4-symmetric bundle over  $M$  which are all globally 4-symmetric bundle over  $M$ .

Moreover, we have  $G^\tau = \text{com}(J_0) \cap SO(2n) = U(\mathbb{R}^{2n}, J_0)$ . Hence  $G^\tau = (G^\tau)^0 = G_0$  thus  $S_0 = H/G_0 = \text{Int}(SO(2n))(J_0) = \Sigma^\varepsilon(\mathbb{R}^{2n})$  and thus  $N = G/G_0 = \Sigma^\varepsilon(S^{2n})$ .

## 5.2 Real grassmannian

More generally, let  $p, q \in \mathbb{N}^*$  such that  $pq$  is even and let us consider  $M = SO(p+q)/SO(p) \times SO(q) = Gr_p(\mathbb{R}^{p+q})$  (oriented  $p$ -plans in  $\mathbb{R}^{p+q}$ ). Since  $p$  and  $q$  play symmetric roles, we will suppose that  $p$  is even and that it has the biggest divisor in the form  $2^r$ . We have  $\dim M = pq$  and the following setting

$$\begin{aligned} G &= SO(p+q), \quad H = SO(p) \times SO(q); \quad \sigma = \text{Int}(\text{diag}(\text{Id}_p, -\text{Id}_q)) \text{ and} \\ G^\sigma &= SO(p) \times SO(q) \sqcup O^-(p) \times O^-(q). \end{aligned}$$

Then  $H = (G^\sigma)^0$  so that  $M_{\min} = Gr_p^*(\mathbb{R}^{p+q})$  (non oriented  $p$ -plan in  $\mathbb{R}^{p+q}$ ) and

$$M_{\max} = Gr_p(\mathbb{R}^{p+q}) = M. \text{ Besides } \mathfrak{h} = \mathfrak{so}(p) \oplus \mathfrak{so}(q), \text{ and } \mathfrak{m} = \left\{ \begin{pmatrix} 0 & B \\ -B^t & 0 \end{pmatrix}, B \in \mathfrak{gl}_{p,q}(\mathbb{R}) \right\} =$$

$i_{\mathfrak{m}}(\mathfrak{gl}_{p,q}(\mathbb{R}))$  ( $i_{\mathfrak{m}}$  defined in an obvious way).

Now let us compute  $\text{Ad}_{\mathfrak{m}}$ . For  $h = \text{diag}(A, C)$  and  $\xi = i_{\mathfrak{m}}(B)$ , we have:

$$\text{Ad}_{\mathfrak{m}} h(\xi) = i_{\mathfrak{m}}(ABC^{-1}).$$

Under the identification  $i_{\mathfrak{m}}$  we have  $\text{Ad}_{\mathfrak{m}}(A, C) = L_A R_{C^{-1}} = \chi(A, C)$ , by introducing the morphism  $\chi: (A, C) \in GL_p(\mathbb{R}) \times GL_q(\mathbb{R}) \mapsto L(A)R(C^{-1}) \in GL(\mathfrak{gl}_{p,q}(\mathbb{R}))$ . Hence  $K = \ker \text{Ad}_{\mathfrak{m}} = \{\pm \text{Id}\}$  if  $q$  is even and  $K = \{\text{Id}\}$  if not. Thus the connected isometry group of  $M$ ,  $\text{Is}(M)^0$ , is  $G' = G/K = PSO(p+q)$  if  $q$  is even and  $G' = G = SO(p+q)$  if not. Let us compute  $\text{Aut}(\mathfrak{m})$ : we already know that  $\text{Aut}(\mathfrak{m}) \subset H \subset \text{Aut}(\mathfrak{m})^0$ . But, it is known that the automorphisms of  $\mathfrak{so}(n+1)$  are all inner automorphisms by  $O(n+1)$  so we have  $\text{Aut}(\mathfrak{m}) = \{L_A R_{C^{-1}}, (A, C) \in O(p) \times O(q)\}$ . Thus  $J_0 = L(J_1)R(J_2^{-1}) \in \text{Aut}(\mathfrak{m})$  is in  $\Sigma(\mathfrak{m})$  if and only if :

$$\begin{cases} (J_1^2, J_2^2) &= \pm(-\text{Id}_p, \text{Id}_q) & \text{if } q \text{ is even,} \\ (J_1^2, J_2^2) &= (-\text{Id}_p, \text{Id}_q) & \text{if } q \text{ is odd.} \end{cases}$$

Then the associated order four automorphism is  $\tau = \text{Int}(\text{diag}(J_1, J_2))$ . In particular,  $\tau(H) = H$  and  $\tau_H = \text{Int} J_1 \times \text{Int} J_2$ . Besides,  $\text{Aut}(\mathfrak{m}) \cap \Sigma(\mathfrak{m})$  has respectively  $2(p+q+2)$  or  $2(q+1)$  connected components if  $q$  is even or  $q$  is odd respectively.

Each connected component is an  $\text{Ad}_m H$ -orbit and corresponds to the fiber of a different maximal 4-symmetric bundle over  $M$ .

Moreover to fix idea let us suppose that we have  $J_1 \in \Sigma(\mathbb{R}^p), J_2 \in OS(\mathbb{R}^q)$ , the set of orthogonal symmetry in  $\mathbb{R}^q$ , then  $G^\tau = U(\mathbb{R}^p, J_1) \times S(O(E_1) \times O(E_2))$  with  $E_1 = \ker(J_2 - \text{Id}), E_2 = \ker(J_2 + \text{Id})$ . We have  $G^\tau \subset H$ . Let  $OS_r(\mathbb{R}^q) = \text{Int}(SO(q))(\text{Id}_r, -\text{Id}_{q-r})$  be the set of orthogonal symmetry in  $\mathbb{R}^q$  with  $\dim E_1 = r$ . Then  $H/G^\tau = \text{Int}(H)(J_1, J_2) = \Sigma^\varepsilon(\mathbb{R}^p) \times OS_r(\mathbb{R}^q)$  ( $\varepsilon$  being determined by  $J_1$ ) and

$$G/G^\tau = \{(x, J), x \in M, J \in \Sigma^\varepsilon(x) \times OS_r(x^\perp)\}. \quad (21)$$

Now let us compute  $G_0$  according to (11):  $h = (A, C) \in H$  si in  $G_0$  if and only if  $\text{Ad}_m \tau(h) = \text{Ad}_m h$  i.e.: if  $q$  is odd,  $\tau(h) = h$ , and  $G_0 = G^\tau \cap H = G^\tau$ ; if  $q$  is even,  $\tau(h) = \pm h$  (and  $G_0 = \pi_K^{-1}(G'_0)$  with  $G'_0 = G'^{\tau'} \cap H'$ ), i.e.  $h \in G^\tau$  or  $\tau(h) = -h$ . The existence of solutions of this last equation depends on  $p, q$  and  $r$  (we remark that if  $h_1$  is a solution then the set of solution is  $h_1 G^\tau$ ). One find that the equation  $\tau(h) = -h$  ( $q$  is even) has a solution in  $G^\sigma$  if and only if  $\dim E_1 = \dim E_2 = q/2$  and that this solution is in  $H$  if  $p/2$  is even and in  $O^-(p) \times O^-(q)$  (the other component of  $G^\sigma$ ) if  $p/2$  is odd. Hence, if  $p$  is divisible by 4,  $q$  is even and  $r = q/2$  (i.e.  $J_0 \in \chi(\Sigma(\mathbb{R}^p) \times OS_{q/2}(\mathbb{R}^q))$ ), we have  $G_0 = G^\tau \sqcup h_1 G^\tau$ . In all the other cases we have  $G_0 = G^\tau$ .

In conclusion, let us denote by  $N^L(r, \varepsilon) := N(J_0)$  (resp.  $N^R(r, \varepsilon)$ ) the maximal 4-symmetric bundle over  $M$  corresponding to  $J_0 \in \chi(\Sigma^\varepsilon(\mathbb{R}^p) \times OS_r(\mathbb{R}^q))$  (resp.  $\text{chi}(OS_r(\mathbb{R}^p) \times \Sigma^\varepsilon(\mathbb{R}^q))$ ). Then:

- if  $p$  is not divisible by 4 or  $q$  is odd,  $N^\alpha(r, \varepsilon)$  is given by (21), for all  $(\alpha, r, \varepsilon)$ ,
- if  $p$  is divisible by 4,  $q$  not divisible by 4 then for  $(\alpha, r) \neq (L, q/2)$ ,  $N^\alpha(r, \varepsilon)$  is given by (21) and for  $(\alpha, r) = (L, q/2)$  it is given by (22), below,
- if  $p$  and  $q$  are divisible by 4, then for  $(\alpha, r) \in \{(L, q/2), (R, p/2)\}$ ,  $N^\alpha(r, \varepsilon)$  is given by (22), and for the other choices it is given by (21),

$$N^\alpha(r, \varepsilon) = \{(x, J), x \in M, J \in P(\Sigma^\varepsilon(x) \times OS_r(x^\perp))\} \quad (22)$$

where  $P(\Sigma^\varepsilon(x) \times OS_r(x^\perp)) = \Sigma^\varepsilon(x) \times OS_r(x^\perp) / \{\pm \text{Id}\}$ . In the cases described by (22),  $G/G^\tau$  is not a submanifold of  $\Sigma(M)$ .

### 5.3 Complex Grassmannian

Let us consider  $M = SU(p+q)/S(U(p) \times U(q)) = Gr_{p,\mathbb{C}}(\mathbb{C}^{p+q})$ . We have  $\dim M = 2pq$  and the following setting

$$G = SU(p+q), H = S(U(p) \times U(q)); \sigma = \text{Int}(\text{diag}(\text{Id}_p, -\text{Id}_q)) \text{ and } G^\sigma = H = (G^\sigma)^0.$$

Besides  $\mathfrak{h} = \mathfrak{s}(\mathfrak{u}(p) \oplus \mathfrak{u}(q))$  and  $\mathfrak{m} = \left\{ \begin{pmatrix} 0 & B \\ -B^* & 0 \end{pmatrix}, B \in \mathfrak{gl}_{p,q}(\mathbb{C}) \right\} = i_{\mathfrak{m}}(\mathfrak{gl}_{p,q}(\mathbb{C}))$ .

Let us compute  $\text{Ad}_m$ . For  $h = \text{diag}(A, C)$  and  $\xi = i_{\mathfrak{m}}(B)$ , we have:

$$\text{Ad}_m h(\xi) = i_{\mathfrak{m}}(ABC^{-1}).$$

Under the identification  $i_{\mathfrak{m}}$  we have  $\text{Ad}_{\mathfrak{m}}(A, C) = L_A R_{C^{-1}} = \chi(A, C)$ , by introducing the morphism  $\chi: (A, C) \in GL_p(\mathbb{C}) \times GL_q(\mathbb{C}) \mapsto L(A)R(C^{-1}) \in GL(\mathfrak{gl}_{p,q}(\mathbb{C}))$ <sup>15</sup>. Hence  $K = \ker \text{Ad}_{\mathfrak{m}} = \{(\lambda \text{Id}_p, \lambda \text{Id}_q), \lambda \in \mathbb{C}, \lambda^{p+q} = 1\} = \hat{U}_{p+q} \text{Id} \simeq \mathbb{Z}_{p+q}$  (with  $\hat{U}_{p+q} = \exp(\frac{2i\pi}{p+q}\mathbb{Z})$ ). Thus  $G' = G/K = PSU(p+q)$  and  $H' = S(U(p) \times U(q))/\hat{U}_{p+q} \simeq S(U(p) \times U(q))$ . The connected isometry group is the unitary group of  $M$ :  $\text{Is}(M)^0 = U(M) = G' = PSU(p+q)$ .

It is well known that the group of automorphism of  $SU(p+q)$  have two components (the  $\mathbb{C}$ -linear one and the anti  $\mathbb{C}$ -linear one) and is generated by the inner automorphisms and the complex conjugaison:  $g \in SU(p+q) \mapsto \bar{g} \in SU(p+q)$ . In particular,  $\text{Aut}(\mathfrak{m}) = \text{Ad}_{\mathfrak{m}} H \rtimes \{\text{Id}, c\} = \chi(S(U(p) \times U(q)) \cdot \{(\text{Id}, \text{Id}), (b_p, b_q)\})$  with  $c = L(b_p)R(b_q^{-1}) : B \in \mathfrak{gl}_{p,q}(\mathbb{C}) \mapsto \bar{B} \in \mathfrak{gl}_{p,q}(\mathbb{C})$ ,  $b_n : v \in \mathbb{C}^n \mapsto \bar{v} \in \mathbb{C}^n$ .

The complex structure in  $\mathfrak{m} = \mathfrak{gl}_{p,q}(\mathbb{C})$  is defined by  $L(I_p) = R(I_q)$  where  $I_n = i\text{Id}_n$  the canonical complex structure in  $\mathbb{C}^n$ , and the two connected components of  $\text{Aut}(\mathfrak{m})$  are respectively the elements in  $\text{Aut}(\mathfrak{m})$  which commute and those which anticommute with this complex structure.

Moreover,  $J_0 = L(J_1)R(J_2^{-1}) \in \text{Aut}(\mathfrak{m})^0 = \text{Ad}_{\mathfrak{m}} H$  is in  $\Sigma(\mathfrak{m})$  if and only if  $(J_1^2, J_2^2) \in (-\text{Id}_p, \text{Id}_q)U(1)$ . Then let us set  $\Sigma_\lambda = \{(J_1, J_2) \in U(p) \times U(q) \mid (J_1^2, J_2^2) = \lambda(-\text{Id}_p, \text{Id}_q)\}$ . Then we have  $\chi(\Sigma_\lambda) = \chi(\Sigma_0)$  for all  $\lambda \in U(1)$  since  $\Sigma_\lambda = \lambda^{\frac{1}{2}}\Sigma_0$  with  $\lambda^{\frac{1}{2}}$  a root of  $\lambda$ . Thus according to the following lemma,  $\text{Aut}(\mathfrak{m})^0 \cap \Sigma(\mathfrak{m})$  has  $(p+1)(q+1)$  connected components (which are  $\text{Ad}_{\mathfrak{m}} H$ -orbits and corresponds to the fibers of different maximal 4-symmetric bundles over  $M$ ).

**Lemma 1** *Let  $J \in U(n)$ , then  $J^2 = -\text{Id}$  (resp.  $J^2 = \text{Id}$ ) if and only if there exists  $h \in U(n)$  such that  $hJh^{-1} = \text{diag}(i\text{Id}_l, -i\text{Id}_{n-l})$  for some  $l \in \{0, \dots, n\}$  (resp.  $hJh^{-1} = \text{diag}(\text{Id}_r, -\text{Id}_{n-r})$  for some  $r \in \{0, \dots, n\}$ ).*

Then the order four automorphism corresponding to  $J_0$  is  $\tau = \text{Int}(\text{diag}(J_1, J_2)$ , with<sup>16</sup>  $J_1 \in \text{Ad}U(p)(i\text{I}_{l,p-l}) \cong iGr_{l,\mathbb{C}}(\mathbb{C}^p)$ ,  $J_2 \in \text{Ad}U(q)(\text{I}_{r,q-r}) \cong Gr_{r,\mathbb{C}}(\mathbb{C}^q)$ . Hence  $G^\tau = S(U(l) \times U(p-l) \times U(r) \times U(q-r))$ , the fiber of the 4-symmetric space  $G/G^\tau$  is  $H/G^\tau = Gr_{l,\mathbb{C}}(\mathbb{C}^p) \times Gr_{r,\mathbb{C}}(\mathbb{C}^q)$ , and

$$G/G^\tau = \{(x, J), x \in Gr_{p,\mathbb{C}}(\mathbb{C}^{p+q}), J \in Gr_{l,\mathbb{C}}(x) \times Gr_{r,\mathbb{C}}(x^\perp)\}. \quad (23)$$

Further,  $G_0$  is defined:  $\text{Ad}_{\mathfrak{m}}\tau(h) = h, h \in H$ , i.e.  $(J_1 A J_1^{-1}, J_2 C J_2^{-1}) = \lambda(A, C)$  for some  $\lambda \in K$ . But it is easy to see that we must have  $\lambda = 1$  and thus  $G_0 = G^\tau$ . Finally, in the  $\mathbb{C}$ -linear case, the maximal 4-symmetric bundle  $N = G/G_0$  is given by (23).

In the antilinear case,  $J_0 = L(J_1)R(J_2^{-1}) \in \text{Aut}(\mathfrak{m})^0.c$ , with  $(J_1, J_2) = (J'_1 b_p, J'_2 b_q)$ , is in  $\Sigma(\mathfrak{m})$  if and only if  $(J_1^2, J_2^2) = (J'_1 \bar{J}'_1, J'_2 \bar{J}'_2) \in (-\text{Id}_p, \text{Id}_q)U(1)$ . It is easy to see that we can only have

$$(J_1^2, J_2^2) = \pm(-\text{Id}_p, \text{Id}_q). \quad (24)$$

<sup>15</sup>For the following it useful to keep in mind that we have  $\text{Ad}_{\mathfrak{m}} H = \chi(S(U(p) \times U(q))) = \chi(U(p) \times U(q))$  and  $\ker \chi = \mathbb{C}^* \text{Id}$ .

<sup>16</sup> $\text{I}_{l,p-l} = \text{diag}(\text{Id}_l, -\text{Id}_{p-l})$

Hence according to the following lemma:

- if  $p, q$  are odd then  $\Sigma(\mathfrak{m}) \cap \text{Aut}(\mathfrak{m})^0.c = \emptyset$ ,
- if  $p, q$  are even then the two signs  $\pm$  are realized in (24) and thus  $\Sigma(\mathfrak{m}) \cap \text{Aut}(\mathfrak{m})^0.c$  has 2 connected components,
- if  $p, q$  have opposite parities, then only one sign is realized in (24) and  $\Sigma(\mathfrak{m}) \cap \text{Aut}(\mathfrak{m})^0.c$  has one component.

**Lemma 2** *Let  $E \subset \mathbb{C}^n$  be a Lagrangian  $n$ -plan, i.e.  $E \oplus iE = \mathbb{C}^n$  and let  $b_E$  be the associated conjugaison:  $v + iw \mapsto v - iw$  for  $v, w \in E$ . then  $U(N).b_E = b_E.U(n)$  does not depend on  $E$  and is the of anti  $\mathbb{C}$ -linear isometries in  $\mathbb{C}^n$  (the elements in  $O(\mathbb{R}^{2n})$  which anticommute with the complex structure  $I = i\text{Id}$ ). Moreover for any  $J$  in this set there exists a lagrangian  $n$ -plan  $E$  such that  $J = J_E.b_E = b_E.J_E$  with  $J_E \in O(E)$ . Besides  $J \in \Sigma(\mathbb{R}^{2n}$  (resp:  $OS(\mathbb{R}^{2n})$ ) if and only if  $J_E \in \Sigma(E)$  (resp.  $OS(E)$ ) and in particular  $\Sigma(\mathbb{R}^{2n}) \cap U(n).b_E \neq \emptyset$  only if  $n$  is even, and  $\Sigma(\mathbb{R}^{2n}) \cap U(n).b_E \subset \Sigma^+(\mathbb{R}^{2n})$ . Further given any  $J_1 \in \Sigma(\mathbb{R}^n)$  (resp.  $OS(\mathbb{R}^n)$ ) there exists  $h \in U(n)$  such that  $h.E = \mathbb{R}^n$ ,  $hJ_1h^{-1} = J_1$  and thus  $hJ_1h^{-1} = J_1.b_{\mathbb{R}^n}$ .*

Then the order four automorphism corresponding to  $J_0$  is  $\tau = \text{Int}(\text{diag}(J_1, J_2))$  with  $J_1 \in \text{Ad}U(p)(J_{\frac{p}{2}}.b_p)$ ,  $J_2 \in \text{Ad}U(q)(b_q)$  and  $J_{\frac{p}{2}} = \begin{pmatrix} 0 & \text{Id}_{\frac{p}{2}} \\ -\text{Id}_{\frac{p}{2}} & 0 \end{pmatrix}$ . In other words  $J_1$  is any complex structure in  $\mathbb{R}^{2p}$  anticommuting with  $I_p$  and  $J_2$  is any orthogonal conjugaison in  $\mathbb{C}^q$ . Hence, we have  $G^\tau = Sp(p/2) \times SO(q)$ . Hence  $U(p) \times U(q)/G^\tau = \Sigma^+(\mathbb{C}^p)_- \times \text{Lag}(\mathbb{C}^q)$  where  $\Sigma^+(\mathbb{C}^p)_- = \Sigma(\mathbb{R}^{2p}) \cap \text{Ant}(I_p)$  are the complex structures in  $\mathbb{R}^{2p}$  anticommuting with  $I_p$  and  $\text{Lag}(\mathbb{C}^q)$  are the oriented Lagrangian plan in  $\mathbb{C}^q$ . Thus we have:

$$H/G^\tau = S(\Sigma^+(\mathbb{C}^p)_- \times \text{Lag}(\mathbb{C}^q)) := \{(J, P) \in \Sigma^+(\mathbb{C}^p)_- \times \text{Lag}(\mathbb{C}^q) \mid \det_{\mathbb{C}}(J)\det_{\mathbb{C}}(P) = 1\}.$$

It is easy to define  $\det_{\mathbb{C}}$  on  $\text{Lag}(\mathbb{C}^q)$ , and for  $\Sigma^+(\mathbb{C}^p)_-$ , we set  $\det_{\mathbb{C}}(J) = \det_{\mathbb{C}}(E)$  for  $E$  any Lagrangian  $n$ -plan stable by  $J$  (definition independant on the choice of  $E$ ). Then

$$G/G^\tau = \{(x, J, P), x \in Gr_{p, \mathbb{C}}(\mathbb{C}^{p+q}), (J, P) \in \Sigma^+(x)_- \times \text{Lag}(x^\perp)\}.$$

Let us compute  $G/G_0$ . We have to solve for  $(A, C) \in U(p) \times U(q)$ :  $(J_{\frac{p}{2}} \bar{A} J_{\frac{p}{2}}^{-1}, \bar{C}) = \lambda(A, C)$  for  $\lambda \in U(1)$  whose the solutions are  $\pm \lambda^{\frac{1}{2}}(Sp(p/2) \times O(q))$ . Hence we have  $G'_0 = G_0/K = \chi(U(1)(Sp(p/2) \times O(q))) = \chi(Sp(p/2) \times O(q)) =$

$$\begin{cases} \chi(Sp(p/2) \times SO(q)) & \text{if } q \text{ is odd} \\ \chi(G^\tau) \sqcup h_1 \chi(G^\tau) & \text{if } q \text{ is even.} \end{cases}$$

Then  $G'/G'_0 = G/G_0 = U(p+q)/(U(1)(Sp(p/2) \times O(q))) = PSU(p+q)/P(Sp(p/2) \times O(q))$  hence  $N = G/G_0$  is equal to  $(G/G^\tau)/\mathbb{Z}_{p+q}$  if  $q$  is odd and to  $(G/G^\tau)/\mathbb{Z}_{2(p+q)}$  if  $q$  is even.

## 6 Appendix

**Theorem 20** *Let  $G$  be a connected Lie group with an involution  $\sigma$ . If  $\text{Ad}_m(G^\sigma)^0$  is compact (resp. relatively compact) then  $\text{Ad}_m H$  is compact (resp. relatively compact) for any  $H$  such that  $(G^\sigma)^0 \subset H \subset G^\sigma$ .*

**Proof.** According to [1],  $(G^\sigma)/(G^\sigma)^0$  is finite hence  $H/(G^\sigma)^0$  is finite and the theorem follows. ■

**Corollary 3** *We give ourself the same setting and notations as in remark 7. If  $\tilde{H} = (\tilde{G}^\sigma)^0$  satisfies  $\text{Ad}_m \tilde{H}$  is compact (resp. relatively compact) then for any symmetric pair  $(G, H)$ ,  $\text{Ad}_m H$  is compact (resp. relatively compact). In other words if one symmetric pair (associated to  $(\mathfrak{g}, \sigma)$ ) is Riemannian then all the others are also.*

**Proof.** Since  $\tilde{G}$  is simply connected, it is the universal covering of  $G$  and we have a covering  $\pi: \tilde{G} \rightarrow G$ . Then  $\text{Ad}_m \tilde{H} = \text{Ad}_m H^0$  (there are connected with the same Lie algebra) hence  $\text{Ad}_m H^0$  is compact and then according to the previous theorem,  $\text{Ad}_m H$  is compact. ■

**Corollary 4** *Let  $(G, H)$  be a symmetric pair with involution  $\sigma$  and  $\tau: G \rightarrow G$  an order four automorphism such that  $\tau^2 = \sigma$ . Then if  $\text{Ad}_m H$  is compact (resp. relatively compact) then the subgroup generated by  $\text{Ad}_m H$  and  $\tau|_m$ ,  $\text{Gr}(\text{Ad}_m H, \tau|_m)$  is compact (resp. relatively compact).*

**Proof.** We have  $\tau|_m(\text{Ad}_m G^\sigma)\tau|_m^{-1} = \text{Ad}_m \tau(G^\sigma) = \text{Ad}_m G^\sigma$ . Hence  $\text{Gr}(\text{Ad}_m G^\sigma, \tau|_m) = (\text{Ad}_m G^\sigma)\text{Gr}(\tau|_m)$  which is (relatively) compact because so is  $\text{Ad}_m G^\sigma$ , according to theorem 20 and then  $\text{Gr}(\text{Ad}_m G^\sigma, \tau|_m)$  is (relatively) compact because since  $\text{Gr}(\text{Ad}_m H, \tau|_m) \subset (\text{Ad}_m H)\text{Gr}(\tau|_m)$  then  $\text{Ad}_m G^\sigma/\text{Ad}_m H$  is a covering of  $\text{Gr}(\text{Ad}_m G^\sigma, \tau|_m)/\text{Gr}(\text{Ad}_m H, \tau|_m)$  which is consequently finite. ■

**Theorem 21** *Let  $(G, H)$  be a symmetric pair with involution  $\sigma$  and  $\tau: \mathfrak{g} \rightarrow \mathfrak{g}$  an order four automorphism such that  $\tau^2 = T_1\sigma$ . Then if  $\text{Ad}_m H$  is relatively compact then the subgroup generated by  $\text{Ad}_m H$  and  $\tau|_m$ ,  $\text{Gr}(\text{Ad}_m H, \tau|_m)$  is relatively compact.*

**Proof.** Let  $G' = \text{Ad}G$ , then  $C := \ker \text{Ad} = \text{center of } G$  and we can identify  $\text{Ad}$  to the covering  $\pi: G \rightarrow G/C$  and  $G'$  to  $G/C$ . The automorphism  $\sigma$  gives rise to  $\sigma': G' \rightarrow G'$  such that  $\sigma' \circ \pi = \pi \circ \sigma$ . Besides the automorphism  $\tau$  integrates in  $G'$  into  $\tau'$  defined by  $\tau' = \text{Int}\tau: \text{Ad}g \in G' \mapsto \tau \circ \text{Ad}g \circ \tau^{-1}$  and we have  $\tau' \circ \pi = \pi \circ \tau$  and  $\tau'^2 = \sigma'$ . Then according to corollary 4,  $\text{Gr}(\text{Ad}_m G'^{\sigma'}, \tau|_m)$  is relatively compact since according to corollary 3,  $\text{Ad}_m G'^{\sigma'}$  is relatively compact because  $\text{Ad}_m H$  is. Moreover we have  $G'^{\sigma'} \subset \pi(G^\sigma)$  then (since  $\text{Ad}\pi(g) = \text{Ad}g \forall g \in G$ )  $\text{Ad}_m G'^{\sigma'} \subset \text{Ad}_m G^\sigma \subset \text{Ad}_m H$  thus  $\text{Gr}(\text{Ad}_m H, \tau|_m)$  is relatively compact. ■

**Theorem 22** Let  $(\mathfrak{g}, \sigma)$  be an orthogonal symmetric Lie algebra<sup>17</sup> such that  $\mathfrak{h} = \mathfrak{g}^\sigma$  contains no ideal  $\neq 0$  in  $\mathfrak{g}$ . Then for any symmetric pair  $(G, H)$  associated with  $(\mathfrak{g}, \mathfrak{h})$ , the associated symmetric spaces  $M = G/H$  is Riemannian. Moreover let  $\tilde{G}$  be the simply connected Lie group with Lie algebra  $\mathfrak{g}$ ,  $\tilde{\sigma}$  integrating  $\sigma$ ,  $\tilde{H} = (\tilde{G}^{\tilde{\sigma}})^0$  and  $\tilde{C}$  the center of  $\tilde{G}$ . Then we have  $\tilde{H} = \tilde{G}^{\tilde{\sigma}}$ . Further, for any subgroup  $S$  of  $\tilde{C}$  put

$$H_S = \{g \in \tilde{G} \mid \tilde{\sigma}(g) \in g.S\}.$$

The symmetric spaces  $M$  associated with  $(\mathfrak{g}, \sigma)$  (i.e.  $(G, H)$  is associated with  $(\mathfrak{g}, \mathfrak{h})$ ) are exactly the spaces  $M = G/H$  with

$$G = \tilde{G}/S \quad \text{and} \quad H = H^*/S \tag{25}$$

where  $S$  varies through all  $\tilde{\sigma}$ -invariant subgroups of  $\tilde{C}$  and  $H^*$  varies through all  $\tilde{\sigma}$ -invariant subgroups of  $\tilde{G}$  such that  $\tilde{H}S \subset H^* \subset H_S$ . Hence, all the symmetric spaces  $M = G/H = \tilde{G}/H^*$  associated with  $(\mathfrak{g}, \sigma)$  cover the adjoint space of  $(\mathfrak{g}, \sigma)$ :  $M' = G'/G'^{\sigma'} = \tilde{G}/H_{\tilde{C}}^{\tilde{\sigma}}$ <sup>18</sup> and are covered by  $\tilde{M} = \tilde{G}/\tilde{H}$  (the universal covering):

$$\tilde{M} \rightarrow M \rightarrow M'. \tag{26}$$

Besides if  $\langle \cdot, \cdot \rangle$  is an  $\text{Ad}_m G'^{\sigma'}$ -invariant inner product then it is invariant by  $\text{ad}_m H = \text{Ad}_m H^*$  for any  $H$  described above, and the coverings (26) are Riemannian, when  $M, \tilde{M}, M'$  are endowed with the corresponding metrics.

**Proof.** We have only to prove  $\tilde{H} = \tilde{G}^{\tilde{\sigma}}$ , since all the rest is an adaptation of [15] (Ch. VII, thm 9.1) using what precedes. According to [1],  $\tilde{G}^{\tilde{\sigma}}$  is connected if  $\pi_1(\tilde{G})$  is finite with odd order, in particular if  $\tilde{G}$  is simply connected. This completes the proof. ■

## References

- [1] J. An, Z. Wang, *On the realization of Riemannian symmetric spaces in Lie groups II* preprint arXiv: math/0504120.
- [2] A.L. Besse, *Einstein Manifolds*, Springer-Verlag, Berlin, Heidelberg, New York, 1987.
- [3] F.E. Burstall, F. Pedit, *Harmonic maps via Adler-Kostant-Symes Theory*, Harmonic maps and integrable systems, A.P. Fordy, J.C. Wood (Eds.), Vieweg (1994), 221-272.

<sup>17</sup>i.e.  $\sigma$  is an involutive automorphism and  $\mathfrak{h} = \mathfrak{g}^\sigma$  is compactly embedded in  $\mathfrak{g}$  (see [15])

<sup>18</sup>with the notation of the proof of theorem 21. For any  $(G, H)$  symmetric pair associated with  $(\mathfrak{g}, \sigma)$ , we have  $G' = \text{Ad}G = \text{Int}(\mathfrak{g})$  the group of inner automorphism of  $\mathfrak{g}$  (see [15]) and  $\sigma$  induces an automorphism  $\sigma'$  of  $G' = \text{Int}(\mathfrak{g})$ .

- [4] F.E. Burstall and J.H. Rawnsley, *Twistor theory for Riemannian Symmetric Spaces with applications to harmonic maps of Riemann Surfaces* Lect. Notes in Math., vol. 1424, Springer, Berlin, Heidelberg, New York, 1990.
- [5] J. Dieudonné, *Éléments d'analyse, Tome 2*, Gauthiers-Villars.
- [6] J. Dorfmeister, F. Pedit and H.-Y. Wu, *Weierstrass type representation of harmonic maps into symmetric spaces*, Comm. in Analysis and Geometry, 6(4) (1998), p. 633-668.
- [7] R. Harvey, *Spinors and Calibrations*, Academic Press Inc., 1990.
- [8] R. Harvey and H. B. Lawson, *Calibrated geometries*, Acta Mathematica, 148 (1982), p. 47-157.
- [9] F. Hélein, *Applications harmoniques, lois de conservations et repères mobiles*, Diderot éditeur, Paris 1996; or *Harmonic maps, conservation laws and moving frames*, Cambridge University Press 2002.
- [10] F. Hélein, *Constant mean Curvature Surfaces, Harmonic maps and Integrable Systems*, Lecture in Mathematics, ETH Zürich, Birkhäuser 2001.
- [11] F. Hélein, *Willmore immersions and loop groups*, J. Diff. Geometry, 50(2) (1998), p.331-338.
- [12] F. Hélein and P. Romon, *Hamiltonian stationary Lagrangian surfaces in  $\mathbb{C}^2$* , Comm. in Analysis and Geometry Vol. 10, N. 1, 2002, p. 79-126.
- [13] F. Hélein and P. Romon, *Weierstrass representation of lagrangian surfaces in four dimensional spaces using spinors and quaternions*, Comment. Math. Helv., 75 (2000), p. 668-680.
- [14] F. Hélein and P. Romon, *Hamiltonian stationnary Lagrangian surfaces in Hermitian symmetric spaces*, in *Differential Geometry and Integrable Systems*, Martin Guest, Reiko Miyaoka, and Yoshihiro Ohnita, Editors-AMS, 2002.
- [15] S. Helgason, *Differential geometry, Lie group and symmetric spaces*, Academic Press, Inc., 1978.
- [16] J.A. Jimenez, *Riemannian 4-symmetric spaces*, Transactions of the American Mathematical Society, Vol.306, No.2. (Apr.,1988), pp. 715-734.
- [17] I. Khemar, *Surfaces isotropes de  $\mathbb{O}$  et systèmes intégrables.*, preprint arXiv:math.DG/0511258, to appear in Journal of Differential Geometry.
- [18] I. Khemar, *Supersymmetric Harmonic Maps into Symmetric Spaces*, Journal of Geometry and Physics 57 (2007) 1601-1630, doi:10.1016/j.geomphys.2007.01.005.



- [19] A. Pressley and G. Segal, *Loop groups*, Oxford Mathematical Monographs, Clarendon Press, Oxford, 1986.
- [20] C.L. Terng, *Geometries and Symmetries of Soliton Equations and Integrable Elliptic Equations*, preprint arXiv:math.DG/0212372.
- [21] J.A. Wolf and A. Gray, *Homogeneous spaces defined by Lie group automorphisms*. I, II, *J. Differential Geom.* **2** (1968), 77-159.

**Idrisse Khemar**  
**khemar@math.jussieu.fr**