



Uniform Eberlein spaces and the finite axiom of choice

Marianne Morillon

► To cite this version:

| Marianne Morillon. Uniform Eberlein spaces and the finite axiom of choice. 2008. hal-00268797

HAL Id: hal-00268797

<https://hal.science/hal-00268797>

Preprint submitted on 1 Apr 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

UNIFORM EBERLEIN SPACES AND THE FINITE AXIOM OF CHOICE

MARIANNE MORILLON

ABSTRACT. We work in set-theory without choice **ZF**. Given a closed subset F of $[0, 1]^I$ which is a bounded subset of $\ell^1(I)$ (*resp.* such that $F \subseteq \ell^0(I)$), we show that the countable axiom of choice for finite subsets of I , (*resp.* the countable axiom of choice **AC**_N) implies that F is compact. This enhances previous results where **AC**_N (*resp.* the axiom of Dependent Choices **DC**) was required. Moreover, if I is linearly orderable (for example $I = \mathbb{R}$), the closed unit ball of $\ell^2(I)$ is weakly compact (in **ZF**).



EQUIPE RÉUNIONNAISE DE MATHÉMATIQUES ET INFORMATIQUE
THÉORIQUE (ERMIT)

CONTENTS

1. Introduction	2
2. Some weak forms of AC	3
2.1. Restricted axioms of choice	3
2.2. Well-orderable union of finite sets	4
2.3. Dependent Choices	4
2.4. The “Tychonov” axiom	4
3. Some classes of closed subsets of $[0, 1]^I$	4
3.1. Eberlein closed subsets of $[0, 1]^I$	5
3.2. Uniform Eberlein closed subsets of $[0, 1]^I$	5
3.3. Strongly Eberlein closed subsets of $[0, 1]^I$	7
4. Compactness (in ZF)	7
4.1. Lattices and filters	7
4.2. Continuous image of a compact space	7
4.3. Sequential compactness	8
4.4. AC _N and countable products of compact spaces	8
5. One-point compactifications and related spaces	9

Date: April 1, 2008.

2000 *Mathematics Subject Classification.* Primary 03E25 ; Secondary 54B10, 54D30, 46B26.

Key words and phrases. Axiom of Choice, product topology, compactness, Eberlein spaces, uniform Eberlein spaces.

5.1.	The one-point compactification of a set	9
5.2.	Various notions of compactness for \hat{X}^α , α ordinal	9
5.3.	Spaces \hat{X}^α , α ordinal	10
5.4.	Spaces $\sigma_n(X)$, n integer ≥ 1	10
6.	$\mathbf{AC}^{fin(I)}$ and closed compactness of $B_1(I)$	12
6.1.	Dyadic representations	12
6.2.	Another equivalent of $\mathbf{AC}^{fin(I)}$	12
6.3.	Consequences	13
7.	$\mathbf{AC}_\mathbb{N}$ and Eberlein spaces	14
7.1.	Sequential compactness of I -Eberlein spaces	14
7.2.	Countable product of finitely restricted spaces	14
7.3.	Countable products of strong Eberlein spaces	14
7.4.	$\mathbf{AC}_\mathbb{N}$ and Eberlein closed subsets of $[0, 1]^I$	15
7.5.	Convex-compactness and the Hahn-Banach property	15
	References	16

1. INTRODUCTION

We work in the set-theory without the Axiom of Choice **ZF**. It is a well known theorem of Kelley (see [11]) that, in **ZF**, the Axiom of Choice (for short **AC**) is equivalent to the Tychonov axiom **T**: “Every family $(X_i)_{i \in I}$ of compact topological spaces has a compact product.” Here, a topological space X is *compact* if every family $(F_i)_{i \in I}$ of closed subsets of X satisfying the finite intersection property (*FIP*) has a non-empty intersection. However, some particular cases of the Tychonov axiom are provable in **ZF**, for example:

Remark 1. A finite product of compact spaces is compact (in **ZF**).

Say that the topological space X is *closely-compact* if there is a mapping Φ associating to every family $(F_i)_{i \in I}$ of closed subsets of X satisfying the *FIP* an element $\Phi((F_i)_{i \in I})$ of $\bigcap_{i \in I} F_i$: the mapping Φ is a *witness of closed-compactness* on X . Notice that a compact topological space X is closely-compact if and only if there exists a mapping Ψ associating to every non-empty closed subset F of X an element of F .

Example 1. Given a linear order (X, \leq) which is complete (every non-empty subset of X has a least upper bound), then the order topology on X is closely compact. In particular, the closed bounded interval $[0, 1]$ of \mathbb{R} is closely compact.

Proof. The space X is compact (the classical proof is valid in **ZF**). Moreover X is closely compact since one can consider the choice function associating to every non-empty closed subset its first element. \square

The following Theorem is provable in **ZF**:

Theorem ([8]). *Let α be an ordinal. If $(X_i, \Phi_i)_{i \in \alpha}$ is a family of witnessed closely-compact spaces, then $\prod_{i \in \alpha} X_i$ is closely-compact, and has a witness of closed-compactness which is definable from $(X_i, \Phi_i)_{i \in \alpha}$.*

Example 2. For every ordinal α , the product topological space $[0, 1]^\alpha$ is closely compact in **ZF**.

Given a set I , denote by $B_1(I)$ the set of $x = (x_i)_{i \in I} \in \mathbb{R}^I$ such that $\sum_{i \in I} |x_i| \leq 1$: then $B_1(I)$ is a closed subset of $[-1, 1]^I$. In this paper, we shall prove that $B_1(I)$ is compact using the *countable axiom of choice for finite subsets of I* (see Theorem 2 in Section 6.2). This enhances Corollary 1 of [14] and partially solves Question 2 in [14]. We shall deduce (see Corollary 3) that, if I is linearly orderable, every closed subset of $[0, 1]^I$ which is contained in $B_1(I)$ is closely compact. In particular, the closed unit ball of the Hilbert space $\ell^2(\mathbb{R})$ is compact in **ZF**, and this solves Question 3 of [14]. Notice that $\{0, 1\}^{\mathbb{R}}$ (and $[0, 1]^{\mathbb{R}}$) is not compact in **ZF** (see [12]). We shall also prove that *Eberlein* closed subsets of $[0, 1]^I$ are compact using the *countable axiom of choice for subsets of I* (see Corollary 4) of Section 7.4. This enhances Corollary 3 in [14] where the same result was proved using the axiom of *Dependent Choices* **DC**. This also solves Questions 4 and 5 thereof.

The paper is organized as follows: in Section 2 we review various consequences of **AC** (in particular the *countable axiom of choice* **AC** _{\mathbb{N}} and the axiom of choice *restricted to finite subsets* **AC** _{\mathbb{N}} ^{fin}) and the known links between them. In Section 3 we present definitions of uniform Eberlein spaces, strong Eberlein spaces and Eberlein spaces. In Section 4 we give some tools for compactness or sequential compactness in **ZF**. In Section 5, we recall the one-point compactification \hat{X} of a discrete space X , and we show that for every ordinal $\alpha \geq 1$, the closed-compactness of \hat{X}^α is equivalent to the axiom of choice restricted to finite subsets of X . Finally, in Section 6 (*resp.* 7) we prove that the countable axiom of choice for finite sets (*resp.* the countable axiom of choice) implies that uniform Eberlein spaces (*resp.* Eberlein spaces) are closely compact (*resp.* compact.) A basic tool for these two last Sections is a “dyadic representation” of elements of powers of $[0, 1]$ (see the Theorem in Section 6.1) which we found in [3, Lemma 1.1], and for which the authors cite [15].

2. SOME WEAK FORMS OF **AC**

In this Section, we review some weak forms of the Axiom of Choice which will be used in this paper and some known links between them. For detailed references and much information on this subject, see [10].

2.1. Restricted axioms of choice. Given a formula ϕ of set-theory with one free variable x , consider the following consequence of **AC**, denoted by **AC**(ϕ): “For every non-empty family $A = (A_i)_{i \in I}$ of non-empty sets such that $\phi[x/A]$ holds, then $\prod_{i \in I} A(i)$ is non-empty.”

Notation 1. In the particular case where the formula ϕ says that “ x is a mapping with domain I with values in some **ZF**-definable class \mathcal{C} ”, the statement **AC**(ϕ) is denoted by **AC** _{I} ^{\mathcal{C}} .

The statement $\forall I \mathbf{AC}_I^{\mathcal{C}}$ is denoted by **AC** ^{\mathcal{C}} . The statement **AC** _{I} ^{\mathcal{C}} where \mathcal{C} is the collection of all sets is denoted by **AC** _{I} .

Notation 2. For every set X , we denote by $\text{fin}(X)$ the set of finite subsets of X . We denote by fin the (definable) class of finite sets.

So, given a set X , **AC** ^{$\text{fin}(X)$} is the following statement: “For every non-empty family $(F_i)_{i \in I}$ of non-empty finite subsets of X , $\prod_{i \in I} F_i$ is non-empty.”, and **AC** ^{fin} is the following statement: “For every non-empty family $(F_i)_{i \in I}$ of non-empty finite sets, $\prod_{i \in I} F_i$ is non-empty.” The *countable Axiom of Choice* says that:

AC_ℕ: If $(A_n)_{n \in \mathbb{N}}$ is a family of non-empty sets, then there exists a mapping $f : \mathbb{N} \rightarrow \bigcup_{n \in \mathbb{N}} A_n$ associating to every $n \in \mathbb{N}$ an element $f(n) \in A_n$.

And the *countable Axiom of Choice for finite sets* says that:

AC_ℕ^{fin}: If $(A_n)_{n \in \mathbb{N}}$ is a family of finite non-empty sets, then there exists a mapping $f : \mathbb{N} \rightarrow \bigcup_{n \in \mathbb{N}} A_n$ associating to every $n \in \mathbb{N}$ an element $f(n) \in A_n$.

2.2. Well-orderable union of finite sets. Given an infinite ordinal α , and a class \mathcal{C} of sets, we consider the following consequence of **AC^C**:

Uwo_α^C: For every family $(F_i)_{i \in \alpha}$ of elements of \mathcal{C} , the set $\bigcup_{i \in \alpha} F_i$ is well-orderable.

Remark 2. **AC^{fin(X)}** implies **Uwo_α^{fin(X)}**.

2.3. Dependent Choices. The axiom of *Dependent Choices* says that:

DC: Given a non-empty set X and a binary relation R on X such that $\forall x \in X \exists y \in X xRy$, then there exists a sequence $(x_n)_{n \in \mathbb{N}}$ of X such that for every $n \in \mathbb{N}$, $x_n R x_{n+1}$.

Of course, **AC** \Rightarrow **DC** \Rightarrow **AC_ℕ** \Rightarrow **AC_ℕ^{fin}**. However, the converse statements are not provable in **ZF**, and **AC_ℕ^{fin}** is not provable in **ZF** (see references in [10]).

2.4. The “Tychonov” axiom. Given a class \mathcal{C} of compact topological spaces and a set I , we consider the following consequence of the Tychonov axiom:

T_I^C: Every family $(X_i)_{i \in I}$ of spaces belonging to the class \mathcal{C} has a compact product.

For example **T_ℕ^{fin(X)}** is the statement “Every sequence of finite discrete subsets of X has a compact product.”

Remark 3. (i) Given a set X , for every ordinal α ,

$$\mathbf{AC}^{fin(X)} \Rightarrow \mathbf{Uwo}_\alpha^{fin(X)} \Rightarrow \mathbf{T}_\alpha^{fin(X)} \Rightarrow \mathbf{AC}_\alpha^{fin(X)}$$

(ii) For every ordinal α , **Uwo_α^{fin}** \Leftrightarrow **T_α^{fin}** \Leftrightarrow **AC_α^{fin}**.

Proof. (i) **Uwo_α^{fin(X)}** \Rightarrow **T_α^{fin(X)}**: Given a family $(F_i)_{i \in \alpha}$ of finite subsets of X , the statement **Uwo_α^{fin(X)}** implies the existence of a family $(\Phi_i)_{i \in \alpha}$ such that for each $i \in \alpha$, the discrete space is closely compact with witness Φ_i . Using the Theorem of Section 1, it follows that $\prod_{i \in \alpha} F_i$ is (closely) compact. **T_α^{fin(X)}** \Rightarrow **AC_α^{fin(X)}**: one can use Kelley’s argument (see [11]). (ii) For **AC_α^{fin}** \Rightarrow **Uwo_α^{fin}**: given some family $(F_i)_{i \in \alpha}$ of finite non-empty sets, then, for each $i \in \alpha$, denote by $c_i := \{0..c_{i-1}\}$ the (finite) cardinal of F_i ; thus set G_i of one-to-one mappings from F_i to c_i is finitel, and, by **AC_α^{fin}**, the set $\prod_{i \in \alpha} G_i$ is non-empty. This implies a well-order on the set $\bigcup_{i \in \alpha} F_i$. \square

3. SOME CLASSES OF CLOSED SUBSETS OF $[0, 1]^I$

Notation 3. Let I be a set. Given some element $x = (x_i)_{i \in I} \in \mathbb{R}^I$, denote by $\text{supp}(x)$ the support $\{i \in I : x_i \neq 0\}$. Given some subset A of \mathbb{R} containing 0, denote by $A^{(I)}$ the set of elements of A^I with *finite* support.

We endow the space \mathbb{R}^I with the product topology, which we denote by \mathcal{T}_I .

3.1. Eberlein closed subsets of $[0, 1]^I$. Given a set I , we denote by $\ell^\infty(I)$ the Banach space of bounded mappings $f : I \rightarrow \mathbb{R}$, endowed with the “sup” norm. If I is infinite, we denote by $c_0(I)$ the closed subspace of $\ell^\infty(I)$ consisting of $f \in \ell^\infty(I)$ such that f converges to 0 according to the Fréchet filter on I (i.e. the set of cofinite subsets of I). Thus

$$\ell^0(I) := \{x = (x_i)_{i \in I} : \forall \varepsilon > 0 \exists F_0 \in \mathcal{P}_f(I) \forall i \in I \setminus F_0 |x_i| \leq \varepsilon\}$$

If I is finite, then we define $c_0(I) := \ell^\infty(I) = \mathbb{R}^I$.

Definition 1. A topological space F is *I-Eberlein* if F is a closed subset of $[0, 1]^I$ and if $F \subseteq c_0(I)$. A topological space X is *Eberlein* if X is homeomorphic with some *I-Eberlein* space.

Remark 4. Amir and Lindenstrauss ([1]) proved in **ZFC** that every weakly compact subset of a normed space is an Eberlein space. This result relies on the existence of a Markhushevich basis in every weakly compactly generated Banach space, and the proof of the existence of such a basis (see [7]) relies on (much) Axiom of Choice.

Remark 5. Consider the compact topological space $X := [0, 1]^\mathbb{N}$. Then, the closed subset X of $[0, 1]^\mathbb{N}$ is not \mathbb{N} -Eberlein. However, the mapping $f : X \rightarrow [0, 1]^\mathbb{N} \cap c_0(\mathbb{N})$ associating to each $x = (x_n)_{n \in \mathbb{N}} \in X$ the element $(\frac{x_n}{n+1})_{n \in \mathbb{N}}$ is continuous and one-to-one, so X is homeomorphic with the compact (hence closed) subset $f[X]$ of $[0, 1]^\mathbb{N} \cap c_0(\mathbb{N})$. It follows that X is homeomorphic with some \mathbb{N} -Eberlein space.

Proposition 1. (i) Every closed subset of a *I-Eberlein* (resp. *Eberlein*) space is *I-Eberlein* (resp. *Eberlein*).

(ii) Let $(I_n)_{n \in \mathbb{N}}$ be a sequence of pairwise disjoint sets, and denote by I the set $\sqcup_{n \in \mathbb{N}} I_n$. Let $(F_n)_{n \in \mathbb{N}}$ be a sequence of topological spaces such that each F_n is I_n -Eberlein. Then the closed subset $\prod_{n \in \mathbb{N}} F_n$ of $[0, 1]^I$ is homeomorphic with a *I-Eberlein* space.

Proof. (i) is trivial. We prove (ii). For every $n \in \mathbb{N}$, let $f_n : F_n \rightarrow [0, 1]^{I_n}$ be the mapping associating to each $x \in F_n$ the element $\frac{1}{n+1}f_n(x)$ of $[0, 1]^{I_n}$. Let $f := \prod_{n \in \mathbb{N}} f_n : \prod_{n \in \mathbb{N}} F_n \rightarrow [0, 1]^I$. Then f is one-to-one and continuous. Moreover, the subset $F := \text{Im}(f)$ of $[0, 1]^I$ is closed since F is the product $\prod_{n \in \mathbb{N}} \tilde{F}_n$ where for each $n \in \mathbb{N}$, \tilde{F}_n is the closed subset $\frac{1}{n+1} \cdot F_n$ of $[0, 1]^{I_n}$. Finally, it can be easily checked that $F \subseteq c_0(I)$. \square

3.2. Uniform Eberlein closed subsets of $[0, 1]^I$.

3.2.1. The ball $B_p(I)$, for $1 \leq p < +\infty$. For every real number $p \geq 1$, define as usual the normed space $\ell^p(I) := \{(x_i)_{i \in I} : \sum_i |x_i|^p < +\infty\}$ endowed with the norm $N_p : x = (x_i)_{i \in I} \mapsto (\sum_i |x_i|^p)^{1/p}$. We denote by $B_p(I)$ the large unit ball $\{x \in \mathbb{R}^I : \sum_i |x_i|^p \leq 1\}$ of $\ell^p(I)$. Notice that for $p = 1$ (resp. $1 < p < +\infty$) the topology induced by \mathcal{T}_I on $B_p(I)$ is the topology induced by the weak* topology $\sigma(\ell^1(I), \ell^0(I))$ (resp. the topology induced by the weak topology $\sigma(\ell^p(I), \ell^q(I))$ where $q = \frac{p}{p-1}$ is the conjugate of p). Also notice that for $1 \leq p < +\infty$, $B_p(I)$ is a closed subset of $[0, 1]^I$.

Proposition 2. If $1 \leq p < +\infty$, then $B_p(I)$ is homeomorphic with $B_1(I)$.

Proof. Consider the mapping $h_p : B_1(I) \rightarrow B_p(I)$ associating to every $x = (x_i)_i \in B_1(I)$ the family $(\text{sgn}(x_i)|x_i|^{1/p})_{i \in I}$. \square

It follows that for every $p, q \in [1, +\infty[$, spaces $B_p(I)$ and $B_q(I)$ are homeomorphic via $h_{p,q} := h_q \circ h_p^{-1} : B_p(I) \rightarrow B_q(I)$.

3.2.2. Uniform Eberlein spaces. Given a set I , and some real number $p \in [1, +\infty[$, we denote by $B_p^+(I)$ the positive ball of $\ell^p(I)$:

$$B_p^+(I) := \{x = (x_i)_{i \in I} \in [0, 1]^I : \sum_{i \in I} x_i^p \leq 1\}$$

Definition 2. A topological space F is *I-uniform Eberlein* if there exists a real number $p \in [1, +\infty[$ such that F is a closed subset of $B_p^+(I)$. A topological space X is *uniform Eberlein* if X is homeomorphic with some *I-uniform Eberlein* space.

Of course, every *I-uniform Eberlein* space is *I-Eberlein*. Moreover, using Proposition 2, every *I-uniform Eberlein* space is homeomorphic with a closed subset of $B_1^+(I)$.

Proposition 3. (i) For every set I , every closed subset of a *I-uniform Eberlein* space is *I-uniform Eberlein*.

(ii) Let $(I_n)_{n \in \mathbb{N}}$ be a sequence of pairwise disjoint sets, and denote by I the set $\sqcup_{n \in \mathbb{N}} I_n$. Let $(F_n)_{n \in \mathbb{N}}$ be a sequence of topological spaces such that each F_n is a I_n -uniform Eberlein space. Then the closed subset $F := \prod_{n \in \mathbb{N}} F_n$ of $[0, 1]^I$ is *I-uniform Eberlein*.

Proof. (i) is easy. The proof of (ii) is similar to the proof of Proposition 1-(ii). \square

In particular, the compact space $[0, 1]^{\mathbb{N}}$ (and thus every metrisable compact space) is \mathbb{N} -uniform Eberlein. For every set I , $B_1^+(I)^{\mathbb{N}}$ is $(I \times \mathbb{N})$ -uniform Eberlein.

Remark 6. Let $Z := \cap_{i \in I} \{(x, y) \in B_1^+(I) \times B_1^+(I) : x_i \cdot y_i = 0\}$: then Z is a closed subset of $B_1^+(I) \times B_1^+(I)$, and the mapping $- : Z \rightarrow B_1(I)$ is an homeomorphism; it follows that $B_1(I)$ is homeomorphic with a $(I \times \{0, 1\})$ -uniform Eberlein space.

3.2.3. Weakly closed bounded subsets of a Hilbert space.

Remark 7. Given a Hilbert space H with a Hilbert basis $(e_i)_{i \in I}$, then its closed unit ball (and thus every bounded weakly closed subset of H) is (linearly) homeomorphic with the uniform Eberlein space $B_2(I)$.

Consider the following statements (the first two ones were introduced in [5] and [13] and are consequences of the Alaoglu theorem):

- **A1:** The closed unit ball (and thus every bounded subset which is closed in the convex topology) of a uniformly convex Banach space is compact in the convex topology.
- **A2:** (Hilbert) The closed unit ball (and thus every bounded weakly closed subset) of a Hilbert space is weakly compact.
- **A3:** (Hilbert with hilbertian basis) For every set I , the closed unit ball of $\ell^2(I)$ is weakly compact.
- **A4:** For every sequence $(F_n)_{n \in \mathbb{N}}$ of finite sets, the closed unit ball of $\ell^2(\cup_{n \in \mathbb{N}} F_n)$ is weakly compact.

Of course, **A1** \Rightarrow **A2** \Rightarrow **A3** \Rightarrow **A4**.

Theorem ([8], [14]). (i) **AC** $_{\mathbb{N}} \Rightarrow$ **A1**.

(ii) **A1** $\not\Rightarrow$ **AC** $_{\mathbb{N}}$.

(iii) $\mathbf{A4} \Rightarrow \mathbf{AC}_{\mathbb{N}}^{\text{fin}}$.

In this paper, we will prove that the following statements are equivalent: $\mathbf{A3}$, $\mathbf{A4}$, $\mathbf{AC}_{\omega}^{\text{fin}}$ (see Corollary 2).

Question 1. Does $\mathbf{A2}$ imply $\mathbf{A1}$? Does $\mathbf{A3}$ imply $\mathbf{A2}$?

Remark 8. If a Hilbert space H has a well orderable dense subset, then H has a well orderable hilbertian basis, thus H is isometrically isomorphic with some $\ell^2(\alpha)$ where α is an ordinal. In this case, the closed unit ball of H endowed with the weak topology is homeomorphic with a closed subset of $[-1, 1]^{\alpha}$, so this ball is weakly compact.

3.3. Strongly Eberlein closed subsets of $[0, 1]^I$.

Definition 3. A topological space F is *I-strong Eberlein* if F is a closed subset of $[0, 1]^I$ which is contained in $\{0, 1\}^{(I)}$. A topological space X is *strong Eberlein* if X is homeomorphic with some *I-strong Eberlein* space.

Of course, every *I-strong Eberlein* set is *I-Eberlein*.

Remark 9. For every set I , every closed subset of a *I-strong Eberlein* space is *I-strong Eberlein*.

4. COMPACTNESS (IN \mathbf{ZF})

4.1. Lattices and filters. Given a lattice \mathcal{L} of subsets of a set X , say that a non-empty proper subset \mathcal{F} of \mathcal{L} is a *filter* if it satisfies the two following conditions:

- (i) $\forall A, B \in \mathcal{F}, A \cap B \in \mathcal{F}$
- (ii) $\forall A \in \mathcal{F}, \forall B \in \mathcal{L}, (A \subseteq B \Rightarrow B \in \mathcal{F})$

Say that an element $A \in \mathcal{L}$ is *\mathcal{F} -stationar* if for every $F \in \mathcal{F}$, $A \cap F \neq \emptyset$.

Remark 10. Let X be a topological space, let \mathcal{L} be a lattice of closed subsets of X , and let \mathcal{F} be a filter of \mathcal{L} . Let $K \in \mathcal{L}$. If K is a compact subset of X and if K is \mathcal{F} -stationar, then $\cap \mathcal{F}$ is non-empty.

Definition 4. Given a family $(X_i)_{i \in I}$ of topological spaces, and denoting by X the topological product of this family, a closed subset F of X is *elementary* if F is a finite union of sets of the form $\prod_{i \neq i_0} X_i \times C$ where $i_0 \in I$ and C is a closed subset of X_{i_0} .

Given a family $(X_i)_{i \in I}$ of topological spaces with product X , the set of elementary closed subsets of X is a lattice of subsets of X that we denote by \mathcal{L}_X . Notice that given a elementary closed subset F of X , and some subset J of I , the projection $p_J[F]$ is a closed subset of $\prod_{j \in J} X_j$.

4.2. Continuous image of a compact space. The following Proposition is easy:

Proposition 4. *Let X, Y be topological spaces and let $f : X \twoheadrightarrow Y$ be a continuous onto mapping. If X is compact (resp. closely-compact), then Y is also compact (resp. closely compact). If Φ is a witness of closed-compactness on X , then Y is closely-compact, and has a witness of closed-compactness which is definable from f and Φ .*

4.3. Sequential compactness. We denote by $[\mathbb{N}]^\omega$ the set of infinite subsets of \mathbb{N} .

Definition 5. A topological space X is *sequentially compact* if every sequence $(x_n)_{n \in \mathbb{N}}$ of X has an infinite subsequence which converges in X . A *witness of sequential compactness* on X is a mapping $\Phi : X^\mathbb{N} \rightarrow [\omega]^\omega \times X$ associating to each sequence $(x_n)_{n \in \mathbb{N}}$ of X an element $(A, l) \in [\omega]^\omega \times X$ such that $(x_n)_{n \in A}$ converges to l .

Example 3. If (X, \leq) is a complete linear order, then X is sequentially compact, with a witness definable from (X, \leq) : given a sequence $(x_n)_{n \in \mathbb{N}}$, build some infinite subset A of \mathbb{N} such that $(x_n)_{n \in A}$ is monotone; then if $(x_n)_{n \in A}$ is ascending (*resp.* descending), then $(x_n)_{n \in A}$ converges to $\sup_{n \in A} x_n$ (*resp.* $\inf_{n \in A} x_n$).

Example 4. Given an infinite set X , and some set $\infty \notin X$, consider the topology on $\tilde{X} := X \cup \{\infty\}$ generated by cofinite subsets of \tilde{X} and $\{\infty\}$. This topology is compact and T_1 but it is not T_2 . This topology is sequentially compact, and, given a point $a \in X$, there is a witness of sequential compactness which is definable from X and a : given a sequence $(x_n)_{n \in \mathbb{N}}$ of \tilde{X} , either the set of terms $\{x_n : n \in \mathbb{N}\}$ is finite, and then one can define by induction an infinite subset A of \mathbb{N} such that $\{x_n : n \in A\}$ is constant; else one can define by induction an infinite subset A of \mathbb{N} such that $\{x_n : n \in A\}$ is one-to-one, thus it converges to a (and also to every point in X).

Notice that the topology in Example 4 is the one used by Kelley (see [11]) to prove that “Tychonov implies \mathbf{AC} ”. The following Lemma is easy:

Lemma 1. *Let X, Y be two topological spaces and let $f : X \twoheadrightarrow Y$ be an onto continuous mapping which has a section j (for example if f is one-to-one). If X is sequentially compact, then Y is also sequentially compact. Moreover, if there is a witness Φ of sequential compactness on X , there also exists a witness of sequential compactness on Y which is definable from f, Φ and j .*

Lemma 2. *Let $(X_n, \phi_n)_{n \in \mathbb{N}}$ be a sequence of witnessed sequentially compact spaces. The space $\prod_{n \in \mathbb{N}} X_n$ is sequentially compact, and has a witness definable from $(X_n, \phi_n)_{n \in \mathbb{N}}$.*

Proof. Usual diagonalization. □

Example 5. If D is a countable set, then the topological space $[0, 1]^D$ is sequentially compact, a witness of sequential compactness being definable from every well order on D .

Say that a sequentially compact topological space X is *witnessable* if there exists a witness of sequential compactness on X . It follows from Lemma 2, that with $\mathbf{AC}_\mathbb{N}$, every sequence $(K_n)_{n \in \mathbb{N}}$ of witnessable sequentially compact spaces has a product which is sequentially compact.

4.4. $\mathbf{AC}_\mathbb{N}$ and countable products of compact spaces. Denote by T_ω^{comp} the following statement: “Every sequence of compact spaces has a compact product.” Then Kelley’s argument shows that $T_\omega^{comp} \Rightarrow \mathbf{AC}_\mathbb{N}$. However, it is an open question (see [4], [9]) to know whether $\mathbf{AC}_\mathbb{N}$ implies T_ω^{comp} .

Definition 6. A topological space X is ω -compact if every descending sequence $(F_n)_{n \in \mathbb{N}}$ of non-empty closed subsets of X has a non-empty intersection. Say that the space X is *cluster-compact* if every sequence $(x_n)_{n \in \mathbb{N}}$ of X has a *cluster point* i.e. the set $\bigcap_{n \in \mathbb{N}} \overline{\{x_k : k \geq n\}}$ is non-empty.

- Remark 11.* (i) Notice that sequentially compact \Rightarrow “cluster-compact”. Also notice that “ ω -compact” \Rightarrow “cluster-compact” and that the converse holds with $\mathbf{AC}_{\mathbb{N}}$ (see [9, Lemma 1]).
- (ii) Given a sequence $(K_n)_{n \in \mathbb{N}}$ of compact spaces, then, denoting by K the product of this family, K is compact *iff* K is ω -compact (see [9, Theorem 6]).
- (iii) If the product K of a sequence $(K_n)_{n \in \mathbb{N}}$ of compact spaces is sequentially compact, then $\mathbf{AC}_{\mathbb{N}}$ implies that K is compact.

Proposition 5. $\mathbf{AC}_{\mathbb{N}}$ is equivalent to the following statement: “Every sequence $(K_n)_{n \in \mathbb{N}}$ of witnessable sequentially compact spaces which are also compact has a compact product.”

Proof. \Rightarrow : Given a sequence $(K_n)_{n \in \mathbb{N}}$ of witnessable sequentially compact spaces which are also compact, then, using $\mathbf{AC}_{\mathbb{N}}$, one can choose a witness of sequential compactness on every space K_n . It follows by Lemma 2 that K is sequentially compact, whence K is compact by Remark 11-(iii).

\Leftarrow : We use Kelley’s argument (see [11]). Let $(A_n)_{n \in \mathbb{N}}$ be a sequence of non-empty sets. Consider some element $\infty \notin \bigcup_{n \in \mathbb{N}} A_n$, and for every $n \in \mathbb{N}$, denote by K_n the set $A_n \cup \{\infty\}$ endowed with the topology generated by $\{\infty\}$ and cofinite subsets of K_n (see Example 4). Then each K_n is compact and sequentially compact; moreover, given an element $a \in A_n$, there is a witness of sequential compactness on K_n which is definable from A_n, ∞ and a . So each K_n is a witnessable sequentially compact space. It follows from the hypothesis that the product $K := \prod_{n \in \mathbb{N}} K_n$ is compact. We end as in Kelley’s proof: for every $n \in \mathbb{N}$, let F_n be the closed set $A_n \times \prod_{i \neq n} K_i$. By compactness of K , the set $\bigcap_{n \in \mathbb{N}} F_n$ is non-empty. This yields an element of $\prod_{n \in \mathbb{N}} A_n$. \square

5. ONE-POINT COMPACTIFICATIONS AND RELATED SPACES

5.1. The one-point compactification of a set. Given a set X , we denote by \hat{X} the Alexandrov compactification of the (Hausdorff locally compact) discrete space X : $\hat{X} := X \cup \{\infty\}$ where ∞ is some set $\notin X$ (for example $\infty := \{x \in X : x \notin x\}$; if X is finite, then \hat{X} is discrete else open subsets of the space \hat{X} are subsets of X or cofinite subsets of \hat{X} containing ∞). Notice that the space \hat{X} is compact and Hausdorff in \mathbf{ZF} .

Example 6. Given a discrete topological space X , the one-point compactification \hat{X} of X is X -uniform Eberlein: consider the Hilbert space $\ell^2(X)$; and denote by $(e_i)_{i \in X}$ the canonical basis of the vector space $\mathbb{R}^{(X)}$; then the subspace $X = \{e_i : i \in X\}$ of $\mathbb{R}^{(X)}$ is discrete and the weakly closed and bounded subset $X \cup 0_{\mathbb{R}^X}$ is the one-point compactification \hat{X} of X .

5.2. Various notions of compactness for \hat{X}^α , α ordinal.

5.2.1. $\hat{X}^{\mathbb{N}}$ is sequentially compact.

Proposition 6. Let X be an infinite set.

- (i) The space \hat{X} is sequentially compact and has a witness of sequential compactness, definable from X .
- (ii) The space $\hat{X}^{\mathbb{N}}$ is sequentially compact with a witness definable from X .

Proof. (i) We define a witness Φ of sequential compactness on X as follows: given a sequence $x = (x_n)_{n \in \mathbb{N}}$ of \hat{X} , if the set $T := \{x_k : k \in \mathbb{N}\}$ is infinite, we build (by induction) some

infinite subset A of \mathbb{N} such that $\{x_k : k \in A\}$ is one-to-one, and we define $\Phi(x) := (A, \infty)$; else the set T is finite, so we build by induction some infinite subset A of \mathbb{N} such that the sequence $\{x_k : k \in A\}$ is a singleton $\{l\}$, and we define $\Phi(x) := (A, l)$.

(ii) We apply (i) and Lemma 2 in Section 4.3. \square

5.2.2. \mathbf{AC}^{fin} and closed-compactness.

Proposition 7. *Let X be a set.*

- (i) *There is a mapping associating to every non-empty closed subset F of \hat{X} , a finite non-empty closed subset \tilde{F} of F .*
- (ii) $\mathbf{AC}^{fin(X)} \Leftrightarrow$ *The space \hat{X} is closely compact.*

Proof. We may assume that X is infinite.

(i) Given a non-empty closed subset F of \hat{X} , define $\tilde{F} := \{\infty\}$ if $\infty \in F$ and $\tilde{F} := F$ if F is finite and $\infty \notin F$.

(ii) Use (i). \square

5.3. Spaces \hat{X}^α , α ordinal.

Remark 12. For every set X , the space \hat{X} is X -uniform Eberlein, so, given an ordinal α , \hat{X}^α is $X \times \alpha$ -uniform Eberlein (see Proposition 3-(ii)).

Proposition 8. *Let X be a set. Let α be an ordinal ≥ 1 .*

- (i) $\mathbf{T}_\alpha^{fin(X)} \Leftrightarrow$ *" \hat{X}^α is compact".*
- (ii) $\mathbf{AC}^{fin(X)} \Leftrightarrow$ *" \hat{X}^α is closely compact".*

Proof. (i) \Rightarrow : Let P be the topological product space \hat{X}^α . Let \mathcal{F} be a filter of the lattice \mathcal{L}_X of elementary closed subsets of P . We are going to define by transfinite recursion a family $(G_n)_{n \in \alpha}$ of finite subsets of \hat{X} such that, denoting for every $n \in \alpha$ by Z_n the elementary closed subset $G_n \times \hat{X}^{\alpha \setminus \{n\}}$ of P , the set $\mathcal{F} \cup \{Z_i : i < n\}$ satisfies the finite intersection property. Given some $n \in \alpha$, we define G_n in function of $(G_i)_{i < n}$ as follows: denote by \mathcal{G} the filter generated by $\mathcal{F} \cup \{Z_i : i < n\}$; since \hat{X} is compact, the closed subset $F_n := \cap p_{\{n\}}[\mathcal{G}]$ is non-empty, so let $G_n := \tilde{F}_n$ and let $Z_n := G_n \times \hat{X}^{\alpha \setminus \{n\}}$. Denote by $\tilde{\mathcal{F}}$ the filter generated by $\mathcal{F} \cup \{Z_i : i < \alpha\}$. Using $\mathbf{T}_\alpha^{fin(X)}$, the product space $F := \prod_{n \in \alpha} G_n$ is compact, and non-empty since $\mathbf{T}_\alpha^{fin(X)}$ implies $\mathbf{T}_\alpha^{fin(X)}$ (see Remark 3). Moreover, the closed subset F of P is $\tilde{\mathcal{F}}$ -stationnar: it follows from Remark 10 of Section 4.1 that $\cap \tilde{\mathcal{F}}$ is non-empty, whence $\cap \mathcal{F} \neq \emptyset$. \Leftarrow : Let $(F_i)_{i < \alpha}$ be a family of finite subsets of X , endowed with the discrete topology. Then $\prod_{i < \alpha} F_i$ is compact because it is a closed subset of the compact Hausdorff space \hat{X}^α .

(ii) \Rightarrow : Use Proposition 7-(ii) and the Theorem of Section 4.2. \Leftarrow : If \hat{X}^α is closely compact, then so is its continuous image \hat{X} , whence $\mathbf{AC}^{fin(X)}$ holds (using Proposition 7-(ii)). \square

5.4. Spaces $\sigma_n(X)$, n integer ≥ 1 .

Notation 4. Given a set X , for every integer $n \geq 1$, let

$$\sigma_n(X) := \{x \in \{0, 1\}^{(X)} : |\text{supp}(x)| \leq n\}$$

Thus $\sigma_n(X)$ is the set of elements of $\mathbb{R}^{(X)}$ with support having at most n elements. Notice that the space $\sigma_n(X)$ is strong Eberlein.

Remark 13. (i) The space $\sigma_1(X)$ is the one-point compactification of the discrete space X (thus $\sigma_1(X)$ is uniform Eberlein).

(ii) The mapping $\mathcal{U}_n : (\sigma_1(X))^n \rightarrow \sigma_n(X)$ associating to each (x_1, \dots, x_n) the set $\cup_{1 \leq i \leq n} x_i$ is continuous.

Proof. (i) Use Example 6. (ii): easy. □

5.4.1. Compactness and closed compactness of $\sigma_n(X)$.

Proposition 9. *Let X be a set, and let n be some integer ≥ 1 .*

- (i) *Both spaces $(\sigma_1(X))^n$ and $\sigma_n(X)$ are compact.*
- (ii) *With $\mathbf{AC}^{fin(X)}$, both spaces $(\sigma_1(X))^n$ and $\sigma_n(X)$ are closely compact (with witnesses of closed compactness definable from X , n and some choice function on non-empty finite subsets of X).*

Proof. The results (i) and (ii) for $\sigma_n(X)$ follow from the result on $(\sigma_1(X))^n$ thanks to Proposition 4 and the continuous onto mapping $\mathcal{U}_n : (\sigma_1(X))^n \rightarrow \sigma_n(X)$ defined in Remark 13. The result (i) for $(\sigma_1(X))^n$ comes from Remark 1. We prove (ii) for $(\sigma_1(X))^n$: with $\mathbf{AC}^{fin(X)}$, $\sigma_1(X)$ is closely compact, so the space $(\sigma_1(X))^n$ is also closely compact because it is a finite power of a closely compact space (use the Theorem of Section 1). □

5.4.2. Sequential compactness of $\prod_{n \in \mathbb{N}} \sigma_n(X)$.

Proposition 10. *Let X be a set, and let n be some integer ≥ 1 .*

- (i) *The space $\sigma_n(X)$ is sequentially compact, with a witness definable from X and n .*
- (ii) *The space $\prod_{k \in \mathbb{N}} \sigma_k(X)$ is sequentially compact, with a witness definable from X .*

Proof. (i) The proof is by induction on n . For $n = 1$, we already know that $\sigma_1(X) = \hat{X}$ is sequentially compact with a witness definable from X (use Proposition 6-(i)). We now assume that for some integer $n \geq 1$, each space $\sigma_k(X)$ ($1 \leq k \leq n$) is sequentially compact with a witness Φ_k definable from X and k . Let $(F_k)_{k \in \mathbb{N}}$ be a sequence of $\sigma_{n+1}(X)$. For every $\nu \in \mathbb{N}$, let $\mathcal{A}_\nu := \{A \in [\mathbb{N}]^\omega : \forall i \neq j \in A \mid |F_i \cap F_j| = \nu\}$. Let ν_0 be the first element of \mathbb{N} such that the set \mathcal{A}_{ν_0} is non-empty. One can build by induction some element $A \in \mathcal{A}_{\nu_0}$, which is definable from X and $(F_n)_{n \in \mathbb{N}}$. If $\nu = 0$, then the subsequence $(F_n)_{n \in A}$ converges to ∞ . Else, there exists $a \in X$ such that the set $D_a := \{n \in A : a \in F_n\}$ is infinite. Build by induction some infinite subset B of A such that there exists some element $a \in X$ satisfying $\forall n \in B \ a \in F_n$. Let R be the non-empty finite set $\cap_{n \in B} F_n$; let p be the cardinal of R . The sequence $(F_n \setminus R)_{n \in B}$ lives in $\sigma_{n+1-p}(X)$ thus, using the witness Φ_{n+1-p} , it has an infinite subsequence $(F_n \setminus R)_{n \in C}$ which converges to some $L \in \sigma_{n+1-p}(X)$. It follows that $(F_n \cup R)_{n \in C}$ converges to $L \cup R$ in $\sigma_{n+1}(X)$.

(ii) Use Proposition 6-(ii) or Lemma 2. □

5.4.3. $\mathbf{AC}^{fin(X)}$ and closed-compactness of the space $\prod_{n \in \mathbb{N}} \sigma_n(X)$.

Theorem 1. *Let X be a set.*

- (i) $\mathbf{AC}^{fin(X)} \Leftrightarrow \text{“}\prod_{n \in \mathbb{N}} \sigma_n(X) \text{ is closely compact”}$.
- (ii) $\mathbf{T}_\mathbb{N}^{fin(X)} \Leftrightarrow \text{“}\prod_{n \in \mathbb{N}} \sigma_n(X) \text{ is compact”}$.

Proof. We may assume that X is infinite. In both cases, we use Proposition 4 and the fact that the space $\prod_{n \in \mathbb{N}} \sigma_n(X)$ is a continuous image of $\prod_{n \in \mathbb{N}} \sigma_1(X)^n$, which is homeomorphic with $\hat{X}^{\mathbb{N}}$.

(i) \Rightarrow : with $\mathbf{AC}^{fin(X)}$, $\hat{X}^{\mathbb{N}}$ is closely compact (see Proposition 8), and so is its continuous image $\prod_{n \in \mathbb{N}} \sigma_n(X)$. \Leftarrow : if $\prod_{n \in \mathbb{N}} \sigma_n(X)$ is closely compact, then so is its continuous image $\sigma_1(X) = \hat{X}$, thus $\mathbf{AC}^{fin(X)}$ holds.

(ii) \Rightarrow : Using Proposition 8-(i), $\mathbf{T}_{\mathbb{N}}^{fin(X)}$ implies that “ $\hat{X}^{\mathbb{N}}$ is compact”. Using Remark 13, it follows that “ $\prod_{n \in \mathbb{N}} \sigma_n(X)$ is compact”. \Leftarrow : if $\prod_{n \in \mathbb{N}} \sigma_n(X)$ is compact, then its closed subset $\sigma_1(X)^{\mathbb{N}}$ is also compact, thus $\mathbf{T}_{\mathbb{N}}^{fin(X)}$ holds by Proposition 8-(i). \square

6. $\mathbf{AC}^{fin(I)}$ AND CLOSED COMPACTNESS OF $B_1(I)$

6.1. Dyadic representations.

Notation 5 (binary expansion of a real number). For every $n \in \mathbb{N}$, let $\varepsilon_n := \frac{1}{2^{n+1}}$. Then the mapping $\phi : \{0, 1\}^{\mathbb{N}} \rightarrow [0, 1]$ associating to every $(x_n)_{n \in \mathbb{N}}$ the real number $\sum_n \varepsilon_n x_n$ is continuous (a uniformly convergent series of continuous functions), onto, and ϕ has a (definable) section.

Theorem. [3, Lemma 1.1] *Let I be a set, and for every $n \in \mathbb{N}$, let $I_n := \{n\} \times I$. Let F be a closed subset of $[0, 1]^I$. Consider the power mapping $g := \phi^I : \{0, 1\}^{\mathbb{N} \times I} \rightarrow [0, 1]^I$. For every $n \in \mathbb{N}$, let $j_n : \{0, 1\}^{I_n} \rightarrow \{0, 1\}^{\mathbb{N} \times I}$ be the canonical inclusion mapping. Let $Z := g^{-1}[F]$ and, for every $n \in \mathbb{N}$, let $Z_n := j_n^{-1}[Z]$: thus Z is a closed subset of $\prod_{n \in \mathbb{N}} Z_n$ and $g : Z \rightarrow F$ is continuous, onto, with a definable section*

(i) *If $F \subseteq B_1^+(I)$, then for every $n_0 \in \mathbb{N}$, $Z_{n_0} \subseteq \sigma_{M_{n_0}}(I_{n_0})$ where $M_{n_0} := \lfloor \frac{1}{\varepsilon_{n_0}} \rfloor$ (the integral part of $\frac{1}{\varepsilon_{n_0}}$), thus F is the continuous image of some closed subset of $\hat{I}^{\mathbb{N}}$.*

(ii) *If $F \subseteq \ell^0(I)$, then for every $n_0 \in \mathbb{N}$, $Z_{n_0} \subseteq \{0, 1\}^{(I_{n_0})}$*

Proof. Let $n_0 \in \mathbb{N}$ and let $(x_i^{n_0})_{i \in I} \in Z_{n_0}$; let $x = (x_i^n)_{n \in \mathbb{N}, i \in I} \in Z$ such that $j_{n_0}((x_i^{n_0})_{i \in I}) = x$.

(i) Since $F \subseteq B_1^+(I)$, $\sum_{i \in I} \varepsilon_n x_i^n \leq 1$, thus $\sum_{i \in I} \varepsilon_{n_0} x_i^{n_0} \leq 1$; it follows that the set $\{i \in I : x_i^{n_0} = 1\}$ has a cardinal $\leq \frac{1}{\varepsilon_{n_0}}$.

(ii) Since $F \subseteq \ell^0(I)$, $(x_i^{n_0})_{i \in I} \in \ell^0(I) \cap \{0, 1\}^I$ thus $Z_{n_0} \subseteq \{0, 1\}^{(I_{n_0})}$. \square

Remark 14. The mapping $- : B_1^+(I) \times B_1^+(I) \rightarrow B_1(I)$ is continuous and onto thus $B_1(I)$ is also the continuous image of a closed subset of $\hat{I}^{\mathbb{N}}$.

Remark 15. Aviles ([2]) proved that $B_1^+(I)$ -and thus $B_1(I)$ - is a continuous image of $\hat{I}^{\mathbb{N}}$ (and not only of a closed subset of $\hat{I}^{\mathbb{N}}$).

6.2. Another equivalent of $\mathbf{AC}^{fin(I)}$.

Theorem 2. *Let I be a set.*

(i) $\mathbf{AC}^{fin(I)} \Leftrightarrow$ “ $B_1(I)$ is closely compact.” Moreover, a witness of closed compactness on $B_1(I)$ is definable from I and a choice function for non-empty finite subsets of I and conversely.

(ii) $\mathbf{T}_{\mathbb{N}}^{fin(I)}$ implies that $B_1(I)$ is compact.

(iii) The space $B_1(I)$ is sequentially compact, with a witness definable from I .

Proof. Using the previous Theorem and Remark 14, consider some sequence $(M_n)_{n \in \mathbb{N}}$, some closed subset Z of $\prod_{n \in \mathbb{N}} \sigma_{M_n}(I)$ and some continuous onto mapping $g : Z \rightarrow B_1(I)$, with a (definable) section.

(i) \Rightarrow : Using $\mathbf{AC}^{fin(I)}$ and Theorem 1-(i), Z is closely compact thus $B_1(I) = g[F]$ is closely compact. \Leftarrow : If $B_1(I)$ is closely compact, then \hat{I} (which is a closed subset of $B_1(I)$ -see Example 6 in Section 5.1-) is also closely compact.

(ii) Using $\mathbf{T}_{\mathbb{N}}^{fin(I)}$ and Theorem 1-(ii), Z is compact thus $B_1(I) = g[F]$ is also compact.

(iii) By Proposition 10-(ii), Z is sequentially compact and g is continuous with a section, thus Lemma 1 implies that $B_1(I)$ is also sequentially compact with a witness definable from I . \square

Corollary 1. *Given a set I , the following statements are equivalent:*

- (i) $\mathbf{Uwo}_{\mathbb{N}}^{fin(I)}$
- (ii) $\mathbf{T}_{\mathbb{N}}^{fin(I)}$
- (iii) *The space $B_1(I)$ is compact.*
- (iv) *For every sequence $(F_n)_{n \in \mathbb{N}}$ of finite subsets of I , the space $B_1(\cup_{n \in \mathbb{N}} F_n)$ is compact.*

Proof. (i) \Rightarrow (ii) is easy and (ii) \Rightarrow (iii) follows from Theorem 2. (iii) \Rightarrow (iv) is easy. We show that (iv) \Rightarrow (i). The idea of the implication is in [8, th. 9 p. 16]: we sketch it for sake of completeness. Let $(F_n)_{n \in \mathbb{N}}$ be a disjoint sequence of non-empty finite sets of I . Let us show that $D := \cup_{n \in \mathbb{N}} F_n$ is countable. The Hilbert spaces $H := \ell^2(D)$ and $\oplus_{\ell^2(\mathbb{N})} \ell^2(F_n)$ are isometrically isomorph. For every $n \in \mathbb{N}$, let $\varepsilon_n : |F_n| \rightarrow]0, 1[$ be a strictly increasing mapping such that $\sum_{n \in \mathbb{N}} \sum_{0 \leq i < |F_n|} \varepsilon_n(i)^2 = 1$. For every $n \in \mathbb{N}$, let $\tilde{F}_n := \{x \in B_H : \forall m \neq n, x|_{F_m} = 0 \text{ and } x|_{F_n} \text{ is one-to-one from } F_n \text{ onto } rg(\varepsilon_n)\}$. Then each \tilde{F}_n is a weakly closed subset of $B_{\ell^2(D)}$ and the sequence $(\tilde{F}_n)_{n \in \mathbb{N}}$ satisfies the finite intersection property. The compactness of $B_2(D)$ implies that $Z := \cap_{n \in \mathbb{N}} \tilde{F}_n$ is non-empty. Given an element $f = (f_n)_{n \in \mathbb{N}}$ of Z , each f_n defines a well-order on the finite set F_n , thus $\cup_{n \in \mathbb{N}} F_n$ is countable. \square

Remark 16. For $I = \mathcal{P}(\mathbb{R})$, none of the equivalent statements in Corollary 1 is provable in \mathbf{ZF} . Indeed, there is a model of \mathbf{ZF} where there exists a sequence $(P_n)_{n \in \mathbb{N}}$ of pairs of subsets of \mathbb{R} such that $\prod_{n \in \mathbb{N}} P_n$ is empty.

Thus, the statement “The closed unit ball of $\ell^2(\mathcal{P}(\mathbb{R}))$ is weakly compact.” is not provable in \mathbf{ZF} .

Corollary 2. *The following statements are equivalent: **A3**, **A4**, $\mathbf{AC}_{\mathbb{N}}^{fin}$.*

6.3. Consequences.

Corollary 3. *If a set I is linearly orderable, then $B_1(I)$ is compact.*

Proof. If I is linearly orderable, then $\mathbf{AC}^{fin(I)}$ holds. \square

For every ordinal α , the set $\mathcal{P}(\alpha)$ is linearly orderable, thus $\mathbf{AC}^{fin(\mathcal{P}(\alpha))}$ holds. In particular, \mathbb{R} is equipotent with $\mathcal{P}(\mathbb{N})$ so the closed unit ball of $\ell^2(\mathbb{R})$ is closely compact. This solves Question 3 of [14].

Question 2. Does \mathbf{AC}^{fin} imply **A2**? What is the power of the statement “Every Hilbert space has a hilbertian basis”? Is this statement provable in \mathbf{ZF} ? Does it imply \mathbf{AC} ?

7. $\mathbf{AC}_{\mathbb{N}}$ AND EBERLEIN SPACES

Given a set I , a closed subset F of $[0, 1]^I$ is I -Corson if every element $x \in F$ has a countable support.

7.1. Sequential compactness of I -Eberlein spaces. Given a set I , denote by $\text{count}(I)$ the set of finite or countable subsets of I . Consider the following consequence of $\mathbf{AC}_{\mathbb{N}}$:

$\mathbf{Uwo}_{\mathbb{N}}^{\text{count}(I)}$: “Every countable union of countable subsets of I is countable.”

Proposition 11. *Let I be a set and let F be a closed subset of $[0, 1]^I$.*

- (i) $\mathbf{Uwo}_{\mathbb{N}}^{\text{fin}(I)}$ implies that:
 - if $F \subseteq [0, 1]^{(I)}$, then F is sequentially compact.
 - if $F \subseteq c_0(I)$, then F is I -Corson.
- (ii) $\mathbf{Uwo}_{\mathbb{N}}^{\text{count}(I)}$ implies that if F is I -Corson, then F is sequentially compact. Thus $\mathbf{AC}_{\mathbb{N}}$ implies that every Eberlein space is sequentially compact.
- (iii) $\mathbf{AC}^{\text{fin}(I)}$ implies that if $F \subseteq [0, 1]^{(I)}$, then F is sequentially compact and has a witness of sequential compactness.

Proof. (i) If $F \subseteq [0, 1]^{(I)}$, then F is sequentially compact using the fact that $[0, 1]^{\mathbb{N}}$ is sequentially compact (see Example 5). \square

7.2. Countable product of finitely restricted spaces.

Theorem 3. *Let F be a closed subset of $[0, 1]^I$ which is contained in $[0, 1]^{(I)}$. Then $\mathbf{AC}_{\mathbb{N}}$ implies that F is compact. In particular, $\mathbf{AC}_{\mathbb{N}}$ implies that every strong Eberlein space is compact.*

Proof. For every $n \in \mathbb{N}$, recall that the subset $\sigma_n(I) := \{x \in [0, 1]^{(I)} : |\text{supp}(x)| \leq n\}$ is compact (see Proposition 9). Let \mathcal{F} be a filter of the lattice of closed subsets of F . If there exists an integer n such that $\sigma_n(I)$ is \mathcal{F} -stationary, then $\bigcap \mathcal{F}$ is non-empty by compactness of $\sigma_n(I)$ and using Remark 10. Else, using $\mathbf{AC}_{\mathbb{N}}$, consider a sequence $(F_n)_{n \in \mathbb{N}}$ of closed subsets of F belonging to \mathcal{F} such that for every $n \in \mathbb{N}$, $F_n \cap \sigma_n(I) = \emptyset$. Re-using $\mathbf{AC}_{\mathbb{N}}$, choose for every $n \in \mathbb{N}$ an element $x_n \in F_n$. A new use of $\mathbf{AC}_{\mathbb{N}}$ and Proposition 11-(i) implies the existence of some infinite subset A of \mathbb{N} such that $(x_n)_{n \in A}$ converges to some element $x \in F$. Then, for every $n \in \mathbb{N}$, $x \in F_n$ (which is disjoint with $\sigma_n(I)$) so the element x of F has an infinite support: this is contradictory! \square

Remark 17. It does not seem provable in \mathbf{ZF} that every closed subset of $[0, 1]^I$ contained in $[0, 1]^{(I)}$ is compact, or has a witness of sequential compactness.

7.3. Countable products of strong Eberlein spaces.

Theorem 4. *Let $(I_n)_{n \in \mathbb{N}}$ be a sequence of pairwise disjoint sets and let $I := \bigcup_{n \in \mathbb{N}} I_n$. For every $n \in \mathbb{N}$, let F_n be a closed subset of $\{0, 1\}^{(I_n)}$. Let F be the closed subset $\prod_{n \in \mathbb{N}} F_n$ of $\{0, 1\}^I$. Then $\mathbf{AC}_{\mathbb{N}}$ implies that F is sequentially compact and compact.*

Proof. Using $\mathbf{AC}_{\mathbb{N}}$, F is sequentially compact: given a sequence $(x_n)_{n \in \mathbb{N}}$ of F , $\mathbf{AC}_{\mathbb{N}}$ implies that for every $n \in \mathbb{N}$, the support D_n of x_n is countable, thus re-using $\mathbf{AC}_{\mathbb{N}}$, the set $D := \bigcup_{n \in \mathbb{N}} D_n$ is also countable; since each x_n belongs to $[0, 1]^D \times \{0\}^{I \setminus D}$, and since $[0, 1]^D$ is sequentially compact (see Example 5), it follows that $(x_n)_{n \in \mathbb{N}}$ has an infinite subsequence

which converges in F . Using $\mathbf{AC}_{\mathbb{N}}$ and Theorem 3, each F_n is compact. Using $\mathbf{AC}_{\mathbb{N}}$ and Remark 11-(iii), it follows that F is compact. \square

7.4. $\mathbf{AC}_{\mathbb{N}}$ and Eberlein closed subsets of $[0, 1]^I$.

Corollary 4. $\mathbf{AC}_{\mathbb{N}}$ implies that every Eberlein space is both sequentially compact and compact.

Proof. Let X be an Eberlein space. Then X is sequentially compact by Proposition 11-(i). Let I be a set such that X is homeomorphic with a closed subset F of $[0, 1]^I$, with $F \subseteq [0, 1]^{(I)}$. Using the Theorem of Section 6.1, there exists a family $(Z_k)_{k \in \mathbb{N}}$ such that for each $k \in \mathbb{N}$, Z_k is I -strong Eberlein, and such that F is the continuous image of a closed subset Z of $\prod_{k \in \mathbb{N}} Z_k$. Using $\mathbf{AC}_{\mathbb{N}}$ and Theorem 4, the space $\prod_{k \in \mathbb{N}} Z_k$ is sequentially compact. With $\mathbf{AC}_{\mathbb{N}}$, each Z_k is compact; with Remark 11-(iii) and $\mathbf{AC}_{\mathbb{N}}$, it follows that $\text{prod}_{k \in \mathbb{N}} Z_k$ (and thus its continuous image F) is also compact. \square

Recall that (see Theorem 2-(iii)) every I -uniform Eberlein space is sequentially compact, with a witness definable from I .

Question 3. Does $\mathbf{AC}_{\mathbb{N}}$ implies that every Eberlein space is sequentially compact with a witness?

7.5. Convex-compactness and the Hahn-Banach property. Given a set I , say that a subset F of \mathbb{R}^I is *convex-compact* if for every set \mathcal{C} of closed convex subsets of \mathbb{R}^I such that $\{F \cap C : C \in \mathcal{C}\}$ satisfies the *FIP*, $C \cap \bigcap \mathcal{C}$ is non-empty; moreover, if there is a mapping associating to each such \mathcal{C} an element of $C \cap \bigcap \mathcal{C}$, then say that F is *closely convex-compact*. Given a topological vector space E , say that E satisfies the *continuous Hahn-Banach property* if, for every continuous sublinear functional $p : E \rightarrow \mathbb{R}$, for every vector subspace F of E , and every linear functional $f : F \rightarrow \mathbb{R}$ such that $f \leq p|_F$, there exists a linear functional $g : E \rightarrow \mathbb{R}$ that extends f and such that $g \leq p$. Moreover, if there is a mapping associating to each such f some g satisfying the previous conditions, then say that E satisfies the *effective continuous Hahn-Banach property*.

Theorem ([8], [6]). *Given a set I , the normed space $\ell^0(I)$ satisfies the effective continuous Hahn-Banach property. For every real number $p \in [1, +\infty[$, $\ell^p(I)$ satisfies the effective continuous Hahn-Banach property.*

Corollary. *For every set I , and every real number $p \in [1, +\infty[$, $B_p(I)$ is closely convex-compact.*

Proof. The continuous dual of $\ell^0(I)$ is (isometrically isomorphic with) $\ell^1(I)$ and, for every $p \in]1, +\infty[$, the continuous dual of $\ell^p(I)$ is $\ell^q(I)$ where $\frac{1}{p} + \frac{1}{q} = 1$ thus $1 < q < +\infty$. We end the proof using the fact (see [8]) that if a normed space E satisfies the (effective) continuous Hahn-Banach property, then the closed unit ball of the continuous dual E' is (*closely*) weak* compact. \square

Question 4. Given a set I , and a closed convex subset C of $[0, 1]^I$ which is I -Eberlein, is C convex-compact (in \mathbf{ZF})?

Question 5. Same question if the closed convex subset C of $[0, 1]^I$ is I -Corson.

REFERENCES

- [1] D. Amir and J. Lindenstrauss. The structure of weakly compact sets in Banach spaces. *Ann. of Math.* (2), 88:35–46, 1968.
- [2] A. Avilés. The unit ball of the Hilbert space in its weak topology. *Proc. Amer. Math. Soc.*, 135(3):833–836 (electronic), 2007.
- [3] Y. Benyamini, M. E. Rudin, and M. Wage. Continuous images of weakly compact subsets of Banach spaces. *Pacific J. Math.*, 70(2):309–324, 1977.
- [4] N. Brunner. Products of compact spaces in the least permutation model. *Z. Math. Logik Grundlag. Math.*, 31(5):441–448, 1985.
- [5] C. Delhommé and M. Morillon. Dependent choices and weak compactness. *Notre Dame J. Formal Logic*, 40(4):568–573, 1999.
- [6] J. Dodu and M. Morillon. The Hahn-Banach property and the axiom of choice. *Math. Log. Q.*, 45(3):299–314, 1999.
- [7] M. Fabian, P. Habala, P. Hájek, V. Montesinos Santalucía, J. Pelant, and V. Zizler. *Functional analysis and infinite-dimensional geometry*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 8. Springer-Verlag, New York, 2001.
- [8] Fossey, J. and Morillon, M. The Baire category property and some notions of compactness. *J. Lond. Math. Soc., II. Ser.*, 57(1):1–19, 1998.
- [9] P. Howard, K. Keremedis, J. E. Rubin, and A. Stanley. Compactness in countable Tychonoff products and choice. *MLQ Math. Log. Q.*, 46(1):3–16, 2000.
- [10] P. Howard and J. E. Rubin. *Consequences of the Axiom of Choice.*, volume 59. American Mathematical Society, Providence, RI, 1998.
- [11] Kelley. The Tychonov product theorem implies the Axiom of Choice. *Fund. Math.*, pages 75–76, 1950.
- [12] K. Keremedis. The compactness of $2^{\mathbf{R}}$ and the axiom of choice. *MLQ Math. Log. Q.*, 46(4):569–571, 2000.
- [13] M. Morillon. James sequences and Dependent Choices. *Math. Log. Quart.*, 51(2):171–186, 2005.
- [14] Morillon, M. Countable choice and compactness. *Topology and applications*, 2008.
- [15] P. Simon. On continuous images of Eberlein compacts. *Comment. Math. Univ. Carolinae*, 17(1):179–194, 1976.

ERMIT, DÉPARTEMENT DE MATHÉMATIQUES ET INFORMATIQUE, UNIVERSITÉ DE LA RÉUNION, 15
 AVENUE RENÉ CASSIN - BP 7151 - 97715 SAINT-DENIS MESSAG. CEDEX 9 FRANCE
E-mail address, Marianne Morillon: `mar@univ-reunion.fr`
URL: `http://personnel.univ-reunion.fr/mar`