



**HAL**  
open science

## On the Mullineux involution for Ariki-Koike algebras

Nicolas Jacon, Cédric Lecouvey

► **To cite this version:**

Nicolas Jacon, Cédric Lecouvey. On the Mullineux involution for Ariki-Koike algebras. 2008. hal-00269840v2

**HAL Id: hal-00269840**

**<https://hal.science/hal-00269840v2>**

Preprint submitted on 23 Apr 2008 (v2), last revised 15 Sep 2008 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON THE MULLINEUX INVOLUTION FOR ARIKI-KOIKE ALGEBRAS

NICOLAS JACON AND CÉDRIC LECOUEY

ABSTRACT. This note is concerned with a natural generalization of the Mullineux involution for Ariki-Koike algebras. By using a result of Fayers together with previous results by the authors, we give an efficient algorithm for computing this generalized Mullineux involution. Our algorithm notably does not involve the determination of paths in affine crystals.

## 1. INTRODUCTION

It is known that the set of simple modules of the symmetric group over a field of characteristic  $p$  can be labelled by the set  $\mathcal{P}_p$  of  $p$ -regular partitions, that is, the partitions where no non zero part is repeated  $p$  or more times. The tensor product of one of these simple modules with the sign representation is again simple. This gives a natural involution  $I$  on  $\mathcal{P}_p$ . In 1979, Mullineux [20] introduced a combinatorial algorithm yielding an involution  $m$  on  $\mathcal{P}_p$  and conjectured the equality  $m = I$ . In 1994, Kleshchev [16] obtained the first combinatorial description of the involution  $I$  by using sequences of  $p$ -regular partitions. In particular, the corresponding algorithm  $k$  is different from the combinatorial procedure  $m$  of Mullineux. Finally, the conjecture  $m = I$  (and thus the equality  $m = k$ ) was proved by Ford and Kleshchev in [12] (see also [5] and [7] for different proofs). For the sake of simplicity, we will refer to the involution  $I=m$  as the Mullineux involution in the sequel.

A natural quantum analogue of this problem has been considered by Brundan [6] in the context of the Hecke algebra of type  $A$  (see also [7] and [17]). The Mullineux involution can then be interpreted as a skew-isomorphism of  $A_{e-1}^{(1)}$ -crystals, that is as an isomorphism of oriented graphs switching the sign of each arrow. The resulting algorithm then coincides with that introduced by Kleshchev in [16].

In this note, we study an analogue of the Mullineux involution for the Ariki-Koike algebras (also called cyclotomic Hecke algebras of type  $G(l, 1, n)$  in the literature). Let  $l, n$  be positive integers,  $R$  be a field of arbitrary characteristic and consider  $(q, Q_0, Q_1, \dots, Q_{l-1})$  an  $l + 1$ -tuple of invertible elements in  $R$ . The Ariki-Koike algebra  $\mathcal{H}_{R,n} := \mathcal{H}_{R,n}(q; Q_0, Q_1, \dots, Q_{l-1})$  over  $R$  is the unital associative  $R$ -algebra presented by:

---

*Date:* April, 2008.

- generators:  $T_0, T_1, \dots, T_{n-1}$ ,
- relations:

$$\begin{aligned}
T_0 T_1 T_0 T_1 &= T_1 T_0 T_1 T_0, \\
T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1} \quad (i = 1, \dots, n-2), \\
T_i T_j &= T_j T_i \quad (|j - i| > 1), \\
(T_0 - Q_0)(T_0 - Q_1) \dots (T_0 - Q_{l-1}) &= 0, \\
(T_i - q)(T_i + 1) &= 0 \quad (i = 1, \dots, n-1).
\end{aligned}$$

If  $l = 1$  (resp.  $l = 2$ ), we obtain a Hecke algebra of type  $A$  (resp. of type  $B$ ). To study these algebras, by a theorem of Dipper, James and Mathas [8], it is sufficient to consider the case where  $Q_0 = q^{s_0}$ ,  $Q_1 = q^{s_1}$ , ...,  $Q_{l-1} = q^{s_{l-1}}$  for  $\mathbf{s} = (s_0, s_1, \dots, s_{l-1})$  an  $l$ -tuple of integers. The algebra  $\mathcal{H}_{R,n}$  is then a cellular algebra in the sense of Graham and Lehrer [11]. As a consequence, we can define Specht modules which are indexed by the  $l$ -partitions of rank  $n$ . Recall that an  $l$ -partition  $\boldsymbol{\lambda}$  of rank  $n$  is a sequence of  $l$  partitions  $\boldsymbol{\lambda} = (\lambda^{(0)}, \dots, \lambda^{(l-1)})$  such that  $\sum_{k=0}^{l-1} |\lambda^{(k)}| = n$ . We denote by

$\Pi_{l,n}$  the set of  $l$ -partitions of rank  $n$ . Hence, to each  $\boldsymbol{\lambda} \in \Pi_{l,n}$  is associated a Specht module  $S^\boldsymbol{\lambda}$ . Let  $e$  be the multiplicative order of  $q$  (we put  $e = \infty$  if  $q$  is generic). The Specht modules are in general not irreducible. Nevertheless, there exists a natural bilinear form,  $\mathcal{H}_{R,n}$ -invariant, on each of these modules and an associated radical such that the quotients  $D^\boldsymbol{\lambda} := S^\boldsymbol{\lambda} / \text{rad}(S^\boldsymbol{\lambda})$  are 0 or irreducible. Moreover, the non-zero  $D^\boldsymbol{\lambda}$  provide a complete set of non isomorphic simple modules. Let  $\Phi_e^\mathbf{s}(n)$  be the set of  $l$ -partitions such that  $D^\boldsymbol{\lambda}$  is non zero (where we set  $\mathbf{s} = (s_0 \pmod{e}, \dots, s_{l-1} \pmod{e})$ ). It has been shown by Ariki [1] and Ariki-Mathas [3], that this set corresponds to the set of Kleshchev  $l$ -partitions. These  $l$ -partitions are generalizations of the  $e$ -regular partitions. They appear as the vertices of a distinguished realization of the abstract  $A_{e-1}^{(1)}$ -crystal  $B_e(\Lambda_\mathbf{s})$  associated to the irreducible  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ -module of highest weight  $\Lambda_\mathbf{s}$  (see section 2.1).

Following [9], denote by  $\widetilde{\mathcal{H}}_{R,n}$  the algebra  $\mathcal{H}_{R,n}(q^{-1}; s_{l-1}, \dots, s_0)$  and write  $\widetilde{T}_0, \dots, \widetilde{T}_{l-1}$  for the standard generators of  $\widetilde{\mathcal{H}}_{R,n}$ . If  $\boldsymbol{\lambda} \in \Pi_{l,n}$ , we write  $\widetilde{S}^\boldsymbol{\lambda}$  for the corresponding Specht module on  $\widetilde{\mathcal{H}}_{R,n}$  and  $\widetilde{D}^\boldsymbol{\lambda}$  for the corresponding irreducible one. We have an isomorphism  $\theta : \mathcal{H}_{R,n} \rightarrow \widetilde{\mathcal{H}}_{R,n}$  given by

$$T_0 \mapsto \widetilde{T}_0 \quad T_i \mapsto -q\widetilde{T}_i \quad (i = 1, \dots, n-1).$$

Put  $\mathbf{s}^\sharp = (-s_{l-1} \pmod{e}, \dots, -s_0 \pmod{e})$ . Then,  $\theta$  induces a functor  $F_l$  from the category of  $\mathcal{H}_{R,n}$ -modules to the category of  $\widetilde{\mathcal{H}}_{R,n}$  modules. As a consequence, we obtain a bijective map

$$m_l : \Phi_e^\mathbf{s} \rightarrow \Phi_e^{\mathbf{s}^\sharp},$$

satisfying

$$F_l(D^\lambda) \simeq \widetilde{D}^{m_l(\lambda)},$$

for all  $\lambda \in \Phi_e^s$ . This map can be viewed as a generalization of the Mullineux involution. In [9], Fayers describes this generalized Mullineux involution by using skew-isomorphisms of  $A_{e-1}^{(1)}$ -crystals. This description is similar to a former one obtained in [18] for the analogue of the Mullineux involution, namely the Zelevinsky involution, in affine Hecke algebras of type  $A$ . Both rest on the determination of paths in crystal graphs. The aim of this note is to give an alternative efficient description of the map  $m_l$  in the spirit of the original procedure of Mullineux, that is without using paths in crystals. We obtain an algorithm to compute the Mullineux involution for all positive integer  $l$ . This is achieved by combining the known description of the original Mullineux involution (i.e. for  $l = 1$ ) with results of [15] on isomorphisms of  $A_{e-1}^{(1)}$ -crystals.

The following sections are structured as follows. Section 2 is devoted to a brief review on  $A_{e-1}^{(1)}$ -crystals and their labellings by Uglov and Kleshchev  $l$ -partitions. In Section 3, we mainly recall and reformulate in an appropriate language results of [14] and [15]. Section 4 contains the main result of this paper, namely the description of the map  $m_l$  on Kleshchev  $l$ -partitions.

## 2. $A_{e-1}^{(1)}$ -CRYSTALS AND THE MULLINEUX INVOLUTION

Let  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$  be the quantum group of affine type  $A_{e-1}^{(1)}$ . We begin this section, by reviewing some background on the crystal graph theory of the irreducible highest weight  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ -modules. In particular, we recall the notion of Uglov and Kleshchev  $l$ -partitions and expose Fayers work on the Mullineux involution.

**2.1. Realizations of abstract  $A_{e-1}^{(1)}$ -crystals.** By slightly abuse of notation, we identify the elements of  $\mathbb{Z}/e\mathbb{Z}$  with their corresponding labels in  $\{0, \dots, e-1\}$  when there is no risk of confusion. Let  $v$  be an indeterminate and  $e > 1$  be an integer. Write  $\{\Lambda_0, \dots, \Lambda_{e-1}\}$  for the set of dominant weights of  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ . Let  $l \geq 1 \in \mathbb{N}$  and consider  $\mathfrak{s} = (s_0, \dots, s_{l-1}) \in \mathbb{Z}^l$ . We set  $\mathfrak{s} = (s_0(\bmod e), \dots, s_{l-1}(\bmod e)) \in (\mathbb{Z}/e\mathbb{Z})^l$ . One can then associate to  $\mathfrak{s}$  the abstract  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ -irreducible module  $V_e(\Lambda_{\mathfrak{s}})$  with highest weight  $\Lambda_{\mathfrak{s}} := \Lambda_{s_0(\bmod e)} + \dots + \Lambda_{s_{l-1}(\bmod e)}$ . There exist distinct realizations of  $V_e(\Lambda_{\mathfrak{s}})$  as an irreducible component of a Fock space. These Fock spaces can be defined in two different ways. The first definition yields  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ -modules  $\mathfrak{F}_e^{\mathfrak{s}}$  whose structure depends on the choice of a representant  $\mathfrak{s} \in \mathfrak{s}$ . The second one yields a  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ -module  $\mathfrak{F}_e^{\mathfrak{s}}$  whose structure only depends on  $\mathfrak{s} \in (\mathbb{Z}/e\mathbb{Z})^l$ . Each of these modules will provide us with a realization  $V_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}}) \subset \mathfrak{F}_e^{\mathfrak{s}}$  and  $V_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}}) \subset \mathfrak{F}_e^{\mathfrak{s}}$  of the abstract module  $V_e(\Lambda_{\mathfrak{s}})$ . As a  $\mathbb{C}(v)$ -vector space, the

Fock spaces  $\mathfrak{F}_e^{\mathfrak{s}}$  and  $\mathfrak{F}_e^{\mathfrak{s}}$  of level  $l$  admit the set of all  $l$ -partitions as a natural basis. Namely the underlying vector space is

$$\mathfrak{F}_e = \bigoplus_{n \geq 0} \bigoplus_{\lambda \in \Pi_{l,n}} \mathbb{C}(v)\lambda,$$

where  $\Pi_{l,n}$  is the set of  $l$ -partitions with rank  $n$ . To describe the different structures of  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ -modules on  $\mathfrak{F}_e$ , we need some combinatorics.

Let  $\lambda$  be an  $l$ -partition (identified with its Young diagram). Then, the nodes of  $\lambda$  are the triplets  $\gamma = (a, b, c)$  where  $c \in \{0, \dots, l-1\}$  and  $a, b$  are respectively the row and column indices of the node  $\gamma$  in  $\lambda^{(c)}$ . The content of  $\gamma$  is the integer  $c(\gamma) = b - a + s_c$  and the residue  $\text{res}(\gamma)$  of  $\gamma$  is the element of  $\mathbb{Z}/e\mathbb{Z}$  such that

$$(1) \quad \text{res}(\gamma) \equiv c(\gamma) \pmod{e}.$$

We will say that  $\gamma$  is an  $i$ -node of  $\lambda$  when  $\text{res}(\gamma) \equiv i \pmod{e}$ . Finally, We say that  $\gamma$  is removable when  $\gamma = (a, b, c) \in \lambda$  and  $\lambda \setminus \{\gamma\}$  is an  $l$ -partition. Similarly  $\gamma$  is said addable when  $\gamma = (a, b, c) \notin \lambda$  and  $\lambda \cup \{\gamma\}$  is an  $l$ -partition.

The  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ -module structures  $\mathfrak{F}_e^{\mathfrak{s}}$  and  $\mathfrak{F}_e^{\mathfrak{s}}$  on  $\mathfrak{F}_e$  are now derived from distinct total orders on the set of addable and removable  $i$ -nodes of the multipartitions. Consider  $\gamma_1 = (a_1, b_1, c_1)$  and  $\gamma_2 = (a_2, b_2, c_2)$  two  $i$ -nodes in  $\lambda$ . We define the orders  $\prec_{\mathfrak{s}}$  and  $\prec_{\mathfrak{s}}$  by setting

$$\gamma_1 \prec_{\mathfrak{s}} \gamma_2 \iff \begin{cases} c(\gamma_1) < c(\gamma_2) \text{ or} \\ c(\gamma_1) = c(\gamma_2) \text{ and } c_1 > c_2 \end{cases}$$

and

$$\gamma_1 \prec_{\mathfrak{s}} \gamma_2 \iff \begin{cases} c_1 > c_2 \text{ or} \\ c_1 = c_2 \text{ and } c(\gamma_1) < c(\gamma_2) \end{cases}.$$

Observe that the order  $\prec_{\mathfrak{s}}$  (res.  $\prec_{\mathfrak{s}}$ ) depends (resp. does not depend) on the choice of  $\mathfrak{s} \in \mathfrak{s}$ . Using  $\prec_{\mathfrak{s}}$  or  $\prec_{\mathfrak{s}}$ , it is possible to define two actions of  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$  on  $\mathfrak{F}_e$  yielding to the Fock spaces  $\mathfrak{F}_e^{\mathfrak{s}}$  and  $\mathfrak{F}_e^{\mathfrak{s}}$ . We do not use the explicit description of these actions in the sequel and refer the reader to [13] for a complete exposition. Nevertheless, we shall need the following:

*Remark 2.1.*

- (1) For  $l = 1$ , the Fock spaces  $\mathfrak{F}_e^{\mathfrak{s}}$  and  $\mathfrak{F}_e^{\mathfrak{s}}$  coincide.
- (2) With  $\mathfrak{s} = (s_0 \pmod{e}, \dots, s_{l-1} \pmod{e})$ , we have

$$(2) \quad \mathfrak{F}_e^{\mathfrak{s}} = \mathfrak{F}_e^{s_0} \otimes \dots \otimes \mathfrak{F}_e^{s_{l-1}}.$$

The Fock space  $\mathfrak{F}_e^{\mathfrak{s}}$  is a tensor product of Fock spaces of level 1.

As  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ -modules,  $\mathfrak{F}_e^{\mathfrak{s}}$ ,  $\mathfrak{s} \in \mathfrak{s}$  and  $\mathfrak{F}_e^{\mathfrak{s}}$  are all isomorphic but with distinct actions for  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ . For each of these actions, the empty multipartition  $\emptyset = (\emptyset, \dots, \emptyset)$  is a highest weight vector of highest weight  $\Lambda_{\mathfrak{s}}$ . We denote by  $V_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  (resp.  $V_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$ ) the irreducible component with highest weight vector  $\emptyset$  in  $\mathfrak{F}_e^{\mathfrak{s}}$  (resp. in  $\mathfrak{F}_e^{\mathfrak{s}}$ ).

The following observation will be crucial in the rest of the paper. Fix  $\mathfrak{s} \in (\mathbb{Z}/e\mathbb{Z})^l$ . The modules  $V_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  and  $V_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  when  $\mathfrak{s}$  runs over  $\mathfrak{s}$  are all isomorphic to the abstract module  $V_e(\Lambda_{\mathfrak{s}})$ . However, the corresponding actions of the Chevalley operators on these modules do not coincide in general.


The modules  $\mathfrak{F}_e^{\mathfrak{s}}$  and  $\mathfrak{F}_e^{\mathfrak{s}}$  are integrable. Hence, by the general theory of Kashiwara crystal bases,  $\mathfrak{F}_e^{\mathfrak{s}}$  (resp.  $\mathfrak{F}_e^{\mathfrak{s}}$ ) has a crystal graph  $B_e^{\mathfrak{s}}$  (resp.  $B_e^{\mathfrak{s}}$ ). Since the parametrizations of the crystals  $B_e^{\mathfrak{s}}$  (resp.  $B_e^{\mathfrak{s}}$ ) depend on the total order  $\prec_{\mathfrak{s}}$  (resp.  $\prec_{\mathfrak{s}}$ ), their underlying graph structures do not coincide in general although they all admit the set of  $l$ -partitions as set of vertices. To describe the graph structure of  $B_e^{\mathfrak{s}}$  (for a fixed  $\mathfrak{s} \in \mathfrak{s}$ ) and  $B_e^{\mathfrak{s}}$  we proceed as follows. Starting from any  $l$ -partition  $\lambda$ , we can consider its set of addable and removable  $i$ -nodes. Let  $w_i$  be the word obtained first by writing the addable and removable  $i$ -nodes of  $\lambda$  in increasing order with respect to  $\prec_{\mathfrak{s}}$  (resp.  $\prec_{\mathfrak{s}}$ ) next by encoding each addable  $i$ -node by the letter  $A$  and each removable  $i$ -node by the letter  $R$ . Write  $\tilde{w}_i = A^p R^q$  for the word derived from  $w_i$  by deleting as many of the factors  $RA$  as possible. If  $p > 0$ , let  $\gamma$  be the rightmost addable  $i$ -node in  $\tilde{w}_i$ . When  $\tilde{w}_i \neq \emptyset$ , the node  $\gamma$  is called the good  $i$ -node.

Now, the crystal graph  $B_e^{\mathfrak{s}}$  (resp.  $B_e^{\mathfrak{s}}$ ) of  $\mathfrak{F}_e^{\mathfrak{s}}$  (resp.  $\mathfrak{F}_e^{\mathfrak{s}}$ ) is the graph with

- vertices: the  $l$ -partitions,
- edges:  $\lambda \xrightarrow{i} \mu$  if and only if  $\mu$  is obtained by adding to  $\lambda$  a good  $i$ -node.

**2.2. Uglov and Kleshchev multipartitions.** Since  $V_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  and  $V_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  are the irreducible components with highest weight vector  $\emptyset$  in  $\mathfrak{F}_e^{\mathfrak{s}}$  and  $\mathfrak{F}_e^{\mathfrak{s}}$ , their crystal graphs  $B_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  and  $B_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  can be realized respectively as the connected components of highest weight vertex  $\emptyset$  in  $B_e^{\mathfrak{s}}$  and  $B_e^{\mathfrak{s}}$ .


**Example.** The graph below is the subgraph of  $B_4^{(0,0,1)}(2\Lambda_0 + \Lambda_1)$  containing the 3-partitions of rank less or equal to 4.



We denote by  $\Phi_e^{\mathfrak{s}}$  the set of vertices in  $B_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$ . The elements of  $\Phi_e^{\mathfrak{s}}$  are called the Uglov  $l$ -partitions. For completion, we also write  $\Phi_e^{\mathfrak{s}}(n)$  for the subset of  $\Phi_e^{\mathfrak{s}}$  containing the Uglov  $l$ -partitions with rank  $n$ . In general the set  $\Phi_e^{\mathfrak{s}}$  is very difficult to describe without computing the underlying crystal. Nevertheless, in the case where  $0 \leq s_0 \leq \dots \leq s_{l-1} \leq e-1$ , Foda et al. have given a simple non recursive description of the Uglov  $l$ -partitions (see [10]).

We denote by  $\Phi_e^{\mathfrak{s}}$  the set of vertices in  $B_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$ . The elements of  $\Phi_e^{\mathfrak{s}}$  are called the Kleshchev  $l$ -partitions. We define  $\Phi_e^{\mathfrak{s}}(n)$  as we have done in the Uglov case. Note that there exists only a partial combinatorial description of the Kleshchev  $l$ -partitions  $\Phi_e^{\mathfrak{s}}$ , namely when the number of fundamental weights appearing in the decomposition of  $\Lambda_{\mathfrak{s}}$  is less or equal to 2, see [2] and [4]. In the other cases, they can only be obtained by computing the underlying crystal or using the procedure described in [15, §5.4].

**Example 2.2.** The graph below is the subgraph of  $B_4^{\mathfrak{s}}(2\Lambda_0 + \Lambda_1)$  with  $\mathfrak{s} = (0 \pmod{4}, 0 \pmod{4}, 1 \pmod{4})$  containing the 3-partitions of rank less than 4.



Let  $\mathfrak{s}' := (s'_0, \dots, s'_{l-1}) \in \mathbb{Z}^l$  be such that  $\mathfrak{s}' \in \mathfrak{s}$ . Then the two modules  $V_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  and  $V_e^{\mathfrak{s}'}(\Lambda_{\mathfrak{s}'})$  are both isomorphic to  $V_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$ . Hence the corresponding three crystal graphs are isomorphic. These crystal isomorphisms define the following natural bijections

$$\Psi_{e,n}^{\mathfrak{s} \rightarrow \mathfrak{s}'} : \Phi_e^{\mathfrak{s}}(n) \rightarrow \Phi_e^{\mathfrak{s}'}(n)$$

$$\Psi_{e,n}^{\mathfrak{s} \rightarrow \mathfrak{s}} : \Phi_e^{\mathfrak{s}}(n) \rightarrow \Phi_e^{\mathfrak{s}}(n).$$

As it can be easily checked on the previous examples, the sets  $\Phi_e^{\mathfrak{s}}(n)$  and  $\Phi_e^{\mathfrak{s}'}(n)$  do not coincide in general. However, if we fix  $n$ , it is possible to realize the set  $\Phi_e^{\mathfrak{s}}(n)$  of Kleshchev  $l$ -partitions with rank  $n$  as a special set  $\Phi_e^{\mathfrak{s}_{\infty}}(n)$ ,  $\mathfrak{s}_{\infty} \in \mathfrak{s}$  of Uglov  $l$ -partitions. We refer the reader to Lemma 5.4.1 in [15] for the proof of the following proposition.

**Proposition 2.3.** *Let  $n \in \mathbb{N}$  and consider  $\mathbf{s}_\infty := (s_0, \dots, s_{l-1}) \in \mathbb{Z}^l$  such that*

$$(3) \quad s_{i+1} - s_i > n - 1$$

for all  $i = 0, \dots, l - 2$ . Then  $\Psi_{e,n}^{\mathbf{s}_\infty \rightarrow \mathbf{s}}$  is the identity. Hence, we have :

$$\Phi_e^{\mathbf{s}}(n) = \Phi_e^{\mathbf{s}_\infty}(n).$$

In particular,  $\Phi_e^{\mathbf{s}_\infty}(n)$  does not depend on the multicharge  $\mathbf{s}_\infty$  provided (3) is satisfied.

This shows that the description of the bijections  $\Psi_{e,n}^{\mathbf{s} \rightarrow \mathbf{s}'}$  for all  $\mathbf{s}, \mathbf{s}' \in \mathfrak{s}$  will lead that of  $\Psi_{e,n}^{\mathbf{s} \rightarrow \mathbf{s}}$  as a special case. A multicharge verifying (3) is called asymptotic (this is called  $n$ -dominant in [21]). This justifies the notation  $\Phi_e^{\mathbf{s}_\infty}(n)$  we have adopted.

### 3. ISOMORPHISMS OF $A_{e-1}^{(1)}$ -CRYSTALS

We now review some results obtained in [14] and [15] making explicit the bijections  $\Psi_{e,n}^{\mathbf{s} \rightarrow \mathbf{s}'}$ . In [15], we have chosen to explain our results in the language of column shaped tableaux to make apparent the link with the combinatorics of  $\mathfrak{sl}_n$ . Here, we rather adopt the equivalent language of symbols also used in [14] which permits to give a very simple description of the elementary steps involved in our algorithms.

**3.1. Elementary transformations in  $\widehat{S}_l$ .** We write as usual  $\widehat{S}_l$  for the extended affine symmetric group in type  $A_{l-1}$ . This group is generated by the elements  $\sigma_1, \dots, \sigma_{l-1}$  and  $y_0, \dots, y_{l-1}$  together with the relations

$$\sigma_c \sigma_{c+1} \sigma_c = \sigma_{c+1} \sigma_c \sigma_{c+1}, \quad \sigma_c \sigma_d = \sigma_d \sigma_c \text{ for } |c - d| > 1, \quad \sigma_c^2 = 1,$$

with all indices in  $\{1, \dots, l - 1\}$  and

$$y_c y_d = y_d y_c, \quad \sigma_c y_d = y_d \sigma_c \text{ for } d \neq c, c + 1, \quad \sigma_c y_c \sigma_c = y_{c+1}.$$

For any  $c \in \{0, \dots, l - 1\}$ , we set  $z_c = y_1 \cdots y_c$ . Write also  $\xi = \sigma_{l-1} \cdots \sigma_1$  and  $\tau = y_l \xi$ . Since  $y_c = z_{c-1}^{-1} z_c$ ,  $\widehat{S}_l$  is generated by the transpositions  $\sigma_c$  with  $c \in \{1, \dots, l - 1\}$  and the elements  $z_c$  with  $c \in \{1, \dots, l\}$ . Observe that for any  $c \in \{1, \dots, l - 1\}$ , we have

$$(4) \quad z_c = \xi^{l-c} \tau^c.$$

This implies that  $\widehat{S}_l$  is generated by the transpositions  $\sigma_c$  with  $c \in \{1, \dots, l - 1\}$  and  $\tau$ . Consider  $e$  a fixed positive integer. We obtain a faithful action of  $\widehat{S}_l$  on  $\mathbb{Z}^l$  by setting for any  $\mathbf{s} = (s_0, \dots, s_{l-1}) \in \mathbb{Z}^l$

$$\sigma_c(\mathbf{s}) = (s_0, \dots, s_c, s_{c-1}, \dots, s_{l-1}) \text{ and } y_c(\mathbf{s}) = (s_0, \dots, s_{c-1}, s_c + e, \dots, s_{l-1}).$$

Then  $\tau(\mathbf{s}) = (s_1, s_2, \dots, s_{l-1}, s_0 + e)$ .



*Remark 3.1.* From the preceding considerations,  $\{s_1, \dots, s_{l-1}, \tau\}$  is a minimal set of generators for  $\widehat{S}_l$ . Hence, the bijections  $\Psi_{e,n}^{\mathbf{s} \rightarrow \mathbf{s}'} : \Phi_e^{\mathbf{s}}(n) \rightarrow \Phi_e^{\mathbf{s}'}(n)$  with  $\mathbf{s} \in \mathfrak{s}$  and  $\Psi_{e,n}^{\mathbf{s} \rightarrow \mathbf{s}} : \Phi_e^{\mathbf{s}}(n) \rightarrow \Phi_e^{\mathbf{s}}(n)$  can be obtained by composing bijections corresponding to the cases  $\mathbf{s}' = \tau(\mathbf{s})$  and  $\mathbf{s}' = \sigma_c(\mathbf{s})$  with  $c = 1, \dots, l-1$ .

**3.2. Elementary crystals isomorphisms.** The following proposition is stated in [15, Prop. 5.2.1]. We give the proof for the convenience of the reader.

**Proposition 3.2.** *Let  $\mathbf{s} \in \mathbb{Z}^l$  and  $e \in \mathbb{N}$ . Let  $\boldsymbol{\lambda} = (\lambda^{(0)}, \dots, \lambda^{(l-1)}) \in \Phi_e^{\mathbf{s}}(n)$ . Then*

$$\Psi_{e,n}^{\mathbf{s} \rightarrow \tau(\mathbf{s})} = (\lambda^{(1)}, \dots, \lambda^{(l-1)}, \lambda^{(0)}).$$

*Proof.* Set  $\mathbf{s} = (s_0, \dots, s_{\ell-1})$ . Then  $\tau(\mathbf{s}) = (s_1, \dots, s_{\ell-1}, s_0 + e)$ . Consider  $\boldsymbol{\lambda} = (\lambda^{(0)}, \dots, \lambda^{(\ell-1)})$  a multipartition and set  $\boldsymbol{\lambda}^\diamond = (\lambda^{(1)}, \dots, \lambda^{(\ell-1)}, \lambda^{(0)})$ . Consider  $i \in \{0, 1, \dots, e-1\}$  and  $\gamma_1 = (a_1, b_1, c_1), \gamma_2 = (a_2, b_2, c_2)$  two  $i$ -nodes of  $\boldsymbol{\lambda}$ . Then  $\gamma_1^\diamond = (a_1, b_1, c_1 - 1 \pmod{e})$  and  $\gamma_2^\diamond = (a_2, b_2, c_2 - 1 \pmod{e})$  are two  $i$ -nodes of  $\boldsymbol{\lambda}^\diamond$ . We then easily check that  $\gamma_2 \prec_{\mathbf{s}} \gamma_1$  if and only if  $\gamma_2^\diamond \prec_{\tau(\mathbf{s})} \gamma_1^\diamond$ . This implies that  $\Psi_{e,n}^{\mathbf{s} \rightarrow \tau(\mathbf{s})}(\boldsymbol{\lambda}) = \boldsymbol{\lambda}^\diamond$ .  $\square$

The description of the bijections  $\Psi_{e,n}^{\mathbf{s} \rightarrow \sigma_c(\mathbf{s})}$ ,  $c = 1, \dots, l-1$  are more complicated. It essentially rests on the following basic procedure on pairs  $(U, D)$  of strictly increasing sequences of integers. let  $U = [x_1, \dots, x_r]$  and  $D = [y_1, \dots, y_s]$  two strictly increasing sequences of integers. We compute from  $(U, D)$  a new pair  $(U', D')$  with  $U' = [x'_1, \dots, x'_r]$  and  $D' = [y'_1, \dots, y'_s]$  of such sequences by applying the following algorithm :

**Algorithm.**

- Assume  $r \geq s$ . We associate to  $y_1$  the integer  $x_{i_1} \in U$  such that

$$(5) \quad x_{i_1} = \begin{cases} \max\{x \in U \mid y_1 \geq x\} & \text{if } y_1 \geq x_1 \\ x_r & \text{otherwise} \end{cases} .$$

We repeat the same procedure to the ordered pair  $(U \setminus \{x_{i_1}\}, D \setminus \{y_1\})$ . By induction this yields a subset  $\{x_{i_1}, \dots, x_{i_s}\} \subset U$ . Then we define  $D'$  as the increasing reordering  $\{x_{i_1}, \dots, x_{i_s}\}$  and  $U'$  as the increasing reordering of  $U \setminus \{x_{i_1}, \dots, x_{i_s}\} \sqcup D$ .

- Assume  $r < s$ . We associate to  $x_1$  the integer  $y_{i_1} \in D$  such that

$$(6) \quad y_{i_1} = \begin{cases} \min\{y \in D \mid x_1 \leq y\} & \text{if } x_1 \leq y_s \\ y_1 & \text{otherwise} \end{cases} .$$

We repeat the same procedure to the ordered sequences  $U \setminus \{x_1\}$  and  $D \setminus \{y_{i_1}\}$  and obtain a subset  $\{y_{i_1}, \dots, y_{i_r}\} \subset D$ . Then we define  $U'$  as the increasing reordering  $\{y_{i_1}, \dots, y_{i_r}\}$  and  $D'$  as the increasing reordering of  $D \setminus \{y_{i_1}, \dots, y_{i_r}\} \sqcup U$ .

**Example 3.3.** Take  $U = [0, 1, 3]$  and  $D = [0, 1, 2, 4, 7]$ . The subset of  $U$  determined by the previous algorithm is  $\{0, 1, 4\}$ . This gives  $U' = [0, 1, 4]$  and  $D' = [0, 1, 2, 3, 7]$ .

Now consider  $c \in \{1, \dots, e-1\}$  and  $\boldsymbol{\lambda} = (\lambda^{(0)}, \dots, \lambda^{(l-1)}) \in \Phi_e^{\mathbf{s}}(n)$ . In order to compute  $\Psi_{e,n}^{\mathbf{s} \rightarrow \sigma_c(\mathbf{s})}(\boldsymbol{\lambda})$ , we need the symbol  $S_c$  of the bipartition  $(\lambda^{(c-1)}, \lambda^{(c)})$ . Let  $d_{c-1}$  and  $d_c$  be the number of nonzero parts in  $\lambda^{(c-1)}$  and  $\lambda^{(c)}$  and put  $m = \max(d_{c-1} - s_{c-1}, d_c - s_c) + 1$ . Set

$$\begin{cases} \beta_i^{(c-1)} = \lambda_i^{(c-1)} - i + s_{c-1} + m \text{ for any } i = 1, \dots, m + s_{c-1} \\ \beta_i^{(c)} = \lambda_i^{(c)} - i + s_c + m \text{ for any } i = 1, \dots, m + s_c \end{cases}$$

where, by convention the partitions  $\lambda^{(c-1)}$  and  $\lambda^{(c)}$  are taken with an infinite number of zero parts. The symbol  $S_c$  is the two-row tableau

$$S_c = \begin{pmatrix} \beta_{m+s_c}^{(c)} & \beta_{m+s_{c-1}}^{(c)} & \cdots & \cdots & \cdots & \beta_1^{(c)} \\ \beta_{m+s_{c-1}}^{(c-1)} & \beta_{m+s_{c-1}-1}^{(c-1)} & \cdots & \cdots & \beta_1^{(c-1)} & \end{pmatrix} = \begin{pmatrix} U \\ D \end{pmatrix}$$

Observe that  $\beta_i^{(k)}, k \in \{c-1, c\}$  is nothing but the content of the node  $\gamma = (i, \lambda_i^{(k)}, k) \in \boldsymbol{\lambda}$  translated by  $m$ . This translation normalizes the symbol  $S_c$  so that  $\beta_{m+s_c}^{(c)} = \beta_{m+s_{c-1}}^{(c-1)} = 0$ . In fact, for a fixed pair  $(s_{c-1}, s_c)$  and  $m$  as above, the map  $\Sigma_{(s_{c-1}, s_c)}$  which associates to each bipartition its symbol is bijective. It is easy, from a symbol  $S_c$ , to recover the unique bipartition  $(\lambda^{(c-1)}, \lambda^{(c)})$  such that  $S_c$  is the symbol of  $(\lambda^{(c-1)}, \lambda^{(c)})$ . We are now ready to describe the bijection  $\Psi_{e,n}^{\mathbf{s} \rightarrow \sigma_c(\mathbf{s})} : \Phi_e^{\mathbf{s}}(n) \rightarrow \Phi_e^{\sigma_c(\mathbf{s})}(n)$ .

Let  $\mathbf{s} \in \mathbb{Z}^l$  and  $e \in \mathbb{N}$ . Consider  $\boldsymbol{\lambda} = (\lambda^{(0)}, \dots, \lambda^{(l-1)}) \in \Phi_e^{\mathbf{s}}(n)$  and  $c \in \{1, \dots, l-1\}$ . Write  $S_c = \begin{pmatrix} U \\ D \end{pmatrix}$  for the symbol corresponding to the bipartition  $(\lambda^{(c-1)}, \lambda^{(c)})$ . Denote by  $\tilde{S}_c = \begin{pmatrix} \tilde{U} \\ \tilde{D} \end{pmatrix}$  the symbol obtained by applying to  $S_c$  the previous algorithm. Then we compute the bipartition  $(\tilde{\lambda}^{(c-1)}, \tilde{\lambda}^{(c)}) = \Sigma_{(s_{c-1}, s_c)}^{-1}(\tilde{S}_c)$ . Finally we consider the  $l$ -partition

$$\tilde{\boldsymbol{\lambda}} = (\lambda^{(0)}, \dots, \tilde{\lambda}^{(c)}, \tilde{\lambda}^{(c-1)}, \dots, \lambda^{(l-1)})$$

obtained by replacing in  $\boldsymbol{\lambda}$ ,  $\lambda^{(c-1)}$  by  $\tilde{\lambda}^{(c)}$  and  $\lambda^{(c)}$  by  $\tilde{\lambda}^{(c-1)}$ .

**Proposition 3.4.** *With the above notation, we have*

$$\Psi_{e,n}^{\mathbf{s} \rightarrow \sigma_{c-1}(\mathbf{s})}(\boldsymbol{\lambda}) = \tilde{\boldsymbol{\lambda}}.$$

*Proof.* The above algorithm is the traduction in terms of symbols of Proposition 5.2.2 in [15].  $\square$

**Example 3.5.** Let  $e = 3$  and let  $\mathbf{s} = (0, 2, 0)$ . Then one can check that the 3-partition  $(4.1, 3.1, 1)$  belongs to  $\Phi_e^{\mathbf{s}}(10)$ . We want to determine the 3-partition  $\Psi_{e,n}^{\mathbf{s} \rightarrow \sigma_2(\mathbf{s})}(4.1, 3.1, 1)$ . We first write the symbol  $S_2$  of  $(3.1, 1)$  with respect to  $\mathbf{s}$  and  $m = 3$  :

$$S_2 = \begin{pmatrix} 0 & 1 & 3 \\ 0 & 1 & 2 & 4 & 7 \end{pmatrix}.$$

By using the previous example, we obtain

$$\tilde{S}_2 = \begin{pmatrix} 0 & 1 & 4 & & \\ 0 & 1 & 2 & 3 & 7 \end{pmatrix}.$$

This thus gives  $\tilde{\lambda}^{(c)} = (2)$  and  $\tilde{\lambda}^{(c-1)} = (3)$ . Hence  $\Psi_{e,n}^{\mathfrak{s} \rightarrow \sigma_2(\mathfrak{s})}(4.1, 3.1.1, \emptyset) = (4.1, 2, 3)$ .

#### 4. MULLINEUX INVOLUTION FOR KLESHCHEV $l$ -PARTITIONS

**4.1. Mullineux involution as a skew crystal isomorphism.** We here keep the setting of the introduction and review a result of Fayers [9] which shows how the Mullineux involution on Kleshchev  $l$ -partitions can be computed by using paths in crystal graphs. The generalized Mullineux involution can be regarded as a bijection

$$m_l : \Phi_e^{\mathfrak{s}} \rightarrow \Phi_e^{\mathfrak{s}^\sharp}$$

where  $\mathfrak{s} = (s_0(\bmod e), \dots, s_{l-1}(\bmod e))$  and  $\mathfrak{s}^\sharp = (-s_{l-1}(\bmod e), \dots, -s_0(\bmod e))$ .

**Theorem 4.1** (Fayers [9]). *Let  $\lambda \in \Phi_e^{\mathfrak{s}}$  and consider in  $B_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  a path*

$$\emptyset \xrightarrow{i_1} \dots \xrightarrow{i_2} \dots \xrightarrow{i_3} \dots \xrightarrow{i_n} \lambda$$

*from the empty  $l$ -partition to  $\lambda$ . There exists a corresponding path*

$$\emptyset \xrightarrow{-i_1} \dots \xrightarrow{-i_2} \dots \xrightarrow{-i_3} \dots \xrightarrow{-i_n} \mu$$

*in  $B_e^{\mathfrak{s}^\sharp}(\Lambda_{\mathfrak{s}^\sharp})$  from the empty  $l$ -partition to an  $l$ -partition  $\mu \in \Phi_e^{\mathfrak{s}^\sharp}$ . We have then*

$$m_l(\lambda) = \mu.$$

In the level 1 case (i.e. for  $l = 1$ ), the map  $m_1$  can be described independently of the notion of crystal graph by using an algorithm originally due to Mullineux. We refer for example to [12] for a complete exposition of this procedure. For all  $s \in \mathbb{Z}$ , it gives the map

$$m_1 : \Phi_e^{\mathfrak{s}} \rightarrow \Phi_e^{-\mathfrak{s}}.$$

where  $\mathfrak{s} = s(\bmod e)$ . Our aim is now to show how the generalized Mullineux involution (i.e. in level  $l > 1$ ) can be computed from the map  $m_1$  and the results of Section 3. For any  $\mathfrak{s} = (s_0(\bmod e), \dots, s_{l-1}(\bmod e)) \in (\mathbb{Z}/e\mathbb{Z})^l$ , set

$$\begin{cases} -\mathfrak{s} = (-s_0(\bmod e), \dots, -s_{l-1}(\bmod e)), \\ \mathfrak{s}^\sharp = (-s_{l-1}(\bmod e), \dots, -s_0(\bmod e)). \end{cases}$$

Our strategy is as follows. Let  $\lambda$  be in  $\Phi_e^{\mathfrak{s}}$  and consider a path from the empty  $l$ -partition to  $\lambda$  in  $B_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  with arrows successively labelled by  $i_1, \dots, i_n$ .

- (1) We first describe the  $l$ -partition  $\nu$  in  $\Phi_e^{-\mathfrak{s}}$  defined by considering the path

$$\emptyset \xrightarrow{-i_1} \dots \xrightarrow{-i_2} \dots \xrightarrow{-i_3} \dots \xrightarrow{-i_n} \nu$$

in  $B_e^{-\mathfrak{s}}(\Lambda_{-\mathfrak{s}})$ .

- (2) Next, we describe the bijection  $\Phi_e^{-\mathfrak{s}} \rightarrow \Phi_e^{\mathfrak{s}\sharp}$  defined by the crystal isomorphism between  $B_e^{-\mathfrak{s}}(\Lambda_{-\mathfrak{s}})$  and  $B_e^{\mathfrak{s}\sharp}(\Lambda_{-\mathfrak{s}})$ . This permits to compute  $\boldsymbol{\mu}$  from  $\boldsymbol{\nu}$ .

#### 4.2. Computing $\boldsymbol{\nu}$ from $\boldsymbol{\lambda}$ .

**Proposition 4.2.** *Let  $\boldsymbol{\lambda} \in \Phi_e^{\mathfrak{s}}$  and consider in  $B_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  a path*

$$\emptyset \xrightarrow{i_1} \cdot \xrightarrow{i_2} \cdot \xrightarrow{i_3} \dots \xrightarrow{i_n} \boldsymbol{\lambda}$$

*from the empty  $l$ -partition to  $\boldsymbol{\lambda}$ . Then there exists a corresponding path*

$$\emptyset \xrightarrow{-i_1} \cdot \xrightarrow{-i_2} \cdot \xrightarrow{-i_3} \dots \xrightarrow{-i_n} \boldsymbol{\nu}$$

*in  $B_e^{-\mathfrak{s}}(\Lambda_{-\mathfrak{s}})$  from the empty  $l$ -partition to an  $l$ -partition  $\boldsymbol{\nu} \in \Phi_e^{-\mathfrak{s}}$ . We have then*

$$\boldsymbol{\nu} = (m_1(\lambda^0), \dots, m_1(\lambda^{l-1})).$$

*Proof.* We know by (2) that the Fock spaces  $\mathfrak{F}_e^{\mathfrak{s}}$  and  $\mathfrak{F}_e^{-\mathfrak{s}}$  can be regarded as tensor products of level 1 Fock spaces. Therefore, each  $l$ -partition  $\mathbf{b} = (b^{(0)}, \dots, b^{(l-1)})$  in the crystals  $B_e^{\mathfrak{s}}$  and  $B_e^{-\mathfrak{s}}$  of the Fock spaces  $\mathfrak{F}_e^{\mathfrak{s}}$  and  $\mathfrak{F}_e^{-\mathfrak{s}}$  can be written  $\mathbf{b} = b^{(0)} \otimes \dots \otimes b^{(l-1)}$ . Recall that in Kashiwara crystal basis theory, the action of each crystal operator  $\tilde{e}_i, \tilde{f}_i, i \in \{0, \dots, l-1\}$  on  $\mathbf{b} = b_0 \otimes \dots \otimes b_{l-1}$  is determined by the intergers  $\varepsilon_i(b_k) = \max\{r \in \mathbb{N} \mid \tilde{e}_i^r(b_k) \neq 0\}$  and  $\varphi_i(b_k) = \max\{r \mid \tilde{f}_i^r(b_k) \neq 0\}$  when  $k$  runs over  $\{0, \dots, l-1\}$ .

We now proceed by induction on  $n$ . For  $n = 0$ , the proposition is clear. Assume that the result is proved for  $n - 1$  and set  $\tilde{\boldsymbol{\lambda}} = \tilde{f}_{i_n} \dots \tilde{f}_{i_1}(\emptyset)$  in  $B_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$ . Consider  $\boldsymbol{\lambda}_b = \tilde{f}_{i_{n-1}} \dots \tilde{f}_{i_1}(\emptyset) \in B_e^{\mathfrak{s}}(\Lambda_{\mathfrak{s}})$  and  $\boldsymbol{\nu}_b = \tilde{f}_{-i_{n-1}} \dots \tilde{f}_{-i_1}(\emptyset) \in B_e^{-\mathfrak{s}}(\Lambda_{-\mathfrak{s}})$ . By induction, we have then  $\boldsymbol{\nu}_b = (m_1(\lambda_b^{(0)}), \dots, m_1(\lambda_b^{(l-1)}))$ . The Mullineux involution  $m_1$  switches the signs of the arrows. Thus  $\varepsilon_i(m_1(\lambda_b^{(k)})) = \varepsilon_{-i}(\lambda_b^{(k)})$  and  $\varphi_i(m_1(\lambda_b^{(k)})) = \varphi_{-i}(\lambda_b^{(k)})$  for any  $k = 0, \dots, l-1$ . Since  $\tilde{f}_{i_n}(\boldsymbol{\lambda}_b) = \boldsymbol{\lambda} \neq 0$ , we have  $\tilde{f}_{i_n}(\boldsymbol{\nu}_b) = \boldsymbol{\nu} \neq 0$  by the above arguments. Moreover  $\tilde{f}_{i_n}$  and  $\tilde{f}_{-i_n}$  act on the same component of  $\boldsymbol{\lambda}_b$  and  $\boldsymbol{\nu}_b$  considered as tensor products. Namely, there exists an integer  $a \in \{0, \dots, l-1\}$  such that

$$\begin{cases} \boldsymbol{\lambda} = (\lambda_b^{(0)}, \dots, \tilde{f}_{i_n}(\lambda_b^{(a)}), \dots, \lambda_b^{(l-1)}) \\ \boldsymbol{\nu} = (m_1(\lambda_b^{(0)}), \dots, \tilde{f}_{-i_n}(m_1(\lambda_b^{(a)})), \dots, m_1(\lambda_b^{(l-1)})) \end{cases}.$$

We have  $m_1(\tilde{f}_{i_n}(\lambda_b^{(a)})) = \tilde{f}_{-i_n}(m_1(\lambda_b^{(a)}))$  since  $m_1$  is a skew-isomorphism. This gives  $\boldsymbol{\nu} = (m_1(\lambda^0), \dots, m_1(\lambda^{l-1}))$  which establishes the proposition by induction.  $\square$

**4.3. Computing  $\boldsymbol{\mu}$  from  $\boldsymbol{\nu}$ .** Now, assume we have computed the Kleshchev  $l$ -partition  $\boldsymbol{\nu} \in \Phi_e^{-\mathfrak{s}}$  from the Kleshchev  $l$ -partition  $\boldsymbol{\lambda} \in \Phi_e^{\mathfrak{s}}$  as in the previous section. To compute  $\boldsymbol{\mu} \in \Phi_e^{\mathfrak{s}\sharp}$  from the required crystal isomorphism, we need to realize  $\boldsymbol{\nu}$  as an Uglov  $l$ -partition. To do this, we consider an asymptotic multicharge  $-\mathfrak{s} \in -\mathfrak{s}$ . The multicharge  $\mathfrak{s}$  can be written  $\mathfrak{s} = (s_0 \pmod$

$e), \dots, s_{l-1} \pmod{e})$  with  $s_c$  in  $\{0, \dots, e-1\}$  for any  $c = 0, \dots, l-1$ . Now for any  $c = 1, \dots, l-1$ , write  $p_c$  for the minimal nonnegative integer such that

$$s_{c-1} - s_c + p_c e > n - 1.$$

Put then :

$$(7) \quad -\mathbf{s} := (-s_0, -s_1 + p_1 e, -s_2 + p_1 e + p_2 e, \dots, -s_{l-1} + \sum_{i=1}^{l-1} p_i e).$$

Since  $-\mathbf{s}$  is asymptotic, we have  $\boldsymbol{\nu} \in \Phi_e^{-\mathbf{s}}$  by Prop 2.3 and

$$\Psi_{e,n}^{-\mathbf{s} \rightarrow -\mathbf{s}}(\boldsymbol{\nu}) = \boldsymbol{\nu}.$$

Now, with the notation of Section 2.1 for the generators of the affine Weyl group of type  $A$ , we set

$$w_{(c,d)} = \sigma_c \sigma_{c-1} \dots \sigma_d \text{ for any } c \geq d.$$

Given  $p \in \{1, \dots, l-1\}$ , we also introduce

$$\gamma_p = w_{(l-p,1)} \dots w_{(l-2,p-1)} w_{(l-1,p)}.$$

Note that for  $\mathbf{v} := (v_0, \dots, v_{l-1}) \in \mathbb{Z}^l$  and  $p = 1, \dots, l-1$ , we have:

$$\gamma_p \cdot \mathbf{v} = (v_p, \dots, v_{l-1}, v_0, \dots, v_{p-1}).$$

For each  $p = 1, \dots, l-1$ , set

$$\alpha_p = \tau^{l-p} \gamma_p.$$

We have thus, for  $\mathbf{v} = (v_0, \dots, v_{l-1}) \in \mathbb{Z}^l$

$$\alpha_p \cdot \mathbf{v} = (v_0, \dots, v_{p-1}, v_p + e, \dots, v_{l-1} + e).$$

Finally, let  $w_0 = w_{(1,1)} w_{(2,1)} \dots w_{(l-1,1)}$  be the longest element of the symmetric group  $S_l$  and put

$$(8) \quad \eta := \alpha_1^{2(p_{l-1}+1)} \dots \alpha_{l-1}^{2(p_1+1)} w_0.$$

**Proposition 4.3.** *Consider the multicharge  $-\mathbf{s}$  of (7) and set*

$$(9) \quad \mathbf{s}^\sharp = (s_0^\sharp, \dots, s_{l-1}^\sharp) := \eta \cdot (-\mathbf{s})$$

*Then for any  $i = 0, \dots, l-1$ , we have*

$$s_i^\sharp \equiv -s_{l-i-1} \pmod{e}.$$

*Therefore  $\mathbf{s}^\sharp$  belongs to  $\mathfrak{s}^\sharp$ . Moreover, the multicharge  $\mathbf{s}^\sharp$  is asymptotic.*

*Proof.* By definition of  $\mathbf{s}^\sharp = (s_0^\sharp, \dots, s_{l-1}^\sharp)$ , we have for all  $d = 0, \dots, l-2$  :

$$(10) \quad s_{d+1}^\sharp - s_d^\sharp = -s_{l-d-2} + s_{l-d-1} + p_{l-d-1} e + 2e.$$

Now, since the integers  $s_c$ ,  $c = 0, \dots, l-1$  belong to  $\{0, \dots, l-1\}$ , we can write

$$-s_{l-d-2} + s_{l-d-1} + 2e \geq s_{l-d-2} - s_{l-d-1}.$$

Hence, we derive from (10) and the definition of the integers  $p_d$  that

$$s_{d+1}^\# - s_d^\# \geq s_{l-d-2} - s_{l-d-1} + p_{l-d-1}e > n - 1.$$

Thus, the multicharge  $\eta(\mathbf{s})$  is asymptotic which proves our proposition.  $\square$

Recall that we have a Kleshchev  $l$ -partition  $\nu \in \Phi_e^{-s}$ . Using Section 3, we can compute

$$\widehat{\nu} := \Psi_{e,n}^{-s \rightarrow s^\#}(\nu)$$

where  $s^\#$  is defined by (9). Since  $s^\#$  is asymptotic and belongs to  $s^\#$ , we derive from Proposition 2.3

$$\Psi_{e,n}^{s^\# \rightarrow s^\#}(\widehat{\nu}) = \widehat{\nu}.$$

Let us summarize the transformations we have computed from  $\lambda \in \Phi_e^s$ .

$$(11) \quad \lambda \in \Phi_e^s \xrightarrow{\Psi_{e,n}^{s \rightarrow -s}} \nu \in \Phi_e^{-s} \xrightarrow{\Psi_{e,n}^{-s \rightarrow -s}} \nu \in \Phi_e^{-s} \xrightarrow{\Psi_{e,n}^{-s \rightarrow s^\#}} \widehat{\nu} \in \Phi_e^{s^\#} \xrightarrow{\Psi_{e,n}^{s^\# \rightarrow s^\#}} \widehat{\nu} \in \Phi_e^{s^\#}$$

Recall now that we have a path

$$\emptyset \xrightarrow{i_1} \cdot \xrightarrow{i_2} \cdot \xrightarrow{i_3} \dots \xrightarrow{i_n} \lambda$$

in  $B_e^s$ . By (11), we have the path

$$\emptyset \xrightarrow{-i_1} \cdot \xrightarrow{-i_2} \cdot \xrightarrow{-i_3} \dots \xrightarrow{-i_n} \widehat{\nu}$$

in  $B_e^{s^\#}$ . This gives

$$m_l(\lambda) = \mu = \widehat{\nu}$$

Hence, the procedure (17) gives a purely combinatorial algorithm for computing the Mullineux involution which does not use the crystal (and thus the notion of good nodes) of irreducible highest weight affine modules.

**4.4. The case  $e = \infty$ .** We now briefly focus on the case  $e = \infty$ . Our algorithm considerably simplifies because in this case, the Kleshchev and FLOTW  $l$ -partitions coincide. Moreover, the Mullineux involution  $m_1$  reduces to the ordinary conjugation of partitions. Consider  $\lambda \in \Phi_\infty^s$ . The computation of  $m_l(\lambda)$  can be described as follows :

- First, by Proposition 4.2, the skew-isomorphism of crystals from  $B_e^s(\Lambda_s)$  to  $B_e^{-s}(\Lambda_{-s})$  which switches the sign of the arrows sends  $\lambda$  to

$$\mu = ((\lambda^0)', \dots, (\lambda^{l-1})')$$

where  $m_1$  just corresponds to the usual conjugation of partitions.

- Next,  $w_0$  being the longest element of the symmetric group, we have

$$w_0 \cdot (-s_0, \dots, -s_{l-1}) = (-s_{l-1}, \dots, -s_0).$$

Thus we derive

$$m_l(\lambda) = \nu := \Psi_n^{s \rightarrow w_0 \cdot s}(\mu).$$

**4.5. Conclusive remarks.** The algorithm of Section 3.2 also permits to compute the bijections  $\Psi_{e,n}^{\mathfrak{s} \rightarrow \sigma(\mathfrak{s})}$  between  $\Phi_e^{\mathfrak{s}}$  and  $\Phi_e^{\sigma(\mathfrak{s})}$  for any  $\sigma \in S_n$ . By decomposing  $\sigma$  in terms of the generators  $s_c, c = 1, \dots, l-1$ , it suffices to compute the isomorphisms  $\Psi_{e,n}^{\mathfrak{s} \rightarrow s_c(\mathfrak{s})}$ . Since  $B_e^{\mathfrak{s}}$  can be regarded as a tensor product of crystals, we can assume  $l = 2$  and describe the bijection

$$\Psi_{e,n}^{(\mathfrak{s}_0, \mathfrak{s}_1) \rightarrow (\mathfrak{s}_1, \mathfrak{s}_0)} : \Phi_e^{(\mathfrak{s}_0, \mathfrak{s}_1)} \rightarrow \Phi_e^{(\mathfrak{s}_1, \mathfrak{s}_0)}.$$

Consider  $\lambda \in \Phi_e^{(\mathfrak{s}_0, \mathfrak{s}_1)}$  and choose  $(a_0, a_1)$  an asymptotic charge with  $a_0 \in \mathfrak{s}_0$  and  $a_1 \in \mathfrak{s}_1$ . We have  $\lambda \in \Phi_e^{(a_0, a_1)}$ . Then one computes  $\Psi_{e,n}^{(a_0, a_1) \rightarrow (a_1, a_0)}(\lambda) = \nu$  as in Proposition 3.4. Now the problem is that  $(a_1, a_0)$  is not asymptotic in general. To make it asymptotic, we need to compose translations by  $e$  in  $\mathbb{Z}^2$  acting on the right coordinate. Observe that for any  $(v_0, v_1) \in \mathbb{Z}^2$ , we have  $\tau s_1(v_0, v_1) = (v_0, v_1 + e)$ . So if we set  $\kappa = \tau s_1$ , we have to compute the bipartitions

$$(12) \quad \mu^{(k)} = \Psi_{e,n}^{(a_1, a_0) \rightarrow \tau^k(a_1, a_0)}(\nu)$$

with  $k \in \{0, \dots, m\}$  where  $m \in \mathbb{N}$  is minimal such that  $a_1 - a_0 + me > n - 1$ . Nevertheless, in practice one can restrict our computations and take  $m$  minimal such  $\mu^{(m)} = \mu^{(m+1)}$ . This is because the Kleshchev bipartition is fixed by the bijection  $\Psi_{e,n}^{(\mathfrak{s}_1, \mathfrak{s}_0) \rightarrow \tau(\mathfrak{s}_1, \mathfrak{s}_0)}$ . Finally, one has

$$\Psi_{e,n}^{(\mathfrak{s}_0, \mathfrak{s}_1) \rightarrow (\mathfrak{s}_1, \mathfrak{s}_0)}(\lambda) = \mu^{(m)}.$$

Such a stabilization phenomenon also happens during the computation of  $\mu$  from  $\nu$  (see Section 4.3). This means that, in practice, we have

$$\Psi_{e,n}^{-\mathfrak{s} \rightarrow \eta(-\mathfrak{s})}(\nu) = \Psi_{e,n}^{-\mathfrak{s} \rightarrow \eta^b(-\mathfrak{s})}(\nu)$$

where  $\eta := \alpha_1^{2(p_{l-1}+1)} \dots \alpha_{l-1}^{2(p_1+1)} w_0$  is defined as in (8) and where  $\eta^b := \alpha_1^{2(p_{l-1}^b+1)} \dots \alpha_{l-1}^{2(p_1^b+1)} w_0$  with  $p_c^b \leq p_c$  for  $c \in \{0, \dots, l-1\}$ .

The previous description of the bijections  $\Psi_{e,n}^{\mathfrak{s} \rightarrow \sigma(\mathfrak{s})}$ ,  $\sigma \in S_l$  yields an alternative for computing  $\Psi_{e,n}^{-\mathfrak{s} \rightarrow \mathfrak{s}^\sharp}(\nu) = \mu$  in Section 4.3. Nevertheless, one can verify that the number of elementary transformations it necessitates (i.e. the number of bijections  $\Psi_{e,n}^{\mathfrak{s} \rightarrow s_c(\mathfrak{s})}$ ,  $c = 1, \dots, l-1$  or  $\Psi_{e,n}^{\mathfrak{s} \rightarrow \tau(\mathfrak{s})}$  to compute) is in general greater than the number of transformations used in  $\Psi_{e,n}^{-\mathfrak{s} \rightarrow \eta(-\mathfrak{s})}$ .

**Example 4.4.** We conclude this paper with a computation of the Mullineux involution on a 3-partition. Take  $e = 4$  and  $\mathfrak{s} = (0 \pmod{4}, 1 \pmod{4}, 3 \pmod{4})$ . Then, one can check that the 3-partition  $\lambda := (4, 3, 1.1.1)$  is a Kleshchev 3-partition of rank  $n = 10$  associated to  $\mathfrak{s}$ . Now we have

$$m_1(4) = (2.1.1), \quad m_1(3) = (1.1.1), \quad m_1(1.1.1) = (3).$$

Hence, to  $\lambda$ , one can attach the Kleshchev 3-partition  $\nu = (2.1.1, 1.1.1, 3)$  which is Kleshchev for  $-\mathfrak{s} = (0 \pmod{4}, -1 \pmod{4}, -3 \pmod{4})$ . Since

$p_1 = p_2 = 3$ , This 3-partition is an Uglov 3-partition associated with the multicharge

$$-\mathbf{s} = (0, -1 + 3 \times 4, -3 + 3 \times 4 + 3 \times 4) = (0, 11, 21).$$

Following §4.3, we put :

$$\eta = \alpha_1^8 \alpha_2^8 w_0.$$

We compute

$$\Psi_{e,n}^{-\mathbf{s} \rightarrow \eta \cdot (-\mathbf{s})}(\boldsymbol{\nu}) = (\emptyset, 1.1.1, 4.1.1.1).$$

Hence, we obtain

$$m_3(\boldsymbol{\lambda}) = (\emptyset, 1.1.1, 4.1.1.1).$$

**Acknowledgements.** The authors are grateful to A. Kleshchev for useful comments and for having pointed out several references. Part of this work has been written while the authors were visiting the Mathematical Research Institute of Mathematics in Berkeley for the programs “Combinatorial Representation Theory” and “Representation Theory of Finite Groups and Related Topics”. The authors want to thank the organizers of these programs and the MSRI for financial supports.

#### REFERENCES

- [1] S. ARIKI, On the classification of simple modules for cyclotomic Hecke algebra of type  $G(m, 1, n)$  and Kleshchev multipartitions. *Osaka J. Math.* **38** (2001), 827-837.
- [2] S. ARIKI AND N. JACON, Dipper-James-Murphy’s conjecture for Hecke algebras of type  $B_n$ , *Progress in Math.*, Birkhauser, “Representation Theory of Algebraic Groups and Quantum Groups”, to appear, [arXiv:math/0703447](https://arxiv.org/abs/math/0703447).
- [3] S. ARIKI AND A. MATHAS, The number of simple modules of the Hecke algebras of type  $G(r, 1, n)$ . *Math. Z.* **233** (2000), 601-623.
- [4] S. ARIKI, V. KREIMAN, S. TSUCHIOKA, On the tensor product of two basis representations of  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ , *Adv. Math.* **218** (2008), 28-86.
- [5] C. BESSENRODT AND J.B. OLSSON, On residue symbols and the Mullineux conjecture. *J. Algebraic Combin.* **7** (1998), no. 3, 227–251.
- [6] J. BRUNDAN, Modular Branching rules and the Mullineux map for Hecke algebras of type  $A$ , *Proc. London Math.* **77** (1998), 551-581.
- [7] J. BRUNDAN AND J. KUJAWA, A new proof of the Mullineux conjecture. *J. Algebraic Combin.* **18** (2003), no. 1, 13–39.
- [8] R. DIPPER AND A. MATHAS, Morita equivalences of Ariki-Koike algebras. *Math. Z.* **240** (2002), no. 3, 579–610.
- [9] M. FAYERS, Weights of multipartitions and representations of Ariki-Koike algebras II : canonical bases., *Journal of Algebra* **319** (2008), 2963–2978.
- [10] O. FODA, B. LECLERC, M. OKADO, J.-Y. THIBON AND T. WELSH, Branching functions of  $A_{n-1}^{(1)}$  and Jantzen-Seitz problem for Ariki-Koike algebras, *Advances in Math.* **141** (1999), 322–365
- [11] J.J. GRAHAM, G.I. LEHRER, Cellular algebras, *Invent. Math.* **123** (1996) 1–34.
- [12] B.. FORD AND A. KLESHCHEV, A proof of the Mullineux conjecture. *Math. Z.* **226** (1997), no. 2, 267–308.
- [13] M. JIMBO, K. C. MISRA, T. MIWA AND M. OKADO, Combinatorics of representations of  $\mathcal{U}_q(\widehat{\mathfrak{sl}}(n))$  at  $q = 0$ , *Communication in Mathematical Physics* **136** (1991), 543-566.
- [14] N. JACON, Crystal graphs of irreducible highest weight  $\mathcal{U}_v(\widehat{\mathfrak{sl}}_e)$ -modules of level two and Uglov bipartitions, *J. Alg. Comb.*, **27**, no. 2 (2008), 143-162,



- [15] N. JACON, C. LECOUEY, Crystal isomorphisms for irreducible highest weight  $\mathcal{U}_v(\widehat{\mathfrak{sl}_e})$ -modules of higher level, preprint, [arXiv:math/0706.0680](https://arxiv.org/abs/math/0706.0680).
- [16] A. KLESHCHEV, Branching rules for modular representations of symmetric groups III: some corollaries and a problem of Mullineux, *J. Lond. Math. Soc.* **54** (1996), 25-38.
- [17] A. LASCoux, B. LECLERC, J-Y THIBON, Hecke algebras at roots of unity and crystal bases of quantum affine algebras. *Comm. Math. Phys.*, **181** no. 1, 205-263 : 1996.
- [18] B. LECLERC, J-Y. THIBON, E.. VASSEROT, Zelevinsky's involution at roots of unity. *J. Reine Angew. Math.* **513** (1999), 33-51.
- [19] A. MATHAS, The representation theory of the Ariki-Koike and cyclotomic  $q$ -Schur algebras. Representation theory of algebraic groups and quantum groups, 261-320, *Adv. Stud. Pure Math.*, **40**, Math. Soc. Japan, Tokyo, 2004.
- [20] G. MULLINEUX, Bijections of  $p$ -regular partitions and  $p$ -modular irreducibles of the symmetric groups. *J. London Math. Soc. (2)* **20** (1979), no. 1, 60-66.
- [21] X. YVONNE, A conjecture for  $q$ -decomposition matrices of cyclotomic  $v$ -Schur algebras, *J. Algebra*, **304** (2006), 419-456.

N.J.:UNIVERSITÉ DE FRANCHE-COMTÉ, UFR SCIENCES ET TECHNIQUES, 16 ROUTE DE GRAY, 25 030 BESANÇON, FRANCE.

*E-mail address:* [njacon@univ-fcomte.fr](mailto:njacon@univ-fcomte.fr)

C. L. LABORATOIRE DE MATHÉMATIQUES PURES ET APPLIQUÉES JOSEPH LIOUVILLE CENTRE UNIVERSITAIRE DE LA MI-VOIX B.P. 699 62228 CALAIS FRANCE

*E-mail address:* [Cedric.Lecouvey@lmpa.univ-littoral.fr](mailto:Cedric.Lecouvey@lmpa.univ-littoral.fr)