



HAL
open science

Hilbert domains quasi-isometric to normed vector spaces

Bruno Colbois, Patrick Verovic

► **To cite this version:**

Bruno Colbois, Patrick Verovic. Hilbert domains quasi-isometric to normed vector spaces. 2008.
hal-00271381v1

HAL Id: hal-00271381

<https://hal.science/hal-00271381v1>

Preprint submitted on 9 Apr 2008 (v1), last revised 27 May 2009 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HILBERT DOMAINS QUASI-ISOMETRIC TO NORMED VECTOR SPACES

BRUNO COLBOIS AND PATRICK VEROVIC

ABSTRACT. We prove that a Hilbert domain which is quasi-isometric to a normed vector space is actually a convex polytope.

1. INTRODUCTION

A *Hilbert domain* in \mathbf{R}^m is a metric space $(\mathcal{C}, d_{\mathcal{C}})$, where \mathcal{C} is an *open bounded convex* set in \mathbf{R}^m and $d_{\mathcal{C}}$ is the distance function on \mathcal{C} — called the *Hilbert metric* — defined as follows.

Given two distinct points p and q in \mathcal{C} , let a and b be the intersection points of the straight line defined by p and q with $\partial\mathcal{C}$ so that $p = (1 - s)a + sb$ and $q = (1 - t)a + tb$ with $0 < s < t < 1$. Then


$$d_{\mathcal{C}}(p, q) := \frac{1}{2} \ln[a, p, q, b],$$

where

$$[a, p, q, b] := \frac{1 - s}{s} \times \frac{t}{1 - t} > 1$$

is the cross ratio of the 4-tuple of ordered collinear points (a, p, q, b) .

We complete the definition by setting $d_{\mathcal{C}}(p, p) := 0$.



The metric space $(\mathcal{C}, d_{\mathcal{C}})$ thus obtained is a complete non-compact geodesic metric space whose topology is the one induced by the canonical topology of \mathbf{R}^m and in which the affine open segments joining two points of the boundary $\partial\mathcal{C}$ are geodesics that are isometric to $(\mathbf{R}, |\cdot|)$.

For further information about Hilbert geometry, we refer to [3, 4, 8, 10] and the excellent introduction [14] by Socié-Méthou.

The two fundamental examples of Hilbert domains $(\mathcal{C}, d_{\mathcal{C}})$ in \mathbf{R}^m correspond to the case when \mathcal{C} is an ellipsoid, which gives the Klein model of m -dimensional hyperbolic geometry (see for

Date: April 10, 2008.

2000 Mathematics Subject Classification. Primary: global Finsler geometry, Secondary: convexity.

example [14, first chapter]), and the case when $\bar{\mathcal{C}}$ is a m -simplex, for which there exists a norm $\|\cdot\|_{\mathcal{C}}$ on \mathbf{R}^m such that $(\mathcal{C}, d_{\mathcal{C}})$ is isometric to the normed vector space $(\mathbf{R}^m, \|\cdot\|_{\mathcal{C}})$ (see [7, pages 110–113] or [13, pages 22–23]).

Much has been done to study the similarities between Hilbert and hyperbolic geometries (see for example [6], [15] or [1]), but little literature deals with the question of knowing to what extent a Hilbert geometry is close to that of a normed vector space. So let us mention two results in this latter direction which are relevant for our present work.

Theorem 1.1 ([9], Theorem 2). *A Hilbert domain $(\mathcal{C}, d_{\mathcal{C}})$ in \mathbf{R}^m is isometric to a normed vector space if and only if \mathcal{C} is the interior of a m -simplex.*

Theorem 1.2 ([5], Theorem 3.1). *If \mathcal{C} is an open bounded convex polygonal set in \mathbf{R}^2 , then $(\mathcal{C}, d_{\mathcal{C}})$ is Lipschitz equivalent to Euclidean plane.*

In light of these two results, it is natural to ask whether the converse of Theorem 1.2 holds. In other words, if a Hilbert domain $(\mathcal{C}, d_{\mathcal{C}})$ in \mathbf{R}^m is quasi-isometric to a normed vector space, what can be said about \mathcal{C} ? Here, by *quasi-isometric* we mean the following (see [2]):

Definition 1.1. Given real numbers $A \geq 1$ and $B \geq 0$, a metric space (S, d) is said to be (A, B) -quasi-isometric to a normed vector space $(V, \|\cdot\|)$ if and only if there exists a map $f : S \rightarrow V$ such that

$$\frac{1}{A}d(p, q) - B \leq \|f(p) - f(q)\| \leq Ad(p, q) + B$$

for all $p, q \in S$.

Recalling that a convex *polytope* in \mathbf{R}^m is the convex hull of a finite set of points whose linear span is the whole space \mathbf{R}^m , we then show

Theorem 1.3. *If a Hilbert domain $(\mathcal{C}, d_{\mathcal{C}})$ in \mathbf{R}^m is (A, B) -quasi-isometric to a normed vector space $(V, \|\cdot\|)$ for some real constants $A \geq 1$ and $B \geq 0$, then \mathcal{C} is the interior of a convex polytope.*

2. PROOF OF THEOREM 1.3

The proof of Theorem 1.3 is based on an idea developed by Förtsch and Karlsson in their paper [9].

It needs the following fact due to Karlsson and Noskov:

Theorem 2.1 ([11], Theorem 5.2). *Let $(\mathcal{C}, d_{\mathcal{C}})$ be a Hilbert domain in \mathbf{R}^m and $x, y \in \partial\mathcal{C}$ such that $[x, y] \not\subseteq \partial\mathcal{C}$. Then, given a point $p_0 \in \mathcal{C}$, there exists a constant $K = K(p_0, x, y) > 0$ satisfying*

$$d_{\mathcal{C}}(x_n, y_n) \geq d_{\mathcal{C}}(x_n, p_0) + d_{\mathcal{C}}(y_n, p_0) - K$$

for all sequences $(x_n)_{n \in \mathbf{N}}$ and $(y_n)_{n \in \mathbf{N}}$ in \mathcal{C} that converge respectively to x and y in \mathbf{R}^m .

Now, here is the key result which gives the proof of Theorem 1.3:

Proposition 2.1. *Let $(\mathcal{C}, d_{\mathcal{C}})$ be a Hilbert domain in \mathbf{R}^m which is (A, B) -quasi-isometric to a normed vector space $(V, \|\cdot\|)$ for some real constants $A \geq 1$ and $B \geq 0$.*

Then, if $N = N(A, \|\cdot\|)$ denotes the maximum number of points in the ball $\{v \in V \mid \|v\| \leq 2A\}$ whose pairwise distances with respect to $\|\cdot\|$ are greater than or equal to $1/(2A)$, and if $X \subseteq \partial\mathcal{C}$ is such that $[x, y] \not\subseteq \partial\mathcal{C}$ for all $x, y \in X$ with $x \neq y$, we have

$$\text{card}(X) \leq N.$$

Proof.

Let $f : \mathcal{C} \rightarrow V$ such that

$$(2.1) \quad \frac{1}{A}d_{\mathcal{C}}(p, q) - B \leq \|f(p) - f(q)\| \leq Ad_{\mathcal{C}}(p, q) + B$$

for all $p, q \in \mathcal{C}$.

First of all, up to translations, we may assume that $0 \in \mathcal{C}$ and $f(0) = 0$.

Then suppose that there exists a subset X of the boundary $\partial\mathcal{C}$ such that $[x, y] \not\subseteq \partial\mathcal{C}$ for all $x, y \in X$ with $x \neq y$ and $\text{card}(X) \geq N + 1$. So, pick $N + 1$ distinct points x_1, \dots, x_{N+1} in X , and for each $k \in \{1, \dots, N + 1\}$, let $\gamma_k : [0, +\infty) \rightarrow \mathcal{C}$ be a geodesic of $(\mathcal{C}, d_{\mathcal{C}})$ that satisfies $\gamma_k(0) = 0$, $\lim_{t \rightarrow +\infty} \gamma_k(t) = x_k$ in \mathbf{R}^m and $d_{\mathcal{C}}(0, \gamma_k(t)) = t$ for all $t \geq 0$.

This implies that for all integers $n \geq 1$ and every $k \in \{1, \dots, N + 1\}$, we have

$$(2.2) \quad \left\| \frac{f(\gamma_k(n))}{n} \right\| \leq A + \frac{B}{n}$$

from the second inequality in Equation 2.1 with $p = \gamma_k(n)$ and $q = 0$.

On the other hand, Theorem 2.1 yields

$$d_{\mathcal{C}}(\gamma_i(n), \gamma_j(n)) \geq 2n - K(0, x_i, x_j)$$

for all integers $n \geq 1$ and every $i, j \in \{1, \dots, N + 1\}$ with $i \neq j$, and hence

$$(2.3) \quad \left\| \frac{f(\gamma_i(n))}{n} - \frac{f(\gamma_j(n))}{n} \right\| \geq \frac{2}{A} - \frac{1}{n} \left(\frac{K(0, x_i, x_j)}{A} - B \right)$$

from the first inequality in Equation 2.1 with $p = \gamma_i(n)$ and $q = \gamma_j(n)$.

Now, fixing an integer $n \geq \max\{B/A, K(0, x_i, x_j) - B/A\}$, we get

$$\left\| \frac{f(\gamma_k(n))}{n} \right\| \leq 2A$$

for all $k \in \{1, \dots, N + 1\}$ by Equation 2.2 together with

$$\left\| \frac{f(\gamma_i(n))}{n} - \frac{f(\gamma_j(n))}{n} \right\| \geq \frac{1}{2A}$$

for all $i, j \in \{1, \dots, N + 1\}$ with $i \neq j$ by Equation 2.3.

But this contradicts the definition of $N = N(A, \|\cdot\|)$.

Therefore, Proposition 2.1 is proved. □

Remark. Given $v \in V$ such that $\|v\| = 2A$, we have $\|-v\| = 2A$ and $\|v - (-v)\| = 2\|v\| = 4A \geq 1/(2A)$, which proves that $N \geq 2$.

The second ingredient we will need for the proof of Theorem 1.3 is the following:

Proposition 2.2. *Let \mathcal{C} be an open bounded convex set in \mathbf{R}^2 .*

If there exists a nonempty finite subset Y of the boundary $\partial\mathcal{C}$ such that for every $x \in \partial\mathcal{C}$ one can find $y \in Y$ with $[x, y] \subseteq \partial\mathcal{C}$, then $\overline{\mathcal{C}}$ is a polygon.

Proof.

Assume $0 \in \mathcal{C}$ and let us consider the continuous map $\pi : \mathbf{R} \longrightarrow \partial\mathcal{C}$ which assigns to each $\theta \in \mathbf{R}$ the unique intersection point $\pi(\theta)$ of $\partial\mathcal{C}$ with the half-line $\mathbf{R}_+^*(\cos \theta, \sin \theta)$.

For each pair $(x_1, x_2) \in \partial\mathcal{C} \times \partial\mathcal{C}$, denote by $A(x_1, x_2) \subseteq \partial\mathcal{C}$ the arc segment defined by $A(x_1, x_2) := \pi([\theta_1, \theta_2])$, where θ_1 and θ_2 are the unique real numbers such that $\pi(\theta_1) = x_1$ with $\theta_1 \in [0, 2\pi)$ and $\pi(\theta_2) = x_2$ with $\theta_1 \leq \theta_2 < \theta_1 + 2\pi$.

We shall prove Proposition 2.2 by induction on $n := \text{card}(Y) \geq 1$.

Before doing this, notice that adding a point of $\partial\mathcal{C}$ to Y does not change Y 's property, and therefore we may assume that $n \geq 2$.

If $n = 2$, writing $Y = \{x_1, x_2\}$, we have that for all $x \in \partial\mathcal{C}$, $[x, x_1] \subseteq \partial\mathcal{C}$ or $[x, x_2] \subseteq \partial\mathcal{C}$.

Let $\theta_1 \in [0, 2\pi)$ and $\theta_2 \in [\theta_1, \theta_1 + 2\pi)$ such that $\pi(\theta_1) = x_1$ and $\pi(\theta_2) = x_2$, and define $\alpha_0 := \max\{\theta \in [\theta_1, \theta_2] \mid [x_1, \pi(\theta)] \subseteq \partial\mathcal{C}\}$.

Then $A(x_1, x_2) = [x_1, \pi(\alpha_0)] \cup [\pi(\alpha_0), x_2]$. Hence, $A(x_1, x_2)$ is the union of two affine segments, and the same holds for $A(x_2, x_1)$.

Since $\partial\mathcal{C} = A(x_1, x_2) \cup A(x_2, x_1)$, this implies that $\partial\mathcal{C}$ is a union of four affine segments in \mathbf{R}^2 , and therefore $\overline{\mathcal{C}}$ is a polygon (quadrilateral or triangle).

Now, assume Proposition 2.2 is true for some $n \geq 2$, and let Y such that $\text{card}(Y) = n + 1$.

Write $Y = \{x_1, \dots, x_{n+1}\}$ with $x_1 = \pi(\theta_1)$, $x_2 = \pi(\theta_2)$, \dots , $x_{n+1} = \pi(\theta_{n+1})$, where $\theta_1 \in [0, 2\pi)$ and $\theta_1 < \theta_2 < \dots < \theta_{n+1} < \theta_{n+2} := \theta_1 + 2\pi$.

Then, defining $x_{n+2} := \pi(\theta_{n+2}) = x_1$, we shall consider two cases.

• **Case 1:** Suppose for every $k \in \{1, \dots, n+1\}$ and all $x \in A(x_k, x_{k+1})$, we have $[x, x_k] \subseteq \partial\mathcal{C}$ or $[x, x_{k+1}] \subseteq \partial\mathcal{C}$.

Considering $\alpha_k := \max\{\theta \in [\theta_k, \theta_{k+1}] \mid [x_k, \pi(\theta)] \subseteq \partial\mathcal{C}\}$, then we get $A(x_k, x_{k+1}) = [x_k, \pi(\alpha_k)] \cup [\pi(\alpha_k), x_{k+1}]$.

Since $\partial\mathcal{C} = \bigcup_{k=1}^{n+1} A(x_k, x_{k+1})$, this implies that $\partial\mathcal{C}$ is a finite union of affine segments in \mathbf{R}^2 , and thus $\overline{\mathcal{C}}$ is a polygon.

• **Case 2:** Suppose there exist $k \in \{1, \dots, n+1\}$ — we may assume to be equal to $n+1$ up to permutations — and $x_0 \in A(x_k, x_{k+1}) = A(x_{n+1}, x_1)$ such that $x_0 \notin \{x_1, x_{n+1}\}$ and $[x_0, x_i] \subseteq \partial\mathcal{C}$ for some $i \in \{2, \dots, n\}$.

Then \mathcal{C} is contained in one of the two open half-planes in \mathbf{R}^2 bounded by the line passing through the points x_0 and x_i , and hence either $A(x_0, x_i) = [x_0, x_i]$, or $A(x_i, x_0) = [x_0, x_i]$.

But $x_1 \in A(x_0, x_i)$ and $x_{n+1} \in A(x_i, x_0)$, so we get either $x_1 \in [x_0, x_i] \subseteq \partial\mathcal{C}$, or $x_{n+1} \in [x_0, x_i] \subseteq \partial\mathcal{C}$. And this implies that Y can be replaced by $Y \setminus \{x_1\}$ or $Y \setminus \{x_{n+1}\}$ in the hypothesis of Proposition 2.2.

Since $\text{card}(Y \setminus \{x_1\}) = \text{card}(Y \setminus \{x_{n+1}\}) = n$, we get that Proposition 2.2 is true for $n + 1$.

This finishes the proof of Proposition 2.2 by induction. \square

Before proving Theorem 1.3, let us recall the following useful result, where a convex *polyhedron* in \mathbf{R}^m is the intersection of a finite number of closed half-spaces:

Theorem 2.2 ([12], Theorem 4.7). *Let K be a convex set in \mathbf{R}^m and $p \in \overset{\circ}{K}$.*

Then K is a convex polyhedron if and only if all its plane sections containing p are convex polyhedra.

Proof of Theorem 1.3.

Let $(\mathcal{C}, d_{\mathcal{C}})$ be a nonempty Hilbert domain in \mathbf{R}^m that is (A, B) -quasi-isometric to a normed vector space $(V, \|\cdot\|)$ for some real constants $A \geq 1$ and $B \geq 0$.

According to Theorem 2.2, it suffices to prove Theorem 1.3 for $m = 2$ since any plane section of \mathcal{C} gives rise to a 2-dimensional Hilbert domain which is also (A, B) -quasi-isometric to $(V, \|\cdot\|)$.

So, let $m = 2$, and consider the set $\mathcal{E} := \{X \subseteq \partial\mathcal{C} \mid [x, y] \not\subseteq \partial\mathcal{C} \text{ for all } x, y \in X \text{ with } x \neq y\}$.

It is not empty since $\{x, y\} \in \mathcal{E}$ for some $x, y \in \partial\mathcal{C}$ with $x \neq y$ (indeed, \mathcal{C} is a nonempty open set in \mathbf{R}^2), which implies together with Proposition 2.1 that $n := \max\{\text{card}(X) \mid X \in \mathcal{E}\}$ does exist and satisfies $2 \leq n \leq N$ (recall that $N \geq 2$).

Then pick $Y \in \mathcal{E}$ such that $\text{card}(Y) = n$, write $Y = \{x_1, \dots, x_n\}$, and prove that for every $x \in \partial\mathcal{C}$ one can find $k \in \{1, \dots, n\}$ such that $[x, x_k] \subseteq \partial\mathcal{C}$.

Owing to Proposition 2.2, this will show that $\overline{\mathcal{C}}$ is a polygon.

So, suppose that there exists $x_0 \in \partial\mathcal{C}$ satisfying $[x_0, x_k] \not\subseteq \partial\mathcal{C}$ for all $k \in \{1, \dots, n\}$, and let us find a contradiction by considering $Z := Y \cup \{x_0\}$.

First, since $x_0 \neq x_k$ for all $k \in \{1, \dots, n\}$ (if not, we would get an index $k \in \{1, \dots, n\}$ such that $[x_0, x_k] = \{x_0\} \subseteq \partial\mathcal{C}$, which is false), we have $x_0 \notin Y$. Hence $\text{card}(Z) = n + 1$.

Next, since $Y \in \mathcal{E}$ and $[x_0, x_k] \not\subseteq \partial\mathcal{C}$ for all $k \in \{1, \dots, n\}$, we have $Z \in \mathcal{E}$.

Therefore, the assumption of the existence of x_0 yields a set $Z \in \mathcal{E}$ whose cardinality is greater than that of Y , which contradicts the very definition of Y .

Conclusion: $\overline{\mathcal{C}}$ is a polygon, and this proves Theorem 1.3. \square

Let us conclude by a natural question: Considering Theorem 1.1 by Förtsch and Karlsson, we may conjecture that if the quasi-isometry in Theorem 1.3 is close to an isometry, then \mathcal{C} is the interior of a m -simplex. More precisely, does there exist a sufficiently small constant $A_0 = A_0(m) \geq 1$ such that if a Hilbert domain $(\mathcal{C}, d_{\mathcal{C}})$ in \mathbf{R}^m is (A, B) -quasi-isometric to a normed vector space for some constants $A = A(m, \mathcal{C}) \in [1, A_0]$ and $B = B(m, \mathcal{C}) \geq 0$, then \mathcal{C} is the interior of a m -simplex?

REFERENCES

- [1] BENOIST, Y. A survey on convex divisible sets. Lecture notes, International Conference and Instructional Workshop on Discrete Groups in Beijing, 2006.
- [2] BURAGO, D., BURAGO, Y., AND IVANOV, S. *A course in metric geometry*. AMS, 2001.
- [3] BUSEMANN, H. *The geometry of geodesics*. Academic Press, 1955.
- [4] BUSEMANN, H., AND KELLY, P. *Projective geometry and projective metrics*. Academic Press, 1953.
- [5] COLBOIS, B., VERNICOS, C., AND VEROVIC, P. Hilbert geometry for convex polygonal domains. Tech. rep., University of Neuchâtel, 2008.
- [6] COLBOIS, B., AND VEROVIC, P. Hilbert geometry for strictly convex domains. *Geom. Dedicata* 105 (2004), 29–42.
- [7] DE LA HARPE, P. On Hilbert’s metric for simplices. *Lond. Math. Soc. Lect. Note Ser.* 1, 181 (1993), 97–119.
- [8] EGLOFF, D. Uniform Finsler Hadamard manifolds. *Ann. Inst. Henri Poincaré, Phys. Théor.* 66 (1997), 323–357.
- [9] FÖRTSCH, T., AND KARLSSON, A. Hilbert metrics and Minkowski norms. *J. Geom.* 83, 1-2 (2005), 22–31.
- [10] GOLDMAN, W. Projective geometry on manifolds. Lecture notes, University of Maryland, 1988.
- [11] KARLSSON, A., AND NOSKOV, G. Some groups having only elementary actions on metric spaces with hyperbolic boundaries. *Geom. Dedicata* 104 (2004), 119–137.
- [12] KLEE, V. Some characterizations of convex polyhedra. *Acta Math.* 102 (1959), 79–107.
- [13] NUSSBAUM, R. Hilbert’s projective metric and iterated nonlinear maps. *Mem. Am. Math. Soc.* 75, 391 (1988).

- [14] SOCIÉ-MÉTHOU, É. *Comportements asymptotiques et rigidités des géométries de Hilbert*. PhD thesis, University of Strasbourg, 2000.
- [15] VERNICOS, C. Introduction aux géométries de Hilbert. *Sémin. Théor. Spectr. Géom.* 23 (2005), 145–168.

BRUNO COLBOIS, UNIVERSITÉ DE NEUCHÂTEL, INSTITUT DE MATHÉMATIQUE, RUE ÉMILE ARGAND 11,
CASE POSTALE 158, CH-2009 NEUCHÂTEL, SWITZERLAND
E-mail address: `bruno.colbois@unine.ch`

PATRICK VEROVIC, UMR 5127 DU CNRS & UNIVERSITÉ DE SAVOIE, LABORATOIRE DE MATHÉMATIQUE,
CAMPUS SCIENTIFIQUE, 73376 LE BOURGET-DU-LAC CEDEX, FRANCE
E-mail address: `verovic@univ-savoie.fr`