



Continuous crystals and Duistermaat-Heckman measure for Coxeter groups.

Philippe Biane, Philippe Bougerol, Neil O'Connell

► To cite this version:

Philippe Biane, Philippe Bougerol, Neil O'Connell. Continuous crystals and Duistermaat-Heckman measure for Coxeter groups.. 2008. hal-00273468v1

HAL Id: hal-00273468

<https://hal.science/hal-00273468v1>

Preprint submitted on 15 Apr 2008 (v1), last revised 14 Jan 2009 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Continuous crystal and Duistermaat-Heckmann measure for Coxeter groups.

Philippe Biane, Philippe Bougerol, and Neil O’Connell

ABSTRACT. We introduce a notion of continuous crystal analogous, for general Coxeter groups, to the combinatorial crystals introduced by Kashiwara in representation theory of Lie algebras. We explore their main properties in the case of finite Coxeter groups, where we use a generalization of the Littelmann path model to show the existence of the crystals. We introduce a remarkable measure, analogous to the Duistermaat-Heckman measure, which we interpret in terms of Brownian motion. We also show that the Littelmann path operators can be derived from simple considerations on Sturm-Liouville equations.

1. Introduction

The aim of this paper is to introduce a notion of continuous crystals for Coxeter groups, which are not necessarily Weyl groups. Crystals are combinatorial objects, which have been associated by Kashiwara to Kac-Moody algebras, in order to provide a combinatorial model for the representation theory of these algebras, see e.g. [16], [18], [19], [22] for an introduction to this theory. The crystal graphs defined by Kashiwara turn out to be equivalent to certain other graphs, constructed independently by Littelmann, using his path model. The approach of Kashiwara to the crystals is through representations of quantum groups and their “crystallization”, which is the process of letting the parameter q in the quantum group go to zero. This requires representation theory and therefore does not make sense for realizations of arbitrary Coxeter groups. On the other hand, as it was realized in a previous paper [2], Littelmann’s model can be adapted to fit with non-crystallographic Coxeter groups, but the price to pay is that, since there is no lattice invariant under the action of the group, one can only define a continuous version of the path model, namely of the Littelmann path operators. In this continuous model, instead of the the Littelmann path operators e_i, f_i we have continuous semigroups e_i^t, f_i^t indexed by nonnegative real numbers $t \geq 0$. In the crystallographic case it is possible to think of these continuous crystals as “semi-classical limits” of the combinatorial crystals, in much the same way as the coadjoint orbits arise as semi-classical limits of the representations of a compact semi-simple Lie group. These continuous path operators, and the closely related Pitman transforms, were used in [2] to investigate

1991 *Mathematics Subject Classification.* Primary ; Secondary .

Research of the third author supported in part by Science Foundation Ireland Grant No. SFI04/RP1/I512.

symmetry properties of Brownian motion in a space where a finite Coxeter group acts, with applications in particular to the motion of eigenvalues of matrix-valued Brownian motions. In this paper, which is a sequel to [2], but can for the most part be read independently, we define continuous crystals and start investigating their main properties. As for now the theory works well for finite Coxeter groups, there are still several difficulties to extend the theory to infinite groups. This theory allows us to define objects which are analogues to simplified versions of the Schubert varieties (or Demazure-Littelmann modules) associated with semi-simple Lie groups. We hope these objects might help in certain questions concerning Coxeter groups, such as, for example, the Kazhdan-Lusztig polynomials.

This paper is organized as follows. The next section contains the main definition, that of a continuous crystal associated with a realization of a Coxeter group. We establish the main properties of these objects, following closely the exposition of Joseph in [19]. It would have been possible to just refer to [19] for the most part of this section, however, for the convenience of the reader, and also for convincing ourselves that everything from the crystallographic situation goes smoothly to the continuous context, we have preferred to write everything down. The main body of the proof is relegated to an appendix in order to ease the reading of the paper. In section 3 we introduce the path operators and establish their most important properties. The path operators are used in section 4 to introduce the path model, which provides a concrete realization of continuous crystals and proves their existence, at least in the finite Coxeter group case. It remains an interesting and challenging problem to extend these properties to the general case. Our approach to the path model is different from that in Littelmann [25], in that we base our exposition on the Pitman transforms, which are defined from scratch, and whose braid relations play a prominent role. We investigate the analogues of the Berenstein-Zelevinsky polytopes (see [3]) which contain the Kashiwara coordinates on the crystals, we also study concatenation of paths, and the action of the Coxeter group, as well as the Schützenberger involution. We think that even in the crystallographic case our treatment sheds some light on these topics. In section 5, we introduce an analogue of the Duistermaat-Heckman measure, motivated by a result of Alexeev and Brion [1]. We prove several interesting properties of this measure, and show that it is intimately linked with Brownian motion. In particular (corollary 5.3) we give a Brownian proof of the fact that the crystal defined by the path model depends only on the final position of the path. The final section is of a quite different nature, and somewhat independent of the rest of the paper. The Littelmann path operators have been introduced as a generalization, for arbitrary root systems, of combinatorial operations on Young tableaux. Here we show how, using some simple considerations on Sturm-Liouville equations, the Littelmann path operators appear naturally. In particular this gives a concrete geometric basis to the theory of geometric lifting which has been introduced by Berenstein and Zelevinsky in [3] in a purely formal way.

2. Continuous crystal

2.1. Basic definition. We use the standard references [4], [17] on Coxeter groups and their realizations. A Coxeter system (W, S) is a group W generated by a finite set of involutions S such that, if $m(s, s')$ is the order of ss' then the

relations

$$(ss')^{m(s,s')} = 1$$

for $m(s, s')$ finite, give a presentation of W .

A realization of (W, S) is given by a real vector space V with dual V^\vee , an action of W on V , and a subset $\{(\alpha_s, \alpha_s^\vee), s \in S\}$ of $V \times V^\vee$ such that each $s \in S$ acts on V by the reflection given by

$$s(x) = x - \alpha_s^\vee(x)\alpha_s, \quad x \in V,$$

so $\alpha_s^\vee(\alpha_s) = 2$. One calls α_s the simple root associated with $s \in S$ and α_s^\vee its coroot.

We consider a realization of a Coxeter system (W, S) in a real vector space V , and the associated simple roots $\Sigma = \{\alpha_s, s \in S\}$ in V and coroots $\{\alpha_s^\vee, s \in S\}$ in V^\vee . The closed Weyl chamber is the convex cone

$$\overline{C} = \{v \in V; \alpha_s^\vee(v) \geq 0, \text{ for all } \alpha \in \Sigma\}$$

thus the simple roots are positive on \overline{C} . There is an order relation on V induced by this cone, namely $\lambda \leq \mu$ if and only if $\mu - \lambda \in \overline{C}$.

We adapt the definition of crystals due to Kashiwara (see, e.g., Kashiwara [20], [22], Joseph [18]) to a continuous setting.

DEFINITION 2.1. *A continuous crystal is a set B equipped with maps*

$$\begin{aligned} wt &: B \rightarrow V, \\ \varepsilon_\alpha, \varphi_\alpha &: B \rightarrow \mathbb{R} \cup \{-\infty\}, \alpha \in \Sigma, \\ e_\alpha^r &: B \cup \{\mathbf{0}\} \rightarrow B \cup \{\mathbf{0}\}, \alpha \in \Sigma, r \in \mathbb{R}, \end{aligned}$$

where $\mathbf{0}$ is a ghost element, such that the following properties hold, for all $\alpha \in \Sigma$, and $b \in B$

$$(C1) \quad \varphi_\alpha(b) = \varepsilon_\alpha(b) + \alpha^\vee(wt(b)),$$

$$(C2) \quad \text{If } e_\alpha^r(b) \neq \mathbf{0} \text{ then}$$

$$\begin{aligned} \varepsilon_\alpha(e_\alpha^r b) &= \varepsilon_\alpha(b) - r, \\ \varphi_\alpha(e_\alpha^r b) &= \varphi_\alpha(b) + r, \\ wt(e_\alpha^r b) &= wt(b) + r\alpha, \end{aligned}$$

(C3) For all $r \in \mathbb{R}, b \in B$ one has $e_\alpha^r(\mathbf{0}) = \mathbf{0}, e_\alpha^0(b) = b$. If $e_\alpha^r(b) \neq \mathbf{0}$ then, for all $s \in \mathbb{R}$,

$$e_\alpha^{s+r}(b) = e_\alpha^s(e_\alpha^r(b)),$$

$$(C4) \quad \text{If } \varphi_\alpha(b) = -\infty \text{ then } e_\alpha^r(b) = \mathbf{0}, \text{ for all } r \in \mathbb{R}, r \neq 0.$$

Sometimes we write, for $r \geq 0$,

$$f_\alpha^r = e_\alpha^{-r}.$$

EXAMPLE 2.2 (The crystal B_α). For each $\alpha \in \Sigma$, we define the crystal B_α as the set $\{b_\alpha(t), t \leq 0\}$ with the maps given by

$$wt(b_\alpha(t)) = t\alpha, \quad \varepsilon_\alpha(b_\alpha(t)) = -t, \quad \varphi_\alpha(b_\alpha(t)) = t,$$

$$e_\alpha^r(b_\alpha(t)) = b_\alpha(t+r) \text{ if } r \leq -t \text{ and } e_\alpha^r(b_\alpha(t)) = \mathbf{0} \text{ otherwise,}$$

and, if $\alpha' \neq \alpha$, $\varepsilon_{\alpha'}(b_\alpha(t)) = -\infty$, $\varphi_{\alpha'}(b_\alpha(t)) = -\infty$, $e_{\alpha'}^r(b_\alpha(t)) = \mathbf{0}$, when $r \neq 0$.

2.2. Morphisms.

DEFINITION 2.3. Let B_1 and B_2 be continuous crystals.

1. A morphism of crystals is a map $\psi : B_1 \cup \{\mathbf{0}\} \rightarrow B_2 \cup \{\mathbf{0}\}$ such that $\psi(\mathbf{0}) = \mathbf{0}$ and for all $\alpha \in \Sigma$ and $b \in B_1$,

$$wt(\psi(b)) = wt(b), \varepsilon_\alpha(\psi(b)) = \varepsilon_\alpha(b), \varphi_\alpha(\psi(b)) = \varphi_\alpha(b)$$

and $e_\alpha^r(\psi(b)) = \psi(e_\alpha^r(b))$ when $e_\alpha^r(b) \in B_1$.

2. A strict morphism is a morphism $\psi : B_1 \rightarrow B_2$ such that $e_\alpha^r(\psi(b)) = \psi(e_\alpha^r(b))$ for all $b \in B_1$.

3. A crystal embedding is an injective strict morphism.

The morphism ψ is called a *crystal isomorphism* if there exists a crystal morphism $\phi : B_2 \rightarrow B_1$ such that $\phi \circ \psi = id_{B_1 \cup \{\mathbf{0}\}}$, and $\psi \circ \phi = id_{B_2 \cup \{\mathbf{0}\}}$. It is then an embedding.

2.3. Tensor product. Consider two continuous crystals B_1 and B_2 associated with (W, S, Σ) . We define the tensor product $B_1 \otimes B_2$ as the continuous crystal with set $B = B_1 \times B_2$, whose elements are denoted $b_1 \otimes b_2$, for $b_1 \in B_1, b_2 \in B_2$.

Let $\sigma = \varphi_\alpha(b_1) - \varepsilon_\alpha(b_2)$ where $(-\infty) - (-\infty) = 0$, let $\sigma^+ = \max(0, \sigma)$ and $\sigma^- = \max(0, -\sigma)$, then the maps defining the tensor product are given by the following formulas:

$$\begin{aligned} wt(b_1 \otimes b_2) &= wt(b_1) + wt(b_2) \\ \varepsilon_\alpha(b_1 \otimes b_2) &= \varepsilon_\alpha(b_1) + \sigma^- \\ \phi_\alpha(b_1 \otimes b_2) &= \phi_\alpha(b_2) + \sigma^+ \\ e_\alpha^r(b_1 \otimes b_2) &= e_\alpha^{\max(r, -\sigma) - \sigma^-} b_1 \otimes e_\alpha^{\min(r, -\sigma) + \sigma^-} b_2, \end{aligned}$$

Here $b_1 \otimes \mathbf{0}$ and $\mathbf{0} \otimes b_2$ are understood to be $\mathbf{0}$. Notice that when $\sigma \geq 0$, one has $\varepsilon_\alpha(b_1 \otimes b_2) = \varepsilon_\alpha(b_1)$ and

$$(2.1) \quad e_\alpha^r(b_1 \otimes b_2) = e_\alpha^r b_1 \otimes b_2, \text{ for all } r \in [-\sigma, +\infty[.$$

As in the discrete case, one can check that the tensor product is associative (but not commutative) so we can define without ambiguity the tensor product of several crystals.

2.4. Highest weight crystal. A crystal B is called upper-normal when, for all $b \in B$,

$$\varepsilon_\alpha(b) = \max\{r \geq 0; e_\alpha^r(b) \neq \mathbf{0}\}$$

and is called lower-normal if

$$\varphi_\alpha(b) = \max\{r \geq 0; e_\alpha^{-r}(b) \neq \mathbf{0}\}.$$

We call it normal (this is sometimes called semi-normal by Kashiwara) when it is lower and upper normal. Notice that this implies that $\varepsilon_\alpha(b) \geq 0$ and $\varphi_\alpha(b) \geq 0$.

We introduce the semigroup \mathcal{F} generated by the $\{f_\alpha^r, \alpha \text{ simple root}, r \geq 0\}$:

$$\mathcal{F} = \{f_{\alpha_1}^{r_1} \cdots f_{\alpha_k}^{r_k}, k \in \mathbb{N}^*, r_1, \dots, r_k \geq 0, \alpha_1, \dots, \alpha_k \in \Sigma\},$$

and, if b is an element of a continuous crystal B , the subset $\mathcal{F}(b) = \{f(b), f \in \mathcal{F}\}$ of B .

DEFINITION 2.4. *Let $\lambda \in V$, a continuous crystal $B(\lambda)$ is said to be of highest weight λ if there exists $b_\lambda \in B(\lambda)$ such that $wt(b_\lambda) = \lambda$, $e_\alpha^r(b_\lambda) = \mathbf{0}$, for all $r > 0$ and $\alpha \in \Sigma$ and such that $B(\lambda) = \mathcal{F}(b_\lambda)$.*

For a continuous crystal with highest weight λ , such an element b_λ is unique, and called the primitive element of $B(\lambda)$. If the crystal is normal then λ must be in the Weyl chamber \bar{C} . The vector λ is a highest weight in the sense that, for all $b \in B(\lambda)$, $wt(b) \leq \lambda$.

2.5. Uniqueness. The crystal $B(\infty)$. Following Joseph [18], [19] we introduce the following definition.

DEFINITION 2.5. *Let $(B(\lambda), \lambda \in \bar{C})$, be a family of highest weight continuous crystals. The family is closed if, for each $\lambda, \mu \in \bar{C}$, the subset $\mathcal{F}(b_\lambda \otimes b_\mu)$ of $B(\lambda) \otimes B(\mu)$ is a crystal isomorphic to $B(\lambda + \mu)$*

Joseph ([18], 6.4.21) has shown in the Weyl group case, for discrete crystals, that a closed family of highest weight normal crystals is unique. The analogue holds in our situation.

THEOREM 2.6. *For a realization of a Coxeter system (W, S) , if a closed family $B(\lambda), \lambda \in \bar{C}$, of highest weight continuous normal crystals exists, then it is unique.*

The proof of the theorem, which follows closely Joseph [19], is in the appendix 7.1.

3. Pitman transforms and Littelmann path operators for Coxeter groups

3.1. The Pitman transform. Let V be a real vector space, with dual space V^\vee . Let $\alpha \in V$ and $\alpha^\vee \in V^\vee$ be such that $\alpha^\vee(\alpha) = 2$. The reflection $s_\alpha : V \rightarrow V$ associated to (α, α^\vee) is the linear map defined, for $x \in V$, by

$$s_\alpha(x) = x - \alpha^\vee(x)\alpha.$$

For $T > 0$, let $C_T^0(V)$ be the set of continuous path $\eta : [0, T] \rightarrow V$ such that $\eta(0) = 0$, with the topology of uniform convergence. We have introduced and studied in [2] the following path transformation, similar to the one defined by Pitman in [32].

DEFINITION 3.1. *The Pitman transform \mathcal{P}_α associated with (α, α^\vee) is defined on $C_T^0(V)$ by the formula:*

$$\mathcal{P}_\alpha \eta(t) = \eta(t) - \inf_{t \geq s \geq 0} \alpha^\vee(\eta(s))\alpha, \quad T \geq t \geq 0.$$

A path $\eta \in C_T^0(V)$ is called α -dominant when $\alpha^\vee(\eta(t)) \geq 0$ for all $t \in [0, T]$. The following properties of the Pitman transform are easily established.

PROPOSITION 3.2. *(i) The transformation $\mathcal{P}_\alpha : C_T^0(V) \rightarrow C_T^0(V)$ is continuous.
(ii) For all $\eta \in C_T^0(V)$, the path $\mathcal{P}_\alpha \eta$ is α -dominant and $\mathcal{P}_\alpha \eta = \eta$ if and only if η is α -dominant.
(iii) The transformation \mathcal{P}_α is an idempotent, i.e. $\mathcal{P}_\alpha \mathcal{P}_\alpha \eta = \mathcal{P}_\alpha \eta$ for all $\eta \in C_T^0(V)$.
(iv) If π is α -dominant, and $0 \leq x \leq \alpha^\vee(\pi(T))$, then there exists a unique path η in $C_T^0(V)$ such that*

$$\mathcal{P}_\alpha \eta = \pi$$

and $\eta(T) = \pi(T) - x\alpha$. Moreover for $0 \leq t \leq T$

$$\eta(t) = \pi(t) - \min[x, \inf_{T \geq s \geq t} \alpha^\vee(\pi(s))]\alpha.$$

3.2. Littelmann path operators. Let $V, V^\vee, \alpha, \alpha^\vee$ be as above. Using proposition 3.2, as in [2], we can define generalized Littelmann path operators (see [25]).

DEFINITION 3.3. Let $\eta \in C_T^0(V)$, and $x \in \mathbb{R}$, then we define $\mathcal{E}_\alpha^x \eta$ as the unique path such that

$$\mathcal{P}_\alpha \mathcal{E}_\alpha^x \eta = \mathcal{P}_\alpha \eta \quad \text{and} \quad \mathcal{E}_\alpha^x \eta(T) = \eta(T) + x\alpha$$

if $-\alpha^\vee(\eta(T)) + \inf_{0 \leq t \leq T} \alpha^\vee(\eta(t)) \leq x \leq -\inf_{0 \leq t \leq T} \alpha^\vee(\eta(t))$ and $\mathcal{E}_\alpha^x \eta = \mathbf{0}$ otherwise. The following formula holds

$$\mathcal{E}_\alpha^x \eta(t) = \eta(t) - \min(-x, \inf_{t \leq s \leq T} \alpha^\vee(\eta(s)) - \inf_{0 \leq s \leq T} \alpha^\vee(\eta(s)))\alpha$$

if $-\alpha^\vee(T) + \inf_{0 \leq t \leq T} \alpha^\vee(\eta(t)) \leq x \leq 0$, and

$$\mathcal{E}_\alpha^x \eta(t) = \eta(t) - \min(0, -x - \inf_{0 \leq s \leq T} \alpha^\vee(\eta(s)) + \inf_{0 \leq s \leq t} \alpha^\vee(\eta(s)))\alpha$$

if $0 \leq x \leq -\inf_{0 \leq t \leq T} \alpha^\vee(\eta(t))$.

Here, as in the definition of crystals, $\mathbf{0}$ is a ghost element. The following result is immediate from the definition of the Littelmann operators.

PROPOSITION 3.4. $\mathcal{E}_\alpha^0 \eta = \eta$ and $\mathcal{E}_\alpha^x \mathcal{E}_\alpha^y \eta = \mathcal{E}_\alpha^{x+y} \eta$ as long as $\mathcal{E}_\alpha^y \eta \neq \mathbf{0}$.

We shall also use the notation $\mathcal{F}_\alpha^x = \mathcal{E}_\alpha^{-x}$ for $x \geq 0$, and denote by \mathcal{H}_α^x the restriction of the operator \mathcal{F}_α^x to α -dominant paths. Let π be an α -dominant path in $C_T^0(V)$ and $0 \leq x \leq \alpha^\vee(T)$, then $\mathcal{H}_\alpha^x \pi$ is the unique path in $C_T^0(V)$ such that

$$\mathcal{P}_\alpha \mathcal{H}_\alpha^x \pi = \pi$$

and

$$\mathcal{H}_\alpha^x \pi(T) = \pi(T) - x\alpha.$$

Observe that in this equality

$$x = -\inf_{0 \leq t \leq T} \alpha^\vee(\mathcal{H}_\alpha^x \pi(t)).$$

3.3. Product of Pitman transforms. Let $\alpha, \beta \in V$ and $\alpha^\vee, \beta^\vee \in V^\vee$ be such that $\alpha^\vee(\beta) < 0$ and $\beta^\vee(\alpha) < 0$. Replacing if necessary $(\alpha, \alpha^\vee, \beta, \beta^\vee)$ by $(t\alpha, \alpha^\vee/t, \beta/t, t\beta^\vee)$, which does not change \mathcal{P}_α and \mathcal{P}_β , we will assume that $\alpha^\vee(\beta) = \beta^\vee(\alpha)$. We use the notations

$$\rho = -\frac{1}{2}\alpha^\vee(\beta) = -\frac{1}{2}\beta^\vee(\alpha), \quad X(s) = \alpha^\vee(\pi(s)), \quad Y(s) = \beta^\vee(\pi(s)).$$

The following result is proved in [2].

THEOREM 3.5. Let n be a positive integer, then if $\rho \geq \cos \frac{\pi}{n}$,

$$\begin{aligned} \underbrace{(\mathcal{P}_\alpha \mathcal{P}_\beta \mathcal{P}_\alpha \dots)}_{n \text{ terms}} \pi(t) &= \pi(t) - \inf_{t \geq s_0 \geq s_1 \geq \dots \geq s_{n-1} \geq 0} \left(\sum_{i=0}^{n-1} T_i(\rho) Z^{(i)}(s_i) \right) \alpha \\ &\quad - \inf_{t \geq s_0 \geq s_1 \geq \dots \geq s_{n-2} \geq 0} \left(\sum_{i=0}^{n-2} T_i(\rho) Z^{(i+1)}(s_i) \right) \beta \end{aligned} \tag{3.1}$$

where $Z^{(k)} = X$ if k is even and $Z^{(k)} = Y$ if k is odd. The $T_k(x)$ are the Tchebycheff polynomials defined by

$$(3.2) \quad T_0(x) = 1, T_1(x) = 2x, 2xT_k(x) = T_{k-1}(x) + T_{k+1}(x) \text{ for } k \geq 1.$$

The Tchebycheff polynomials satisfy $T_k(\cos \theta) = \frac{\sin(k+1)\theta}{\sin \theta}$ and, in particular, under the assumptions on ρ and n , $T_k(\rho) \geq 0$ for all $k \leq n-1$. An important property of the Pitman transforms is the following corollary (see [2]).

THEOREM 3.6. (*Generalized braid relations for the Pitman transforms.*) Let $\alpha, \beta \in V$ and $\alpha^\vee, \beta^\vee \in V^\vee$ be such that $\alpha^\vee(\alpha) = \beta^\vee(\beta) = 2$, and $\alpha^\vee(\beta) < 0, \beta^\vee(\alpha) < 0$ and $\alpha^\vee(\beta)\beta^\vee(\alpha) = 4 \cos^2 \frac{\pi}{n}$, where $n \geq 2$ is some integer. Then

$$\mathcal{P}_\alpha \mathcal{P}_\beta \mathcal{P}_\alpha \dots = \mathcal{P}_\beta \mathcal{P}_\alpha \mathcal{P}_\beta \dots$$

where there are n factors in each product.

3.4. Pitman transforms for Coxeter groups. Let (W, S) be a Coxeter system, with a realization in the space V . For a simple reflection s , denote by \mathcal{P}_{α_s} or \mathcal{P}_s the Pitman transform associated with the pair $(\alpha_s, \alpha_s^\vee)$. From theorem 3.6 and Matsumoto's lemma [[4], Ch. IV, No. 1.5. Prop.5] we deduce:

THEOREM 3.7. Let $w = s_1 \dots s_r$ be a reduced decomposition of $w \in W$, with $s_1, \dots, s_r \in S$. Then

$$\mathcal{P}_w := \mathcal{P}_{s_1} \dots \mathcal{P}_{s_r}$$

depends only on w and not on the chosen decomposition.

When W is finite, it has a unique longest element, denoted by w_0 . The transformation \mathcal{P}_{w_0} plays a fundamental role in the sequel. The following result is proved in [2].

PROPOSITION 3.8. For any path $\eta \in C_T^0(V)$, the path $\mathcal{P}_{w_0}\eta$ takes values in the closed Weyl chamber \overline{C} . Furthermore \mathcal{P}_{w_0} is an idempotent and $\mathcal{P}_w \mathcal{P}_{w_0} = \mathcal{P}_{w_0} \mathcal{P}_w = \mathcal{P}_{w_0}$ for all $w \in W$.

3.5. Braid relations for the \mathcal{H} operators. Let $w \in W$ and fix a reduced decomposition $w = s_1 \dots s_p$. For any path η in $C_T^0(V)$, denote $\eta_p = \eta$ and for $k = 2, \dots, p$,

$$\eta_{k-1} = \mathcal{P}_{s_k} \dots \mathcal{P}_{s_p} \eta.$$

Then $\eta_{k-1} = \mathcal{P}_{s_k} \eta_k$ is α_{s_k} -dominant, by proposition 3.2 (ii) and

$$\eta_k = \mathcal{F}_{s_k}^{x_k} \eta_{k-1} = \mathcal{H}_{s_k}^{x_k} \eta_{k-1}$$

where

$$(3.3) \quad x_k = - \inf_{0 \leq t \leq T} \alpha_{s_k}^\vee(\eta_k(t)).$$

Observe that

$$(3.4) \quad x_k \in [0, \alpha_{s_k}^\vee(\eta_{k-1}(T))]$$

and

$$\eta_{k-1}(T) = \eta_k(T) - x_k \alpha_{s_k};$$

thus

$$\eta_{k-1}(T) = \pi(T) - \sum_{i=k}^p x_i \alpha_{s_i}.$$

Furthermore,

$$(3.5) \quad \eta_k = \mathcal{H}_{s_k}^{x_k} \mathcal{H}_{s_{k-1}}^{x_{k-1}} \cdots \mathcal{H}_{s_1}^{x_1} \mathcal{P}_w \eta,$$

and the numbers (x_1, \dots, x_k) are uniquely determined by this equation.

Let $w \in W$ be an element of the Coxeter group. We consider two reduced decompositions

$$w = s_1 \cdots s_p, w = s'_1 \cdots s'_p$$

of w . Let $\mathbf{i} = (s_1, \dots, s_p)$ and $\mathbf{j} = (s'_1, \dots, s'_p)$.

Let $\eta : [0, T] \rightarrow V$ be a continuous path such that $\eta(0) = 0$, and let (x_1, \dots, x_p) , respectively (y_1, \dots, y_p) , be the numbers determined by equation (3.5) for the two decompositions \mathbf{i} and \mathbf{j} . The following theorem states that the correspondance between the x_n 's and the y_n 's actually does not depend on the path η . In other words, we have the following braid relation for the operators \mathcal{H} .

$$(3.6) \quad \mathcal{H}_{s_p}^{x_p} \cdots \mathcal{H}_{s_2}^{x_2} \mathcal{H}_{s_1}^{x_1} = \mathcal{H}_{s'_p}^{y_p} \cdots \mathcal{H}_{s'_2}^{y_2} \mathcal{H}_{s'_1}^{y_1}.$$

THEOREM 3.9. *There exists a piecewise linear continuous map $\phi_{\mathbf{i}}^{\mathbf{j}} : \mathbb{R}^p \rightarrow \mathbb{R}^p$ such that for all paths $\eta \in C_T^0(V)$,*

$$(y_1, \dots, y_p) = \phi_{\mathbf{i}}^{\mathbf{j}}(x_1, \dots, x_p).$$

PROOF. First step: If the roots α, β generate a system of type $A_1 \times A_1$ and $w = s_\alpha s_\beta = s_\beta s_\alpha$, then \mathcal{P}_α and \mathcal{P}_β commute, and it is immediate that $x_1 = y_2$, $x_2 = y_1$. Let α, α^\vee and β, β^\vee be such that

$$\alpha^\vee(\alpha) = \beta^\vee(\beta) = 2, \quad \alpha^\vee(\beta) = \beta^\vee(\alpha) = -1,$$

then α and β generate a root system of type A_2 and the braid relation is

$$w_0 = s_\alpha s_\beta s_\alpha = s_\beta s_\alpha s_\beta.$$

We prove that the following map

$$(3.7) \quad \begin{aligned} x_1 &= (y_2 - y_1) \wedge y_3 & y_1 &= (x_2 - x_1) \wedge x_3 \\ x_2 &= y_1 + y_3 & y_2 &= x_1 + x_3 \\ x_3 &= y_1 \vee (y_2 - y_3) & y_3 &= x_1 \vee (x_2 - x_3) \end{aligned}$$

satisfies the required properties. Assume that, for $\pi = \mathcal{P}_{w_0} \eta$,

$$\eta = \mathcal{H}_\alpha^{x_3} \mathcal{H}_\beta^{x_2} \mathcal{H}_\alpha^{x_1} \pi.$$

Then define $\eta_2 = \mathcal{P}_\alpha \eta$, $\eta_1 = \mathcal{P}_\beta \mathcal{P}_\alpha \eta$, $\eta_0 = \pi = \mathcal{P}_\alpha \mathcal{P}_\beta \mathcal{P}_\alpha \eta$. Using theorem 3.5 for computing the paths η_i one gets the explicit formulas.

$$\begin{aligned} x_3 &= -\inf_{0 \leq s \leq T} \alpha^\vee(\eta(s)) \\ x_2 &= -\inf_{0 \leq s_2 \leq s_1 \leq T} (\beta^\vee(\eta(s_1)) + \alpha^\vee(\eta(s_2))) \\ x_1 &= -\inf_{0 \leq s_2 \leq s_1 \leq T} (\alpha^\vee(\eta(s_1)) + \beta^\vee(\eta(s_2))) - x_3. \end{aligned}$$

Similar formulas are obtained for the y_i coming from the other reduced decomposition, by exchanging the roles of α and β . The formula (3.7) follows by inspection.

In the context of crystals, this result is well known and first appeared in Lusztig [27] and Kashiwara [20]. We observe that it can also be obtained from the considerations of section 6.6, see especially 6.7.

Second step: When the roots generate a root system of type A_n , using Matsumoto's lemma, one can pass from one reduced decomposition to another by a sequence of braid relations corresponding to the two cases of the first step.

Third step: We consider now the case where the roots generate the dihedral group $I(m)$, and $w = s_\alpha s_\beta \dots = s_\beta s_\alpha \dots$ is the longest element in W . We will use an embedding of the dihedral group $I(m)$ in the Weyl group of the system A_{m-1} , see e.g. Bourbaki [4], ch. V, 6, Lemme 2. Recall the Tchebicheff polynomials T_k defined in (3.2). Let $\lambda = \cos(2\pi/m)$, $a_1 = a_2 = 1$ and, for $k \geq 1$,

$$a_{2k} = T_{k-1}(\lambda), \quad a_{2k+1} = T_k(\lambda) + T_{k-1}(\lambda)$$

then,

$$(3.8) \quad a_{2k} + a_{2k+2} = a_{2k+1}, \quad a_{2k+1}a_{2k-1} + a_{2k+1} = (1 + a_3)a_{2k},$$

Moreover $a_k > 0$ when $k < m$ and $a_m = 0$.

In the Euclidean space $V = \mathbb{R}^{m-1}$ we choose simple roots $\alpha_1, \dots, \alpha_{m-1}$ which satisfy $\langle \alpha_i, \alpha_j \rangle = a_{ij}$ where $a_{ij} = 2$ if $i = j$, $a_{ij} = -1$ if $|i - j| = 1$, $a_{ij} = 0$ otherwise. Let $\alpha_i^\vee = \alpha_i$ and $s_i = s_{\alpha_i}$. These generate a root system of type A_{m-1} .

Let Π be the two dimensional plane defined as the set of $x \in V$ such that for all $n < m$,

$$\langle \alpha_n, x \rangle = a_n \langle \alpha_1, x \rangle$$

if n is odd, and

$$\langle \alpha_n, x \rangle = a_n \langle \alpha_2, x \rangle$$

if n is even. It follows from the relation (3.8) that the vectors

$$\alpha = \sum_{n \text{ odd}, n < m} a_n \alpha_n, \quad \beta = \sum_{n \text{ even}, n < m} a_n \alpha_n$$

are in Π . Let $\alpha^\vee = 2\alpha/||\alpha||^2$, $\beta^\vee = 2\beta/||\beta||^2$ and

$$\tau_1 = s_1 s_3 s_5 \dots s_{2p-1},$$

$$\tau_2 = s_2 s_4 s_6 \dots s_{2r},$$

where $2p = m - 1$, $r = p$ when m is odd and $2p = m$, $r = p - 1$ when m is even. Let w_0 be the longest element in the Weyl group of A_{m-1} . Its length is $q = (m - 1)m/2$. We first consider the case where m is odd, $m = 2p + 1$, $q = pm$. Then

$$w_0 = (\tau_1 \tau_2)^p \tau_1, \quad \text{and} \quad w_0 = \tau_2 (\tau_1 \tau_2)^p$$

are two reduced decompositions of w_0 . Since $(\tau_1 \tau_2)^m = Id$ the angle between α and $-\beta$ is π/m and these vectors are the simple roots of the dihedral system $I(m)$.

Let γ be a continuous path in Π , let $\gamma_p = \gamma$ and for $1 < k \leq p$, $\gamma_{k-1} = \mathcal{P}_{\alpha_{2k-1}} \gamma_k$ and

$$z_k(t) = - \inf_{0 \leq s \leq t} \alpha_{2k-1}^\vee(\gamma_k(s)).$$

LEMMA 3.10. *Let γ be a continuous path with values in Π and let*

$$x(t) = - \inf_{0 \leq s \leq t} \alpha^\vee(\gamma(s)).$$

Then, for all k , $z_k(t) = a_{2k-1}x(t)$ and

$$\mathcal{P}_{\tau_1} \gamma(t) = \mathcal{P}_{\alpha_1} \mathcal{P}_{\alpha_3} \mathcal{P}_{\alpha_5} \dots \mathcal{P}_{\alpha_{2p-1}} \gamma(t) = \gamma(t) - \inf_{s \leq t} \alpha^\vee(\gamma(s))\alpha = \mathcal{P}_\alpha \gamma(t).$$

PROOF. First, notice that $\alpha^\vee(\gamma(t)) = \alpha_1^\vee(\gamma(t))$. Since γ is in Π , one has

$$z_p(t) = - \inf_{0 \leq s \leq t} \alpha_{2p-1}^\vee(\gamma(s)) = - \inf_{0 \leq s \leq t} a_{2p-1} \alpha_1^\vee(\gamma(s)) = a_{2p-1} x(t)$$

where we use the positivity of a_{2p-1} . Therefore

$$\gamma_{p-1}(t) = \mathcal{P}_{\alpha_{2p-1}} \gamma(t) = \gamma(t) + z_p(t) \alpha_{2p-1} = \gamma(t) + a_{2p-1} x(t) \alpha_{2p-1}.$$

Now, since the α_{2i+1} are orthogonal,

$$z_{p-1}(t) = - \inf_{0 \leq s \leq t} \alpha_{2p-3}^\vee(\gamma_{p-1}(s)) = - \inf_{0 \leq s \leq t} \alpha_{2p-3}^\vee(\gamma(s)) = a_{2p-3} x(t),$$

and

$$\begin{aligned} \gamma_{p-2}(t) &= \mathcal{P}_{\alpha_{2p-3}} \gamma_{p-1}(t) = \gamma_{p-1}(t) + z_{p-1}(t) \alpha_{2p-3} \\ &= \gamma(t) + x(t) (a_{2p-3} \alpha_{2p-3} + a_{2p-1} \alpha_{2p-1}). \end{aligned}$$

Continuing, we obtain that

$$\begin{aligned} z_k(t) &= a_{2k-1} x(t) \\ \gamma_k(t) &= \gamma(t) + x(t) (a_{2k-1} \alpha_{2k-1} + \cdots + a_{2p-1} \alpha_{2p-1}) \end{aligned}$$

Since $\alpha = \alpha_1 + a_3 \alpha_3 + a_5 \alpha_5 + \cdots + a_{2p-1} \alpha_{2p-1}$ we obtain the lemma. \square

We have similarly, if γ is a path in Π ,

$$\mathcal{P}_{\tau_2} \gamma(t) = \mathcal{P}_{\alpha_2} \mathcal{P}_{\alpha_4} \mathcal{P}_{\alpha_6} \cdots \mathcal{P}_{\alpha_{2r}} \gamma(t) = \gamma(t) - \inf_{s \leq t} \beta^\vee(\gamma(s)) \beta = \mathcal{P}_\beta \gamma(t).$$

Let $\mathbf{i} = (s_{i_1}, \dots, s_{i_q}) = (\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_m)$ and $\mathbf{j} = (s_{j_1}, \dots, s_{j_q}) = (\mathbf{j}_1, \mathbf{j}_2, \dots, \mathbf{j}_m)$ where $\mathbf{i}_k = \mathbf{j}_{k+1} = (s_1, s_3, \dots, s_{2p-1})$ when k is odd and $\mathbf{i}_k = \mathbf{j}_{k+1} = (s_2, s_4, \dots, s_{2p})$ when k is even. We write explicitly

$$w_0 = (\tau_1 \tau_2)^p \tau_1 = s_{i_1} \cdots s_{i_q}, w_0 = \tau_2 (\tau_1 \tau_2)^p = s_{j_1} \cdots s_{j_q}.$$

Let us denote by $\phi_1^{\mathbf{j}} : \mathbb{R}^q \rightarrow \mathbb{R}^q$ the mapping given by the second step corresponding to these two reduced decompositions of w_0 in the Weyl group of A_{m-1} .

Let γ be a path with values in Π . If we consider it as a path in V we can set $\eta_q = \tilde{\eta}_q = \gamma$ and, for $n = 1, 2, \dots, q$,

$$\eta_{n-1} = \mathcal{P}_{\alpha_{i_n}} \eta_n, \quad z_n = - \inf_{0 \leq t \leq T} \alpha_{i_n}^\vee(\eta_n(t))$$

$$\tilde{\eta}_{n-1} = \mathcal{P}_{\alpha_{j_n}} \tilde{\eta}_n, \quad \tilde{z}_n = - \inf_{0 \leq t \leq T} \alpha_{j_n}^\vee(\tilde{\eta}_n(t)).$$

Then, by definition,

$$(\tilde{z}_1, \dots, \tilde{z}_q) = \phi_1^{\mathbf{j}}(z_1, \dots, z_q).$$

We now consider γ as a path in Π . We let

$$(u_1, u_2, \dots, u_m) = (\alpha, \beta, \alpha, \beta, \dots, \alpha)$$

and

$$(v_1, v_2, \dots, v_m) = (\beta, \alpha, \beta, \alpha, \dots, \beta).$$

In $I(m)$ the two reduced decompositions of the longest element are

$$s_{u_1} \cdots s_{u_m} = s_{v_1} \cdots s_{v_m}.$$

We introduce $\gamma_m = \tilde{\gamma}_m = \gamma$, and, for $n = 1, 2, \dots, m$,

$$\begin{aligned} \gamma_{n-1} &= \mathcal{P}_{u_n} \cdots \mathcal{P}_{u_m} \gamma_m, \tilde{\gamma}_{n-1} = \mathcal{P}_{v_n} \cdots \mathcal{P}_{v_m} \tilde{\gamma}_m \\ x_n &= - \inf_{0 \leq t \leq T} u_n^\vee(\gamma_n(t)), \tilde{x}_n = - \inf_{0 \leq t \leq T} v_n^\vee(\tilde{\gamma}_n(t)). \end{aligned}$$

It follows from lemma 3.10 and from its analogue with α replaced by β that

$$z_1 = a_1 x_1, z_2 = a_3 x_1, \dots, z_p = a_{2p-1} x_1$$

$$z_{p+1} = a_2 x_2, z_{p+2} = a_4 x_2, \dots, z_{2p} = a_{2p} x_2$$

and more generally, for $k = 0, \dots$

$$a_1^{-1} z_{2kp+1} = a_3^{-1} z_{2kp+2} = \dots = a_{2p-1}^{-1} z_{2kp+p} = x_{k+1}$$

$$a_2^{-1} z_{(2k+1)p+1} = a_4^{-1} z_{(2k+1)p+2} = \dots = a_{2p}^{-1} z_{(2k+2)p} = x_{k+2}.$$

This defines a linear map

$$(x_1, \dots, x_m) = g(z_1, z_2, \dots, z_q).$$

Analogously exchanging the role of α and β we define a similar map

$$(\tilde{x}_1, \dots, \tilde{x}_m) = \tilde{g}(\tilde{z}_1, \tilde{z}_2, \dots, \tilde{z}_q)$$

(for instance $\tilde{z}_1 = a_2 \tilde{x}_1, \tilde{z}_2 = a_3 \tilde{x}_1, \dots$). Then we see that

$$(x_1, \dots, x_m) = \phi(\tilde{x}_1, \dots, \tilde{x}_m)$$

where

$$\phi = \tilde{g} \circ \phi_1^j \circ g^{-1}.$$

The proof when m is even is similar (when $m = 2p$, $w_0 = (\tau_1 \tau_2)^p$ and $w_0 = (\tau_2 \tau_1)^p$ are two reduced decompositions of w_0). This proves the theorem in the dihedral case.

Fourth step. We use Matsumoto's lemma to reduce the general case to the dihedral case.

This ends the proof of theorem 3.9. \square

REMARK 3.11. Although the given proof is constructive, it gives a complicated expression for ϕ_1^j which can sometimes be simplified. In the dihedral case $I(m)$, for the Weyl group case, i.e. $m = 3, 4, 6$, these expressions are given in Littelman [26]. For $m = 5$ it can be shown by a tedious verification that it is given when α, β have the same length, by a similar formula. Thus for $m = 2, 3, 4, 5, 6$ let $c_0 = 1, c_1 = 2 \cos(\pi/m), c_{n+1} + c_{n-1} = c_1 c_n$ for $n \geq 1$, and

$$u = \max(c_k x_{k+1} - c_{k-1} x_{k+2}, 0 \leq k \leq m-3), \quad v = \min(c_k x_{k+2} - c_{k+1} x_{k+1}, 1 \leq k \leq m)$$

Then the expressions are given by

$$\begin{aligned} y_m &= \max(x_{m-1} - c_1 x_m, u) \\ y_{m-1} &= x_m + \max(x_{m-2} - c_2 x_m, c_1 u) \\ y_2 &= x_1 + \min(x_3 - c_2 x_1, c_1 v) \\ y_1 &= \min(x_2 - c_1 x_1, v) \end{aligned}$$

and

$$\begin{aligned} y_1 + y_3 + \dots &= x_2 + x_4 + \dots \\ y_2 + y_4 + \dots &= x_1 + x_3 + \dots \end{aligned}$$

This determines completely (y_1, \dots, y_m) as a function of (x_1, \dots, x_m) when $m \leq 6$. For $m = 7$ we think (and made a computer check) that we have to add that

$$\begin{aligned} y_7 + y_5 &= x_6 + \max(c_2 x_1, x_4 - c_3 x_7, w) \\ w &= \min(c_2 u, x_4 - c_2 v, \max(x_6 - c_1 x_5 + x_4 + c_2 u, c_1 x_3 - x_2 - c_2 v)). \end{aligned}$$

We do not know of similar formulas for $m \geq 8$.

REMARK 3.12. The map given by theorem 3.9 is unique on the set of all possible coordinates of paths. We will see in the next section that this set is a convex cone. Since the value of the map $\phi_{\mathbf{i}}^{\mathbf{j}}$ is irrelevant outside this cone, we will assume that there exists a unique such map for each pair of reduced decompositions \mathbf{i}, \mathbf{j} .

4. Continuous Littelmann module

4.1. String parametrization of $C_T^0(V)$. Let (W, S, V, V^\vee) be a realization of the Coxeter system (W, S) . We say that a path is dominant if it takes its values in the closed Weyl chamber \overline{C} . Thus a path is dominant if and only if it is α -dominant for all simple roots α .

DEFINITION 4.1. Let $\pi \in C_T^0(V)$ be a dominant path, and $w \in W$. We define

$$L_\pi^w = \{\eta \in C_T^0(V); \mathcal{P}_w \eta = \pi\}.$$

These sets are defined for arbitrary Coxeter groups. We shall establish their main properties in the case of finite Coxeter groups, where they are analogues of Demazure-Littelmann modules. It remains an interesting problem to establish similar properties in the general case.

From now on we assume that W is finite, with longest element w_0 , and we denote $L_\pi = L_\pi^{w_0}$, which we call the Littelmann module associated with π . We will see that $L_\pi \cup \{\mathbf{0}\}$ is invariant under the Littelmann operators.

For notational convenience, we sometimes write $\alpha^\vee \eta$ instead of $\alpha^\vee(\eta)$.

Let $\eta \in L_\pi$, where π is dominant and $w_0 = s_1 \dots s_q$ be a reduced decomposition, then we have seen that

$$\eta = \mathcal{H}_{s_q}^{x_q} \mathcal{H}_{s_{q-1}}^{x_{q-1}} \dots \mathcal{H}_{s_1}^{x_1} \pi$$

for a unique sequence

$$\varrho_{\mathbf{i}}(\eta) = (x_1, \dots, x_q).$$

Following Berenstein and Zelevinsky [3], we call $\varrho_{\mathbf{i}}(\eta)$ the \mathbf{i} -string parametrization of η , or the string parametrization if no confusion is possible.

We let

$$C_{\mathbf{i}}^\pi = \varrho_{\mathbf{i}}(L_\pi),$$

this is the set of all the $(x_1, \dots, x_q) \in \mathbb{R}^q$ which occur in the string parametrizations of the elements of L_π .

PROPOSITION 4.2. *The set L_π is compact and the map $\varrho_{\mathbf{i}}$ is a bicontinuous bijection from L_π onto its image $C_{\mathbf{i}}^\pi$.*

PROOF. The map $\varrho_{\mathbf{i}}$ has an inverse

$$\varrho_{\mathbf{i}}^{-1}(x_1, \dots, x_q) = \mathcal{H}_{s_q}^{x_q} \mathcal{H}_{s_{q-1}}^{x_{q-1}} \dots \mathcal{H}_{s_1}^{x_1} \pi,$$

hence it is bijective. It is clear that $\varrho_{\mathbf{i}}$ and $\varrho_{\mathbf{i}}^{-1}$ are continuous. Since \mathcal{P}_{w_0} is continuous, $L_\pi = \{\eta; \mathcal{P}_{w_0}(\eta) = \pi\}$ is closed. Using $\varrho_{\mathbf{i}}^{-1}$ we easily see that L_π is equicontinuous, it is thus compact by Ascoli's theorem. \square

We will study $C_{\mathbf{i}}^\pi$ in detail in the following sections.

4.2. The crystallographic case. In this section we consider the case of a Weyl group W with a crystallographic root system. When α is a root and α^\vee its coroot, then \mathcal{E}_α^1 and \mathcal{E}_α^{-1} from definition 3.3 coincide with the Littelmann operators e_α and f_α , defined in [25]. Recall that a path η is called integral in [25] if its endpoint $\eta(T)$ is in the weight lattice and if, for each simple root α , the minimum of the function $\alpha^\vee(\eta(t))$ over $[0, T]$ is an integer. The class of integral paths is invariant under the Littelmann operators.

Let π be a dominant integral path. The discrete Littelmann module B_π is defined as the orbit of π under the semigroup generated by all the transformations e_α, f_α , for all simple roots α .

Let $\mathbf{i} = (s_1, \dots, s_q)$ where $w_0 = s_1 \cdots s_q$ is a reduced decomposition, then it follows from Littelmann's theory that

$$B_\pi = \{\eta \in L_\pi; x_1, \dots, x_q \in \mathbb{N}\} = \varrho_{\mathbf{i}}^{-1}(\{(x_1, \dots, x_q) \in C_{\mathbf{i}}^\pi; x_1 \in \mathbb{N}, \dots, x_q \in \mathbb{N}\}).$$

Furthermore, the set B_π has a crystal structure isomorphic to the Kashiwara crystal associated with the highest weight $\pi(T)$. On B_π the coordinates (x_1, \dots, x_q) are called the string parametrization of the dual canonical basis by Berenstein and Zelevinsky. They are described in Littelmann [26] and Berenstein and Zelevinsky [3].

When restricted to B_π , the Pitman operator \mathcal{P}_α coincides with e_α^{max} , i.e. the operator sending η to $e_\alpha^n \eta$, where $n = \max(k, e_\alpha^k \eta \neq \mathbf{0})$.

For any path $\eta : [0, T] \rightarrow V$ and $\lambda > 0$ let $\lambda\eta$ be the path defined by $(\lambda\eta)(t) = \lambda\eta(t)$ for $0 \leq t \leq T$. The following results are immediate.

PROPOSITION 4.3 (Scaling property). *For any $\lambda > 0$, $\lambda L_\pi = L_{\lambda\pi}$.*

PROPOSITION 4.4. *If π is a dominant integral path, then the set*

$$B_\pi(\mathbb{Q}) = \cup_{n \in \mathbb{N}} \frac{1}{n} B_{n\pi}$$

is dense in L_π .

PROPOSITION 4.5. *Let $\eta \in C_T^0(V)$, $r \in \mathbb{R}, u > 0$, then*

$$\mathcal{E}_\alpha^{ru}(u\eta) = u\mathcal{E}_\alpha^r(\eta).$$

PROPOSITION 4.6. *Let π be a dominant path and $a > 0$ then $C_{\mathbf{i}}^{a\pi} = aC_{\mathbf{i}}^\pi$ and, for each $\eta \in L_\pi$, $\varrho_{a\pi}(a\eta) = a\varrho_\pi(\eta)$.*

Actually a good interpretation of L_π in the Weyl group case is as the "limit" of $\frac{1}{n}B_{n\pi}$ when $n \rightarrow \infty$. In the general Coxeter case only the limiting object is defined.

4.3. Polyhedral nature of the crystal for a Weyl group. Let W be a finite Weyl group, associated to a crystallographic root system. Let B_π be the discrete Littelmann module associated with an integral dominant path π . Recall that it is the image of π under all the products of the Littelmann operators e_α, f_α , for α simple roots. We fix a reduced decomposition $w_0 = s_1 \cdots s_q$ of the longest element and let $\mathbf{i} = (s_1, \dots, s_q)$. We have seen that if $\rho_{\mathbf{i}} : L_\pi \rightarrow C_{\mathbf{i}}^\pi$ is the string parametrization of the continuous module L_π , then

$$B_\pi = \{\eta \in L_\pi; x_1, \dots, x_q \in \mathbb{N}\} = \varrho_{\mathbf{i}}^{-1}(\{(x_1, \dots, x_q) \in C_{\mathbf{i}}^\pi; x_1 \in \mathbb{N}, \dots, x_q \in \mathbb{N}\}).$$

Therefore the set

$$\tilde{C}_{\mathbf{i}}^\pi = C_{\mathbf{i}}^\pi \cap \mathbb{N}^q$$

is the image of the discrete Littelmann module B_π , or equivalently, the image of the Kashiwara crystal with highest weight $\pi(T)$, under the string parametrization of Littelmann [26] and Berenstein and Zelevinsky [3]. Let

$$K_\pi = \{(x_1, \dots, x_q) \in \mathbb{R}^q; 0 \leq x_r \leq \alpha_{i_r}^\vee(\pi(T) - \sum_{n=1}^{r-1} x_n \alpha_{i_n}), r = 1, \dots, q\}.$$

It is shown in Littelmann [26] that there exists a convex rational polyhedral cone $C_{\mathbf{i}}$ in \mathbb{R}^q , depending only on \mathbf{i} such that, for all dominant integral paths π ,

$$\tilde{C}_{\mathbf{i}}^\pi = C_{\mathbf{i}} \cap \mathbb{N}^q \cap K_\pi.$$

This cone is described explicitly in Berenstein and Zelevinsky [3]. Recall that $C_{\mathbf{i}}^\pi = \varrho_{\mathbf{i}}(L_\pi)$. Using propositions 4.3, 4.6, 4.5 it is easy to see that the following holds.

PROPOSITION 4.7. *For all dominant paths π ,*

$$C_{\mathbf{i}}^\pi = C_{\mathbf{i}} \cap K_\pi.$$

4.4. The cone in the general case. We now consider a general Coxeter system (W, S) , with W finite, realized in V .

THEOREM 4.8. *Let \mathbf{i} be a reduced decomposition of w_0 , then there exists a unique polyhedral cone $C_{\mathbf{i}}$ in \mathbb{R}^q such that for any dominant path π*

$$C_{\mathbf{i}}^\pi = C_{\mathbf{i}} \cap K_\pi.$$

In particular $C_{\mathbf{i}}^\pi$ depends only on $\lambda = \pi(T)$.

PROOF. It remains to consider the non crystallographic Coxeter systems. It is clearly enough to consider reduced systems. We use their classification: W is either a dihedral group $I(m)$ or H_3 or H_4 (see Humphreys [17]), and the same trick as the one used in the proof of theorem 3.9.

We first consider the case $I(m)$ where $m = 2p + 1$ and we use the notation of the proof of theorem 3.9. Let $\mathbf{i} = (i_1, \dots, i_q)$ be as in that proof, and write

$$w_0 = (\tau_1 \tau_2)^p \tau_1 = s_{i_1} \cdots s_{i_q}$$

for the longest word in A_{m-1} . Let γ be a path with values in the plane Π . If we consider γ as a path in $V = \mathbb{R}^{m-1}$ we can set, for $q = (m-1)m/2$,

$$\eta_q = \gamma$$

and, for $n = 1, 2, \dots, q$

$$\eta_{n-1} = \mathcal{P}_{\alpha_{i_n}} \eta_n, \quad z_n = - \inf_{0 \leq t \leq T} \alpha_{i_n}^\vee(\eta_n(t)).$$

We can also consider γ as a path in Π , with the realization of $I(m)$. Let

$$\mathbf{u} = (u_1, u_2, \dots, u_m) = (\alpha, \beta, \alpha, \beta, \dots, \alpha).$$

Let $\tilde{\eta}_m = \gamma$ and, for $n = 1, 2, \dots, m$,

$$\tilde{\eta}_{n-1} = \mathcal{P}_{u_n} \dots \mathcal{P}_{u_m} \tilde{\eta}_m, \quad x_n = - \inf_{0 \leq t \leq T} u_n^\vee(\tilde{\eta}_m(t)).$$

We have seen that the map


$$(x_1, \dots, x_m) = g(z_1, z_2, \dots, z_q),$$

is linear. Let $C_{\mathbf{i}}$ be the cone associated with \mathbf{i} in A_{m-1} , then $C_{\mathbf{u}} = g(C_{\mathbf{i}})$ is the cone in \mathbb{R}^m associated with the reduced decomposition $\alpha\beta\cdots\alpha$ of the longest word in $I(m)$. Furthermore, for any dominant path π in Π , $C_{\mathbf{u}}^\pi = C_{\mathbf{u}} \cap K_\pi$.

The proof when m is even is similar.

In order to deal with the cases H_3 and H_4 it is enough, using an analogous proof to embed these systems in some Weyl groups.

Let us first consider the case of H_4 . We use the embedding of H_4 in E_8 (see [29]). Consider the following indexation of the simple roots of the system E_8 :



In the euclidean space $V = \mathbb{R}^8$ the roots $\alpha_1, \dots, \alpha_8$, satisfy $\langle \alpha_i, \alpha_j \rangle = -1$ or 0 depending whether they are linked or not. Let $\phi = (1 + \sqrt{5})/2$. We consider the 4-dimensional subspace Π of V defined as the set of $x \in V$ orthogonal to $\alpha_8 - \phi\alpha_1, \alpha_7 - \phi\alpha_2, \alpha_6 - \phi\alpha_3$ and $\phi\alpha_5 - \alpha_4$. Let s_i be the reflection which corresponds to α_i and

$$\tau_1 = s_1 s_8, \tau_2 = s_2 s_7, \tau_3 = s_3 s_6, \tau_4 = s_4 s_5.$$


One checks easily that $\tau_1, \tau_2, \tau_3, \tau_4$ generate H_4 and that the vectors

$$\tilde{\alpha}_1 = \alpha_1 + \phi\alpha_8, \tilde{\alpha}_2 = \alpha_2 + \phi\alpha_7, \tilde{\alpha}_3 = \alpha_3 + \phi\alpha_6, \tilde{\alpha}_4 = \alpha_4 + \phi^{-1}\alpha_5$$

are in Π . If π is a continuous path in Π , then, for $i = 1, \dots, 4$, if $\tilde{\alpha}_i^\vee = \tilde{\alpha}_i / (2\|\tilde{\alpha}_i\|^2)$,

$$\mathcal{P}_{\tau_i} \pi(t) = \pi(t) - \inf_{0 \leq s \leq t} \tilde{\alpha}_i^\vee(\pi(s)) \tilde{\alpha}_i.$$

The case of H_3 is similar by using D_6 :



In $V = \mathbb{R}^6$ we choose the roots $\alpha_1, \dots, \alpha_6$ with $\langle \alpha_i, \alpha_j \rangle = -1$ if they are linked. We define a 3-dimensional subspace Π defined as the set of $x \in V$ orthogonal to $\alpha_5 - \phi\alpha_1, \alpha_4 - \phi\alpha_2$ and $\phi\alpha_6 - \alpha_3$. Then the reflections

$$(4.1) \quad \tau_1 = s_1 s_5, \tau_2 = s_2 s_4, \tau_3 = s_3 s_6,$$

generate H_3 and

$$\tilde{\alpha}_1 = \alpha_1 + a\alpha_5, \tilde{\alpha}_2 = \alpha_2 + a\alpha_4, \tilde{\alpha}_3 = \alpha_3 + b\alpha_6$$

are in Π . \square

We will prove in corollary 5.3 that the cones C_i have the following description: for any simple root α , let $\mathbf{j}(\alpha)$ be a reduced decomposition of w_0 which begins by s_α . Then

$$C_i = \{x \in \mathbb{R}^q; \phi_i^{\mathbf{j}(\alpha)}(x)_1 \geq 0, \text{ for all simple roots } \alpha\}.$$

4.5. The cone in the dihedral case. In this section we provide explicit equations for the cone, in the dihedral case, following the approach of Littellmann [26] in the Weyl group case.

LEMMA 4.9. *Let $\alpha, \beta \in V$, $\alpha^\vee, \beta^\vee \in V^\vee$ and $c = -\beta^\vee(\alpha)$. Consider a continuous path $\eta \in C_T^0(V)$ and $\pi = \mathcal{P}_\alpha \eta$. Let*

$$\begin{aligned} U &= \min_{T \geq t \geq 0} [a\beta^\vee(\eta(t)) + b \min_{t \geq s \geq 0} \alpha^\vee(\eta(s))], \\ V &= \min_{T \geq t \geq 0} [a \min_{t \geq s \geq 0} \beta^\vee(\pi(s)) + (ac - b)\alpha^\vee(\pi(t))], \\ W &= a \min_{T \geq t \geq 0} \beta^\vee(\pi(t)) - (ac - b) \min_{T \geq t \geq 0} \alpha^\vee(\eta(t)), \end{aligned}$$

where a, b are real numbers such that $a \geq 0, ac - b \geq 0$. Then $U = \min(V, W)$.

PROOF. Since $\pi = \mathcal{P}_\alpha \eta$,

$$\beta^\vee(\eta(t)) = \beta^\vee(\pi(t)) - c \min_{t \geq s \geq 0} \alpha^\vee(\eta(s)),$$

thus

$$\begin{aligned} U &= \min_{T \geq t \geq 0} [a\beta^\vee(\pi(t)) + (b - ac) \min_{t \geq s \geq 0} \alpha^\vee(\eta(s))] \\ &= \min_{T \geq t \geq 0} [\min_{t \geq s \geq 0} a\beta^\vee(\pi(s)) + (b - ac) \min_{t \geq s \geq 0} \alpha^\vee(\eta(s))]. \end{aligned}$$

where we have used the fact that, if $f, g : [0, T] \rightarrow \mathbb{R}$ are two continuous functions, and if g is non decreasing, then

$$\min_{T \geq t \geq 0} [f(t) + g(t)] = \min_{T \geq t \geq 0} [\min_{t \geq s \geq 0} f(s) + g(t)].$$

Since $\alpha^\vee(\pi(t)) \geq -\min_{t \geq s \geq 0} \alpha^\vee(\eta(s))$,

$$\min_{t \geq s \geq 0} a\beta^\vee(\pi(s)) + (ac - b)\alpha^\vee(\pi(t)) \geq \min_{t \geq s \geq 0} a\beta^\vee(\pi(s)) - (ac - b) \min_{t \geq s \geq 0} \alpha^\vee(\eta(s)).$$

Let t_0 be the largest $t \leq T$ where the minimum of the right hand side is achieved. Suppose that $t_0 < T$. If $\alpha^\vee(\pi(t_0)) > -\min_{t_0 \geq s \geq 0} \alpha^\vee(\eta(s))$ then $\min_{t \geq s \geq 0} \alpha^\vee(\eta(s))$ is locally constant on the right of t_0 . Since $\min_{t \geq s \geq 0} a\beta^\vee(\pi(s))$ is non increasing, it follows that t_0 is not maximal. Therefore, when $t_0 < T$,

$$\alpha^\vee(\pi(t_0)) = -\min_{t_0 \geq s \geq 0} \alpha^\vee(\eta(s))$$

and

$$U = \min_{T \geq t \geq 0} [\min_{t \geq s \geq 0} a\beta^\vee(\pi(s)) - (ac - b) \inf_{t \geq s \geq 0} \alpha^\vee(\eta(s))] = V \leq W.$$

When $t_0 = T$, then $U = W \leq V$. Thus $U = \min(V, W)$. \square

We consider a realization of the dihedral system $I(m)$ with two simple roots α, β and $c := -\alpha^\vee(\beta) = -\beta^\vee(\alpha) = 2 \cos \frac{\pi}{m}$. Let

$$a_n = \frac{\sin(n\pi/m)}{\sin(\pi/m)}.$$

Then $a_0 = 0, a_1 = 1$, and $a_{n+1} + a_{n-1} = ca_n$, $a_n > 0$ if $1 \leq n \leq m-1$ and $a_m = 0$. Let $w_0 = s_1 \dots s_m$ be a reduced decomposition of the longest element $w_0 \in W$, $\mathbf{i} = (s_1, \dots, s_m)$ and $\alpha_1, \dots, \alpha_m$ be the simple roots associated with s_1, \dots, s_m . This sequence is either $(\alpha, \beta, \alpha, \dots)$ or $(\beta, \alpha, \beta, \dots)$. Clearly the two roots play a symmetric role, and the cones associated with these two decompositions are the same. We define α_0 as the simple root not equal to α_1 . As before, when $\eta \in C_T^0(V)$, we define $\eta_m = \eta$ and for $k = 0, \dots, m-1$, $\eta_k = \mathcal{P}_{s_{k+1}} \dots \mathcal{P}_{s_m} \eta$, and

$$x_k = - \min_{0 \leq t \leq T} \alpha_k^\vee(\eta_k(t)) \quad \text{for } k = 1, \dots, m.$$

PROPOSITION 4.10. *The cone for the dihedral system $I(m)$ is given by*

$$C_{\mathbf{i}} = \{(x_1, \dots, x_m) \in \mathbb{R}_+^m; \frac{x_{m-1}}{a_{m-1}} \geq \frac{x_{m-2}}{a_{m-2}} \geq \dots \geq \frac{x_1}{a_1}\}.$$

PROOF. For any p, k such that $0 \leq p \leq m, 0 \leq k \leq p$, let

$$\begin{aligned} V_k &= \min_{T \geq t \geq 0} [a_{k+1} \alpha_{p+1-k}^\vee(\eta_{p-k}(t)) + a_k \min_{t \geq s \geq 0} \alpha_{p-k}^\vee(\eta_{p-k}(s))], \\ W_k &= a_k \min_{T \geq t \geq 0} \alpha_{p-k}^\vee(\eta_{p-k}(t)) - a_{k+1} \min_{T \geq t \geq 0} \alpha_{p+1-k}^\vee(\eta_{p+1-k}(t)). \end{aligned}$$

Since $a_{k-1} + a_{k+1} = ca_k$, the lemma above gives that $V_k = \min(W_{k+1}, V_{k+1})$. Therefore

$$V_0 = \min(W_1, W_2, \dots, W_p, V_p).$$

Notice that

$$V_p = \min_{T \geq t \geq 0} [a_{p+1} \alpha_1^\vee(\eta_0(t)) + a_p \min_{t \geq s \geq 0} \alpha_0^\vee(\eta_0(s))] = 0$$

and

$$W_p = a_{p+1} x_1$$

since $\eta_0 = \mathcal{P}_{w_0} \eta$ is dominant. Furthermore

$$V_0 = \min_{0 \leq t \leq T} \alpha_{p+1}^\vee(\eta_p(t))$$

since $a_0 = 0$ and $a_1 = 1$. Hence,

$$(4.2) \quad \min_{0 \leq t \leq T} \alpha_{p+1}^\vee(\eta_p(t)) = \min(a_2 x_p - a_1 x_{p-1}, \dots, a_p x_2 - a_{p-1} x_1, a_{p+1} x_1, 0).$$

The path $\eta_{m-1} = \mathcal{P}_{\alpha_m} \eta$ is α_m -dominant, therefore $\alpha_m^\vee(\eta_{m-1}(t)) \geq 0$ and it follows from (4.2) applied with $p = m-1$ that for $k = 1, \dots, m-2$

$$a_{m-k} x_{k+1} - a_{m-k-1} x_k \geq 0,$$

which is equivalent, since $a_{m-k} = a_k$ to

$$\frac{x_{m-1}}{a_{m-1}} \geq \frac{x_{m-2}}{a_{m-2}} \geq \dots \geq \frac{x_1}{a_1} \geq 0.$$

Conversely, we suppose that these inequalities hold, i.e. that for $k = 1, \dots, m-2$

$$(4.3) \quad a_{k+1} x_{m-k} - a_k x_{m-k-1} \geq 0,$$

$$a_{m-k} x_{k+1} - a_{m-k-1} x_k \geq 0,$$

and that $(x_1, \dots, x_m) \in K_\pi$ for some dominant path π . Let us show that

$$\eta = \mathcal{H}_{\alpha_m}^{x_m} \dots \mathcal{H}_{\alpha_1}^{x_1} \pi$$

is well defined. Since the string parametrization of η is x this will prove the proposition. It is enough to show, by induction on $p = 0, \dots, m$ that

$$\eta_p := \mathcal{H}_{\alpha_p}^{x_p} \mathcal{H}_{\alpha_{p-1}}^{x_{p-1}} \dots \mathcal{F}_{\alpha_1}^{x_1} \pi$$

is α_{p+1} -dominant. This is clear for $p = 0$ since $\eta_0 = \pi$ is dominant. If we suppose that this is true until $p - 1$ can apply (4.2) and write that

$$\min_{0 \leq t \leq T} \alpha_{p+1}^\vee(\eta_p(t)) = \min(a_2 x_p - a_1 x_{p-1}, \dots, a_p x_2 - a_{p-1} x_1, a_{p+1} x_1, 0)$$

Since $c \leq 2$, it is easy to see that

$$\frac{a_{n-1}}{a_n} \geq \frac{a_{n-2}}{a_{n-1}}$$

for $n \leq m - 1$. Therefore,

$$\frac{x_{k+1}}{x_k} \geq \frac{a_{m-k-1}}{a_{m-k}} \geq \frac{a_{p-k}}{a_{p-k+1}}$$

and $\alpha_{p+1}^\vee(\eta_p(t)) \geq 0$ for all $0 \leq t \leq T$. \square

In the definition of V_k and W_k in the proof above, replace the sequence (a_k) by the sequence (a_{k+1}) . We obtain the following formula.

PROPOSITION 4.11. *If $y_m = -\min_{T \geq t \geq 0} \alpha_{m-1}^\vee(\eta_m(t))$, then*

$$y_m = \max\{0, a_{m-1}x_{m-1} - a_{m-2}x_m, a_{m-2}x_{m-2} - a_{m-3}x_{m-1}, \dots, a_2x_2 - a_1x_3, a_1x_1\}$$

4.6. Remark on Gelfand Tsetlin cones. In the Weyl group case, the continuous cone appears in the description of toric degeneration (see Caldero [5], Alexeev and Brion [1]). The polytopes $C_i(\lambda)$ are called the string polytopes in Alexeev and Brion [1]. Notice that they have shown that the classical Duistermaat Heckman measure coincides with the one given below in Definition 5.4. Explicit inequalities for the string cone C_i (and therefore for the string polytopes) in the Weyl group case are given in full generality in Berenstein and Zelevinsky in [3, Thm.3.12]. Before, Littelmann [26, Thm.4.2] has described it for the so called "nice decompositions" of w_0 . As explained in this paper they were introduced to generalize the Gelfand Tsetlin cones.

For the convenience of the reader let us reproduce the description in the A_n case, considered explicitly in Alexeev and Brion [1]. The standard reduced decomposition of the longest element in the symmetric group $W = S_{n+1}$ is

$$w_0 = (s_1)(s_2s_1)(s_3s_2s_1) \dots (s_ns_{n-1} \dots s_1),$$

where s_i denotes the transposition exchanging i with $i + 1$. Let us use the coordinates $x_{i,j}$ with $i, j \geq 1, i + j \leq n + 1$. The string cone is defined by

$$x_{n,1} \geq 0; \quad x_{n-1,2} \geq x_{n-1,1} \geq 0; \quad \dots \quad x_{1,n} \geq \dots \geq x_{1,1} \geq 0,$$

and to define the polyhedron C_i^λ one has to add the inequalities

$$x_{i,j} \leq \alpha_j^\vee(\lambda) - x_{i,j-1} + \sum_{k=1}^{i-1} (-x_{k,j-1} + 2x_{k,j} - x_{k,j+1}).$$

A more familiar description of this cone is in terms of Gelfand-Tsetlin patterns:

$$g_{i,j} \geq g_{i+1,j} \geq g_{i,j+1}$$

in variables $g_{i,j}$, $i, j \geq 1$, $i + j \leq n + 1$ and with variables $g_{0,i}$ corresponding to λ :

$$\lambda = \sum_{i=0}^{n+1} g_{0,i} \epsilon_i = \sum_{i=0}^{n+1} \lambda_i \epsilon_i, \quad \text{so that} \quad \alpha_j^\vee(\lambda) = \langle \lambda, \epsilon_j - \epsilon_{j+1} \rangle = \lambda_j - \lambda_{j+1}.$$

The linear change of coordinates is given by $g_{i,j} = \lambda_j + \sum_{k=1}^i (x_{k,j-1} - x_{k,j})$.

4.7. Crystal structure of the Littelmann module. We now return to the general case of a finite Coxeter group.

Let π be a dominant path in $C_T^0(V)$. For $\eta \in L_\pi$ we define $wt(\eta) = \eta(T)$. Let e_α^r be the generalized Littelmann operator \mathcal{E}_α^r defined in Definition 3.3, and

$$\varepsilon_\alpha(\eta) = \max\{r \geq 0; \mathcal{E}_\alpha^r(\eta) \neq 0\} = - \inf_{0 \leq t \leq T} \alpha^\vee(\eta(t))$$

$$\varphi_\alpha(\eta) = \max\{r \geq 0; \mathcal{E}_\alpha^{-r}(\eta) \neq 0\} = \alpha^\vee(T) - \inf_{0 \leq t \leq T} \alpha^\vee(\eta(t)).$$

THEOREM 4.12. *With the above definitions, L_π is a normal continuous crystal with highest weight $\pi(T)$.*

PROOF. All properties have been verified in the preceding sections, except the highest weight property, but this follows from the fact that any $\eta \in L_\pi$ can be written as

$$\eta = \mathcal{H}_{s_q}^{x_q} \mathcal{H}_{s_{q-1}}^{x_{q-1}} \cdots \mathcal{H}_{s_1}^{x_1} \pi,$$

see proposition 4.2. □

The geometry of this crystal is easy to describe, using the sets C_i^π which parametrize L_π . We have seen (cf. theorem 4.8) that C_i^π depend on the path π only through $\pi(T)$. We put on C_i^π a continuous crystal structure in the following way. Let $\mathbf{i} = (s_1, \dots, s_q)$ where $w_0 = s_1 \cdots s_q$ is a reduced decomposition. If $x = (x_1, \dots, x_q) \in C_i^\pi$ we set

$$wt(x) = \pi(T) - \sum_{k=1}^q x_k \alpha_k.$$

If the simple root α is α_{s_1} then we define $e_{\alpha, \mathbf{i}}^r$ for $r \in \mathbb{R}$ by

$$e_{\alpha, \mathbf{i}}^r(x_1, x_2, \dots, x_q) = (x_1 + r, x_2, \dots, x_q) \text{ or } \mathbf{0}$$

depending whether $(x_1 + r, \dots, x_q)$ is in C_i^π or not. We let, for $b \in C_i^\pi$,

$$\varepsilon_\alpha(b) = \max\{r \geq 0; e_{\alpha, \mathbf{i}}^r(b) \neq \mathbf{0}\}$$

and

$$\varphi_\alpha(b) = \max\{r \geq 0; e_{\alpha, \mathbf{i}}^{-r}(b) \neq \mathbf{0}\}.$$

We now consider the case where α is not α_1 . We choose a reduced decomposition $w_0 = s'_1 s'_2 \cdots s'_q$ with $\alpha_{s'_1} = \alpha$ and let $\mathbf{j} = (s'_1, s'_2, \dots, s'_q)$. We can define $e_{\alpha, \mathbf{j}}^r$ on C_j^π , ε_α , ϕ_α as above and transport this action on C_i^π by the piecewise linear map $\phi_i^{\mathbf{j}}$ introduced in theorem 3.9. In other words

$$e_{\alpha, \mathbf{i}}^r = \phi_i^{\mathbf{j}} \circ e_{\alpha, \mathbf{j}}^r \circ \phi_j^{\mathbf{i}}.$$

Then $\rho_i : L_\pi \rightarrow C_i^\pi$ is an isomorphism of crystal. This first shows that our construction does not depend on the chosen decompositions $w_0 = s'_1 s'_2 \cdots s'_q$ and then that the crystal structure on L_π depends only on the extremity $\pi(T)$ of the path π :

THEOREM 4.13. *If π and $\bar{\pi}$ are two dominant paths such that $\pi(T) = \bar{\pi}(T)$ then the crystals on L_π and $L_{\bar{\pi}}$ are isomorphic.*

This is the analogue of Littelmann independence theorem (see [25]).

DEFINITION 4.14. *When W is finite, for $\lambda \in \bar{C}$, we denote $B(\lambda)$ the class of the continuous crystals isomorphic to L_π where π is a dominant path such that $\pi(T) = \lambda$.*

4.8. Concatenation and closed crystals. The concatenation $\pi \star \eta$ of two paths $\pi : [0, T] \rightarrow V$, $\eta : [0, T] \rightarrow V$ is defined in Littelmann [25] as the path $\pi \star \eta : [0, T] \rightarrow V$ given by $(\pi \star \eta)(t) = \pi(2t)$, and $(\pi \star \eta)(t + T/2) = \pi(T) + \eta(t)$ when $0 \leq t \leq T/2$. Recall that L_π is the crystal contained in $C_T^0(V)$ generated by a dominant path π . If π_1 and π_2 are two dominant paths, let $D(\pi_1, \pi_2) = \mathcal{P}_{w_0}(L_{\pi_1} \star L_{\pi_2})$ and

$$L(\pi_1, \pi_2) = \cup \{L_\pi; \pi \in D(\pi_1, \pi_2)\}.$$

Since $L(\pi_1, \pi_2)$ is a union of crystals it is itself a crystal.

THEOREM 4.15. *If π_1 and π_2 are two dominant paths, there exists a crystal embedding*

$$\Theta : L_{\pi_1} \otimes L_{\pi_2} \rightarrow L(\pi_1, \pi_2) \subset C_T^0(V)$$

defined by $\Theta(\eta_1 \otimes \eta_2) = \eta_1 \star \eta_2$.

PROOF. We have to show that, for simple roots α , for $\eta_1 \in L_{\pi_1}$, $\eta_2 \in L_{\pi_2}$, for all $s \in \mathbb{R}$,

$$\Theta[e_\alpha^s(\eta_1 \otimes \eta_2)] = \mathcal{E}_\alpha^s(\eta_1 \star \eta_2).$$

This is a purely one-dimensional statement, which uses only one root, hence it follows from the similar fact for Littelmann and Kashiwara crystals. For the convenience of the reader we provide a proof below.

For any $x \geq 0$, let

$$\mathcal{P}_\alpha^x \eta(t) = \eta(t) - \min(0, x + \inf_{0 \leq s \leq t} \alpha^\vee \eta(s)) \alpha.$$

Thus

$$\mathcal{P}_\alpha^x \eta = \eta \quad \text{if} \quad \inf_{0 \leq s \leq t} \alpha^\vee \eta(s) \geq -x$$

and

$$(4.4) \quad \mathcal{P}_\alpha^x \eta = \mathcal{E}_\alpha^y \eta$$

with $y = (-\inf_{0 \leq s \leq T} \alpha^\vee \eta(s) - x) \vee 0$.

LEMMA 4.16. *Let $\eta_1, \eta_2 \in C_T^0(V)$, $x \geq 0$, $y \in [0, \alpha^\vee \pi(T)]$, and π be an α -dominant path, then*

- (i) $\mathcal{P}_\alpha(\eta_1 \star \eta_2) = \mathcal{P}_\alpha \eta_1 \star \mathcal{P}_\alpha^x \eta_2$ where $x = \alpha^\vee \eta_1(T) - \inf_{0 \leq t \leq T} \alpha^\vee \eta_1(t)$;
- (ii) $\mathcal{P}_\alpha \mathcal{P}_\alpha^x = \mathcal{P}_\alpha$;
- (iii) $\mathcal{P}_\alpha^x \mathcal{H}_\alpha^y \pi = \mathcal{H}_\alpha^{x \wedge y} \pi$.

PROOF. For all $t \in [0, T/2]$

$$\mathcal{P}_\alpha(\eta_1 \star \eta_2)(t) = \mathcal{P}_\alpha \eta_1(t).$$

Furthermore,

$$\begin{aligned}
& \mathcal{P}_\alpha(\eta_1 \star \eta_2)((T+t)/2) \\
&= (\eta_1 \star \eta_2)((T+t)/2) - \min[\inf_{0 \leq s \leq T} \alpha^\vee \eta_1(s), \alpha^\vee \eta_1(T) + \inf_{0 \leq s \leq t} \alpha^\vee \eta_2(s)]\alpha \\
&= \eta_1(T) - \inf_{0 \leq s \leq T} \alpha^\vee \eta_1(s)\alpha + \\
&\quad \eta_2(t) - \min[0, \inf_{0 \leq s \leq t} \alpha^\vee \eta_2(s) + \alpha^\vee \eta_1(T) - \inf_{0 \leq s \leq T} \alpha^\vee \eta_1(s)]\alpha \\
&= \mathcal{P}_\alpha \eta_1(T) + \mathcal{P}_\alpha \eta_2(t).
\end{aligned}$$

This proves (i), and (ii) follows from 4.4. Furthermore, $\inf_{0 \leq s \leq T} \alpha^\vee(\mathcal{H}_\alpha^y \pi(s)) = -y$, therefore (iii) follows also from 4.4. \square

PROPOSITION 4.17. *Let π_1, π_2 be α -dominant, $x \in [0, \alpha^\vee \pi_1(T)]$, $y \in [0, \alpha^\vee \pi_2(T)]$, $z = \min(y, \alpha^\vee \pi_1(T) - x)$ and $r = x + y - z$, then*

$$\mathcal{H}_\alpha^x \pi_1 \star \mathcal{H}_\alpha^y \pi_2 = \mathcal{H}_\alpha^r (\pi_1 \star \mathcal{H}_\alpha^z \pi_2),$$

PROOF. Let $s = \alpha^\vee(\mathcal{H}_\alpha^x \pi_1(T)) - \inf_{0 \leq t \leq T} \alpha^\vee(\mathcal{H}_\alpha^x \pi_1)(t)$. By lemma 4.16:

$$\mathcal{P}_\alpha(\mathcal{H}_\alpha^x \pi_1 \star \mathcal{H}_\alpha^y \pi_2) = \mathcal{P}_\alpha(\mathcal{H}_\alpha^x \pi_1) \star \mathcal{P}_\alpha^s(\mathcal{H}_\alpha^y \pi_2)$$

and $\mathcal{P}_\alpha^s \mathcal{H}_\alpha^y \pi_2 = \mathcal{H}_\alpha^{s \wedge y} \pi_2$. Since $\mathcal{P}_\alpha \mathcal{H}_\alpha^x \pi_1 = \pi_1$ one has

$$\mathcal{P}_\alpha(\mathcal{H}_\alpha^x \pi_1 \star \mathcal{H}_\alpha^y \pi_2) = \pi_1 \star \mathcal{H}_\alpha^{s \wedge y} \pi_2.$$

Notice that $s = \alpha^\vee(\pi_1(T)) - x$. On the other hand,

$$\begin{aligned}
(\mathcal{H}_\alpha^x \pi_1 \star \mathcal{H}_\alpha^y \pi_2)(T) &= \mathcal{H}_\alpha^x \pi_1(T) + \mathcal{H}_\alpha^y \pi_2(T) = \pi_1(T) + \pi_2(T) - (x + y)\alpha \\
(\pi_1 \star \mathcal{H}_\alpha^{s \wedge y} \pi_2)(T) &= \pi_1(T) + \pi_2(T) - (s \wedge y)\alpha
\end{aligned}$$

and we know that $\eta = \mathcal{H}_\alpha^r \pi$ is characterized by the properties $\mathcal{P}_\alpha \eta = \pi$ and $\eta(T) = \pi(T) - r\alpha$. Therefore the proposition holds for $r + s \wedge y = x + y$. \square

We now prove that, for $\alpha \in \Sigma$, $\eta_1 \in L_{\pi_1}$, $\eta_2 \in L_{\pi_2}$, for all $s \in \mathbb{R}$,

$$\Theta[e_\alpha^s(\eta_1 \otimes \eta_2)] = \mathcal{E}_\alpha^s(\eta_1 \star \eta_2).$$

Since $e_\alpha^s e_\alpha^t = e_\alpha^{s+t}$ and $\mathcal{E}_\alpha^s \mathcal{E}_\alpha^t = \mathcal{E}_\alpha^{s+t}$ it is sufficient to check this for s near 0. We write $\eta_1 = \mathcal{H}_\alpha^x \pi_1$ and $\eta_2 = \mathcal{H}_\alpha^y \pi_2$ where $\pi_1 = \mathcal{P}_\alpha(\eta_1)$, $\pi_2 = \mathcal{P}_\alpha(\eta_2)$ are α -dominant. By proposition 4.17, if $z = \min(y, \alpha^\vee \pi_1(T) - x)$ and $r = x + y - z$, then

$$\mathcal{E}_\alpha^s(\eta_1 \star \eta_2) = \mathcal{E}_\alpha^s(\mathcal{H}_\alpha^x \pi_1 \star \mathcal{H}_\alpha^y \pi_2) = \mathcal{E}_\alpha^s \mathcal{H}_\alpha^r (\pi_1 \star \mathcal{H}_\alpha^z \pi_2).$$

We first show that if

$$(4.5) \quad \mathcal{E}_\alpha^s(\eta_1 \star \eta_2) = \mathbf{0}$$

then $e_\alpha^s(\eta_1 \otimes \eta_2) = \mathbf{0}$. For $|s|$ small enough (4.5) holds only when $r = 0$ and $s > 0$ or when $s < 0$ and

$$(4.6) \quad r = \alpha^\vee((\pi_1 \star \mathcal{H}_\alpha^z \pi_2)(T)) = \alpha^\vee \pi_1(T) + \alpha^\vee \pi_2(T) - 2z.$$

If $r = 0$, then $z = \min(y, \alpha^\vee \pi_1(T) - x) = x + y$ hence $x = 0$ and $y \leq \alpha^\vee \pi_1(T)$. But

$$\varepsilon_\alpha(\eta_1 \otimes \eta_2) = \varepsilon_\alpha(\eta_1) - \min(\varphi_\alpha(\eta_1) - \varepsilon_\alpha(\eta_2), 0) = \max(2x + y - \alpha^\vee \pi_1(T), x).$$

(notice that, in general, when π is α -dominant, $\varepsilon_\alpha(\mathcal{H}_\alpha^x \pi) = x$ and $\varphi_\alpha(\mathcal{H}_\alpha^x \pi) = \alpha^\vee \pi(T) - x$). Therefore $\varepsilon_\alpha(\eta_1 \otimes \eta_2) = 0$ and $e_\alpha^s(\eta_1 \otimes \eta_2) = \mathbf{0}$. Now, if r is given by (4.6), then

$$z = \alpha^\vee \pi_1(T) - x + \alpha^\vee \pi_2(T) - y$$

since $r = x + y - z$. We know that $\alpha^\vee \pi_2(T) - y \geq 0$, hence $z = \min(y, \alpha^\vee \pi_1(T) - x)$ only if

$$z = \alpha^\vee \pi_1(T) - x, \alpha^\vee \pi_2(T) = y, y \geq \alpha^\vee \pi_1(T) - x.$$

Then

$$\varepsilon_\alpha(\eta_1 \otimes \eta_2) = 2x + y - \alpha^\vee \pi_1(T).$$

On the other hand,

$$wt(\eta_1 \otimes \eta_2) = wt(\eta_1) + wt(\eta_2) = \pi_1(T) - x\alpha + \pi_2(T) - y\alpha,$$

thus, using $y = \alpha^\vee \pi_2(T)$,

$$\varphi_\alpha(\eta_1 \otimes \eta_2) = \varepsilon_\alpha(\eta_1 \otimes \eta_2) + \alpha^\vee(wt(\eta_1 \otimes \eta_2)) = 0$$

and $e_\alpha^s(\eta_1 \otimes \eta_2) = \mathbf{0}$ when $s < 0$.

We now consider the case where (4.5) does not hold. Then for s small enough,

$$\mathcal{E}_\alpha^s(\eta_1 \star \eta_2) = \mathcal{E}_\alpha^s \mathcal{H}_\alpha^r(\pi_1 \star \mathcal{H}_\alpha^z \pi_2) = \mathcal{H}_\alpha^{r-s}(\pi_1 \star \mathcal{H}_\alpha^z \pi_2).$$

Using proposition 4.17, if s is small enough, and $y > \alpha^\vee \pi_1(T) - x$, then

$$\mathcal{H}_\alpha^{r-s}(\pi_1 \star \mathcal{H}_\alpha^z \pi_2) = \mathcal{H}_\alpha^{x-s} \pi_1 \star \mathcal{H}_\alpha^y \pi_2 = \Theta(e_\alpha^s(\mathcal{H}_\alpha^x \pi_1 \otimes \mathcal{H}_\alpha^y \pi_2))$$

and if $y < \alpha^\vee \pi_1(T) - x$, then

$$\mathcal{H}_\alpha^{r-s}(\pi_1 \star \mathcal{H}_\alpha^z \pi_2) = \mathcal{H}_\alpha^x \pi_1 \star \mathcal{H}_\alpha^{y-s} \pi_2 = \Theta(e_\alpha^s(\mathcal{H}_\alpha^x \pi_1 \otimes \mathcal{H}_\alpha^y \pi_2)).$$

The end of the proof is straightforward. \square

This proves that the family of crystals $B(\lambda), \lambda \in \bar{C}$ is closed. From theorem 4.12 and theorem 2.6, we get

THEOREM 4.18. *When W is a finite Coxeter group, there exists one and only one closed family of highest weight normal continuous crystals $B(\lambda), \lambda \in \bar{C}$.*

4.9. Action of W on the Littelmann crystal. Following Kashiwara [20], [22] and Littelmann [25], we show that we can define an action of the Coxeter group on each crystal L_π . We first notice that for each simple root α , we can define an involution S_α on the set of paths by

$$S_\alpha \eta = \mathcal{E}_\alpha^x \eta \quad \text{for } x = -\alpha^\vee(\eta(T)).$$

In particular,

$$(4.7) \quad S_\alpha \eta(T) = s_\alpha(\eta(T)).$$

LEMMA 4.19. *Let $\eta \in C_T^0(V)$ and $\alpha \in \Sigma$ such that $\alpha^\vee(\eta(T)) < 0$. For each $\gamma \in C_T^0(V)$ there exists $m \in \mathbb{N}$ such that, for all $n \geq 0$,*

$$\mathcal{P}_\alpha(\gamma \star \eta^{*(m+n)}) = \mathcal{P}_\alpha(\gamma \star \eta^{*m}) \star S_\alpha(\eta)^{*n}.$$

PROOF. By lemma 4.16,

$$\mathcal{P}_\alpha(\gamma \star \eta^{*(n+1)}) = \mathcal{P}_\alpha(\gamma \star \eta^{*n}) \star \mathcal{P}_\alpha^x(\eta)$$

where

$$x = \alpha^\vee(\gamma \star \eta^{*n})(T) - \min_{0 \leq s \leq T} \alpha^\vee(\gamma \star \eta^{*n})(s).$$

Let $\gamma_{\min} = \min_{0 \leq s \leq T} \alpha^\vee \gamma(s)$ and $\eta_{\min} = \min_{0 \leq s \leq T} \alpha^\vee \eta(s)$. Since $\alpha^\vee \gamma(T) < 0$, there exists $m > 0$ such that for $n \geq m$ one has,

$$\begin{aligned} \min_{0 \leq s \leq T} \alpha^\vee(\gamma \star \eta^{*n})(s) &= \min(\gamma_{\min}, \alpha^\vee(\gamma(T) + k\eta(T)) + \eta_{\min}; 0 \leq k \leq n-1) \\ &= \alpha^\vee(\gamma(T) + (n-1)\eta(T)) + \eta_{\min}. \end{aligned}$$

Using that $(\gamma \star \eta^{*m})(T) = \gamma(T) + m\eta(T)$ we have $x = \alpha^\vee \eta(T) - \eta_{\min}$. In this case, $\mathcal{P}_\alpha^x(\eta) = S_\alpha(\eta)$, which proves the lemma by induction on $n \geq m$. \square

THEOREM 4.20. *There is an action $\{S_w, w \in W\}$ of the Coxeter group W on each L_π such that $S_{s_\alpha} = S_\alpha$ when α is a simple root.*

PROOF. By Matsumoto's lemma, it suffices to prove that the transformations S_α satisfy to the braid relations. Therefore we can assume that W is a dihedral group $I(q)$. Consider two roots α, β generating W . Let η be a path, there exists a sequence $(\alpha_i) = \alpha, \beta, \alpha, \dots$ or $\beta, \alpha, \beta, \dots$ such that $s_{\alpha_1} s_{\alpha_2} \dots s_{\alpha_r} \eta(T) \in -\bar{C}$. Let $\tilde{\eta} = S_{\alpha_1} S_{\alpha_2} \dots S_{\alpha_r} \eta$. Let $s_{\alpha_q} \dots s_{\alpha_1}$ be a reduced decomposition. We show by induction on $k \leq q$ that there exists $m_k \geq 0$ and a path γ_k such that

$$(4.8) \quad \mathcal{P}_{\alpha_k} \dots \mathcal{P}_{\alpha_1}(\tilde{\eta}^{*(m_k+n)}) = \gamma_k \star (S_{\alpha_k} \dots S_{\alpha_1} \tilde{\eta})^{\star n}$$

For $k = 1$, this is the preceding lemma. Suppose that this holds for some k . Then

$$\alpha_{k+1}^\vee(S_{\alpha_k} \dots S_{\alpha_1} \tilde{\eta}(T)) \leq 0$$

(cf. Bourbaki, [4], ch.5, no.4, Th. 1). Thus, by the lemma, there exists m such that, for $n \geq 0$,

$$\mathcal{P}_{\alpha_{k+1}}(\gamma_k \star (S_{\alpha_k} \dots S_{\alpha_1} \tilde{\eta})^{\star(m+n)}) = \mathcal{P}_{\alpha_{k+1}}(\gamma_k \star (S_{\alpha_k} \dots S_{\alpha_1} \tilde{\eta})^{\star m}) \star (S_{\alpha_{k+1}} S_{\alpha_k} \dots S_{\alpha_1} \tilde{\eta})^{\star n}$$

Hence, by the induction hypothesis, if $\gamma_{k+1} = \mathcal{P}_{\alpha_{k+1}}(\gamma_k \star (S_{\alpha_k} \dots S_{\alpha_1} \tilde{\eta})^{\star m})$, then

$$\mathcal{P}_{\alpha_{k+1}} \mathcal{P}_{\alpha_k} \dots \mathcal{P}_{\alpha_1}((\tilde{\eta}^{\star(m_k+m+n)})) = \gamma_{k+1} \star (S_{\alpha_{k+1}} S_{\alpha_k} \dots S_{\alpha_1} \tilde{\eta})^{\star n}$$

We apply (4.8) with $k = q$, then there exists two reduced decompositions, and we see that $S_{\alpha_q} S_{\alpha_{q-1}} \dots S_{\alpha_1} \tilde{\eta}$ does not depend on the reduced decomposition because the left hand side does not, by the braid relations for the \mathcal{P}_α . This implies easily that $S_{\alpha_q} S_{\alpha_{q-1}} \dots S_{\alpha_1} \eta$ also does not depend on the reduced decomposition. \square .

Using the crystal isomorphism between L_π and the crystal $B(\pi(T))$ we see that

COROLLARY 4.21. *The Coxeter group W acts on each crystal $B(\lambda)$, where $\lambda \in \bar{C}$, in such a way that, for $s = s_\alpha$ in S , and $b \in B(\lambda)$,*

$$S_\alpha(b) = e_\alpha^x(b), \text{ where } x = -\alpha^\vee(wt(b)).$$

Notice that these S_α are not crystal morphisms.

4.10. Schützenberger involution. The classical Schützenberger involution associates to a Young tableau T another Young tableau \hat{T} of the same shape. If (P, Q) is the pair associated by Robinson-Schensted-Knuth (RSK) algorithm to the word $u_1 \dots u_n$ in the letters $1, \dots, k$, then (\hat{P}, \hat{Q}) is the pair associated with $u_1^* \dots u_n^*$ where $i^* = k + 1 - i$, see e.g. Fulton [8]. It is remarkable that \hat{P} depends only on P , and that \hat{Q} depends only on Q . We will establish an analogous property for the analogue of the Schützenberger involution defined in [2] for finite Coxeter groups. The crystallographic case has been recently investigated by Henriques and Kamnitzer [14], [15], Morier-Genoud [28], and Lenart [24].

For any path $\eta \in C_T^0(V)$, let

$$\kappa\eta(t) = \eta(T - t) - \eta(T), \quad 0 \leq t \leq T.$$

Let us show that

$$S\eta = -w_0\kappa\eta,$$

suitably interpreted, is the analogue of Schützenberger involution. Since $w_0^2 = id$, S is indeed an involution. The following is proved in [2].

PROPOSITION 4.22. *For any $\eta \in C_T^0(V)$,*

$$\mathcal{P}_{w_0} S\eta(T) = \mathcal{P}_{w_0} \eta(T).$$

As remarked there, this implies that the transformation on dominant paths

$$\pi \mapsto I\pi = \mathcal{P}_{w_0} S\pi$$

gives the analogue of the Schützenberger involution on the Q 's. We will consider the action on the crystal itself, i.e. the analogue of the Schützenberger involution on the P 's.

Let π be a dominant path and $\lambda = \pi(T)$. Notice that the transformation S on $C_T^0(V)$ does not preserve the Littelmann module L_π . It sends L_π onto $L_{I\pi}$. By the preceding proposition $\lambda = \pi(T) = I\pi(T)$, therefore L_π and $L_{I\pi}$ are crystals isomorphic to $B(\lambda)$. More precisely, there is a unique strict morphism $G : L_\pi \rightarrow L_{I\pi}$ such that $G(\pi) = I\pi$.

If α is a simple root, then $\tilde{\alpha} = -w_0\alpha$ is also a simple root and $\tilde{\alpha}^\vee = -\alpha^\vee w_0$. The following property is straightforward. In the A_n case, it was shown by Lascoux, Leclerc and Thibon [23] and Henriques and Kamnitzer [14] that it characterizes the Schützenberger involution on the crystals $B(\lambda)$.

LEMMA 4.23. *For any path η in $C_T^0(V)$, any $r \in \mathbb{R}$, and any simple root α , one has*

$$\begin{aligned} \mathcal{E}_\alpha^r S\eta &= S\mathcal{E}_{\tilde{\alpha}}^{-r} \eta \\ \varepsilon_{\tilde{\alpha}}(S\eta) &= \varphi_\alpha(\eta), \varphi_{\tilde{\alpha}}(S\eta) = \varepsilon_\alpha(\eta) \\ S\eta(T) &= w_0\eta(T). \end{aligned}$$

For each simple root α , we define the co-Pitman operators \mathcal{F}_α by $\mathcal{F}_\alpha = \kappa \mathcal{P}_\alpha \kappa$. Note that these operators were defined in [2], section 4.2, where the unfortunate notation \mathcal{E}_α was used, however we prefer the notation here. If π is α -dominant, then

$$\mathcal{F}_\alpha \mathcal{H}_\alpha^x \pi = \mathcal{H}_\alpha^z \pi$$

where $z = \alpha^\vee \pi(T)$. In other words, $\mathcal{F}_\alpha \mathcal{H}_\alpha^x \pi$ is the “lowest” element in the α -string of π .

PROPOSITION 4.24. *If $\sigma \in C_T^0(V)$, the string parameters of $\mathcal{F}_\alpha \sigma$ and $\mathcal{F}_\alpha \sigma(T)$ depend only on the string parameters of σ and $\sigma(T)$.*

PROOF. Choose a reduced decomposition $\mathbf{i}(\alpha)$ of w_0 which ends with s_α (i.e. $\alpha_{s_q} = \alpha$) and let $\varrho_{\mathbf{i}(\alpha)}(\sigma) = (x_1, x_2, \dots, x_{q-1}, x_q)$. Then $\sigma = \mathcal{H}_\alpha^{x_q} \mathcal{P}_\alpha \sigma$, therefore $\mathcal{F}_\alpha \sigma = \mathcal{H}_\alpha^z \mathcal{P}_\alpha \sigma$ for $z = \alpha^\vee(\mathcal{P}_\alpha \sigma(T)) = \alpha^\vee(\sigma(T)) + 2x_q$, hence

$$\varrho_{\mathbf{i}(\alpha)}(\mathcal{F}_\alpha \sigma) = (x_1, x_2, \dots, x_{q-1}, z).$$

On the other hand, $\mathcal{F}_\alpha \sigma(T) = \mathcal{P}_\alpha \sigma(T) - z\alpha = \sigma(T) + (x_q - z)\alpha$. \square

PROPOSITION 4.25. *Let $\eta \in C_T^0(V)$, then $S\eta(T)$ and the string parameters of $S\eta$ depend only on $\eta(T)$ and on the string parameters of η .*

PROOF. One has $(-w_0)\mathcal{P}_\alpha(-w_0) = \mathcal{P}_{\tilde{\alpha}}$, (recall that $\tilde{\alpha} = -w_0\alpha$) and,

$$(4.9) \quad S\mathcal{F}_\alpha = \mathcal{P}_{\tilde{\alpha}} S.$$

Let $\mathbf{i} = (s_1, \dots, s_q)$, where $w_0 = s_1 \cdots s_q$ is a reduced decomposition and $\mathbf{i}' = (\tilde{s}_1, \dots, \tilde{s}_q)$. Then $w_0 = \tilde{s}_1 \cdots \tilde{s}_q$ is also a reduced decomposition. Let us compute the \mathbf{i}' string coordinates $\varrho_{\mathbf{i}'}(S\eta) = (y_1, \dots, y_q)$ of $S\eta$, for which

$$S\eta = \mathcal{H}_{\alpha_q}^{y_q} \mathcal{H}_{\alpha_{q-1}}^{y_{q-1}} \cdots \mathcal{H}_{\alpha_1}^{y_1} \mathcal{P}_{w_0} S\eta.$$

Let $\sigma_q = \eta$ and, for any $k = 0, \dots, q$,

$$\sigma_k = \mathcal{F}_{\alpha_{k+1}} \cdots \mathcal{F}_{\alpha_q} \eta$$

then, by (4.9),

$$S\sigma_k = \mathcal{P}_{\tilde{\alpha}_{k+1}} \cdots \mathcal{P}_{\tilde{\alpha}_q} S\eta = \mathcal{H}_{\tilde{s}_k}^{y_k} \cdots \mathcal{H}_{\tilde{s}_1}^{y_1} \mathcal{P}_{w_0} S\eta.$$

It follows from the proposition above that $\sigma_k(T)$ and the string coordinates of σ_k depend only on $\eta(T)$ and the string coordinates of η . In particular this is the case of $\alpha_k^\vee \sigma_k(T)$ and $\min_{0 \leq s \leq T} \alpha_k^\vee \sigma_k(s)$ and therefore of

$$\begin{aligned} y_k &= - \inf_{0 \leq s \leq T} \tilde{\alpha}_k^\vee(S\sigma_k)(s) = - \inf_{0 \leq s \leq T} \alpha_k^\vee(-w_0 S\sigma_k)(s) \\ &= - \inf_{0 \leq s \leq T} \alpha_k^\vee(\kappa \sigma_k)(s) = - \inf_{0 \leq s \leq T} \alpha_k^\vee(\sigma_k(T-s) - \sigma_k(T)) \\ &= \alpha_k^\vee \sigma_k(T) - \inf_{0 \leq s \leq T} \alpha_k^\vee \sigma_k(s). \quad \square \end{aligned}$$

Let $\eta \in C_T^0(V)$. Fix a reduced decomposition $\mathbf{i} = (\alpha_1, \dots, \alpha_q)$ of w_0 and let $(y_1, \dots, y_q) = \varrho_{\mathbf{i}}(S\eta)$ be the string parametrization of $S\eta$. One has, by definition,

$$(4.10) \quad S\eta = \mathcal{H}_{\alpha_q}^{y_q} \cdots \mathcal{H}_{\alpha_1}^{y_1} \mathcal{P}_{w_0} S\eta.$$

Since $\mathcal{P}_{w_0} S\eta(T) = \mathcal{P}_{w_0} \eta(T)$, by proposition 4.22, we know that (y_1, \dots, y_q) is in the polyhedron $C_{\mathbf{i}}^\lambda$ where $\lambda = \mathcal{P}_{w_0} \eta(T)$. Therefore we can define

$$\tilde{S}\eta = \mathcal{H}_{\alpha_q}^{y_q} \cdots \mathcal{H}_{\alpha_1}^{y_1} \mathcal{P}_{w_0} \eta.$$

In the A_n case, in the RSK interpretation the two paths $S\eta$ and $\tilde{S}\eta$ have the same P but not the same Q . It follows from lemma 4.23 that

LEMMA 4.26. *Let $\eta \in C_T^0(V)$, then*

$$\begin{aligned} \tilde{S}(\mathcal{E}_\alpha^r(\eta)) &= \mathcal{E}_{\tilde{\alpha}}^{-r}(\tilde{S}(\eta)), \quad wt(\tilde{S}(\eta)) = w_0 wt(\eta), \\ \varphi_\alpha(\tilde{S}(\eta)) &= \varepsilon_{\tilde{\alpha}}(\eta), \quad \varepsilon_\alpha(\tilde{S}(\eta)) = \phi_{\tilde{\alpha}}(\eta). \end{aligned}$$

THEOREM 4.27. *The map $\tau : C_T^0(V) \rightarrow C_T^0(V)$ defined by*

$$\tau(\eta_1 \star \eta_2) = \tilde{S}(S\eta_2 \star S\eta_1)$$

is a crystal isomorphism.

PROOF. Let $\eta_1, \eta_2 \in C_T^0(V)$, $\sigma = \varphi_\alpha(\eta_1) - \varepsilon_\alpha(\eta_2)$ and $\tilde{\sigma} = -\sigma$. Then

$$S(\mathcal{E}_\alpha^{\min(r, -\sigma) + \sigma^+} \eta_2) = \mathcal{E}_{\tilde{\alpha}}^{-\min(r, -\sigma) - \sigma^+} S\eta_2 = \mathcal{E}_{\tilde{\alpha}}^{\max(-r, \tilde{\sigma}) - \tilde{\sigma}^-} S\eta_2$$

and

$$S(\mathcal{E}_\alpha^{\max(r, -\sigma) - \sigma^-} \eta_1) = \mathcal{E}_{\tilde{\alpha}}^{-\max(r, -\sigma) + \sigma^-} S\eta_1 = \mathcal{E}_{\tilde{\alpha}}^{\min(-r, -\tilde{\sigma}) + \tilde{\sigma}^+} S\eta_1$$

Therefore, using lemma 4.23,

$$S(\mathcal{E}_\alpha^{\min(r, -\sigma) + \sigma^+} \eta_2) \star S(\mathcal{E}_\alpha^{\max(r, -\sigma) - \sigma^-} \eta_1) = \mathcal{E}_{\tilde{\alpha}}^{-r}(S\eta_2 \star S\eta_1)$$

hence

$$\begin{aligned} \tau(\mathcal{E}_\alpha^r(\eta_1 \star \eta_2)) &= \tau(\mathcal{E}_\alpha^{\max(r, -\sigma) - \sigma^-} \eta_1 \star \mathcal{E}_\alpha^{\min(r, -\sigma) + \sigma^+} \eta_2) \\ &= \tilde{S}(S(\mathcal{E}_\alpha^{\min(r, -\sigma) + \sigma^+} \eta_2) \star S(\mathcal{E}_\alpha^{\max(r, -\sigma) - \sigma^-} \eta_1)) \\ &= \tilde{S}(\mathcal{E}_{\tilde{\alpha}}^{-r}(S\eta_2 \star S\eta_1)) \\ &= \mathcal{E}_\alpha^r \tilde{S}(S\eta_2 \star S\eta_1) \\ &= \mathcal{E}_\alpha^r \tau(\eta_1 \star \eta_2). \end{aligned}$$

On the other hand, using lemma 4.26, and the fact that $S\eta(T) = w_0\eta(T)$,

$$\begin{aligned} wt(\tau(\eta_1 \star \eta_2)) &= wt(\tilde{S}(S\eta_2 \star S\eta_1)) = w_0 wt(S\eta_2 \star S\eta_1) \\ w_0(S\eta_2 \star S\eta_1)(T) &= w_0(w_0(\eta_2(T) + \eta_1(T))) = wt(\eta_1 \star \eta_2) \end{aligned}$$

One checks analogously the intertwining relations $\varepsilon_\alpha \circ \tau = \tau \circ \varepsilon_\alpha$ and $\varphi_\alpha \circ \tau = \tau \circ \varphi_\alpha$. \square

We can now define an involution S_λ on each continuous crystal of the family $\{B(\lambda), \lambda \in \bar{C}\}$ by transporting the action of S on $C_T^0(V)$:

$$\begin{array}{ccc} L_\pi & \xrightarrow{S} & L_{I\pi} \\ \varrho_i^\pi \downarrow & & \downarrow \varrho_i^{I\pi} \\ B(\lambda) & \xrightarrow{S_\lambda} & B(\lambda) \end{array}$$

where π is a dominant path such that $\pi(T) = \lambda$. It does not depend on π itself by proposition 4.25. Similarly we can define \tilde{S}_λ by transporting the action of \tilde{S}

$$\begin{array}{ccc} L_\pi & \xrightarrow{\tilde{S}} & L_\pi \\ \varrho_i^\pi \downarrow & & \downarrow \varrho_i^\pi \\ B(\lambda) & \xrightarrow{\tilde{S}_\lambda} & B(\lambda) \end{array}$$

Let $\lambda, \mu \in \bar{C}$. For $b_1 \in B(\lambda)$ and $b_2 \in B(\mu)$ let

$$\tau_{\lambda,\mu}(b_1 \otimes b_2) = \tilde{S}_\gamma(S_\mu b_2 \otimes S_\lambda b_1)$$

where $\gamma \in \bar{C}$ is such that $S_\mu b_2 \otimes S_\lambda b_1 \in B(\gamma)$.

THEOREM 4.28. *For $\lambda, \mu \in \bar{C}$, the map*

$$\tau_{\lambda,\mu} : B(\lambda) \otimes B(\mu) \rightarrow B(\mu) \otimes B(\lambda)$$

is a crystal isomorphism.

This follows from theorem 4.27. As in the construction of Henriques and Kamnitzer [14], [15] these isomorphisms do not obey the axioms for a braided monoidal category, but instead we have that:

- (1) $\tau_{\mu,\lambda} \circ \tau_{\lambda,\mu} = 1$
- (2) The following diagram commutes:

$$\begin{array}{ccc} B(\lambda) \otimes B(\mu) \otimes B(\sigma) & \xrightarrow{1 \otimes \tau_{(\mu,\sigma)}} & B(\lambda) \otimes B(\sigma) \otimes B(\mu) \\ \tau_{(\lambda,\mu)} \otimes 1 \downarrow & & \downarrow \tau_{(\lambda,(\sigma,\mu))} \\ B(\mu) \otimes B(\lambda) \otimes B(\sigma) & \xrightarrow{\tau_{((\mu,\lambda),\sigma)}} & B(\sigma) \otimes B(\mu) \otimes B(\lambda) \end{array}$$

which makes of $B(\lambda), \lambda \in \bar{C}$, a coboundary category.

5. The Duistermaat-Heckman measure and Brownian motion

5.1. In this section, we consider a finite Coxeter group, with a realization in some Euclidean space V identified with its dual so that, for each root α , $\alpha^\vee = \frac{2\alpha}{\|\alpha\|^2}$.

5.2. Brownian motion and the Pitman transform. Fix a reduced decomposition of the longest word

$$w_0 = s_1 s_2 \cdots s_q$$

and let $\mathbf{i} = (s_1, \dots, s_q)$. Recall that for any $\eta \in C_T^0(V)$, its string parameters $x = (x_1, \dots, x_q) = \varrho_{\mathbf{i}}(\eta)$ satisfy

$$(5.1) \quad 0 \leq x_i \leq \alpha_{s_i}^\vee \left(\lambda - \sum_{j=1}^{i-1} x_j \alpha_{s_j} \right), \quad i \leq q;$$

where $\lambda = \mathcal{P}_{w_0} \eta(T)$. For each simple root α choose a reduced decomposition $\mathbf{i}_\alpha = (s_1^\alpha, \dots, s_q^\alpha)$ such that $s_1^\alpha = s_\alpha$ and denote the corresponding string parameters $\varrho_{\mathbf{i}_\alpha}(\eta)$ by $(x_1^\alpha, \dots, x_q^\alpha)$. Using the map $\phi_{\mathbf{i}}^{\mathbf{i}_\alpha}$ given by theorem 3.9 we obtain a continuous piecewise linear function $\Psi_\alpha^{\mathbf{i}} : \mathbb{R}^q \rightarrow \mathbb{R}$ such that

$$(5.2) \quad x_1^\alpha = \Psi_\alpha^{\mathbf{i}}(x).$$

Of course

$$(5.3) \quad \Psi_\alpha^{\mathbf{i}}(x) \geq 0, \quad \text{for all } \alpha \in \Sigma.$$

Denote by $M_{\mathbf{i}}$ the set of $(x, \lambda) \in \mathbb{R}_+^q \times C$ which satisfy the inequalities (5.1) and (5.3), and set

$$M_{\mathbf{i}}^\lambda = \{x \in \mathbb{R}_+^q : (x, \lambda) \in M_{\mathbf{i}}\}.$$

Let \mathbb{P} be a probability measure on $C_T^0(V)$ under which η is a standard Brownian motion in V . We recall the following theorem from [2].

THEOREM 5.1. *The stochastic process $\mathcal{P}_{w_0} \eta$ is a Brownian motion in V conditioned, in Doob's sense, to stay in the Weyl chamber \bar{C} .*

This means that $\mathcal{P}_{w_0} \eta$ is the h -process of the standard Brownian motion in V killed when it exits \bar{C} , for the harmonic function

$$h(\lambda) = \prod_{\alpha \in R_+} \alpha^\vee(\lambda),$$

for $\lambda \in V$, where R_+ is the set of all positive roots. Let $c_t = t^{q/2} \int_V e^{-\|\lambda\|^2/2t} d\lambda$ and

$$k = c_1^{-1} \int_C h(\lambda)^2 e^{-\|\lambda\|^2/2} d\lambda.$$

THEOREM 5.2. *For $(\sigma, \lambda) \in M_{\mathbf{i}}$,*

$$(5.4) \quad \mathbb{P}(\varrho_{\mathbf{i}}(\eta) \in d\sigma, \mathcal{P}_{w_0} \eta(T) \in d\lambda) = c_T^{-1} h(\lambda) e^{-\|\lambda\|^2/2T} d\sigma d\lambda.$$

The conditional law of $\varrho_{\mathbf{i}}(\eta)$, given $(\mathcal{P}_{w_0} \eta(s), s \leq T)$ and $\mathcal{P}_{w_0} \eta(T) = \lambda$, is the normalized Lebesgue measure on $M_{\mathbf{i}}^\lambda$, and the volume of $M_{\mathbf{i}}^\lambda$ is $k^{-1} h(\lambda)$.

This theorem has the following interesting corollary, which gives a new proof of the fact that the set $C_{\mathbf{i}}^\pi$ depends only on $\pi(T)$, and is polyhedral.

COROLLARY 5.3. *For any dominant path π , let $\lambda = \pi(T)$, then $C_{\mathbf{i}}^\pi = M_{\mathbf{i}}^\lambda$, and*

$$C_{\mathbf{i}} = \{x \in \mathbb{R}_+^q; \Psi_\alpha^{\mathbf{i}}(x) \geq 0, \text{ for all } \alpha \in \Sigma\}.$$

PROOF. It is clear that C_i^π is contained in M_i^λ and the theorem implies that C_i^π , equal by definition to the set of $\varrho_i(\eta)$ when $\mathcal{P}_{w_0}\eta = \pi$, contains M_i^λ . The description of C_i follows, since $C_i = \cup\{C_i^\pi, \pi \text{ dominant path}\}$. \square

Theorem 5.2 is proved in section 5.4.

5.3. The Duistermaat-Heckman measure. Let G be a compact semi-simple Lie group with maximal torus T . If \mathcal{O}_λ is a coadjoint orbit of G , corresponding to a dominant regular weight, endowed with its canonical symplectic structure ω , then this maximal torus acts on the symplectic manifold $(\mathcal{O}_\lambda, \omega)$, and the image of the Liouville measure on \mathcal{O}_λ by the moment map, which takes values in the dual of the Lie algebra of T , is called the Duistermaat-Heckman measure. It is proved in [1] that this measure is the image of the Lebesgue measure on the Berenstein-Zelevinsky polytope by an affine map. In analogy with this case, we define for a realization of a finite Coxeter group, the Duistermaat-Heckman measure, and prove some properties which generalize the case of crystallographic groups.

DEFINITION 5.4. *For any $\lambda \in C$, the Duistermaat-Heckman measure m_{DH}^λ on V is the image of the Lebesgue measure on M_i^λ by the map*

$$(5.5) \quad x = (x_1, \dots, x_q) \in M_i^\lambda \mapsto \lambda - \sum_{j=1}^q x_j \alpha_j \in V.$$

THEOREM 5.5. *The Laplace transform of the Duistermaat-Heckman measure is given, for $z \in V^*$, by*

$$\int_V e^{\langle z, v \rangle} m_{DH}^\lambda(dv) = \frac{\sum_{w \in W} \varepsilon(w) e^{\langle z, w\lambda \rangle}}{h(z)},$$

where $\varepsilon(w)$ is the signature of $w \in W$.

With the notations of theorem 5.2, the conditional law of $\eta(T)$, given $(\mathcal{P}_{w_0}\eta(s), 0 \leq s \leq T)$ and $\mathcal{P}_{w_0}\eta(T) = \lambda$, is the probability measure $\mu_{DH}^\lambda = k m_{DH}^\lambda / h(\lambda)$.

Theorem 5.5 is proved in section 5.5

PROPOSITION 5.6. *The Duistermaat-Heckman measure m_{DH}^λ has a continuous piecewise polynomial density, invariant under W and with support equal to the convex hull $co(W\lambda)$ of $W\lambda$.*

PROOF. The measure m_{DH}^λ is the image by an affine map of the Lebesgue measure on the convex polytope C_i^π when $\pi(T) = \lambda$. Therefore it has a piecewise polynomial density and a convex support. Its Laplace transform is invariant under W so m_{DH}^λ itself is invariant under W . The support $S(\lambda)$ is equal to $\{\eta(T); \eta \in L_\pi\}$. Notice that if η is in L_π , then when $x = \alpha^\vee(\eta(T))$, $\mathcal{E}_\alpha^x \eta$ is in L_π and $\mathcal{E}_\alpha^x \eta(T) = s_\alpha \eta(T)$. Starting from $\pi(T) = \lambda$ we thus see that $W\lambda$ is contained in $S(\lambda)$. So $co(W\lambda)$ is contained in $S(\lambda)$. The components of $x \in M_i^\pi$ are non negative, therefore $co(W\lambda)$ contains $S(\lambda) \cap \bar{C}$ and, by W -invariance it contains $S(\lambda)$ itself. \square

5.4. Proof of theorem 5.2. First we recall some further path transformations which were introduced in [2]. For any positive root β (not necessarily simple), define $\mathcal{Q}_\beta = \mathcal{P}_\beta s_\beta$. Then, for $\psi \in C_T^0(V)$,

$$\mathcal{Q}_\beta \psi(t) = \psi(t) - \inf_{t \geq s \geq 0} \beta^\vee(\psi(t) - \psi(s))\beta, \quad T \geq t \geq 0.$$

Let $w_0 = s_1 s_2 \dots s_q$ be a reduced decomposition, and let $\alpha_i = \alpha_{s_i}$. Since $s_\alpha \mathcal{P}_\beta = \mathcal{P}_{s_\alpha \beta} s_\alpha$, for roots $\alpha \neq \beta$, the following holds

$$\mathcal{Q}_{w_0} := \mathcal{P}_{w_0} w_0 = \mathcal{Q}_{\beta_1} \dots \mathcal{Q}_{\beta_q},$$

where $\beta_1 = \alpha_1$, $\beta_i = s_1 \dots s_{i-1} \alpha_i$, when $i \leq q$. Set $\psi_q = \psi$ and,

$$(5.6) \quad \psi_{i-1} = \mathcal{Q}_{\beta_i} \dots \mathcal{Q}_{\beta_q} \psi \quad y_i = - \inf_{T \geq t \geq 0} \beta_i^\vee(\psi_i(T) - \psi_i(t)).$$

Then $\psi_0 = \mathcal{Q}_{w_0} \psi$ and, for each $i \leq q$,

$$\mathcal{Q}_{w_0} \psi(T) = \psi_i(T) + \sum_{j=1}^i y_j \beta_j.$$

Define $\varsigma_i(\psi) := (y_1, y_2, \dots, y_q)$. Now let $\eta = w_0 \psi$, so that $\mathcal{Q}_{w_0} \psi = \mathcal{P}_{w_0} \eta$. Set $\eta_q = \eta$ and, for $i \leq q$,

$$(5.7) \quad \eta_{i-1} = \mathcal{P}_{\alpha_i} \dots \mathcal{P}_{\alpha_q} \eta \quad x_i = - \inf_{T \geq t \geq 0} \alpha_i^\vee(\eta_i(t)).$$

Then $\eta_0 = \mathcal{P}_{w_0} \eta$ and, for each $i \leq q$,

$$\mathcal{P}_{w_0} \eta(T) = \eta_i(T) + \sum_{j=1}^i x_j \alpha_j.$$

The parameters $\varrho_i(\eta) = (x_1, \dots, x_q)$ are related to $\varsigma_i(\psi) = (y_1, y_2, \dots, y_q)$ as follows.

LEMMA 5.7. *For each $i \leq q$, we have:*

- (i) $\eta_i = s_i \dots s_1 \psi_i$,
- (ii)

$$x_i = y_i + \beta_i^\vee(\psi_i(T)) = \beta_i^\vee(\mathcal{Q}_{w_0} \psi(T) - \sum_{j=1}^{i-1} y_j \beta_j) - y_i,$$

- (iii)

$$y_i = x_i + \alpha_i^\vee(\eta_i(T)) = \alpha_i^\vee(\mathcal{P}_{w_0} \eta(T) - \sum_{j=1}^{i-1} x_j \alpha_j) - x_i.$$

PROOF. We prove (i) by induction on $i \leq q$. For $i = q$ it holds because $\eta_q = \eta = w_0 \psi = w_0 \psi_q$ and $s_q \dots s_1 = w_0$. Note that, for each $i \leq q$, we can write

$$\mathcal{Q}_{\beta_i} = \mathcal{P}_{\beta_i} s_{\beta_i} = s_1 \dots s_{i-1} \mathcal{P}_{\alpha_i} s_i \dots s_1.$$

Therefore, assuming the induction hypothesis $\eta_i = s_i \dots s_1 \psi_i$,

$$\begin{aligned} \eta_{i-1} &= \mathcal{P}_{\alpha_i} \eta_i = \mathcal{P}_{\alpha_i} s_i \dots s_1 \psi_i \\ &= s_{i-1} \dots s_1 \mathcal{Q}_{\beta_i} \psi_i \\ &= s_{i-1} \dots s_1 \psi_{i-1}, \end{aligned}$$

as required. This implies (ii), using $\eta_{i-1}(T) = \eta_i(T) + x_i \alpha_i$ and $\psi_{i-1}(T) = \psi_i(T) + y_i \beta_i$:

$$\begin{aligned} 2x_i &= \alpha_i^\vee(\eta_{i-1}(T) - \eta_i(T)) \\ &= \alpha_i^\vee(s_{i-1} \dots s_1 \psi_{i-1}(T) - s_i \dots s_1 \psi_i(T)) \\ &= \alpha_i^\vee(s_{i-1} \dots s_1 (\psi_i(T) + y_i \beta_i) - s_i \dots s_1 \psi_i(T)) \\ &= 2y_i + \alpha_i^\vee(\alpha_i^\vee(s_{i-1} \dots s_1 \psi_i(T)) \alpha_i) \\ &= 2y_i + 2\beta_i^\vee(\psi_i(T)). \end{aligned}$$

Finally, (iii) follows immediately from (ii) and (i). \square

By (iii) of the preceding theorem, we can define a mapping $F : M_{\mathbf{i}} \rightarrow \mathbb{R}_+^q \times C$ such that

$$(\mathfrak{G}(\psi), \mathcal{Q}_{w_0}\psi(T)) = F(\varrho_{\mathbf{i}}(\eta), \mathcal{P}_{w_0}\eta(T)).$$

Let $L_{\mathbf{i}} = F(M_{\mathbf{i}})$. It follows from (ii) that $F^{-1}(y, \lambda) = (G(y, \lambda), \lambda)$, where

$$G(y, \lambda) = \beta_i^\vee(\lambda - \sum_{j=1}^{i-1} y_j \beta_j) - y_i.$$

Thus, $L_{\mathbf{i}}$ is the set of $(y, \lambda) \in \mathbb{R}_+^q \times C$ which satisfy

$$(5.8) \quad 0 \leq y_i \leq \beta_i^\vee(\lambda - \sum_{j=1}^{i-1} y_j \beta_j) \quad (i \leq q)$$

and

$$(5.9) \quad \Psi_{\alpha}^{\mathbf{i}}(G(y, \lambda)) \geq 0 \quad \alpha \in \Sigma.$$

We can interpret the last inequality as follows. Set

$$(5.10) \quad \Phi_{\alpha}^{\mathbf{i}}(y, \lambda) = \alpha^\vee(\lambda) - \Psi_{\alpha}^{\mathbf{i}}(G(y, \lambda)).$$

For each simple root α choose $\mathbf{i}_{\alpha} = (s_1^{\alpha}, \dots, s_q^{\alpha})$ such that $s_1^{\alpha} = s_{\alpha}$ and write $\varsigma_{\mathbf{i}_{\alpha}}(\psi) = (y_1^{\alpha}, \dots, y_q^{\alpha})$. Denote the corresponding ψ_i 's by ψ_i^{α} . It follows from lemma 5.7 that $y_1^{\alpha} = \Phi_{\alpha}^{\mathbf{i}}(y, \mathcal{Q}_{w_0}\psi(T))$. To see that the inequality (5.9) holds directly in this context, observe that, if $\mathcal{Q}_{w_0}\psi(T) = \lambda$ then, for each simple α ,

$$\Psi_{\alpha}^{\mathbf{i}}(G(y, \lambda)) = \alpha^\vee(\lambda) - \Phi_{\alpha}^{\mathbf{i}}(y, \lambda) = y_1^{\alpha} + \alpha^\vee(\psi_1^{\alpha}(T)) \geq 0.$$

As in [2], we extend the definition of \mathcal{Q}_{β} to two-sided paths. Denote by $C_{\mathbb{R}}^0(V)$ the set of continuous paths $\pi : \mathbb{R} \rightarrow V$ such that $\pi(0) = 0$ and $\alpha^\vee(\pi(t)) \rightarrow \pm\infty$ as $t \rightarrow \pm\infty$ for all simple α . For $\pi \in C_{\mathbb{R}}^0(V)$ and β a positive root, define $\mathcal{Q}_{\beta}\pi$ by

$$\mathcal{Q}_{\beta}\pi(t) = \pi(t) + [\omega(t) - \omega(0)]\beta,$$

where

$$\omega(t) = - \inf_{t \geq s > -\infty} \beta^\vee(\pi(t) - \pi(s)).$$

It is easy to see that $\mathcal{Q}_{\beta}\pi \in C_{\mathbb{R}}^0(V)$. Thus, we can set $\pi_q = \pi$ and, for $i \leq q$,

$$\pi_{i-1} = \mathcal{Q}_{\beta_i} \dots \mathcal{Q}_{\beta_q} \pi \quad \omega_i(t) = - \inf_{s \leq t} \beta_i^\vee(\pi_i(t) - \pi_i(s)).$$

Then

$$\pi_0 = \mathcal{Q}_{w_0}\pi := \mathcal{Q}_{\beta_1} \dots \mathcal{Q}_{\beta_q} \pi$$

and, for each $i \leq q$,

$$\mathcal{Q}_{w_0}\pi(t) = \pi_i(t) + \sum_{j=1}^i [\omega_j(t) - \omega_j(0)]\beta_j.$$

The following is an immediate consequence of [2, Lemma 5.5].

LEMMA 5.8. *If $\mathcal{Q}_{w_0}\pi(t) = \lambda$ and $\omega(t) = y$ then*

$$\inf_{u \geq t} \alpha^\vee(\mathcal{Q}_{w_0}\pi(u)) = \alpha^\vee(\lambda) - \Phi_{\alpha}^{\mathbf{i}}(y, \lambda).$$

Introduce a probability measure \mathbb{P}_μ under which π is a two-sided Brownian motion in V with drift $\mu \in C$, and set

$$\psi = (\pi(t), t \geq 0).$$

Set $\omega(t) = (\omega_1(t), \dots, \omega_q(t))$.

PROPOSITION 5.9. *Under \mathbb{P}_μ , the following statements hold:*

- (1) $\mathcal{Q}_{w_0}\pi$ has the same law as π .
- (2) For each $T \in \mathbb{R}$, the random variables $\omega_1(T), \dots, \omega_q(T)$ are mutually independent and exponentially distributed with respective parameters $2\beta_1^\vee(\mu), \dots, 2\beta_q^\vee(\mu)$.
- (3) For each $T \in \mathbb{R}$, $\omega(T)$ is independent of $(\mathcal{Q}_{w_0}\pi(s), -\infty < s \leq T)$.
- (4) The random variables $\inf_{u \geq 0} \alpha^\vee(\mathcal{Q}_{w_0}\pi(u))$, α a simple root, are independent of the σ -algebra generated by $(\pi(t), t \geq 0)$.

PROOF. We see by backward induction on $k = q, \dots, 1$ that $\mathcal{Q}_{\beta_k} \cdots \mathcal{Q}_{\beta_q} \pi(s)$, $s \leq T$ has the same distribution as $\mathcal{Q}_{\beta_{k-1}} \cdots \mathcal{Q}_{\beta_q} \pi(s)$, $s \leq T$ and is independent of $\omega_k(T)$ and that $\omega_k(T)$ has an exponential distribution with parameter $2\beta_k^\vee(\mu)$. At each step, this is a one dimensional statement which can be checked directly or as a consequence of the classical Burke output theorem for the single queue, see [30]. This implies that (1), (2), and (3) hold. Moreover

$$\inf_{t \geq 0} \beta_1^\vee(\mathcal{Q}_{w_0}\pi(t)) = - \inf_{s \leq 0} \beta_1^\vee(\mathcal{Q}_{\beta_2} \cdots \mathcal{Q}_{\beta_q} \pi(s))$$

is independent of $\pi(t)$, $t \geq 0$. Since β_1 can be chosen as any simple root α , this proves (4). \square

For $\xi \in C$, denote by E_ξ the event that $\mathcal{Q}_{w_0}\pi(s) \in C - \xi$ for all $s \geq 0$ and by $E_{\xi,T}$ the event that $\mathcal{Q}_{w_0}\pi(s) \in C - \xi$ for all $T \geq s \geq 0$. By proposition 5.9, E_ξ is independent of ψ .

For $r > 0$, define

$$B(\lambda, r) = \{\zeta \in V : \|\zeta - \lambda\| < r\}$$

and

$$R(z, r) = (z_1 - r, z_1 + r) \times \cdots \times (z_q - r, z_q + r).$$

Fix (z, λ) in the interior of L_i and choose $\epsilon > 0$ sufficiently small so that $R(z, \epsilon)$ is contained in $L_i \times B(\lambda, \epsilon)$ and

$$(5.11) \quad \inf_{\lambda' \in B(\lambda, \epsilon), z' \in R(z, \epsilon)} \alpha^\vee(\lambda') - \Phi_\alpha^i(z', \lambda') \geq 0.$$

LEMMA 5.10.

$$\begin{aligned} & \mathbb{P}_\mu(\mathcal{Q}_{w_0}\psi(T) \in B(\lambda, \epsilon), \varsigma_i(\psi) \in R(z, \epsilon)) \\ &= \lim_{C \ni \xi \rightarrow 0} \mathbb{P}_\mu(E_\xi)^{-1} \mathbb{P}_\mu(\mathcal{Q}_{w_0}\pi(T) \in B(\lambda, \epsilon), \omega(T) \in R(z, \epsilon), E_{\xi,T}). \end{aligned}$$

PROOF. An elementary induction argument on the recursive construction of \mathcal{Q}_{w_0} shows that, on the event E_ξ , there is a constant C for which

$$\max_{i \leq q} \|y_i - \omega_i(T)\| \vee \|\mathcal{Q}_{w_0}\psi(T) - \mathcal{Q}_{w_0}\pi(T)\| \leq C\|\xi\|.$$

Hence, for ξ sufficiently small,

$$\begin{aligned} & \mathbb{P}_\mu(\mathcal{Q}_{w_0}\psi(T) \in B(\lambda, \epsilon - C\|\xi\|), \varsigma_i(\psi) \in R(z, \epsilon - C\|\xi\|), E_\xi) \\ & \leq \mathbb{P}_\mu(\mathcal{Q}_{w_0}\pi(T) \in B(\lambda, \epsilon), \omega(T) \in R(z, \epsilon), E_\xi) \\ & \leq \mathbb{P}_\mu(\mathcal{Q}_{w_0}\psi(T) \in B(\lambda, \epsilon + C\|\xi\|), \varsigma_i(\psi) \in R(z, \epsilon + C\|\xi\|), E_\xi). \end{aligned}$$

Now E_ξ is independent of ψ , and so

$$\begin{aligned} & \mathbb{P}_\mu(\mathcal{Q}_{w_0}\psi(T) \in B(\lambda, \epsilon - C\|\xi\|), \varsigma_1(\psi) \in R(z, \epsilon - C\|\xi\|)) \\ & \leq \mathbb{P}_\mu(E_\xi)^{-1} \mathbb{P}_\mu(\mathcal{Q}_{w_0}\pi(T) \in B(\lambda, \epsilon), \omega(T) \in R(z, \epsilon), E_\xi) \\ & \leq \mathbb{P}_\mu(\mathcal{Q}_{w_0}\psi(T) \in B(\lambda, \epsilon + C\|\xi\|), \varsigma_1(\psi) \in R(z, \epsilon + C\|\xi\|)). \end{aligned}$$

Letting $\xi \rightarrow 0$, we obtain that

$$(5.12) \quad \begin{aligned} & \mathbb{P}_\mu(\mathcal{Q}_{w_0}\psi(T) \in B(\lambda, \epsilon), \varsigma_1(\psi) \in R(z, \epsilon)) \\ & = \lim_{C\exists\xi\rightarrow 0} \mathbb{P}_\mu(E_\xi)^{-1} \mathbb{P}_\mu(\mathcal{Q}_{w_0}\pi(T) \in B(\lambda, \epsilon), \omega(T) \in R(z, \epsilon), E_\xi). \end{aligned}$$

Finally observe that, on the event

$$\{\mathcal{Q}_{w_0}\pi(T) \in B(\lambda, \epsilon), \omega(T) \in R(z, \epsilon)\},$$

we have, by Lemma 5.8 and (5.11),

$$\begin{aligned} \inf_{u \geq T} \alpha^\vee(\mathcal{Q}_{w_0}\pi(u)) &= \alpha^\vee(\mathcal{Q}_{w_0}\pi(T)) - \Phi_\alpha^1(\omega(T), \mathcal{Q}_{w_0}\pi(T)) \\ &\geq \inf_{\lambda' \in B(\lambda, \epsilon), z' \in R(z, \epsilon)} \alpha^\vee(\lambda') - \Phi_\alpha^1(z', \lambda') \geq 0. \end{aligned}$$

Thus, we can replace E_ξ by $E_{\xi, T}$ on the right hand side of (5.12), and this concludes the proof of the lemma. \square

For $a, b \in C$, define $\phi(a, b) = \sum_{w \in W} \varepsilon(w) e^{\langle wa, b \rangle}$.

LEMMA 5.11. *Fix $\mu \in C$. The functions $f(a, b) = \phi(a, b)/[h(a)h(b)]$ and $g_\mu(a, b) = \phi(a, b)/\phi(a, \mu)$ have unique analytic extensions to $V \times V$. Moreover, $f(0, b) = k^{-1}$ and $g_\mu(0, b) = h(b)/h(\mu)$.*

PROOF. It is clear that the function ϕ is analytic in (a, b) , furthermore it vanishes on the hyperplanes $\langle \beta, a \rangle = 0, \langle \beta, b \rangle = 0$, for all roots β . The first claim follows from an elementary analytic functions argument. In the expansion of ϕ as an entire function, the term of homogeneous degree d is a polynomial in a, b which is antisymmetric under W , therefore a multiple of $h(a)h(b)$. In particular the term of lowest degree is a constant multiple of $h(a)h(b)$. This constant is nonzero, as can be seen by taking derivatives in the definition of ϕ . By l'Hôpital's rule, $\lim_{a \rightarrow 0} g_\mu(a, b) = h(b)/h(\mu)$. It follows that $\lim_{a \rightarrow 0} f(a, b)$ is a constant. To evaluate this constant, note that, since h is harmonic and vanishes at the boundary of C ,

$$\int_C h(\lambda)^2 e^{-\|\lambda\|^2/2} f(a, \lambda) d\lambda = e^{|a|^2/2} \int_V e^{-\|\lambda\|^2/2} d\lambda.$$

Letting $a \rightarrow 0$, we deduce that $f(0, \lambda) = k^{-1}$, as required. \square

Denote by F_ξ the event that $\psi(s) \in C - \xi$ for all $s \geq 0$ and by $F_{\xi, T}$ the event that $\psi(s) \in C - \xi$ for all $T \geq s \geq 0$.

LEMMA 5.12. *For $B \subset C$, bounded and measurable,*

$$\begin{aligned} & \lim_{C\exists\xi\rightarrow 0} \mathbb{P}_\mu(F_\xi)^{-1} \mathbb{P}_\mu(\psi(T) \in B, F_{\xi, T}) \\ &= c_T^{-1} h(\mu)^{-1} \int_B e^{\langle \mu, \lambda' \rangle - \|\mu\|^2 T/2} e^{-\|\lambda'\|^2/2 T} h(\lambda') d\lambda'. \end{aligned}$$

PROOF. Set $z_T = \int_V e^{-\|\lambda\|^2/2T} d\lambda$. By the reflection principle,

$$\mathbb{P}_\mu(\psi(T) \in d\lambda, F_{\xi,T}) = e^{\langle \mu, \lambda \rangle - \|\mu\|^2 T/2} \sum_{w \in W} \varepsilon(w) p_T(w\xi, \xi + \lambda) d\lambda,$$

where $p_t(a, b) = z_t^{-1} e^{-\|b-a\|^2/2t}$ is the transition density of a standard Brownian motion in V . Integrating over λ and letting $T \rightarrow \infty$, we obtain (see [2])

$$\mathbb{P}_\mu(F_\xi) = \sum_{w \in W} \varepsilon(w) e^{\langle w\xi - \xi, \mu \rangle}.$$

Thus, using lemma 5.11 and the bounded convergence theorem,

$$\begin{aligned} & \lim_{C \ni \xi \rightarrow 0} \mathbb{P}_\mu(F_\xi)^{-1} \mathbb{P}_\mu(\psi(T) \in B, F_{\xi,T}) \\ &= z_T^{-1} \lim_{C \ni \xi \rightarrow 0} \int_B e^{\langle \mu, \lambda' \rangle - \|\mu\|^2 T/2} e^{-(|\xi|^2 + |\xi + \lambda'|^2)/2T} \phi(\xi, \mu)^{-1} \phi\left(\xi, \frac{\xi + \lambda'}{T}\right) d\lambda' \\ &= z_T^{-1} \lim_{C \ni \xi \rightarrow 0} \int_B e^{\langle \mu, \lambda' \rangle - \|\mu\|^2 T/2} e^{-(\|\xi\|^2 + \|\xi + \lambda'\|^2)/2T} g_\mu\left(\xi, \frac{\xi + \lambda'}{T}\right) d\lambda' \\ &= z_T^{-1} h(\mu)^{-1} \int_B e^{\langle \mu, \lambda' \rangle - \|\mu\|^2 T/2} e^{-|\lambda'|^2/2T} h(\lambda'/T) d\lambda' \\ &= c_T^{-1} h(\mu)^{-1} \int_B e^{\langle \mu, \lambda' \rangle - \|\mu\|^2 T/2} e^{-\|\lambda'\|^2/2T} h(\lambda') d\lambda', \end{aligned}$$

as required. □

Applying lemmas 5.10, 5.12 and proposition 5.9, we obtain

$$\begin{aligned} & \mathbb{P}_\mu(\mathcal{Q}_{w_0}\psi(T) \in B(\lambda, \epsilon), \mathfrak{g}_1(\psi) \in R(z, \epsilon)) \\ &= \lim_{C \ni \xi \rightarrow 0} \mathbb{P}_\mu(E_\xi)^{-1} \mathbb{P}_\mu(\mathcal{Q}_{w_0}\pi(T) \in B(\lambda, \epsilon), \omega(T) \in R(z, \epsilon), E_{\xi,T}) \quad (\text{lemma 5.11}) \\ &= \lim_{C \ni \xi \rightarrow 0} \mathbb{P}_\mu(E_\xi)^{-1} \mathbb{P}_\mu(\omega(T) \in R(z, \epsilon)) \mathbb{P}_\mu(\mathcal{Q}_{w_0}\pi(T) \in B(\lambda, \epsilon), E_{\xi,T}) \quad (\text{lemma 5.9(3)}) \\ &= \lim_{C \ni \xi \rightarrow 0} \mathbb{P}_\mu(E_\xi)^{-1} \mathbb{P}_\mu(\omega(T) \in R(z, \epsilon)) \mathbb{P}_\mu(\psi(T) \in B(\lambda, \epsilon), F_{\xi,T}) \\ &= \prod_{i=1}^q e^{-\beta_i^\vee(\mu)z_i} [e^{\epsilon\beta_i^\vee(\mu)} - e^{-\epsilon\beta_i^\vee(\mu)}] \lim_{C \ni \xi \rightarrow 0} \mathbb{P}_\mu(E_\xi)^{-1} \mathbb{P}_\mu(\psi(T) \in B(\lambda, \epsilon), F_{\xi,T}) \\ &\quad (\text{lemma 5.9 (2)}) \\ &= \prod_{i=1}^q e^{-\beta_i^\vee(\mu)z_i} [e^{\epsilon\beta_i^\vee(\mu)} - e^{-\epsilon\beta_i^\vee(\mu)}] \\ &\quad \times c_T^{-1} h(\mu)^{-1} \int_{B_V(\lambda, \epsilon)} e^{\mu(\lambda') - \|\mu\|^2 T/2} e^{-\|\lambda'\|^2/2T} h(\lambda') d\lambda'. \quad (\text{lemma 5.12}) \end{aligned}$$

Now divide by $\|B(y, \epsilon)\|(2\epsilon)^q$ and let ϵ tend to zero to obtain

$$\begin{aligned} & \mathbb{P}_\mu(\mathcal{Q}_{w_0}\psi(T) \in d\lambda, \mathfrak{g}_1(\psi) \in dz) \\ &= \prod_{i=1}^q e^{-\beta_i^\vee(\mu)z_i} e^{\langle \mu, \lambda \rangle - \|\mu\|^2 T/2} c_T^{-1} h(\lambda) e^{-\|\lambda\|^2/2T} d\lambda dz. \end{aligned}$$

Letting $\mu \rightarrow 0$ this becomes, writing $\mathbb{P} = \mathbb{P}_0$,

$$(5.13) \quad \mathbb{P}(\mathcal{Q}_{w_0}\psi(T) \in d\lambda, \mathfrak{g}_1(\psi) \in dz) = c_T^{-1} h(\lambda) e^{-\|\lambda\|^2/2T} d\lambda dz.$$

Using lemma 5.7, it follows that, for (w, λ) in the interior of M_i ,

$$(5.14) \quad \mathbb{P}(\varrho_i(\eta) \in dw, \mathcal{P}_{w_0}\eta(T) \in d\lambda) = c_T^{-1} h(\lambda) e^{-\|\lambda\|^2/2T} dw d\lambda.$$

Under the probability measure \mathbb{P} , η is a standard Brownian motion in V with transition density given by $p_t(a, b) = z_t^{-1} e^{-\|b-a\|^2/2t}$. By theorem 5.1 under \mathbb{P} , $\mathcal{P}_{w_0}\eta$ is a Brownian motion in C . Its transition density is given, for $\xi, \lambda \in C$, by

$$q_t(\xi, \lambda) = \frac{h(\lambda)}{h(\xi)} \sum_{w \in W} \varepsilon(w) p_t(w\xi, \lambda).$$

As remarked in [2], this transition density can be extended by continuity to the boundary of C . From lemma 5.11 we see that $q_T(0, \lambda) = k^{-1} h(\lambda)^2 e^{-\|\lambda\|^2/2T}$. Thus,

$$(5.15) \quad \mathbb{P}(\mathcal{P}_{w_0}\eta(T) \in d\lambda) = k^{-1} h(\lambda)^2 e^{-\|\lambda\|^2/2T} d\lambda.$$

To complete the proof of the theorem, first note that since $\varsigma_i(\psi)$ is measurable with respect to the σ -algebra generated by $(\mathcal{Q}_{w_0}\psi(u), u \geq T)$, $\varrho_i(\eta)$ is measurable with respect to the σ -algebra generated by $(\mathcal{P}_{w_0}\eta(u), u \geq T)$. Thus, by the Markov property of $\mathcal{P}_{w_0}\eta$, the conditional distribution of $\varrho_i(\eta)$, given $(\mathcal{P}_{w_0}\eta(s), s \leq T)$, is measurable with respect to the σ -algebra generated by $\mathcal{P}_{w_0}\eta(T)$. Combining this with (5.14) and (5.15), we conclude that the conditional law of $\varrho_i(\eta)$, given $(\mathcal{P}_{w_0}\eta(s), s \leq T)$ and $\mathcal{P}_{w_0}\eta(T) = \lambda$, is almost surely uniform on M_i^λ , and that the Euclidean volume of M_i^λ is $k^{-1} h(\lambda)$, as required.

5.5. Proof of theorem 5.5. Let $\psi = w_0\eta$ and $\mathcal{Q}_{w_0} = \mathcal{P}_{w_0}w_0$. Denote by P_t (respectively Q_t) the semigroup of Brownian motion in V (respectively C). Under \mathbb{P} , by [2, Theorem 5.6], $\mathcal{Q}_{w_0}\psi$ is a Brownian motion in C . Let $\delta \in C$. The function $e_\delta(v) = e^{\langle \delta, v \rangle}$ is an eigenfunction of P_t and the e_δ -transform of P_t is a Brownian motion with drift δ . Setting $\phi_\delta(v) = \sum_{w \in W} \varepsilon(w) e^{\langle w\delta, v \rangle}$, the function ϕ_δ/h is an eigenfunction of Q_t and the (ϕ_δ/h) -transform of Q_t is a Brownian motion with drift δ conditioned never to exit C (see [2, Section 5.2] for a definition of this process). By theorem 5.2, the conditional law of $\eta(T)$, given $(\mathcal{P}_{w_0}\eta(s), s \leq T)$ and $\mathcal{P}_{w_0}\eta(T) = \lambda$, is almost surely given by μ_{DH}^λ . It follows that the conditional law of $\psi(T)$, given $(\mathcal{Q}_{w_0}\psi(s), s \leq T)$ and $\mathcal{Q}_{w_0}\psi(T) = \lambda$, is almost surely given by μ_{DH}^λ . Denote the corresponding Markov operator by $K(\lambda, \cdot) = \mu_{DH}^\lambda(\cdot)$. By [2, Theorem 5.6] we automatically have the intertwining $KP_t = Q_tK$. Note that Ke_δ is an eigenfunction of Q_t . By construction, the Ke_δ -transform of Q_t , started from the origin, has the same law as $\mathcal{Q}_{w_0}\psi^{(\delta)}$, where $\psi^{(\delta)}$ is a Brownian motion in V with drift δ . Recalling the proof of [2, Theorem 5.6] we note that $\mathcal{Q}_{w_0}\psi^{(\delta)}$ has the same law as a Brownian motion with drift δ conditioned never to exit C . It follows that $Ke_\delta = \phi_\delta/(c(\delta)h)$, for some $c(\delta) \neq 0$. Now observe (using lemma 5.11 for example) that $\lim_{\xi \rightarrow 0} Ke_\delta(\xi) = 1$. Thus, by lemma 5.11, $c(\delta) = \lim_{\xi \rightarrow 0} \phi_\delta(\xi)/h(\xi) = k^{-1}h(\delta)$. We conclude that

$$\int_V e^{\langle \delta, v \rangle} \mu_{DH}^\lambda(dv) = k \frac{\sum_{w \in W} \varepsilon(w) e^{\langle w\delta, \lambda \rangle}}{h(\delta)h(\lambda)}.$$

This formula extends to $\delta \in V^*$ by analytic continuation (see lemma 5.11 again), and the proof is complete.

5.6. A Littlewood-Richardson property. Let $\mathbf{i} = (s_1, \dots, s_q)$ where $w_0 = s_1 \dots s_q$ is a reduced decomposition. For $\eta \in C_T^0(V)$, let $x = \rho_{\mathbf{i}}(\eta)$.

For each simple root α choose now $\mathbf{j}_\alpha = (s_1^\alpha, \dots, s_q^\alpha)$, a reduced decomposition of w_0 , such that $s_q^\alpha = s_\alpha$, and denote the corresponding string parameters of the path η by $(\tilde{x}_1^\alpha, \dots, \tilde{x}_q^\alpha) = \varrho_{\mathbf{j}_\alpha}(\eta)$. As in (5.2), there is a continuous function $\Psi'_\alpha : \mathbb{R}^q \rightarrow \mathbb{R}$ such that $\tilde{x}_q^\alpha = \Psi'_\alpha(x)$. Fix $\lambda, \mu \in C$ and suppose that $\lambda + \eta(s) \in C$ for $0 \leq s \leq T$. Then $\tilde{x}_q^\alpha = -\inf_{s \leq T} \alpha^\vee(\eta(s)) \leq \alpha^\vee(\lambda)$. In other words,

$$(5.16) \quad \Psi'_\alpha(x) \leq \alpha^\vee(\lambda), \quad \alpha \in \Sigma.$$

Let $M_{\mathbf{i}}^{\lambda, \mu}$ denote the set of $x \in M_{\mathbf{i}}^\lambda$ which satisfy the additional constraints (5.16). This is a compact convex polytope. Let $\nu^{\lambda, \mu}$ be the uniform probability distribution on $M_{\mathbf{i}}^{\lambda, \mu}$ and let $\nu_{\lambda, \mu}$ be its image on V by the map

$$x = (x_1, \dots, x_q) \in M_{\mathbf{i}}^{\lambda, \mu} \mapsto \lambda + \mu - \sum_{j=1}^q x_j \alpha_j \in V.$$

Let η be the Brownian motion in V starting from 0. Observe that, by theorem 3.9, the event $\{\eta(s) \in C - \lambda, 0 \leq s \leq T\}$ is measurable with respect to the σ -algebra generated by $\rho_{\mathbf{i}}(\eta)$. Combining this with theorem 5.2 we obtain:

COROLLARY 5.13. *The conditional law of $\rho_{\mathbf{i}}(\eta)$, given $\mathcal{P}_{w_0}\eta(s), s \leq T, \mathcal{P}_{w_0}\eta(T) = \mu$ and $\lambda + \eta(s) \in C$ for $0 \leq s \leq T$, is $\nu^{\lambda, \mu}$ and the conditional law of $\lambda + \eta(T)$ is $\nu_{\lambda, \mu}$.*

For $s, t \geq 0$ let

$$(\tau_s \eta)(t) = \eta(s+t) - \eta(s), (\tau_s \mathcal{P}_{w_0} \eta)(t) = \mathcal{P}_{w_0} \eta(s+t) - \mathcal{P}_{w_0} \eta(s).$$

LEMMA 5.14. *For all $s \geq 0$,*

$$\mathcal{P}_{w_0}(\tau_s \mathcal{P}_{w_0} \eta) = \mathcal{P}_{w_0} \tau_s \eta.$$

PROOF. If $\pi_1, \pi_2 : \mathbb{R}^+ \rightarrow V$ are continuous path starting at 0, let $\pi_1 \star_s \pi_2$ be the path defined by $\pi_1 \star_s \pi_2(r) = \pi_1(r)$ when $0 \leq r \leq s$ and $\pi_1 \star_s \pi_2(r) = \pi_1(s) + \pi_2(r-s)$ when $s \leq r$. By lemma 4.16, $\mathcal{P}_{w_0}(\pi_1 \star_s \pi_2) = \mathcal{P}_{w_0}(\pi_1) \star_s \tilde{\pi}_2$ where $\tilde{\pi}_2$ is a path such that $\mathcal{P}_{w_0}(\tilde{\pi}_2) = \mathcal{P}_{w_0}(\pi_2)$. Since $\tau_s(\pi_1 \star_s \pi_2) = \pi_2$, this gives the lemma. \square

Let $\gamma_{\lambda, \mu}$ be the measure on C given by

$$\gamma_{\lambda, \mu}(dx) = \frac{h(x)}{h(\lambda)} \nu_{\lambda, \mu}(dx).$$

It will follow from theorem 5.15 that this is a probability measure. Consider the following σ -algebra

$$\mathcal{G}_{s, t} = \sigma(\mathcal{P}_{w_0} \eta(a), a \leq s, \mathcal{P}_{w_0} \tau_s \eta(r), r \leq t).$$

The following result is a continuous analogue of the Littelmann interpretation of the Littlewood-Richardson decomposition of a tensor product.

THEOREM 5.15. *For $s, t > 0$, $\gamma_{\lambda, \mu}$ is the conditional distribution of $\mathcal{P}_{w_0} \eta(s+t)$ given $\mathcal{G}_{s, t}$, $\mathcal{P}_{w_0} \eta(s) = \lambda$ and $\mathcal{P}_{w_0} \tau_s \eta(t) = \mu$.*

PROOF. When $(X_t, (\theta_t), P_x)$ is a Markov process with shift θ_t (i.e. $X_{s+t} = X_s \circ \theta_t$), for any $\sigma(X_r, r \geq 0)$ -measurable random variables $Z, Y \geq 0$, one has

$$\mathbb{E}(Z \circ \theta_t | \sigma(X_s, s \leq t, Y \circ \theta_t)) = \mathbb{E}_{X_0}(Z | \sigma(Y)) \circ \theta_t.$$

Let us apply this relation to the Markov process $X = \mathcal{P}_{w_0}\eta$ (see [2]). Notice that since $\mathcal{P}_{w_0}(\tau_s X) = \mathcal{P}_{w_0}(\tau_0 X) \circ \theta_s$, it follows from the lemma that

$$\mathcal{G}_{s,t} = \sigma(X_a, \mathcal{P}_{w_0}(\tau_0 X)(r) \circ \theta_s, a \leq s, r \leq t).$$

Therefore, for any Borel nonnegative function $f : V \rightarrow \mathbb{R}$,

$$\mathbb{E}[f(\mathcal{P}_{w_0}\eta(s+t) | \mathcal{G}_{s,t})] = \mathbb{E}_{X_0}[f(X_t) | \sigma(\mathcal{P}_{w_0}(\tau_0 X)(r), r \leq t)] \circ \theta_s.$$

One knows (Theorem 5.1 in [2]) that X is the h -process of the Brownian motion killed at the boundary of C . In other words, starting from $X_0 = \lambda$, X is the h -process of $\lambda + \eta(t)$ conditionally on $\lambda + \eta(s) \in C$, for $0 \leq s \leq t$. It thus follows from corollary 5.13 that

$$\mathbb{E}_\lambda[f(X_t) | \sigma(\mathcal{P}_{w_0}(\tau_0 X)(r), r \leq t)] = \frac{1}{h(\lambda)} \int f(x) h(x) d\nu_{\lambda,\mu}(x)$$

when $\mathcal{P}_{w_0}(\tau_0 X)(t) = \mu$. This proves that

$$\mathbb{E}[f(\mathcal{P}_{w_0}\eta(s+t) | \mathcal{G}_{s,t})] = \int f(x) d\mu_{\lambda,\mu}(x)$$

when $\mathcal{P}_{w_0}\eta(s) = \lambda$ and $\mathcal{P}_{w_0}\tau_s\eta(t) = \mu$.

5.7. A product formula. Consider the Laplace transform of μ_{DH}^λ given, for $\lambda \in C, z \in V^*$, by

$$(5.17) \quad J_\lambda(z) = k \frac{\sum_W \varepsilon(w) e^{\langle z, w\lambda \rangle}}{h(z)h(\lambda)}.$$

This is an example of a generalized Bessel function, following the terminology of Helgason [13] in the Weyl group case and Opdam [31] in the general Coxeter case. It was a conjecture in Gross and Richards [10] that these are Laplace transform of positive measures (this also follows from Rösler [33]). They are positive eigenfunctions of the Laplace and of the Dunkl operators on the Weyl chamber C with eigenvalue $\|\lambda\|^2$ and Dirichlet boundary conditions and $J_\lambda(0) = 1$. Let f_λ be the density of the probability measure μ_{DH}^λ . One has

$$(5.18) \quad \int_V e^{\langle z, v \rangle} f_\lambda(v) dv = J_\lambda(z).$$

Let, for $v \in C$,

$$f_{\lambda,\mu}(v) = \frac{1}{h(\mu)} \sum_{w \in W} h(wv) f_\lambda(wv - \mu).$$

It follows from the next result that $f_{\lambda,\mu}(v) \geq 0$.

THEOREM 5.16. (i) For $\lambda, \mu \in C$ and $z \in V^*$,

$$J_\lambda(z) J_\mu(z) = \int_C J_v(z) f_{\lambda,\mu}(v) dv.$$

(ii)

$$\gamma_{\lambda,\mu}(dx) = f_{\lambda,\mu}(x) dx.$$

PROOF. The first part is given by the following computation, similar to the one in Dooley et al [6], we give it for the convenience of the reader. It follows from (5.17) and (5.18) that

$$J_\lambda(z)J_\mu(z) = \int_V e^{\langle z, v \rangle} J_\mu(z) f_\lambda(v) dv = k \sum_W \varepsilon(w) \int_V \frac{e^{\langle z, w\mu+v \rangle}}{h(\mu)h(z)} f_\lambda(v) dv.$$

Using the invariance of the measure μ_{DH}^λ under W , $f_\lambda(wv) = f_\lambda(v)$ for $w \in W$. One has

$$\begin{aligned} J_\lambda(z)J_\mu(z) &= k \sum_W \varepsilon(w) \int_V \frac{e^{\langle z, w(\mu+v) \rangle}}{h(\mu)h(z)} f_\lambda(v) dv \\ &= k \sum_W \varepsilon(w) \int_V \frac{e^{\langle z, wv \rangle}}{h(\mu)h(z)} f_\lambda(v - \mu) dv \\ &= \frac{1}{h(\mu)} \int_V J_v(z) h(v) f_\lambda(v - \mu) dv \\ &= \frac{1}{h(\mu)} \sum_{w \in W} \int_{w^{-1}C} J_v(z) h(v) f_\lambda(v - \mu) dv \\ &= \frac{1}{h(\mu)} \sum_{w \in W} \int_C J_v(z) h(wv) f_\lambda(wv - \mu) dv \\ &= \int_C J_z(v) f_{\lambda, \mu}(v) dv \end{aligned}$$

where we have used that, up to a set of measure zero, $V = \cup_{w \in W} w^{-1}C$. This proves (i).

Let us now prove (ii), using theorem 5.15. Since η is a standard Brownian motion in V , $\{\eta(r), r \leq s\}$ and $\tau_s \eta$ are independent, hence, for $z \in V^*$,

$$\begin{aligned} \mathbb{E}(e^{\langle z, \eta(s+t) \rangle} | \mathcal{G}_{s,t}) &= \mathbb{E}(e^{\langle z, \eta(s) \rangle} e^{\langle z, \tau_s \eta(t) \rangle} | \mathcal{G}_{s,t}) \\ &= \mathbb{E}(e^{\langle z, \eta(s) \rangle} | \sigma(\mathcal{P}_{w_0} \eta(a), a \leq s)) \mathbb{E}(e^{\langle z, \tau_s \eta(t) \rangle} | \sigma(\mathcal{P}_{w_0} \tau_s \eta(b), b \leq t)). \end{aligned}$$

By theorem 5.5,

$$J_\lambda(z) = \mathbb{E}(e^{\langle z, \eta(s) \rangle} | \sigma(\mathcal{P}_{w_0} \eta(a), a \leq s))$$

when $\mathcal{P}_{w_0} \eta(s) = \lambda$ and, since $\tau_s \eta$ and η have the same law,

$$J_\mu(z) = \mathbb{E}(e^{\langle z, \tau_s \eta(t) \rangle} | \sigma(\mathcal{P}_{w_0} \tau_s \eta(b), b \leq t))$$

when $\mathcal{P}_{w_0} \tau_s \eta(t) = \mu$. Therefore

$$\mathbb{E}(e^{\langle z, \eta(s+t) \rangle} | \mathcal{G}_{s,t}) = J_\lambda(z)J_\mu(z).$$

On the other hand, by lemma 4.16, $\mathcal{G}_{s,t}$ is contained in $\sigma(\mathcal{P}_{w_0} \eta(r), r \leq s+t)$, thus

$$\begin{aligned} \mathbb{E}(e^{\langle z, \eta(s+t) \rangle} | \mathcal{G}_{s,t}) &= \mathbb{E}(\mathbb{E}(e^{\langle z, \eta(s+t) \rangle} | \sigma(\mathcal{P}_{w_0} \eta(r), r \leq s+t)) | \mathcal{G}_{s,t}) \\ &= \mathbb{E}(J_z(\mathcal{P}_{w_0} \eta(s+t)) | \mathcal{G}_{s,t}). \end{aligned}$$

It thus follows from theorem 5.15 that

$$J_\lambda(z)J_\mu(z) = \int J_v(z) d\gamma_{\lambda, \mu}(v).$$

Therefore, for all $z \in V^*$,

$$\int J_v(z) f_{\lambda, \mu}(v) dv = \int J_v(z) d\gamma_{\lambda, \mu}(v).$$

By injectivity of the Fourier-Laplace, transform this implies that

$$d\gamma_{\lambda,\mu}(v) = f_{\lambda,\mu}(v) dv. \quad \square$$

The positive product formula gives a positive answer to a question of Rösler [34] for the radial Dunkl kernel. It shows that one can generalize the structure of Bessel-Kingman hypergroup to any Weyl chamber, for the so called geometric parameter.

6. Littelmann modules and geometric lifting.

6.1. It was observed some time ago by G. Lusztig that the combinatorics of the canonical basis is closely related to the geometry of the totally positive varieties. This connection was made precise by Berenstein and Zelevinsky in [3], in terms of transformations called "tropicalization" and "geometric lifting". In this section we show how some simple considerations on Sturm-Liouville equations lead to a natural way of lifting Littelmann paths, which take values in a Cartan algebra, to the corresponding Borel group. Using this lift, an application of Laplace's method explains the connection between the canonical basis and the totally positive varieties.

This section is organized as follows. We first recall the notions of tropicalization and geometric lifting in the next subsection, as well as the connection between the totally positive varieties and the canonical basis. Then we make some observations on Sturm-Liouville equations and their relation to Pitman transformations and the Littelmann path model in type A_1 . We extend these observations to higher rank in the next subsections then we show how they explain the link between string parametrization of the canonical basis and the totally positive varieties.

6.2. Tropicalization and geometric lifting. A subtraction free rational expression is a rational function in several variables, with positive real coefficients and without minus sign, e.g.

$$t_1 + 2t_2/t_3, (1 - t^3)/(1 - t) \text{ or } 1/(t_1t_2 + 3t_3t_4)$$

are such expressions, but not $t_1 - t_2$. Any such expression $F(t_1, \dots, t_n)$ can be tropicalized, which means that

$$F_{trop}(x_1, \dots, x_n) = \lim_{\varepsilon \rightarrow 0+} \varepsilon \log(F(e^{x_1/\varepsilon}, \dots, e^{x_n/\varepsilon}))$$

exists as a piecewise linear function of the real variables (x_1, \dots, x_n) , and is given by an expression in the maxplus algebra over the variables x_1, \dots, x_n . More precisely, the tropicalization $F \rightarrow F_{trop}$ replaces each occurrence of $+$ by \vee (the max sign $x \vee y = \max(x, y)$), each product by a $+$, and each fraction by a $-$, and each positive real number by 0. For example the three expressions above give

$$(t_1 + t_2/t_3)_{trop} = x_1 \vee (x_2 - x_3), ((1 - x^3)/(1 - x))_{trop} = 0 \vee x \vee 2x,$$

and

$$(1/(x_1x_2 + 3x_3x_4))_{trop} = -((x_1 + x_2) \vee (x_3 + x_4)).$$

Tropicalization is not a one to one transformation, and there exists in general many subtraction free rational expressions which have the same tropicalization. Given some expression G in the maxplus algebra, any subtraction free rational expression whose tropicalization is G is called a geometric lifting of G , cf [3].

6.3. Double Bruhat cells and string coordinates. We recall some standard terminology, using the notations of [3]. We consider a simply connected complex semisimple Lie group G , associated with a root system R . Let H be a maximal torus, and B, B_- be corresponding opposite Borel subgroups with unipotent radicals N, N_- . Let $\alpha_i, i \in I$, and $\alpha_i^\vee, i \in I$, be the simple positive roots and coroots, and s_i the corresponding reflections in the Weyl group W . Let $e_i, f_i, h_i, i \in I$, be Chevalley generators of the Lie algebra of G . One can choose representatives $\bar{w} \in G$ for $w \in W$ by putting $\bar{s}_i = \exp(-e_i) \exp(f_i) \exp(-e_i)$ and $\bar{v}\bar{w} = \bar{v}\bar{w}$ if $l(v) + l(w) = l(vw)$ (see [9] (1.8), (1.9)). The Lie algebra of H , denoted by \mathfrak{h} has a Cartan decomposition $\mathfrak{h} = \mathfrak{a} + i\mathfrak{a}$ such that the roots α_i take real values on the real vector space \mathfrak{a} . Thus \mathfrak{a} is generated by $\alpha_i^\vee, i \in I$ and its dual \mathfrak{a}^* by $\alpha_i, i \in I$.

A double Bruhat cell is associated with each pair $u, v \in W$ as

$$L^{u,v} = N\bar{u}N \cap B_-\bar{v}B_-.$$

We will be mainly interested here in the double Bruhat cells $L^{w,e}$. As shown in [3], given a reduced decomposition $w = s_{i_1} \dots s_{i_q}$ every element $g \in L^{w,e}$ has a unique decomposition $g = x_{-i_1}(r_1) \dots x_{-i_q}(r_q)$ with non zero complex numbers (r_1, \dots, r_q) , where $x_{-i}(s) = \varphi_i \begin{pmatrix} s & 0 \\ 1 & s^{-1} \end{pmatrix}$ (where φ_i is the embedding of SL_2 into G given by e_i, f_i, h_i). The totally positive part of the double Bruhat cell corresponds to the set of elements with positive real coordinates. For two different reduced decompositions, the transition map between two sets of coordinates of the form (r_1, \dots, r_q) is given by a subtraction free rational map, which is therefore subject to tropicalization.

As a simple example consider the case of type A_2 . Let the coordinates on the double Bruhat cell $L^{w_0,e}$ for the reduced decompositions $w_0 = s_1 s_2 s_1$, and $w_0 = s_2 s_1 s_2$ be (u_1, u_2, u_3) and (t_1, t_2, t_3) respectively, then

$$(6.1) \quad \begin{pmatrix} t_2 & 0 & 0 \\ t_1 & t_1 t_3 / t_2 & 0 \\ 1 & t_3 / t_2 + 1 / t_1 & 1 / t_1 t_3 \end{pmatrix} = \begin{pmatrix} u_1 u_3 & 0 & 0 \\ u_3 + u_2 / u_1 & u_2 / u_1 u_3 & 0 \\ 1 & 1 / u_3 & 1 / u_2 \end{pmatrix}$$

which yields transition maps

$$\begin{aligned} t_1 &= u_3 + u_2 / u_1 \\ t_2 &= u_1 u_3 \\ t_3 &= u_1 u_2 / (u_2 + u_1 u_3). \end{aligned}$$

On the other hand, for each reduced expression $w_0 = s_{i_1} \dots s_{i_q}$ we can consider the parametrization of the canonical basis by means of string coordinates. For any two such reduced decompositions, the transition maps between the two sets of string coordinates are given by piecewise linear expressions. As shown by Berenstein and Zelevinsky, these expressions are the tropicalizations of the transition maps between the two parametrizations of the double Bruhat cell $L^{w_0,e}$, associated to the Langlands dual group. For example, in type A_2 (which is its own Langlands dual) let (x_1, x_2, x_3) be the Kashiwara coordinates of the canonical basis, using the reduced decomposition $w_0 = s_1 s_2 s_1$, and (y_1, y_2, y_3) the ones corresponding to $w_0 = s_2 s_1 s_2$. The transition map between the two is given by

$$\begin{aligned} y_1 &= x_3 \vee (x_2 - x_1) \\ y_2 &= x_1 + x_3 \\ y_3 &= x_1 \wedge (x_2 - x_3) \end{aligned}$$

which is the tropicalization of (6.3).

We will show how some elementary considerations on the Sturm-Liouville equation, and the method of variation of constants, together with the Littelmann path model explain these connections.

6.4. Sturm-Liouville equations. We consider the Sturm-Liouville equation

$$(6.2) \quad \varphi'' + q\varphi = \lambda\varphi$$

on some interval of the real line, say $[0, T]$ to fix notations. In general there exists no closed form for the solution to such an equation. However, if one solution φ_0 is known, which does not vanish in the interval then all the solutions can be found by quadrature. Indeed using for example the "method of variation of constants" one sees that every other solution φ of this equation in the same interval can be written in the form

$$\varphi(t) = u\varphi_0(t) + v\varphi_0(t) \int_0^t \frac{1}{\varphi_0^2(s)} ds$$

for some constants u, v . If this new solution does not vanish in the interval I , we can use it to generate other solutions of the equation by the same kind of formula. This leads us to investigate the composition of two maps of the form

$$E_{u,v} : \varphi \mapsto u\varphi(t) + v\varphi(t) \int_0^t \frac{1}{\varphi^2(s)} ds$$

acting on non vanishing continuous functions. It is easy to see, using integration by parts, that whenever the composition is well defined, one has

$$E_{u,v} \circ E_{u',v'} = E_{uu', uv' + v/u'}$$

therefore these maps define a partial right action of the group of unimodular lower triangular matrices

$$\begin{pmatrix} u & 0 \\ v & u^{-1} \end{pmatrix}$$

on the set of continuous paths which do not vanish in I . Of course this is equivalently a partial left action of the upper triangular group, but for reasons which will soon appear we choose this formulation. In particular if we start from φ and construct

$$\psi(t) = u\varphi(t) + v\varphi(t) \int_0^t \frac{1}{\varphi^2(s)} ds$$

which does not vanish on $[0, T]$, then φ can be recovered from ψ by the formula

$$\varphi(t) = u^{-1}\psi(t) - v\psi(t) \int_0^t \frac{1}{\psi^2(s)} ds.$$

Coming back to the Sturm-Liouville equation, let η, ρ be a fundamental basis of solutions at 0, namely $\eta(0) = \rho'(0) = 1$, $\eta'(0) = \rho(0) = 0$. Then in the two-dimensional space spanned by ρ, η the transformation is given by

$$(x, y) \mapsto (ux, uy + v/x)$$

and it is defined on $x \neq 0$. Again it is easy to check, using this formula, that this defines a right action of the lower triangular group.

Let us now investigate the limiting case $u = 0$, which gives (assuming $v = 1$ for simplicity)

$$(6.3) \quad \mathcal{T}\varphi(t) = \varphi(t) \int_0^t \frac{ds}{\varphi(s)^2}.$$

This map provides a “geometric lifting” of the one-dimensional Pitman transformation. Indeed set $\varphi(t) = e^{a(t)}$, then using Laplace’s method

$$(6.4) \quad \lim_{\varepsilon \rightarrow 0_+} \varepsilon \log \left(e^{a(t)/\varepsilon} \int_0^t e^{-2a(s)/\varepsilon} ds \right) = a(t) - 2 \inf_{0 \leq s \leq t} a(s).$$

This time the function φ cannot be recovered from $\mathcal{T}\varphi$. If we compute the same transformation with $\varphi_v(t) := \varphi(t)(1 + v \int_0^t \frac{1}{\varphi(s)^2} ds)$ we get

$$\begin{aligned} \mathcal{T}\varphi_v(t) &= \varphi_v(t) \int_0^t \frac{1}{\varphi_v(s)^2} ds \\ &= \varphi(t) \left(1 + v \int_0^t \frac{1}{\varphi(s)^2} ds\right) \left(\frac{1}{v} - \frac{1}{v(1 + v \int_0^t \frac{1}{\varphi(s)^2} ds)} \right) \\ &= \varphi(t) \int_0^t \frac{1}{\varphi(s)^2} ds \\ &= \mathcal{T}\varphi(t). \end{aligned}$$

This is of course not surprising, since $\mathcal{T}\varphi$ vanishes at 0, it thus belongs to a one-dimensional subspace of the space of solutions to the Sturm-Liouville equation, and \mathcal{T} is not invertible. In order to recover the function φ from $\psi = \mathcal{T}\varphi$ we thus need to specify some real number. A convenient choice is to impose the value of

$$\xi = \int_0^T \frac{1}{\varphi(s)^2} ds = \frac{\psi(T)}{\varphi(T)}.$$

With this we can compute

$$\int_t^T \frac{1}{\psi(s)^2} ds = \frac{1}{\int_0^t \frac{1}{\varphi(s)^2} ds} - \frac{1}{\int_0^T \frac{1}{\varphi(s)^2} ds} = \frac{\varphi(t)}{\psi(t)} - \frac{1}{\xi}$$

PROPOSITION 6.1. *Assume that $\psi = \mathcal{T}\varphi$ for some nonvanishing φ , then the set $\mathcal{T}^{-1}(\psi)$ can be parametrized by $\xi \in]0, +\infty[$. For each such ξ there exists a unique $\varphi_\xi \in \mathcal{T}^{-1}(\psi)$ such that $\xi = \int_0^T \frac{1}{\varphi_\xi(s)^2} ds$, given by*

$$\varphi_\xi(t) = \psi(t) \left(\frac{1}{\xi} + \int_t^T \frac{1}{\psi(s)^2} ds \right).$$

Identifying the positive halfline with the Weyl chamber for SL_2 , we see that sets of the form $\mathcal{T}^{-1}(\psi)$ are geometric liftings of the Littelmann modules for SL_2 . The formula in the proposition gives a geometric lifting of the operator \mathcal{H}^x since

$$\mathcal{H}^x a(t) = a(t) - x \wedge 2 \inf_{t \leq s \leq T} a(s) = \lim_{\varepsilon \rightarrow 0_+} \varepsilon \log \left(e^{a(t)/\varepsilon} (e^{-x/\varepsilon} + \int_t^T e^{-2a(s)/\varepsilon} ds) \right).$$

We shall now find the geometric liftings of the Littelmann operators. For this we have, knowing an element $\varphi_{\xi_1} \in \mathcal{T}^{-1}(\psi)$, to find the solution corresponding to ξ_2 . Since

$$\varphi_{\xi_i}(t) = \psi(t) \left(\frac{1}{\xi_i} + \int_t^T \frac{1}{\psi(s)^2} ds \right) \quad i = 1, 2$$

one has

$$\varphi_{\xi_1} = \varphi_{\xi_2} + \psi\left(\frac{1}{\xi_1} - \frac{1}{\xi_2}\right) = \varphi_{\xi_2} \left(1 + \left(\frac{1}{\xi_1} - \frac{1}{\xi_2}\right) \int_0^t \frac{1}{\varphi_{\xi_2}(s)^2} ds\right)$$

Using Laplace method again one can recover the formula for the operators \mathcal{E}_α^x , see definition 3.3.

6.5. A 2×2 matrix interpretation. We shall now recast the above computations using a 2×2 matrix differential equation of order one, and the Gauss decomposition of matrices. This will allow us in the next section to extend these constructions to higher rank groups.

Let N_+ be the nilpotent group of upper triangular invertible 2×2 matrices, let N_- be the corresponding group of lower triangular matrices, and A the group of diagonal matrices, then an invertible 2×2 matrix g has a Gauss decomposition if it can be written as $g = [g]_- [g]_0 [g]_+$ with $[g]_- \in N_-$, $[g]_0 \in A$ and $[g]_+ \in N_+$. We will use also the decomposition $g = [g]_- [g]_{0+}$ with $[g]_{0+} = [g]_0 [g]_+ \in B = AN_+$. The condition for such a decomposition to exist is exactly that the upper left coefficient of the matrix g be non zero.

Let us consider a smooth path $a : [0, T] \rightarrow \mathbf{R}$, such that $a(0) = 0$, and let the matrix $b(t)$ be the solution to

$$(6.5) \quad \frac{db}{dt} = \begin{pmatrix} \frac{da}{dt} & 1 \\ 0 & -\frac{da}{dt} \end{pmatrix} b; \quad b(0) = Id.$$

Then one has

$$b(t) = \begin{pmatrix} e^{a(t)} & e^{a(t)} \int_0^t e^{-2a(s)} ds \\ 0 & e^{-a(t)} \end{pmatrix}$$

Now let $g = \begin{pmatrix} u & 0 \\ v & u^{-1} \end{pmatrix}$ and consider the Gauss decomposition of the matrix

$$bg = \begin{pmatrix} ue^{a(t)} + ve^{a(t)} \int_0^t e^{-2a(s)} ds & u^{-1} e^{a(t)} \int_0^t e^{-2a(s)} ds \\ ve^{-a(t)} & u^{-1} e^{-a(t)} \end{pmatrix}$$

One finds that

$$[bg]_- = \begin{pmatrix} 1 & 0 \\ \frac{ve^{-a(t)}}{ue^{a(t)} + ve^{a(t)} \int_0^t e^{-2a(s)} ds} & 1 \end{pmatrix}$$

and

$$[bg]_{0+} = \begin{pmatrix} ue^{a(t)} + ve^{a(t)} \int_0^t e^{-2a(s)} ds & u^{-1} e^{a(t)} \int_0^t e^{-2a(s)} ds \\ 0 & (ue^{a(t)} + ve^{a(t)} \int_0^t e^{-2a(s)} ds)^{-1} \end{pmatrix}$$

One can check the following proposition.

PROPOSITION 6.2. *The upper triangular matrix $[bg]_{0+}$ satisfies the differential equation*

$$\frac{d}{dt}[bg]_{0+} = \begin{pmatrix} \frac{d}{dt} T_{u,v} a(t) & 1 \\ 0 & -\frac{d}{dt} T_{u,v} a(t) \end{pmatrix} [bg]_{0+}$$

where $T_{u,v} a(t) = \log(E_{u,v} e^{a(t)})$.

This equation is of the same kind as the equation (6.5) satisfied by the original matrix b , but with a different initial point. The right action $E_{u,v}$ is thus obtained by taking the matrix solution to (6.5), multiplying it on the right by $g = \begin{pmatrix} u & 0 \\ v & u^{-1} \end{pmatrix}$ and looking at the diagonal part of the Gauss decomposition of the resulting matrix. Actually in this way the partial action $T_{u,v}$ extends to a partial action T_g of the whole group of invertible real 2×2 matrices. One starts from the path a , constructs the matrix b by the differential equation and then takes the 0-part in the Gauss decomposition of bg . This yields a path $T_g a$. The statement of the proposition above remains true for $[bg]_{0+}$. The importance of this statement is that one can iterate the procedure and see that $T_{g_1 g_2} = T_{g_2} \circ T_{g_1}$ when defined.

Consider now the element $s = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, then

$$T_s a(t) = a(t) + \log \left(\int_0^t e^{-2a(s)} ds \right).$$

This is the geometric lifting of the Pitman operator obtained in (6.3). In the next section we shall extend these considerations to groups of higher rank.

6.6. Paths in the Cartan algebra. We work now in the general framework of the beginning of section 6.3.

One has the usual decomposition $\mathfrak{g} = \mathfrak{n}_- + \mathfrak{a} + \mathfrak{n}_+$. Correspondingly there is a Gauss decomposition $g = [g]_- [g]_0 [g]_+$ with $[g]_- \in N_-$, $[g]_0 \in A$, $[g]_+ \in N$, defined on an open dense subset. We denote by $[g]_{0+} = [g]_0 [g]_+$ the $B = AN_+$ part of the decomposition.

The following is easy to check, and provides a useful characterization of the vector space generated by the e_i .

LEMMA 6.3. *Let $n \in \mathfrak{n}_+$, then one has $[h^{-1}nh]_+ = n$ for all $h \in N_-$ if and only if n belongs to the vector space generated by the e_i .*

Let a be a path in the Cartan algebra \mathfrak{a} and let b be a solution to the equation

$$\frac{d}{dt} b = \left(\frac{d}{dt} a + n \right) b$$

where $n \in \oplus_i \mathbb{C} e_i$.

PROPOSITION 6.4. *Let $g \in G$, and assume that bg has a Gauss decomposition, then the upper part $[bg]_{0+}$ in the Gauss decomposition of bg satisfies the equation*

$$(6.6) \quad \frac{d}{dt} [bg]_{0+} = \left(\frac{d}{dt} T_g a + n \right) [bg]_{0+}$$

where $T_g a(t)$ is a path in the Cartan algebra.

Proof.

Let us write the equation

$$\frac{d}{dt} ([bg]_- [bg]_{0+}) = \left(\frac{d}{dt} a + n \right) [bg]_- [bg]_{0+}$$

in the form

$$[bg]_-^{-1} \frac{d}{dt} [bg]_- = [bg]_-^{-1} \left(\frac{d}{dt} a + n \right) [bg]_- - \frac{d}{dt} [bg]_{0+} [bg]_{0+}^{-1}$$

Since the left hand side of this equation is lower triangular, the right hand side has zero upper triangular part therefore, by lemma 6.3

$$n = \left[[bg]^{-1} \left(\frac{d}{dt} a + n \right) [bg]_- \right]_+ = \left[\frac{d}{dt} [bg]_{0+} [bg]_{0+}^{-1} \right]_+$$

therefore there exists a path $T_g a$ such that equation (6.6) holds. \square

We now assume that

$$n = \sum_i n_i e_i$$

with all $n_i > 0$. When $g = \bar{s}_i$ is a fundamental reflection, one gets a geometric lifting of the Pitman operator

$$T_{s_i} a(t) = a(t) + \log \left(\int_0^t e^{-\alpha_i(a(s))} ds \right) \alpha_i^\vee$$

associated with the dual root system, i.e.

$$\lim_{\varepsilon \rightarrow 0} T_{s_i} \left(\frac{1}{\varepsilon} a \right) = \mathcal{P}_{\alpha_i^\vee} a.$$

Thanks to the above proposition, one can prove that these geometric liftings satisfy the braid relations, and T_w provides a geometric lifting of the Pitman operator \mathcal{P}_w for all $w \in W$.

Analogously the Littelmann raising and lowering operators also have geometric liftings.

6.7. Reduced double Bruhat cells. In this section we show how our considerations on Littelmann's path model allow us to make the connection with the work of Berenstein and Zelevinsky [3]. We consider a path a on the Cartan Lie algebra, and the solution b to $\frac{d}{dt} b = (\frac{d}{dt} a + n)b$. When we multiply b on the right by \bar{s}_{i_1} , and take its Gauss decomposition

$$[bs_{i_1}]_- [bs_{i_1}]_0 [bs_{i_1}]_+ = [b]_0 [b]_+ s_{i_1}$$

then

$$[b]_+ s_{i_1} [bs_{i_1}]_+^{-1} = [b]_0^{-1} [bs_{i_1}]_- [bs_{i_1}]_0 \in N s_{i_1} N \cap B_- L^{s_{i_1}, e}$$

and

$$[b]_+ s_{i_1} [bs_{i_1}]_+^{-1} = x_{-i_1}(r_1)$$

for some r_1 . In fact, using our formula for Littelmann operators,

$$r_1 = e^{\alpha_1(a(T))} \int_0^T e^{-\alpha_1(a(s))} ds.$$

Comparing with (3.3) we see that $r_1 e^{-\alpha_1(a(T))}$ gives a geometric lifting of the Kashiwara coordinate for the Littelmann module. We can continue the process starting from $[bs_{i_1}]_+$, to get

$$[bs_{i_1}]_+ s_{i_2} [bs_{i_1} s_{i_2}]_+^{-1} = x_{-i_2}(r_2)$$

(using the fact that $[g_1 g_2]_+ = [[g_1]_+ g_2]_+$ for $g_1, g_2 \in G$) obtaining successive decompositions

$$[b]_+ s_{i_1} \dots s_{i_k} [bs_{i_1} \dots s_{i_k}]_+^{-1} = x_{-i_1}(r_1) \dots x_{-i_k}(r_k)$$

This gives the Kashiwara coordinates of $[[b]_+ w_0]_- \in L^{w_0, e}$, which are thus seen to correspond to the coordinates on the cone by a geometric lifting.

7. Appendix

This appendix is devoted to the proof of theorem 2.6.

LEMMA 7.1. *If $B(\lambda), \lambda \in \bar{C}$, is a closed normal family of highest weight continuous crystals then for each $\lambda, \mu \in \bar{C}$ such that $\lambda \leq \mu$ there exists an injective map $\Psi_{\lambda, \mu} : B(\lambda) \rightarrow B(\mu)$ with the following properties*

- (i) $\Psi_{\lambda, \mu}(b_\lambda) = b_\mu$,
- (ii) $\Psi_{\lambda, \mu} e_\alpha^r(b) = e_\alpha^r \Psi_{\lambda, \mu}(b)$, for all $b \in B(\lambda), \alpha \in \Sigma, r \geq 0$,
- (iii) $\Psi_{\lambda, \mu} f_\alpha^r(b) = f_\alpha^r \Psi_{\lambda, \mu}(b)$ if $f_\alpha^r(b) \in B(\lambda)$.

PROOF. Let $\nu = \mu - \lambda$. First consider the map $\phi_{\lambda, \mu} : B(\lambda) \rightarrow B(\lambda) \otimes B(\nu)$ given by $\phi_{\lambda, \mu}(b) = b \otimes b_\nu$, when $b \in B(\lambda)$. Since b_ν is a highest weight $\varepsilon_\alpha(b_\nu) = 0$. By normality, for all $b \in B(\lambda), \varphi_\alpha(b) \geq 0$. Therefore $\sigma := \varphi_\alpha(b) - \varepsilon_\alpha(b_\nu) = \varphi_\alpha(b) \geq 0$. By definition, this implies that $\varepsilon_\alpha(b \otimes b_\nu) = \varepsilon_\alpha(b)$, $\varphi_\alpha(b \otimes b_\nu) = \varphi_\alpha(b)$, $wt(b \otimes b_\nu) = wt(b) + \nu$. Using (2.1) we see also that, for $r \geq 0$, $e_\alpha^r(b \otimes b_\nu) = e_\alpha^r b \otimes b_\nu$ and that, when $f_\alpha^r(b) \in B(\lambda)$, $r \leq \varphi_\alpha(b) = \sigma$ by normality, and therefore $f_\alpha^r(b \otimes b_\nu) = f_\alpha^r b \otimes b_\nu$. Since the family is closed there is an isomorphism $i_{\lambda, \mu} : \mathcal{F}(b_\lambda \otimes b_\nu) \rightarrow B(\mu)$. One has $i_{\lambda, \mu}(b_\lambda \otimes b_\nu) = b_\mu$. One can take $\Psi_{\lambda, \mu} = i_{\lambda, \mu} \circ \phi_{\lambda, \mu}$. \square

The family $\Psi_{\lambda, \mu}$ constructed above satisfies $\Psi_{\lambda, \lambda} = id$ and, when $\lambda \leq \mu \leq \nu$, $\Psi_{\mu, \nu} \circ \Psi_{\lambda, \mu} = \Psi_{\lambda, \nu}$, so that we can consider the direct limit $B(\infty)$ of the family $B(\lambda), \lambda \in \bar{C}$, with the injective maps $\Psi_{\lambda, \mu} : B(\lambda) \rightarrow B(\mu), \lambda \leq \mu$. Still following Joseph [19], we define a crystal structure on $B(\infty)$.

PROPOSITION 7.2. *The direct limit $B(\infty)$ is a highest weight upper normal continuous crystal with highest weight 0.*

PROOF. By definition, the direct limit $B(\infty)$ is the quotient set B / \sim where $B = \cup_{\lambda \in \bar{C}} B(\lambda)$ is the disjoint union of the $B(\lambda)$'s and where $b_1 \sim b_2$ for $b_1 \in B(\lambda), b_2 \in B(\mu)$, when there exists a $\nu \in \bar{C}$ such that $\nu \geq \lambda, \nu \geq \mu$ and $\Psi_{\lambda, \nu}(b_1) = \Psi_{\mu, \nu}(b_2)$. Let \bar{b} be the image in $B(\infty)$ of $b \in B$. If $b \in B(\lambda)$, then we define $wt(\bar{b}) = wt(b) - \lambda$, $\varepsilon_\alpha(\bar{b}) = \varepsilon_\alpha(b)$, $\varphi_\alpha(\bar{b}) = \varepsilon_\alpha(\bar{b}) + \alpha^\vee(wt(\bar{b}))$ and, when $r \geq 0$, $e_\alpha^r(\bar{b}) = \overline{e_\alpha^r(b)}$. These do not depend on λ , since if $\mu \geq \lambda$ and $b' = \Psi_{\lambda, \mu}(b)$, then one has $\bar{b}' = \bar{b}$ and $wt(b') = wt(b) + \mu - \lambda$. In order to define $f_\alpha^r(\bar{b})$ for $r \geq 0$, let us choose $\mu \geq \lambda$ large enough to ensure that

$$\varphi_\alpha(b') = \varepsilon_\alpha(b') + \alpha^\vee(wt(b)) + \alpha^\vee(\mu - \lambda) \geq r.$$

Then $f_\alpha^r b' \neq \mathbf{0}$ by normality and we define $f_\alpha^r \bar{b} = \overline{f_\alpha^r b'}$. Again this does not depend on μ . Using the lemma we check that this defines a crystal structure on $B(\infty)$. Each $\Psi_{\lambda, \mu}, \lambda \leq \mu$, commutes with the $e_\alpha^r, r \geq 0$. This implies that $B(\infty)$ is upper normal. Since each $B(\lambda)$ is a highest weight crystal, $B(\infty)$ has also this property. \square

We will denote b_∞ the unique element of $B(\infty)$ of weight 0. Note that $B(\infty)$ is not lower normal. For instance,

$$(7.1) \quad \varphi_\alpha(b_\infty) = 0, f(b_\infty) \neq \mathbf{0}, \text{ for all } f \in \mathcal{F}.$$

For $\lambda \in \bar{C}$ we define the crystal $S(\lambda)$ as the set with a unique element $\{s_\lambda\}$ and the maps $wt(s_\lambda) = \lambda, \varepsilon_\alpha(s_\lambda) = -\alpha^\vee(\lambda), \varphi_\alpha(s_\lambda) = 0$ and $e_\alpha^r(s_\lambda) = \mathbf{0}$ when $r \neq 0$.

LEMMA 7.3. *The map*

$$\Psi_\lambda : b \in B(\lambda) \mapsto \bar{b} \otimes s_\lambda \in B(\infty) \otimes S(\lambda)$$

is a crystal embedding.

PROOF. Let $b \in B(\lambda)$, then

$$wt(\Psi_\lambda(b)) = wt(\bar{b} \otimes s_\lambda) = wt(\bar{b}) + wt(s_\lambda) = wt(b) - \lambda + \lambda = wt(b).$$

Let $\sigma = \varphi_\alpha(\bar{b}) - \varepsilon_\alpha(s_\lambda)$. Then $\sigma = \varphi_\alpha(b)$ since $\varepsilon_\alpha(s_\lambda) = -\alpha^\vee(\lambda)$ and $\varphi_\alpha(\bar{b}) = \varphi_\alpha(b) - \alpha^\vee(\lambda)$. Thus $\sigma \geq 0$ by normality of $B(\lambda)$. By the definition of the tensor product, this implies that

$$\varepsilon_\alpha(\Psi_\lambda(b)) = \varepsilon_\alpha(\bar{b} \otimes s_\lambda) = \varepsilon_\alpha(\bar{b}) = \varepsilon_\alpha(b),$$

thus $\varphi_\alpha(\Psi_\lambda(b)) = \varphi_\alpha(b)$. Furthermore,

$$e_\alpha^r(\Psi_\lambda(b)) = e_\alpha^r(\bar{b} \otimes s_\lambda) = e_\alpha^{\max(r, -\sigma) - \sigma^-}(\bar{b}) \otimes e^{\min(r, -\sigma) + \sigma^-} s_\lambda.$$

When $r \geq -\sigma$, this is equal to $e_\alpha^r(b) \otimes s_\lambda = \Psi_\lambda(e_\alpha^r(b))$. If $r < -\sigma$ then $e_\alpha^r(\Psi_\lambda(b)) = e_\alpha^{-\sigma}(\bar{b}) \otimes e_\alpha^{r+\sigma}(s_\lambda) = \mathbf{0}$, since $e_\alpha^s(s_\lambda) = \mathbf{0}$ when $s \neq 0$, and on the other hand, $e_\alpha^r(b) = \mathbf{0}$ by normality. Thus $\Psi_\lambda(e_\alpha^r(b)) = \mathbf{0}$. \square

If $f = f_{\alpha_n}^{r_n} \cdots f_{\alpha_1}^{r_1} \in \mathcal{F}$, we say that $f' \in F$ is extracted from f if $f' = f_{\alpha_n}^{r'_n} \cdots f_{\alpha_1}^{r'_1}$ with $0 \leq r'_k \leq r_k, k = 1, \dots, n$. Recall the definition of $B_\alpha = \{b_\alpha(t), t \leq 0\}$ given in Example 2.2.

LEMMA 7.4. *Let $f \in \mathcal{F}$ and $\alpha \in \Sigma$, then there exists f' extracted from f and $t \geq 0$ such that*

$$f(b_\infty \otimes b_\alpha(0)) = f' b_\infty \otimes b_\alpha(-t).$$

Moreover if $\lambda \in \bar{C}$ is such that $\alpha^\vee(\lambda) = 0$ and $\beta^\vee(\lambda)$ large enough for all $\beta \in \Sigma - \{\alpha\}$, then for $\mu \in \bar{C}$, for the same $f' \in \mathcal{F}$ and $t \geq 0$,

$$f(b_\lambda \otimes b_\mu) = f' b_\lambda \otimes f_\alpha^t b_\mu.$$

PROOF. The first part follows easily from the definition of the tensor product. Let $\lambda \in \bar{C}$ such that $\alpha^\vee(\lambda) = 0$, $\mu \in \bar{C}, \beta \in \Sigma - \{\alpha\}$ and $r \geq 0$. If, for some $s > 0$, one has $e_\beta^s(f_\alpha^r b_\mu) \neq \mathbf{0}$ then $wt(e_\beta^s(f_\alpha^r b_\mu)) = \mu + s\beta - r\alpha$ is in $\mu - \bar{C}$ (since μ is a highest weight). This is not possible because $\beta^\vee(s\beta - r\alpha) \geq s\beta^\vee(\beta) > 0$. Therefore, by normality, $\varepsilon_\beta(f_\alpha^r b_\mu) = 0$. On the other hand, for all $f = f_{\alpha_n}^{r_n} \cdots f_{\alpha_1}^{r_1} \in \mathcal{F}$,

$$\varphi_\beta(fb_\lambda) = \beta^\vee(wt(fb_\lambda)) + \varepsilon_\beta(fb_\lambda) \geq \beta^\vee(wt(fb_\lambda)) = \beta^\vee(\lambda) - \sum_{k=1}^n r_k \beta^\vee(\alpha_k).$$

Let $\sigma = \varphi_\beta(fb_\lambda) - \varepsilon_\beta(f_\alpha^r b_\mu) = \varphi_\beta(fb_\lambda)$ and $s \geq 0$. Then

$$\sigma = \varphi_\beta(fb_\lambda) \geq \beta^\vee(\lambda) - \sum_{k=1}^n r_k \beta^\vee(\alpha_k).$$

If $\beta^\vee(\lambda)$ is large enough, then $\sigma \geq \max(s, 0)$ which implies, see (2.1), that

$$(7.2) \quad f_\beta^s(fb_\lambda \otimes f_\alpha^r b_\mu) = (f_\beta^s fb_\lambda) \otimes f_\alpha^r b_\mu.$$

On the other hand, $\varphi_\alpha(b_\lambda) = \alpha^\vee(\lambda) + \varepsilon_\alpha(b_\lambda) = 0$, since $\varepsilon_\alpha(b_\lambda) = 0$ by normality. We also know that $\varphi_\alpha(b_\infty) = 0$, see (7.1), hence

$$\varphi_\alpha(fb_\lambda) = \varphi_\alpha(b_\lambda) - \sum_{k=1}^n r_k \alpha^\vee(\alpha_k) = \varphi_\alpha(b_\infty) - \sum_{k=1}^n r_k \alpha^\vee(\alpha_k) = \varphi_\alpha(fb_\infty).$$

Thus $\sigma = \varphi_\alpha(fb_\infty)$ and does not depend on λ . It follows that the following decomposition is independent of λ :

$$(7.3) \quad f_\alpha^s(fb_\lambda \otimes f_\alpha^r b_\mu) = f_\alpha^{\sigma \wedge s} fb_\lambda \otimes f_\alpha^{r+s-\sigma \wedge s} b_\mu.$$

Using (7.2) and (7.3), it is now easy to prove the lemma by induction on n , proving first the second assertion. \square

PROPOSITION 7.5. *For each simple root α , there is a crystal embedding $\Gamma_\alpha : B(\infty) \rightarrow B(\infty) \otimes B_\alpha$ such that $\Gamma_\alpha(b_\infty) = b_\infty \otimes b_\alpha(0)$.*

PROOF. Let us show that the expression

$$(7.4) \quad \Gamma_\alpha(fb_\infty) = f(b_\infty \otimes b_\alpha(0)), \quad f \in \mathcal{F},$$

defines the morphism Γ_α . First we check that it is well defined. By definition, $fb_\infty = \overline{fb}_\nu$ for all $\nu \in \bar{C}$ such that $\overline{fb}_\nu \neq \mathbf{0}$.

Let us choose λ as in lemma 7.4. For $\mu \in \bar{C}$ large enough, $\overline{fb}_{\lambda+\mu} \neq \mathbf{0}$. Let us write

$$\overline{fb}_{\lambda+\mu} = f(\bar{b}_\lambda \otimes \bar{b}_\mu) = \overline{f'b}_\lambda \otimes \overline{f^t b}_\mu.$$

Then f' and t depend only on $fb_{\lambda+\mu}$, which by definition depends only on fb_∞ . By lemma 7.4,

$$f(b_\infty \otimes b_\alpha(0)) = f'b_\infty \otimes b_\alpha(-t)$$

which depends only on fb_∞ (and not on f itself), showing that Γ_α is well defined on $\mathcal{F}b_\infty$, and thus on $B(\infty)$, since $\mathcal{F}b_\infty = B(\infty)$. Notice that $f(b_\infty \otimes b_\alpha(0)) \neq \mathbf{0}$ since $f'b_\infty \neq \mathbf{0}$.

Let us prove that Γ_α is injective. Suppose that $f(b_\infty \otimes b_\alpha(0)) = \tilde{f}(b_\infty \otimes b_\alpha(0))$ for some $f, \tilde{f} \in \mathcal{F}$. Using lemma 7.4,

$$f(b_\infty \otimes b_\alpha(0)) = f'b_\infty \otimes b_\alpha(-t) \text{ and } \tilde{f}(b_\infty \otimes b_\alpha(0)) = \tilde{f}'b_\infty \otimes b_\alpha(-\tilde{t}).$$

If $\lambda \in \bar{C}$ is as in this lemma, then

$$f(b_\lambda \otimes b_\mu) = f'b_\lambda \otimes f_\alpha^t(b_\mu) = \tilde{f}'b_\lambda \otimes \tilde{f}_\alpha^t b_\mu = \tilde{f}(b_\lambda \otimes b_\mu),$$

therefore $fb_{\lambda+\mu} = \tilde{f}b_{\lambda+\mu}$, thus $fb_\infty = \tilde{f}b_\infty$. It is clear that Γ_α commutes with $f_\alpha^r, r \geq 0$. Since $\varepsilon_\alpha(b_\alpha(0)) = \varphi_\alpha(b_\infty) = 0$,

$$\varepsilon_\alpha(\Gamma_\alpha(b_\infty)) = \varepsilon_\alpha(b_\infty \otimes b_\alpha(0)) = \varepsilon_\alpha(b_\infty),$$

hence, if $f = f_{\alpha_n}^{r_n} \cdots f_{\alpha_1}^{r_1} \in \mathcal{F}$,

$$\varepsilon_\alpha(\Gamma_\alpha(fb_\infty)) = \varepsilon_\alpha(f\Gamma_\alpha(b_\infty)) = \varepsilon_\alpha(\Gamma_\alpha(b_\infty)) - \sum_{k=1}^n r_k \beta^\vee(\alpha_k) = \varepsilon_\alpha(fb_\infty).$$

Therefore Γ_α commutes with ε_α . It also commutes with wt since $wt(b_\infty) = 0$. Let us now consider $e_\alpha^r, r \geq 0$. Let $b \in B(\infty)$. If $e_\alpha^r(b) \neq \mathbf{0}$, then

$$\Gamma_\alpha(b) = \Gamma_\alpha(f_\alpha^r e_\alpha^r(b)) = f_\alpha^r(\Gamma_\alpha(e_\alpha^r(b))) \neq \mathbf{0}$$

hence $\Gamma_\alpha(e_\alpha^r(b)) = e_\alpha^r(\Gamma_\alpha(b))$. Suppose now that $e_\alpha^r(b) = \mathbf{0}$. Since $B(\infty)$ is upper normal, one has $\varepsilon_\alpha(b) = 0$, hence $\varepsilon_\alpha(\Gamma_\alpha(b)) = 0$. By the lemma, there is $f' \in \mathcal{F}$ and $t \geq 0$ such that $\Gamma_\alpha(b) = \Gamma_\alpha(b) = f'b_\infty \otimes b_\alpha(-t)$. Therefore

$$0 = \varepsilon_\alpha(\Gamma_\alpha(b)) \geq \varepsilon_\alpha(f'b_\infty) \geq 0.$$

By upper normality this implies that $e_\alpha^r(f'b_\infty) = \mathbf{0}$, hence

$$e_\alpha^r(\Gamma_\alpha(b)) = e_\alpha^r(f'b_\infty \otimes b_\alpha(-t)) = (e_\alpha^r f'b_\infty) \otimes b_\alpha(-t) = \mathbf{0}. \quad \square$$

The following lemma is clear.

LEMMA 7.6. *Let B_1, B_2 and C be three continuous crystals and $\psi : B_1 \rightarrow B_2$ be crystal embeddings. Then $\tilde{\psi} : B_1 \otimes C \rightarrow B_2 \otimes C$ defined by $\tilde{\psi}(b \otimes c) = \psi(b) \otimes c$ is a crystal embedding.*

7.1. Uniqueness. Proof of theorem 2.6. Recall that Σ is the set of simple roots. Fix a sequence $A = (\dots, \alpha_2, \alpha_1)$ of elements of Σ such that each simple root occurs infinitely many times and $\alpha_n \neq \alpha_{n+1}$ for all $n \geq 1$. Let $\hat{B}(A)$ be the subset of $\dots B_{\alpha_2} \otimes B_{\alpha_1}$ in which the k -th entry differs from $b_{\alpha_k}(0)$ for only finitely many k . One checks that the rules given for the multiple tensor give $\hat{B}(A)$ the structure of a continuous crystal (see, e.g., Kashiwara, [22], 7.2, Joseph [18], [19]). Let b_A be the element of $\hat{B}(A)$ with entries $b_{\alpha_n}(0)$ for all $n \geq 1$. We denote $B(A) = \mathcal{F}b_A$.

PROPOSITION 7.7. *There exists a crystal embedding Γ from $B(\infty)$ onto $B(A)$ such that $\Gamma(b_\infty) = b_A$.*

PROOF. Let $f \in \mathcal{F}$. We can write $f = f_{\alpha_k}^{r_k} \dots f_{\alpha_1}^{r_1}$ where $(\dots, \alpha_2, \alpha_1) = A$ and $r_n \geq 0$ for all $n \geq 1$. By lemma 7.4

$$\Gamma_{\alpha_1}(f_{\alpha_1}^{r_1}(b_\infty)) = f_{\alpha_1}^{r_1}(\Gamma_{\alpha_1}b_\infty) = f_{\alpha_1}^{r_1}(b_\infty \otimes b_{\alpha_1}(0)) = b_\infty \otimes b_{\alpha_1}(-r_1)$$

therefore

$$\Gamma_{\alpha_1}(f_{\alpha_k}^{r_k} \dots f_{\alpha_1}^{r_1}b_\infty) = (f_{\alpha_k}^{r'_k} \dots f_{\alpha_2}^{r'_2}b_\infty) \otimes b_{\alpha_1}(-r'_1)$$

for some $r'_1, \dots, r'_k \geq 0$. Similarly,

$$\Gamma_{\alpha_2}(f_{\alpha_k}^{r'_k} \dots f_{\alpha_2}^{r'_2}b_\infty) = (f_{\alpha_k}^{r''_k} \dots f_{\alpha_3}^{r''_3}b_\infty) \otimes b_{\alpha_2}(-r''_2)$$

for some $r''_2, r''_3, \dots, r''_k$. If we apply lemma 7.6 to $B_1 = B(\infty), B_2 = B(\infty) \otimes B_{\alpha_2}, \psi = \Gamma_{\alpha_2}, C = B_{\alpha_1}$, we obtain a crystal embedding

$$\tilde{\Gamma}_{\alpha_2} : B(\infty) \otimes B_{\alpha_1} \rightarrow B(\infty) \otimes B_{\alpha_2} \otimes B_{\alpha_1}$$

such that, for $b \in B(\infty), b_1 \in B_{\alpha_1}$

$$\tilde{\Gamma}_{\alpha_2}(b \otimes b_1) = \Gamma_{\alpha_2}b \otimes b_1.$$

Let $\Gamma_{\alpha_2, \alpha_1} = \tilde{\Gamma}_{\alpha_2} \circ \Gamma_{\alpha_1} : B(\infty) \rightarrow B(\infty) \otimes B_{\alpha_2} \otimes B_{\alpha_1}$, then

$$\begin{aligned} \Gamma_{\alpha_2, \alpha_1}(f_{\alpha_k}^{r_k} \dots f_{\alpha_1}^{r_1}b_\infty) &= \tilde{\Gamma}_{\alpha_2}(f_{\alpha_k}^{r'_k} \dots f_{\alpha_2}^{r'_2}b_\infty \otimes b_{\alpha_1}(-r'_1)) \\ &= \Gamma_{\alpha_2}(f_{\alpha_k}^{r'_k} \dots f_{\alpha_2}^{r'_2}b_\infty) \otimes b_{\alpha_1}(-r'_1) \\ &= (f_{\alpha_k}^{r''_k} \dots f_{\alpha_3}^{r''_3}b_\infty) \otimes b_{\alpha_2}(-r''_2) \otimes b_{\alpha_1}(-r'_1). \end{aligned}$$

Again, with Γ_{α_3} we build $\Gamma_{\alpha_3, \alpha_2, \alpha_1} = \tilde{\Gamma}_{\alpha_3} \circ \Gamma_{\alpha_2, \alpha_1}$. Inductively we obtain strict morphisms

$$\Gamma_{\alpha_k, \dots, \alpha_1} : B(\infty) \rightarrow B(\infty) \otimes B_{\alpha_k} \otimes \dots \otimes B_{\alpha_2} \otimes B_{\alpha_1}$$

such that for some s_k, \dots, s_1

$$\Gamma_{\alpha_k, \dots, \alpha_1}(f_{\alpha_k}^{r_k} \dots f_{\alpha_1}^{r_1}b_\infty) = b_\infty \otimes b_{\alpha_k}(-s_k) \otimes \dots \otimes b_{\alpha_1}(-s_1).$$

Now we can define $\Gamma : B(\infty) \rightarrow B(A)$ by the formula

$$\Gamma(f_{\alpha_k}^{r_k} \dots f_{\alpha_1}^{r_1}b_\infty) = \dots \otimes b_{\alpha_{k+n}}(0) \otimes \dots \otimes b_{\alpha_{k+1}}(0) \otimes b_{\alpha_k}(-s_k) \otimes \dots \otimes b_{\alpha_1}(-s_1).$$

One checks that this is a crystal embedding. \square

This shows that $B(\infty)$ is isomorphic to $B(A)$, which does not depend on the chosen closed family of crystals, and thus proves the uniqueness. It also shows that $B(A)$ does not depend on A , as soon as a closed family exists.

CONTENTS

1. Introduction	1
2. Continuous crystal	2
3. Pitman transforms and Littelmann path operators for Coxeter groups	5
4. Continuous Littelmann module	12
5. The Duistermaat-Heckman measure and Brownian motion	26
6. Littelmann modules and geometric lifting.	38
7. Appendix	45
References	49

References

- [1] V. Alexeev and M. Brion. Toric Degenerations of spherical varieties. *Selecta Math. (N.S.)*, 10, (2004) no.4, 453-478.
- [2] Ph. Biane, Ph. Bougerol and N. O’Connell. Littelmann paths and Brownian paths. *Duke Math. J.* 130 (2005), no. 1, 127-167.
- [3] A. Berenstein and A. Zelevinsky. Tensor product multiplicities, canonical bases and totally positive varieties. *Invent. Math.* 143 (2001), no. 1, 77-128.
- [4] N. Bourbaki. *Groupes et algèbres de Lie*, Ch. IV-VI: Hermann, Paris, 1975.
- [5] P. Caldero, Toric degenerations of Schubert varieties, *Transform. Groups* **7** (2002), no. 1, 51-60.
- [6] A. H. Dooley, J. Repka, N. J. Wildberger, Sums of adjoint orbits, *Linear and Multilinear Algebra*, 36 (1993), 79-101.
- [7] J. J. Duistermaat and G. J. Heckman. On the variation in the cohomology of the symplectic form of the reduced phase space. *Invent. Math.* 69 (1982), no. 2, 259-268.
- [8] W. Fulton. *Young Tableaux*. London Mathematical Society student texts: 35. Cambridge University Press, 1997.
- [9] S. Fomin and A. Zelevinsky. Double Bruhat cells and total positivity. *J. Amer. Math. Soc.* 12 (1999), no. 2, 335-380.
- [10] K. Gross, D. Richards, Total positivity, finite reflection groups, and a formula of Harish-Chandra. *J. Approx. Th.* 82 (1995) 60-87.
- [11] V. Guillemin and S. Sternberg. *Symplectic techniques in physics*. Second edition. Cambridge University Press, Cambridge, 1990.
- [12] G. J. Heckman, Projections of orbits and asymptotic behavior of multiplicities for compact connected Lie groups. *Invent. Math.* 67 (1982), no. 2, 333-356.
- [13] S. Helgason, *Geometric analysis on symmetric spaces*. Math surveys and monographs, 39, AMS, Providence, 1994.
- [14] A. Henriques and J. Kamnitzer, The octahedron recurrence and $\mathfrak{gl}(n)$ crystals. *Adv. in Math.* 206, no. 1 (2006), 211-249.
- [15] A. Henriques and J. Kamnitzer, Crystals and coboundary categories. *Duke Math. J.*, 132 (2006) no. 2, 191-216.
- [16] J. Hong, S-J. Kang, *Introduction to Quantum Groups and Crystal Bases* Graduate studies in Mathematics, Vol. 42, AMS, Providence, 2002.
- [17] J. E. Humphreys. *Reflection Groups and Coxeter Groups*. Cambridge Studies in Advanced Mathematics 29. Cambridge University Press, 1990.
- [18] A. Joseph, *Quantum groups and their primitive ideals*, Springer, Berlin 1995
- [19] A. Joseph, Lie algebras, their representations and crystals,
<http://www.wisdom.weizmann.ac.il/~gorelik/agrt.htm>

- [20] M. Kashiwara. The crystal base and Littelmann's refined Demazure character formula. *Duke Math. J.* 71, no. 3 (1993), 839-858
- [21] M. Kashiwara. Crystal bases of modified quantized enveloping algebra. *Duke Math. J.* 73 (1994), no. 2, 383-413.
- [22] M. Kashiwara, *Bases cristallines des groupes quantiques*, rédigé par Charles Cochet, Cours Spécialisés SMF, 9 (2002), 115 pages
- [23] A. Lascoux, B. Leclerc, J.-Y. Thibon, Crystal graphs and q -analogues of weight multiplicities for the root system A_n . *Lett. Math. Phys.* 35 (1995) no. 4 395-374.
- [24] C. Lenart, On the combinatorics of crystal Graphs II. The Crystal Commutor, C. Lenart, Preprint, 2006, math.RT/0611444.
- [25] P. Littelmann. Paths and root operators in representation theory, *Annals of Mathematics* 142, 1995, pp. 499-525.
- [26] P. Littelmann. Cones, crystals, and patterns. *Transform. Groups* 3 (1998), no. 2, 145-179.
- [27] G. Lusztig, *Introduction to quantum groups*, Birkhauser, Boston-Basel-Berlin, 1994
- [28] S. Morier-Genoud, Geometric lifting of the canonical basis and semitoric degenerations of the Richardson varieties, to appear in *Trans. Amer. Math. Soc.*, math.RT/0504538
- [29] R.V. Moody, J. Patera, Quasicrystals and icosians, *J. Phys. A: Math. Gen.* 26, (1993), 2829-2853.
- [30] N. O'Connell, M. Yor, Brownian analogues of Burke's theorem, *Stochastic Process. Appl.* 96 (2001), 285-304.
- [31] E.M. Opdam, Dunkl operators, Bessel functions and the discriminant of a finite Coxeter group, *Compositio Math.* 85 (1993), 333-373.
- [32] J.W. Pitman. One-dimensional Brownian motion and the three-dimensional Bessel process, *Adv. Appl. Probab.* 7 (1975) 511-526.
- [33] M. Rösler, Positivity of Dunkl intertwining operator. *Duke Math. J.* 98 (1999), 445- 463.
- [34] M. Rösler, A positive radial product formula for the Dunkl kernel. *Trans. Amer. Math. Soc.* 355 (2003), 2413-2438.

CNRS, IGM, UNIVERSITÉ PARIS-EST, 77454 MARNE-LA-VALLÉE CEDEX 2
E-mail address: `Philippe.Biane@univ-mlv.fr`

LABORATOIRE DE PROBABILITÉS ET MODÈLES ALÉATOIRES, UNIVERSITÉ PIERRE ET MARIE CURIE, 4, PLACE JUSSIEU, 75005 PARIS, FRANCE
E-mail address: `philippe.bougerol@upmc.fr`

MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY CV4 7AL, UK
E-mail address: `n.m.o-connell@warwick.ac.uk`