



HAL
open science

Spectral extrema and Lifshitz tails for non monotonous alloy type models

Frédéric Klopp, Shu Nakamura

► **To cite this version:**

Frédéric Klopp, Shu Nakamura. Spectral extrema and Lifshitz tails for non monotonous alloy type models. 2008. hal-00275774v1

HAL Id: hal-00275774

<https://hal.science/hal-00275774v1>

Preprint submitted on 25 Apr 2008 (v1), last revised 12 Aug 2008 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Spectral extrema and Lifshitz tails for non monotonous alloy type models

Frédéric Klopp* and Shu Nakamura†

April 25, 2008

Abstract

In the present note, we determine the ground state energy and study the existence of Lifshitz tails near this energy for some non monotonous alloy type models. Here, non monotonous means that the single site potential coming into the alloy random potential changes sign. In particular, the random operator is not a monotonous function of the random variables.

RÉSUMÉ. Cet article est consacré à la détermination de l'énergie de l'état fondamental et à l'étude de possibles asymptotiques de Lifshitz au voisinage de cette énergie pour certains modèles d'Anderson continus non monotones. Ici, non monotone signifie que le potentiel de simple site entrant dans la composition du potentiel aléatoire change de signe. En particulier, l'opérateur aléatoire n'est pas une fonction monotone des variables aléatoires.

0 Introduction and results

In this paper, we consider the continuous alloy type (or Anderson) random Schrödinger operator:

$$H_\omega = -\Delta + V_\omega \text{ where } V_\omega(x) = \sum_{\gamma \in \mathbb{Z}^d} \omega_\gamma V(x - \gamma) \quad (0.1)$$

on \mathbb{R}^d , $d \geq 1$, where V is the site potential, and $(\omega_\gamma)_{\gamma \in \mathbb{Z}^d}$ are the random coupling constants. Throughout this paper, we assume

(H1) (1) $V : \mathbb{R}^d \rightarrow \mathbb{R}$ is L^p (where $p = 2$ if $d \leq 3$ and $p > d/2$ if $d > 3$), non identically vanishing and supported in $(-1/2, 1/2)^d$;

*LAGA, U.M.R. 7539 C.N.R.S, Institut Galilée, Université de Paris-Nord, 99 Avenue J.-B. Clément, F-93430 Villetaneuse, France et Institut Universitaire de France. Email: klopp@math.univ-paris13.fr

†Graduate School of Mathematical Sciences, University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, Japan 153-8914. Email: shu@ms.u-tokyo.ac.jp

- (2) $(\omega_\gamma)_\gamma$ are independent identically distributed (i.i.d.) random variables distributed in $[a, b]$ ($a < b$) with essential infimum a and essential supremum b .

Let Σ be the almost sure spectrum of H_ω and $E_- = \inf \Sigma$. When V has a fixed sign, it is well known that the $E_- = \inf(\sigma(-\Delta + V_{\bar{b}}))$ if $V \leq 0$ and $E_- = \inf(\sigma(-\Delta + V_{\bar{a}}))$ if $V \geq 0$. Here, \bar{x} is the constant vector $\bar{x} = (x)_{\gamma \in \mathbb{Z}^d}$. Moreover, in this case, it is well known that the integrated density of states of the Hamiltonian (see e.g. (0.2)) admit a Lifshitz tail near E_- , i.e., that the integrated density of states at energy E decays exponentially fast as E goes to E_- from above. We refer to [9, 7, 22, 20, 6, 5, 11] for precise statements.

In the present paper, we address the case when V changes sign, i.e., there may exist $x_+ \neq x_-$ such that

$$V(x_-) \cdot V(x_+) < 0.$$

The basic difficulty this property introduces is that the variations of the potential V_ω as a function of ω is not monotonous. In the monotonous case, to get the minimum, one can simply minimize with respect to each of the random variables individually. In the non monotonous case, this uncoupling between the different random variables may fail. Our results concern reflection symmetric potential since, as we will see, for these potentials we also have analogous decoupling between the different random variables. Thus we make the following symmetry assumption on V :

(H2) V is reflection symmetric i.e. for any $\sigma = (\sigma_1, \dots, \sigma_d) \in \{0, 1\}^d$ and any $x = (x_1, \dots, x_d) \in \mathbb{R}^d$,

$$V(x_1, \dots, x_d) = V((-1)^{\sigma_1} x_1, \dots, (-1)^{\sigma_d} x_d).$$

We now consider the operator $H_\lambda^N = -\Delta + \lambda V$ with Neumann boundary conditions on the cube $[-1/2, 1/2]^d$. Its spectrum is discrete, and we let $E_-(\lambda)$ be its ground state energy. It is a simple eigenvalue and $\lambda \mapsto E_-(\lambda)$ is a real analytic concave function defined on \mathbb{R} . We first observe:

Proposition 0.1. *Under the above assumptions (H1) and (H2),*

$$E_- = \inf(E_-(a), E_-(b)).$$

For a and b sufficiently small, this result was proven in [17] without the assumption (H2) but with an additional assumption on the sign of $\int_{\mathbb{R}^d} V(x) dx$. The method used by Najar relies on a small coupling constant expansion for the infimum of Σ . These ideas were first used in [3] to treat other non

monotonous perturbations, in this case magnetic ones, of the Laplace operator. In [1], the authors study the minimum of the almost sure spectrum for a random displacement model i.e. the random potential is defined as $V_\omega(x) = \sum_{\gamma \in \mathbb{Z}^d} V(x - \gamma - \xi_\gamma)$ where $(\xi_\gamma)_\gamma$ are i.i.d. random variables supported in a sufficiently small compact.

We now turn to the results on Lifshitz tails. We denote by $N(E)$ the integrated density of states of H_ω , i.e., it is defined by the limit

$$N(E) = \lim_{L \rightarrow +\infty} \frac{\#\{\text{eigenvalues of } H_{\omega,L}^N \leq E\}}{(2L+1)^d} \quad (0.2)$$

where $H_{\omega,L}^N$ is the operator H_ω restricted to the cube $[-L - 1/2, L + 1/2]^d$ with Neumann boundary conditions. This quantity has been the source of a lot of studies and we refer to [20, 23] for extensive review.

We first give an upper bound on the integrate density of states. In the applications of the Lifshitz tails asymptotics, in particular, to localization, this side of the bound is the most important and also the difficult one to obtain.

Theorem 0.1. *Suppose assumptions (H1) and (H2) are satisfied. Assume moreover that*

$$E_-(a) \neq E_-(b). \quad (0.3)$$

then

$$\limsup_{E \rightarrow E_-^+} \frac{\log |\log N(E)|}{\log(E - E_-)} \leq -\frac{d}{2} - \alpha_+ \quad (0.4)$$

where we have set $c = a$ if $E_-(a) < E_-(b)$ and $c = b$ if $E_-(a) > E_-(b)$, and

$$\alpha_+ = -\limsup_{\varepsilon \rightarrow 0} \frac{\log |\log \mathbb{P}(\{|c - \omega_0| \leq \varepsilon\})|}{\log \varepsilon} \geq 0.$$

As will be clear from the proofs, we could also consider the model $H_\omega = H_0 + V_\omega$ where V_ω is above and $H_0 = -\Delta + W$ where W is a \mathbb{Z}^d -periodic potential that satisfies the symmetry assumption (H2).

We now study a lower bound for the integrated density of states that we will prove in a more general case than the upper bound, i.e., we don't need to assume (H2). The assumption is as follows:

(HP) there exists $\omega^P \in [a, b]^{\mathbb{Z}^d}$ that is periodic (, i.e., for some $L_0 \in \mathbb{N}$, for all $\gamma \in \mathbb{Z}^d$ and $\beta \in \mathbb{Z}^d$, $\omega_{\gamma+L_0\beta}^P = \omega_\gamma^P$) such that $\inf \Sigma = \inf \sigma(H_{\omega^P})$.

Under this assumption, we have

Theorem 0.2. *Let H_ω be defined as above, and assume (H1) and (HP) hold. Then*

$$-\frac{d}{2} - \alpha_- \leq \liminf_{E \rightarrow E_-^+} \frac{\log |\log N(E)|}{\log(E - E_-)}, \quad (0.5)$$

where

$$\alpha_- = -\liminf_{\varepsilon \rightarrow 0} \frac{\log |\log \mathbb{P}(\{\forall \gamma \in \mathbb{Z}^d / L_0 \mathbb{Z}^d; |\omega_\gamma^P - \omega_\gamma| \leq \varepsilon\})|}{\log \varepsilon}.$$

Assume now that (H1) and (H2) hold, hence, (HP) holds with $\omega^P = \bar{a}$ or $\omega^P = \bar{b}$. Indeed it is not difficult to see that, under assumption (H2), $E_-(a)$ is also the bottom of the spectrum of the periodic operator $H_{\bar{a}}$ (see section 1). If we assume that $\alpha_+ = \alpha_- = 0$ and $E_-(a) \neq E_-(b)$, we obtain the following corollary

Theorem 0.3. *Assume that (H1) and (H2) and (0.3) hold and that $\alpha_- = \alpha_+ = 0$ then*

$$\lim_{E \rightarrow E_-^+} \frac{\log |\log N(E)|}{\log(E - E_-)} = -\frac{d}{2}.$$

Combining Theorem 0.1 with the Wegner estimates obtained in [10, 4] and the multiscale analysis as developed in [2], we learn

Theorem 0.4. *Assume (H1), (H2) and (0.3) hold. Assume, moreover, that the common distribution of the random variables admits an absolutely continuous density. Then, the bottom edge of the spectrum of H_ω exhibits complete localization in the sense of [2].*

Lifshitz tail have already been proved for various non monotonous random models, mainly models with a random magnetic fields (see e.g. [3, 15, 18, 19]). In the models we consider, we will now see that Lifshitz tail do not always happen.

In a companion paper (see [14]), we study the case when $E_-(a) = E_-(b)$. This requires techniques different from the ones used in the present paper and gets particularly interesting when the random variables are Bernoulli distributed. However, it is quite easy to see that, when $E_-(a) = E_-(b)$, Lifshitz tails may fail; the density of states can even exhibit a van Hove singularity.

Theorem 0.5. *There exists potentials V and random variables $(\omega_\gamma)_\gamma$ satisfying (H1) and (H2) such that*

- $E_-(a) = E_-(b) = 0$
- there exists $C > 0$ such that, for $E \geq 0$, one has

$$\frac{1}{C} E^{d/2} \leq N(E) \leq C E^{d/2}.$$

This paper is constructed as follows. In Section 1, we determine the bottom of the almost sure spectrum, and prove Proposition 0.1. In Section 2,

we prove our main theorems, Theorem 0.2 and Theorem 0.4. Finally, in section 3, we prove Theorem 0.5.

Acknowledgment. FK would like to thank the University of Tokyo where part of this work was done. A part of this research was done when SN was invited to Univ. Paris 13 in 2007, and he would like to thank it. SN is partially supported by JSPS Research Grant, Kiban (B) 17340033.

1 Determining the bottom of the spectrum

We denote by $E_-(t)$ be the ground state energy of the operator $H_{t,0}^N$, i.e., $-\Delta + tV$ on $[-1/2, 1/2]^d$ with Neumann boundary conditions. This function is concave and, hence, for $t \in [a, b]$, we have $E_-(t) \geq \min(E_-(a), E_-(b))$. Then, partitioning \mathbb{R}^d into the cubes $\gamma + [-1/2, 1/2]^d$ for $\gamma \in \mathbb{Z}^d$, and, restricting H_ω to each of these cubes with Neumann boundary conditions, we obtain

$$H_\omega \geq \bigoplus_{\gamma \in \mathbb{Z}^d} H_{\omega_\gamma, 0}^N$$

Hence we learn

$$H_\omega \geq \min(E_-(a), E_-(b)).$$

We let $L \geq 1$, and consider $H_{\omega, L}^P$, the operator H_ω restricted to the cube $[-L - 1/2, L + 1/2]^d$ with periodic boundary conditions. Clearly, this operator depends only on finitely many random variables. We prove

Lemma 1.1.

$$\Sigma = \overline{\bigcup_{L \geq 1} \bigcup_{\omega \text{ admissible}} \sigma(H_{\omega, L}^P)},$$

where ω is called admissible if all the components of ω are in the support of the distribution of the random variables defining the alloy type random operator.

This lemma is a variant of a standard characterization of the almost sure spectrum of an alloy type model. To prove Proposition 0.1, i.e., that $E_- = \min(E_-(a), E_-(b))$, it is hence enough to prove that, for any L sufficiently large,

$$\inf_{\omega \in [a, b]^{C_L^d}} \inf \sigma(H_{\omega, L}^P) \leq \min(E_-(a), E_-(b)) \quad (1.1)$$

where $C_L^d = \mathbb{Z}^d \cap [-L - 1/2, L + 1/2]^d$. To prove (1.1), we will use the assumption (H.2). For the sake of definiteness, let us assume $E_-(a) \leq E_-(b)$.

The ground state of $H_{a, 0}^N$, say ψ , is simple and can be chosen uniquely as a normalized positive function. The reflection symmetry of the potential V guarantees that ψ is reflection symmetric. For $\gamma \in \mathbb{Z}^d$ such that

$|\gamma| = |\gamma_1| + \dots + |\gamma_d| = 1$, we can continue ψ to the $\gamma + [-1/2, 1/2]^d$ by reflection symmetry with respect to the common boundary of $[-1/2, 1/2]^d$ and $\gamma + [-1/2, 1/2]^d$. As ψ is reflection symmetric, we continue this process of reflection with respect to the boundaries of the new cubes to obtain a continuation of ψ that is \mathbb{Z}^d -periodic, positive and reflection symmetric with respect to any plane that is common boundary to two cubes of the form $\gamma + [-1/2, 1/2]^d$. Moreover ψ satisfies, for any $L \geq 0$, $H_{a,L}^P \psi = H_{a,0}^P \psi = H_{a,0}^N \psi = E_-(a)\psi$. This proves that $E_-(a) \geq \inf \sigma(H_{a,L}^P)$. Hence, (1.1) holds. This completes the proof of Proposition 0.1. \square

Proof of Lemma 1.1. Recall the well known characterization of the almost sure spectrum of an alloy type model in terms of periodic approximations (see e.g. [20, 5]). Therefore, for $L \geq 1$, define the $L\mathbb{Z}^d$ -periodic operator

$$H_{\omega,L} = -\Delta + V_{\omega,L}, \quad V_{\omega,L}(\cdot) = \sum_{\beta \in L\mathbb{Z}^d} \sum_{\gamma \in \mathbb{Z}^d / (L\mathbb{Z}^d)} \omega_\gamma V(\cdot - \beta - \gamma). \quad (1.2)$$

Then, it is well known (see e.g. [20]) that

$$\Sigma = \overline{\bigcup_{L \geq 1} \bigcup_{(\omega_\gamma)_{\gamma \in \mathbb{Z}^d / (L\mathbb{Z}^d)} \text{ admissible}} \sigma(H_{\omega,L})}$$

where $(\omega_\gamma)_{\gamma \in \mathbb{Z}^d / (L\mathbb{Z}^d)}$ is admissible if all its components belong to the support of the random variables defining the alloy type model.

Floquet theory (see e.g. [21]) guarantees that $\sigma(H_{\omega,L}^P) \subset \sigma(H_{\omega,L})$. So, in order to prove Lemma 1.1, it is sufficient to prove that

$$\sigma(H_{\omega,L}) \subset \overline{\bigcup_{n \geq 1} \sigma(H_{\omega^L, nL}^P)} \quad (1.3)$$

for some well chosen admissible ω^L .

Consider ω^L defined by $\omega_{\gamma+L\beta}^L = \omega_\gamma$ for $\gamma \in \mathbb{Z}^d / (L\mathbb{Z}^d)$ and $\beta \in \mathbb{Z}^d$. Clearly, $V_{\omega^L} = V_{\omega,L}$; hence, if $E_n(\theta)$ are the Floquet eigenvalues of $H_{\omega,L}$, the spectrum of the operator $H_{\omega^L, nL}^P$ is given by the values $E_n(2\pi\gamma/n)$ for $\gamma \in \mathbb{Z}^d / (nL\mathbb{Z}^d)$ (see e.g.). (1.3) then follows from the continuity of the Floquet eigenvalues as function of the Floquet parameter (see e.g Lemma 7.1 in [16]). \square

2 Lifshitz tails

To fix ideas let us assume $E_-(a) < E_-(b)$. The two bounds in 0.1 will be proved separately.

2.1 The upper bound

The upper bound on the integrated density of states, Theorem 0.1, will be an immediate consequence of the following result.

Theorem 2.1. *Suppose assumptions (H1) and (H2) are satisfied, and, that $E_-(a) < E_-(b)$. Then, there exists $C > 0$ such that, for E sufficiently close to $E_-(a)$, one has*

$$N(E) \leq N_m(C(E - E_-(a))) \quad (2.1)$$

where N_m is the integrated density of states of the random operator

$$H_\omega^m = H_{\bar{a}} - E_-(a) + \sum_{\gamma \in \mathbb{Z}^d} (\omega_\gamma - a) \mathbf{1}_{[-1/2, 1/2]^d}(x - \gamma) \quad (2.2)$$

and $H_{\bar{a}}$ is defined above.

The upper bound is then deduced from the same bound for the integrated density of states of H_ω^m which is standard (see e.g. [13, 12, 11]).

Proof. We first note it is well known that, at E , a continuity point of $N(E)$, the sequence

$$N_L^N(E) = \mathbb{E} \left(\frac{\#\{\text{eigenvalues of } H_{\omega, L}^N \leq E\}}{(2L)^d} \right) \quad (2.3)$$

is decreasing and converges to $N(E)$ (see e.g. [20, 5]). So to prove Theorem 2.1, it suffices to prove that, there exists $c > 0$ such that, for E real and L large

$$N_L^N(E) \leq N_L^m(C(E - E_-(a)))$$

where $N_L^m(E)$ is defined by (2.3) where $H_{\omega, L}^N$ is replaced by $H_{\omega, L}^m$, i.e., the restriction of H_ω^m to $[-L-1/2, L+1/2]^d$ with Neumann boundary conditions. By the Rayleigh-Ritz principle ([21]), this follows from the quadratic form inequality

$$H_{\omega, L}^m \leq C(H_{\omega, L}^N - E_-(a)).$$

Under our assumptions on V , the form domain of both of these operators are $H^1([-L-1/2, L+1/2]^d)$. Now, as for $\psi \in H^1([-L-1/2, L+1/2]^d)$ and $\gamma \in \mathbb{Z}^d \cap +[-L-1/2, L+1/2]^d$, $\psi \mathbf{1}_{\gamma+[-1/2, 1/2]^d} \in H^1(\gamma+[-1/2, 1/2]^d)$, this inequality in turns follows from the inequalities

$$\begin{aligned} \forall \gamma \in \mathbb{Z}^d \cap +[-L-1/2, L+1/2]^d, \forall \psi \in H^1(\gamma+[-1/2, 1/2]^d), \\ \langle H_{\omega, L}^m \psi, \psi \rangle_{\gamma+[-1/2, 1/2]^d} \leq C \langle (H_{\omega, L}^N - E_-(a)) \psi, \psi \rangle_{\gamma+[-1/2, 1/2]^d}. \end{aligned}$$

Taking into account the structure of our random potentials, we see that this will follow from the operator inequality

$$(H_{a, 0}^N - E_-(a)) + (t - a) \leq C(H_{t, 0}^N - E_-(a)). \quad (2.4)$$

where $H_{t,0}^N = -\Delta + tV$ on $[-1/2, 1/2]^d$ with Neumann boundary conditions as before.

Lemma 2.1. *Let H_0 be self-adjoint on \mathcal{H} a separable Hilbert space such that $0 = \inf \sigma(H_0)$. Let V_1 be a non trivial closed symmetric operator relatively bounded with respect to H_0 with bound 0. Set $H_1 = H_0 + V_1$ and $E_1 = \inf \sigma(H_1)$. Assume $E_1 > 0$. Then, there exists $C > 0$ such that, for $t \in [0, 1]$, one has*

$$C(H_0 + tV_1) \geq H_0 + t$$

Lemma 2.1 applies to our case and, as a result, we obtain (2.4). This completes the proof of Theorem 2.1. \square

Proof of Lemma 2.1. Regular perturbation theory ensures that $\inf \sigma(H_0 + \beta V_1) = \delta > 0$ for some $\beta > 1$. Then, for $t \in [0, 1]$, we have

$$\begin{aligned} H_0 + tV_1 &= (1 - t/\beta)H_0 + (t/\beta)(H_0 + \beta V_1) \\ &\geq (1 - 1/\beta)H_0 + (t/\beta)\delta \geq \frac{1}{C}(H_0 + t) \end{aligned}$$

where $C^{-1} = (1 - 1/\beta) \cdot \inf(1, \delta/\beta)$. \square

2.2 The lower bound

We will use the techniques set up in [12, 13, 16]. We first recall two results from these papers which are valid in the generality of the present work. Consider a random Schrödinger operator of the form (0.1) where

(HL1) V is a not identically vanishing real valued, compactly supported function that is in L^p (where $p = 2$ if $d \leq 3$ and $p > d/2$ if $d > 3$);

(HL2) the random variables $(\omega_\gamma)_{\gamma \in \Gamma}$ are independent, identically distributed, non trivial and bounded.

Clearly these two assumptions are consequences of assumption (H1).

The existence of $N(E)$, the integrated density of states defined by (0.2) is known ([20, 23]).

Theorem 2.2 ([12]). *Assume (HL1) and (HL2) hold. Pick $\eta_0 > 0$ and $I \subset \mathbb{R}$, a compact interval. Then, there exists $\nu_0 > 0$ and $\varepsilon_0 > 0$ such that, for $0 < \varepsilon < \varepsilon_0$, $E_0 \in I$ and $L \geq \varepsilon^{-\nu_0}$, we have*

$$\begin{aligned} &\mathbb{E}(N_{\omega,L}(E_0 + \varepsilon/2) - N_{\omega,L}(E_0 - \varepsilon/2)) - e^{-(L\varepsilon^{\nu_0})^{-\eta_0}} \\ &\leq N(E_0 + \varepsilon) - N(E_0) \\ &\leq \mathbb{E}(N_{\omega,L}(E_0 + 2\varepsilon) - N_{\omega,L}(E_0 - 2\varepsilon)) + e^{-(L\varepsilon^{\nu_0})^{-\eta_0}}. \end{aligned} \tag{2.5}$$

where $N_{\omega,L}$ is the integrated density of states of the periodic operator $H_{\omega,L}$ defined in (1.2).

In [12], Theorem 2.2 is not stated in exactly the same form and under slightly stronger but unnecessary assumptions. The modifications necessary to obtain the form given here are simple and left to the reader.

The second result we use is the following

Lemma 2.2 ([12]). *Consider a periodic Schrödinger operator of the form H_{ω^P} where ω^P satisfies (HP). Let $E_0 = \inf \sigma(H_{\omega^P})$. Then, for any $\alpha \in (0, 1)$ and any ε sufficiently small, there exists a function w_ε with the following properties*

- (1) w_ε is supported in a ball of center 0 and radius $2\varepsilon^{-(1+2\alpha)/2}$;
- (2) $1 \leq \|w_\varepsilon\|_{L^2}^2 \leq 2$;
- (3) $\|(H_{\omega^P} - E_0)w_\varepsilon\|_{L^2}^2 \leq C\varepsilon^{2(1+\alpha)}$ for some $C > 0$ (independent of ε).

Though not formulated as a lemma, this result is proved in section 2 of [12].

We now prove the lower bound. Pick $\alpha \in (0, 1)$ arbitrary. Let Γ' be the lattice with respect to which ω^P is periodic. Pick $n \in \mathbb{N}^*$ such that $n \sim \varepsilon^{-\nu}$ for $\nu > \nu_0$ (ν_0 given by Theorem 2.2), ν to be chosen sufficiently large. As w_ε constructed in Lemma 2.2 has compact support in the interior of $C(0, n)$, it can be “periodized” to satisfy quasi-periodic boundary conditions; for $\theta \in \mathbb{R}^d$, we set

$$w_{\varepsilon, \theta}(\cdot) = \sum_{\beta \in (2n+1)\Gamma'} e^{-i\beta\theta} w_\varepsilon(\cdot + \beta).$$

Then, $w_{\varepsilon, \theta}$ satisfies $w_{\varepsilon, \theta}(x + \beta) = e^{i\beta\theta} w_{\varepsilon, \theta}(x)$ for $x \in \mathbb{R}^d$ and $\beta \in (2n+1)\Gamma'$, and we have

$$\|w_{\varepsilon, \theta}\|_{L^2(C(0, n))}^2 \geq 1. \quad (2.6)$$

We define

$$\Lambda_\alpha(\varepsilon) = \left\{ \gamma \in \Gamma'; \text{ for } 1 \leq j \leq d, |\gamma_j| \leq \varepsilon^{-(1+3\alpha)/2} \right\}.$$

Since V satisfies (HL1), if we assume $\omega_\gamma \leq \varepsilon^{1+\alpha}$ for $\gamma \in \Lambda_\alpha(\varepsilon)$, then

$$\|V_{\omega, n}(H_{\omega^P} - E_0 - 1)^{-1}\| \leq C\varepsilon^{1+\alpha}.$$

This, (2.6) and point (3) of Lemma 2.2 imply that, for some $C > 0$ and ε sufficiently small, we have

$$\|(H_{\omega, n, \theta} - E_0)w_{\varepsilon, \theta}\| \leq C\varepsilon^{1+\alpha}\|w_{\varepsilon, \theta}\|.$$

This proves that, for ε sufficiently small, if $\omega_\gamma \leq \varepsilon^{1+\alpha}$ for $\gamma \in \Lambda_\alpha(\varepsilon)$, then for all $\theta \in \mathbb{T}_n^*$, $H_{\omega, n, \theta}$ has an eigenvalue in $[E_0 - \varepsilon/2, E_0\varepsilon/2]$. Hence, we learn that

$$\mathbb{P}(N_{\omega, n}(E_0 + \varepsilon/2) - N_{\omega, n}(E_0 - \varepsilon/2)) \geq P(\varepsilon, \alpha)$$

where $P(\varepsilon, \alpha)$ is the probability of the event $\{\omega; \forall \gamma \in \Lambda_\alpha(\varepsilon), \omega_\gamma \leq \varepsilon^{1+\alpha}\}$. By assumption (HL2), we have

$$\liminf_{\substack{\varepsilon \rightarrow 0 \\ \varepsilon > 0}} \frac{\log |\log(P(\varepsilon, \alpha))|}{\log \varepsilon} \geq -\frac{d}{2}(1 + 3\alpha) - \alpha_-$$

where α_- is defined in Theorem 0.2. Since $\alpha > 0$ is arbitrary, this completes the proof of Theorem 0.2. \square

3 An example where Lifshitz tails fail to exist

We now will construct an example of the type described in Theorem 0.5.

Let $\varphi \in \mathcal{C}^\infty((-1/2, 1/2)^d)$ be positive, reflection symmetric, constant near the boundary of $[-1/2, 1/2]^d$ and normalized in the cube. Let $V = \Delta\varphi/\varphi$. Then, φ is the positive normalized ground state of $-\Delta + V$ on $[-1/2, 1/2]^d$ with Neumann boundary conditions. Let $(\omega_\gamma)_{\gamma \in \mathbb{Z}^d}$ be Bernoulli r.v. with support $\{0, 1\}$. Pick $L \geq 1$ and let φ_L be ground state of the operator $H_{\omega, L}^N$ acting on $[-1/2, L - 1/2]^d$ with Neumann boundary defined in section 1. Then, this ground state can be described as follows

- in $\gamma + [-1/2, 1/2]^d$, $\varphi_L(\cdot) = \varphi(\cdot - \gamma)$ if $\omega_\gamma = 1$;
- in $\gamma + [-1/2, 1/2]^d$, $\varphi_L(\cdot) = \text{cst}$ if $\omega_\gamma = 0$.

The constants here are chosen to be equal to the constant value of φ near the boundary of $[-1/2, 1/2]^d$, and so that, the thus constructed ground state be normalized.

We can now use the results of [8] to obtain a lower bound on the second eigenvalue of $H_{\omega, L}^N$ that is independent of ω . Indeed, the construction above shows that the maximum and minimum of the (positive normalized) ground state of $H_{\omega, L}^N$ does not depend on ω . Then, Theorem 1.4 of [8] applied to $H_{\omega, L}^N$ and the Neumann Laplacian on the same cube guarantees that the second eigenvalue of $H_{\omega, L}^N$ is larger than cL^{-2} where the constant c does not depend on ω . Hence, the standard upper bound for the integrated density of states by the normalized Neumann counting function (see e.g. [20, 5]) yields

$$N(E) \leq \mathbb{E} \left(\frac{\#\{\text{eigenvalues of } H_{\omega, L}^N \leq E\}}{L^d} \right).$$

If we pick $L = cE^{-1/2}$ for some $c > 0$ sufficiently small, the second eigenvalue of $H_{\omega, L}^N$ is larger than $2E$, this for any realization of ω , we obtain

$$N(E) \leq CE^{d/2}.$$

The standard lower bound for the integrated density of states by the normalized Neumann counting function (see e.g. [20, 5]) yields

$$\mathbb{E} \left(\frac{\#\{\text{eigenvalues of } H_{\omega,L}^D \leq E\}}{L^d} \right) \leq N(E) \quad (3.1)$$

where $H_{\omega,L}^D$ is the Dirichlet restriction of H_ω to $[-1/2, L - 1/2]^d$. Let ψ_L be the (positive normalized) ground state of the Dirichlet Laplacian on $[-1/2, L - 1/2]^d$. And define $\phi_L = \psi_L \cdot \varphi_L$. This smooth function clearly satisfies Dirichlet boundary conditions and one computes

$$\begin{aligned} \langle H_{\omega,L}^D \phi_L, \phi_L \rangle &= \langle (-\Delta + V_\omega) \phi_L, \phi_L \rangle \\ &= \langle (-\Delta \psi_L) \varphi_L, \phi_L \rangle - 2 \langle \nabla \psi_L \cdot \nabla \varphi_L, \phi_L \rangle \\ &= 2 \frac{\pi^2}{L^2} \|\phi_L\|^2 + \|\varphi_L |\nabla \psi_L|\|^2 \leq \frac{C}{L^2} \|\phi_L\|^2. \end{aligned}$$

where in the final step we integrated by parts and used the explicit form of the (positive normalized) ground state of the Dirichlet Laplacian.

Hence, taking $L = CE^{-1/2}$ for some $C > 0$ sufficiently large, (3.1) gives

$$\frac{1}{C} E^{d/2} \leq N(E).$$

Finally, combining the two estimates, we have proved that, for the above random model, the density of states exhibits a van Hove singularity at the bottom of the spectrum that is, for $E \geq 0$, one has

$$\frac{1}{C} E^{d/2} \leq N(E) \leq CE^{d/2}.$$

This completes the proof of Theorem 0.5. □

References

- [1] J. Baker, M. Loss, and G. Stolz. Minimizing the ground state energy of an electron in a randomly deformed lattice. Preprint arXiv:0707.3988.
- [2] F. Germinet and A. Klein. Bootstrap multiscale analysis and localization in random media. *Comm. Math. Phys.*, 222(2):415–448, 2001.
- [3] F. Ghribi. Internal Lifshits tails for random magnetic Schrödinger operators. *J. Funct. Anal.*, 248(2):387–427, 2007.
- [4] P. Hislop and F. Klopp. The integrated density of states for some random operators with nonsign definite potentials. *Jour. Func. Anal.*, 195:12–47, 2002.

- [5] W. Kirsch. Random Schrödinger operators. In A. Jensen H. Holden, editor, *Schrödinger Operators*, number 345 in Lecture Notes in Physics, Berlin, 1989. Springer Verlag. Proceedings, Sonderborg, Denmark 1988.
- [6] W. Kirsch and F. Martinelli. Large deviations and Lifshitz singularities of the integrated density of states of random hamiltonians. *Communications in Mathematical Physics*, 89:27–40, 1983.
- [7] W. Kirsch. Random Schrödinger operators and the density of states. In *Stochastic aspects of classical and quantum systems (Marseille, 1983)*, volume 1109 of *Lecture Notes in Math.*, pages 68–102. Springer, Berlin, 1985.
- [8] W. Kirsch and B. Simon. Comparison theorems for the gap of Schrödinger operators. *J. Funct. Anal.*, 75(2):396–410, 1987.
- [9] W. Kirsch, P. Stollmann, and G. Stolz. Localization for random perturbations of periodic Schrödinger operators. *Random Oper. Stochastic Equations*, 6(3):241–268, 1998.
- [10] F. Klopp. Localization for some continuous random Schrödinger operators. *Communications in Mathematical Physics*, 167:553–570, 1995.
- [11] F. Klopp. Internal Lifshitz tails for random perturbations of periodic Schrödinger operators. *Duke Math. J.*, 98(2):335–396, 1999.
- [12] F. Klopp. Internal Lifshitz tails for Schrödinger operators with random potentials. *J. Math. Phys.*, 43(6):2948–2958, 2002.
- [13] F. Klopp. Une remarque à propos des asymptotiques de Lifshitz internes. *C. R. Math. Acad. Sci. Paris*, 335(1):87–92, 2002.
- [14] F. Klopp and S. Nakamura. In progress.
- [15] F. Klopp, S. Nakamura, F. Nakano, and Y. Nomura. Anderson localization for 2D discrete Schrödinger operators with random magnetic fields. *Ann. Henri Poincaré*, 4(4):795–811, 2003.
- [16] F. Klopp and T. Wolff. Lifshitz tails for 2-dimensional random Schrödinger operators. *J. Anal. Math.*, 88:63–147, 2002. Dedicated to the memory of Tom Wolff.
- [17] H. Najjar. The spectrum minimum for random Schrödinger operators with indefinite sign potentials. *J. Math. Phys.*, 47(1):013515, 13, 2006.
- [18] S. Nakamura. Lifshitz tail for 2D discrete Schrödinger operator with random magnetic field. *Ann. Henri Poincaré*, 1(5):823–835, 2000.

- [19] S. Nakamura. Lifshitz tail for Schrödinger operator with random magnetic field. *Comm. Math. Phys.*, 214(3):565–572, 2000.
- [20] L. Pastur and A. Figotin. *Spectra of random and almost-periodic operators*, volume 297 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1992.
- [21] M. Reed and B. Simon. *Methods of modern mathematical physics. IV. Analysis of operators*. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1978.
- [22] P. Stollmann. *Caught by disorder*, volume 20 of *Progress in Mathematical Physics*. Birkhäuser Boston Inc., Boston, MA, 2001. Bound states in random media.
- [23] I. Veselić. Integrated density of states and Wegner estimates for random Schrödinger operators. In *Spectral theory of Schrödinger operators*, volume 340 of *Contemp. Math.*, pages 97–183. Amer. Math. Soc., Providence, RI, 2004.