



HAL
open science

Josephson Effect through a Magnetic Metallofullerene Molecule

Minchul Lee, Thibaut Jonckheere, Thierry Martin

► **To cite this version:**

Minchul Lee, Thibaut Jonckheere, Thierry Martin. Josephson Effect through a Magnetic Metallofullerene Molecule. 2008. hal-00276921v1

HAL Id: hal-00276921

<https://hal.science/hal-00276921v1>

Preprint submitted on 2 May 2008 (v1), last revised 27 Aug 2008 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Josephson Effect through a Magnetic Metallofullerene Molecule

Minchul Lee,¹ Thibaut Jonckheere,¹ and Thierry Martin^{1,2}

¹*Centre de Physique Théorique, UMR6207, Case 907, Luminy, 13288 Marseille Cedex 9, France*

²*Université de la Méditerranée, 13288 Marseille Cedex 9, France*

We investigate the Josephson effect through a molecular quantum dot magnet connected to superconducting leads. The molecule depicts a metallofullerene where the spin of the magnetic atom inside the carbon shell is assumed to be isotropic. It is coupled to the electron spin on the dot via exchange coupling. Using the numerical renormalization group method we calculate the Andreev levels and the supercurrent and examine intertwined effect of the exchange coupling, Kondo correlation, and superconductivity on the current. Exchange coupling typically suppresses the Kondo correlation so that the system undergoes a phase transition from 0 to π state as $|J|$ increases, where J is the exchange coupling constant. Antiferromagnetic coupling is found to drive exotic transitions: the reentrance to the π state for a small superconducting gap and the restoration of 0 state for large J . We suggest that the asymmetric dependence of supercurrent on J could be used as to detect the sign of J in experiments.

PACS numbers: 73.63.-b, 74.50.+r, 72.15.Qm, 73.63.Kv

Molecular spintronics [1] aims at exploring spin-dependent electronic transport through molecules with intrinsic degrees of freedom such as spin, connected to leads of various nature. On the theoretical and experimental side, recent advances have concerned both coherent [2] and incoherent [3, 4, 5] transport through these molecular quantum dot magnets (MQDM). They consist of a magnetic molecule with either a large [6] or a small anisotropy, as is the case for a metallofullerene.

Here, we provide the first exact computation of the low temperature transport properties of a MQDM connected to superconducting leads using a numerical renormalization group (NRG) approach. The Josephson current allows a diagnosis of the interaction between the intrinsic spin of the molecule, its itinerant electron spin, and the polarization of the leads. It has been known for some time [7, 8, 9, 10], and recently analyzed in experiments [11], that a quantum dot sandwiched between superconducting leads can show a π junction behavior [12]. At the same time, a quantum dot connected to leads at low enough temperatures exhibits the Kondo effect [13]. It was shown [8, 14, 15] that with superconducting leads, at low temperature the zero junction state of the Josephson current is restored when the Kondo temperature exceeds the superconducting gap. The stability of this Kondo phase is put in question in the presence of additional spin degrees for freedom [16] which may compete with Kondo screening. Here the Josephson current flows through an isotropic MQDM which describes a metallofullerene molecule [17]. The electron spin in the carbon shell and the magnetic ion inside it interact via an exchange coupling [4]. We calculate the Andreev level spectrum and the supercurrent and determine the spin of the ground state. We find that the exchange coupling typically suppresses the Kondo effect and drives a transition from 0 to π state. Moreover, antiferromagnetic coupling is found to drive exotic transitions: the reentrance to π state for

small superconducting gap and the restoration of 0 state for large J .

The MQDM connected to two s -wave superconducting leads (inset of Fig. 1) is modeled by a single-impurity Anderson model: $\mathcal{H} = \mathcal{H}_M + \mathcal{H}_L + \mathcal{H}_T$, where

$$\mathcal{H}_M = \epsilon_0 n + U n_\uparrow n_\downarrow + \mathbf{J} \mathbf{S} \cdot \mathbf{S}_e \quad (1)$$

$$\mathcal{H}_L = \sum_{\ell \mathbf{k}} \left[\epsilon_{\mathbf{k}} n_{\ell \mathbf{k}} - \left(\Delta e^{i\phi_\ell} c_{\ell \mathbf{k} \uparrow}^\dagger c_{\ell - \mathbf{k} \downarrow}^\dagger + (h.c.) \right) \right] \quad (2)$$

$$\mathcal{H}_T = \sum_{\ell \mathbf{k} \mu} \left[t d_\mu^\dagger c_{\ell \mathbf{k} \mu} + (h.c.) \right]. \quad (3)$$

Here $c_{\ell \mathbf{k} \mu}$ (d_μ) destroys an electron with energy $\epsilon_{\mathbf{k}}$, and spin μ on lead $\ell = L, R$ (on the carbon cell); $n_{\ell \mathbf{k}}$ and n are occupation operators for the leads and the cell. The single-particle energy ϵ_0 can be tuned by gate voltages. J denotes the exchange energy between the metal ion spin \mathbf{S} and the electron spin $\mathbf{S}_e = \frac{1}{2} \sum_{\mu \mu'} d_\mu^\dagger \boldsymbol{\sigma}_{\mu \mu'} d_{\mu'}$. Δ is the superconducting gap. Except for the finite phase difference $\phi = \phi_L - \phi_R$, the leads are identical and their coupling to the MQDM is symmetric. The hybridization between the molecule and the leads is well characterized by a tunneling rate $\Gamma = \pi \rho_0 |t|^2$, where ρ_0 is the density of states of the leads at the Fermi energy. As we are interested in the low temperature behavior, we concentrate for the most part on the Kondo regime with a localized level $-\epsilon_0 \gg \Gamma$ with large charging energy $U \gg |\epsilon_0|$. Specifically, we choose $\epsilon_0 = -0.1D$ (the band width D is taken as a unit of energy), $\Gamma = 0.01D$, and $U = \infty$ and introduce the bare Kondo temperature $T_K^0 = \sqrt{D\Gamma}/2 \exp[\frac{\pi \epsilon_0}{2\Gamma} (1 + \frac{\epsilon_0}{U})]$ (at $J = \Delta = 0$). The energy spectrum is found with the NRG method [18] extended to superconducting leads [14, 19].

Fig. 1 shows the schematic phase diagram of our system, which constitutes the main result of our study. The junction property switches between 0 and π state, depending on the strengths of J and Δ with respect to T_K^0 .




FIG. 1: (color online) Schematic phase diagram of a metallofullerene superconducting junction system [see inset] at zero phase difference ($\phi = 0$) indicating the 0 state (shaded area) and the π state regions. Each region is divided into two subregions according to the ground-state spin: S and $S-1/2$ for 0_1 and 0_2 regions and $S-1/2$ and $S+1/2$ for π_1 and π_2 regions, respectively. For larger molecular spin $S' > S$ (see the dotted lines), the phase boundaries between 0_1 and $\pi_{1/2}$ are shifted toward smaller $|J|$, and one between 0_2 and π_1 moves toward larger J .

In the absence of exchange coupling ($J = 0$), the system undergoes the Kondo-driven phase transition [8, 14, 15]: The ground-state wave function is of spin singlet kind for $\Delta < \Delta_c \approx 1.84T_K^0$ and of spin doublet for $\Delta > \Delta_c$. In the strong coupling limit ($\Delta < \Delta_c$) the Kondo correlations screen out the localized spin and Cooper pairs tunnel through the Kondo resonance state, resulting in a 0-junction [10, 14, 15]. In the weak coupling limit ($\Delta > \Delta_c$), strong superconductivity in the leads leaves the local spin unscreened and the tunneling of Cooper pairs subject to strong Coulomb interaction acquires an additional phase π , making a π -junction [7, 8, 9, 14, 15].

Finite exchange coupling between electron spins and the metal ion spin introduces another electronic correlation and affects Cooper pair transport. Fig. 2 shows typical variations of Andreev levels and supercurrents with J along the line aa' (see Fig. 1) in the strong coupling limit ($\Delta/T_K^0 = 0.1$). Any finite J clearly induces a splitting in the Andreev bound states and consequently causes them to cross at the Fermi level at $\phi \neq \pi$ (at least for $|J/T_K^0| \lesssim O(1)$); the level crossing otherwise takes place only at $\phi = \pi$. It reflects a sudden change in the physical properties of the ground state: for example, the ground state spin is changed from S to $S-1/2$ for $J > 0$ and to $S+1/2$ for $J < 0$. In parallel with an abrupt change in Andreev levels, the supercurrent-phase relation (SPR) shows a discontinuous sign change (note that $I \propto -\partial E_A/\partial\phi$), culminating in a transition




FIG. 2: (color online) (a) Andreev bound spectra in units of Δ and (b) supercurrents I in units of $I_c^0 \equiv e\Delta/\hbar$ as functions of the phase ϕ in the strong coupling limit ($\Delta/T_K^0 = 0.1$) for various values of J/T_K^0 : see the line aa' in Fig. 1. Here the metal ion spin S is set to $1/2$.

from 0 to π state: two π states, $\pi_{1,2}$ are identified according to the ground-state spin $S \mp 1/2$, respectively. The curve of $I(\phi)$ then has three distinct segments [14]. The central segment resembles that of a short ballistic junction, while the two surrounding segments are parts of π -junction curve. As J grows in magnitude the central segment shrinks and eventually vanishes. The SPR then becomes sinusoidal like in a tunnel junction. It should be noted that the 0- π transition is asymmetric with respect to the sign of J : the transition for $J > 0$ takes place at $\delta E_S \sim T_K^0$, where $\delta E_S = \frac{J}{2}(2S+1)$ is the exchange-coupling energy gap, while the 0 state survives much larger ferromagnetic coupling ($J < 0$). Once the π -junction is fully established, stronger ferromagnetic coupling does not lead to any qualitative change in the SPR, while a second transition back to 0 state is observed for large antiferromagnetic coupling ($J \gg \Delta$). The NRG results distinguish the second 0 state (0_2) from the former one (0_1) in three points: (1) the ground state has spin $S-1/2$ like π_1 phase, (2) the SPR is that of a tunneling junction, and (3) the π_1 - 0_2 transition boundary is independent of ϕ . Figure 3 (c) shows that the critical current has its maximum at $J = 0$ and decreases with $|J|$ rapidly across the phase boundary for $J > 0$ or rather gradually

for $J < 0$. The critical current totally vanishes at the π_1 - 0_2 boundary and increases again slowly with J in 0_2 phase (see the curve for $\Delta/T_K^0 = 0.01$).

The 0 - π transitions (0_1 - π_1 and 0_1 - π_2) can be attributed to the competition between superconducting and Kondo correlations as in the absence of exchange coupling. The relevant parameters are then the Kondo temperature T_K and the superconducting gap Δ , and the 0 - π phase transition occurs when they are comparable to each other: In our choice of parameters the transition happens at $\Delta_c/T_K \approx 1.84$. The exchange coupling manifests itself by renormalizing the Kondo temperature $T_K(J)$. To see this, we applied the poor man's scaling theory to a corresponding Kondo Hamiltonian with no superconductivity and $S = 1/2$: $\mathcal{H}_{\text{KM}} = \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} n_{\mathbf{k}} + J\mathbf{S} \cdot \mathbf{S}_e + (J_K \mathbf{S}_e + J_M \mathbf{S}) \cdot \mathbf{S}_L$, where \mathbf{S}_L is the spin operator for the lead electrons at molecule site. The last term $\mathbf{S} \cdot \mathbf{S}_L$ describing direct coupling between spins of the metal ion and the lead electrons arises during the scaling process. The renormalization group analysis leads to the following scaling equations: together with $J \approx J(\Lambda = D)$,

$$\frac{dJ_K}{d\ln \Lambda} \approx -\rho_0 J_K^2 + \frac{J}{4D} (2J_K J_M - J_M^2) \quad (4a)$$

$$\frac{dJ_M}{d\ln \Lambda} \approx -\rho_0 J_M^2 + \frac{J}{4D} (2J_K J_M - J_K^2). \quad (4b)$$

As the band width Λ is decreased from D to T_K , the coefficient J_K , responsible for the Kondo correlation, diverges and the scaling breaks down. In the presence of finite exchange coupling, however, since $JJ_K J_M > 0$ with $J_M(\Lambda = D) = 0$ and $|J_M| \ll J_K$, the term proportional to J in Eq. (4a) turns out to slow down the flow of J_K and accordingly lowers the Kondo temperature. This point is confirmed by NRG calculations applied in the absence of superconductivity. As can be seen in Fig. 3 (a) and (b), the width of the spectral density for shell electrons, identified as the Kondo temperature $T_K(J)$, decreases with $|J|$ (for $J < 0$ this decrease, being marginal, is not clearly shown with the logarithmic scale). We find out that for the ferromagnetic case the ratio $T_K(J)/T_K^0$ matches well with $\Delta_c(J)/\Delta_c(J=0)$. For the antiferromagnetic case, the Kondo correlation is observed to be suppressed not only by the Kondo peak narrowing but by lowering the peak height.

Antiferromagnetic exchange coupling can, on the other hand, exert a more profound effect than simply renormalizing the Kondo temperature: it gives rise to a reentrant transition to the π state at small Δ and restoration of the 0 state for large J . It is known that small antiferromagnetic exchange coupling ($J \lesssim T_K^0$), studied in the context of side-coupled quantum dot systems [20], can produce a two-stage Kondo effect. After the magnetic moment of carbon shell is screened by conduction electrons below T_K , at a much lower energy scale (denoted as T_K^J) the metal ion spin is screened by the local Fermi liquid that is formed on the shell. T_K^J is then the Kondo




FIG. 3: (color online) Spectral weights $\mathcal{A}_d(\omega)$ for shell electrons coupled to normal leads with antiferromagnetic [(a)] and ferromagnetic [(b)] exchange coupling to metal ion spin for various values of J/T_K^0 (as annotated). (c) Critical currents as functions of J/T_K^0 for different values of Δ/T_K^0 (see the annotations). The arrows locate transition points corresponding to data with the same color. Here we have used $S = 1/2$.

temperature of a magnetic moment screened by electrons of a bandwidth $\sim T_K$ and density of states $\sim 1/(\pi T_K)$ [20]: $T_K^J \sim T_K \exp[-\frac{\pi T_K}{J}]$. The second Kondo effect leads to a Fano resonance and makes a dip in the shell electron density of states as shown in Fig. 3 (a). The dip becomes widened with J and overrides the Kondo peak when $T_K^J \approx T_K$ so that the Kondo effect is completely overridden. As long as $\Delta > T_K^J$, the second Kondo effect does not appear since the superconducting gap blocks any quasi-particle excitation with energy less than Δ . For $\Delta \lesssim T_K^J$, however, Cooper pairs notice the suppression of the Kondo resonance level, and their tunneling is governed by cotunneling under strong Coulomb interaction, forming a π -junction again. Since T_K^J decreases with decreasing J , Δ_c decreases to zero as $J \rightarrow 0$. Note that the extremely small $T_K^J \ll T_K$ (unless $\delta E_S \sim T_K^0$) might make it hard to detect the reentrance even under rather weak thermal fluctuations with $T_K > T > T_K^J$.

The revival of the 0 -state for strong antiferromagnetic coupling can be explained in the picture of cotunneling of Cooper pairs [9]. In weak coupling limit, the fourth-order perturbation theory leads to the supercurrent given by

$$I = \frac{4e}{\hbar} \sin \phi \sum_{\mathbf{k}\mathbf{k}'} t_L^2 t_R^2 \frac{u_{\mathbf{k}} u_{\mathbf{k}'} v_{\mathbf{k}} v_{\mathbf{k}'}}{\mathcal{E}_{\mathbf{k}} \mathcal{E}_{\mathbf{k}'}} \quad (5)$$

$$\times \frac{1}{2S+1} \left(\frac{1}{E_{\mathbf{k}} + E_{\mathbf{k}'}} - \frac{2S+2}{\delta E_S + E_{\mathbf{k}} + E_{\mathbf{k}'}} \right),$$

where $E_{\mathbf{k}} = \sqrt{\Delta^2 + \epsilon_{\mathbf{k}}^2}$, $u_{\mathbf{k}} = \sqrt{(1 + \epsilon_{\mathbf{k}}/E_{\mathbf{k}})/2}$, $v_{\mathbf{k}} =$




FIG. 4: (color online) (a) Andreev bound states in units of Δ and (b) supercurrents I in units of I_C^s as functions of the phase ϕ with $J/T_K^0 = 10$ and $\Delta/T_K^0 = 0.02$ (at $\epsilon_d = -0.1$) while the gate voltage ϵ_d is tuned from -0.1 (bottom) to 0 (top). See the line bb' in Fig. 1.

$\sqrt{(1 - \epsilon_k/E_k)/2}$, and $\mathcal{E}_k = -\epsilon_d - \frac{J}{2}(S+1) - E_k < 0$. For antiferromagnetic coupling, the ground state for the uncoupled system has spin $S - 1/2$. After one electron tunnels through the molecule the system can be in spin eigenstate of either $S - 1/2$ and $S + 1/2$. The latter virtual process, costing more energy by the gap δE_S , turns out to acquire a π phase, contributing to a negative supercurrent. The larger amplitude of this process by a factor $2S+2$ (degeneracy of the spin state $S+1/2$) dominates over spin-preserving process as long as the gap δE_S is small. For a large gap δE_S , however, this process becomes negligible and the sign of the supercurrent is reversed. Note that according to Eq. (5) the SPR is always sinusoidal and the current should vanish at the transition, which is also confirmed in our NRG calculations.

The physical arguments for the $0-\pi$ transitions discussed so far are valid for arbitrary values of the metal ion S , while the phase boundaries are shifted with changing S as shown in Fig. 1. The exchange-coupling energy gap δE_S that is supposed to compete with T_K increases with S so that for larger S the transitions can occur at smaller J . On the other hand, we have observed that the π_1-0_2 transition takes place at slightly larger J for larger S . This is because the increase in the degeneracy factor $2S+2$ overwhelms the decrease in matrix elements due to a larger energy cost by δE_S [see Eq. (5)].

Finally, we present potential experimental manifestations of exchange-coupling-driven $0-\pi$ transition. While the direct control of exchange coupling in molecules is

difficult to achieve, the relative strength J/T_K^0 can be controlled by the gate voltage which can tune the Kondo temperature. Fig. 4 proposes a possibility to observe a double transition (along the line bb' in Fig. 1) as the gate voltage is swept. Note that the double transition is an evidence of strong exchange coupling ($J \gg T_K^0 \gg \Delta$): for examples, with $T_K^0 \sim 3\text{K}$ measured in a recent C_{60} single-molecular transistor, one estimates $J \sim 30\text{K}$. Asymmetry of the phase diagram enables the sign and possibly the amplitude of J to be determined without ambiguity by observing the evolution of the SPR or the critical current.

The authors thank W. Wernsdorfer, F. Balestro, and Mahn-Soo Choi for helpful discussions. This work is supported by ANR-PNANO Contract MolSpintronics No. ANR-06-NANO-27.

-
- [1] A. R. Rocha *et al.*, Nature Materials **4**, 335 (2005).
 - [2] C. Romeike, M. R. Wegewijs, W. Hofstetter, and H. Schoeller, Phys. Rev. Lett. **96**, 196601 (2006); *ibid.* **97**, 206601 (2006).
 - [3] C. Romeike, M. R. Wegewijs, W. Hofstetter, and H. Schoeller, Phys. Rev. Lett. **96**, 196805 (2006).
 - [4] F. Elste, and C. Timm, Phys. Rev. B **71**, 155403 (2005); *ibid.* **73**, 235304 (2006); *ibid.* **73**, 235305 (2006).
 - [5] H. B. Heersche, *et al.*, Phys. Rev. Lett. **96**, 206801 (2006).
 - [6] R. Sessoli, D. Gatteschi, A. Caneschi, and M. A. Novak, Nature (London) **365**, 141 (1993).
 - [7] H. Shiba and T. Soda, Prog. Theor. Phys. **41**, 25 (1969).
 - [8] L. I. Glazman and K. A. Matveev, Pis'ma Zh. Teor. Fiz. **49**, 570 (1989) [JETP Lett. **49**, 659 (1989)].
 - [9] B. I. Spivak and S. A. Kivelson, Phys. Rev. B **43**, 3740 (1991).
 - [10] A. V. Rozhkov and Daniel P. Arovas, Phys. Rev. Lett. **82**, 2788 (1999); Phys. Rev. B **62**, 6687 (2000).
 - [11] J. A. van Dam *et al.*, Nature (London) **442**, 667 (2006).
 - [12] V. V. Ryazanov *et al.*, Phys. Rev. Lett. **86**, 2427 (2001); T. Kontos *et al.*, Phys. Rev. Lett. **89**, 137007 (2002).
 - [13] D. Goldhaber-Gordon *et al.*, Nature (London) **391**, 156 (1998); S. M. Cronenwett, T. H. Oosterkamp, and L. P. Kouwenhoven, Science **281**, 540 (1998).
 - [14] M.-S. Choi, M. Lee, K. Kang, and W. Belzig, Phys. Rev. B **70**, 020502 (2004).
 - [15] F. Siano and R. Egger, Phys. Rev. Lett. **93**, 047002 (2004).
 - [16] F. S. Bergeret, A. Levy-Yeyati, and A. Martin-Rodero, Phys. Rev. B **74**, 132505 (2006).
 - [17] A. Yu. Kasumov, *et al.*, Phys. Rev. B **72**, 033414 (2005).
 - [18] K. G. Wilson, Rev. Mod. Phys. **47**, 773 (1975); H. R. Krishnamurthy, J. W. Wilkins, and K. G. Wilson, Phys. Rev. B **21**, 1003 (1980); *ibid.*, **21**, 1044 (1980).
 - [19] T. Yoshioka and Y. Ohashi, J. Phys. Soc. Jpn. **69**, 1812 (2000).
 - [20] P. S. Cornaglia and D. R. Grempel, Phys. Rev. B **71**, 75305 (2005); R. Zitko and J. Bonca, Phys. Rev. B **73**, 35332 (2006).
 - [21] N. Roch *et al.*, to appear in Nature.