



HAL
open science

On a semilinear elliptic equation with inverse square potential

Haïm Brezis, Louis Dupaigne, Alberto Tesei

► **To cite this version:**

Haïm Brezis, Louis Dupaigne, Alberto Tesei. On a semilinear elliptic equation with inverse square potential. 2008. hal-00288567

HAL Id: hal-00288567

<https://hal.science/hal-00288567>

Preprint submitted on 17 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On a semilinear elliptic equation with inverse-square potential

Haïm Brezis ^{*†}, Louis Dupaigne ^{*} and Alberto Tesei [‡]

In this paper we study existence and nonexistence of solutions $u \geq 0$ of the equation :

$$-\Delta u = \frac{c}{r^2}u + u^p \quad (1)$$

in a ball $B(0, R)$ of \mathbb{R}^N , $N \geq 3$. Here $r = |x|$, $p > 1$ and the coefficient c satisfies the inequality $0 < c \leq c_0$, where $c_0 = (N - 2)^2/4$ is the best constant in the Hardy inequality.

In this study an important role is played by the roots

$$\alpha = \alpha^\pm := (N - 2)/2 \pm \sqrt{c_0 - c}$$

of the equation

$$\alpha^2 - (N - 2)\alpha + c = 0. \quad (2)$$

Observe that $\alpha^+ > \alpha^- > 0$.

Our main result asserts that nontrivial solutions of equation (1) exist if and only if $p < p^+$ where

$$p^+ = 1 + 2/\alpha^-.$$

Theorem 1. *Let $0 \leq c \leq c_0$. For any $p \in (1, p^+)$, there exists a nontrivial solution to equation (1) with u^p and u/r^2 belonging to $L^1(B_R)$ and (1) holds in $\mathcal{D}'(B_R)$.*

The proof of Theorem 1 is straightforward and elementary, except for the limiting value $c = c_0$. The conclusion of Theorem 1 was known in many—but presumably not all—cases (see e.g. [12]). Concerning nonexistence we have

Theorem 2. *Let $0 < c \leq c_0$, $p \geq p^+$. Assume $u \in L^p_{loc}(B_R \setminus \{0\})$, $u \geq 0$ satisfies*

$$-\Delta u - \frac{c}{r^2}u \geq u^p$$

in $\mathcal{D}'(B_R \setminus \{0\})$. Then $u \equiv 0$.

^{*} Laboratoire Jacques-Louis Lions, Université Pierre et Marie Curie, Boîte Courrier 187, 4 place Jussieu, 75252 Paris Cedex 05, France.

brezis@ann.jussieu.fr, dupaigne@ann.jussieu.fr

[†] Department of Mathematics – Hill Center, Rutgers Univ., 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019 USA.

brezis@math.rutgers.edu

[‡] Dipartimento di Matematica “G. Castelnuovo”, Università di Roma “La Sapienza”, P.le A. Moro 5, I-00185 Roma, Italia.

tesei@mat.uniroma1.it

Theorem 2 is reminiscent of the nonexistence results of Brezis-Cabré [2] concerning the so-called very weak solutions to the inequality

$$-\Delta u \geq \frac{u^p}{r^2}, \quad u \geq 0, \quad u \in L_{loc}^p(B_R \setminus \{0\}),$$

for any $p > 1$. The nonexistence aspect in (1) when $p \geq p^+$ was first investigated by Pohozaev-Tesei [11]. However the concept of solution used there was stronger; our concept is the weakest possible.

We also observe that Theorem 2 seems (formally) to contradict the Implicit Function Theorem since there is no solution of $-\Delta u = (c/|x|^2)u + u^p + t$, even when $t > 0$ is small. As observed in [1], this is due to the lack of an appropriate functional space in which to apply the IFT.

Proof of Theorem 1.

Set $p^- = 1 + 2/\alpha^+$ and observe that

$$1 < p^- < \frac{N+2}{N-2} < p^+ \quad \text{for any } 0 < c < c_0 \text{ and}$$

$$\lim_{c \rightarrow 0} p^- = \frac{N}{N-2}, \quad \lim_{c \rightarrow c_0} p^- = \frac{N+2}{N-2},$$

$$\lim_{c \rightarrow 0} p^+ = +\infty, \quad \lim_{c \rightarrow c_0} p^+ = \frac{N+2}{N-2}.$$

We distinguish three cases :

Case 1 : $0 \leq c < c_0$ and $p < \frac{N+2}{N-2}$.

Here the existence of a positive solution $u \in H_0^1(B_R)$ of (1) is a standard and straightforward consequence of the Mountain Pass Theorem. In fact, one can find a radial solution by working in the class of radial functions.

Case 2 : $0 \leq c < c_0$ and $p^- < p < p^+$.

Here we have an explicit solution of (1) of the form $u = A/r^\beta$ with $\beta = 2/(p-1)$, $A > 0$ given by

$$A^{p-1} = -\beta^2 + (N-2)\beta - c > 0,$$

because α^-, α^+ are the roots of (2) and the restriction $\alpha^- < \beta < \alpha^+$ is equivalent to the condition $p^- < p < p^+$. Since $\beta < N-2$, u satisfies (1) in the sense of $\mathcal{D}'(B_R)$.

Case 3 : $c = c_0$ and $1 < p < p^+ = \frac{N+2}{N-2}$.

This case is a little more delicate : here we need the improved Hardy inequality which asserts that

$$\int_{B_R} |\nabla u|^2 \geq c_0 \int_{B_R} \frac{u^2}{r^2} + c_q \|u\|_{L^q(B_R)}^2,$$

for any $1 \leq q < \frac{2N}{N-2}$ and $u \in C_0^\infty(B_R)$. See [5]. Let H be the Hilbert space obtained by completing $C_0^\infty(B_R)$ with respect to the scalar product

$$a(u, v) = \int_{B_R} \nabla u \cdot \nabla v - c_0 \int_{B_R} uv/r^2.$$

Clearly H is contained in every $L^q(B_R)$ with $1 \leq q < \frac{2N}{N-2}$ with continuous injection. Moreover the injection is compact. This fact is due to H. Brezis and the proof is presented in Lemmas 3.2, 3.3 of [7]. We may then use the Mountain Pass Theorem in H and the (PS) condition is satisfied.

Proof of Theorem 2.

We will use the following lemma :

Lemma 1. *Let $\Sigma \subset\subset \Omega$ be a closed set of zero (newtonian) capacity and assume that $u, f \in L^1_{loc}(\Omega \setminus \Sigma)$ are two nonnegative functions such that*

$$-\Delta u \geq f \quad \text{in } \mathcal{D}'(\Omega \setminus \Sigma).$$

Then $u, f \in L^1_{loc}(\Omega)$ and

$$-\Delta u \geq f \quad \text{in } \mathcal{D}'(\Omega).$$

Furthermore given any smooth subdomain $\Omega' \subset\subset \Omega$, if $v \in L^1(\Omega')$ is the solution of

$$\begin{cases} -\Delta v = f & \text{in } \Omega' \\ v = 0 & \text{on } \partial\Omega', \end{cases}$$

in the sense that

$$\int v(-\Delta\phi) = \int f\phi \quad \forall \phi \in C^2(\bar{\Omega}') \quad \text{such that} \quad \phi|_{\partial\Omega'} \equiv 0,$$

then

$$u \geq v \quad \text{a.e. in } \Omega'. \tag{3}$$

Proof of Lemma 1

This lemma can be seen as a fairly easy consequence of Theorem 7.7 in [9]. It is also closely related to a result in [4]. We provide a proof for completeness. Let $u_k = \min(u, k)$, $k > 0$, which by Kato's Lemma (see [10]) satisfies

$$-\Delta u_k \geq f_k \quad \text{in } \mathcal{D}'(\Omega \setminus \Sigma), \quad (4)$$

where $f_k := f\chi_{\{u < k\}}$. Since $-\Delta u_k$ is a nonnegative distribution on $\Omega \setminus \Sigma$, it extends to a nonnegative measure on $\Omega \setminus \Sigma$. Since u_k is bounded, it follows from a Gagliardo-Nirenberg-type inequality that $u_k \in H_{loc}^1(\Omega \setminus \Sigma)$. We show next that in fact $u_k \in H_{loc}^1(\Omega)$. We first take a nonnegative function $\phi \in C_0^\infty(\Omega)$ and a sequence $\phi_n \in C_0^\infty(\Omega \setminus \Sigma)$ converging to ϕ in $H^1(\Omega)$. This is always possible since $\text{cap}_\Omega(\Sigma) = 0$ (take e.g. $\phi_n = \phi(1 - \chi_n)$ where $\chi_n = 1$ near Σ and $\|\chi_n\|_{H^1} \rightarrow 0 = \text{cap}_\Omega(\Sigma)$). We then have, with $C_k = e^k$,

$$\begin{aligned} \int |\nabla u_k|^2 \phi_n^2 &\leq C_k \int e^{-u_k} |\nabla u_k|^2 \phi_n^2 = -C_k \int \phi_n^2 \nabla(e^{-u_k}) \cdot \nabla u_k \\ &= C_k \left(2 \int e^{-u_k} \phi_n \nabla \phi_n \cdot \nabla u_k + \int e^{-u_k} \Delta u_k \phi_n^2 \right) \\ &\leq 2C_k^2 \int e^{-u_k} |\nabla \phi_n|^2 + \frac{1}{2} \int e^{-u_k} |\nabla u_k|^2 \phi_n^2, \end{aligned}$$

so that

$$\int |\nabla(u_k \phi_n)|^2 \leq C'_k \int |\nabla \phi_n|^2.$$

Passing to the limit as $n \rightarrow \infty$ in the above inequality implies that $u_k \in H_{loc}^1(\Omega)$.

We next show that

$$-\Delta u_k \geq f_k \quad \text{in } \mathcal{D}'(\Omega). \quad (5)$$

Take ϕ and ϕ_n as above so that by (4),

$$\int u_k(-\Delta \phi_n) \geq \int f_k \phi_n. \quad (6)$$

Now, as $n \rightarrow \infty$,

$$\int u_k(-\Delta \phi_n) = \int \nabla u_k \nabla \phi_n \rightarrow \int \nabla u_k \nabla \phi = - \int u_k \Delta \phi.$$

Passing to the limit in (6) as $n \rightarrow \infty$, we thus obtain (5).

In particular u_k is superharmonic in Ω and given almost any $x \in \Omega$ and any ball $B \subset \Omega$ centered at x , we have

$$u_k(x) \geq \frac{1}{|B|} \int_B u_k(y) dy. \quad (7)$$

Now, since $u \in L_{loc}^1(\Omega \setminus \Sigma)$ (and $|\Sigma| = 0$), $u_k \rightarrow u$ a.e. in Ω as $k \rightarrow \infty$ and u is finite almost everywhere. By Fatou's Lemma we then conclude from (7) that for almost every ball B ,

$$\int_B u < \infty,$$

which means that $u \in L^1_{loc}(\Omega)$. Using this information, we can now easily pass to the limit in (5) and conclude that $f \in L^1_{loc}(\Omega)$ and that

$$-\Delta u \geq f \quad \text{in } \mathcal{D}'(\Omega).$$

It only remains to prove (3). We let ρ_n be a standard smooth mollifier and let $u_n = u * \rho_n$, $f_n = f * \rho_n$ so that for n large enough $-\Delta u_n \geq f_n$ and $u_n \geq 0$ in Ω' . By the Maximum Principle

$$u_n \geq v_n \quad \text{in } \Omega',$$

where v_n solves

$$\begin{cases} -\Delta v_n = f_n & \text{in } \Omega' \\ v_n = 0 & \text{on } \partial\Omega'. \end{cases}$$

As $n \rightarrow \infty$, $u_n \rightarrow u$ in $L^1(\Omega')$, $f_n \rightarrow f$ in $L^1(\Omega')$ and (by Lemma 4 in [3]) $v_n \rightarrow v$ in $L^1(\Omega')$, which yields the desired conclusion. \square

Proof of Theorem 2

We argue by contradiction and assume that $u \not\equiv 0$. By Lemma 1, $u \in L^p_{loc}(B_R)$, $u/r^2 \in L^1_{loc}(B_R)$ and by the mean-value formula for superharmonic functions, given $R' \in (0, R)$, there exists $\epsilon > 0$ such that $u \geq \epsilon$ a.e. in $B_{R'}$. Let $\lambda := \epsilon^{q-1}/2 > 0$ and v_0 be the solution of

$$\begin{cases} -\Delta v_0 = \lambda & \text{in } B_{R'} \\ v_0 = 0 & \text{on } \partial B_{R'}. \end{cases}$$

Once more by Lemma 1, we have

$$0 \leq v_0 \leq u. \tag{8}$$

Next, for $n \geq 1$, define inductively v_n by

$$\begin{cases} -\Delta v_n = \frac{c}{|x|^2} v_{n-1} + \frac{1}{2} v_{n-1}^p + \lambda & \text{in } B_{R'} \\ v_n = 0 & \text{on } \partial B_{R'}. \end{cases}$$

In order to have a well-defined solution v_n (in the sense of Lemma 4 in [3]) it suffices to prove that $f := \frac{c}{|x|^2} v_{n-1} + \frac{1}{2} v_{n-1}^p \in L^1(B_{R'})$. When $n = 0$, this follows from (8) and Lemma 1 which implies that $\frac{c}{|x|^2} u + \frac{1}{2} u^p \in L^1(B_{R'})$. Assume now that $v_{n-1} \in L^1(B_{R'})$ is well-defined. Using the Maximum Principle, it is easy to see that

$$0 \leq v_0 \leq v_1 \leq \dots \leq v_{n-1} \leq u,$$

whence $f \in L^1(B_{R'})$ and by the Maximum Principle again $0 \leq v_{n-1} \leq v_n \leq u$.

By monotone convergence, letting $v := \lim_{n \rightarrow \infty} v_n$, we have that

$$\begin{cases} -\Delta v = \frac{c}{|x|^2} v + \frac{1}{2} v^p + \lambda & \text{in } B_{R'} \\ v = 0 & \text{on } \partial B_{R'}, \end{cases}$$

in the sense that given any $\phi \in C^2(\bar{B}_{R'})$ such that $\phi|_{\partial B_{R'}} \equiv 0$,

$$\int v(-\Delta \phi) = \int \frac{c}{|x|^2} v \phi + \frac{1}{2} \int v^p \phi + \lambda \int \phi.$$

This contradicts Theorem 1 of [6].

□

Remark 1. Theorems 1 and 2 extend to more general situations— for example, when u^p is replaced by $|x|^{-\beta} u^q$. Assume $0 < c \leq c_0$ and set $q^+ = 1 + \frac{2-\beta}{\alpha^-}$, where α^- is as above. The conclusions of Theorems 1 and 2 remain valid with p^+ replaced by q^+ .

Remark 2. The argument presented in the proof of Theorem 2 may be used to provide a slightly simpler proof of Theorem 1 in [2].

Remark 3. Theorem 2 can be extended to problems of the type

$$-\Delta u = \frac{c}{\text{dist}(x, \Sigma)^2} u + u^p,$$

where $c > 0$ is a small constant, Σ is a smooth compact manifold of codimension $k \geq 3$ and p is larger than some critical exponent, which can be computed explicitly in terms of k and c . The argument is the same as in the proof of Theorem 2 except that the result of [6] is replaced by a result from [8].

Acknowledgments. The three authors are partially supported by an EC Grant through the RTN Program "Fronts-Singularities", HPRN-CT-2002-00274. The first author (H.B.) is also a member of the Institut Universitaire de France.

References

- [1] H. Brezis, *Is there failure of the Inverse Function Theorem ?*, Proceedings of the Workshop held at the Morningside Center of Mathematics, Chinese Academy of Science, Beijing, June 1999.
- [2] H. Brezis and X. Cabré, *Some simple nonlinear PDE's without solutions*, Bull UMI, **1** (1998), 223–262.
- [3] H. Brezis, X. Cabré, Y. Martel and A. Ramiandrisoa, *Blow-up for $u_t - \Delta u = g(u)$ revisited*, Adv. Diff. Eq., **1** (1996), 73–90.

- [4] H. Brezis and P.L. Lions, *A note on isolated singularities for linear elliptic equations*, Mathematical analysis and applications, Adv. in Math. Suppl. Stud., **7a**, Academic Press, New York–London, 1981, 263–266.
- [5] H. Brezis and J.L. Vazquez, *Blow-up solutions of some nonlinear elliptic problems*, Rev. Mat. Univ. Complut. Madrid, **10** (1997), 443–469.
- [6] L. Dupaigne, *A nonlinear elliptic PDE with the inverse-square potential*, J. d'Analyse Mathématique, **86** (2002), 359–398.
- [7] J. Dávila and L. Dupaigne, *Comparison principles for PDE's with a singular potential*, Proc. Roy. Soc. Edinburgh, **133A** (2003), 61–83.
- [8] L. Dupaigne and G. Nedev, *Semilinear elliptic PDE's with a singular potential*, Adv. Differential Equations, **7** (2002), 973–1002.
- [9] L.L. Helms, *Introduction to potential theory*, New York, Wiley and Sons, 1969.
- [10] T. Kato, *Schrödinger operators with singular potentials*, Israel J. Math., **13** (1972), 135–148.
- [11] S.I. Pohozaev and A. Tesei, *Nonexistence of local solutions to semilinear partial differential inequalities*, Nota Scientifica 01/28, Dip. Mat. Università "La Sapienza", Roma (2001).
- [12] S. Terracini, *On positive entire solutions to a class of equations with a singular coefficient and critical exponent*, Adv. Differential Equations, **1** (1996), 241–264.