



HAL
open science

Renewal series and square-root boundaries for Bessel processes

Nathanael Enriquez, Christophe Sabot, Marc Yor

► **To cite this version:**

Nathanael Enriquez, Christophe Sabot, Marc Yor. Renewal series and square-root boundaries for Bessel processes. 2008. hal-00289151

HAL Id: hal-00289151

<https://hal.science/hal-00289151>

Preprint submitted on 19 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RENEWAL SERIES AND SQUARE-ROOT BOUNDARIES FOR BESSEL PROCESSES

NATHANAËL ENRIQUEZ, CHRISTOPHE SABOT, AND MARC YOR

Abstract. We show how a description of Brownian exponential functionals as a renewal series gives access to the law of the hitting time of a square-root boundary by a Bessel process. This extends classical results by Breiman and Shepp, concerning Brownian motion, and recovers by different means, extensions for Bessel processes, obtained independently by Delong and Yor.

Let B_t be the standard real valued Brownian motion and for $\nu > 0$, introduce the geometric Brownian motion $\mathcal{E}_t^{(-\nu)}$ and its exponential functional $\mathcal{A}_t^{(-\nu)}$

$$\begin{aligned}\mathcal{E}_t^{(-\nu)} &:= \exp(B_t - \nu t) \\ \mathcal{A}_t^{(-\nu)} &:= \int_0^t (\mathcal{E}_s^{(-\nu)})^2 ds.\end{aligned}$$

Lamperti's representation theorem [5] applied to $\mathcal{E}_t^{(-\nu)}$ states

$$(0.1) \quad \mathcal{E}_t^{(-\nu)} = R_{\mathcal{A}_t^{(-\nu)}}^{(-\nu)}$$

where $(R_u^{(-\nu)}, u \leq T_0(R^{(-\nu)}))$ denotes the Bessel process of index $(-\nu)$ (equivalently of dimension $\delta = 2(1 - \nu)$), starting at 1, which is an \mathbb{R}_+ -valued diffusion with infinitesimal generator $\mathcal{L}^{(-\nu)}$ given by

$$\mathcal{L}^{(-\nu)} f(x) = \frac{1}{2} f''(x) + \frac{1 - 2\nu}{2x} f'(x), \quad f \in C_b^2(\mathbb{R}_+^*).$$

Let us remark that, in the special case $\nu = 1/2$, equation (0.1) is nothing else but the Dubins-Schwarz representation of the exponential martingale $\mathcal{E}_t^{(-1/2)}$ as Brownian motion time changed with $\mathcal{A}_t^{(-1/2)}$.

For a short summary of relations between Bessel processes and exponentials of Brownian motion, see e.g. Yor [10].

Let us consider now the following random variable Z , which is often called a perpetuity in the mathematical finance literature:

$$Z := \mathcal{A}_\infty^{(-\nu)} = \int_0^\infty (\mathcal{E}_s^{(-\nu)})^2 ds$$

We deduce directly from (0.1) that

$$\mathcal{A}_\infty^{(-\nu)} = T_0(R^{(-\nu)})$$

2000 *Mathematics Subject Classification.* 60G40, 60J57.

Key words and phrases. Bessel processes, renewal series, exponential functionals, square-root boundaries.

where $T_0 := \inf\{u : X_u = 0\}$, and it is well-known (see [4], [11]), that

$$(0.2) \quad \mathcal{A}_\infty^{(-\nu)} \stackrel{(law)}{=} \frac{1}{2\gamma_\nu}$$

where γ_ν is a gamma variable with parameter ν (i.e. with density $\frac{1}{\Gamma(\nu)}x^{\nu-1}e^{-x}\mathbf{1}_{\mathbb{R}_+}$).

Our main result characterizes the law of the hitting time of a parabolic boundary by $R_u^{(-\nu)}$ which corresponds to a Bessel process of dimension $d < 2$.

Theorem 1. *Let $0 < b < c$, and $\sigma := \inf\{u : (R_u^{(-\nu)})^2 = \frac{1}{c}(b + u)\}$ with $R_0^{(-\nu)} = 1$.*

$$(0.3) \quad E[(b + \sigma)^{-s}] = c^{-s} \frac{E[(1 + 2b\gamma_{\nu+s})^{-s}]}{E[(1 + 2c\gamma_{\nu+s})^{-s}]}, \quad \text{for any } s \geq 0$$

Proof: using the strong Markov property and the stationarity of the increments of Brownian motion, we obtain that for any stopping time τ of the Brownian motion

$$\mathcal{A}_\infty^{(-\nu)} =: Z = \mathcal{A}_\tau^{(-\nu)} + (\mathcal{E}_\tau^{(-\nu)})^2 Z'$$

where Z' is independent of $(\mathcal{A}_\tau^{(-\nu)}, \mathcal{E}_\tau^{(-\nu)})$ and $Z \stackrel{(law)}{=} Z'$.

This implies, by (0.1), that Z satisfies the following affine equation (see [8] for a survey about these equations)

$$(0.4) \quad \mathcal{A}_\infty^{(-\nu)} =: Z = \mathcal{A}_\tau^{(-\nu)} + (R_{\mathcal{A}_\tau^{(-\nu)}}^{(-\nu)})^2 Z'$$

where Z' is independent of $(\mathcal{A}_\tau^{(-\nu)}, R_{\mathcal{A}_\tau^{(-\nu)}}^{(-\nu)})$ and $Z \stackrel{(law)}{=} Z'$.

Obviously, $\sigma < T_0(R^{(-\nu)})$. Taking now :

$$\tau = \inf\{t : (R_{\mathcal{A}_t^{(-\nu)}}^{(-\nu)})^2 = \frac{1}{c}(b + \mathcal{A}_t^{(-\nu)})\}$$

we get $\mathcal{A}_\tau^{(-\nu)} = \sigma$, and the identity in law

$$(0.5) \quad b + Z \stackrel{(law)}{=} (b + \sigma)\left(1 + \frac{Z}{c}\right)$$

where the variables σ and Z on the right-hand side are independent.

As a result, we obtain the Mellin-Stieltjes transform of σ :

$$E[(b + \sigma)^{-s}] = c^{-s} \frac{E[(b + Z)^{-s}]}{E[(c + Z)^{-s}]}$$

But, from (0.2)

$$E[(b + \sigma)^{-s}] = c^{-s} \frac{E[(2\gamma_\nu)^s \frac{1}{(1+2b\gamma_\nu)^s}]}{E[(2\gamma_\nu)^s \frac{1}{(1+2c\gamma_\nu)^s}]}$$

which gives the result. \square

One can now use the duality between the laws of Bessel processes of dimension d and $4 - d$ to get the analogous result of Theorem 1, and recover the result of Delong [2], [3], and Yor [9] which deals with the case $d \geq 2$.

Theorem 2. *Let $0 < b < c$, and $\sigma := \inf\{u : (R_u^{(\nu)})^2 = \frac{1}{c}(b + u)\}$ with $R_0^{(\nu)} = 1$.*

$$(0.6) \quad E[(b + \sigma)^{-s}] = c^{-s} \frac{E[(1 + 2b\gamma_s)^{-s+\nu}]}{E[(1 + 2c\gamma_s)^{-s+\nu}]}, \quad \text{for any } s \geq 0.$$

Proof : it is based on the following classical relation between the laws of the Bessel processes with indices ν and $-\nu$:

$$(0.7) \quad \mathcal{P}_x^{(\nu)}|_{\mathcal{F}_t} = \frac{(X_{t \wedge T_0})^{2\nu}}{x^{2\nu}} \cdot \mathcal{P}_x^{(-\nu)}|_{\mathcal{F}_t}$$

which implies that

$$E_1^{(\nu)}[(b + \sigma)^{-s}] = E_1^{(-\nu)}[X_\sigma^{2\nu}(b + \sigma)^{-s}] = \frac{1}{c^\nu} E_1^{(-\nu)}[(b + \sigma)^{-s+\nu}]$$

Theorem 1 gives the result. □

Finally, it is easily shown, thanks to the classical representations of the Whittaker functions (see Lebedev [6]), that the right-hand sides of (0.3) and (0.6) are expressed in terms of ratios of Whittaker functions.

Acknowledgement: We would like to thank Daniel Dufresne for useful and enjoyable discussions on the subject.

REFERENCES

- [1] Breiman, L. First exit times from a square root boundary. (1967) Proc. Fifth Berkeley Sympos. Math. Statist. and Probability (Berkeley, Calif., 1965/66), Vol. II: Contributions to Probability Theory, Part 2 pp. 9–16 Univ. California Press, Berkeley, Calif.
- [2] Delong, D.M. Crossing probabilities for a square-root boundary for a Bessel process. Comm. Stat. A - Theory Methods. 10 (1981), no 21, 2197–2213.
- [3] Delong, D. M. Erratum: "Crossing probabilities for a square root boundary by a Bessel process" Comm. Statist. A - Theory Methods 12 (1983), no. 14, 1699.
- [4] Dufresne, D. The distribution of a perpetuity, with applications to risk theory and pension funding. Scand. Actuarial J. (1990) 39–79.
- [5] Lamperti, J. Semi-stable Markov processes. I. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete 22 (1972), 205–225.
- [6] Lebedev, N. N. Special functions and their applications. Dover (1972).
- [7] Shepp, L. A. A first passage problem for the Wiener process. Ann. Math. Statist. 38 (1967) 1912–1914.
- [8] Vervaat, W. On a stochastic difference equation and a representation of nonnegative infinitely divisible random variables. Adv. in Appl. Probab. 11 (1979), no. 4, 750–783.
- [9] Yor, M. On square-root boundaries for Bessel processes, and pole-seeking Brownian motion. Stochastic analysis and applications (Swansea, 1983), 100–107, Lecture Notes in Math., 1095, Springer, Berlin, 1984.
- [10] Yor, M. On some exponential functionals of Brownian motion. Adv. Appl. Prob. 24 (1992), 509–531.
- [11] Yor, M. Sur certaines fonctionnelles exponentielles du mouvement brownien réel. J. Appl. Prob 29 (1992), 202–208.

LABORATOIRE MODAL'X, UNIVERSITÉ PARIS 10, 200 AVENUE DE LA RÉPUBLIQUE, 92000
NANTERRE, FRANCE

E-mail address: `nenriquez@u-paris10.fr`

UNIVERSITÉ DE LYON, UNIVERSITÉ LYON 1, INSTITUT CAMILLE JORDAN, CNRS UMR 5208,
43, BOULEVARD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX, FRANCE

E-mail address: `sabot@math.univ-lyon1.fr`

LABORATOIRE DE PROBABILITÉS ET MODÈLES ALÉATOIRES, CNRS UMR 7599, UNIVERSITÉ
PARIS 6, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE

E-mail address: `deaproba@proba.jussieu.fr`