



**HAL**  
open science

## Young measures, Cartesian maps, and polyconvexity

Patrick Bernard, Ugo Bessi

► **To cite this version:**

Patrick Bernard, Ugo Bessi. Young measures, Cartesian maps, and polyconvexity. Journal of the Korean Mathematical Society, 2010, 47 (2), pp.331-350. 10.4134/JKMS.2010.47.2.331 . hal-00294644v1

**HAL Id: hal-00294644**

**<https://hal.science/hal-00294644v1>**

Submitted on 10 Jul 2008 (v1), last revised 6 May 2010 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Young measures, Cartesian maps, and polyconvexity

Patrick Bernard\* and Ugo Bessi

April 2008

---

Patrick Bernard, Université Paris-Dauphine,  
CEREMADE, UMR CNRS 7534  
Pl. du Maréchal de Lattre de Tassigny  
75775 Paris Cedex 16, France  
patrick.bernard@ceremade.dauphine.fr

Ugo Bessi, Università di Roma 3  
Dipartimento di Matematica  
Largo S. Leonardo Murialdo 1  
00146 Roma, Italia  
bessi@matrm3.mat.uniroma3.it

---

**Abstract.** We consider the variational problem consisting of minimizing a polyconvex integrand for maps between manifolds. We offer a simple and direct proof of the existence of a minimizing map. The proof is based on Young measures.

---

**Résumé.** On considère le problème variationnel consistant à minimiser un intégrand polyconvexe pour des applications entre variétés. On donne une preuve simple et directe de l'existence d'une application minimisante en utilisant les mesures d'Young.

---

**Riassunto.** Consideriamo il problema di minimizzare un funzionale policonvesso tra varietà. Diamo una dimostrazione semplice e diretta dell'esistenza di un minimo usando le misure di Young.

---

MSC: 49Q20, 49J45.

---

\*membre de l'IUF

# 1 Introduction

Let  $\bar{N}$  be a compact Riemannian manifold with boundary of dimension  $n$  and  $M$  be a complete Riemannian manifold of dimension  $m$ . We denote by  $N$  the interior of  $\bar{N}$ , and by  $dt$  the non-negative Borel measure on  $N$  associated to the metric. It can be defined as the  $n$ -Hausdorff measure associated to the Riemannian distance, and it exists also if  $N$  is not orientable. This measure is also characterized by the fact that

$$\int_D dt = \left| \int_D \Omega \right|$$

for each embedded disk  $D \subset M$ , where  $\Omega$  is any of the two unitary volume forms on  $D$ . We set  $p = \min\{m, n\}$ . We want to discuss the classical question of minimizing the quantity

$$\int_N L(t, u(t), du_t) dt$$

on appropriate classes of maps  $u : N \rightarrow M$ . We recall some terminology about Lagrangians:

**Convexity :** Given  $k \in \{1, \dots, \min(m, n)\}$ , we say that the integrand  $L(t, x, v)$  is  $k$ -convex if it can be written in the form

$$L(t, x, v) = \mathbf{L}(t, x, v, \wedge_2 v, \dots, \wedge_k v) = \mathbf{L}_{t,x}(v, \wedge_2 v, \dots, \wedge_k v)$$

with a Borel measurable function  $\mathbf{L}$  such that

$$\mathbf{L}_{t,x} : \mathcal{L}(T_t N, T_x M) \times \mathcal{L}(\wedge_2 T_t N, \wedge_2 T_x M) \cdots \times \mathcal{L}(\wedge_k T_t N, \wedge_k T_x M) \rightarrow \mathbb{R} \cup \{+\infty\}$$

is convex for each  $t$  and  $x$ . When  $k = 1$ , this just says that  $L$  is convex in  $v$ ; when  $k = p$ , this hypothesis is usually called polyconvexity. Let us explain the notations. We denote by  $\mathcal{L}(E, F)$  the set of linear maps between  $E$  and  $F$ . The space  $\wedge_i E$  is the vector space  $E \wedge E \dots \wedge E$  generated by elements of the form  $e_1 \wedge e_2 \dots \wedge e_i$ ,  $e_j \in E$ . Given  $v \in \mathcal{L}(E, F)$ , we denote by  $\wedge_i v \in \mathcal{L}(\wedge_i E, \wedge_i F)$  the linear map such that

$$\wedge_i v(e_1 \wedge \dots \wedge e_i) = v(e_1) \wedge v(e_2) \dots \wedge v(e_i).$$

If  $E$  has dimension  $n$  and  $F$  has dimension  $m$ , the dimension of  $\wedge_i E$  is  $C_n^i$  (binomial coefficient) and  $\wedge_i v$  can be represented by a  $C_n^i \times C_m^i$  matrix whose coefficients are the determinants of all  $i \times i$  sub-matrices that can be extracted from the matrix representing  $v$ .

**Regularity :** We say that  $L : J^1(N, M) \rightarrow \mathbb{R} \cup \{+\infty\}$  is a normal integrand if it is a Borel measurable function and if, for almost all  $t \in N$ , the function  $(x, v) \mapsto L(t, x, v)$  is lower semi-continuous.

**Coercivity :** We say that  $L$  is  $k$ -superlinear if there exists a superlinear function  $l : [0, \infty) \rightarrow \mathbb{R}$  such that

$$L(t, x, v) \geq l(\|v\| + \|\wedge_2 v\| + \dots + \|\wedge_k v\|)$$

for all  $(t, x, v) \in J^1(N, M)$ .

Our goal is to explain a compact and simple approach to this kind of problems. We do not present real novelties, and much of the techniques we will use can be found in [14] or [12, 13]. Yet we believe it is not useless to present the short path to Theorem 1 that follows. This work started with an attempt to extend the methods of [7] to higher dimension.

We will define, by studying relevant sets of Young measures, sets  $\text{cart}_k(N, M)$  of maps  $u : N \rightarrow M$  such that

$$W^{1,n}(N, M) \subset \text{cart}_{\min\{n,m\}}(N, M) \subset \text{cart}_k(N, M) \subset \text{cart}_1(N, M) = W^{1,1}(N, M).$$

In the case  $k = \min\{n, m\}$  the set that we denote  $\text{cart}_k(N, M)$  is similar to the set denoted  $\text{cart}^1(N, M)$  in [12, 13], but our presentation is quite different.

In order to state appropriately a variational problem, it is useful to specify boundary conditions. We assume that the boundary  $\partial N$  of  $\bar{N}$  is not empty and we fix a map  $u_0 \in \text{cart}_k(N, M)$ . We denote by  $\text{cart}_k(N, M; u_0)$  the set of maps  $u$  in  $\text{cart}_k(N, M)$  such that the trace of  $u$  in  $L^1(\partial N, M)$  is equal to the trace of  $u_0$ . These traces are well defined (at least in the case where  $M$  is a Euclidean space) because  $\text{cart}_k(N, M) \subset W^{1,1}(N, M)$ , and each element of  $W^{1,1}(N, M)$  has a unique boundary trace in  $L^1$  (see for instance [11] for the definition); we recall that the integration by parts formula holds for this trace. In the case where  $M$  is a manifold, we shall give the precise definition of  $\text{cart}_k(N, M; u_0)$  in Section 3. Our goal is to provide a short and direct proof of the following result:

**Theorem 1.** *Let  $k \in \{1, \dots, \min(m, n)\}$  be given and let  $L : J^1(N, M) \longrightarrow \mathbb{R} \cup \{+\infty\}$  be a  $k$ -convex and  $k$ -superlinear normal integrand. Let  $u_0 \in \text{cart}_k(N, M)$  be given, such that*

$$\int_N L(t, u_0(t), du_0(t)) dt < \infty.$$

*Then there exists a map  $u \in \text{cart}_k(N, M; u_0)$  which minimizes the integral*

$$\int_N L(t, u(t), du(t)) dt$$

*in  $\text{cart}_k(N, M; u_0)$ .*

Note that, in general, we may have

$$\inf_{u \in W^{1,1}(N, M; u_0)} \int_N L(t, u(t), du_t) dt < \min_{u \in \text{cart}_k(N, M; u_0)} \int_N L(t, u(t), du_t) dt$$

and, even if  $u_0$  is smooth,

$$\min_{u \in \text{cart}_k(N, M; u_0)} \int_N L(t, u(t), du_t) dt < \inf_{u \in C^1(N, M; u_0)} \int_N L(t, u(t), du_t) dt.$$

The paper [5] contains an example in which  $N$  has dimension 1 and the minimum on  $W^{1,1}(N, \mathbb{R}^n) = \text{cart}_1(N, \mathbb{R}^n; u_0)$  is smaller than the infimum on  $C^1$ . We shall see an example of the first gap phenomenon in section 4.2.

Theorem 1 is a slight extension on the seminal results of Ball, [3]. Compared to this work and to classical papers on polyconvexity, our proof works under slightly weaker coercivity. Our Theorem reduces to the famous Tonelli theorem in the case  $n = 1$ , and to the famous De Giorgi Theorem in the case  $m = 1$ . Several extensions are known, which go much beyond what we plan to expose. First, the kind of convexity hypothesis can be relaxed to the so-called quasi-convexity, but then one has to add more stringent growth conditions, see [10, 14, 9]. Second, one can, in certain circumstances, relax the coercivity condition to the case when  $L$  has only linear growth, by using cartesian currents and functions of bounded variations. Excellent surveys of these methods are in [12, 13, 9].

Our approach is based on Young measures, also called parametrized measures. A survey on the use of Young measure is the book [14]. Many of our techniques are adapted from this book. In section 2, we define the various sets of Young measures that are useful, in particular the set of Generalized Maps, on which it is appropriate to relax the variational problem. We study the structure of generalized maps and conclude that minimizing generalized maps correspond to minimizing maps. In section 3, we prove a compactness results under boundary conditions. In section 4, we briefly expose how the various tools exposed in sections 2 and 3 lead to a proof

of Theorem 1. We also collect various related remarks. Finally, we recall in the Appendix some relevant facts on the topology of some spaces of measures.

We end this introductory section collecting some notation and material on  $n$ -vectors and  $n$ -forms we shall need in the following.

### 1.1 Some algebra

Let  $E$  and  $F$  be two Euclidean vector spaces of dimension  $n$  and  $m$ . It will be convenient to set  $p := \min(m, n)$ . We denote by  $\mathcal{L}(E, F)$  the set of linear maps between  $E$  and  $F$ . Recall that the vector space  $\wedge_l E$  is endowed with a natural inner product (which is induced from the inner product of  $E$ ). This inner product can be characterized by the property that

$$\langle v_1 \wedge \dots \wedge v_l, w_1 \wedge \dots \wedge w_l \rangle = \det(G)$$

where  $G \in M^{l,l}(\mathbb{R})$  is the Gram matrix  $G_{i,j} = \langle v_i, w_j \rangle$  and  $\det G$  is the determinant of  $G$ . Note then that

$$\|v_1 \wedge \dots \wedge v_l\| := \langle v_1 \wedge \dots \wedge v_l, v_1 \wedge \dots \wedge v_l \rangle^{1/2} = 1$$

if  $(v_1, \dots, v_l)$  is orthonormal in  $E$ . Given  $a \in \mathcal{L}(E, F)$ , we denote by  $\|a\| := \sup_{\|x\|_E \leq 1} \|a(x)\|_F$  its norm and by  $\wedge_l a \in \mathcal{L}(\wedge_l E, \wedge_l F)$  the unique linear map which satisfies

$$\wedge_k a(v_1 \wedge v_2 \dots \wedge v_l) = a(v_1) \wedge a(v_2) \wedge \dots \wedge a(v_l)$$

for each  $v_1, \dots, v_l$  in  $E$ . The map  $\wedge_l a$  is called the  $l$ -adjoint of  $a$ . In coordinates, this map is represented by a matrix whose coefficients are the  $l$ -minors of  $a$ .

If  $\omega$  is a  $k$ -form on  $E$  and  $U$  is an  $l$ -vector,  $l \leq k$ , we denote by  $i_U \omega$  the  $(k-l)$ -form defined by

$$i_U \omega \cdot v = \omega \cdot (U \wedge v)$$

for any  $(k-l)$ -vector  $v$ .

**Lemma 1.** *Let  $\Omega$  be a volume form on  $E$ , let  $\lambda$  be the unique  $n$ -vector on  $E$  such that  $\Omega \cdot \lambda = 1$ , let  $U$  be a  $k$ -vector on  $E$  and let  $a \in \mathcal{L}(E, F)$ . Then, for any  $k$ -form  $\chi$  on  $F$ , we have*

$$((i_U \Omega) \wedge \chi) \circ \wedge_n (Id \oplus a) \cdot \lambda = (-1)^{k(n-k)} \chi \circ \wedge_k a \cdot U,$$

where  $Id \oplus a : E \rightarrow E \oplus F$  is the map  $v \mapsto (v, a(v))$ .

PROOF. We make a proof in coordinates. Let  $(e_1, \dots, e_n)$  be a base of  $E$  such that  $\lambda = e_1 \wedge e_2 \wedge \dots \wedge e_n$ . If  $e_j^*$  is the dual base of  $E^*$ , then we have  $\Omega = e_1^* \wedge \dots \wedge e_n^*$ . If  $I$  is a subset of  $\{1, \dots, n\}$ , we denote by  $e_I$  the product  $e_{\alpha_1} \wedge \dots \wedge e_{\alpha_i}$ , where  $i$  is the cardinal of  $I$ , and  $\alpha_j, 1 \leq j \leq i$  are the elements of  $I$  in increasing order. We denote by  $\sigma(I)$  the sign such that  $e_I \wedge e_{I^c} = \sigma(I) \lambda$ , where  $I^c$  is the complement of  $I$ . Note then that  $\Omega = \sigma(I) e_I^* \wedge e_{I^c}^*$ , so that  $i_{e_I} \Omega = \sigma(I) e_{I^c}^*$ . Note that  $\sigma(I^c) = (-1)^{k(n-k)} \sigma(I)$ . It is sufficient to prove the Lemma for  $U = \sigma(J^c) e_{J^c}$ , where  $J$  has cardinal  $n-k$ , in which case  $i_U \Omega = e_J^*$ . We have

$$\wedge_n (Id \oplus a) \cdot \lambda = \sum_I \sigma(I) e_I \wedge (\wedge_{|I^c|} a \cdot e_{I^c})$$

where the sum is taken on all subsets  $I$  of  $\{1, \dots, n\}$  and  $|I|$  is the cardinal of  $I$ . We get

$$\begin{aligned} ((i_U \Omega) \wedge \chi) \circ \wedge_n (Id \oplus a) \cdot \lambda &= (e_J^* \wedge \chi) \circ \wedge_n (Id \oplus a) \cdot \lambda = \\ &= \sigma(J) \chi \circ \wedge_k a \cdot e_{J^c} = (-1)^{k(n-k)} \chi \circ \wedge_k a \cdot U. \end{aligned}$$

□

## 2 Generalized maps

Let us first recall the definitions of the Sobolev space  $W^{1,q}(N, M)$ . We say that  $u$  belongs to this space if there exists a  $v(t) \in \mathcal{L}(T_t N, T_{u(t)} M)$ , depending measurably on  $t$ , such that

$$\int_N \|v(t)\|_{t,u(t)}^q dt < \infty$$

and

$$d(\chi \circ u)_t = d\chi_{u(t)} \circ v(t)$$

in the sense of distributions for all bounded smooth functions  $\chi : M \rightarrow \mathbb{R}$  with bounded derivative. This can be written intrinsically on the manifold  $N$  by requiring that, for each smooth vectorfield  $U(t)$  on  $N$  compactly supported in the interior of  $N$ , we have

$$\int_N d\chi_{u(t)} \circ v(t) \cdot U(t) dt + \int_N \chi(u(t)) \cdot \operatorname{div} U(t) dt = 0.$$

It is not hard to see that, if  $M$  is a Riemannian submanifold of a Euclidean space  $E$ , then  $W^{1,q}(N, M)$  is just the set of the functions  $u \in W^{1,q}(N, E)$  which satisfy  $u(t) \in M$  for almost every  $t$ . We recall that smooth functions are not necessarily dense in these spaces if  $q < n$ .

### 2.1 Young measures

Let us denote by  $J^1(N, M)$  the set of 1-jets of maps  $u : N \rightarrow M$ . In many examples,  $N$  is an open subset of  $\mathbb{R}^n$ ,  $M$  is  $\mathbb{R}^m$ , and then

$$J^1(N, M) = N \times \mathbb{R}^m \times \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m).$$

We shall usually denote by  $(t, x, v)$  the points of  $J^1(N, M)$ . We define the function

$$r_k(t, x, v) = 1 + \|v\| + \|\wedge_2 v\| + \cdots + \|\wedge_k v\|$$

and associate to it the complete metric space  $\mathcal{P}_{r_k}(J^1(N, M))$  as in the appendix. This is the space of Borel probability measures  $\eta$  on  $J^1(N, M)$  such that  $\int r_k d\eta < \infty$ . We note that the measure on  $N$  induced by the Riemann metric, which we have denoted by  $dt$ , is finite, since  $N$  is compact; to simplify the following definitions, we shall suppose that the measure of  $N$  is 1.

Let  $t : J^1(N, M) \rightarrow N$  denote the natural projection; we denote by  $\mathcal{Y}_k(N, M)$  the set of non-negative Borel measures  $\eta \in \mathcal{P}_{r_k}(J^1(N, M))$  such that  $t_{\#}\eta$  coincides with the measure  $dt$ .

**Proposition 2.** *Assume that  $L$  is a normal integrand which is bounded from below (or more generally such that  $L/r_k$  is bounded from below), then  $\eta \mapsto \int L d\eta$  is lower semi-continuous on  $\mathcal{Y}_k$ .*

PROOF. Assume first that  $L$  is continuous and that  $L/r_k$  is bounded. Then, the functional is continuous by definition of the topology on  $\mathcal{P}_{r_k}$ .

As an intermediate step, assume that  $L(t, x, v)$  is a Caratheodory integrand (measurable in  $t$  and continuous in  $(x, v)$ ) and that  $L/r_k$  is bounded. By the Scorza-Dragnoni Theorem, (see [4], Theorem I.1.1, p 132) there exists an increasing sequence  $K_i$  of compact subsets on  $N$  such that  $L$  is continuous on  $J^1(N, M)|_{K_i}$  (the set of points  $(t, x, v)$  such that  $t \in K_i$ ) and such that  $\cup_i(K_i)$  has full measure in  $N$ . Then, there exists a sequence of continuous functions  $L_i$  such that  $|L_i|/r_k$  is bounded, independently of  $i$ , and such that  $L_i = L$  on  $J^1(N, M)|_{K_i}$ . It follows that the map  $\eta \mapsto \int L d\eta$  is the uniform limit on  $\mathcal{Y}_k(N, M)$  of the continuous maps  $\eta \mapsto \int L_i d\eta$ , and therefore it is continuous on  $\mathcal{Y}_k(N, M)$ .

In the general case, we first write the integrand  $L(t, x, v) = r_k(t, x, v)g(t, x, v)$  with a normal integrand  $g$  which is bounded from below. Then  $g$  is the increasing pointwise limit of a sequence  $g_i$  of bounded Caratheodory integrands, see [4], Theorem I.1.2, p 138. Finally, the map  $\eta \longmapsto \int Ld\eta$  is the increasing limit of the continuous maps  $\eta \longmapsto \int r_k g_i d\eta$ , and therefore it is lower semi-continuous.

□

## 2.2 Closed measures

It is a fundamental and well known observation that there exists many null-Lagrangians, that is functions  $F(t, x, v) : J^1(N, M) \longrightarrow \mathbb{R}$  such that

$$\int_N F(t, u(t), du_t)dt = 0$$

for all  $C^1$  maps  $u : N \longrightarrow M$ . We define  $\mathcal{N}_k(N, M)$  as the sets of continuous functions  $F(t, x, v)$  such that

- $F/r_k$  is bounded.
- $\int_N F(t, u(t), du_t)dt = 0$  for each  $C^1$  map  $u$ .
- There exists a compact  $K \subset N$  such that  $F(t, x, v) = 0$  if  $t \notin K$ .
- We have  $F(t, x, v) = F_{t,x}(v, \wedge_2 v, \dots, \wedge_k v) = F(t, x, v, \wedge_2 v, \dots, \wedge_k v)$ , where  $F$  is continuous and where the functions  $F_{t,x}(v, v_2, \dots, v_k)$  are affine (We say that  $F_{t,x}$  is  $k$ -affine).

By extension, we shall also denote by  $\mathcal{N}_k(N, M)$  the set of functions  $F(t, x, v_1, \dots, v_k)$  associated to the elements  $F \in \mathcal{N}_k(N, M)$ . Note that the set  $\mathcal{N}_k(N, M)$  may depend on the metric on  $M$  if  $M$  is not compact.

**Definition 3.** A Young measure  $\eta \in \mathcal{Y}_k$  is called closed if  $\int Fd\eta = 0$  for all  $F \in \mathcal{N}_k$ . The set  $\mathcal{C}_k$  of closed measures is closed in  $\mathcal{Y}_k$ , and contains the Young measures  $\hat{u}$  associated to maps  $u \in W^{1,n}(N, M)$ .

Let us explain how to build null-Lagrangians. Given a field of  $l$ -vectors  $U$ , we will denote by  $\dot{U}(t)$  the field of  $(l-1)$ -vectors which satisfies

$$d(i_U \Omega) = (-1)^{l+1} i_{\dot{U}} \Omega$$

for any volume form  $\Omega$  on  $N$  which is compatible with the Riemannian metric (meaning that the volume of an orthonormal base is  $\pm 1$ ). Notice that there are exactly two such volume forms on  $N$  if it is orientable, and that they lead to the same  $\dot{U}$ . If  $N$  is not orientable, then no global volume form  $\Omega$  exists, but we can still define  $\dot{U}$  by using volume forms defined on orientable open subsets of  $N$  (for example discs). If  $l = 1$ , for example,  $U$  is a vector-field, and  $\dot{U} = \text{div}U$ .

**Lemma 4.** For each  $l \in 1, \dots, k$ , each smooth  $(l-1)$ -form  $\chi$  on  $M$  such that both  $\chi$  and  $d\chi$  are bounded, and each compactly supported smooth field  $U(t)$  of  $l$ -vectors on  $N$ , the function

$$F(t, x, v) := \chi_x \circ \wedge_{l-1} v \cdot \dot{U}(t) + d\chi_x \circ \wedge_l v \cdot U(t)$$

belongs to  $\mathcal{N}_k(N, M)$ . In the case  $l = 1$ , the form  $\chi$  is just a function  $\chi(x)$  on  $M$ , and the function  $F$  can be rewritten more clearly

$$F(t, x, v) = \chi(x) \text{div}U(t) + d\chi_x \circ v \cdot U(t).$$

PROOF. Let  $u : N \rightarrow \mathbb{R}^m$  be a  $C^1$  function. Let us still denote by  $i_U\Omega$  and  $\chi$  the pull-backs of  $i_U\Omega$  and  $\chi$  by the projections  $N \times \mathbb{R}^m \rightarrow N$  and  $N \times \mathbb{R}^m \rightarrow \mathbb{R}^m$  respectively. This allows us to define on  $N \times \mathbb{R}^m$  the  $(n-1)$ -form  $\xi = i_U\Omega \wedge \chi$ . We have

$$\begin{aligned} 0 &= \int_N (Id \times u)^* d\xi = \int_N d\xi_{(t,u(t))} \circ \wedge_n (Id \times du_t) \cdot \lambda dt \\ &= (-1)^{l+1} \int_N (i_{\dot{U}}\Omega \wedge \chi) \circ \wedge_n (Id \times du_t) \cdot \lambda dt + (-1)^{n-l} \int_N (i_U\Omega \wedge d\chi) \circ \wedge_n (Id \times du_t) \cdot \lambda dt \end{aligned}$$

Using Lemma 1 of section 1.1, we obtain

$$0 = (-1)^{(l-1)(n-l)} \int_N \chi_{u(t)} \circ \wedge_{l-1} du_t \cdot \dot{U}(t) dt + (-1)^{(l+1)(n-l)} \int_N d\chi_{u(t)} \circ \wedge_l du_t \cdot U(t) dt.$$

After simplifying the signs, we obtain

$$\int_N \chi_{u(t)} \circ \wedge_{l-1} du_t \cdot \dot{U}(t) dt + \int_N d\chi_{u(t)} \circ \wedge_l du_t \cdot U(t) dt = 0.$$

This is the required equality.  $\square$

### 2.3 Generalized maps and Cartesian maps

The closed measure  $\eta \in \mathcal{C}_k$  is called a generalized map if there exists a measurable map  $u : N \rightarrow M$  such that the marginal of  $\eta$  on  $N \times M$  is concentrated on the graph of  $u$ . We then say that  $\eta$  is a generalized map over  $u$ .

**Definition 5.** We denote by  $\text{cart}_k(N, M)$  the set of measurable maps  $u$  such that there exists a generalized map over  $u$ . We call these maps cartesian maps. We have a natural projection  $\pi$  from the set  $\mathcal{G}_k$  of generalized maps to the set  $\text{cart}_k(N, M)$  of cartesian maps.

The generalized maps have a remarkable structure:

**Theorem 2.** Let  $\eta$  be a generalized map over  $u$ . Then, there exists a measurable family  $\Gamma_t$  of probability measures on  $\mathcal{L}(T_t N, T_{u(t)} M)$  such that  $\eta = dt \otimes \delta_{u(t)} \otimes \Gamma_t$ . Setting

$$g_i(t) := \int_{\mathcal{L}(T_t N, T_{u(t)} M)} \wedge_j v \, d\Gamma_t(v),$$

we have  $u \in W^{1,1}(N, M)$ ,  $g_1(t) = du_t$  and  $g_i(t) = \wedge_i g_1(t)$  for almost all  $t$ .

By Jensen's inequality, we immediately obtain:

**Corollary 6.** If  $\eta$  is a generalized map above  $u$ , and if  $L$  is  $k$ -convex, then

$$\int_{J^1(N, M)} L d\eta \geq \int_N L(t, u(t), du_t) dt.$$

PROOF.

$$\int_{J^1(N, M)} L d\eta = \int_N \int_{\mathcal{L}(T_t N, T_{u(t)} M)} L(t, u(t), v) d\Gamma_t(v) dt$$

But we have, for each  $t$ ,

$$\int_{\mathcal{L}(T_t N, T_{u(t)} M)} L(t, u(t), v) d\Gamma_t(v) = \int_{\mathcal{L}(T_t N, T_{u(t)} M)} \mathbf{L}(t, u(t), v, \wedge_2 v, \dots, \wedge_k v) d\Gamma_t(v)$$



$$\geq L(t, u(t), du(t), \wedge_2 du(t), \dots, \wedge_k du(t))$$

by Jensen's inequality, because  $\int_{\mathcal{L}(T_t N, T_{u(t)} M)} \wedge_j v d\Gamma_t(v) = \wedge_j du(t)$  by theorem 2.  $\square$

The proof of Theorem 2 will occupy the end of the present section. The functions  $g_i(t)$  depend only on the map  $u$ , not on  $\eta$ . This is a consequence of the following:

**Lemma 7.** *Let  $u : N \rightarrow M$  be a given measurable function. Then there exists at most one family of functions  $g_1(t), \dots, g_k(t)$  such that*

$$\int_N F(t, u(t), g_1(t), g_2(t), \dots, g_k(t)) dt = 0$$

for each  $F \in \mathcal{N}_k$ . We call these functions the *distributional minors* of  $u$  if they exist. The map  $u$  belongs to  $\text{cart}_k(N, M)$  if and only if it admits distributional minors.

PROOF. The maps  $g_l$  satisfy the following equations:

$$\int_N d\chi_{u(t)} \circ g_1(t) \cdot U(t) dt + \int_N \chi(u(t)) \cdot \dot{U}(t) dt = 0. \quad (\text{E1})$$

for all smooth field of  $n$ -vectors on  $N$  supported in the interior of  $N$ , and all smooth function  $\chi : M \rightarrow \mathbb{R}$ , and

$$\int_N d\chi_{u(t)} \circ g_l(t) \cdot U(t) dt + \int_N \chi(u(t)) \circ g_{l-1}(t) \cdot \dot{U}(t) dt = 0 \quad (\text{El})$$

for all  $l \in 2, \dots, k$ , for all compactly supported smooth field of  $l$ -vectors  $U(t)$  on  $N$ , and for all smooth  $l-1$ -form  $\chi$  on  $M$  which is bounded as well as  $d\chi$ . Now assume that  $g'_l(t)$  are other maps satisfying the same equation. Then, we have

$$\int_N d\chi_{u(t)} \circ (g_l(t) - g'_l(t)) \cdot U(t) = 0$$

for each  $l$ , each  $\chi$  and each  $U$ . We claim that this implies the  $g_l(t) - g'_l(t) = 0$  almost everywhere. Since we have the freedom of choosing  $U$ , we conclude easily that  $d\chi_{u(t)} \circ (g_l(t) - g'_l(t)) = 0$  for almost all  $t$ . If the claim did not hold, we could find a compact set  $K \subset N$  of positive measure, such that  $u$  and  $g_l - g'_l$  are continuous on  $K$  and  $g_l - g'_l$  does not vanish on  $K$ . Let  $t_0$  be a point of density of  $K$ , and let  $\chi$  be a compactly supported  $(l-1)$ -form on  $M$  such that

$$d\chi_{u(t_0)} \circ (g_l(t_0) - g'_l(t_0)) \neq 0.$$

Since  $t_0$  is a density point of  $K$ , and since all the involved functions are continuous on  $K$ , there exists a compact subset  $K'$  of  $K$  of positive measure such that the relation  $d\chi_{u(t)} \circ (g_l(t) - g'_l(t)) \neq 0$  holds for all  $t \in K'$ . This is a contradiction.  $\square$

**Lemma 8.** *If  $u \in \text{cart}_k(N, M)$ , then  $u \in W^{1,1}(N, M)$  and the first distributional minor  $g_1(t)$  of  $u(t)$  is the weak differential of  $u$ .*

PROOF. This is a direct consequence of (E1).  $\square$

The following remark can be applied for example when  $f$  is an embedding of  $M$  into some Euclidean space, and  $h$  is a chart of  $N$ :

**Proposition 9.** *Let  $\tilde{N}$  and  $\tilde{M}$  be other manifolds and let  $f : M \rightarrow \tilde{M}$  and  $h : \tilde{N} \rightarrow N$ . Assume that  $\tilde{M}$  is endowed with a complete metric. If  $f$  is smooth with bounded differential,  $h$  is a smooth diffeomorphism onto its image  $h(\tilde{N}) \subset N$ , and  $u \in \text{cart}_k(N, M)$ , then  $f \circ u \circ h \in \text{cart}_k(\tilde{N}, \tilde{M})$ . Moreover, the distributional minors  $\tilde{g}_i$  of  $f \circ u \circ h$  are :*

$$\tilde{g}_i(\tilde{t}) = \wedge_i df_{u(h(\tilde{t}))} \circ g_i(h(\tilde{t})) \circ \wedge_i dh_{\tilde{t}}$$

where  $g_i$  are the distributional minors of  $u$ .

PROOF. Let us endow  $\tilde{N}$  with the metric such that  $h$  is an isometry. Let  $\tilde{F}(t, x, v)$  be an element of  $\mathcal{N}_k(\tilde{N}, \tilde{M})$ . We want to prove that

$$\int_{\tilde{N}} \tilde{F}(\tilde{t}, f \circ u \circ h(\tilde{t}), df_{u(h(\tilde{t}))} \circ du_{h(\tilde{t})} \circ dh_{\tilde{t}}) d\tilde{t} = 0. \quad (1)$$

Setting

$$F(t, x, v) := \tilde{F}(h^{-1}(t), f(x), df_x \circ v \circ dh_t)$$

when  $t \in h(\tilde{N})$  and  $F(t, x, v) = 0$  when  $t \notin h(\tilde{N})$ , we observe that (1) is equivalent to

$$\int_N F(t, u(t), du_t) dt = 0. \quad (2)$$

This relation, on the other hand, holds if  $F \in \mathcal{N}_k(N, M)$  by definition of  $\text{cart}_k(N, M)$ .

We prove that  $F \in \mathcal{N}_k(N, M)$ . We begin to note that there is  $\tilde{K} \subset \tilde{N}$ ,  $\tilde{K}$  compact, such that  $\tilde{F}(\tilde{t}, \tilde{x}, \tilde{v}) = 0$  if  $\tilde{t} \notin \tilde{K}$ ; thus,  $F(t, x, v) = 0$  if  $t$  does not belong to the compact set  $h(\tilde{K})$ . Moreover, (2) holds for all  $C^1$  maps  $u$ . This is true because (2) is equivalent to (1), and (1) for  $C^1$  maps follows because  $\tilde{F}$  is a null Lagrangian, and thus it sends the  $C^1$  map  $f \circ u \circ h$  into zero; this amounts to (1) by the chain rule.

In order to prove the equality between distributional minors, we expand (1) to

$$\int_{\tilde{N}} \tilde{F}(\tilde{t}, f \circ u \circ h(\tilde{t}), \dots, \wedge_i df_{u(h(\tilde{t}))} \circ \tilde{g}_i(h(\tilde{t})) \circ \wedge_i dh_{\tilde{t}}, \dots) d\tilde{t} = 0$$

and use lemma 7. □

**Lemma 10.** *We have  $g_l(t) = \wedge_l g_1(t)$  for almost every  $t \in N$ .*

PROOF. If  $M$  is a Riemannian submanifold of the Euclidean space  $E$ , then every map in  $\text{cart}_k(N, M)$  belongs to  $\text{cart}_k(N, E)$ . Therefore, using the embedding theorem of Nash, we can assume for this proof that  $M$  is a Euclidean space. The set of points  $t_0$  which are simultaneously Lebesgue points of the function  $u$  and of all the functions  $g_l$ , have total measure. Let  $t_0$  be such a point. By taking a chart in  $N$ , we can suppose that  $N$  is the ball  $B$  of radius one in  $\mathbb{R}^n$ , that  $t_0 = 0$ , and that  $dt$  is the Lebesgue measure. Translating in  $\mathbb{R}^n$ , we can suppose that  $u(0) = 0$ . Let us consider, for  $s \geq 1$  the maps

$$u^s(t) := su(t/s), \quad g_l^s := g_l(t/s)$$

on  $N$ . By Proposition 9,  $u^s$  is a cartesian map on  $\tilde{N}$ , the ball of radius  $s$ , and  $g_l^s$  are its distributional minors. Our hypothesis on the point  $t_0$  can be rephrased by saying that, strongly in  $L^1(N)$ , we have

$$u^s(t) \rightarrow u^\infty(t) = g_1(0)t, \quad g_l^s(t) \rightarrow g_l^\infty(t) = g_l(0)$$

when  $s \rightarrow \infty$ . We can take a subsequence in order that these limits also hold almost everywhere. Let  $F$  be a null Lagrangian on the ball of radius 1; in particular, when trivially extended, it is a null Lagrangian on the ball of radius  $s$ , so that

$$\int_N F(t, u^s(t), du^s(t)) dt = 0.$$

Passing to the limit, we obtain

$$\int_N F(t, u^\infty(t), g_1^\infty(t), \dots, g_k^\infty(t)) dt = 0.$$

In other words, the limit function  $u^\infty$  has  $g_l^\infty$  as distributional minors. On the other hand, since the function  $u^\infty$  is smooth, we know that its distributional minors are  $\wedge_l du^\infty(t)$ , which here are just the constant functions  $\wedge_l g_l(0)$ . Therefore, by uniqueness of the distributional minors, we have proved that  $\wedge_l g_l(0) = g_l(0)$ .  $\square$

We have proved Theorem 2. We can reformulate it as follows: A function  $u$  belongs to  $\text{cart}_k(N, M)$  if and only if the minors  $\wedge_l du$  belong to  $L^1$  and are distributional, which means that they satisfy the equation

$$\int_N F(t, u(t), du_t, \wedge_2 du_t, \dots, \wedge_k du_t) dt = 0.$$

for all  $F \in \mathcal{N}_k$ . Note that  $\text{cart}_1(N, M) = W^{1,1}(N, M)$ .

## 2.4 Topology

The set  $\mathcal{G}_k$  of generalized maps is endowed with the topology of  $\mathcal{Y}_k$ .

**Proposition 11.** *The set  $\mathcal{G}_k(N, M)$  of generalized maps is closed in  $\mathcal{Y}_k(N, M)$ .*

PROOF. Let  $\eta^j$  be a sequence of generalized maps above  $u^j$ . Let us assume that the sequence  $\eta^j$  is converging to  $\eta$  in  $\mathcal{C}_k(N, M)$ . We have to prove that there exists a map  $u \in W^{1,1}(N, M)$  such that the marginal of  $\eta$  on  $N \times M$  is concentrated on the graph of  $u$ . It is enough to prove that, for each embedded ball  $B \subset N$ , the marginal of  $\eta|_{J^1(B, M)}$  on  $B \times M$  is concentrated on the graph of a map  $u$ . As a consequence, we can suppose that  $N$  is the open unit ball in  $\mathbb{R}^n$ . We consider  $M$  as a Riemannian submanifold of a Euclidean space  $E$ , so that we see  $u^j$  as elements of  $W^{1,1}(N, E)$  with values in  $M$ . Let  $m^j \in E$  be the average of  $u^j$ ,  $m^j = \int_N u_j(t) dt$ . Since the sequence  $\eta^j$  is  $r_k$ -tight, see appendix, the derivatives  $du^j$  are bounded in  $L^1$ . Therefore, by Poincaré inequality, the sequence  $(u^j - m^j)$  is bounded in  $W^{1,1}$ . By the compactness of the embedding  $W^{1,1} \rightarrow L^1$ , this sequence is strongly compact in  $L^1$ . We assume, taking a subsequence, that it has a limit  $u^\infty$ , and that the convergence holds almost everywhere. By Lusin and Egorov Theorems, for all  $\epsilon > 0$ , there exists a compact subset  $K \subset N$  such that  $dt(N - K) \leq \epsilon$  and such that  $u^j$  is continuous on  $K$  and  $(u^j - m^j)$  is converging uniformly on  $K$  to  $u^\infty$ . It is clear at this point that the unboundedness of  $m^j$  would contradict the tightness of  $\eta^j$ , and therefore we can assume that the averages  $m^j$  have a limit  $m^\infty$ . Setting  $u := u^\infty + m^\infty$ , we see that  $u^j$  is converging uniformly to the continuous function  $u$  on  $K$ . Denoting by  $\mu$  the marginal of  $\eta$  on  $N \times M$ , we conclude that the  $\mu$ -measure of the graph of  $u$  is greater than  $1 - \epsilon$ . Since this holds for all  $\epsilon > 0$ , we conclude that the measure  $\mu$  is concentrated on the graph of  $u$ .  $\square$

### 3 Boundary conditions and compactness

In most applications, the manifold  $M$  is not compact, and it is necessary to introduce boundary conditions in order to get compactness. We fix, as explained in the introduction, a map  $u_0 \in \text{cart}_k(N, M)$ . We define the set  $\mathcal{C}_k(N, M; u_0) \subset \mathcal{C}_k(N, M)$  of closed measures with boundary  $u_0$  as the set of measures  $\eta \in \mathcal{C}_k(N, M)$  such that

$$\int_{J^1(N, M)} d\chi_x \circ v \cdot U(t) + \chi(x) \cdot \text{div}U(t) d\eta(t, x, v) = \int_N d\chi_{u_0(t)} \circ du_0(t) \cdot U(t) + \chi(u_0(t)) \cdot \text{div}U(t) dt.$$

for each smooth vectorfield  $U(t)$  on  $\bar{N}$  (not necessarily supported in  $N$ ) and each bounded smooth function  $\chi(x)$  on  $M$  with bounded derivative. We can also define the set of generalized maps with boundary value  $u_0$ :

$$\mathcal{G}_k(N, M; u_0) := \mathcal{G}_k(N, M) \cap \mathcal{C}_k(N, M; u_0).$$

The space  $\text{cart}_k(N, M; u_0)$  is the space of maps  $u$  such that there exists a generalized map  $\eta \in \mathcal{G}_k(N, M; u_0)$  above  $u$ , or in other words the maps  $u$  whose Young measure  $\hat{u}$  belongs to  $\mathcal{G}_k(N, M; u_0)$ . In [12], the functions in  $\text{cart}_k(N, M; u_0)$  are said to satisfy a weak anchorage condition. In the case where  $M$  is a Euclidean space, the functions in  $\text{cart}_k(N, M; u_0)$  are just the functions in  $\text{cart}_k(N, M)$  which have the same trace on  $\partial N$  as  $u_0$  in the  $W^{1,1}$  sense.

**Proposition 12.** *Let  $L$  be a  $k$ -convex and coercive Lagrangian. For each  $c > 0$ , the set of measures  $\eta \in \mathcal{C}_k(N, M; u_0)$  which satisfy*

$$\int_{J^1(N, M)} L(t, x, v) d\eta(t, x, v) \leq c \tag{3}$$

*is compact.*

PROOF. Let us denote by  $\mathcal{C}(c)$  the set of measures  $\eta \in \mathcal{C}_k(N, M; u_0)$  which satisfy (3). Since the functional

$$\eta \longmapsto \int L d\eta$$

is lower semi-continuous on  $\mathcal{Y}_k(N, M)$  (by Proposition 2), and since  $\mathcal{C}_k(N, M; u_0)$ , is closed in  $\mathcal{Y}_k(N, M)$ , the set  $\mathcal{C}(c)$  is closed in  $\mathcal{Y}_k(N, M)$ . So it is enough to prove that it is relatively compact. By the appendix, this follows if we can prove that it is  $r_k$ -tight. In other words, we have to show that for each  $\epsilon > 0$  there exists a compact subset  $Z \in J^1(N, M)$  such that

$$\int_{J^1(N, M) \setminus Z(R)} r_k(t, x, v) d\eta(t, x, v) \leq 2\epsilon$$

for each measure  $\eta \in \mathcal{C}(c)$ . We shall prove that this holds for

$$Z(R) = \{(t, x, v) \in J^1(N, M) \quad : \quad d(x_0, x) \leq R, \|v\| \leq R\}$$

when  $R$  is large enough ( $x_0$  is a point in  $M$  that we have fixed once and for all). At this point it is convenient to assume, without loss of generality, that  $L \geq 0$ . We define

$$\tilde{Z}(R) = \{(t, x, v) \in J^1(N, M) \quad : \quad \|v\| \leq R\}$$

and we see that there exists  $A(R) > 0$  with  $A(R) \rightarrow +\infty$  as  $R \rightarrow +\infty$  such that, for all  $\eta \in \mathcal{C}(c)$ ,

$$c \geq \int_{J^1(N, M) \setminus \tilde{Z}(R)} L(t, x, v) d\eta(t, x, v) \geq A(R) \int_{J^1(N, M) \setminus \tilde{Z}(R)} r_k(t, x, v) d\eta(t, x, v).$$

Taking  $R$  sufficiently large, we get from the inequality above that

$$\int_{J^1(N,M) \setminus \tilde{Z}(R)} r_k(t, x, v) d\eta(t, x, v) \leq \epsilon \quad (4)$$

for each  $\eta \in \mathcal{C}(c)$ . Setting now

$$\hat{Z}(R) = \{(t, x, v) \in J^1(N, M) \quad : \quad d(x_0, x) \leq R\}$$

we see that the desired inequality follows if we prove that

$$\eta(J^1(N, M) - \hat{Z}(R)) \leq \epsilon(R) \quad \forall \eta \in \mathcal{C}(c) \quad (5)$$

for all  $R$ , with  $\epsilon(R) \rightarrow 0$  as  $R \rightarrow \infty$ . Indeed, taking  $R_0$  such that (4) holds, and then setting  $S = \max_{\tilde{Z}(R_0)} r_k$ , we get

$$\int_{J^1(N,M) \setminus Z(R)} r_k d\eta \leq \int_{J^1(N,M) \setminus \tilde{Z}(R_0)} r_k d\eta + S\eta(J^1(N, M) \setminus \hat{Z}(R)) \leq \epsilon + S\epsilon(R).$$

In order to prove (5), we consider, for each  $R > 0$ , a function  $g \in C^1(M, \mathbb{R})$  such that

$$\begin{aligned} 0 \leq g_R(x) \leq 1, \quad g(x) = 1 \quad \text{if} \quad d(x_0, x) \geq R, \\ g(x) = 0 \quad \text{if} \quad d(x_0, x) \leq R/2 \\ |dg_x| \leq \delta(R) \quad \forall x \end{aligned}$$

where  $\delta(R) \rightarrow 0$  as  $R \rightarrow \infty$ ; and a smooth vector-field  $U(t)$  on  $\bar{N}$  such that  $\dot{U} = 1$  on  $N$  or equivalently such that  $\text{div} U = 1$  on  $N$ . The existence of such a vector-field is given by Lemma 13 below. We note that  $U$  is bounded, since  $\bar{N}$  is compact. Since  $\mathcal{C}(c) \subset \mathcal{C}_k(N, M; u_0)$ , we have, for  $\eta \in \mathcal{C}(c)$ ,

$$\int_{J^1(N,M)} g(x) d\eta(t, x, v) = \int_N g(u_0(t)) + dg_{u_0(t)} \circ du_0(t) \cdot U(t) dt - \int_{J^1(N,M)} dg_x \circ v \cdot U(t) d\eta(t, x, v).$$

The last formula and the definition of  $g$  imply that there exists  $C > 0$  such that

$$\int_{J^1(N,M) \setminus Z(R)} d\eta(t, x, v) \leq C\delta(R) + \int_N g(u_0(t)) dt$$

for all  $R$  and all  $\eta \in \mathcal{C}(c)$ . The term on the right converges to zero as  $R \rightarrow \infty$ , this ends the proof.  $\square$

**Lemma 13.** *Let  $\bar{N}$  be a compact Riemannian manifold with a non-empty boundary. There exists a smooth vector-field  $U(t)$  on  $\bar{N}$  such that  $\dot{U} = 1$  on  $N$  or equivalently such that  $\text{div} U = 1$  on  $N$ .*

PROOF. In the case where  $N$  is a ball in  $\mathbb{R}^n$ , this is obvious, just take  $U(t) = t/n$ . In general, one can build  $U$  as the gradient of a function  $h$  which solves  $\Delta h = 1$  on  $N$ .  $\square$

## 4 Conclusion

We now collect the tools we have introduced to prove Theorem 1. We also add some discussions and variations.

## 4.1 Proof of Theorem 1

By Propositions 11 and 12, there exists a generalized map  $\eta$  over some  $u \in W^1(N, M)$  such that  $\eta$  minimizes  $\int L d\eta$  on  $\mathcal{G}_k(N, M; u_0)$ . We want to show that  $u$  minimizes in  $\text{cart}_k(N, M; u_0)$ .

If  $v \in \text{cart}_k(N, M; u_0)$  is another map, we have

$$\int_{J^1(N, M)} L(t, u(t), du_t) dt \leq \int L d\eta \leq \int L d\hat{v} = \int_{J^1(N, M)} L(t, v(t), dv_t) dt$$

where the first inequality comes from Corollary 6, and where  $\hat{v} \in \mathcal{G}_k(N, M; u_0)$  is the Young measure associated to  $v$ . This proves that  $u$  is minimizing in  $\text{cart}_k(N, M; u_0)$ .

## 4.2 An example

We consider  $M = \mathbb{R}^2$ ;  $N = B$ , the unit open ball of  $\mathbb{R}^2$ , and the Lagrangian

$$L(t, x, v) = \epsilon(|v|^p + |t|^4|v|^4) + |\det v|^2,$$

with  $p \in ]1, 2[$ . We claim that

$$\inf_{u \in W^{1,1}(N, M; Id)} \int_N L(t, u(t), du_t) dt < \min_{u \in \text{cart}_2(N, M; Id)} \int_N L(t, u(t), du_t) dt$$

when  $\epsilon > 0$  is small enough. Indeed, taking  $\underline{u}(t) = t/|t|$ , and observing that  $\det d\underline{u} = 0$ , we get a constant  $C > 0$  such that

$$\inf_{u \in W^{1,1}(N, M; Id)} \int_N L(t, u(t), du_t) dt \leq \int_N L(t, \underline{u}(t), d\underline{u}_t) dt \leq C\epsilon.$$

On the other hand, if  $u \in \text{cart}_2(B, \mathbb{R}^2; Id)$  is a minimizer we have

$$\int_B L(t, u(t), du_t) dt \geq \int_B |\det du_t|^2 dt \geq \frac{1}{|B|} \left( \int_B |\det du_t| dt \right)^2 \geq \pi,$$

where the last inequality follows from the following Lemma:

**Lemma 14.** *If  $u \in \text{cart}_2(B, \mathbb{R}^2; Id)$  minimizes  $\int_B L(t, u(t), du_t) dt < \infty$ , then  $\int_B \det du(t) dt = \pi$ .*

PROOF. We claim that  $u(B) \subset \bar{B}$ . Indeed, let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a smooth diffeomorphism such that  $f = Id$  on  $B$  and  $|df| < 1$  outside of  $\bar{B}$ . By Proposition 9, the map  $f \circ u$  belongs to  $\text{cart}_2(B, \mathbb{R}^2)$ , and it has the same boundary condition as  $u$ . Since  $|df(u)| \leq 1$  we have that  $|d(f \circ u)| \leq |du|$  and  $|\det d(f \circ u)| \leq |\det du|$ ; the first inequality is strict if  $|u| > 1$  and  $du \neq 0$ . If we did not have  $u(t) \subset \bar{B}$  for almost every  $t$ , the action of  $f \circ u$  would be strictly smaller than the action of  $u$ , which would contradict the assumption that  $u$  is a minimizer.

Let us denote by  $A$  the annulus  $1/2 < |t| < 1$ . We have

$$u \in W^{1,p}(B, \mathbb{R}^2) \cap W^{1,4}(A, \mathbb{R}^2),$$

so that  $u$  is continuous on  $A$ , and extends by continuity to  $\partial B$ , where it takes the value  $u|_{\partial B} = Id$ . Finally, recall that  $u(B) \subset \bar{B}$ . Define

$$u_i(t) := i^2 \int_B \tau(is) u((1 - 1/i)t - s) ds$$

where  $\tau : B \rightarrow [0, 1]$  is a smooth convolution kernel. It is classical that  $u_i \rightarrow u$  in  $W^{1,p}(B, \mathbb{R}^2)$ , and in  $W^{1,4}(A, \mathbb{R})$ . As a consequence,  $u_i|_{\partial B}$  converges uniformly to the identity; since  $u_i$  is smooth, this implies

$$\int_B \det du_i dt \rightarrow \pi.$$

Thus it is enough to prove that

$$\int_B \det du_i dt \rightarrow \int_B \det du dt. \quad (6)$$

Let  $r \in ]1/2, 1[$  and let  $\phi_r \in C_0^\infty(B, \mathbb{R})$  be such that  $0 \leq \phi_r \leq 1$  and  $\phi_r = 1$  on  $B(0, r)$ . Since

$$\int_B \det du \cdot \phi_r dt \rightarrow \int_B \det du dt$$

as  $r \rightarrow 1$ , the formula (6) follows if we prove

$$\int_B \det du_i \cdot \phi_r dt \rightarrow \int_B \det du \cdot \phi_r dt \quad \forall r \in ]1/2, 1[ \quad (7)$$

and

$$\left| \int_B \det du_i \cdot (1 - \phi_r) dt \right| < \epsilon \quad \forall i \quad \text{if } r \geq 1 - \delta. \quad (8)$$

Note that (8) follows from the boundedness of  $\det du_i$  in  $L^2(A)$ . In order to prove (7), we set  $u_i = (u_i^1, u_i^2)$  and  $u = (u^1, u^2)$ , we call  $(x_1, x_2)$  the coordinates on the target space  $\mathbb{R}^2$  and we assert that

$$\int_B \det du \cdot \phi_r dt = \int_B u^1(\partial_2 u^2, -\partial_1 u^2) \cdot \nabla \phi_r dt.$$

Indeed, this formula is just (El) with  $l = 2$ ,  $\chi = a(x^1)dx^2$  and  $U = \phi_r e_1 \wedge e_2$ , where  $a : \mathbb{R} \rightarrow [-3, 3]$  is a smooth function such that  $a(x^1) = x^1$  on  $[-2, 2]$ . Here we use that  $u(B) \subset \bar{B}$ . Similarly, by Lemma 4,

$$\int_B \det du_i \cdot \phi_r dt = \int_B u_i^1(\partial_2 u_i^2, -\partial_1 u_i^2) \cdot \nabla \phi_r dt.$$

As a consequence (7) is equivalent to

$$\int_B [u_i^1(\partial_2 u_i^2, -\partial_1 u_i^2)] \cdot \nabla \phi_r dt \rightarrow \int_B [u^1(\partial_2 u^2, -\partial_1 u^2)] \cdot \nabla \phi_r dt$$

which holds because the integrand is converging almost everywhere and is bounded in  $L^2$ .  $\square$

### 4.3 Weak continuity of minors

Let us mention the following classical result which follows from our tools (see [12], 3.3.1 or [10], 8.3):

**Proposition 15.** *Let  $N$  be a bounded disc in  $\mathbb{R}^n$ . Let  $u_i$  be a sequence of maps in  $\text{cart}_k(N, \mathbb{R}^m)$ , and let  $u \in W^{1,1}(N, M)$  and  $g_j(t) \in L^1(N, \mathcal{L}(\wedge_j \mathbb{R}^n, \wedge_j \mathbb{R}^m))$  be such that  $u_i \rightarrow u$  weakly in  $W^{1,1}$  and*

$$\wedge_2 du_i \rightarrow g_2, \quad \dots \quad \wedge_k du_i \rightarrow g_k$$

*weakly in  $L^1$ . Then  $g_2 = \wedge_2 du, \dots, g_k = \wedge_k du$ .*

PROOF. We consider the Young measures  $\hat{u}_i$  in  $\mathcal{G}_k(N, M)$  associated to the functions  $u_i$ . Now weak convergence implies uniform integrability, which translates to the fact that  $\hat{u}_i$  is  $r_k$ -tight, and therefore compact in  $\mathcal{Y}_k(N, \mathbb{R}^m)$ . We can suppose that it has a limit  $\eta$ , which is a generalized map above  $u$ . If  $F(t, v) = \mathbb{F}(t, v, v_2, \dots, v_k)$  is a continuous function which is affine in  $(v, v_2, \dots, v_k)$ , then we have

$$\int F d\hat{u}_i \longrightarrow \int F d\eta$$

on one hand, because  $\hat{u}_i \longrightarrow \eta$ , and

$$\int \mathbb{F}(t, du_i(t), \wedge_2 du_i(t), \dots, \wedge_k du_i(t)) dt \longrightarrow \int \mathbb{F}(t, du(t), g_2(t), \dots, g_k(t)) dt$$

on the other hand, because  $\wedge_j du_i \longrightarrow g_j$  weakly; we recall that  $F$  is affine in the minors. We conclude that

$$\int F d\eta = \int \mathbb{F}(t, du(t), g_2(t), \dots, g_k(t)) dt.$$

This implies that, for almost all  $t$ ,

$$g_j(t) = \int \wedge_j v d\Gamma_t = \wedge_j du(t)$$

by Theorem 2. □

#### 4.4 On Null-Lagrangians

It may seem unnatural in the definitions of the sets  $\mathcal{N}_k(N, M)$  to require that the null-Lagrangians  $F(t, x, v)$  be  $k$ -affine functions of  $v$ . Indeed, working with a larger set  $\mathcal{N}(N, M)$  of null-Lagrangians would make the result stronger, and may allow to relax somewhat the  $k$ -convexity hypothesis on  $L$ . The following result, however, shows that there is not much hope in that direction:

**Proposition 16.** *Let us assume that  $N = \mathbb{R}^n$ , and that  $M = \mathbb{R}^m$ . If  $F$  is a null-Lagrangian such that  $F/r_k$  is bounded, then  $F_{t,x}$  is  $k$ -affine for each  $t$  and  $x$ .*

PROOF. We just give an idea of the proof. It follows from proposition 9 that, for  $\lambda > 0$  and  $(t_0, x_0) \in \mathbb{R}^n \times \mathbb{R}^m$ , the function

$$F_\lambda(t, x, v) := F(t_0 + \lambda t, x_0 + \lambda x, v)$$

is a null-Lagrangian. But then  $F_0$  is also a null-Lagrangian, which means that  $F_{t_0, x_0}$  is quasi-affine in the sense of [10], section 4.1; but in the same section of [10] it is proven that quasi-affine functions are poly-affine. In other words, there exists an affine function  $\mathbb{F}(v, v_2, \dots, v_p)$  such that  $F_{t_0, x_0}(v) = \mathbb{F}(v, \wedge_2 v, \dots, \wedge_p v)$ , where  $p = \min\{m, n\}$ . But the bound implies that  $\mathbb{F}$  does not depend on  $\wedge_j v$  for  $j > k$ . □

#### 4.5 More general setting

The heart of the matters is the Jensen's inequality obtained in Corollary 6. This inequality is the result of an equilibrium between the known properties of the measures  $\Gamma_t$  appearing in the disintegration of generalized maps and the convexity assumed on the integrand  $L$ .



Other, but less explicit equilibria might be obtained as follows. Let  $r(t, x, v)$  be a continuous function on  $J^1(N, M)$  such that  $r(t, x, v) \geq 1 + \|v\|$ . We define the associated Kantorovich-Rubinstein space  $\mathcal{P}_r(J^1(N, M))$ , which is the set of Borel probabilities  $\eta$  on  $J^1(N, M)$  such that  $\int r d\eta < \infty$ . We also define the set  $\mathcal{Y}_r(N, M)$  of those elements  $\eta$  of  $\mathcal{P}_r(J^1(N, M))$  such that  $t_{\#}\eta = dt$ .

Now let  $\hat{\mathcal{G}}_r(N, M)$  be the closure, in  $\mathcal{Y}_r(N, M)$  of the set of Young measures associated to smooth maps. We can prove as in Proposition 11 that, to each  $\eta \in \hat{\mathcal{G}}_r(N, M)$  is associated a map  $u \in W^{1,1}(N, M)$  such that

$$\eta = dt \otimes \delta_{u(t)} \otimes \Gamma_t \quad (9)$$

and such that  $\int v d\Gamma_t = du(t)$  for almost all  $t$ . We can define  $\text{cart}_r(N, M)$  as the set of maps which appear in this way. In this setting, we can fix boundary conditions as before by taking  $u_0 \in \text{cart}_r(N, M)$ . If the coercivity condition of the Lagrangian is modified to

$$L(t, x, v) \geq l(r(t, x, v)),$$

with  $l$  super-linear, we still have compactness: Proposition 12 still holds, with the same proof. So if  $L$  is a normal integrand satisfying the modified coercivity condition, then there exists a Young measure  $\eta \in \hat{\mathcal{G}}_r(N, M; u_0)$  which minimizes the integral  $\int L d\eta$  in this set.

In order to prove the existence of minimizers in  $\text{cart}_r(\tilde{N}, M; u_0)$ , it is enough to adapt the convexity condition, in such a way that Corollary 6 holds for the elements of  $\hat{\mathcal{G}}_r(N, M)$ .

Let  $\mathcal{P}_{t,x}$  be the set of Borel probability measures  $\Gamma$  on  $\mathcal{L}(T_t N, T_x M)$  such that  $\int r_{t,x}(v) d\Gamma(v) < \infty$ . In short, we have

$$\mathcal{P}_{t,x} := \mathcal{P}_{r_{t,x}}(\mathcal{L}(T_t N, T_x M))$$

(see the Appendix below). Let  $B$  be a closed ball of volume one in  $T_t N$ . Let  $\mathbf{P}_{t,x}$  be the closure, in  $\mathcal{P}_{t,x}$  of the measures of the form

$$\Gamma = (du)_{\#}(dt|_B)$$

where  $u : T_t N \rightarrow T_x M$  is a smooth map supported in  $B$ . Note that if  $\Gamma \in \mathbf{P}_{t,x}$ , then  $\int_{\mathcal{L}(T_t N, T_x M)} v d\Gamma = 0$ . A last notation is necessary: we denote by  $\tau_z$  the translation of vector  $z$ . Then, possibly under some mild assumption on the function  $r$ , the following result can be proved by a blow-up argument called localisation procedure in [14]:

**Structure Theorem:** *The measures  $\eta \in \hat{\mathcal{G}}_r(N, M)$  can be written in the form (9), with*

$$(\tau_{-du_t})_{\#} \Gamma_t \in \mathbf{P}_{t,x}$$

for almost all  $t$ .

As a consequence, the convexity condition that has to be assumed in order that Corollary 6, and then Theorem 1 hold in this more general setting is

$$\int_{\mathcal{L}(T_t N, T_x M)} L_{t,x}(a + v) d\Gamma(v) \geq L(t, x, a)$$

for all  $(t, x) \in N \times M$ , for all  $a \in \mathcal{L}(T_t N, T_x M)$  and for all  $\Gamma \in \mathbf{P}_{t,x}$ . This is not an easy condition to check on examples.

## A Kantorovich-Rubinstein space

Let us recall some standard facts on probability measures, see [2, 16]. Let  $(X, d)$  be a complete and separable metric space, and let  $r : X \rightarrow [1, \infty)$  be a continuous function. Let  $\mathcal{P}_r(X)$  be the set of Borel probability measures  $\mu$  on  $X$  which satisfy

$$\int_X r(x) d\mu(x) < \infty.$$

Let us denote by  $C_r(X)$  the set of continuous functions  $f$  on  $X$  such that

$$\sup_{x \in X} \frac{|f(x)|}{r(x)} < \infty.$$

There exists a distance  $d$  on  $\mathcal{P}_r(X)$  such that  $d(\mu_n, \mu) \rightarrow 0$  if and only if

$$\int f d\mu_n \rightarrow \int f d\mu$$

for all  $f \in C_r(X)$ . This distance can be chosen such that, in addition, the metric space  $(\mathcal{P}_r, d)$  is a complete and separable metric space.

In order to define such a distance  $d$  on  $\mathcal{P}_r(X)$  one can define first the distance

$$d_r(x, y) := \min(d(x, y), 1) + |r(y) - r(x)|.$$

on  $X$ , which is complete and equivalent to  $d$ . Then, we can define the distance  $d$  on  $\mathcal{P}_r(X)$  as the Kantorovich-Rubinstein (also called 1-Wasserstein) distance of  $(X, d_r)$ .

The relatively compact subsets of  $(\mathcal{P}_r(X), d)$  are those which are  $r$ -tight:

**Definition 17.** *The subset  $Y \subset \mathcal{P}_r(X)$  is called  $r$ -tight if one of the following equivalent properties holds:*

- *For each  $\epsilon > 0$ , there exists a compact set  $K \subset X$  such that  $\int_{X-K} r(x) d\mu \leq \epsilon$  for each  $\mu \in Y$ .*
- *There exists a function  $f : X \rightarrow [0, \infty]$  whose sublevels are compact and a constant  $C$  such that  $\int_X r(x) f(x) d\mu \leq C$  for each  $\mu \in Y$ .*
- *The family  $Y$  is tight and  $r$  is  $Y$ -uniformly integrable. The first means that, for each  $\epsilon > 0$ , there exists a compact set  $K \subset X$  such that  $\mu(X - K) \leq \epsilon$  for each  $\mu \in Y$ . The second means that for each  $\epsilon > 0$ , there exists a ball  $B$  in  $X$  such that  $\int_{X-B} r(x) d\mu \leq \epsilon$  for each  $\mu \in Y$ .*

Note that 1-tightness is just tightness if  $r \equiv 1$ . If  $r$  is proper, then  $Y$  is  $r$ -tight if and only if there exists a constant  $C$  and a superlinear function  $f : [0, \infty) \rightarrow \mathbb{R}$  such that

$$\int_X f \circ r d\eta \leq C$$

for all  $\eta \in Y$ .

## References

- [1] R. A. Adams, Sobolev spaces, Academic press, New York, 1975.
- [2] L. Ambrosio, N. Gigli and G. Savaré, Gradient flows, *Lectures in Math. ETH Zürich*, Birkhäuser (2005).
- [3] J. Ball, *Convexity conditions and existence theorems in nonlinear elasticity*, Arch. Rat. Mech. Anal. **63** (1977), no. 3, 337–403.
- [4] H. Berliocchi and J.-M. Lasry, *Intégrales normales et mesures paramétrées en calcul des variations*, Bull. Soc. Math. France **101** (1973), 129–184.
- [5] J. Ball, V. J. Mizel, *One-dimensional variational problems whose minimizers do not satisfy the Euler-Lagrange equation*, Arch. Rat. Mech. Anal. **90** (1985), 325–388.

- [6] P. Bernard and B. Buffoni, *Optimal mass transportation and Mather theory*, JEMS **9** (2007), no. 1, 85–121.
- [7] P. Bernard, *Young measures, superposition and transport* Indiana Math. Journal, (2008)
- [8] Y. Bernier, *Minimal geodesics on groups of volume-preserving maps and generalized solutions of the Euler equations*, Comm. Pure. Ap. Math. **52** (1999), 411–452.
- [9] G. Buttazzo, *Semicontinuity and relaxation and integral representation in the calculus of variations*, Longman Scientific and Technical, 207 (1989).
- [10] B. Dacorogna, *Direct methods in the calculus of variations (second edition)* App. Math. Sciences 78 (2008).
- [11] L. C. Evans, R. F. Gariepy, *Measure theory and fine properties of functions*, CRC press, Boca Raton, 1992.
- [12] M. Giaquinta, G. Modica, J. Soucek, *Cartesian currents in the calculus of variations I*, Springer, Berlin, 1998.
- [13] M. Giaquinta, G. Modica, J. Soucek, *Cartesian currents in the calculus of variations II*, Springer, Berlin, 1998.
- [14] P. Pedregal, *Parametrized Measures and Variational Principles*, *Progress in nonlinear Partial Differential Equations and their Applications* Birkhauser (1997).
- [15] M. Valadier, *A course on Young measures*, Rend. Istit. Mat. Univ. Trieste **24** (1994), 349–394.
- [16] C. Villani, *Topics in optimal transportation*, American Mathematical Society, Providence, Rhode Island, 2003.
- [17] L. C. Young, *Lectures on the calculus of variations and optimal control theory*, second edition, Chelsea (1980).