



## Carbon isotope anomaly in the major plant C1 pool and its global biogeochemical implications

F. Keppler, R. M. Kalin, D. B. Harper, W. C. Mcroberts, J. T. G. Hamilton

### ► To cite this version:

F. Keppler, R. M. Kalin, D. B. Harper, W. C. Mcroberts, J. T. G. Hamilton. Carbon isotope anomaly in the major plant C1 pool and its global biogeochemical implications. *Biogeosciences Discussions*, 2004, 1 (1), pp.393-412. hal-00297756

**HAL Id: hal-00297756**

**<https://hal.science/hal-00297756>**

Submitted on 18 Jun 2008

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*Biogeosciences Discussions* is the access reviewed discussion forum of *Biogeosciences*

# Carbon isotope anomaly in the major plant C<sub>1</sub> pool and its global biogeochemical implications

F. Keppler<sup>1,2</sup>, R. M. Kalin<sup>2</sup>, D. B. Harper<sup>1</sup>, W. C. McRoberts<sup>1,3</sup>, and J. T. G. Hamilton<sup>1,3</sup>

<sup>1</sup>School of Agriculture and Food Science, Queen's University Belfast, Newforge Lane, Belfast BT9 5PX, United Kingdom

<sup>2</sup>Environmental Engineering Research Centres, Queen's University Belfast, Belfast BT9 5AG, United Kingdom

<sup>3</sup>Department of Agriculture and Rural Development for Northern Ireland, Newforge Lane, Belfast BT9 5PX, United Kingdom

Received: 26 July 2004 – Accepted: 10 August 2004 – Published: 18 August 2004

Correspondence to: F. Keppler (f.keppler@qub.ac.uk)

## Carbon isotope anomaly in the major plant C<sub>1</sub> pool

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

## Abstract

We report that the most abundant  $C_1$  units of terrestrial plants, the methoxyl groups of pectin and lignin, have a unique carbon isotope signature exceptionally depleted in  $^{13}C$ . Plant-derived  $C_1$  volatile organic compounds (VOCs) are also anomalously depleted in  $^{13}C$  compared with  $C_{n+1}$  VOCs. The results confirm that the plant methoxyl pool is the predominant source of biospheric  $C_1$  compounds of plant origin such as methanol, chloromethane and bromomethane. Furthermore this pool, comprising ca. 2.5% of carbon in plant biomass, represents an important substrate for methanogenesis and could be a significant source of isotopically light methane entering the atmosphere. Our findings have significant implications for the use of carbon isotope ratios in elucidation of global carbon cycling. Moreover methoxyl groups could act as markers for biological activity in organic matter of terrestrial and extraterrestrial origin.

## 1. Introduction

Stable isotope analysis has become a powerful tool for environmental scientists, plant biologists, ecologists and geochemists studying global elemental cycles or past climatic conditions (e.g. Ehleringer et al., 2002; Yakir, 2002; Hayes, 2001; Griffiths, 1998; Lajtha and Michener, 1994). Thus plant species have been photosynthetically characterised as Calvin cycle ( $C_3$ ), Slack-Hatch cycle ( $C_4$ ) and Crassulacean acid metabolism (CAM) categories using carbon isotope signatures (Griffiths, 1998; O'Leary, 1981). Moreover variations in the carbon isotope composition ( $\delta^{13}C$  values) of compounds, produced and destroyed in the global carbon cycle, are often used to investigate biogeochemical cycles and global source-sink relationships, as well as the underlying mechanisms (e.g. Cerling et al., 1997; Sherwood Lollar et al., 2002; Michaelis et al., 2002). Stable isotope techniques are increasingly applied to the study of atmospheric budgets of volatile organic compounds (VOCs). Many  $C_1$  VOCs, such as methanol ( $CH_3OH$ ), chloromethane ( $CH_3Cl$ ), bromomethane ( $CH_3Br$ ), iodomethane ( $CH_3I$ ), cyanomethane

**BGD**

1, 393–412, 2004

## Carbon isotope anomaly in the major plant $C_1$ pool

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

(CH<sub>3</sub>CN) and methane (CH<sub>4</sub>), play an important role in atmospheric chemistry and possibly climate change (see, for example, Heikes et al., 2002; Montzka et al., 2003; O'Dowd et al., 2002 ; Sanhueza et al., 2004; Wuebbles and Hayhoe, 2002). Numerous investigations into the atmospheric budget of such C<sub>1</sub> compounds, some employing stable isotope techniques (Goldstein, 2003; Bill et al., 2004; Harper et al., 2003; Thompson et al., 2002; Whiticar, 1999; Kalin et al., 2001) have been reported but many questions regarding their origin and fate remain unresolved.

Most CH<sub>3</sub>OH released from plants is derived from the ubiquitous plant component pectin by both enzymic and abiotic processes (Fall and Benson, 1996, Warneke et al., 1999; Galbally and Kirstine, 2002). Pectin which normally comprises between 7 and 35% of cell wall material in leaves is composed of galacturonic acid monomer units. Between 50 and 90% of the carboxyl groups of the latter are methyl esterified and provide the methyl pool for the reaction. We have also identified pectin as the source of CH<sub>3</sub>Cl and other monohalomethanes produced abiotically by senescent and dead leaf material (Hamilton et al., 2003). However, little consideration has been given to the stable isotope signature of the methoxyl pool of pectin, or indeed that of another important plant component lignin, and the impact these might have on the  $\delta^{13}\text{C}$  values of C<sub>1</sub> compounds in the biosphere.

## 2. Materials and methods

**Stable carbon isotope measurements:** Carbon isotopic ratios of VOCs were measured by compound specific isotope analysis (GC-MS-IRMS) using a Thermo Finnigan Delta+ isotope ratio mass spectrometer interfaced with a Finnigan DSQ gas chromatograph trace mass spectrometer. Bulk  $\delta^{13}\text{C}$  signatures of dried plant samples were determined using a Eurovector elemental analyzer coupled to a Micromass PRISM III isotope ratio mass spectrometer. Internal precision of  $\delta^{13}\text{C}$  was  $\pm 0.2$  (‰). Values of  $\delta^{13}\text{C}$  (‰) relative to that for the Vienna-PDB are defined by the equation  $\delta^{13}\text{C}$  (‰) =  $(R_{\text{sample}}/R_{\text{standard}} - 1) \times 1000$ ‰ with  $R = {}^{13}\text{C}/{}^{12}\text{C}$ . The isotope difference ( $\Delta$ ) between

### Carbon isotope anomaly in the major plant C<sub>1</sub> pool

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

two pools is defined as  $\Delta = \delta^{13}\text{C}_{\text{pool 1}} - \delta^{13}\text{C}_{\text{pool 2}}$ .

**Heating experiments:** For experiments shown in Fig. 1 freeze-dried milled leaf biomass (250–1000 mg) was heated in a glass vessel according to the method of Hamilton et al. (2003) except that temperature programming increments were 12.5°C instead of 25°C.  $\delta^{13}\text{C}$  values of volatile organic compounds were measured at the end of each temperature increment by GC-MS-IRMS. Results shown in Table 1 are for isothermal heating for 20 min at 225°C of dried plant biomass.

**Incubation experiments with fresh plant tissue:** Fresh leaves (15–30 g) were detached from the plant and immediately placed in glass vials (44 ml) and sealed with caps containing a PTFE lined silica septa. Samples ( $n=3-6$ ) were incubated in the dark for 18 h at 25°C and VOCs were measured by GC-MS-IRMS. Table 2 shows the mean values  $\pm$ SD. Results for  $\text{C}_3$  plants are also mean and SD between species given.

**Pectin methoxyl groups:** Carbon isotope signatures of the pectin methoxyl pool were assessed by measuring  $\delta^{13}\text{C}$  values of methanol released by alkaline hydrolysis of freeze-dried biomass. Molar NaOH (1 ml) was added to biomass (200 mg) in a 5 ml reaction vial. The vials were sealed with caps containing PTFE lined silicone septa and incubated for 12 h at 50°C to quantitatively hydrolyse ester methoxyl groups to methanol. Control experiments indicated that no chemical fractionation of carbon isotopes in methanol occurred during the analytical procedure.

**Lignin methoxyl groups:** Carbon isotope signatures of lignin methoxyl groups were assessed by measuring  $\delta^{13}\text{C}$  values of  $\text{CH}_3\text{I}$  released by HI treatment (for 30 min at 100°C) of the biomass fraction remaining after removal of the pectin methoxyl pool by alkaline hydrolysis. Control experiments with aromatic and aliphatic methyl esters indicated the procedure resulted in quantitative conversion of OCH<sub>3</sub> groups to  $\text{CH}_3\text{I}$ . No significant chemical fractionation of carbon isotopes in  $\text{CH}_3\text{I}$  occurred during the analytical procedure conducted as described.

**Sample collection:** The origin of the investigated plant tissues is shown below the tables.

**Carbon isotope  
anomaly in the major  
plant  $\text{C}_1$  pool**

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

### 3. Results

In this study we have employed compound-specific carbon isotope ratio/mass spectrometry (GC-MS/IR-MS) to measure the  $\delta^{13}\text{C}$  of two plant  $\text{C}_1$  pools, ester methoxyl (largely present as pectin), and aromatic ether methoxyl (predominantly present as lignin), and also the  $\delta^{13}\text{C}$  of several VOCs derived from fresh plant material at ambient and elevated temperatures. We initially assessed carbon isotope fractionation on the pectin methyl pool in leaf tissue from ash (*Fraxinus excelsior*) by measuring  $\delta^{13}\text{C}$  in  $\text{CH}_3\text{OH}$  released on alkaline hydrolysis. The  $\delta^{13}\text{C}$  observed for this esterified methyl pool was  $-77.2\text{‰}$ , a remarkably large  $^{13}\text{C}$  fractionation ( $\Delta \sim -45\text{‰}$ ) compared with the overall  $\delta^{13}\text{C}$  of leaf biomass of  $-31.8\text{‰}$ . A biochemical rationale for this striking depletion is however possible. Carboxyl groups in pectin are esterified by the enzyme pectin *O*-methyltransferase (PMT) using S-adenosylmethionine (SAM) as methyl donor. Work on purine alkaloids in several plant species (Weilacher et al., 1996) has suggested that the methyl pool in SAM is significantly depleted ( $\delta^{13}\text{C} \leq -39\text{‰}$ ) relative to the carbohydrate pool ( $\delta^{13}\text{C} = -27\text{‰}$ ). Moreover, enzymic transmethylation involving SAM can entail a substantial kinetic isotope effect (KIE); thus the reaction catalysed by catechol *O*-methyltransferase displays a large fractionation ( $\epsilon = 90$ ) (Hegazi et al., 1979). A similar KIE in the enzymic methylation of pectin by PMT utilising  $^{13}\text{C}$ -depleted SAM as the methyl donor could account for the magnitude of the  $^{13}\text{C}$  depletion observed in the pectin methyl pool.

We next investigated the effect of progressive heating of leaf tissue of ash from 150 to  $300^\circ\text{C}$  on the  $\delta^{13}\text{C}$  of volatiles released (Fig. 1). The main VOCs produced were  $\text{CH}_3\text{Cl}$ ,  $\text{CH}_3\text{OH}$ , acetaldehyde and acetone (Fig. 1a). The  $\delta^{13}\text{C}$  values for both  $\text{CH}_3\text{Cl}$  and  $\text{CH}_3\text{OH}$  (Fig. 1b) were strikingly depleted with respect to biomass ( $\Delta$  between  $-30$  and  $-100\text{‰}$ ). Emissions of  $\text{CH}_3\text{Cl}$  exhibited a  $\delta^{13}\text{C}$  of  $-128\text{‰}$  at  $150^\circ\text{C}$  ( $-147\text{‰}$  at  $40^\circ\text{C}$  (see supplemental material, Table S1, <http://www.copernicus.org/EGU/bg/bgd/1/393/bgd-1-393-sp1.pdf>),  $^{13}\text{C}$  fractionations, which, to the best of our knowledge, are the lightest isotopic values ever observed in a terrestrial carbon compound produced

## Carbon isotope anomaly in the major plant $\text{C}_1$ pool

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

during natural processes. A comparison of the  $\delta^{13}\text{C}$  of the pectin methyl pool with the composite  $\delta^{13}\text{C}$  value calculated on a molar basis for  $\text{CH}_3\text{Cl}$  and  $\text{CH}_3\text{OH}$  released during heating is displayed in Fig. 1b. This isotopic mass balance clearly shows that when production of  $\text{CH}_3\text{Cl}$  and  $\text{CH}_3\text{OH}$  had ceased the composite  $\delta^{13}\text{C}$  value for these compounds closely corresponded with the  $\delta^{13}\text{C}$  value of the pectin methyl pool showing that pectin methoxyl groups are the major source of both  $\text{CH}_3\text{Cl}$  and  $\text{CH}_3\text{OH}$ . Alkaline hydrolysis of the residual material indicated that the total methoxyl pool had been volatilised by  $300^\circ\text{C}$  (Fig. 1c). Measurements conducted using a model system of purified pectin also confirmed that isotopic mass balance was achieved with respect to  $\text{CH}_3\text{Cl}$ ,  $\text{CH}_3\text{OH}$  and pectin methoxyl during the heating cycle (see supplemental material, Fig. S1, <http://www.copernicus.org/EGU/bg/bgd/1/393/bgd-1-393-sp1.pdf>). Furthermore it is evident that relative to the pectin methoxyl pool  $\text{CH}_3\text{Cl}$  is always highly depleted whilst  $\text{CH}_3\text{OH}$  normally exhibits slight but significant enrichment. The  $\delta^{13}\text{C}$  values determined for acetaldehyde and acetone at all stages of the heating programme clearly reflected the isotope signature of bulk leaf biomass (Fig. 1d), unequivocally distinguishing their origin from that of the  $\text{C}_1$  compounds. An explanation for the unprecedented depletion of  $^{13}\text{C}$  in  $\text{CH}_3\text{Cl}$  released during heating of leaf tissue must await elucidation of the mechanism of the solid state reaction of halide ion with pectin (Hamilton et al., 2003).

To determine whether our findings with ash leaves could be replicated with other species we conducted further studies on leaf tissue from trees, grasses and halophytes including plants from  $\text{C}_3$ ,  $\text{C}_4$  and CAM plant categories involving heating of dried biomass isothermally at  $225^\circ\text{C}$  (Table 1). For all species examined we observed a large fractionation between the pectin methyl pool and bulk biomass ( $\Delta \sim -33\text{‰}$ , range  $-21$  to  $-45\text{‰}$ ). In general the isotope signatures for  $\text{CH}_3\text{OH}$  reflected those of the pectin methyl pool whilst those of  $\text{CH}_3\text{Cl}$  were considerably more depleted in  $^{13}\text{C}$  ( $\Delta \sim -30\text{‰}$ ). Signatures of  $\text{C}_2$  VOCs mirrored in general those of bulk biomass. Although the  $\text{Br}^-$ ,  $\text{I}^-$  and  $\text{CN}^-$  content of most plant tissues are insufficient to permit measurements on emissions of the corresponding substituted methanes, experiments performed with pu-

**Carbon isotope  
anomaly in the major  
plant  $\text{C}_1$  pool**

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

rified apple pectin supplemented with the different ions revealed that  $\text{CH}_3\text{Br}$ ,  $\text{CH}_3\text{I}$  and  $\text{CH}_3\text{CN}$  released on heating were also highly depleted in  $^{13}\text{C}$ .

We extended our measurements to VOCs released from freshly collected plant material at ambient temperatures. Methanol, ethanol, acetaldehyde and acetone were naturally released in sufficient quantities for analytical measurements. The results were similar to those obtained at higher temperatures (Table 2). Thus  $\text{CH}_3\text{OH}$ , like the pectin methyl pool, was highly depleted in  $^{13}\text{C}$  ( $\Delta \sim -33\%$  relative to bulk biomass). Carbon isotope signatures of ethanol and acetone were close to that of bulk biomass whilst acetaldehyde showed slight enrichment in  $^{13}\text{C}$  ( $\Delta \sim 5\%$ ).  $\delta^{13}\text{C}$  values for  $\text{CH}_3\text{Cl}$  could not be measured (except for the halophytes) as amounts were below the detection limit of the analytical method. As has been shown previously  $\text{CH}_3\text{Cl}$  formation in fresh leaves with a high water content is generally low (Hamilton et al., 2003).

In addition to the pectin methyl pool the other important  $\text{C}_1$  pool in plant cell walls is represented by aromatic ether methoxyl groups which can comprise up to 18% of lignin. Lignin is a major component of wood (up to 31%) and is also found in smaller quantities in leaves and grasses ( $\sim 5\%$ ). We therefore measured  $\delta^{13}\text{C}$  values of the lignin methoxyl pool in plant tissue from several species after conversion to  $\text{CH}_3\text{I}$  with HI subsequent to removal of the pectin methyl pool by alkaline hydrolysis (Table 3). Depletion in  $^{13}\text{C}$  of lignin methoxyl groups in wood ( $\Delta \sim -13\%$ ) relative to bulk biomass was substantial although not as dramatic as that observed for lignin methoxyl groups in leaves (mean of all leaves  $\Delta \sim -29\%$ , range  $-20$  to  $-38\%$ ). These findings explains the widely reported  $^{13}\text{C}$  depletion of lignin relative to other major plant components (Benner et al., 1987; Schweizer et al., 1999; Fernandez et al., 2003; Hobbie and Werner, 2004) which has previously been attributed to  $^{13}\text{C}$  fractionation in aromatic amino acids involved in lignin biosynthesis. Thus assuming a methoxyl content of 15–20% a depletion of  $-13\%$  in methoxyl carbon readily explains the observed 2–3% difference between  $\delta^{13}\text{C}$  of lignin and bulk biomass of wood. Similarly the much larger depletion observed in methoxyl carbon in leaf tissue provides an explanation for the 3–7%  $^{13}\text{C}$  depletion of lignin relative to bulk biomass in leaves of both  $\text{C}_3$  and  $\text{C}_4$  plants. Archaeological and

---

**Carbon isotope  
anomaly in the major  
plant  $\text{C}_1$  pool**F. Keppler et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion



fossil wood specimens are often used to provide information on palaeoenvironmental and palaeoclimatic conditions in the geological record (van Bergen and Poole, 2002). Hence alterations to wood which involve cleavage of the isotopically light methoxyl groups will be critically important in interpretation of the significance of the isotope signatures of individual wood components.

#### 4. Conclusions

We have summarised our findings on the  $\delta^{13}\text{C}$  values of methoxyl pools in plants and some plant derived  $\text{C}_1$  VOCs in Fig. 2 where they are related to fractionations reported in the literature for bulk biomass of various categories of plant and other terrestrial carbon sources (Whiticar, 1996). The depletion between bulk plant biomass and plant methoxyl pools ranges from  $-11$  to  $-46\text{‰}$  with the pectin  $\text{C}_1$  pool generally more depleted than the lignin  $\text{C}_1$  pool. The fractionation associated with the methoxyl pools is retained and even further enhanced during their conversion to  $\text{C}_1$  VOCs. For biogenic VOC emissions from living vegetation, the more labile pectin pool rather than the lignin pool is likely to be the major source. However during biomass burning both pectin and lignin  $\text{C}_1$  pools will contribute to  $\text{C}_1$  VOC production. In soils under aerobic conditions demethylation of pectin and lignin, both enzymically (e.g. Fall and Benson, 1996; Ander and Eriksson, 1985) and abiotically (e.g. Dec et al., 2001, Keppler et al., 2000) is very probably an important source of several  $\text{C}_1$  VOCs. Under anoxic conditions  $\text{C}_1$  compounds from both  $\text{C}_1$  pools can act as substrates for methylotrophic methanogenic bacteria forming  $\text{CH}_4$  (Whiticar, 1999; Cicerone and Oremland, 1988). It has been assumed to date that the carbon signature of a substrate for methanogens in a specific environment is broadly similar to that of the bulk organic matter present, provided severe substrate depletion has not occurred (Whiticar, 1999). If, however in the upper horizon of wetlands, peat bogs and rice paddies, methanogens are utilising methanol and other  $\text{C}_1$  substrate from the plant methoxyl pool with  $\delta^{13}\text{C}$  values averaging  $-50\text{‰}$ , the production of  $\text{CH}_4$  more highly depleted than previously envisaged

### Carbon isotope anomaly in the major plant $\text{C}_1$ pool

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

might be expected. Indeed  $\delta^{13}\text{C}$  values for  $\text{CH}_4$  derived from the plant methoxyl pool may be of the same order as the lowest observed for  $\text{CH}_4$  produced by bacterial carbonate reduction (Whiticar, 1999; Cicerone and Oremland, 1988). Since approximately 2.5% of carbon in plant biomass is methoxyl carbon (Galbally and Kirstine, 2002), any isotopically-based discussion of global carbon cycling must give consideration to this isotopically anomalous  $\text{C}_1$  pool.

Our findings may also have some relevance to the search for ancient life on earth and for extraterrestrial life. The striking depletion of  $\delta^{13}\text{C}$  in methoxyl carbon consequent on the biochemistry of  $\text{C}_1$  metabolism in plants may well extend to many other organisms which utilise S-adenosylmethionine as a methyl donor in *O*-methyltransferase reactions and could even serve to distinguish biologically formed methyl esters and ethers from those generated abiotically. Conversely, the exceptional  $^{13}\text{C}$  fractionation during abiotic production of  $\text{CH}_3\text{Cl}$  from biomass (which we have reproduced by heating hydrochlorides of methyl esters of amino acids, (see supplemental material, Fig. S2, <http://www.copernicus.org/EGU/bg/bgd/1/393/bgd-1-393-sp1.pdf>) suggest that caution is necessary in interpreting such fractionation as unequivocal evidence of life.

In conclusion the fractionation of carbon isotopes by the principal *O*-methyltransferase enzymes in plants appears to be of the same order as that achieved by ribulose biphosphate carboxylase-oxygenase (Rubisco) in photosynthesis. Hitherto it has been assumed that within a given photosynthetic category the carbon isotope signature of specific chemical components and specific intramolecular sites within such components does not differ from the carbon isotope signature of bulk biomass by more than 12‰ (Weilacher et al., 1996; Hobbie and Werner, 2004). Our results indicate a  $^{13}\text{C}$  depletion relative to bulk biomass of up to 45‰ for methoxyl carbon in plants, the largest carbon isotope fractionation ever observed in the plant kingdom. This isotope anomaly should prove not only an invaluable tool in tracing the path of such  $\text{C}_1$  carbon in the environment but also provide a new insight into the global cycling of many  $\text{C}_1$  atmospheric trace gases and the biochemical pathways involved.

# Carbon isotope anomaly in the major plant $\text{C}_1$ pool

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

*Acknowledgements.* The authors thank W. Michaelis, J. Raven, and T. Röckmann for helpful advice regarding the manuscript; W. Meier-Augenstein, A. Downey, K. Redeker, E. Tujek, B. Ferguson, T. Kennedy for technical assistance and the European Commission for a Marie Curie-Research Training Grant (MCFI-2002-00022) awarded to F. K. Supported by grants from EPSRC GR/R03099/01, QUB SRIF for Environmental Engineering and Biotechnology, and INI TDP Centre of Excellence.

## References

- Ander, P. and Eriksson, K.-E.: Methanol formation during lignin degradation by *Phanerochaeta chrysosporium*, *Appl. Microbial. Biotechnol.*, 21, 96–102, 1985.
- Benner, R., Fogel, M. L., Sprague, E. K., and Hodson, R. E.: Depletion of  $^{13}\text{C}$  lignin and its implications for stable isotope studies, *Nature*, 329, 708–710 (1987).
- Bill, M., Conrad, M. E., and Goldstein, A. H.: Stable carbon isotope composition of atmospheric methyl bromide, *Geophys. Res. Lett.*, 31, L04109, DOI:10.1029/2003GL018639, 2004.
- Cerling, T. E., Harris, J. M., MacFadden, B. J. et al.: Global vegetation change through the Miocene/Pliocene boundary, *Nature*, 389, 153–158, 1997.
- Cicerone, R. J. and Oremland R. S.: Biogeochemical aspects of atmospheric methane, *Global Biogeochem Cycles*, 2, 299–327, 1988.
- Dec, J., Haider, K., and Bollag, J.-M.: Decarboxylation and demethoxylation of naturally occurring phenols during coupling reactions and polymerization, *Soil Science*, 166, 660–671, 2001.
- Ehleringer, J. R., Bowling, D. R., Flanagan, L. B. et al.: Stable isotopes and carbon cycle processes in forests and grasslands, *Plant. Biol.*, 4, 181–189, 2002.
- Fall, R. and Benson, A. A.: Leaf methanol – the simplest natural product from plants, *Trends in Plant Science*, 1, 296–301, 1996.
- Fernandez, I., Mahieu, N., and Cadisch, G.: Carbon isotopic fractionation during decomposition of plant materials of different quality, *Global Biogeochem Cycles*, 17, DOI:10.1029/2001GB001834, 2003.
- Galbally, I. E. and Kirstine, W.: The production of methanol by flowering plants and the global cycle of methanol, *J. Atmos. Chem.*, 43, 195–229, 2002.

**BGD**

1, 393–412, 2004

## Carbon isotope anomaly in the major plant $\text{C}_1$ pool

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

- Goldstein, A. H. and Shaw, S. L.: Isotopes of volatile organic compounds: an emerging approach for studying atmospheric budgets and chemistry, *Chem. Rev.*, 103, 5025–5048, 2003.
- Griffiths, H. (ed.): *Stable Isotopes and the Integration of Biological, Ecological and Geochemical Processes*, Bios Scientific Publishers, Oxford, 1998.
- 5 Hamilton, J. T. G., McRoberts, W. C., Keppler, F., Kalin, R. M., and Harper, D. B.: Chloride methylation by plant pectin: an efficient environmentally significant process, *Science*, 301, 206–209, 2003.
- Harper, D. B., Hamilton, J. T. G., Ducrocq, V. et al.: The distinctive isotopic signature of plant-derived chloromethane: possible application in constraining the atmospheric chloromethane budget, *Chemosphere*, 52, 433–436, 2003.
- 10 Hayes, J. M.: Fractionation of the isotopes of carbon and hydrogen in biosynthetic processes, *Rev Mineral Geochem*, 43, 225–277, 2001.
- Hegazi, M. F., Borchardt, R. T., and Schowen, R. L.:  $\alpha$ -Deuterium and carbon-13 isotope effects for methyl transfer catalyzed by catechol *O*-methyltransferase.  $S_N2$ -like transition state, *J. Am. Chem. Soc.*, 101, 4359–4364, (1979).
- 15 Heikes, B. G., Chang, W., Pilson, M. E. Q. et al.: Atmospheric methanol budget and ocean implication, *Global Biogeochem Cycles*, 16, DOI:10.1029/2002GB001895, 2002.
- Hobbie, E. A. and Werner, R.: Intramolecular, compound-specific, and bulk carbon isotope patterns in  $C_3$  and  $C_4$  plants: a review and synthesis, *New Phytologist*, 161, 371–385, 2004.
- 20 Kalin, R. M., Hamilton, J. T. G., Harper, D. B. et al.: Continuous flow stable isotope methods for study of  $\delta^{13}C$  fractionation during halomethane production and degradation, *Rapid Commun Mass Spectrom*, 15, 357–363, 2001.
- Keppler, F., Eiden, R., Niedan, V., Pracht, J. and Schöler, H. F.: Halocarbons produced by natural oxidation processes during degradation of organic matter, *Nature*, 403, 298–301, 2000.
- 25 Lajtha, K. and Michener, R. B. (eds.): *Stable Isotopes in Ecology and Environmental Science*, Blackwell Scientific Publications, Oxford, 1994.
- Michaelis, W., Seifert, R., Nauhaus, K. et al.: Microbial reefs in the Black Sea fueled by anaerobic oxidation of methane, *Science*, 297, 1013–1015, 2002.
- 30 Montzka, S. A., Fraser, P. J., Butler, J. H. et al.: “Controlled Substances and Other Source Gases” in WMO Scientific Assessment of Ozone Depletion. Global Ozone Research and Monitoring Project, Report No. 47, World Meteorological Organization, Geneva, [http://www.wmo.ch/web/arep/reports/ozone\\_2002/06.chapter1.pdf](http://www.wmo.ch/web/arep/reports/ozone_2002/06.chapter1.pdf), 2003.

## Carbon isotope anomaly in the major plant $C_1$ pool

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

- O'Dowd, C. D., Jimenez, J. C. Bahreini, R. et al.: Marine aerosol formation from biogenic iodine emissions, *Nature*, 417, 632–636, 2002.
- O'Leary, M. H.: Carbon isotope fractionations in plants, *Phytochemistry*, 20, 553–567, 1981.
- Sanhueza, E., Holzinger, R., Kleiss, B., Donoso, L., and Crutzen, P. J.: New insights in the global cycle of acetonitrile: release from the ocean and dry deposition in the tropical savanna of Venezuela, *Atmos. Chem. Phys. Discuss.*, 4, 275–280, 2004.
- Schweizer, M., Fear, F., and Cadisch, G.: Isotopic ( $^{13}\text{C}$ ) fractionation during plant residue decomposition and its implications for soil organic matter studies, *Rapid Commun. Mass Spectrom.*, 13, 1284–1290, 1999.
- Sherwood Lollar, B., Westgate, T. D., Ward, J. A. et al.: Abiogenic formation of alkanes in the Earth's crust as a minor source for global hydrocarbon reservoirs, *Nature*, 416, 522–524, 2002.
- Thompson, A. E., Anderson R. S., Rudolph, J., and Huang, L.: Stable carbon isotopes signatures of background tropospheric chloromethane and CFC113, *Biogeochemistry*, 60, 191–211, 2002.
- van Bergen, P. F. and Poole, I.: Stable carbon isotopes of wood: a clue to palaeoclimate?, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 182, 31–45, 2002.
- Warneke, C., Karl, T., Judmaier, H. et al.: Acetone, methanol, and other partially oxidised volatile organic emissions from dead plant matter by abiotical processes: significance for atmospheric  $\text{HO}_x$  chemistry, *Global Biogeochem. Cycles*, 13, 9–17, 1999.
- Weilacher, T., Gleixner, G., and Schmidt, H. L.: Carbon isotope pattern in purine alkaloids a key to isotope discriminations in  $\text{C}_1$  compounds, *Phytochemistry*, 41, 1073–1077, 1996.
- Whiticar, M. J.: Stable isotope geochemistry of coals, humic kerogens and related natural gases, *Int. J. Coal. Geol.*, 32, 191–215, 1996.
- Whiticar, M. J.: Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane, *Chem. Geol.*, 161, 291–314, 1999.
- Wuebbles, D. J. and Hayhoe, K.: Atmospheric methane and global change, *Earth-Science Reviews*, 57, 177–210, 2002.
- Yakir, D.: Sphere of influence, *Nature*, 416, 795, 2002.

**Carbon isotope anomaly in the major plant  $\text{C}_1$  pool**

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

## Carbon isotope anomaly in the major plant C<sub>1</sub> pool

F. Keppler et al.

**Table 1.**  $\delta^{13}\text{C}$  values<sup>1</sup> of biomass, pectin methoxyl groups and VOCs produced upon heating (225°C) dried biomass.

Plant common name ( <i>species</i> )	Biomass (B) ( $\delta^{13}\text{C}$ )	Pectin methoxyl (PM) ( $\delta^{13}\text{C}$ )	$\Delta^{13}\text{C}$ (PM-B) ( $\delta^{13}\text{C}_{\text{PM}} - \delta^{13}\text{C}_{\text{B}}$ )	Methanol ( $\delta^{13}\text{C}$ )	Chloromethane ( $\delta^{13}\text{C}$ )	Acetaldehyde ( $\delta^{13}\text{C}$ )	Acetone ( $\delta^{13}\text{C}$ )
<b>C<sub>3</sub>-leaf tissue</b>							
European ash ( <i>Fraxinus excelsior</i> ) <sup>2</sup>	-31.8 ± 0.2	-77.2 ± 0.1	-45.4	-70.4 ± 2.5	-98.4 ± 2.2	-34.0 ± 1.1	-36.2 ± 2.8
Wych Elm ( <i>Ulmus glabra</i> ) <sup>2</sup>	-28.4 ± 0.1	-61.7 ± 0.2	-33.3	-61.1 ± 1.8	-85.3 ± 3.1	-31.2 ± 0.7	-33.2 ± 2.1
Hazelnut ( <i>Corylus avellana</i> ) <sup>2</sup>	-29.1 ± 0.1	-66.3 ± 0.2	-37.2	-64.0 ± 1.8	-96.0 ± 2.1	-33.2 ± 1.3	34.5 ± 1.9
English oak ( <i>Quercus robur</i> ) <sup>2</sup>	-31.4 ± 0.1	-74.4 ± 0.2	-43.0	-75.0	-104.3	-28.9	-31.7
Norway maple ( <i>Acer platanoides</i> ) <sup>2</sup>	-27.6 ± 0.2	-61.3 ± 1.0	-35.7	-58.3	-92.4	-27.0	-21.1
Horse chestnut ( <i>Aesculus hippocastanum</i> ) <sup>2</sup>	-31.7 ± 0.2	-66.3 ± 2.0	-34.6	-60.4	-94.9	-29.5	-29.7
Scots pine ( <i>Pinus sylvestris</i> ) <sup>2</sup>	-27.6 ± 0.1	-53.7 ± 0.2	-26.1	-49.3	-86.8	-24.9	-30.8
Cocksfoot ( <i>Dactylis glomerata</i> ) <sup>3</sup>	-29.3 ± 0.2	-50.7 ± 0.2	-21.4	-52.6	-72.8	-28.1	-34.3
Glasswort ( <i>Salicornia spp</i> ) <sup>4</sup>	-28.6 ± 0.1	-53.7 ± 0.2	-25.1	-42.0	-76.3	-27.1	-34.6
<b>Mean of C<sub>3</sub> plants</b>	<b>-29.5</b>	<b>-62.8</b>	<b>-33.5</b>	<b>-59.2</b>	<b>-89.7</b>	<b>-29.3</b>	<b>-31.8</b>
(SD between C <sub>3</sub> plants)	(± 1.7)	(± 9.2)	(± 8.1)	(± 10.3)	(± 10.3)	(± 3.0)	(± 4.5)
<b>C<sub>4</sub>-leaf tissue</b>							
Maize ( <i>Zea mays</i> ) <sup>4</sup>	-11.0 ± 0.1	-40.5 ± 0.6	-29.5	-39.7	-91.3	-11.6	-16.9
<b>CAM-leaf tissue</b>							
Saltwort ( <i>Batis maritima</i> ) <sup>4</sup>	-25.6 ± 0.4	-63.3 ± 0.4	-37.7	-64.9	-78.3	-22.4	-25.3
Scarlet paintbrush ( <i>Crassula falcata</i> ) <sup>4</sup>	-17.9 ± 0.1	-51.1 ± 0.5	-33.3	-41.1	-81.4	-16.2	-20.8

<sup>1</sup>All values in ‰, either mean of two samples or n=3-5 ± SD, for sample preparation and analytical measurements see Methods (25). <sup>2</sup>Leaves were collected at Crossgar, N. Ireland in October 2002. <sup>3</sup>Leaves were collected at Crossgar, N. Ireland in July 2003. <sup>4</sup>Greenhouse-grown in N. Ireland 2003.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

## Carbon isotope anomaly in the major plant C<sub>1</sub> pool

F. Keppler et al.

**Table 2.**  $\delta^{13}\text{C}$  values<sup>1</sup> of biomass, pectin methoxyl pool and VOCs emitted at 25°C from fresh plant tissue.

Plant common name ( <i>species</i> )	Biomass (B) ( $\delta^{13}\text{C}$ )	Pectin methoxyl (PM) ( $\delta^{13}\text{C}$ )	$\Delta^{13}\text{C}$ (PM-B) ( $\delta^{13}\text{C}_{\text{PM}} - \delta^{13}\text{C}_{\text{B}}$ )	Methanol ( $\delta^{13}\text{C}$ )	Acetaldehyde ( $\delta^{13}\text{C}$ )	Ethanol ( $\delta^{13}\text{C}$ )	Acetone ( $\delta^{13}\text{C}$ )
<b>C<sub>3</sub>-leaf<sup>2</sup></b>							
European ash ( <i>Fraxinus excelsior</i> )	-27.9 ± 0.2	-73.7 ± 1.0	-45.8	-73.5 ± 0.7	-22.7 ± 0.4	-28.5 ± 0.5	-31.3 ± 1.4
Wych elm ( <i>Ulmus glabra</i> )	-28.7 ± 0.1	-68.9 ± 0.1	-40.2	-82.9 ± 5.9	-25.6 ± 1.9	-30.1 ± 2.0	-26.7 ± 3.1
Hazelnut ( <i>Corylus avellana</i> )	-33.6 ± 0.2	-64.6 ± 0.4	-31.0	-63.5 ± 2.8	-25.9 ± 2.5	-30.5 ± 1.2	-26.3 ± 2.4
English oak ( <i>Quercus robur</i> )	-30.8 ± 0.1	-69.2 ± 0.3	-38.4	-76.6 ± 3.9	-23.9 ± 0.5	-29.9 ± 0.3	-28.8 ± 2.9
European beech ( <i>Fagus sylvatica</i> )	-31.8 ± 0.2	-68.2 ± 1.0	-36.4	-84.2 ± 2.6	-25.8 ± 1.4	-31.3 ± 1.1	-27.5 ± 3.2
Norway maple ( <i>Acer platanoides</i> )	-33.6 ± 0.1	-63.1 ± 0.6	-29.5	-70.3 ± 0.7	-27.8 ± 1.4	-33.7 ± 0.5	-26.9 ± 1.2
Horse chestnut ( <i>Aesculus hippocastanum</i> )	-31.7 ± 0.3	-73.4 ± 0.1	-41.7	-71.0 ± 2.4	-20.8 ± 0.8	-26.9 ± 1.2	-23.6 ± 2.4
Scots pine ( <i>Pinus sylvestris</i> )	-28.2 ± 0.2	-57.3 ± 0.3	-29.1	-60.2 ± 1.5	-24.3 ± 1.7	-23.9 ± 1.4	-31.6 ± 2.5
Cocksfoot ( <i>Dactylis glomerata</i> )	-29.3 ± 0.2	-50.7 ± 0.2	-21.4	-51.9 ± 0.2	-22.7 ± 1.1	-29.0 ± 1.4	-30.6 ± 1.4
Yorkshire fog ( <i>Holcus lanata</i> )	-31.3 ± 0.3	-57.1 ± 0.2	-25.8	-65.4 ± 1.3	-27.5 ± 0.9	-31.6 ± 1.0	-26.7 ± 0.5
Glasswort ( <i>Salicornia sp</i> ) <sup>3</sup>	-28.6 ± 0.1	-53.7 ± 0.2	-25.1	-50.4 ± 0.9	-26.8 ± 0.7	-28.5 ± 1.5	-29.4 ± 2.1
<b>Mean of C<sub>3</sub> plants</b>	<b>-30.5</b>	<b>-63.6</b>	<b>-33.1</b>	<b>-68.2</b>	<b>-24.9</b>	<b>-29.4</b>	<b>-28.1</b>
(SD between C <sub>3</sub> plant species)	(± 2.1)	(± 7.9)	(± 7.8)	(± 11.2)	(± 2.2)	(± 2.6)	(± 2.5)
<b>C<sub>4</sub>-leaf</b>							
Maize ( <i>Zea mays</i> ) <sup>3</sup>	-11.0 ± 0.1	-40.5 ± 0.6	-29.5	-52.9 ± 0.8	-9.3 ± 1.3	-15.2 ± 0.5	-16.6 ± 1.2
<b>CAM-leaf</b>							
Saltwort ( <i>Batis maritima</i> ) <sup>3</sup>	-25.6 ± 0.4	-63.3 ± 0.4	-37.7	-60.0 ± 0.9	-24.4 ± 1.7	-23.8 ± 1.2	-27.8 ± 0.3
Scarlet paintbrush ( <i>Crassula falcata</i> ) <sup>3</sup>	-17.9 ± 0.1	-51.1 ± 0.9	-33.2	-55.9 ± 2.1	-10.3 ± 0.5	-17.2 ± 0.2	-19.8 ± 0.7

<sup>1</sup>All values in ‰ ± SD (n=3–5), analytical measurements see Methods. <sup>2</sup>Fresh leaves were collected at Crossgar, N. Ireland in July 2003. <sup>3</sup>Greenhouse-grown in N. Ireland 2003.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

**Table 3.**  $\delta^{13}\text{C}$  values<sup>1</sup> of biomass and the lignin and pectin methoxyl pools of plant tissues.

Plant common name ( <i>species</i> )	Biomass (B) ( $\delta^{13}\text{C}$ )	Lignin methoxyl (LM) ( $\delta^{13}\text{C}$ )	$\text{D}^{13}\text{C}$ (LM-B) ( $\delta^{13}\text{C}_{\text{LM}} - \delta^{13}\text{C}_{\text{B}}$ )	Pectin methoxyl (PM) ( $\delta^{13}\text{C}$ )	$\text{D}^{13}\text{C}$ (PM-B) ( $\delta^{13}\text{C}_{\text{PM}} - \delta^{13}\text{C}_{\text{B}}$ )
<b>C<sub>3</sub>-wood<sup>2</sup></b>					
European ash ( <i>Fraxinus excelsior</i> )	-24.5 ± 0.7	-36.9 ± 1.2	-12.4	-43.1 ± 0.6	-18.6
English oak ( <i>Quercus robur</i> )	-29.4 ± 0.2	-41.1 ± 1.3	-11.7	-44.2 ± 1.2	-14.8
Sweet osmanthus ( <i>Osmanthus fragans</i> )	-27.3 ± 0.1	-41.7 ± 1.2	-14.7	-53.3 ± 1.1	-26.0
Geronggang ( <i>Cratogeomys</i> sp)	-26.5 ± 0.3	-39.5	-13.0	n.d.	-
Tasmanian oak ( <i>Eucalyptus delegatensis</i> )	-26.3 ± 0.2	-37.7	-11.4	-45.4 ± 0.7	-18.1
Dark red meranti ( <i>Shorea</i> sp)	-28.2 ± 0.2	-44.0	-15.8	-45.6 ± 0.3	-17.4
Utile ( <i>Entandrophragma utile</i> )	-27.1 ± 0.2	-39.5	-12.4	n.d.	-
<b>Mean of wood</b>	<b>-27.1</b>	<b>-40.1</b>	<b>-13.0</b>	<b>-46.3</b>	<b>-19.2</b>
<b>C<sub>3</sub>-leaf<sup>3</sup></b>					
European ash ( <i>Fraxinus excelsior</i> )	-27.9 ± 0.2	-65.5	-37.6	-73.7 ± 1.0	-45.8
English oak ( <i>Quercus robur</i> )	-30.8 ± 0.1	-62.2	-31.4	-69.2 ± 0.3	-38.4
European beech ( <i>Fagus sylvatica</i> )	-31.8 ± 0.2	-66.2	-34.4	-68.2 ± 1.0	-36.4
Norway maple ( <i>Acer platanoides</i> )	-33.6 ± 0.2	-61.4	-27.8	-63.1 ± 0.6	-29.5
Scots pine ( <i>Pinus sylvestris</i> )	-27.6 ± 0.1	-51.7	-24.1	-53.7 ± 0.3	-26.1
Cocksfoot grass ( <i>Dactylis glomerata</i> )	-29.3 ± 0.2	-53.5	-24.2	-50.7 ± 0.2	-21.4
<b>Mean of C<sub>3</sub>-leaves</b>	<b>-30.2</b>	<b>-60.1</b>	<b>-29.9</b>	<b>-63.1</b>	<b>-32.9</b>
<b>C<sub>4</sub>-leaf</b>					
Sugar cane ( <i>Saccharum officinarum</i> ) <sup>4</sup>	-11.9 ± 0.1	-42.1	-30.2	-36.0 ± 0.9	-24.1
Savanna grass ( <i>Hyparrhenia</i> sp) <sup>5</sup>	-12.8 ± 0.2	-33.1	-20.2	n.d.	-
Maize ( <i>Zea mays</i> ) <sup>6</sup>	-11.0 ± 0.1	-47.3	-36.3	-40.5 ± 0.6	-29.5
<b>CAM-leaf</b>					
Saltwort ( <i>Batis maritima</i> ) <sup>6</sup>	-25.6 ± 0.4	-52.4	-25.7	-63.3 ± 0.4	-37.3
Scarlet paintbrush ( <i>Crassula falcata</i> ) <sup>6</sup>	-17.9 ± 0.1	-49.6	-31.7	-51.1 ± 0.5	-33.2
<b>Mean of C<sub>3</sub>, C<sub>4</sub> and CAM leaves</b>			<b>-29.0</b>		<b>-32.2</b>

<sup>1</sup>All values in ‰, either mean of two samples or  $n=3-5 \pm \text{SD}$ ; n.d.: - not detectable, for sample preparation and analytical measurements see Methods (25). <sup>2</sup>Wood samples were collected from Cameroon, Indonesia, Malaysia, and N. Ireland. <sup>3</sup>Leaves were collected at Crossgar, N. Ireland in July 2003. <sup>4</sup>Sampled from South Africa. <sup>5</sup>Sampled from Cote d'Ivoire. <sup>6</sup>Greenhouse-grown in N. Ireland 2003.

## Carbon isotope anomaly in the major plant C<sub>1</sub> pool

F. Keppler et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version


Interactive Discussion

© EGU 2004



## Carbon isotope anomaly in the major plant C<sub>1</sub> pool

F. Keppler et al.




**Fig. 1.** Amounts and isotopic signatures of several volatile organic compounds formed during progressive heating of lyophilised ash leaves. **(a)** Cumulative amounts of methanol, chloromethane, acetaldehyde and acetone formed are shown on a molar basis. Each point is the mean of three replicate analyses of independent samples ( $n=3$ ). Error bars shown for CH<sub>3</sub>OH are typical of SDs for all compounds.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2004

# Carbon isotope anomaly in the major plant C<sub>1</sub> pool

F. Keppler et al.



**Fig. 1. (b)** Carbon isotopic composition of accumulated  $\text{CH}_3\text{OH}$  and  $\text{CH}_3\text{Cl}$  at each temperature during progressive heating. Also shown is the composite  $\delta^{13}\text{C}$  values calculated on a molar basis for  $\text{CH}_3\text{OH}$  and  $\text{CH}_3\text{Cl}$  released during heating. For reference the measured initial  $\delta^{13}\text{C}$  of the pectin methoxyl pool is displayed. Vertical bars show SD for triplicate samples.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Print Version

Interactive Discussion

© EGU 2004

# Carbon isotope anomaly in the major plant C<sub>1</sub> pool

F. Keppler et al.




**Fig. 1. (c)** Remaining pectin methoxyl pool (PM) after each heating step in relation to the formation of methanol and chloromethane.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Print Version](#)
[Interactive Discussion](#)

# Carbon isotope anomaly in the major plant C<sub>1</sub> pool

F. Keppler et al.




**Fig. 1. (d)** Carbon isotope composition of accumulated acetaldehyde and acetone at various temperatures during progressive heating of ash leaf biomass with reference to the  $\delta^{13}\text{C}$  value of the original bulk biomass. Error bars shown for acetaldehyde are also typical of those for acetone.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Print Version](#)
[Interactive Discussion](#)

© EGU 2004

# Carbon isotope anomaly in the major plant C<sub>1</sub> pool

F. Keppler et al.



**Fig. 2.** Schematic diagram displaying  $^{13}\text{C}$  depletion of methoxyl groups relative to bulk biomass of terrestrial plants and their relationship to biospheric  $\text{C}_1$  and  $\text{C}_{n+1}$  VOCs. Data for carbon isotopic composition of major carbon pools were taken from Whiticar (1996). Carbon isotope signatures for VOCs are related to sources, degradation steps after formation are not taken into account.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004