



**HAL**  
open science

## Seasonal cycling of phosphorus in the southern bight of the North Sea

C. van Der Zee, L. Chou

► **To cite this version:**

C. van Der Zee, L. Chou. Seasonal cycling of phosphorus in the southern bight of the North Sea. Biogeosciences Discussions, 2004, 1 (1), pp.681-707. hal-00297773

**HAL Id: hal-00297773**

**<https://hal.science/hal-00297773>**

Submitted on 18 Jun 2008

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*Biogeosciences Discussions* is the access reviewed discussion forum of *Biogeosciences*

---

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

---

# Seasonal cycling of phosphorus in the southern bight of the North Sea

C. van der Zee and L. Chou

Laboratoire d'Océanographie Chimique et Géochimie des Eaux, Université Libre de Bruxelles,  
Campus de la Plaine – CP 208, Boulevard du Triomphe, B-1050 Brussels, Belgium

Received: 22 September 2004 – Accepted: 5 October 2004 – Published: 12 October 2004

Correspondence to: C. van der Zee (cvdzee@ulb.ac.be)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

## Abstract

We have investigated the seasonal cycle of nutrients and the phosphorus speciation, i.e. dissolved inorganic and organic phosphorus (DIP and DOP) and particulate inorganic and organic phosphorus (PIP and POP), for 10 stations in the Belgian coastal zone. The Belgian part of the southern North Sea is strongly influenced by the river plumes of the Rhine, Meuse and Scheldt. In winter, high nutrient concentrations are observed, whereas in April–May these have all been consumed during the spring bloom and silica or phosphorus limitation develops. The phosphate concentrations increase rapidly again in summer-fall, whereas nitrate and silicate return to their winter values much later. This shows the efficient phosphorus recycling that takes place in the water column. The DOP concentration exhibits two peaks during a seasonal cycle: one in April–May when the phosphate concentration is at its lowest and a second one in fall when the POP content decreases. This indicates two periods of increased phosphorus recycling activity. The seasonal cycle of the DOP is different from that of dissolved organic nitrogen (DON).

## 1. Introduction

The southern bight of the North Sea receives nutrients from domestic, industrial and agricultural wastewater discharge, river inputs and atmospheric deposition. Most of these nutrients are linked to anthropogenic activities that have significantly increased the riverine flux of nitrogen (N) and phosphorus (P) to the coastal zones of the North Sea (North Sea Task Force, 1993). The flux of another major nutrient, dissolved silicate, has remained relatively constant because its major land source is through chemical weathering of silicate minerals. As a consequence, the nutrient ratios in the riverine flux to the coastal zone have drastically changed. This results in a concomitant modification of phytoplankton species composition and thus of the coastal ecosystem. For example, a reduced importance of the diatom bloom and the frequent occurrence of

**BGD**

1, 681–707, 2004

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

massive *Phaeocystis* blooms have been observed (Lancelot et al., 1987).

Turner et al. (2003a, b) reviewed data for dissolved inorganic nitrogen, dissolved inorganic phosphate and dissolved silicate in the world's largest rivers in order to predict future aquatic nutrient limitations assuming the predicted increase in fertilizer use (which has a N:P ratio of 26). The authors found that the coastal areas in northern Europe are most likely to have both P and Si limitation. At the international conference on the protection of the North Sea (London, 1987), all countries surrounding the North Sea agreed on reducing the anthropogenic input of nutrients by 50% between 1985 and 1995 for areas where nutrients cause pollution. As a result of the active reduction of P in wastewaters, the riverine P inputs have declined steadily, whereas this is much less the case for N (Behrendt et al., 2000). Recently, Skogen et al. (2004) ran model scenarios for the years of 1988 and 1989 with reduced nutrient inputs from the rivers to the North Sea. Simulation results indicated that a 50% decrease in the loads of N and P would lead to a decline in the primary production by 10–30% in the southern North Sea. A reduction only in the P load has shown that the primary production in the southern North Sea is limited by P, while N is limiting in the northern North Sea.

Philippart et al. (2000) showed that the phytoplankton community changed drastically in the eutrophic Marsdiep (westernmost inlet of the Wadden Sea) both between 1976 and 1978 and again between 1987 and 1988, whereas it was stable before (1974–1976), in-between (1978–1987) and thereafter (1988–1994). These major changes coincide with changes in absolute and relative (TN:TP) nutrient concentrations. In 1977, the system shifted from a phosphorus-controlled situation, towards a nitrogen-controlled environment, and re-shifted towards P control in 1987–1988. This coincidence implies a strong relationship between N:P ratios and the phytoplankton community structure. Phosphate concentrations dropped a little further since 1992 in the Marsdiep and since 1994 (until 2000) a decrease in primary productivity, chlorophyll *a* concentration and *Phaeocystis* cell numbers was observed (Cadée and Hegeman, 2002). These are the first signs of de-eutrophication.

The Belgian coastal zone is greatly influenced by the rivers Rhine, Meuse and the

---

**Seasonal P cycling in  
the North Sea**

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

---

**Seasonal P cycling in  
the North Sea**C. van der Zee and  
L. Chou

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2004

Scheldt (Van Bennekom and Wetsteijn, 1990; Lacroix et al., 2004). As observed generally in the North Sea, the phosphorus flux decreased more in recent years than the nitrogen flux from the Scheldt to the Belgian coastal zone. The maximal annual total nitrogen load was reported to be 34 kt in 1993 and the annual phosphate ( $\text{PO}_4$ ) load supplied by the river Scheldt to the Belgian coast was at average 1 kt between 1990 and 1993, whereas about 2 kt were supplied in the 1980s (Lenhart, 1996). Two major phytoplankton communities, the diatoms and *Phaeocystis*, dominate the nutrient removal in the Belgian coastal zone during the spring bloom. Tungaraza et al. (2003) measured the nutrient concentrations at station 330 ( $51^\circ 26' \text{N}$ ,  $2^\circ 48' \text{E}$ ) in the Belgian coastal zone between late winter and early summer in 1996 and 1997. They found that the  $\text{PO}_4$  concentrations ranged from 0.1 to 3.4  $\mu\text{M}$  and 0.3 to 1.7  $\mu\text{M}$  in 1996 and in 1997, respectively, and did not show pronounced seasonal patterns. Silicate and nitrate, on the other hand, showed clear seasonal variations with the lowest concentrations observed at the end of spring and during summer. Rousseau et al. (2002), however, did observe a seasonal pattern in the phosphate concentrations at station 330 in 1995. The  $\text{PO}_4$  concentration started to decrease between February and March with the onset of the early-spring diatom bloom. In the second half of May, the decline in the *Phaeocystis* and *Rhizosolania* spp. bloom corresponded to the lowest DIN (3.3  $\mu\text{M}$ ),  $\text{Si}(\text{OH})_4$  (0.9  $\mu\text{M}$ ) and  $\text{PO}_4$  (0.03  $\mu\text{M}$ ) concentrations recorded during spring. The lowest seasonal  $\text{pCO}_2$  values coincide with the *Phaeocystis* bloom and the lowest nutrient levels (Gypens et al., 2004).

These studies, however, focussed mainly on the role of nitrogen, silicon and carbon in the Belgian coastal zone and the only phosphorus species measured was  $\text{PO}_4$ . Little research has been devoted to the phosphorus cycle in the Belgian coastal zone, while this element is an important and potentially limiting nutrient. Also with the changing N:P ratio of the riverine fluxes, recent data are required to assess the present nutrient situation of the ecosystem. In this context, we visited during one year (from August 2002 to December 2003) 10 stations located on the Belgian Continental Shelf. Particular attention is given to the speciation of phosphorus, as in the past usually only

PO<sub>4</sub> was measured and little is known about the role of dissolved organic phosphorus compounds in eutrophied aquatic systems.

## 2. Materials and methods

### 2.1. Study site and sampling methods

5 The study area comprised the Belgian coastal zone of the southern North Sea (Fig. 1). This part of the North Sea is highly influenced by the eutrophied river Scheldt and oceanic water flowing from the Atlantic through the Channel into the North Sea. The 10 stations can be grouped along three near shore-offshore transects: one located close to the mouth of the Scheldt (stations 700, B07, 710 and 780), a second one near the city of Oostende (stations 130, 230 and 330) and a third most southern transect near Nieuwpoort (stations 120, 215 and ZG02). Surface waters were collected with Niskin bottles from 1 m water depth during almost monthly surveys with the *RV Zeeleeuw* in the Belgian coastal zone (Table 1 and Fig. 1). Samples were filtered for dissolved phosphorus speciation, nitrogen speciation and dissolved silicate determination. The particulate phase was collected on filters for the measurement of chlorophyll *a*, suspended matter, carbon and phosphorus species.

### 2.2. Chlorophyll *a* and particulate organic carbon

Chlorophyll *a* was determined on GF/F filters after filtration of 250 ml seawater under low suction, following the fluorometric method of Yentsch and Menzel (1963). Filters were immediately put in small tubes, wrapped in aluminium foil and stored at -20°C until analysis. For analysis the filters were extracted with 90% acetone at -20°C for 24 h. Samples were centrifuged (10 min, 5500 rpm) and the fluorescence of the extract was measured on a Shimadzu RF-150 fluorometer, using an excitation wavelength of 430 nm and an emission wavelength of 663 nm. The fluorescence was calibrated with a stock solution of pure chlorophyll *a* (Merck).

---

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Particulate organic carbon (POC) measurements were performed on particulate matter collected by filtration of seawater on precombusted (4 h, 500°C) GF/F filters. POC was measured using a Fisons NA-1500 elemental analyser after carbonate removal from the filters by strong acid fumes overnight. Certified reference materials were used for the calibration.

### 2.3. Dissolved Si and N species and dissolved organic carbon

Water samples for silicate were taken in plastic bottles and filtered on a plastic filtration set. Dissolved silicate was measured following the method of Grasshoff et al. (1983). Nitrate, nitrite and ammonium were determined colorimetrically. Nitrate and nitrite were analysed with a Technicon Autoanalyzer system following Grasshoff et al. (1983). Ammonium was measured with the indophenol blue technique according to Koroleff (1969). The dissolved organic nitrogen (DON) was determined as the difference between total dissolved nitrogen and the dissolved inorganic nitrogen (nitrate, nitrite and ammonium). Total dissolved nitrogen was measured by wet oxidation in alkaline persulphate (120°C, 30 min) (Grasshoff et al., 1983). The dissolved organic carbon (DOC) was determined with a Shimadzu TOC-5000 analyser using the widely accepted high-temperature catalytic oxidation (HTCO) technique (Sugimura and Suzuki, 1988; Suzuki, 1993).

### 2.4. Phosphorus speciation

Dissolved phosphorus species were determined after filtration through precombusted (4 h, 500°C) GF/F filters, which were used for the particulate phosphorus speciation. Soluble reactive phosphorus, hereafter phosphate, was measured according to the method of Grasshoff et al. (1983). Dissolved inorganic phosphorus (DIP) was determined as phosphate after digestion with 9N H<sub>2</sub>SO<sub>4</sub> (120°C, 30 min). DIP comprises phosphate and polyphosphates, although hydrolysis of acid-labile organic phosphorus compounds may also contribute to this parameter. Polyphosphate was determined

---

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

as the difference between DIP and phosphate. Total dissolved phosphorus (TDP) was measured by wet oxidation in acid persulphate (120°C, 30 min) (Grasshoff et al., 1983). Dissolved organic phosphorus was subsequently calculated as the difference between TDP and phosphate as upper estimate and the difference between TDP and DIP as lower estimate. Filters for the total particulate phosphorus (TPP) determination were combusted at 500°C for 1.5 h with MgSO<sub>4</sub> and subsequently extracted in 1N HCl for 24 h (Solorzano and Sharp, 1980). Filters for particulate inorganic phosphorus (PIP) were extracted in 1N HCl for 24 h. Particulate organic phosphorus (POP) was calculated as the difference between TPP and PIP. The unfiltered water sample was in addition digested by wet oxidation in acid persulphate, which gives the total phosphorus.

### 3. Results

#### 3.1. Chlorophyll *a*, POC and DOC

The chlorophyll *a* concentration was low from September 2002 until February 2003 after which it increased at all stations (Fig. 2). Peak concentrations in chlorophyll *a* occurred at the end of March and of April 2003. The chlorophyll *a* concentration remained subsequently at a higher level from May to September 2003 compared to the values from September 2002 to February 2003. Finally, the chlorophyll *a* concentration returned to a lower level in December 2003. No data are available between September 2003 and December 2003.

The POC content of the suspended material exhibits a seasonal trend similar to the chlorophyll *a* concentration (Fig. 2). The POC content was relatively low between September 2002 and the end of March/April when it reached its maximum, subsequently it remained relatively high until September 2003 when it returned to its winter values. The highest POC contents are observed at the stations along the Nieuwpoort transect and the lowest at the stations near the mouth of the Scheldt. The measured

---

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

POC probably does not only represent the organic carbon in pelagic material but may also contain resuspended sediment especially in periods of stormy weather. The DOC values were determined for September and November 2002 and January, March, April, May and June 2003. The DOC concentration exhibits an increasing trend from its minimum in November 2003 until April 2003, when it reaches its maximum value (Data not shown).

### 3.2. Dissolved Si and N species

The seasonal pattern in the silicate concentration showed the same trend for all stations (Fig. 3). The silicate concentration was the highest in the months of January and February and reached its lowest values in May. Two general trends can be observed. First, the maximal silicate concentrations observed in winter are higher at the stations on the Scheldt transect close to the mouth of the estuary (700, 710, 780 and B07; 35–48  $\mu\text{M Si(OH)}_4$ ), than those of the Nieuwpoort transect (120, 215, ZG02; 10–28  $\mu\text{M Si(OH)}_4$ ), the most southwestern transect. The maximal silicate values at the stations of the Oostende transect (130, 230 and 330) lie in between (17–39  $\mu\text{M Si(OH)}_4$ ). Secondly, the dissolved silicate concentration decreases going from the near-shore to the offshore stations along each transect.

Nitrate is the predominant dissolved nitrogen species of the four N species measured (Fig. 3). The nitrate plus nitrite (hereafter referred to as nitrate) concentration at all stations increased after October 2002 and remained high from December 2002 to February/March 2003 after which it reached its lowest values in May 2003. Similar to the silicate concentration, the maximal nitrate concentrations are higher at stations located close to the mouth of the Scheldt estuary as compared to those along the Nieuwpoort transect. Along each transect, a trend of decreasing nitrate concentration going from the coast seaward is also apparent. The variation in the ammonium concentration with time is less pronounced than for nitrate, but again a minimum in the concentration occurs around April–May 2003 (Fig. 3). Differences in ammonium concentration between the stations are much less distinct than for the silicate and nitrate

---

**Seasonal P cycling in  
the North Sea**

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

data. The DON concentrations are the highest in wintertime and start to decrease after March 2004 until they reach their minimum in July (Fig. 3). The DON seasonal patterns for the stations on the same transect are very similar.

### 3.3. Phosphorus species

5 Three dissolved P species were determined: phosphate ( $\text{PO}_4$ ), polyphosphates and dissolved organic phosphorus (DOP). The polyphosphate concentration was always very low, generally less than  $0.11 \mu\text{M}$ , and did not show any seasonal or spatial variation (data not shown). The seasonal pattern of the DOP concentration appears to contain two maxima: one around May 2003 and a second one at a higher concentration around August-September 2003 (Fig. 4). The DOP concentration was similar  
10 for all stations and usually low compared to the  $\text{PO}_4$  concentration, although the DOP concentration exceeded the  $\text{PO}_4$  concentration at all stations in the months of April and May 2003. The  $\text{PO}_4$  concentration at the different stations shows the decreasing trend along a transect going from the coast to the open sea (Fig. 4). However, the differences  
15 in concentration are not as pronounced as observed for the silicate and the nitrate concentrations. At the stations of the Nieuwpoort transect (120, 215 and ZG02) and station 330, the strongest decline in  $\text{PO}_4$  concentrations was observed between February and March 2003, whereas the other stations showed a strong decline between March and April 2003. The lowest values for the  $\text{PO}_4$  concentration were measured in April at the  
20 following stations: 120, 130, 700, 710, 780 and B07, whereas the lowest values at stations 215, 230 and 330 were measured in May 2003. No data are available for station ZG02 in April 2003. The  $\text{PO}_4$  concentrations increase in the fall, from August onwards for most stations, reaching the winter values again, in contrast to the seasonal patterns observed for silicate and nitrate.

25 The POP contents are high when the  $\text{PO}_4$  concentrations are low in spring (Fig. 4). The highest values of POP content are observed at the stations further away from the mouth of the Scheldt estuary in contrast to the concentrations of the dissolved nutrients phosphate, nitrate and silicate. The PIP contents at the stations of the "Scheldt" tran-

---

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

sect are very similar to those of POP as well as the seasonal trend (data not shown). However, at the stations of the other two transects the PIP content is similar to the POP content in the wintertime, but PIP content does not increase in spring as the POP content does.

## 4. Discussion

### 4.1. Spatial trends

The spatial distributions of the major nutrients in the Belgian coastal zone result from the mixing of relatively low nutrient-high salinity North Atlantic water flowing northeastward through the Channel into the North Sea with high nutrient freshwater which is discharged from the rivers Rhine, Meuse and Scheldt. The freshwater flows clockwise southwestward along the Dutch and Belgian coast before it mixes with the northeastward flowing Channel water (Lacroix et al., 2004). This water circulation pattern is reflected in the winter nutrient concentrations at the different stations: high nutrient concentrations at the stations on the Scheldt transect as well as the coastal stations 120 and 130, which are located in the Rhine/Meuse and Scheldt plumes and lower nutrient concentrations at the other stations. Basically, it gives a wintertime nutrient gradient in the nearshore-offshore direction and in the northeast-southwest direction.

The lack of a spatial trend in the concentrations of both the DOP and DON at any time suggests that they are produced at sea in the Belgian coastal zone in contrast to the phosphate and nitrate where a concentration gradient going from the northeast towards the southwest indicates the strong influence of the rivers Rhine/Meuse and Scheldt on their winter concentrations. This is also reflected in the DOP and PO<sub>4</sub> concentration versus salinity plots (Fig. 5) where the wintertime PO<sub>4</sub> concentration is dominated by conservative mixing, in contrast to the wintertime DOP concentration.

The intensity of the spring bloom, as reflected by the chlorophyll *a* data (Fig. 2), seems to be evenly distributed over the Belgian coastal zone. There are two pro-

---

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

nounced peak values, namely the peak values of station 230 and station B07 at the end of April 2003. General trends as seen in the nutrient distribution, however, are not apparent for chlorophyll *a*. The spatial distribution of the particulate organic carbon and phosphorus contents indicates an increasing trend going from the northeast to the southwest (Figs. 2 and 4). This trend is caused by the dilution of organic-rich pelagic material with relatively organic-poor resuspended sediment at the more turbid stations near the mouth of the Scheldt estuary. It is further supported by the higher suspended matter concentration at the stations on the Scheldt transect and the similar POP concentration at all stations, when expressed in  $\mu\text{mol P/l}$  seawater (not shown).

#### 4.2. Seasonal trends in phosphate concentration

Besides the minimum in  $\text{PO}_4$  concentration in spring, another minimum is observed in October 2002, which is not accompanied by an increase in POP content or chlorophyll *a* concentration. This minimum is caused by the relative higher contribution of high salinity-low nutrient water from the Channel. Phosphate concentration versus salinity plots for the different months indicate a more or less conservative behaviour in the winter months (October 2002, November 2002, December 2002, January 2003 and December 2003), whereas the biological activity determines the phosphate concentration in spring-summer and summer-fall (Fig. 5).

The largest decrease in the major nutrient concentrations occurs between the end of February and the end of April 2003 coinciding with the largest increase in chlorophyll *a* concentration and particulate organic carbon content (Figs. 2–4). For the northeastern stations (700, 710, 780, B07, 130 and 230) the largest decline in nutrients occurs between the end of March and the end of April 2003, whereas for the southwestern stations the major decrease occurs one month earlier. At the end of the summer the dissolved Si,  $\text{PO}_4$  and DIN concentrations increase again when their supply exceeds their consumption. In contrast to the dissolved Si and DIN, the  $\text{PO}_4$  concentrations reach their pre-bloom levels again already in the fall. This difference in the seasonal evolution of P versus N and Si was also noted in a compilation of nutrient data from

---

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

1993–2000 in the same area (De Galan et al., in press). Thus, the rate of regeneration and/or supply of phosphorus is higher than that for dissolved Si and DIN in the Belgian coastal zone. One source of these nutrients is river run-off from the Rhine/Meuse and Scheldt. It is not likely that river run-off would deliver more phosphate than nitrate in fall, because nitrate concentrations tend to increase as river flow increases, whereas phosphate concentrations tend to decrease (Balls, 1992; 1994). Another source for nitrogen is atmospheric input, but for phosphorus this is negligible in the North Sea (Brion et al., 2004). Both atmospheric and riverine input would deliver more nitrogen than phosphorus and are therefore not likely to cause the earlier replenishment of phosphate as compared to nitrate in the Belgian coastal zone. In the southern bight of the North Sea, the oceanic/riverine inflow rates of phosphate, nitrate and silicate are insufficient to support their seasonal cycle and therefore internal recycling is required (Prandle et al., 1997). Since the supply of phosphate to the Belgian coastal zone is not higher than that of silicate and nitrate, the difference can be attributed to the internal cycling. Preferential regeneration of P over N and C has already been demonstrated both in the hydrolysis of dissolved organic matter (e.g. Hopkinson et al., 1997; Clark et al., 1998; Vidal et al., 1999) and in the remineralisation of particulate organic matter (Bishop et al., 1977; Garber, 1984; Martin et al., 1987).

#### 4.3. Phosphate being the first limiting nutrient

In estuaries and plume zones, turbidity can cause light limitation of primary production, while nutrient levels remain high. In the Thames plume for example, minimal concentrations of nitrate and phosphate were ca.  $7 \mu\text{M}$  and  $1 \mu\text{M}$ , respectively, but silicate concentrations were strongly depleted in summer and spring, presumably by diatom growth (Sanders et al., 2001). The authors hypothesized that the limited development of non-siliceous organisms, in spite of high residual levels of nitrate and phosphate, was due to a more intense light limitation of non-siliceous compared to diatom growth. Primary production was also strongly regulated by irradiance and nutrients in the Wadden Sea (Colijn and Cadée, 2003). Model results from the Humber estuary indicate that in

---

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

the plume area the primary production is controlled by light limitation between October and March and by nutrient availability during the rest of the year (Allen, 1997). Typical nitrate and phosphate half saturation constants for coastal phytoplankton populations are ca. 1–2  $\mu\text{M}$  nitrate and 0.1–0.5  $\mu\text{M}$  phosphate and typical half saturation constants for coastal diatom populations are ca. 1–2  $\mu\text{M}$  silicate (Fisher et al., 1988). In our study area, the lowest nutrient concentrations recorded for the individual stations range from <0.01 to 0.13  $\mu\text{M}$   $\text{PO}_4$ , from 0.27 to 1.53  $\mu\text{M}$   $\text{Si}(\text{OH})_4$  and from 0.37 to 5.48  $\mu\text{M}$   $\text{NO}_3$ . The first nutrient to drop below its half saturation constant is phosphate, then silicate and later nitrate, although not at all stations. The consumption of nutrients resulting in such low levels indicates that it is the nutrient that limits the primary production at the time of the phytoplankton bloom and not the light intensity.

We have compiled a table in attempt to give an indication of what the limiting nutrient for phytoplankton growth can be (Table 2). The concentrations of  $\Sigma\text{DIN}$ , phosphate and silicate are given in the month when they are the lowest for the individual stations. For some stations, all three nutrients reach their lowest levels in the same month; then only the data of that month are listed and the N:P, Si:P and Si:N ratios are given. When N:P and Si:P are larger than 16, phosphate is likely to be the limiting nutrient. In the case where  $\text{Si:P} < 16$  and  $\text{Si:N} < 1$ , then silicate is likely to be the limiting nutrient (for diatom growth). When  $\text{N:P} < 16$  and  $\text{Si:N} > 1$ , nitrogen would be the limiting element. Note that for the four stations near the mouth of the estuary (700, 710, 780 and B07) and station 130, the phytoplankton is first P limited but then becomes Si limited. The stations that are least influenced by the river plumes (ZG02 and 330) are likely to be P limited for phytoplankton growth. Phosphorus limitation of primary production during the early spring bloom was also observed in the Gironde plume (Herbland et al., 1998; Labry et al., 2002), whereas the winter bloom was initiated by light availability (Labry et al., 2001). At no time is the phytoplankton community limited by inorganic nitrogen in the Belgian coastal zone. Also when the N:P ratios are calculated using total dissolved N and P concentrations, e.g. including dissolved organic nitrogen and phosphorus, P is limiting. This is because the DON concentration is more than 16 times that of

**Seasonal P cycling in  
the North Sea**C. van der Zee and  
L. Chou

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

DOP in the months considered. DON always constitutes a large fraction of the total dissolved nitrogen, in contrast to DOP that becomes an important fraction only when the phosphate levels are low. DON representing a major fraction of the total dissolved nitrogen was also reported by De Galan et al. (2004), but a seasonal trend in the DON concentrations was not observed, possibly due to a too low sampling frequency.

#### 4.4. DOP and phosphorus cycling

DOP is an important source of biologically available phosphorus for both bacteria and phytoplankton in marine ecosystems, when the  $\text{PO}_4$  concentration is low (Orrett and Karl, 1987; Björkman and Karl, 1994). DOP is not readily available to phytoplankton and bacteria, but it can be utilized as phosphorus source by means of hydrolytic enzymes: the non-specific alkaline phosphatase (Perry, 1972) and 5'-nucleotidase (Ammerman and Azam, 1985; Flynn et al., 1986). Alkaline phosphatase is induced by low  $\text{PO}_4$  concentrations and cleaves the  $-\text{PO}_4$  moiety from organic phosphorus compounds after which the cell can assimilate the  $\text{PO}_4$ . Van Boekel and Veldhuis (1990) have shown that alkaline phosphatase synthesis in *Phaeocystis* sp. is controlled by the external phosphate concentration and that the threshold concentration for derepression is ca  $0.5 \mu\text{M}$ . Unlike alkaline phosphatase, 5'-nucleotidase recognizes the carbon moiety of nucleotides and is not inhibited at high  $\text{PO}_4$  levels.

The wintertime DOP concentrations in the Belgian coastal zone (Fig. 5) are comparable to those reported for the oligotrophic North Pacific Subtropical Gyre ( $\sim 0.2 \mu\text{M}$ ; Church et al., 2002), the oligotrophic central Atlantic Ocean ( $0.1\text{--}0.3 \mu\text{M}$ ; Vidal et al., 1999) and the northeastern continental shelf of the USA ( $\sim 0.2 \mu\text{M}$ ; Hopkinson et al., 1997). The elevated DOP concentrations in the Belgian coastal zone are of similar magnitude to those reported for surface waters along the north coast of Australia (Mulholland et al., 2002) and the Southern California Bight (Ammerman and Azam, 1991). DOP concentrations of  $\sim 0.2 \mu\text{M}$  are reported for the Baltic Sea, but higher DOP values of  $0.35\text{--}0.75 \mu\text{M}$  were measured in cyanobacterial blooms (Nausch et al., 2004).

When the  $\text{PO}_4$  concentration is low, a significant fraction of the dissolved P is present

---

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

in the form of DOP. Between March and July, when the  $\text{PO}_4$  concentration is at its lowest level (Fig. 4), phytoplankton and bacteria may have to turn to DOP as their phosphorus source. In general, it is assumed that elevated dissolved organic matter (DOM) concentrations in surface waters result from enhanced DOM production associated with primary production (Duursma, 1961). This may be due to DOM excretion from cells (Lancelot, 1979), viral mediated cell lysis (Furhman and Shuttle, 1993), sloppy feeding by zooplankton (Eppley et al., 1981) or from particle solubilization (Smith et al., 1992). The seasonal trend in DOP shows 2 periods of elevated concentrations (Fig. 4). The first period is in April for the most southwestern stations and in May for the most northeastern stations. This period of elevated DOP concentration coincides with the lowest  $\text{PO}_4$  concentration and the highest chlorophyll *a* concentration and POP content, and therefore corresponds most likely to the period of elevated primary production. The second period of elevated DOP concentration occurs in August-September when the POP content is decreasing strongly and the  $\text{PO}_4$  concentration rises again. In this period, the organic phosphorus contained in cells is released upon their lysis and degraded, and the DOP is presumably further remineralised into phosphate. Turk et al. (1992) have demonstrated that bacterial biomass grazed by protozoa plays an important role in the cycling of P, due to the ejection of the consumed bacterial DNA and the subsequent rapid DNA degradation and liberation of  $\text{PO}_4$ . Thus, the elevated DOP concentrations indicate two distinct periods of intensive P cycling. During the first period P is regenerated and sustains the bloom, and during the second period P is recycled as biomass is decaying.

## 5. Conclusions

Nutrient and P speciation data obtained by monthly cruises during one complete year were presented for 10 stations in Belgian coastal zone. Spatial and seasonal trends were studied. The rivers Rhine/Meuse and Scheldt have a strong influence on the distribution of inorganic nutrients, but not on that of the DOP and DON.

**BGD**

1, 681–707, 2004

---

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

---

**Seasonal P cycling in  
the North Sea**

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

At most stations,  $\text{PO}_4$  is the first nutrient to decrease below its half-saturation value during the bloom period, thereafter  $\text{Si}(\text{OH})_4$  and subsequently  $\text{NO}_3$ . The low nutrient levels indicate that phytoplankton growth is limited by the nutrient and not by the light availability during the bloom. The elemental ratios of N, P and Si suggest that either P or Si limitation of phytoplankton growth could develop, and for the stations on the Scheldt transect first P and later Si limitation. The rate of regeneration of  $\text{PO}_4$  is much faster than that of  $\text{NO}_3$  and  $\text{Si}(\text{OH})_4$  so that  $\text{PO}_4$  reaches again its wintertime concentrations often in August/September already.

The DOP concentration is elevated at the time of the lowest  $\text{PO}_4$  concentration and highest primary production as is also frequently observed for the DOC concentration, but not for DON at our stations. A second period of elevated DOP concentration occurs during the decline of the bloom when POP is degraded and  $\text{PO}_4$  levels are restored. Thus, elevated DOP concentrations are indicative for intensive internal cycling of P. Phosphorus regeneration is an important process in the southern bight of the North Sea, which allows phytoplankton growth even at very low  $\text{PO}_4$  levels.

*Acknowledgements.* The authors wish to thank the captain and crew of the “RV Zeeleeuw”, A. Cattrijsse and F. Hernandez (VLIZ), as well as N. Roevros (ULB-LOGCE) for their logistic support. We are grateful to M. Tsagaris and C. de Marneffe (ULB-LOGCE) for their analytical work in the laboratory and to our ULB-LOGCE colleagues for their assistance at sea. This study was supported by the Belgian Federal Science Policy Office under contract no. EV/11/20B (CANOPY). Partial funding provided by the SISCO project (contract no. EV/11/17A) is also acknowledged.

## References

- Ammerman, J. W. and Azam, F.: Bacterial 5'-nucleotidase in aquatic ecosystems: A novel mechanism for phosphorus regeneration, *Science*, 227, 1338–1340, 1985.
- Ammerman, J. W. and Azam, F.: Bacterial 5'-nucleotidase activity in estuarine and coastal marine waters: Role in phosphorus regeneration, *Limnol. Oceanogr.*, 36, 1437–1447, 1991.

---

**Seasonal P cycling in  
the North Sea**C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

- Balls, P. W.: Nutrient behaviour in two contrasting Scottish estuaries, the Forth and Tay, *Oceanologica Acta*, 15, 261–277, 1992
- Balls, P. W.: Nutrient inputs to estuaries from nine Scottish east coast rivers; influence of estuarine processes on inputs to the North Sea, *Est. Coast. Shelf Sci.*, 39, 329–352, 1994.
- 5 Behrendt, H., Huber, P., Kornmilch, M., Opitz, D., Schmoll, O., Scholz, G., and Uebe, R.: Nutrient emissions into river basins of Germany, UBA-Texte, 23/00, Berlin, 261, 2002.
- Bishop, J. K. B., Collier, R. W., Ketten, D. R., and Edmond, J. M.: The chemistry, biology and vertical flux of particulate matter from the upper 400 m of the equatorial Atlantic ocean, *Deep Sea Res.*, 24, 511–548, 1977.
- 10 Björkman, K. and Karl, D. M.: Bioavailability of inorganic and organic phosphorus compounds to natural assemblages of microorganisms in Hawaiian coastal waters, *Mar. Ecol. Progres. Ser.*, 111, 265–273, 1994.
- Brion, N., Baeyens, W., De Galan, S., Elskens, M., and Laane, R. W. P. M.: The North Sea: source or sink for nitrogen and phosphorus to the Atlantic Ocean?, *Biogeochem.*, 68, 277–295, 2004.
- 15 Church, J. J., Ducklow, H. W., and Karl, D. M.: Multiyear increases in dissolved organic matter inventories at station ALOHA in the North Pacific Subtropical Gyre, *Limnol. Oceanogr.*, 47, 1–10, 2002.
- Clark, L. L., Ingall, E. D., and Benner, R.: Marine phosphorus is selectively remineralized, *Nature*, 393, 426, 1998.
- 20 Duursma, E.: Dissolved organic carbon, nitrogen and phosphorus in the sea, *Neth. J. Sea Res.*, 1, 1–148, 1961.
- De Galan, S., Elskens, M., Goeyens, L., Pollentier, A., Brion, N., and Baeyens, W.: Spatial and temporal trends in nutrient concentrations in the Belgian continental area of the North Sea during the period 1993–2000, *Est. Coast. Shelf Sci.*, in press, 2004.
- 25 Eppley, R., Horrigan, S., Fuhrman, J., Brooks, E., Price, C., and Sellner, K.: Origins of dissolved organic mater in Southern California coastal waters: experiments on the role of zooplankton, *Mar. Ecol. Progres. Ser.*, 6, 149–159, 1981.
- Fisher, T. R., Harding, L. W., Stanley, D. W., and Ward, L. G.: Phytoplankton, nutrients and turbidity in the Chesapeake, Delaware and Hudson estuaries, *Est. Coast. Shelf Sci.*, 27, 61–93, 1988.
- 30 Flynn, K. J., Opik, H., and Syrett, P. J.: Localization of the alkaline phosphatase and 5'-nucleotidase activities of the diatom *Phaeodactylum tricornutum*, *J. Gen. Microbiol.*, 132,

289–298, 1986.

Furhman, J. and Shuttle, C.: Viruses in marine planktonic systems, *Oceanography*, 6, 51–63, 1993.

Garber, J.: Laboratory study of nitrogen and phosphorus mineralization during the decomposition of coastal plankton and seston, *Est. Coast. Shelf Sci.*, 18, 685–702, 1984.

Grasshoff, K., Ehrhardt, M., Kremling, K. (eds.): *Methods of Seawater Analysis*, Verlag Chemie, Basel, 1983.

Gypens, N., Lancelot, C., and Borges, A. V.: Carbon dynamics and CO<sub>2</sub> air-sea exchanges in the eutrophied coastal waters of the southern bight of the North Sea: a modeling study, *Biogeosciences Discussions*, 1, 561–589, 2004,

SRef-ID: 1810-6285/bgd/2004-1-561.

Herbland, A., Delmas, D., Laborde, P., Sautour, B., and Artigas, F.: Phytoplankton spring bloom of the Gironde plume waters in the Bay of Biscay: early phosphorus limitation and food-web consequences, *Oceanol. Acta*, 21, 279–291, 1998.

Hopkinson Jr., C. S., Fry, B., and Nolin, A. L.: Stoichiometry of dissolved organic matter dynamics on the continental shelf of the northeastern USA, *Cont. Shelf Res.*, 17, 473–489, 1997.

Koroleff, F.: Direct determination of ammonia in natural waters as indophenol blue, *Int. Council Expl. Sea*, 9, 19–22, 1969.

Lacroix, G., Ruddick, K., Ozer, J., and Lancelot, C.: Modelling the impact of the Scheldt and Rhine/Meuse plumes on the salinity distribution in Belgian waters (southern North Sea), *J. Sea Res.*, 52, 149–163, 2004.

Labry, C., Herbland, A., Delmas, D., Laborde, P., Lazure, P., Froidefond, J. M., Jegou, A. M., and Sautour, B.: Initiation of winter phytoplankton blooms within the Gironde plume waters in the Bay of Biscay, *Mar. Ecol. Progres. Ser.*, 212, 117–130, 2001.

Labry, C., Herbland, A., and Delmas, D.: The role of phosphorus on planktonic production of the Gironde plume waters in the Bay of Biscay, *J. Plankt. Res.*, 24, 97–117, 2001.

Lancelot, C.: Gross excretion rate of natural marine phytoplankton and heterotrophic uptake of excreted products in the southern North Sea, as determined by short-term kinetics, *Mar. Ecol. Progres. Ser.*, 1, 179–186, 1979.

Lancelot, C., Billen, G., Sournia, A., Weisse, T., Colijn, F., Veldhuis, M. J. W., Davies, A., and Wassman, P.: *Phaeocystis* blooms and nutrient enrichment in the continental coastal zones of the North Sea, *Ambio*, 16, 38–46, 1987.

**BGD**

1, 681–707, 2004

---

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGU 2004

---

**Seasonal P cycling in  
the North Sea**C. van der Zee and  
L. Chou

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGU 2004

- Lenhart, H. J., Pätsch, J., and Radach, G.: Daily nutrient loads of the European continental rivers for the years 1977–1993, Institut für Meereskunde Bibliothek, Hamburg, Germany, 1996.
- Martin, J., Knauer, G., Karl, D., and Broenkow, W.: VERTEX: carbon cycling in the northeast Pacific, *Deep-Sea Res.*, 34, 257–285, 1987.
- Mulholland, M. R., Flöge, S., Carpenter, E. J., and Capone, D. G.: Phosphorus dynamics in cultures and natural populations of *Trichodesmium spp.*, *Mar. Ecol. Progress. Ser.*, 239, 45–55, 2002.
- Nausch, M., Nausch, G., and Wasmund, N.: Phosphorus dynamics during the transition from nitrogen to phosphate limitation in the central Baltic Sea, *Mar. Ecol. Progress. Ser.*, 266, 15–25, 2004.
- North Sea Task Force: North Sea Quality Status Report (QSR), Oslo and Paris Commissions, London and International Council for the Exploration of the Sea, Copenhagen, 1993.
- Orrett, K. and Karl, D. M.: Dissolved organic phosphorus production in surface waters, *Limnol. Oceanogr.*, 32, 383–395, 1987.
- Perry, M. J.: Alkaline phosphatase activity in subtropical Central North Pacific waters using a sensitive fluorometric method, *Mar. Biol.*, 15, 113–119, 1972.
- Philippart, C. J. M., Cadée, G. C., van Raaphorst, W., and Riegman, R.: Long-term phytoplankton-nutrient interactions in a shallow coastal sea: Algal community structure, nutrient budgets, and denitrification potential, *Limnol. Oceanogr.*, 45, 131–144, 2000.
- Prandle, D., Hydes, D. J., Jarvis, J., and McManus, J.: The seasonal cycles of temperature, salinity, nutrients and suspended sediment in the southern North Sea in 1988 and 1989, *Est. Coast. Shelf Sci.*, 45, 669–680, 1997.
- Rousseau, V., Leynaert, A., Daoud, N., and Lancelot, C.: Diatom succession, silicification and silicic acid availability in Belgian coastal waters (southern North Sea), *Mar. Ecol. Progress. Ser.*, 236, 61–73, 2002.
- Sanders, R., Jickells, T., and Mills, D.: Nutrients and chlorophyll at two sites in the Thames plume and southern North Sea, *J. Sea Res.*, 46, 13–28, 2001.
- Skogen, M. D., Søiland, H., and Svendsen, E.: Effects of changing nutrient loads to the North Sea, *J. Mar. Sys.*, 46, 23–38, 2004.
- Sugimura, Y. and Suzuki, Y.: A high-temperature catalytic oxidation method for the determination of non-volatile dissolved organic carbon in seawater by direct injection of a liquid sample, *Mar. Chem.*, 24, 105–131, 1988.

- Suzuki, Y.: On the measurement of DOC and DON in seawater, *Mar. Chem.*, 41, 287–288, 1993.
- Smith, D., Simon, M., Alldredge, A., and Azam, F.: Intense hydrolytic enzyme activity on marine aggregates and implications for rapid particle dissolution, *Nature*, 359, 129–142, 1992.
- 5 Tungaraza, C., Rousseau, V., Brion, N., Lancelot, C., Gichuki, J., Baeyens, W., and Goeyens, L.: Contrasting nitrogen uptake by diatom and *Phaeocystis*-dominated phytoplankton assemblages in the North Sea, *J. Exp. Mar. Biol. Ecol.*, 292, 19–41, 2003.
- Turk, V., Rehnstam, A., Lundberg, E., and Hagstrom, A.: Release of bacterial DNA by marine nanoflagellates, an intermediate step in phosphorus regeneration, *Appl. Environ. Microbiol.*, 10 58, 3744–3750, 1992.
- Turner, R. E., Rabalais, N. N., Justic, D., and Dortch, Q.: Future aquatic nutrient limitations, *Mar. Poll. Bul.*, 46, 1032–1034, 2003a.
- Turner, R. E., Rabalais, N. N., Justic, D., and Dortch, Q.: Global patterns of dissolved N, P and Si in large rivers, *Biogeochem.*, 64, 297–317, 2003b.
- 15 Van Bennekom, A. J. and Wetsteijn, F. J.: The winter distribution of nutrients in the southern bight of the North Sea (1961–1978) and in the estuaries of the Scheldt and the Rhine/Meuse, *Neth. J. Sea Res.*, 25, 75–87, 1990.
- Vidal, M., Duarte, C. M., and Agusti, S.: Dissolved organic nitrogen and phosphorus pools and fluxes in the central Atlantic Ocean, *Limnol. Oceanogr.*, 44, 106–115, 1999.
- 20 Yentsch, C. S. and Menzel, D. W.: A method for the determination of phytoplankton chlorophyll and phaeophytin by fluorescence, *Deep-Sea Res.*, 10, 221–231, 1963.

---

**Seasonal P cycling in  
the North Sea**

C. van der Zee and  
L. Chou

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

**Seasonal P cycling in the North Sea**

C. van der Zee and  
L. Chou

**Table 1.** Sampling locations and dates.

Station name	Latitude	Longitude	Sampling dates
Nieuwpoort Transect			23 August 2003
120	51°11.1	2°42.0	27 September 2003
215	51°16.5	2°36.8	29 October 2003
ZG02	51°20.0	2°30.0	21 November 2003 23 December 2003
Oostende Transect			20 January 2004
130	51°16.2	2°54.2	26 February 2004
230	51°18.4	2°51.0	31 March 2004
330	51°26.0	2°48.5	25 April 2004 26 May 2004
Scheldt Transect			30 June 2004
700	51°22.6	3°13.2	31 July 2004
710	51°26.5	3°08.3	28 August 2004
780	51°28.3	3°03.5	26 September 2004
B07	51°25.9	3°17.9	9 December 2004

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Seasonal P cycling in the North Sea**

C. van der Zee and  
L. Chou

**Table 2.** Lowest nutrient concentrations observed, N:P, Si:P and Si:N ratios and possible limiting nutrient for each station.

Station	Date	$\Sigma$ DIN ( $\mu$ M)	DIP ( $\mu$ M)	Si ( $\mu$ M)	N:P ratio	Si:P ratio	Si:N ratio	Limiting nutrient
120	April 2003	14.05	<b>0.07</b>	0.92	211	14	0.1	Si
	May 2003	<b>0.78</b>	0.19	<b>0.41</b>	4	2	0.5	Si
215	May 2003	0.66	<0.01	0.36	>66	>36	0.5	P
ZG02	May 2003	0.73	0.02	0.36	35	17	0.5	P
130	April 2003	16.87	<b>0.07</b>	3.63	253	54	0.2	P
	May 2003	<b>0.60</b>	0.10	<b>0.46</b>	6	5	0.8	Si
230	May 2003	0.37	0.04	0.36	9	9	1.0	?
330	May 2003	1.31	0.01	0.27	116	24	0.2	P
700	April 2003	<b>5.48</b>	<b>0.10</b>	3.72	53	36	0.7	P
	June 2003	16.37	0.57	<b>1.53</b>	29	3	0.1	Si
710	April 2003	1.51	<b>0.09</b>	2.27	18	27	1.5	P
	May 2003	1.15	0.13	0.55	9	4	0.5	Si
780	April 2003	3.76	<b>0.08</b>	1.95	50	26	0.5	P
	May 2003	<b>0.92</b>	0.08	<b>0.36</b>	11	4	0.4	Si
B07	April 2003	6.85	0.13	2.79	52	21	0.4	P
	June 2003	<b>3.30</b>	0.23	<b>1.06</b>	14	5	0.3	Si

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Print Version

Interactive Discussion

**Seasonal P cycling in the North Sea**

C. van der Zee and  
L. Chou



**Fig. 1.** Map of the Belgian coastal zone indicating the location of the ten stations.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Print Version

Interactive Discussion

Seasonal P cycling in the North Sea

C. van der Zee and L. Chou



**Fig. 2.** Seasonal trend in the chlorophyll *a* concentration and in the particulate organic carbon content for the ten stations investigated. The stations are grouped per transect: on the Nieuwpoort transect stations 120, 215 and ZG02, on the Oostende transect stations 130, 230 and 330 and on the Scheldt transect stations B07, 700, 710 and 780.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Print Version

Interactive Discussion

Seasonal P cycling in the North Sea

C. van der Zee and L. Chou



**Fig. 3.** The seasonal trend in the concentrations of silicate, nitrate, ammonium and dissolved organic nitrogen (DON) for the ten stations.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close


Full Screen / Esc

Print Version

Interactive Discussion

## Seasonal P cycling in the North Sea

C. van der Zee and  
L. Chou



**Fig. 4.** Seasonal trends in the  $PO_4$  and DOP concentrations and in the POP content for the Nieuwpoort transect (120, 215 and ZG02), Oostende transect (130, 230 and 330) and Scheldt transect (B07, 700, 710 and 780).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Print Version

Interactive Discussion

© EGU 2004

Seasonal P cycling in the North Sea

C. van der Zee and L. Chou



**Fig. 5.** The PO<sub>4</sub> and DOP concentrations versus salinity, data of all the stations in different months. For the month of May, no salinity data were available. DOP was not determined for the months of August and September 2002.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion