

Controls over N2O, NOx and CO2 fluxes in a calcareous mountain forest soil

B. Kitzler, S. Zechmeister-Boltenstern, C. Holtermann, U. Skiba, K.

Butterbach-Bahl

► To cite this version:

B. Kitzler, S. Zechmeister-Boltenstern, C. Holtermann, U. Skiba, K. Butterbach-Bahl. Controls over N2O, NOx and CO2 fluxes in a calcareous mountain forest soil. Biogeosciences Discussions, 2005, 2 (5), pp.1423-1455. hal-00297816

HAL Id: hal-00297816 https://hal.science/hal-00297816

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Biogeosciences Discussions, 2, 1423–1455, 2005 www.biogeosciences.net/bgd/2/1423/ SRef-ID: 1810-6285/bgd/2005-2-1423 European Geosciences Union



Biogeosciences Discussions is the access reviewed discussion forum of Biogeosciences

Controls over N_2O , NO_x and CO_2 fluxes in a calcareous mountain forest soil

B. Kitzler¹, S. Zechmeister-Boltenstern¹, C. Holtermann², U. Skiba³, and K. Butterbach-Bahl⁴

¹Federal Research and Training Centre for Forests, Natural Hazards and Landscape (BFW), Seckendorff-Gudent-Weg 8, Vienna, Austria

²Sellenyg. 2–4/52, Vienna, Austria

³Institute of Terrestrial Ecology, Bush Estate, Penicuik, Midlothian EH26 OQB, Scotland ⁴Institute for Meteorology and Climate Research, Atmospheric Environmental Research, Forschungszentrum Karlsruhe, Kreuzeckbahnstraße 19, 82467, Garmisch-Partenkirchen, Germany

Received: 1 July 2005 - Accepted: 11 August 2005 - Published: 9 September 2005

Correspondence to: B. Kitzler (barbara.kitzler@bfw.gv.at)

© 2005 Author(s). This work is licensed under a Creative Commons License.

BGD

2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



Abstract

We measured nitrogen oxides (N₂O and NO_x), dinitrogen (N₂) and carbon dioxide (CO₂) emissions from a spruce-fir-beech forest soil in the North Tyrolean limestone Alps in Austria. The site received 12.1 kg nitrogen via wet and dry deposition. Fluxes of nitric oxide (NO) were measured by an automatic dynamic chamber system on an hourly basis over a two year period. Daily N₂O emissions were obtained by a semi-automatic gas measuring system. In order to cover spatial variability biweekly manual measurements of N₂O and CO₂ emissions were carried out, additionally. For acquiring information on the effects of soil and meteorological conditions and of N-deposition

- on N-emissions we chose the autoregression procedure (time-series analysis) as our means of investigation. Hence, we could exclude the data's autocorrelation in the course of the time. We found that soil temperature, soil moisture and wet N-deposition followed by air temperature and precipitation were the most powerful influencing parameters effecting N-emissions. With these variables up to 89% of observed temporal
- ¹⁵ variations of N-emissions could be explained. During the two-year investigation period between 2.5 and 3.5% of deposited N was reemitted in form of N₂O whereas only 0.2% were emitted as NO. At our mountain forest site the main end-product of microbial activity processes was N₂ and trace gases (N₂O and NO) were only of minor importance.

20 1. Introduction

25

Nitrous oxide (N_2O) is a stable greenhouse gas in the troposphere and is involved directly in global warming. In the stratosphere it is responsible for the catalytic decomposition of ozone (Crutzen, 1979). The increase in the atmosphere is reported at present to be about 0.25% per year (IPCC, 2001). As the potential of global warming of N_2O is 300 times higher than that of carbon dioxide (CO_2) a further increase of atmospheric N_2O concentration might affect global climate even more in the future.

BGD

2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



In contrast, nitric oxide (NO) is indirectly involved in global warming and contributes to the net production of radiative tropospheric ozone and photochemical air pollution (Crutzen, 1995; Firestone and Davidson, 1989; Williams et al., 1992).

The production of N₂O, NO, and N₂ in soils is mainly the result of nitrifying and denitrifying bacteria (e.g. Davidson et al., 2000; Conrad, 1996; Lovett et al., 2002). These microbial activity processes – and thus the production rates of NO, N₂O and N₂ are influenced by various factors. These drivers are temperature, precipitation, total N, availability of O₂, mineralization rate, respiration, texture and structure, available nitrogen, pH and tree species composition (Davidson et al., 2000; Ludwig et al., 2001; Simek and Cooper, 2002). As forest soils are generally not treated with N fertilizers, N deposition from the atmosphere which are mainly due to NH₃ release form agricultural practices and NO_x release from combustion processes provides a considerable increase in mineral N input to forest soils. Atmospheric N-deposition is estimated to be

- at least 5 kg N ha⁻¹ y⁻¹ in most regions of Europe, but can reach up to 50 kg N ha⁻¹ y⁻¹ in central and northern Europe (NADP, 2002). Nitrogen surplus in forest soil can be stored as soil N, taken up by microbes and plants or leached into groundwater. It can alter the rates of microbial N- and C- turnover in soil or it can be nitrified and denitrified to gaseous products of NO, N₂O and N₂ (Skiba et al., 2004). Increased N-deposition can lead to changes in soil chemistry, forest composition and forest productivity.
- ²⁰ A deposition of 12 kg ha⁻¹ y⁻¹ of nitrogen at the study site Achenkirch (AK) is moderate but exceeds Critical Loads according to the WHO-Guideline (WHO, 1995) for sensitive coniferous forest ecosystems (Smidt et al., 1996). In other alpine areas an input of up to 30 kg ha⁻¹ y⁻¹ was found (Herman et al., 2002); and increased flux rates of atmospheric greenhouse gas emissions (NO, N₂O, CO₂, CH₄) might be expected.
- ²⁵ However, only limited information is available about effects of atmospheric N deposition on the biosphere-atmosphere exchange of N trace gases in alpine forest ecosystems. During a two-year sampling period of CO₂, N₂O and NO_x emissions, measurements were carried out in order (1) to investigate the trace gas exchange between a typical spruce-beech-fir forest soil in the limestone Alps and the atmosphere. Measurements

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



were conducted in high temporal and spatial resolution (2) to get better estimates of annual emissions, (3) to study the relationships between N-trace gas emissions and meteorological factors and soil parameters, (4) to investigate if N-deposition affects N-emissions in these forest ecosystems and (5) to find an appropriate statistical pro-5 cedure to describe the relationships between N-emissions and their ecological drivers.

Material and methods 2.

2.1. Investigation site and soil

The 127 year old forest is located in the Achental of the North Tyrolean limestone Alps, and was described by Englisch and Starlinger (1995) as a warm, central-montane spruce-fir-beech forest. The Mühleggerköpfl is a completely isolated hill, formed from 10 dolomite, at a height of 895 m above sea level. The soils are mainly Rendzic Leptosols and Chromic Cambisol and are characterized by high clay content and a low percentage of coarse material in the upper 10 cm. The soil properties are characterized by a high spatial heterogeneity. The depth of the A-horizon varies from 10-35 cm. For this horizon the organic matter content is very high (>10%). Mean pH is 6.42. For 15 site description see Table 1. A detailed description of the site is given in Herman et al. (2002).

2.2. N₂O and CO₂ flux measurements

Gas measurements were carried out as described in Kitzler et al. (2005). Gas samples of four manually operated chambers (area: 1 m², volume: 801) were taken biweekly 20 from April 2002 until July 2004. Duplicate air samples from the chambers were taken after 0, 1 and 2 h and injected into gas tight headspace vials (20 ml). Daily (1/day) samples were taken by the automatic gas sampling system (AGPS - patent DE 198 52 859) and vials were collected every two weeks. The system is described in detail

Close

Back

Full Screen / Esc

Print Version

Interactive Discussion

EGU

by Kitzler et al. (2005) and has been shown to allow realistic N₂O emission estimates for forest soils (Zechmeister-Boltenstern et al., 2002). Gas samples were stored at 4°C until analysis. The time of automatic sampling was scheduled for 6 a.m. due to the fact that in a previous experiment highest emissions had been measured at morning dew.

⁵ In order to avoid freezing of the covering case on the sealing plate, the thermostat was set to 1°C and there were no measurements conducted below this temperature. During winter no automatic measurements were carried out, but manual measurements were carried out as long as the site was accessible.

Emissions of N_2O (μ g N m⁻² h⁻¹) and CO₂ (mg C m⁻² h⁻¹) were determined by the linear increase of the mixing ratio within the incubation period. Nitrous oxide and CO₂ fluxes were calculated as described in (IAEA, 1992).

A gas chromatograph (HP 5890 Series II) with a 63 Ni-electron-capture detector (ECD), connected to an automatic sample-injection system (DANI HSS 86.50, HEADSPACE-SAMPLER) was used for N₂O analysis. The oven, injector and detector temperatures were set at 120°C, 120°C and 330°C, respectively. A calibration gas of $5 \,\mu$ II⁻¹ N₂O (Linde Gas) and N₂ in ECD-quality with a flow rate of 30 ml min⁻¹ served as carrier-gas. Carbon dioxide was analysed by using a gas chromatograph (Hewlett-Packard 5890 II series) equipped with a thermal conductivity detector (TCD). Helium was used as carrier-gas (flow rate 10 ml min⁻¹); the CO₂ calibration gas contained 10 ml I⁻¹ CO₂ (Linde Gas).

2.3. NO flux measurements

25

Continuous measurements of NO/NO₂ were completed by a fully automated system (Holtermann, 1996), at an hourly frequency, throughout the vegetation periods of 2002 and 2003. NO and NO₂ were measured with a HORIBA APNA-360 chemoluminescence nitrogen oxide analyzer (detection limit: was 1 ppb; flow rate: 1 I min^{-1}). Calibration was conducted by using a HORIBA calibration unit (Transferstandard, ASGU-360,



BGD

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



with Purafil and activated charcoal, was used in 2002 as no ozone analyzer was available in this year. In year 2003 we used an ozone analyzer (HORIBA APOA-360) for measuring O_3 concentrations within the chambers. Thus, the chemical reaction between NO and O_3 could be considered. It has to be mentioned that the NO₂ measure-

- ⁵ ments with the HORIBA is the sum of NO₂ (most likely the main component) and other nitrogen compounds (PAN, NH₃, HONO). Air samples were taken from 5 chambers (Ø=20 cm; Vol.=3.27 I), plus one reference chamber where the opening to the soil was sealed with a Plexiglas pane. Each chamber was closed for 5 min within which steady state was reached. Fluxes were calculated as described by Schindlbacher et al. (2004)
- for 2002 fluxes and Butterbach-Bahl et al. (1997) for 2003 fluxes. For further details on analysis of gas samples and calculation of fluxes compare Kitzler et al. (2005). In view of the low concentrations and the spatial heterogeneity of NO_x fluxes the uncertainty coming from disregarding wall effects was assumed to be of minor importance (Ludwig, 1994).
- 15 2.4. Soil samples

20

Soil samples were taken approximately every two months. Extractable nitrogen was determined from litter layer (frame: 30×30 cm) and mineral soil (upper 7 cm), and analysed according to Kandeler (1995). Soil moisture was determined gravimetrically. The pH was measured in soil suspensions in 0.01 M CaCl₂ solution using a glass electrode. Additionally to these analyses, an acetylene (C₂H₂) inhibition experiment was carried out twice (May 03 and August 03). Acetylene inhibits the reduction of N₂O to N₂ and is used to quantify the total N-loss (N₂O+N₂) that is produced via denitrification (Klemedtsson et al., 1990).

Per sampling time 16 soil cores from the upper 7 cm mineral soil were incubated ²⁵ in airtight incubation cylinders (volume: 500 ml) and sealed with silicon grease. The headspace of the cylinders was sampled (30 ml) at the beginning of the incubation period and after 4 h at a temperature of 25°C; thus allowing the determination of N₂O production. Afterwards, acetylene was injected into the headspace exposing the soil



1429

core to 10 kPa acetylene. The incubation period lasted again for 4 h at 25°C. Nitrous oxide was analyzed as described in Sect. 2.2.

2.5. Meteorological data

- Soil temperature (in °C, Vitel-thermisters) and soil moisture (in %, Vitel Theta probes)
 were quantified at a soil depth of 5, 15, 25 and 50 cm and stored in a data logger. Air temperature (°C) was measured with a temperature sensor at 2 m above ground. Measuring interval was 4 h. As far as the manual gas sampling time was concerned, soil temperatures at a soil depth of 3 and 10 cm were additionally measured around the chambers. Daily precipitation was recorded by an ombrometer at the nearby meteorological station Pumpwerk Achenkirch. The biweekly input below the crown was calculated from the crown throughfall collectors.
 - 2.6. Deposition measurements
- Dry deposition of NO₂ and NH₃ was captured by three passive diffusion tubes and three CEH ALPHA samplers, respectively (Tang et al., 2001), and were analysed by CEH Edinburgh. They were placed in the canopy at the investigation site and at an open site nearby, at a height of 1.5 m and were changed monthly. Dry deposition velocities were assumed to be 1.5 mm s⁻¹ for NO₂ and 3 mm s⁻¹ for NH₃ (Duyzer, pers. comm.). Wet deposition was collected biweekly using 15 crown throughfall collectors. Wet deposition was analysed for NH₄⁺ and NO₃⁻ (Dionex DX100 and Dionex 120).
- 20 2.7. Statistical analysis

Data were checked for normal distribution. The t-test or the nonparametric Wilcoxontest was used for determining differences in soil emissions, soil chemical and Ndeposition data between the investigation years. Pearson or Spearman rank correlation was used for determining relations between daily or biweekly emission data and

BGD

2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



soil, meteorological or deposition data. A moving average of 2 or 3 was applied for deposition data.

As variables are autocorrelated in the course of time (Durbin-Watson statistic), the autoregression procedure – GARCH (p, q) model – was used to detect influences of soil data, meteorological- or deposition data on N-emissions. Specifications of the GARCH model are described in detail in Kitzler et al. (2005). Variables that emerged to correlate best with N₂O and NO emissions and variables, whose effects on emissions were lagged, were incorporated in the models. Statistical analysis was either completed using SAS Enterpriseguide Version 2 or SAS Version 8. All differences reported were significant at p<0.05 unless otherwise stated.

3. Results

20

3.1. Soil nitrogen and pH

Concentrations of extractable N in the litter layer were at their highest after litterfall in autumn and reached a maximum of $138 \mu g NH_4^+$ -N and $80 \mu g NO_3^-$ -N g⁻¹ dw (dry ¹⁵ weight) in September 2002 (Fig. 1). Mean NH₄⁺ and NO₃⁻ concentrations of the first year were considerably higher than the concentrations measured in the second year (Table 2).

A similar seasonality was also observed with regard to N concentrations in the mineral soil. Here, up to $7.9 \,\mu g \, \text{NO}_3^- \text{N g}^{-1}$ dw and up to $50 \,\mu g \, \text{NH}_4^+ \text{-N g}^{-1}$ dw were reached in autumn 2002 and 2003, respectively. In 2003 considerably lower $\text{NH}_4^+ \text{-N}$ concentrations were detected, particularly in autumn.

Mean soil pH was 6.4 with a maximum of 7.0 and a minimum of 5.4. PH of the litter layer was generally lower (mean=5.9) and was ranging from 5.1-7.1.

BGD

2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



3.2. Nitrogen input

In the two years of investigation, the N-input via throughfall amounted to 10.6 kg and 11.9 kg N ha⁻¹ y⁻¹ (Table 3). Ratio of deposited NO₃⁻:NH₄⁺ is 1.7:1. In general, N from throughfall had its maximum in spring/early summer. In 2003, however, the highest wet
N-deposition was detected in autumn (Fig. 2). No relationship was found between precipitation rates and N-input. Dry deposition was measured from May 2002 to July 2003 and amounted to 1.52 kg N ha⁻¹ y⁻¹, thus amounting to a total of 12.1 kg N ha⁻¹ y⁻¹ through wet and dry deposition in the first year of sampling. NO₂ deposition was at its highest in winter, whereas concentration of NH₃ deposition was at its highest in spring and summer. The forest floor received 21.6 kg N per ha⁻¹ y⁻¹ (1999) through litterfall (Herman et al., 2002).

- 3.3. Gas fluxes
- 3.3.1. Carbon dioxide emissions

Maximum CO_2 emissions were recorded in the summer 2003 and 2004 (Table 4), fol-¹⁵ lowing a seasonal course of air and soil temperature (Fig. 3a). In 2002 the summer peak was not as pronounced and summer emissions were significantly (p<0.01) lower than in the following years. In late autumn emissions decreased steadily and reached a minimum in winter (0.0–0.4 mg CO_2 -C m⁻² h⁻¹).

Mean annual CO₂ emission were $11.6\pm1.1 \text{ mg } \text{CO}_2\text{-}\text{Cm}^{-2}\text{h}^{-1}$ (year 1) and 19.7±2.0 mg CO₂-Cm⁻²h⁻¹ (year 2) showing a significant difference between the two years (Table 4). At our site the cumulative soil respiration rate was $1.3\pm0.3\text{ t}\text{ C}\text{ ha}^{-1}\text{ y}^{-1}$ (Table 5) with a 60% higher annual CO₂ production rate in year 2.

The most influencing parameters controlling CO_2 emissions (Table 6) were soil temperature at a soil depth of 3–25 cm (r^2 =0.86) and air temperature (r^2 =0.70). Furthermore, a positive correlation could be detected with N-deposition (r^2 =0.62). A negative relationship was found between CO_2 emission and soil moisture in the upper 5 cm

BGD

2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



 $(r^2 = -0.59)$. All correlations were highly significant (p<0.001).

3.3.2. N₂O emissions

25

Nitrous oxide emissions were mostly dependent on air and soil temperature, as well as soil moisture. Figure 3b shows the high spatial and temporal variability during the field ⁵ survey from May 2002 until July 2004. Spatial variation between the chambers was high (mean CV: 50%) and reached a maximum in winter (CV: 350%).

Mean annual N₂O emissions averaged $4.4\pm0.3\,\mu$ g N₂O-N m⁻² h⁻¹ (Table 4). In the course of the year highest N₂O emissions were observed in summer 2003 (40.6 μ g N₂O-N m⁻² h⁻¹). Although we missed the summer peak with the manual chamber system in year 2002, we were able to observe high N₂O emissions by the AGPS (Fig. 3b). During winter, when the site was covered with snow, even a significant uptake of atmospheric N₂O (range: -0.3 to -3.5 μ g N₂O-N m⁻² h⁻¹) could be observed. Significant differences in the seasonality of N₂O emissions between both years could only be observed for August (p<0.01).

¹⁵ Significant correlations could be found between C- and N-emissions, and soil, meteorological and N-input data (Table 6). Log-transformed N₂O emission rates measured by the manual system were positively correlated (p<0.001) to soil respiration rates (r^2 =0.58), soil temperature in a depth of 3 and 10 cm (r^2 =0.64) and air temperature (r^2 =0.58). A negative dependency was found with soil moisture (r^2 =-0.52). Correlation analysis between daily N₂O emissions from the AGPS system and soil temperature and soil moisture revealed weaker but significant relationships (r^2 =0.27 and r^2 =-0.32, p<0.001).

Nitrate and NH₄⁺ deposition via throughfall and NO₃⁻ concentration in the mineral soil showed significant positive effects on N₂O emission (r^2 =0.26, r^2 =0.25, p<0.001 and r^2 =0.41, p<0.05), even higher correlation was found with NO₃⁻ concentration in the litter layer (r^2 =0.44, p<0.01).

Simple regression models were insufficient for revealing dependencies of emission

BGD 2, 1423-1455, 2005 N_2O , NO_x and CO_2 fluxes in a calcareous mountain forest soil B. Kitzler et al. **Title Page** Abstract Introduction References Conclusions **Figures** Tables Back Close Full Screen / Esc Print Version

Interactive Discussion

EGU

data as residuals were correlated over time. Two GARCH models could be developed to predict N₂O emissions as measured with the manually operated system. Independent variables in model 1 were (x_{1t}) soil moisture at a soil depth of 5 cm and (x_{2t}) soil temperature at a soil depth of 3 cm. As an effect of N-deposition on N₂O emission was discovered while carrying out the correlation analysis (Table 6), we included the monthly sum of nitrogen input (NO₃⁻ and NH₄⁺) via throughfall within model 2 and achieved an improvement of modelled N₂O emissions in comparison to simple regression analysis. Both models follow the GARCH (2, 1) process and revealed a r² of 0.89 (model 1) and r² of 0.83 (model 2) (Fig. 4). Daily N₂O emissions from the AGPS showed a high temporal variability. Moreover, correlations with soil parameters were lower compared to the N₂O emissions measured by the spatially extended manual system. The GARCH model (Fig. 4, model 3 and Table 7) revealed that precipitation and air temperature on daily N₂O emissions are delayed by 8 days (p<0.001, r²=0.43).

3.3.3. NO_x emissions

- ¹⁵ NO and NO₂ flux rates from soil were measured from July 2002 until November 2003 (Fig. 3c) but not during the winter months. Mean NO emissions in the vegetation period were $0.7\pm0.04\,\mu$ g NO-N m⁻² h⁻¹ and hence significantly higher in year 2002 (p<0.001) compared to the year 2003 ($0.5\pm0.07\,\mu$ g NO-N m⁻² h⁻¹) when low soil N concentrations were measured. Mean NO emissions were generally lower than N₂O emission rates. The ratio NO/N₂O was 1:3. A consistent seasonal trend could not be observed: In the year 2002 highest emissions occurred in autumn, whereas in
- year 2003 maximum emissions were measured in spring (Fig. 3c). Lowest NO emissions or even negative fluxes were observed in November 2003. NO₂ was deposited over the entire observation period with a mean of $-0.4 \pm 0.2 \,\mu$ g NO m⁻² h⁻¹ (2002) and $-0.3 \pm 0.1 \,\mu$ g NO₂ m⁻² h⁻¹ (2003).

Correlation analysis (Table 6) revealed a weak positive relationship between mean daily NO fluxes and soil temperature at the soil depth of 5 cm ($r^2=0.19$, p<0.5), whereas

BGD

2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



a highly significant positive dependency could be observed with soil moisture in 5 cm, $(r^2=0.55, p<0.001)$. No valid statistical explaination could be given to characterize NO emissions by a simple regression model. By using the autoregression procedure, however, a GARCH (1,1) model could be developed to predict daily NO emissions of our site (Fig. 4, model 4). Through a mean of daily soil moisture content (x_{1t}) and

- ⁵ our site (Fig. 4, model 4). Inrough a mean of daily soil moisture content (x_{1t}) and soil temperature (x_{2t}) at a soil depth of 5 cm, predictions of mean daily NO emissions could be carried out with an r² of 0.77 (Table 7). This model revealed a time lag of 2, signifying that the independent variables have an influence on NO emissions over two days.
- 10 3.3.4. Actylen inhibition

By using the C₂H₂ inhibition method the reduction of N₂O to N₂ is inhibited and total N-loss, which is produced via denitrification, can be measured. The increase of N₂O production in the Acetylene-treated samples, in comparison to the samples without added C₂H₂ was between 101% and 4200% with a median of 266%. The ratio of N₂O:N₂ was 1:6. Mean N₂ production of the two experiments (n=16) was 11.6±1.4 μ g N₂ m⁻² h⁻¹

and $19.1 \pm 3.9 \mu g N_2 m^{-2} h^{-1}$.

4. Discussion

15

4.1. CO₂ emissions

Carbon dioxide emissions, a measure for general microbial activity, followed a typical seasonal trend, with highest rates in summer, when mineralization of organic matter occurs. Lowest rates were measured in winter. Organic carbon is converted to CO₂ during mineralization. This process is strongly dependent on soil temperature and soil moisture. At our site, soil temperature at a soil depth of 3 and 10 cm was mostly responsible for temporal variation in soil respiration. This finding is in good agreement

BGD 2, 1423-1455, 2005 N_2O , NO_x and CO_2 fluxes in a calcareous mountain forest soil B. Kitzler et al. **Title Page** Abstract Introduction References Conclusions Figures Tables Back Close Full Screen / Esc Print Version Interactive Discussion EGU

with results from other studies (Epron et al., 1999; Merino et al., 2004). Soil moisture showed also a significant effect on soil respiration rates. Carbon dioxide release was reduced during periods of heavy rain (summer 2002), when the water content was between 50–65%, probably as a consequence of O_2 deficiency in soil due to diffusion restrictions (Howard and Howard, 1993). Further significant relationships could be established between biweekly mean CO_2 emissions and biweekly N-deposition rates. The cumulative soil respiration rates at our site $(1-1.6 \text{ t C ha}^{-1} \text{ y}^{-1})$ are lower than values for temperate coniferous forests reported by Raich and Schlesinger (1992).

4.2. N₂O emissions

5

- At our site, seasonal variations in N₂O emissions followed mainly the annual changes in soil temperature, soil moisture and the availability of N in the soil. Highest emissions were thus observed during summer/autumn. Although, we missed out on the summer peak in year 2002 through the manual chamber system (biweekly sampling), we were able to observe high emissions by the AGPS. Correlation analysis revealed that soil temperature was the most important factor controlling N₂O emissions followed by soil moisture. Those parameters affect microorganisms and their metabolism and, hence, the production and consumption of N trace gases in soils (Conrad, 1996). The air-filled porosity controls the movement of the gases towards and away from the atmosphere;
- it also affects soil aeration, and, thus, indirectly controls the capacity of the soil for producing or consuming soil-produced trace-gases (Smith et al., 2003); (Davidson et al., 2000). In a laboratory parameterisation study, Schindlbacher et al. (2004) found that for the AK site a N₂O emission maximum for a soil moisture range of 60%–90% water filled pore space (WFPS) which corresponds to a water content of 53–80%. In the field we found, beside the peak in August 2003, that maximum N₂O emissions occurred at
- water moisture values in the range of 50%–65%, i.e. in accordance to the laboratory results of Schindlbacher et al. (2004). With the GARCH model 1 N₂O emissions could be predicted best using soil moisture and soil temperature as predictors (r^2 =0.89).

BGD

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



This analysis also revealed that past environmental conditions (up to 12 weeks) can significantly affect actual N₂O emissions. Within the model, soil moisture showed a highly significant effect (p<0.001) whereas the effect of soil temperature was lower (p<0.01). Furthermore, precipitation and air temperature showed a clear influence in daily (AGPS) N₂O emissions (model 3). Here a highly significant effect of conditions during the past 8 days were observed.

Highest emissions were observed in August 2003. There are various parameters that might have contributed to the N_2O peak. Even at this site, where soil moisture is generally high, the water content during the European drought period of summer 2002 degreesed to 200%. After a reinfoll quart pair maintenance in general data and mark

- ¹⁰ 2003 decreased to 30%. After a rainfall event soil moisture increased again and may have caused an increase in microbial activity. On the other hand, the actual level of N-deposition and the NO₃⁻ pool in the soil might have played an important role in the sudden release of N₂O (Davidson et al., 2000). Highest amounts in NO₃⁻ and NH₄⁺ via throughfall reached the forest floor in August. Although, throughout the study wet
- ¹⁵ N-deposition accounted for up to 31% of the variation in N₂O emission (correlation analysis) only, a strong relationship was apparent between the two factors. This high N₂O peak coincided with significant larger available soil N concentrations, in particular soil NO₃⁻. Using model 2 the relationship between biweekly measured N₂O emissions and biweekly sum of nitrogen input (NO₃⁻ and NH₄⁺) via throughfall was found by a
- high model efficiency (r^2 =0.83). The time series analysis revealed a highly significant (p<0.001) lagged effect of wet N-deposition on N₂O emission for 2 weeks, but correlations are expected for a 3-month period.

The soil at the Mühleggerköpfl is, typically, moist (mean 55%) and denitrification may have been the main source of N₂O emissions throughout the year. A high pH value is also one of the factors that provide favourable conditions for denitrification at the site. Previous studies (Härtel et al., 2002) have indicated that complete nitrification takes place at our site and that it is of major importance for N₂O emission. ¹⁵NO₃⁻ was strongly immobilized. Furthermore, close C/N-ratios (16–18) suggest favourable conditions for N-mineralization and nitrification (Herman et al., 2002). It was in Au-

BGD

2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



gust/September that nitrate concentration increased and we suppose that the high N_2O peak in year 2003 is the product of nitrification.

At our site winter emissions were generally low and the site even functioned as a sink for N₂O. The AGPS system, in particular, was a useful means for detecting negative
fluxes. Butterbach-Bahl et al. (2002) detected negative N₂O fluxes from pine forests with moderate N-deposition whereas a pine forest with high N-loads exclusively functioned as a source of N₂O during winter. On the other hand winter emissions, comparable to the ones from the rest of the year (January 2003; 4.3±1.3 µg N₂O-N m⁻² h⁻¹), could be observed after freeze-thaw events. These high emissions were either a result
of enhanced denitrification activity or could be due to the physical release of accumulated N₂O in the snow or in the soil (Teepe et al., 2001).

4.3. NO emissions

The production of NO is strongly dependent on climate and varies considerable with soil temperature and soil moisture (Davidson et al., 2000; Ludwig et al., 2001; Van Dijk
et al., 2002). We found best correlations with soil moisture and only little dependency on soil temperature. The optimum soil temperature, as found in laboratory studies (20°C) by Schindlbacher et al. (2004), could not be confirmed in the field experiment, where highest NO emissions were found at 8–10°C soil temperature; maximum soil temperature in the field was 16°C. Optimum WFPS in the field for NO emission was found to be at 30–45% WFPS (water content 27–40%) and was, thus, higher than in laboratory studies (Schindlbacher et al., 2004). In the field highest NO emissions were found in autumn and in spring, when soil water content was between 45–50%. High spring emissions were detected in May, when soil was still moist and soil temperature increased. At our site soil moisture had a stronger impact on NO emission than soil

temperature. With the GARCH procedure a relationship between NO emissions and soil moisture and soil temperature could be found with a model efficiency of r² of 0.77 (model 4). Autoregressive parameters were estimated for a time-lag of 2 days; i.e. soil moisture and soil temperature affected NO emission over a two days period at our site. 2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



Soil nitrogen content contributed to high NO emission in autumn, where highest NO_3^--N and NH_4^+-N concentrations were measured. As in autumn 2003 the N-pool (especially extractable NH_4^+-N) in the soil and the litter layer was low compared to concentrations measured in autumn 2002 no increase in NO emissions could be found. The effect of rewetting the soil as it was found for N_2O emissions could not be seen for NO emissions.

Soil acidity is supposed to be an important factor promoting N-losses from forest soils influenced by atmospheric nitrogen deposition (Li et al., 2000 and Venterea et al., 2003). Soils with a pH of 7–8 are supposed to produce higher amounts of N_2 (Simek concerner 2002) but revealed on order of magnitude higher NLO emissions then exile

- ¹⁰ and Cooper, 2002) but revealed an order of magnitude higher N₂O emissions than soils with a pH between 4 and 6 (Vermoesen et al., 1996). A soil pH of 6.7–7.1 measured at our limestone site restricts self-decomposition of HNO_2 and might, therefore, be responsible for low NO fluxes (Van Cleemput and Samater, 1996). Beside the high pH and the high microbial biomass (Härtel et al., 2002), the well-balanced community com-
- position of the microflora (Zechmeister-Boltenstern, pers. comm.) could be a reason for low NO-fluxes. Results of PLFA studies of European soils by (Zechmeister-Boltenstern, pers. comm.) showed that at acidic high N-input sites the microbes are under stress which leads to structural changes of their cell membranes. This could cause losses of intermediate products in the nitrification or denitrification pathway and could, thereby, load to higher NO and N. O emissions. The assumption of a well balanced microflora.
- ²⁰ lead to higher NO and N₂O emissions. The assumption of a well-balanced microflora was supported by the acetylene inhibition and simulation runs with the Pnet-N-DNDC where high N₂ emissions were detected and a large portion of N was fully recycled (Butterbach-Bahl, pers. comm.).
 - 4.4. Annual N-losses

5

²⁵ In both years N released in form of N₂O was between 0.3 and 0.4 kg N₂O-N ha⁻¹ y⁻¹, and 0.03 kg NO-N ha⁻¹ (Table 5). These values correspond to a yearly release of actual N input of 2.5–3.5% in form of N₂O and 0.2% in form of NO and are located in the lower 2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



reported percentage of deposited N which is 0–16% (N₂O) and 0–10% (NO) (Skiba et al., 2004). Through the acetylene inhibition we calculated a N₂/N₂O ratio of 6.0 and concluded a rough estimation of 2.2 kg N ha⁻¹ of annual N₂ loss. Simulated mean N₂ production by the model PnetNDNDC for years 1998–2001 (Zechmeister-Boltenstern, unpublished) was 14 kg, which would mean that more than 18% of the deposited N were lost by the complete pathway of microbial soil processes as N₂. As the acetylene inhibition was only conducted twice within the investigation period this value has to be interpreted with caution. However, it can be hypothesized that the deposited N from the atmosphere is mainly emitted as N₂ at this limestone site.

The magnitude of N₂O emissions from our site are consistent with values reported of other coniferous forests (MacDonald et al., 1997; Papen and Butterbach-Bahl, 1999; Butterbach-Bahl et al., 2002) and limed beech forests (Borken and Brumme, 1997). Considerable higher fluxes were measured in more acidic spruce forests, where high loads of N was deposited (Papen and Butterbach-Bahl, 1999; Butterbach-Bahl et al., 2002) and in deciduous forest ecosystems with low pH (Brumme and Beese, 1992; Borken and Brumme, 1997).

Nitric oxide emission measured in our study are low in comparison to other published NO fluxes from temperate coniferous forests. But our rates are in accordance with studies by (Johansson, 1984) who reported annual NO emissions of 0.04 kg NO-N ha⁻¹ from an unfertilized coniferous forest soil in Sweden. The majority of studies, however, revealed significantly higher values. Pilegaard et al. (1999) and Butterbach-Bahl et al. (2002) found in moderately N-effected spruce forests annual NO losses of 3.15 kg NO-N ha⁻¹ and 3.1 μ g NO-N m⁻² h⁻¹, respectively. It has to be taken into account that most of the reported data were measured at forest soils with low pH values,

however, NO fluxes are reported to be significantly higher in more acidic soils compared to N₂O fluxes (Yamulki et al., 1997).

Nitrogen oxide emissions could be predicted with high efficiencies. Nitrous oxide emissions measured by manual chambers revealed an r^2 of 0.89 (model 1) and an r^2 of 0.83 (model 2). Through model 2, where N-deposition is the only independent

BGD

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



variable, annual emission rates could be estimated better than with model 1, where soil moisture and soil temperature were the crucial parameters. Especially in year 2003, where high emissions were detected in August, the predicted emissions with model 2 revealed a better accuracy as model 1. High variations in N₂O emissions measured

⁵ by the AGPS led to an overestimation of N_2O emissions through model 3 (21–27%). Nitric oxide emissions predicted with model 4 could be predicted with an r^2 of 0.77 and annual simulated fluxes were in accordance with measured annual NO emissions.

It can be concluded that with time series analysis lagged effects of the controlling factors soil moisture, soil temperature and N-deposition on N_2O and NO emissions could be revealed. In particular significant two weeks delayed response of N_2O emission to

- ¹⁰ be revealed. In particular significant two weeks delayed response of N_2O emission to N-deposition was observed. Nitric oxide emissions were three times lower compared to N_2O emissions and total losses of N were small compared to N-inputs. This may be attributed to the fact that in the limestone Alps soil pH is high, a large microbial biomass can be found and nitrogen is mainly emitted in the form of N_2 .
- Acknowledgements. We would like to thank the landowners for access on their land. A. Fiege, J. Pausch, J. Plattner, and F. Winter for field work, C. Abo-Elschabaik and M. Kitzler for helping with the sample analysis and S. Smidt for provision of meteorological data. The study is a part of the NOFRETETE project EVK2-CT2001-00106 funded by the European Commission DG Research – Vth Framework Programme.

20 **References**

Borken, W. and Brumme, R.: Liming practice in temperate forest ecosystems and the effects on CO₂, N₂O and CH₄ fluxes, Soil use and management, 13, 251–257, 1997.
Brumme, R. and Beese, F.: Effects of liming and nitrogen fertilization on emissions of CO₂ and N₂O from temperate forest, J. Geophys. Res., 97, 12, 851–858, 1992.

²⁵ Butterbach-Bahl, K., Gasche, R., Breuer, L., and Papen, H.: Fluxes of NO and N₂O from temperate forest soils: impact of forest type, N deposition and of liming on NO and N₂O emissions, Nutr. Cycl. Agroecosyst., 48, 79–90, 1997.

BGD 2, 1423-1455, 2005 N_2O , NO_x and CO_2 fluxes in a calcareous mountain forest soil B. Kitzler et al. **Title Page** Abstract Introduction References Conclusions Figures Tables Back Close Full Screen / Esc Print Version Interactive Discussion EGU

- Butterbach-Bahl, K., Breuer, L., Gasche, R., Willibald, G., and Papen, H.: Exchange of trace gases between soils and the atmosphere in Scots pine forest ecosystems of the northeastern German lowlands: 1. Fluxes of N₂O, NO/NO₂ and CH₄ at forest sites with different N-deposition, Forest Ecology and Management, 167, 1–3, 123–134, 2002.
- ⁵ Conrad, R.: Soil microrganisms as controllers of atmospheric trace tases (H₂, CO, CH₄, OCS, N₂O, NO), Microbiology Reviews, 60, 609–640, 1996.
 - Crutzen, P. J.: Ozone in the troposphere, in: Composition, Chemistry, and Climate of the Atmosphere, edited by: Signh, H. P., Van Nostrand Reinhold, New York, 1995.
 - Crutzen, P. J.: The role of NO and NO₂ in the chemistry of the troposphere and stratosphere, Annu. Rev. Earth Planet. Sci., 7, 443–472, 1979.
- Annu. Rev. Earth Planet. Sci., 7, 443–472, 1979.
 Davidson, E. A., Keller, M., Erickson, H. E., Verchot, L. V., and Veldkamp, E.: Testing a conceptual model of soil emissions of nitrous and nitric oxides, BioScience, 50, 667–680, 2000.
 Englisch, M. and Starlinger, F.: Waldgesellschaften und Standorte im Bereich der Höhenprofile Achenkirch, FBVA-Bericht, 25–54, 1995.
- ¹⁵ Epron, D., Farque, L., Lucot, E., and Badot, P. M.: Soil CO₂ efflux in a beech forest: the contribution of root respiration, Annals of Forest Science, 56, 4, 289–295, 1999.
- Firestone, M. K. and Davidson, E. A.: Microbial Basis of NO and N₂O Production and Consumption in Soil, in: Exchange of Trace Gases between Terrestrial Ecosystems and the Atmosphere, edited by: Andreae, M. O. and Schimel, D. S., John Wiley, New York, 7–21, 1989.
 - Härtel, E., Zechmeister-Boltenstern, S., and Gerzabek, M.: Gaseous nitrogen losses from a forest site in the north Tyrolean limestone alps, Environmental Science and Pollution research, Special Issue 2, 23–30, 2002.

Herman, F., Smidt, S., Englisch, M., Feichtinger, F., Gerzabek, M., Haberhauer, G., Jandl, R.,

- Kalina, M., and Zechmeister-Boltenstern, S.: Investitations of nitrogen fluxes and pools on a limestone site in the Alps, ESPR – Environmental Science and Pollution Research, 46–52, 2002.
 - Holtermann, C.: A transportable system for the on-line-measurement of NO_x (NO, NO_2)emission from soils, Die Bodenkultur, 47, 235–244, 1996.
- ³⁰ Howard, D. M. and Howard, P. J.: Relationships between CO₂ evolution, moisture content and temperature for a range of soil types, Soil Biology & Biochemistry, 25, 1537–1546, 1993.
 - IAEA: Manual on measurements of methane and nitrous oxide emissions from agriculture, International Atomic Energy Agency, Vienna, 90, 1992.

2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil

Title Page						
Abstract	Abstract Introduction					
Conclusions	References					
Tables	Figures					
I	۶I					
•	Þ					
Back	Close					
Full Screen / Esc						
Print Version						
Interactive Discussion						
FGU						

- IPCC: Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel of Climate Change, in: Cambridge University Press, edited by: Houghton, J.T., Ding, Y., Griggs, D. J., Noguer, M., Van der Linden, P. J., and Xiaosu, D., Cambridge, United Kingdom and New York, NY, USA, 881, 2001.
- Johansson, C.: Field measurements of emission of nitric oxide from fertilized and unfertilized forest soils in Sweden, J. Atmos. Chem., 1, 429–442, 1984.

Kandeler, E.: Ammonium and Nitrate, in: Methods of Soil Biology, edited by: Schinner, F., Öhlinger, T., Kandeler, E., and Margesin, R., Springer, Berlin, 406–410, 1995.

Kitzler, B., Zechmeister-Boltenstern, S., Holtermann, C., Skiba, U., and Butterbach-Bahl, K.: Nitrogen oxides emission from two beech forests subjected to different nitrogen loads, Biogeosciences Discuss., 2, 1381–1422, 2005,

SRef-ID: 1810-6285/bgd/2005-2-1381.

5

Klemedtsson, L., Hanson, G., and Mosier, A.: The use of acetylene for the quantification of N₂

- and N₂O production from biological processes in soils, in: Denitrification in soil and sediment, edited by: Sorensen, J. and Revsbech, N. P., Plenum Press, 167–180, 1990.
 - Li, C., Aber, J. D., Stange, F., Butterbach-Bahl, K., and Papen, H.: A process-oriented model of N₂O and NO emission from forest soils,1, Model development, J. Geophys. Res., 105(4) 4369–4384, 2000.
- Lovett, G. M., Weathers, K. C., and Arthur, M. A.: Control of nitrogen loss from forested watersheds by soil carbon: Nitrogen ratio and tree species composition, Ecosys., 5, 712–718, 2002.
 - Ludwig, J.: Untersuchungen zum Austausch von NO und NO₂ zwischen Atmosphäre und Biosphäre, PhD Thesis, University of Bayreuth, 251, 1994.
- ²⁵ Ludwig, J., Meixner, F. X., Vogel, B., and Förstner, J.: Soil-air exchange of nitric oxide: An overview of processes, environmental factors, and modeling studies, Biogeochemistry, 52, 225–257, 2001.
 - MacDonald, J., Skiba, U., Sheppard, L. J., Ball, B. C., Roberts, J. D., Smith, K. A., and Fowler, D.: The effect of nitrogen deposition and seasonal variability on methane oxidation and ni-
- trous oxide emission rates in upland spruce plantation and moorland, Atmos. Environ., 31, 22, 3693–3706, 1997.
 - Merino, A., Pérez-Batallón, P., and Macías, F.: Responses of soil organic matter and greenhouse gas fluxes to soil management and land use changes in a humid temperate region of

2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil

Title Page						
Abstract	Abstract Introduction					
Conclusions	References					
Tables	Figures					
I	۶I					
•	•					
Back	Back Close					
Full Screen / Esc						
Print Version						
Interactive Discussion						
FGU						

southern Europe, Soil Biology & Biochemistry, 36, 917-925, 2004.

- Mutsch, F.: Bodenchemische Charakterisierung des Mühleggerköpfls im Rahmen einer Untersuchung über Stickstoff-Flüsse in den Norditoler Kalkalpen, FBVA Berichte, 116, Wien, 33–40, 2001.
- 5 NADP: National Atmospheric Deposition Program 2001 Summary. NADP Report, Illinois State Water Survey, Champaign, IL, 2002.
 - Papen, H. and Butterbach-Bahl, K.: A 3-year continuos record of N-trace gas fluxes from untreated and limed soil of a N-saturated spruce and beech forest ecosystem in Germany: I. N₂O-emissions, J. Geophys. Res., 104, 18487–18503, 1999.
- ¹⁰ Pilegaard, K., Hummelshoj, P., and Jensen, N. O.: Nitric oxide emission from a Norway spruce forest floor, J. Geophys. Res., 104, 3433–3445, 1999.
 - Raich, J. W. and Schlesinger, W. H.: The global carbon dioxide flux in soil respiration and its relationship to vegetation and climate, Tellus, 44B, 81–99, 1992.

Schindlbacher, A., Zechmeister-Boltenstern, S., and Butterbach-Bahl K.: Effects of soil mois-

- ture and temperature on NO, NO₂ and N₂O emissions from forest soils, J. Geophys. Res., 109, D17302, 1–12, 2004.
 - Simek, M. and Cooper, J. E.: The influence of soil pH on denitrification: progress towards the understanding of this interaction over the last 50 years, European Journal of Soil Science, 53, 3, 345–354, 2002.
- Skiba, U., Pitcairn, C. E. R., Sheppard, L. J., Kennedy, V. H., and Fowler, D.: The influence of atmospheric N deposition on nitrous oxide and nitric oxide fluxes and soil ammonium and nitrate concentrations, Water, Air and Soil Pollution, Focus 4, 37–43, 2004.
 - Smidt, S., Herman, F., Grill, D., and Guttenberger, H.: Studies of ecosystems in the Limestone Alps "Achenkirch Altitude Profiles", Phyton, 36, Horn, Austria, 1996.
- Smith, K. A. Ball, T., Conen, F., Dobbie, K. E., Massheder, J. and Rey, A.: Exchange of greenhouse gases between soil and atmosphere: interactions of soil physical factors and biological processes, Eur. J. Soil Science, 54, 4, 779–791, 2003.
 - Tang, Y. S., Cape, J. N., and Sutton, M. A.: Development and types of passive samplers for monitoring atmospheric NO₂ and NH₃ concentrations, The Scientific World, 1, 513–529, 2001.

30

Teepe, R., Brumme, R., and Beese, F.: Nitrous oxide emissions from soil during freezing and thawing period, Soil Biol Biochem, 33, 1269–1275, 2001.

Van Cleemput, O. and Samater, A. H.: Nitrite in soils: accumulation and role in the formation of

BGD

2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



gaseous N compounds, Fertilizer Research, 45, 81-89, 1996.

- Van Dijk, S. M., Gut, A., Kirkman, G. A., Meixner, F. X. and Andreae, M. O.: Biogenic NO emissions from forest and pasture soils: Relating laboratory studies to field measurements, J. Geophys. Res., 107, D20, 8058, 2002.
- ⁵ Venterea, R. T., Groffman, P. M., Verchot, L. V., Magill, A. H., Aber, J. D., and Streudler, P. A.: Nitrogen oxide gas emissions from temperate forest soils receiving long-term nitrogen inputs, Global Change Biology, 9, 346–357, 2003.
 - Vermoesen, A., de Groot, C., Nollet, L., Boeckx, P., and Van Cleemput, O.: Effect of ammonium and nitrate application on the NO and N₂O emission out of different soils, Plant and Soil, 181, 153–162, 1996.

10

15

WHO: Updating and revision of the air quality guidelines for Europe, Les Diablerets, 1995.

Williams, E. J., Hutchinson, G. L., and Fehsenfeld, F. C.: NO_x and N₂O emissions from soil, Global Biogeochem. Cycles, 6, 351–388, 1992.

Yamulki, S., Harrison, R. M., Goulding, K. W. T., and Webster, C. P.: N₂O, NO and NO₂ fluxes from a grassland: Effect of soil pH, Soil Biol. Biochem., 29, 8, 1199–1208, 1997.

Zechmeister-Boltenstern, S., Hahn, M., Meger, S., and Jandl, R.: Nitrous oxide emissions and nitrate leaching in relation to microbial biomass dynamics in a beech forest soil, Soil Biol. Biochem, 34, 823–832, 2002.

BGD

2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil

Title Page					
Abstract	Introduction				
Conclusions	References				
Tables	Figures				
14	►I				
•	•				
Back	Close				
Full Screen / Esc					
Print Version					
Interactive Discussion					
FGU					

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil

B. Kitzler et al.

Title Page					
Abstract	Introduction				
Conclusions	References				
Tables	Figures				
I	۶I				
•	•				
Back	Close				
Full Screen / Esc					
Print Version					
Interactive Discussion					
EGU					

Table 1. Site and soil characteristics of the investigation site AK.

Location	47°34′50′′ N 11°38′21′′E
Vegetation	Aposerido-Fagetum caricetosum albae –
	Carici albae-Fagetum ^{a)}
Stand age [y]	127
Exposition, elevation [m]	N, 895 m a sl.
Tree height [m]	19.1
Basal area [m ² ha ⁻¹]	40
Soil type	Rendzic Leptosols / Chromic Cambisol
Soil texture	loam
Soil aciditiy 0-7cm (pH CaCl ₂)	5.8-7.1
C:N	16-18
Mean precipitation [mm]	1733 ^{b)}
Mean air temp [°C]	6.52 ^{b)}
Soil density (g cm ⁻³)	0.3
$N_{tot} (mg g^{-1})$	9 ^{c)}
$C_{org} (mg g^{-1})$	150 ^{c)}

^{a)}Englisch and Starlinger 1995, ^{b)} Mean of years 1998-2003, ^{c)} Mutsch 2001.

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil

B. Kitzler et al.

NH +	65 8 (8 5)	431(40)		
1114	05.0 (0.5)	43.1 (4.0)		
NO_3^-	31.4 (4.4)	18.3 (2.2)		
pH (CaCl ₂)	6.1 (0.1)	5.8 (0.1)		
$\mathrm{NH_4}^+$	21.0 (3.0)	8.8 (1.2)		
NO ₃	1.5 (0.3)	2.1 (0.3)		
pН				
$(CaCl_2)$	6.5 (0.1)	6.3 (0.1)		
are means with standard error in				

year 2

Table 2. Mean soil nitrogen $[\mu g N g^{-1} dw]$ and pH (CaCl₂) in the litter layer and in the upper 7 cm mineral soil in year 1 and year 2 at AK.

year 1

Note. Soil data are parenthesis.

Litter layer

Mineral soil



2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil

B. Kitzler et al.

Title Page		
Abstract	Introduction	
Conclusions	References	
Tables	Figures	
I	► I	
•	•	
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		
FGU		

Table 3. Wet and dry deposition $[kg N ha^{-1} y^{-1}]$, precipitation via throughfall, soil nitrogen and pH (CaCl₂) in study year 1 and study year 2 at AK.

	year 1	year 2
Precipitation [mm]	1416	1067
N-input by wet deposition/throug	ghfall	
$\mathrm{NH_4}^+$	3.5	4.6
NO ₃	7.1	7.3
Sum of wet deposition	10.6	11.9
N-input by dry deposition		
NH ₃	0.56	n.d.
NO_2	0.96	n.d.
deposition	1.52	n.d.

n.d. = not determined.

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil

B. Kitzler et al.

Title Page					
Abstract	Abstract Introduction				
Conclusions	References				
Tables	Figures				
[∢	۶I				
•	•				
Back	Close				
Full Screen / Esc					
Print Version					
Interactive Discussion					
EGU					

Table 4. Mean annual CO_2 -C, N_2O -N and mean NO-N losses in the vegetation period, minimum and maximum values and number of observations at AK for the two investigation years.

		CO_2 -C	N_2O	NO
		$[mg C m^{-2} h^{-1}]$	[µg N 1	$n^{-2} h^{-1}$]
year 1	mean	11.6	4.5	0.7
	min-max	(0-56.4)	(-3.5 - 34.0)	(0.4-2.5)
	Ν	95	242	105
year 2	mean	19.7	4.4	0.5
	min-max	(0.2-83.0)	(-2.9-40.6)	(-0.1-7.6)
	Ν	96	288	145

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil

B. Kitzler et al.

Title Page					
Abstract Introduction					
Conclusions	References				
Tables	Figures				
•	►I				
•	•				
Back	Close				
Full Screen / Esc					
Print Version					
Interactive Discussion					
EGU					

Table 5. Measured and predicted annual CO_2 -C, N_2O -N losses (± S.E) from the manual chambers for the two investigation years at the site AK; N_2O -N (AGPS) and NO-N losses are calculated for June–November 2002 (year 1) and May–November 2003 (year 2).

	CO ₂	N ₂ O-manual		N ₂ O-AGPS		NO	
	kg C ha ⁻¹ y ⁻¹		kg N ha ⁻¹ y ⁻¹				
	measured	measured	predicted 1) 2)	measured	predicted 3)	measured	predicted 4)
year 1	1020±241	0.30±0.11	0.36±0.04	0.22 ± 0.02	0.28±0.01	0.03 ± 0.002	0.03±0.001
			0.33±0.03				
year 2	1636±283	0.41±0.09	0.26 ± 0.04	0.19 ± 0.02	0.23±0.01	0.03 ± 0.004	0.03 ± 0.004
-			0.38±0.04				

 $^{(1)-4)}$ Model 1 - Model 4.

Table 6. Matrix of Pearson correlation coefficients for significant relationships of mean C- and N-emissions (M=manual chambers, A=AGPS) and soil-, meteorological- and N-input data at the site AK. Total N-deposition (Total N_{dep}) is calculated from the first investigation year when dry deposition was measured.

	lnCO ₂	lnN ₂ O M	lnN ₂ O A	lnNO
CO ₂		0.58 ***		
N ₂ O	0.58 ***			
NO				
Air temperature	0.70 ***	0.58 ***	0.16 *	
Soil temperature	0.86 ***	0.64 ***	0.27 *** -0.32 ***	0.19 ** 0.55 ***
Soil moisture	-0.59 ***	-0.52 ***		
NO ₃ ⁻ litter layer	0.39 *	0.44 **		
NO ₃ ⁻ mineral soil		0.41 *		
TF NO ₃	0.43 ***	0.26 ***		
$\mathrm{TF}\mathrm{NH_4}^+$	0.21 **	0.25 ***		
TF sumN	0.46 ***	0.31 ***		
DRY_NO ₂	-0.66 *	-0.37 **		
DRY_NH ₃	0.71 ***	0.37 **		
Total N _{dep}	0.62 ***	0.33 *		

TF = throughfall, DRY = dry deposition. A moving average of 2 was applied to deposition data. Asterisk indicates the statistic significance (* p<0.05, **p<0.01, ***p<0.001).

BGD

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil

Title Page					
Abstract	Introduction				
Conclusions	References				
Tables	Figures				
14	N				
•	•				
Back	Close				
Full Screen / Esc					
Print Version					
Interactive Discussion					

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil

B. Kitzler et al.

Title Page					
Abstract	Introduction				
Conclusions	References				
Tables	Figures				
•	•				
Back	Close				
Full Screen / Esc					
Print Version					
Interactive Discussion					
EGU					

1451

Table 7. Parameter estimation for the autoregression models 1 to 4 to predict N_2O and NO emissions from the study site, AK. Independent variables (x_1) and (x_2) for the individual models are shown in the table (regression coefficients). Sample period is from April 2002–May 2004.

Model			1	2	3	4
independent variable (y_t) :		N ₂ O	N_2O	N_2O	NO	
Intercept:	${m eta}_0$		-12.9481 **	1.9971 ***	7.0895 ***	1.0429 ***
regression	coefficie	nts of the inde _l	vendent variables	(x_1) and (x_2) w	ith significance le	evel:
β_1	(x_1)	soil moist.	0.2885 ***			-0.0034 *
	(x_1)	wet N dep.		-0.4061 *		
	(x_1)	precip.			0.0851 ***	
β_2	(x_{2})	soil temp.	0.4753 **			-0.0308 *
	(x_{2})	air temp.			-0.1992 *	
total R^2 :	. 2.	1	0.89	0.83	0.43	0.77
n			147	213	260	217

Regression coefficients and model parameters are statistically significant at the * p<0.05, ** p<0.01, *** p<0.001 level.

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil

B. Kitzler et al.





Fig. 1. Extractable NH_4^+ (circles) and NO_3^- (triangles) in the litter layer and in the mineral soil at AK in the investigation years. Pooled samples (n=4) were taken from around the individual chambers.



Fig. 2. Bar chart: Biweekly N-input (kg N ha⁻¹) at AK measured in the two investigation years. Note: Dry deposition was measured only in the first year. Pie chart: Portion of throughfall (TF NH_4^+ , TF NO_3^-) and dry deposition (DRY NO_2 , DRY NH_3) on annual N-input (kg N ha⁻¹) in the first investigation year.

BGD

2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



2, 1423–1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil





Fig. 3. (a) Mean CO_2 emissions (squares±S.E) measured with the manual chambers and soil temperature [5 cm] (black line), (b) mean N_2O emissions from manual (circles±S.E) and automatic (diamonds) chambers, (c) mean NO emissions (±S.E) and, (d) daily precipitation and soil moisture [5 cm] at the study site AK from May 2002 to July 2004.



2, 1423-1455, 2005

N₂O, NO_x and CO₂ fluxes in a calcareous mountain forest soil



Fig. 4. Predicted versus measured N₂O and NO emissions at AK between April 2002 and July 2004. Independent variables are soil moisture and soil temperature for model 1 (M 1) and model 4 (M 4); N-deposition (NH_4^+ and NO_3^-) for model 2 (M 2) and, precipitation and air temperature for model 3 (M 3). 1455