



Competition for inorganic and organic forms of nitrogen and phosphorous between phytoplankton and bacteria during an *Emiliana huxleyi* spring bloom (PeECE II)

T. Løvdal, C. Eichner, H.-P. Grossart, V. Carbonnel, L. Chou, T. F. Thingstad

► To cite this version:

T. Løvdal, C. Eichner, H.-P. Grossart, V. Carbonnel, L. Chou, et al.. Competition for inorganic and organic forms of nitrogen and phosphorous between phytoplankton and bacteria during an *Emiliana huxleyi* spring bloom (PeECE II). *Biogeosciences Discussions*, 2007, 4 (5), pp.3343-3375. hal-00297923

HAL Id: hal-00297923

<https://hal.science/hal-00297923>

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

Competition for inorganic and organic forms of nitrogen and phosphorous between phytoplankton and bacteria during an *Emiliana huxleyi* spring bloom (PeECE II)

T. Løvdal^{1,*}, C. Eichner^{1,**}, H.-P. Grossart², V. Carbone³, L. Chou³, and
T. F. Thingstad¹

¹Department of Biology, University of Bergen, Jahnebakken 5, P.O. Box 7800, 5020 Bergen, Norway

²Leibniz Institute of Freshwater Ecology and Inland Fisheries, Department of Limnology of Stratified Lakes, Alte Fischehuette 2, 16775 Stechlin, Germany

³Laboratoire d'Océanographie Chimique et Géochimie des Eaux, Université Libre de Bruxelles, Campus Plaine – CP 208, 1050 Brussels, Belgium

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

*present address: Department of Mathematics and Natural Sciences, Faculty of Science and Technology, University of Stavanger, 4036 Stavanger, Norway

**present address: Institute of Marine Research, P.O. Box 1870 Nordnes, 5817 Bergen, Norway

Received: 18 September 2007 – Accepted: 19 September 2007 –

Published: 26 September 2007

Correspondence to: T. Løvdal (trond.lovdal@uis.no)

BGD

4, 3343–3375, 2007

**Algal-bacterial
competition for N and
P**

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

Abstract

Using ^{15}N and ^{33}P , we measured the turnover of organic and inorganic nitrogen (N) and phosphorus (P) substrates, and the partitioning of N and P from these sources into two size fractions of marine osmotrophs during the course of a phytoplankton bloom in a nutrient manipulated mesocosm. The larger size fraction ($>0.8\ \mu\text{m}$), mainly consisting of the coccolithophorid *Emiliania huxleyi*, but also including an increasing amount of large particle-associated bacteria as the bloom proceeded, dominated uptake of the inorganic forms NH_4^+ , NO_3^- , and PO_4^{3-} . The uptake of N from leucine, and P from ATP and dissolved DNA (dDNA), was initially dominated by the $0.8\text{--}0.2\ \mu\text{m}$ size fraction, but shifted towards dominance by the $>0.8\ \mu\text{m}$ size fraction as the system turned to an increasing degree of N-deficiency. Normalizing uptake to biomass of phytoplankton and heterotrophic bacteria revealed that organisms in the $0.8\text{--}0.2\ \mu\text{m}$ size fraction had higher specific affinity for leucine-N than those in the $>0.8\ \mu\text{m}$ size fraction when N was deficient, whereas the opposite was the case for NH_4^+ . There was no such difference regarding the specific affinity for P substrates. Since heterotrophic bacteria seem to acquire N from organic compounds like leucine more efficiently than phytoplankton, our results suggest different structuring of the microbial food chain in N-limited relative to P-limited environments.

1 Introduction

Under conditions of mineral nutrient limitation, heterotrophic bacteria may compete with phytoplankton for inorganic nutrients (e.g. orthophosphate (PO_4^{3-}), ammonium (NH_4^+), and nitrate (NO_3^-)). The outcome of this competition potentially influences the carbon cycle both by heterotrophic microbes indirectly limiting primary production by depriving phytoplankton of nutrients (Joint et al., 2002), and by phytoplankton indirectly limiting the extent of heterotrophic degradation of organic material (Havskum et al., 2003). Understanding how competition, predation, and other trophic interactions structure the

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

flows of C, N, P, and other elements through the microbial food web is central to our understanding of the role and function of this part of the pelagic ecosystem, both in a biological and in a biogeochemical context.

Competition between phytoplankton and bacteria potentially influences the species composition of the phytoplankton and bacterial communities (Jacobsen et al., 1995; Samuelsson et al., 2002). It may also affect the fundamental functioning of the microbial ecosystem by shifting the balance between phytoplankton and bacteria (Bratbak and Thingstad, 1985), as well as the bacterial degradation of organic matter (Pengerud et al., 1987; Grossart et al., 2006b). In principle, the competition between heterotrophic bacteria and phytoplankton for N and P may be different for the organic forms than it is for the inorganic ones. In addition, the case of N- and P-limitation may be different. Bacteria have traditionally been expected to be more superior in competition for dissolved organic N (DON) (reviewed by Antia et al., 1991), than they are for dissolved organic P (DOP), where both phytoplankton and bacteria are known to produce enzymes, such as alkaline phosphatase, splitting orthophosphate off from the organic part before uptake (reviewed by Chróst, 1990). If this is correct, simple mathematical models suggest that the microbial part of the C-cycle may differ substantially between N- and P-limited systems (Thingstad, 2000).

Although heterotrophic uptake by bacteria has long been recognized as the major process removing DON (e.g. as amino acids (Paul, 1983; Billen, 1984)), uptake of amino acids and other DON-compounds by phytoplankton can occur under several environmental conditions (Ietswaart et al., 1994; Pantoja and Lee, 1994; Palenik and Henson, 1997). Proteolytic activity has been found in association with eukaryote algae (Berges and Falkowski, 1996) and cyanobacteria (Martinez and Azam, 1993). One important question is whether cyanobacteria, being a diverse group of prokaryotic algae with cell envelope characteristics similar to those of their eubacterial counterparts (reviewed by Hoiczky and Hansel, 2000), behave more like heterotrophic bacteria or eukaryotic algae in their uptake of mineral nutrients. Measuring amino acid uptake rates in cyanobacteria, Zubkov et al. (2003), and Zubkov and Tarran (2005) found that

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

methionine, leucine and tyrosine could be an important source of N for *Prochlorococcus*, but not for *Synechococcus*. Apart from this, there has been little work to quantify the importance of DON to the nutrition of these organisms in the natural environment. Laboratory experiments leave no doubt that the majority of aquatic algal species are able to utilize common organic compounds as N sources for growth if sufficient substrate concentration is provided and enough time is allowed for metabolic adaptation (reviewed by Berman and Bronk, 2003). Most of these studies, however, used axenic laboratory batch cultures growing on high initial substrate concentrations, thus the ability of organisms to exploit the much lower concentrations of these compounds encountered in the environment in situ still remains unclear (Paul, 1983; Berman and Bronk, 2003).

This paper focuses on algal and bacterial uptake of different N- and P-substrates during an artificially induced phytoplankton bloom dominated by coccolithophorids, mainly *Emiliania huxleyi*. The experiment was done in a mesocosm set-up with a time dependent change from presumably C-stressed bacteria and P-stressed phytoplankton, to N-limitation, at least of autotrophic processes. Bacterial and algal uptake of inorganic N (NH_4^+ , NO_3^-) and of leucine as a model substrate for DON, as well as inorganic P (PO_4^{3-}) and the two organic forms ATP and DNA as model substrates for DOP was measured. Analyses of ^{15}N and ^{33}P uptake from these sources in algal and bacterial size fractions were performed in order to compare algal vs. bacterial competition for organic and inorganic dissolved N and P by means of their biomass-specific affinities.

2 Materials and methods

2.1 Experiment

An outdoor mesocosm experiment was carried out at the Marine Biological Field Station, Espeland, 20 km south of Bergen, western Norway from 4 to 24 May. The study was part of the Pelagic Ecosystem CO_2 Enrichment Study (PeECE). Experimental

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

setup and sampling procedures are described elsewhere (Grossart et al., 2006a). The data referred to here were obtained from enclosure number 4 referred to as “present” in the paper of Grossart et al. (2006a) because it represented the present-day level of 370 ppmV atmospheric CO₂ concentration. The enclosure (volume ~20 m³) was filled with unfiltered, nutrient-poor, post-spring bloom sea water from the fjord and supplemented with 9 μmol L⁻¹ NO₃⁻, 0.5 μmol L⁻¹ PO₄³⁻, and 12 μmol L⁻¹ Si(OH)₄ on day 0 of the experiment to induce a phytoplankton bloom.

Stormy weather from day 4 eventually led to a collapse of the enclosure on day 6. The enclosure was restored to an upright position, and 705 mg of KH₂PO₄ was added to the enclosure on day 7 to maintaining the phytoplankton bloom. The addition of KH₂PO₄ would ideally correspond to a PO₄³⁻ enrichment corresponding to a concentration of 0.3 μmol L⁻¹ PO₄³⁻, but was actually slightly higher since the enclosure was not entirely unfolded at the time. From day 10 on, the weather calmed down, and the enclosure was underlayered with an unknown volume of deep, high salinity water (with a soluble reactive phosphorus (SRP) concentration of 0.77 μmol L⁻¹) in order to unfold the enclosure after the storm.

2.2 Chemical analysis

SRP was measured using the molybdate blue method (Koroleff, 1983). Nitrate and nitrite (NO_x) were determined with a Skalar AutoAnalyser, based on the sulphanilamide colorimetric method (Grasshoff, 1983). Ammonium (NH₄⁺) was analysed by the indophenol blue method according to Koroleff (1969). Dissolved free amino acids (DFAA) were analyzed by High Pressure Liquid Chromatography (HPLC) after ortho-phthaldialdehyde derivatization as described elsewhere (Grossart et al., 2006a).

2.3 Biomass estimations

Free and particle-associated bacteria were enumerated and their volumes calculated as described elsewhere (Grossart et al., 2006a). C, N, and P biomass of free bacteria

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

was calculated assuming a constant biomass of $18.5 \text{ fg C cell}^{-1}$ and a C:N:P molar ratio of 50:10:1 for coccoid bacteria (Fagerbakke et al., 1996). Particle-associated bacteria were significantly bigger than free bacteria, and increased from 0.6 to $1.5 \mu\text{m}^3$ at the end of the experiment (Grossart et al., 2006a). Because of the large and temporally variable size of the particle-associated bacteria, allometric conversion factors between volume (V) and C, N, and P content (in femtograms) of $C=220 \times V^{0.9}$, $N=50 \times V^{0.8}$, and $P=2 \times V^{0.7}$ was applied. The bacterial volume to C conversion factor is comparable to a factor ($C=218 \times V^{0.86}$) for both cultures and bacterioplankton from lakes of different trophic status (Loferer-Kröbächer et al., 1998) and to a factor ($C=224 \times V^{0.89}$) for *Vibrio splendidus* grown along a gradient from C to P limitation (Løvdal et al., 2007a). The volume to N and volume to P conversion factors are adapted from Løvdal et al. (2007a).

Chlorophyll *a* (Chl *a*) measurements (Grossart et al., 2006a) were used for estimating phytoplankton C, N, and P biomass. A C:Chl *a* ratio of 30 (w:w) and a C:N:P molar ratio of 106:16:1 in phytoplankton biomass was assumed. With the conversion factors applied here, we got an estimated “bacterial + algal C-biomass” to Chl *a* ratio (w:w) of 85 ± 22 (mean \pm SD, $n=10$), which is close to the reported constant particulate organic C (POC):Chl *a* ratio (w:w) of 81 (Aure et al., 2000, as cited in Erga et al., 2005) from the waters of the Samnanger Fjord. This site constitutes the fjord system adjacent to the study site.

2.4 Alkaline phosphatase activity (APA)

APA was measured fluorometrically using 3-0-methylfluorescein-phosphate (MFP) as substrate (Perry, 1972). Samples were mixed with MFP solution in 0.1 mol L^{-1} Trizma-HCl pH 8.3 (final concentration $0.1 \mu\text{mol L}^{-1}$). Fluorescence was measured directly after the addition of the reagent and at two subsequent times according to the expected activity using a Perkin Elmer fluorometer LS50B. After correcting fluorescence values of samples to those of autoclaved samples used as blanks, APA ($\text{nmol-P L}^{-1} \text{ h}^{-1}$) was calculated using linear regression of fluorescence values versus incubation time.

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

2.5 Uptake of ^{15}N -compounds

Water for use in ^{15}N uptake studies was collected with a tube covered with $18\text{ }\mu\text{m}$ gaze under stirring into thoroughly cleaned 1000 mL septum glass bottles. ^{15}N enriched NH_4Cl , NaNO_3 , or L-leucine ($>98\text{ atom}\% ^{15}\text{N}$; Larodan Fine Chemicals AB), respectively, in a concentration of about 10% of the ambient concentrations was added. When the ambient NH_4^+ concentrations were below the measurement limit, unlabelled NH_4Cl was added to a final concentration of $0.5\text{ }\mu\text{mol L}^{-1}$ 1 to 4 h before tracer addition. The precise portion of tracer added was calculated after determination of the ambient NH_4^+ , NO_3^- , or DFAA (Grossart et al., 2006a) concentrations. The bottles were closed gas-tight and were incubated at in situ temperature and light. Four time points were taken within 3 to 5 h for each measurement. The reaction was terminated by filtering the samples through silver-membrane filters (Osmonics; pre treated 500°C for 1 h) with pore sizes of 0.2 and $0.8\text{ }\mu\text{m}$. The filtration volume was 400 to 500 mL. Filters were flushed with $0.2\text{ }\mu\text{m}$ filtered sea water to remove adherent tracer-containing water and stored frozen until analysis.

After drying in a drying chamber, the filters were wrapped in silver cups ($6\times6\times12\text{ mm}$; Elementar Analysensysteme) and formed into pellets with a laboratory press. As the amount of N on the $0.2\text{ }\mu\text{m}$ filters was under the measurement limit, $\delta^{15}\text{N}$ values were measured together with a well defined N isotope standard (peptone; Merck). One $\mu\text{mol L}^{-1}$ peptone was pipetted onto the $0.2\text{ }\mu\text{m}$ filters before packing and N isotopes were measured by continuous flow isotope ratio mass spectrometry (CF-IRMS) on a Finnigan MAT Delta plus coupled with a Thermo NA 2500 CHN analyser. The N content was calculated using acetanilide (Fisons) as a standard. The standard deviation of the $\delta^{15}\text{N}$ values measured for the standards was on average 0.15‰ . The stable N isotope ratio was calculated in terms of $\delta^{15}\text{N}$ -values as follows:

$$\delta^{15}\text{N} = \left(\frac{R_{\text{sample}} - R_{\text{standard}}}{R_{\text{standard}}} \right) \times 1000[\text{‰}]$$

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

where $R = \frac{^{15}\text{N}}{^{14}\text{N}}$. The measured $\delta^{15}\text{N}$ values were converted to atom percent ^{15}N after the formula (Montoya et al., 1996):

$$\text{atom}\%^{15}\text{N} = 100 \times \left[\frac{(10^{-3} \times \delta^{15}\text{N} + 1) \times (^{15}\text{N}/^{14}\text{N})_{\text{atmosphere}}}{1 + (10^{-3} \times \delta^{15}\text{N} + 1) \times (^{15}\text{N}/^{14}\text{N})_{\text{atmosphere}}} \right]$$

with $\left(\frac{^{15}\text{N}}{^{14}\text{N}}\right)_{\text{atmosphere}} = 0.003676$ (Junk and Svec 1958). The uptake rates were estimated from the regression relationship between uptake and time and converted to turnover time (T ; see below).

2.6 ^{33}P -labelling of DNA

Radiolabeling of DNA was performed by random oligonucleotide primed synthesis (ROPS) with the DecaLabel DNA labeling kit (Fermentas K0621) in accordance with the manufacturer's instructions. [^{33}P]DNA was prepared for use in uptake studies as described by Løvdaal et al. (2007b). This procedure yields labelled DNA products with an average length of 0.5 kilobase pairs (kb) which is at the lower end of the range for naturally occurring dDNA (0.12–35.2 kb) in aquatic environments (DeFlaun et al., 1987). We did not correct for potential shortening of the DNA chain length by radiochemical decay, but the radiolabelled DNA was used well within one half life of the radioactive precursor in order to avoid significant shortening. The final product had a specific activity of approximately 10^8 counts per minute (cpm) μg^{-1} with $97.0 \pm 0.6\%$ (mean \pm SD of 3 replicates) incorporation of label and a concentration of $12.6 \pm 0.1 \text{ ng } \mu\text{L}^{-1}$ (mean \pm SD of 3 replicates) as calculated by the DE-81 filter-binding assay (Sambrook and Russell, 2001).

2.7 Uptake of ^{33}P -compounds

Uptake of $^{33}\text{PO}_4^{3-}$, AT^{33}P and [^{33}P]DNA was measured according to Thingstad et al. (1993), modified as described by Løvdaal et al. (2007b), except that the samples

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdaal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

were not assayed for inorganic P released from organic substrates which was not taken up by the organisms. All incubations were done in 10 mL subsamples in 15 mL Falcon tubes at subdued light and $15.5 \pm 1.0^\circ\text{C}$. Samples were incubated according to the expected turnover time; for samples incubated with $^{33}\text{PO}_4^{3-}$, AT^{33}P and $[^{33}\text{P}]\text{DNA}$, the respective incubation times varied between 15 min and 1 h, 1–2 h, and 2–5 h, respectively. Incubations were stopped by cold chase (Løvdaal et al., 2007b). Bacteria and phytoplankton were separated into different size fractions by filtration onto polycarbonate filters (Poretics) with pore sizes of 0.2 and $0.8\ \mu\text{m}$ according to Løvdaal et al. (2007b). Subsamples and filters were radioassayed by liquid scintillation counting (Løvdaal et al., 2007b).

2.8 Estimation of turnover times and biomass-specific affinity

Turnover times (T ; h) were calculated by the equation (Thingstad et al., 1993):

$$T = \frac{t}{-\ln(1 - R)}$$

where R is the consumed fraction of added label and t is the incubation time. Hence, T represents here the turnover of substrates into particulate matter and does not include the release of hydrolysis products to the water phase. T for ^{33}P substrates was calculated from one time-point measurements (Thingstad et al., 1993), whereas T for ^{15}N substrates was calculated from the regression line between four time points.

Biomass-specific affinity for N (α_N ; $\text{L nmol-N}^{-1} \text{h}^{-1}$) and P (α_P ; $\text{L nmol-P}^{-1} \text{h}^{-1}$) uptake was estimated according to the procedure proposed by Thingstad and Ras-soulzadegan (1999):

$$\alpha = f / (TB) \quad (1)$$

where f is the fraction of uptake in the respective size fraction and B is the biomass-N (nmol-N L^{-1}) or biomass-P (nmol-P L^{-1}). The biomass of particle-associated bacteria was contributed to the $>0.8\ \mu\text{m}$ size fraction because they could be trapped on a

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdaal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

5 $5\ \mu\text{m}$ filter (Grossart et al., 2006a). Hence, due to the particle-associated bacteria, the mechanical separation of bacteria and phytoplankton by size fractionation was not entirely successful. In our estimates of biomass-specific affinity, this was corrected for by assuming that particle-associated bacteria had the same affinity for uptake as free
 5 bacteria. From this assumption, the fraction of uptake by particle-associated bacteria was calculated by rearranging Eq. (1) to

$$f = \alpha TB \quad (2)$$

inserting the estimated biomass of particle associated bacteria. Corrected estimates corresponding to algae and bacteria was then calculated by subtracting this value from
 10 the $>0.8\ \mu\text{m}$ size fraction, contributing it to the $0.8\text{--}0.2\ \mu\text{m}$ size fraction, respectively.

2.9 Statistical analysis

Statistical analysis was performed by student's t -tests according to Sokal and Rohlf (1995). The confidence level of all analyses was set at 95%.

3 Results

15 3.1 Nutrients, alkaline phosphatase activity (APA), and bloom development

Dissolved inorganic N (DIN), represented as the sum of $\text{NH}_4^+ + \text{NO}_x$ was dominated by NO_x concentrations ranging from $8\ \mu\text{mol L}^{-1}$ on day 1, to $4\ \mu\text{mol L}^{-1}$ on day 7, before it rapidly declined to under the detection limit ($<0.01\ \mu\text{mol L}^{-1}$) on day 16, whereas NH_4^+ concentrations were low ($<0.02\ \mu\text{mol L}^{-1}$) the whole study period. SRP concentrations
 20 declined from approximately $0.4\ \mu\text{mol L}^{-1}$ on day 1 to $\sim 0.04\ \mu\text{mol L}^{-1}$ on day 20. Thus, the DIN:SRP ratio was well above the Redfield ratio of 16 beyond the second addition of PO_4^{3-} on day 7 before it rapidly declined (Fig. 1). DFAA concentrations ranged

between 0.4 and 1.6 $\mu\text{mol L}^{-1}$, with the lowest concentrations after the peak of the bloom (Grossart et al., 2006a).

Alkaline phosphatase activity (APA) ranged from 1 $\text{nmol-P L}^{-1} \text{h}^{-1}$ on day 2 of the experiment, to 5 $\text{nmol-P L}^{-1} \text{h}^{-1}$ on day 20. Specific values for APA (S-APA) and for phosphate affinity ($S\text{-}\alpha_{\text{PO}_4}$) normalized for the summed P-biomass of phytoplankton and bacteria are shown in Fig. 2. Both values increased after day 8, but remained below 0.04 h^{-1} and 0.008 $\text{L nmol-P}^{-1} \text{h}^{-1}$, respectively. According to Tanaka et al. (2006), S-APA above 0.2 h^{-1} or $S\text{-}\alpha_{\text{PO}_4}$ above 0.02 $\text{L nmol-P}^{-1} \text{h}^{-1}$, is indicative of systemic P-limitation. According to these criteria, our data on S-APA and $S\text{-}\alpha_{\text{PO}_4}$ (Fig. 2) do not indicate severe P-limitation during any parts of the study period.

The algal bloom, almost exclusively dominated by coccolithophorids (mainly *Emiliania huxleyi*), was initiated by the addition of inorganic nutrients on day 0 and reached its maximum on day 12 (Grossart et al., 2006a) where the DIN:SRP ratio was ~ 5 ($\text{NO}_x\text{:SRP} \sim 2$). Highest numbers of total as well as free bacteria were recorded on day 8 (Grossart et al., 2006a) when the DIN:SRP ratio was ~ 11 and NO_x accounted for 99% of DIN. Particle-associated bacteria contributed to 40–80% of the total bacterial volume (Grossart et al., 2006a). From our estimates, the contribution of particle-associated bacteria to the total C-biomass of algae and bacteria increased from $\sim 30\%$ on day 0 to $\sim 65\%$ on day 20.

3.2 Turnover times

During the first week of the experiment, T of all substrates were in general long and T of the organic N and P substrates measured were shorter than T of the respective inorganic substrates (Fig. 3). This may indicate C-limitation of heterotrophic processes in this phase (see discussion). The peak of the phytoplankton bloom on day 12 coincided with the shortest T_{NH_4} (4.8 h) and T_{NO_3} (6.2 h). From this point on, NH_4^+ and NO_x concentrations were both in the nanomolar range and T for NH_4^+ was measured as the only DIN substrate. T_{NH_4} stayed stable below 10 h throughout the study period,

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

whereas T_{Leucine} decreased from >100 h to 34 h at the same time (Fig. 3a).

Despite high DIN:SRP ratios (Fig. 1), which could have been interpreted to indicate phosphate limitation, T_{PO_4} was longer than T_{ATP} at the beginning of the experiment (Fig. 3b). Because P limitation was not expected immediately after the addition of PO_4^{3-} on day 7, T for P substrates was not measured between day 6 and day 12, the peak of the phytoplankton bloom. On day 12, T_{PO_4} (5.8 h) was shorter than T_{ATP} and was comparable to T_{NH_4} and T_{NO_3} , before it reached a minimum of 1.4 h at day 16 (Fig. 3b). T_{ATP} and T_{dDNA} declined from 24 and 346 h, respectively, on day 12 before it stabilized on 6 ± 2 h (mean \pm SD, $n=$) and 84 ± 8 h (mean \pm SD, $n=4$), respectively, for the rest of the study period (Fig. 3b).

3.3 Uptake distribution of ^{15}N - and ^{33}P -substrates and biomass-specific affinity

The distribution of added activity taken up by the two size fractions is shown in Fig. 4. The 0.8–0.2 μm size fraction took up most of the organic substrates during the first part of the study period, whereas the >0.8 μm size fraction dominated uptake of the inorganic substrates. In our set of data, both T_{NH_4} and T_{PO_4} dropped below 10 h at the peak of the phytoplankton bloom (Fig. 3). There are strong indications for a transition from a non-N-limited to an N-limited system at this point (see Discussion). Hence, we used T_{NH_4} as an indicator to split the dataset into non-N-limitation ($T_{\text{NH}_4} > 10$ h) and N-limitation ($T_{\text{NH}_4} < 10$ h) which also represents the periods before and after the peak of the phytoplankton bloom, respectively. After the peak of the phytoplankton bloom, the >0.8 μm size fraction dominated uptake of all substrates, and NH_4^+ uptake was almost exclusively by the >0.8 μm size fraction. There were small changes in the distribution of inorganic N and P compared to before the phytoplankton peak, with the >0.8 μm size fraction taking up slightly more after this peak (Fig. 4).

Table 1 is a compilation of data presented in Fig. 4 showing uptake distributions to the >0.8 μm size fraction grouped for the two periods. NO_3^- and NH_4^+ data are pooled as DIN, whereas ATP and dDNA data are pooled as DOP. From DON (i.e. leucine-N) and DOP uptake dominated by the 0.8–0.2 μm size fraction during $T_{\text{NH}_4} > 10$ h, there

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

was a shift towards DON and DOP uptake dominated by the $>0.8\ \mu\text{m}$ size fraction during $T_{\text{NH}_4} < 10\ \text{h}$ (Table 1). There were no drastic shifts in the distribution of inorganic substrates, with the $>0.8\ \mu\text{m}$ size fraction dominating uptake in both periods, but more so during $T_{\text{NH}_4} < 10\ \text{h}$ (Table 1).

5 $S\text{-}\alpha_{\text{NO}_3}$ and $S\text{-}\alpha_{\text{NH}_4}$, and $S\text{-}\alpha_{\text{PO}_4}$ are specific affinities normalized for the summed N and P biomass, respectively, of algae and bacteria, and are shown in Fig. 2. Affinities for all substrates were low during the first half of the study period. Maximum values of $S\text{-}\alpha_{\text{NH}_4}$ paralleled the peak of the phytoplankton bloom, whereas $S\text{-}\alpha_{\text{PO}_4}$ peaked on day 16.

10 Comparing competitive ability of osmotrophs in the $0.8\text{--}0.2\ \mu\text{m}$ and the $>0.8\ \mu\text{m}$ size fraction by means of biomass-specific affinity for N-uptake in the two size fractions, focusing on the N-limited phase ($T_{\text{NH}_4} < 10\ \text{h}$), reveals a significant shift depending on whether N is present in organic or inorganic form (Fig. 5). The $>0.8\ \mu\text{m}$ size fraction had a significantly higher biomass-specific affinity for NH_4^+ than the $0.8\text{--}0.2\ \mu\text{m}$ size fraction ($p=0.005$), and a significantly lower biomass-specific affinity for leucine-N than the $0.8\text{--}0.2\ \mu\text{m}$ size fraction ($p=0.003$) (Figs. 5a–b). No such shift between the competition for DIP and DOP were observed; no significant differences were found between biomass-specific affinity in the two size fractions for PO_4^{3-} ($p=0.449$), ATP-P ($p=0.372$), or dDNA-P ($p=0.494$) (Figs. 5c–e).

20 From Eq. (2), it was estimated that much of the uptake in the $>0.8\ \mu\text{m}$ size fraction was by particle-associated bacteria. Particle-associated bacteria took up larger shares of DON compared to DIN (Table 2). In fact, it was estimated that the majority ($73\pm 6\%$, $n=5$) of total DON uptake was by particle-associated bacteria during N-limitation ($T_{\text{NH}_4} < 10\ \text{h}$), leaving phytoplankton with $<2\%$ (Table 2), and thus free-living bacteria with approximately 25% of total DON uptake. Hence, from the corrected estimates, phytoplankton had a significantly lower biomass-specific affinity for DON than heterotrophic bacteria ($p<0.0001$) (Fig. 5b) but a significantly higher biomass-specific affinity for NH_4^+ ($p<0.0001$) (Fig. 5a). No shift between the competition for DIP and DOP were observed; no significant differences were found between biomass-specific affinity

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

in algae and bacteria for PO_4^{3-} ($p=0.324$), ATP-P ($p=0.237$), or dDNA-P ($p=0.407$) (Figs. 5c–e).

4 Discussion

It has been hypothesized that phosphate limitation, classically indicated by DIN:DIP ratios >16 , is one of the critical factors allowing the coccolithophore *Emiliania huxleyi* to bloom (reviewed by Lessard et al., 2005). This hypothesis is based on physiological studies showing that *E. huxleyi* has an exceptionally high affinity for PO_4^{3-} and is able to use DOP (Kuenzler and Perras, 1965; Riegman et al., 2000). High $\text{NO}_3^-:\text{PO}_4^{3-}$ ratios, however, appear to be the exception rather than the rule in *E. huxleyi* blooms. In fact, in most blooms studied to date, $\text{NO}_3^-:\text{PO}_4^{3-}$ ratios were low, and nitrate was low or undetectable (Lessard et al., 2005). In the current experiment, the addition of NO_3^- and PO_4^{3-} in a molar ratio of 18 initiated a bloom of coccolithophorids, dominated by *E. huxleyi*, reaching its maximum after 12 days when the DIN:SRP ratio was ~ 5 . The first half of the period prior to the peak of the bloom was characterized by high DIN:SRP ratios (Fig. 1), thus the phytoplankton growth may be suspected to have been restricted by the availability of inorganic P. Low S-APA and low $\text{S-}\alpha_{\text{PO}_4}$, $\text{S-}\alpha_{\text{NO}_3}$ and $\text{S-}\alpha_{\text{NH}_4}$ values (Fig. 2), however, does not indicate severe mineral nutrient limitation, and T of the organic N and P substrates measured were shorter than T of the respective inorganic substrates (Fig. 3), possibly indicating C-limitation of heterotrophic processes in this phase. Bacteria hydrolyze specific DON and DOP-compounds in the presence of inorganic mineral nutrients, presumably to obtain other associated mineral nutrients and C (Jørgensen et al., 1993; Benitez-Nelson and Buesseler, 1999). DIN:SRP ratios rapidly declined from day 7 (Fig. 1) due to the PO_4^{3-} enrichment. However, the additional PO_4^{3-} was rapidly consumed followed by increasing values of S-APA and $\text{S-}\alpha_{\text{PO}_4}$ (Fig. 2). $\text{S-}\alpha_{\text{NO}_3}$ and $\text{S-}\alpha_{\text{NH}_4}$ values rapidly increased from day 8 with $\text{S-}\alpha_{\text{NH}_4}$ peaking at day 12 (Fig. 2). Additionally, T_{NO_3} and T_{NH_4} got shorter as the experiment proceeded, with

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

T_{NH_4} reaching minimum values at day 10 before stabilizing, and T_{Leucine} declining from day 10 to the end (Fig. 3). Conclusively, the splitting of the investigation period into a non-N-limited and an N-limited phase can be justified.

From our data, there is evidence for a shift in terms of algal – bacterial competition when N is available in the form of leucine contrary to NH_4^+ (Fig. 5), with organisms in the $>0.8\ \mu\text{m}$ size fraction having significantly lower biomass-specific affinity for leucine-N and significantly higher biomass-specific affinity for NH_4^+ compared to smaller organisms ($0.8\text{--}0.2\ \mu\text{m}$). Correcting for the contribution of particle-associated bacteria in the $>0.8\ \mu\text{m}$ size fraction by assuming they have similar affinity per biomass as free-living bacteria, probably represents a best estimate for the competition between algae and heterotrophic bacteria. However, since specific aminopeptidase (Karner and Herndl, 1992; Middelboe et al., 1995), protease (Becquevort et al., 1998) and phosphatase (Smith et al., 1992; Simon et al., 2002) activities in particle-associated bacteria is often higher than those characteristic for free-living bacteria, our affinity values for heterotrophic bacteria may have been underestimated, and the values for phytoplankton correspondingly overestimated.

In a previous investigation in P-limited estuarine mesocosms, Løvdal et al. (2007b) found no significant shift in algal – bacterial competition for P from ATP and dDNA relative to PO_4^{3-} , in terms of biomass-specific affinity. In the current experiment, using comparable methodology, a shift from bacteria dominating the competition for DON towards algae dominating the competition for DIN was evident (Fig. 5). This does indeed indicate that the structure of the microbial food web in N-limited environments may be different from that in P-limited environments. If amino acids are utilized almost exclusively by bacteria, this changes the food web structure relative to P-limited environments, since there will be a pool of dissolved N for which bacteria do not experience significant competition from phytoplankton to achieve. One effect of this is, theoretically, that the probability for C-limitation of bacteria could be larger in N-deficient regions than in P-deficient regions (Thingstad, 2000).

E. huxleyi has been shown to be a moderate competitor for nitrate and a good com-

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

petitor for phosphate compared to other algal species (Riegman et al., 1992, 2000). During natural *E. huxleyi* blooms, *E. huxleyi* primarily use NH_4^+ and urea (Kristiansen et al., 1994; Fernández et al., 1996; Rees et al., 2002). The high P acquisition capacity and ability to use non-nitrate N has been suggested to (Lessard et al., 2005), at least in part, explain the success of *E. huxleyi* in nutrient-depleted waters. This is supported by our data because the biomass-specific affinity for all P-substrates in *E. huxleyi* was comparable to that in heterotrophic bacteria, whereas *E. huxleyi*, although they seemed to have low amino acid-N acquisition capacity, dominated the competition for NH_4^+ . Our results also agree with previous findings that DFAA are an important N source to marine bacteria (Hollibaugh and Azam, 1983; Jørgensen et al., 1993).

The biomass-specific affinity values estimated in the present study can be compared with those given by diffusion limitation of substrate transport to the cell surface – that is, the theoretical maximum. Assuming that the cell is diffusion-limited, i.e., that the cell's uptake system is so efficient (and the bulk nutrient concentration so low) that all substrate molecules hitting the cell surface are captured, it is possible to derive a theoretical expression for maximum specific affinity (α_{max}) for a spherical cell (Fig. 6; see also Thingstad and Rassoulzadegan, 1999):

$$\alpha_{\text{max}} = 3D/(\sigma r^2)$$

where D is the diffusion constant for the substrate molecules and σ is the volume-specific content of the element in question. Assuming free-living and particle-associated bacteria to have volumes of 0.2 and $1.5 \mu\text{m}^3$, respectively, and particle-associated bacteria to make up 30% of the total bacterial community (cf. Grossart et al., 2006a), their mean equivalent r can be calculated to be $\sim 0.5 \mu\text{m}$. Assuming the phytoplankton community is dominated by N-limited *E. huxleyi* with volumes of $\sim 40 \mu\text{m}^3$ (cf. Riegman et al., 2000), their mean r may be set to $2.1 \mu\text{m}$. One should however keep in mind that genetic and physiological intraspecific diversity in *E. huxleyi* has been documented, and multiple strains can be present within an environment (Paasche, 2002), complicating comparison between laboratory and field studies. The *E. huxleyi* (L-strain) used as a model organism by Riegman et al. (2000) was cultivated

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

in continuous cultures and was more than three times smaller and had lower internal N and P cell quotas than *E. huxleyi* reported elsewhere (Ho et al., 2003). Thus, the results of their study cannot be applied directly to the natural environment. Hence, the cell radius applied here and the lines representing the theoretical maximum for biomass-specific affinity in *E. huxleyi* (Fig. 6) is to be understood as extremes.

The mean biomass-specific affinity for NH_4^+ uptake in phytoplankton was higher than that for bacteria, approaching that predicted by the diffusion model (Fig. 6a), contrary to that estimated for leucine (Fig. 6b). Although *E. huxleyi* is reported to grow fairly on leucine as the sole N-source (Ietswaart et al., 1994), our data suggest that the natural amount of leucine does not serve as an important N-source for *E. huxleyi* in our study.

The reason for the low biomass-specific affinity for N-substrates estimated for bacteria (Figs. 6a–b) is unknown. One possibility, contradictory to our previous conclusion, could be that the growth rate of heterotrophic bacteria was limited by C or other factors, rather than N. Another possibility could be artefacts in our estimation procedure. However, the main reason that biomass-specific affinities for N-compounds, both in bacteria and algae, seem unrealistically low for a presumably N-stressed system, compared to that of P-compounds, stems probably from a methodological limitation. Although the continuous flow isotope ratio mass spectrometry (CF-IRMS) is very sensitive, it requires a certain threshold level of particulate material for the analysis of microbial N uptake. Hence, following long filtration time due to large sample volumes, this technique only allows for analysis of relatively high levels of isotope enrichment to avoid exhaustion of the added tracer. Thus, it is not optimal in environmental labelling studies when the ambient turnover time is extremely short. In our experiment, effort was done to add below 10% of ambient concentrations of tracer ^{15}N isotope. In the case of NH_4^+ uptake studies, this was occasionally achieved only by adding additional unlabelled NH_4Cl to the incubation bottles. During the N-limited phase, this enrichment could increase the DIN:SRP ratio in the incubation bottles by a factor of up to 10 compared to the ambient concentrations (Fig. 1), but the DIN:SRP ratio in the incubation bottles was still below 16. We acknowledge that this treatment presumably relaxed

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

the N-limitation in these bottles, and led to significant overestimates of the ambient NH_4^+ turnover times (Fig. 3a). Additions of ^{15}N -leucine were based on measurements of ambient DFAA concentrations. Most likely, the DFAA pool measured chemically is larger than that utilized by osmotrophs since both bacteria and phytoplankton prefer certain amino acids (including the neutral leucine) over others (Ietswaart et al., 1994). Hence, in the N-limited phase, ^{15}N enrichment presumably exceeded that typically regarded as true tracer levels, leading to an overestimate of turnover times, and hence underestimates of biomass-specific affinities for N-compounds (Figs. 5a–b). The liquid scintillation counting technique on the other hand, applied for ^{33}P uptake studies, is not hampered with this problem because the high specific activity of ^{33}P allows for true tracer level enrichment, e.g. picomolar concentrations. Although this problem with the CF-IRMS technique can be expected to have given significant underestimates of biomass-specific affinity, we believe that it did not affect the relative distribution of N-substrates in the two size fractions considerably. Thus, qualitative differences between the 0.8–0.2 μm and the >0.8 μm size fractions were attributed to the differences between bacterial and algal metabolism.

DNA and ATP contain 16% and 14% N, and 10% and 18% P, respectively. Hence, they contain as much N per dry weight as amino acids of algal proteins (Laws, 1991). Therefore these substrates may serve as potential N and P sources at the same time. The high biomass-specific affinities for dDNA and ATP, considering their diffusion constants, compared to PO_4^{3-} (Figs. 6c–e), may indicate that these substrates were hydrolyzed for their N content rather than their P content. dDNA is mainly utilized as a P source by marine bacteria, and accounts <10 of their N and C requirements (Jørgensen et al., 1993; Jørgensen and Jacobsen, 1996). The possibility then exists, that organic compounds, such as dDNA, are more bioavailable to phytoplankton than DON-compounds, such as amino acids, and are thus utilized as alternative sources for N by phytoplankton when the ambient DIN-concentrations are too low to support their growth. The potential for dDNA and ATP to also have rapid turnover times in non-P-limited environments, and to substantially support algal N-demand, indicates that phy-

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

toplankton may rely on these compounds to support their N-demand, rather than those traditionally looked upon as important DON-sources. Yet, isotope P-labelled substrates may yield limited information about N uptake. Therefore, an N-isotope is preferable in assessing phytoplankton dDNA and ATP utilization.

The main conclusion of our study is that the competitive fitness of bacteria and algae for nutrients (and particularly N) vary dramatically depending on whether the substrates are organic or inorganic. Whereas bacteria appear to be superior competitors for organic forms of N, phytoplankton appear to be superior competitors for inorganic N supplied as NH_4^+ . This observation has major implications for our understanding of marine food webs and effects of nutrient status on community structure.

Acknowledgements. We thank the staff of the Large Scale Facility (LSF) in Bergen and U. Riebesell for coordinating the project. S. Trimborn is thanked for Chl *a* measurements, and E. Foss Skjoldal, T. Tanaka, and M. Allgaier for technical assistance. U. Struck at the Museum für Naturkunde Berlin (Germany) is acknowledged for stable isotope analysis. This work was supported by the University of Bergen and the Research Council of Norway; project 158936/I10 Patterns in Biodiversity.

References

- Antia, N. J., Harrison, P. J., and Oliveira, L.: The role of dissolved organic nitrogen in phytoplankton nutrition, cell biology and ecology, *Phycologia*, 30, 1–89, 1991.
- Aure, J., Erga, S. R., and Asplin, L.: Increased biological production in fjords by artificial upwelling, *Fisken Havet*, 11, 1–30, 2000.
- Becquevort, S., Rousseau, V., and Lancelot, C.: Major and comparable roles for free-living and attached bacteria in the degradation of *Phaeocystis*-derived organic matter in Belgian coastal waters of the North Sea, *Aquat. Microb. Ecol.*, 14, 39–48, 1998.
- Benitez-Nelson, C. R. and Buesseler, K. O.: Variability of inorganic and organic phosphorus turnover rates in the coastal ocean, *Nature*, 398, 502–505, 1999.
- Berges, J. A. and Falkowski, P. G.: Cell-associated proteolytic enzymes from marine phytoplankton, *J. Phycol.*, 32, 566–574, 1996.

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

- Berman, T. and Bronk, D. A.: Dissolved organic nitrogen: a dynamic participant in aquatic systems, *Aquat. Microb. Ecol.*, 31, 279–305, 2003.
- Billen, G.: Heterotrophic utilization and regeneration of nitrogen, in: *Heterotrophic activity in the sea*, edited by: Hobbie, J. E. and Williams, P. J. L. B., Plenum Press, New York, London, 313–355, 1984.
- Bratbak, G. and Thingstad, T. F.: Phytoplankton-bacteria interactions: an apparent paradox? analysis of a model system with both competition and commensalism, *Mar. Ecol. Prog. Ser.*, 25, 23–30, 1985.
- Chróst, R. J.: Microbial ectoenzymes in aquatic environments, in: *Aquatic microbial ecology: Biochemical and molecular approaches*, edited by: Overbeck, J. and Chróst, R. J., Springer Verlag, 47–78, 1990.
- DeFlaun, M. F., Paul, J. H., and Jeffrey, W. H.: Distribution and molecular weight of dissolved DNA in subtropical estuarine and oceanic environments, *Mar. Ecol. Prog. Ser.*, 38, 65–73, 1987.
- Diehl, H., Ihlefeld, H., and Schwegler, H.: *Physik für Biologen*, Springer-Verlag, 391, 1991.
- Erga, S. R., Aursland, K., Frette, Ø., Hamre, B., Lotsberg, J. K., Stamnes, J. J., Aure, J., Rey, F., and Stamnes, K.: UV transmission in Norwegian marine waters: controlling factors and possible effects on primary production and vertical distribution of phytoplankton, *Mar. Ecol. Prog. Ser.*, 305, 79–100, 2005.
- Fagerbakke, K. M., Heldal, M., and Norland, S.: Content of carbon, nitrogen, oxygen, sulfur and phosphorus in native aquatic and cultured bacteria, *Aquat. Microb. Ecol.*, 10, 15–27, 1996.
- Fernández, E., Marañón, E., Harbour, D. S., Kristiansen, S., and Heimdahl, B. R.: Patterns of carbon and nitrogen uptake during blooms of *Emiliania huxleyi* in two Norwegian fjords, *J. Plankton Res.*, 18, 2349–2366, 1996.
- Grasshoff, K.: Determination of nitrate, in: *Methods of seawater analysis*, edited by: Grasshoff, K., Erhardt, M., and Kremling, K., Verlag Chemie, Weinheim, 143–150, 1983.
- Grossart, H.-P., Allgaier, M., Passow, U., and Riebesell, U.: Testing the effect of CO₂ concentration on the dynamics of marine heterotrophic bacterioplankton, *Limnol. Oceanogr.*, 51, 1–11, 2006a.
- Grossart, H.-P., Czub, G., and Simon, M.: Specific interactions of planktonic algae and bacteria: Implications for aggregation and organic matter cycling in the sea, *Environ. Microbiol.*, 8, 1074–1084, 2006b.
- Havskum, H., Thingstad, T. F., Scharek, R., Peters, F., Berdalet, E., Sala, M. M., Alcaraz, M.,

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdaal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

- Bangsholt, J. C., Zweifel, U. L., Hagström, Å., Perez, M., and Dolan, J. R.: Silicate and labile DOC interfere in structuring the microbial food web via algal-bacterial competition for mineral nutrients: Results from a mesocosm experiment, *Limnol. Oceanogr.*, 48, 129–140, 2003.
- Ho, T.-Y., Quigg, A., Finkel, Z. V., Milligan, A. J., Wyman, K., Falkowski, P. G., and Morel, F. M. M.: The elemental composition of some marine phytoplankton, *J. Phycol.*, 39, 1145–1159, 2003.
- Hollibaugh, J. T. and Azam, F.: Microbial degradation of dissolved proteins in seawater, *Limnol. Oceanogr.*, 28, 1104–1116, 1983.
- Hoiczyk, E. and Hansel, A.: Cyanobacterial cell walls: News from an unusual prokaryotic envelope, *J. Bacteriol.*, 182, 1191–1199, 2000.
- Ietswaart, T., Schneider, P. J., and Prins, R. A.: Utilization of organic nitrogen sources by two phytoplankton species and a bacterial isolate in pure and mixed cultures, *Appl. Environ. Microbiol.*, 60, 1554–1560, 1994.
- Jacobsen, A., Egge, J. K., and Heimdal, B. R.: Effects of increased concentration of nitrate and phosphate during a springbloom experiment in mesocosm, *J. Exp. Mar. Biol. Ecol.*, 187, 239–251, 1995.
- Joint, I., Henriksen, P., Fonnes, G. A., Bourne, D., Thingstad, T. F., and Riemann, B.: Competition for inorganic nutrients between phytoplankton and bacterioplankton in nutrient manipulated mesocosms, *Aquat. Microb. Ecol.*, 29, 145–159, 2002.
- Junk, G. and Svec, H. J.: The absolute abundance of the nitrogen isotopes in the atmosphere and compressed gas from various sources, *Geochim. Cosmochim. Acta*, 14, 234–243, 1958.
- Jørgensen, N. O. G. and Jacobsen, C. S.: Bacterial uptake and utilization of dissolved DNA, *Aquat. Microb. Ecol.*, 11, 263–270, 1996.
- Jørgensen, N. O. G., Kroer, N., Coffin, R. B., Yang, X.-H., and Lee, C.: Dissolved free amino acids, combined amino acids, and DNA as sources of carbon and nitrogen to marine bacteria, *Mar. Ecol. Prog. Ser.*, 98, 135–148, 1993.
- Karner, M. and Herndl, G. F.: Extracellular enzymatic activity and secondary production in free-living and marine-snow-associated bacteria, *Mar. Biol.*, 113, 341–347, 1992.
- Koroleff, F.: Direct determination of ammonia in natural waters as indophenol blue, Council Meeting Document – International Council for the Exploration of the Sea, 19–22, 1969.
- Koroleff, F.: Determination of phosphorus, in: *Methods in seawater analysis*, edited by: Grasshoff, K., Erhardt, M., and Kremling, K., Verlag Chemie, 125–131, 1983.

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Kristiansen, S., Thingstad, T. F., van der Wal, P., Farbroth, T., and Skjoldal, E. F.: An *Emiliania huxleyi* dominated subsurface bloom in Samnangerfjorden, western Norway. Importance of hydrography and nutrients, *Sarsia*, 79, 357–368, 1994.

Kuenzler, E. J. and Perras, J. P.: Phosphatases of marine algae, *Biol. Bull.*, 128, 271–284, 1965.

Laws, E. A.: Photosynthetic quotients, new production and net community production in the open ocean, *Deep-Sea Res.*, 38, 143–167, 1991.

Lessard, E. J., Merico, A., and Tyrrell, T.: Nitrate: phosphate ratios and *Emiliania huxleyi* blooms, *Limnol. Oceanogr.*, 50, 1020–1024, 2005.

Loferer-Krößbacher, M., Klima, J., and Psenner, R.: Determination of bacterial cell dry mass by transmission electron microscopy and densitometric image analysis, *Appl. Environ. Microbiol.*, 64, 688–694, 1998.

Løvdaal, T., Skjoldal, E. F., Heldal, M., Norland, S., and Thingstad, T. F.: Changes in morphology and elemental composition of *Vibrio splendidus* along a gradient from carbon-limited to phosphate-limited growth, *Microb. Ecol.*, published online first, doi:10.1007/s00248-007-9262-x, 2007a.

Løvdaal, T., Tanaka, T., and Thingstad, T. F.: Algal-bacterial competition for phosphorus from dissolved DNA, ATP, and orthophosphate in a mesocosm experiment, *Limnol. Oceanogr.*, 52, 1407–1419, 2007b.

Lukacs, G. L., Haggie, P., Seksek, O., Lechardeur, D., Freedman, N., and Verkman, A. S.: Size-dependent DNA mobility in cytoplasm and nucleus, *J. Biol. Chem.*, 275, 1625–1629, 2000.

Martinez, J. and Azam, F.: Aminopeptidase activity in marine chroococcoid cyanobacteria, *Appl. Environ. Microbiol.*, 59, 3701–3707, 1993.

Middelboe, M., Søndergaard, M., Letarte, Y., and Borch, N. H.: Attached and free-living bacteria: Production and polymer hydrolysis during a diatom bloom, *Microb. Ecol.*, 29, 231–248, 1995.

Montoya, J. P., Voss, M., Kähler, P., and Capone, D. G.: A simple, high-precision, high-sensitivity tracer assay for N_2 fixation, *Appl. Environ. Microbiol.*, 62, 986–993, 1996.

Nimer, E., Schneiderman, R., and Maroudas, A.: Diffusion and partition of solutes in cartilage under static load, *Biophys. Chem.*, 106, 125–146, 2003.

Palenik, B. and Henson, S. E.: The use of amides and other organic nitrogen sources by the phytoplankton *Emiliania huxleyi*, *Limnol. Oceanogr.*, 42, 1544–1551, 1997.

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdaal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

- Pantoja, S. and Lee, C.: Cell-surface oxidation of amino acids in seawater, *Limnol. Oceanogr.*, 39, 1718–1726, 1994.
- Paul, J. H.: Uptake of organic nitrogen, in: *Nitrogen in the marine environment*, edited by: Carpenter, E. J. and Capone, D. G., Academic Press, Inc., New York, 275–308, 1983.
- 5 Pengerud, B., Skjoldal, E. F., and Thingstad, T. F.: The reciprocal interactions between degradation of glucose and ecosystem structure. *Studies in mixed chemostat cultures of marine bacteria, algae, and bacterivorous nanoflagellates*, *Mar. Ecol. Prog. Ser.*, 35, 111–117, 1987.
- Perry, M. J.: Alkaline phosphatase activity in subtropical Central North Pacific waters using a sensitive fluorometric method, *Mar. Biol.*, 15, 113–119, 1972.
- 10 Paasche, E.: A review of the coccolithophorid *Emiliania huxleyi* (Prymnesiophyceae), with particular reference to growth, coccolith formation, and calcification-photosynthesis interactions, *Phycologia*, 40, 503–529, 2002.
- Rees, A. P., Woodward, E. M. S., Robinson, C., Cummings, D. G., Tarran, G. A., and Joint, I.: Size-fractionated nitrogen uptake and carbon fixation during a developing coccolithophore bloom in the North Sea during June 1999, *Deep-Sea Res. II*, 49, 2905–2927, 2002.
- 15 Riegman, R., Noordeloos, A. A. M., and Cadée, G. C.: *Phaeocystis* blooms and eutrophication of the continental coastal zones of the North Sea, *Mar. Biol.*, 112, 479–484, 1992.
- Riegman, R., Stolte, W., Noordeloos, A. A. M., and Slezak, D.: Nutrient uptake and alkaline phosphatase (EC 3:1:3:1) activity of *Emiliania huxleyi* (Prymnesiophyceae) during growth under N and P limitation in continuous cultures, *J. Phycol.*, 36, 87–96, 2000.
- 20 Sambrook, J. and Russell, D. W.: *Molecular cloning: A laboratory journal*, Cold Spring Harbour Laboratory Press, 2001.
- Samuelsson, K., Berglund, J., Haecky, P., and Andersson, A.: Structural changes in an aquatic microbial food web caused by inorganic nutrient addition, *Aquat. Microb. Ecol.*, 29, 29–38, 2002.
- 25 Simon, M., Grossart, H.-P., Schweitzer, B., and Ploug, H.: Microbial ecology of organic aggregates in aquatic ecosystems, *Aquat. Microb. Ecol.*, 28, 175–211, 2002.
- Smith, D. C., Simon, M., Alldredge, A. L., and Azam, F.: Intense hydrolytic enzyme activity on marine aggregates and implications for rapid particle dissolution, *Nature*, 359, 139–142, 1992.
- 30 Sokal, R. R. and Rohlf, F. J.: *Biometry*, 3rd ed., W. H. Freeman, 207–271, 1995.
- Tanaka, T., Henriksen, P., Lignell, R., Olli, K., Seppälä, J., Tamminen, T., and Thingstad, T. F.: Specific affinity for phosphate uptake and specific alkaline phosphatase activity as diagnostic

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

tools for detecting P-limited phytoplankton and bacteria, *Estuaries and Coasts*, 29, 1226–1241, 2006.

Thingstad, T. F.: Control of bacterial growth in idealized food webs, in: *Microbial ecology of the oceans*, edited by: Kirchman, D. L., Wiley-Liss, 229–260, 2000.

- 5 Thingstad, T. F. and Rassoulzadegan, F.: Conceptual models for the biogeochemical role of the photic zone microbial food web, with particular reference to the Mediterranean Sea, *Prog. Oceanogr.*, 44, 271–286, 1999.

Thingstad, T. F., Skjoldal, E. F., and Bohne, R. A.: Phosphorus cycling and algal-bacterial competition in Sandsfjord, western Norway, *Mar. Ecol. Prog. Ser.*, 99, 239–259, 1993.

- 10 Zubkov, M. V., Fuchs, B. M., Tarran, G. A., Burkill, P. H., and Amann, R.: High rate of uptake of organic nitrogen compounds by *Prochlorococcus* cyanobacteria as a key to their dominance in oligotrophic waters, *Appl. Environ. Microbiol.*, 69, 1299–1304, 2003.

Zubkov, M. V. and Tarran, G. A.: Amino acid uptake of *Prochlorococcus* spp. in surface waters across the South Atlantic subtropical front, *Aquat. Microb. Ecol.*, 40, 241–249, 2005.

BGD

4, 3343–3375, 2007

Algal-bacterial competition for N and P

T. Løvdal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

**Algal-bacterial
competition for N and
P**T. Løvdal et al.

Table 1. Uptake of labelled substrate in the $>0.8\ \mu\text{m}$ size fraction as percent of total uptake. Pooled data for inorganic and organic N and P substrates, grouped in samples with T_{NH_4} longer than and shorter than 10 h. Means with standard deviation.

Substrate	$T_{\text{NH}_4}>10\ \text{h}$	$T_{\text{NH}_4}<10\ \text{h}$
DIN	83 ± 7	96 ± 1
DON	35 ± 28	74 ± 6
DIP	72 ± 22	80 ± 7
DOP	25 ± 8	77 ± 13

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Algal-bacterial competition for N and P

T. Løvdal et al.

Table 2. Uptake of labelled substrates by particle-associated bacteria and phytoplankton given as percent of total uptake. Estimates from Eq. (2) assuming similar affinity in particle-associated bacteria as in free-living bacteria (see text). Pooled data for inorganic and organic N and P substrates, grouped in samples with T_{NH_4} longer than and shorter than 10 h. Means with standard deviation.

Substrate	$T_{\text{NH}_4} > 10$ h		$T_{\text{NH}_4} < 10$ h	
	particle-associated bacteria	phytoplankton	particle-associated bacteria	phytoplankton
DIN	16±7	67±9	18±7	78±8
DON	28±17	7±13	73±6	1±1
DIP	4±3	68±24	22±6	58±12
DOP	10±2	15±10	23±13	54±25

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




Fig. 1. Ambient DIN:SRP molar ratio through the course of the experiment.

Algal-bacterial competition for N and P

T. Løvdaal et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Algal-bacterial competition for N and P

T. Løvdaal et al.




Fig. 2. APA and affinity for PO_4^{3-} uptake, and affinity for NO_3^- and NH_4^+ uptake, normalized for the summed P-biomass of phytoplankton and bacteria and the summed N-biomass of phytoplankton and bacteria, respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Algal-bacterial
competition for N and
P**

T. Løvdal et al.




Fig. 3. Turnover times (logarithmic scale) of **(A)** N-substrates and **(B)** P-substrates.

Algal-bacterial competition for N and P

T. Løvdal et al.




Fig. 4. Relative proportions of incorporated label in the two size fractions.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Algal-bacterial competition for N and P

T. Løvdaal et al.




Fig. 5. Biomass-specific affinity (logarithmic scale) for the uptake of **(A)** NH_4^+ , **(B)** Leucine-N, **(C)** PO_4^{3-} , **(D)** ATP-P, and **(E)** dDNA-P in the two size fractions (solid lines), and in bacteria and phytoplankton after correcting for particle-associated bacteria (dotted lines) during the N-limited phase. See text for details.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Algal-bacterial competition for N and P

T. Løvdaal et al.




Fig. 6. Comparison of the mean estimated biomass-specific affinity values during the N-limited phase (\pm SD, $n=5$) of phytoplankton and bacteria vs. the theoretical maximum affinity by the diffusion model. The assumption for the diffusion model is that the diffusion constant (D) for small molecules like NH_4^+ and PO_4^{3-} is $10^{-5} \text{ cm}^2 \text{ s}^{-1}$, D for leucine is $3.7 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$ (Nimer et al., 2003), D for ATP is $3.0 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1}$ (Diehl et al., 1991), and D for DNA is $4.9 \times 10^{-6} \text{ cm}^2 \text{ s}^{-1} \times (\text{basepair size})^{-0.72}$ (Lukacs et al., 2000). The solid and dotted lines denote the theoretical maximum for bacteria and phytoplankton, respectively, assuming the cell density is 1.2 g cm^{-3} for bacteria and phytoplankton, dry weight is 50% of wet weight, C weight is 50% of dry weight, and the molar C:N:P ratio is 50:10:1 for bacteria and 106:16:1 for phytoplankton. The dashed line denotes the theoretical maximum for *E. huxleyi*, assuming minimum N and P cell quotas; e.g. 1.1 and $0.05 \text{ fmol } \mu\text{m}^{-3}$, respectively (cf. Riegman et al., 2000).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion