



Branch xylem density variations across Amazonia

S. Patiño, J. Lloyd, R. Paiva, C. A. Quesada, T. R. Baker, A. J. B. Santos, L. M. Mercado, Y. Malhi, O. L. Phillips, A. Aguilar, et al.

► To cite this version:

S. Patiño, J. Lloyd, R. Paiva, C. A. Quesada, T. R. Baker, et al.. Branch xylem density variations across Amazonia. Biogeosciences Discussions, 2008, 5 (3), pp.2003-2047. hal-00298004

HAL Id: hal-00298004

<https://hal.science/hal-00298004>

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

Branch xylem density variations across Amazonia

S. Patiño^{1,2,3}, J. Lloyd², R. Paiva⁴, C. A. Quesada^{2,5}, T. R. Baker², A. J. B. Santos^{5,6,†}, L. M. Mercado⁷, Y. Malhi⁸, O. L. Phillips², A. Aguilar⁹, E. Alvarez¹¹, L. Arroyo¹², D. Bonal³, A. C. L. Costa¹³, C. I. Czimczik¹⁴, J. Gallo¹⁵, R. Herrera¹⁶, N. Higuchi⁵, V. Horna¹⁷, E. J. Hoyos¹⁸, E. M. Jimenez⁹, T. Killeen¹⁹, E. Leal¹⁰, F. Luizão⁵, P. Meir²⁰, A. Monteagudo^{21,22}, D. Neill²³, P. Núñez-Vargas²¹, W. Palomino²¹, J. Peacock², A. Peña-Cruz²², M. C. Peñuela⁹, N. Pitman²⁴, N. Priante Filho²⁵, A. Prieto¹, S. N. Panfil²⁶, A. Rudas²⁷, R. Salomão¹⁰, N. Silva^{28,29}, M. Silveira³⁰, S. Soares de Almeida¹⁰, A. Torres-Lezama³¹, J. D. Turriago¹⁵, R. Vásquez-Martínez²¹, M. Schwarz¹⁶, A. Sota¹⁶, J. Schmerler¹⁶, I. Vieira¹⁰, B. Villanueva¹⁵, and P. Vitzthum⁹

¹Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. Diagonal 27 No. 15-09, Bogotá D.C, Colombia

²Earth and Biosphere Institute, School of Geography, Univ. of Leeds, LS2 9JT, UK

³UMR-ECOFOG, INRA, 97310 Kourou, French Guiana

⁴Secretaria Municipal de Desenvolvimento e Meio Ambiente na Prefeitura Municipal de Maués, Maués, Brazil

⁵Instituto National de Pesquisas Amazônicas, Manaus, Brazil

⁶Departamento de Ecología, Universidad de Brasília, Brazil

⁷Centre for Ecology and Hydrology, Wallingford, UK

⁸Oxford University, Centre for the Environment, Oxford, UK

⁹Universidad Nacional de Colombia Sede Amazonía, Leticia, Colombia

¹⁰Museu Paraense Emílio Goeldi, Belém, Brazil

¹¹Equipo de Gestión Ambiental, Interconexión Eléctrica S.A. ISA., Medellín, Colombia

¹²Museo Noel Kempff Mercado, Santa Cruz, Bolivia

¹³Universidade Federal do Pará, Belém, Brazil

¹⁴Department of Earth System Science, University of California, Irvine, CA, USA

¹⁵Departamento de Biología, Universidad Distrital, Bogotá, Colombia

¹⁶Max-Planck-Institut für Biogeochemie, Jena, Germany

¹⁷Abteilung Ökologie und Ökosystemforschung, Albrecht-von-Haller-Institut für Pflanzenwissenschaften, Universität Göttingen, Göttingen, Germany

¹⁸Departamento de Ciencias Forestales, Univ. Nacional de Colombia, Medellín, Colombia

¹⁹Center for Applied Biodiversity Science, Conservation Int., Washington D.C., WA, USA

²⁰School of Geography, University of Edinburgh, Edinburgh, Scotland, UK

²¹Herbario Vargas, Universidad Nacional San Antonio Abad del Cusco, Cusco, Perú

²²Proyecto Flora del Perú, Jardín Botánico de Missouri, Oxampapa, Perú

²³Herbario Nacional del Ecuador, Quito, Ecuador

²⁴Center for Tropical Conservation, Duke University, Durham, NC, USA

²⁵Universidade Federal do Mato Grosso, Cuiabá, Brazil

²⁶Department of Botany, University of Georgia, Athens, GA, USA

²⁷Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá, Colombia

²⁸CIFOR, Tapajos, Brazil

²⁹EMBRAPA Amazonia Oriental, Belém, Brazil

³⁰Departamento de Ciências da Natureza, Universidade Federal do Acre, Rio Branco, Brazil

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



³¹Facultad de Ciencias Forestales y Ambiental, Universidad de Los Andes, Mérida, Venezuela
†deceased

Received: 27 February 2008 – Accepted: 10 March 2008 – Published: 9 May 2008

Correspondence to: S. Patiño (sanpatiga@gmail.com)

BGD

5, 2003–2047, 2008

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

BGD

5, 2003–2047, 2008

Measurements of branch xylem density, D_x , were made for 1466 trees representing 503 species, sampled from 80 sites across the Amazon basin. Measured values ranged from 240 kg m^{-3} for a *Brosimum parinariooides* from Tapajos in West Pará, Brazil to 1130 kg m^{-3} for an *Aiouea* sp. from Caxiúana, Central Pará, Brazil. Analysis of variance showed significant differences in average D_x across the sample plots as well as significant differences between families, genera and species. A partitioning of the total variance in the dataset showed that geographic location and plot accounted for 33% of the variation with species identity accounting for an additional 27%; the remaining “residual” 40% of the variance accounted for by tree to tree (within species) variation. Variations in plot means, were, however, hardly accountable at all by differences in species composition. Rather, it would seem that variations of xylem density at plot level must be explained by the effects of soils and/or climate. This conclusion is supported by the observation that the xylem density of the more widely distributed species varied systematically from plot to plot. Thus, as well as having a genetic component branch xylem density is a plastic trait that, for any given species, varies according to where the tree is growing and in a predictable manner. Exceptions to this general rule may be some pioneers belonging to *Pourouma* and *Miconia* and some species within the genera *Brosimum*, *Rinorea* and *Trichilia* which seem to be more constrained in terms of this plasticity than most species sampled as part of this study.

1 Introduction

The Amazon Basin remains the home to the most diverse and largest contiguous tropical forest on the planet (Malhi and Grace, 2000; Laurance et al., 2004). Different ecological systems and vegetation formations exist within its boundary creating a mosaic of forests and vegetation types with a floristic complexity the basis of which is still not well understood (Phillips et al., 2003).

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



How and why species are distributed and where trees can grow in this mosaic of tropical forest types is a central question for tropical ecology (Pitman et al., 2001; Phillips et al., 2003). On the one hand water availability has been considered as an important factor determining tree species distributions at global (Woodward, 1987) and regional scales (Borchert, 1998; Veenendaal and Swaine, 1998; Bongers et al., 1999). Tree physiology data also shows that it may be one key factor determining species distributions in tropical rain forest despite the substantial rainfall they receive (Meinzer et al., 1999; Engelbrecht et al., 2002, 2005, 2006, 2007; Tyree et al., 2003; Baltzer et al., 2008). On the other hand, although water shortage (seasonality and rainfall) is not considered by some as an important selective pressure determining the functional composition of moist and wet tropical forest (Baker et al., 2004b; ter Steege and Hammond, 2004), it clearly does help to determine overall diversity and distribution patterns of canopy tree genera across the Amazon basin (ter Steege et al., 2006). Light as a factor explaining coexistence of trees has also been considered (Poorter and Arets, 2003) and climate more than geographic distance in lowland forest seems also to affect species composition (Pitman et al., 2001) and distributions (Bongers et al., 1999; Baltzer et al., 2008). Additionally, even if there clearly is some “distance-decay” i.e. the similarity between two sites declines as the distance between them increases, see (Soininen et al., 2007) in Amazonian forest communities (Condit et al., 2002; Tuomisto et al., 2003), consistent with the view of plant species distribution as a partly random controlled process (Hubbell, 2001), it is also clear that substrate-mediated processes play a significant deterministic role in controlling species distributions across Amazonian landscapes (Gentry, 1988; Tuomisto et al., 1995, 2003; Phillips et al., 2003), this also being the case for other tropical forests (Harms et al., 2001; Robert et al., 2007). These theories are by no means exclusive, and one of the main purposes of the work described in this special issue (Lloyd et al., 2008a¹) was to interface dedicated plant physiological trait data with species abundance, climate and soil type information to

¹Lloyd, J., Grace, J., and Meir, P.: Introducing the “Biogeochemistry and Function of Amazon Forest” project, Biogeosciences Discuss., in preparation, 2008a.

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



help understand the underlying factors controlling species distribution, composition and stand dynamics at the Basin-wide scale. In what is the first of a series of papers, we here analyse in detail geographic and taxonomic patterns of tree branch wood density across Amazonia. Although wood density (D_w) is most often measured on tree boles sampled after destructive harvesting, often in association with commercial operations (Baker et al., 2004b; Chave et al., 2006), here we have used an alternative approach – the sampling of the density of the functional xylem or sapwood of small (ca. 1.5 cm diameter) branches. This measurement, referred here as “xylem density”, D_x , is a plant trait usually considered in physiological processes studies; for example when looking at wood properties in relation to transpiration, stem water storage capacity, cavitation resistance, mechanical support, photosynthesis and/or growth (Niklas, 1997; Enquist et al., 1999; Stratton et al., 2000; Hacke et al., 2001; Meinzer, 2003; Bucci et al., 2004; Santiago et al., 2004; Jacobsen et al., 2007a, 2007b; Pratt et al., 2007; Scholz et al., 2007). We rationalised that these are all processes that should be expected to influence where a given tree species can survive and thus should ultimately contribute to the shaping of species distributions across landscapes.

For any given species D_x and D_w should be related as both reflect an individual species' water transport strategy and the invariable trade-off against mechanical strength (Taneda and Tateno, 2004; Wagner et al., 1998; Asner and Goldstein, 1997). But D_w may well be affected by factors in addition to those modulating D_x . For example it may also reflect differences in the storage of resins and/or increased of secondary compounds of bole heartwood with time, different biomechanical requirements for maintaining vertical position and support of the whole tree and individual branches, also generally reflecting the longer term growth history of the tree. It means that, D_w may be considered to effectively integrate plant ontogeny, edaphic and climatic effects over a plant's lifetime.

It has long known that D_w is a genetically conserved trait, and this characteristic has been used extensively in tree breeding (Yang et al., 2001) also leading to studies of its phylogenetic inheritance (Enquist et al., 1999; Chave et al., 2006). Nevertheless, it has

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



also long been known that for a given tree species, marked variations may be observed attributable to differences in site quality and/or forest type. There have already been several reports of variations with site/growth conditions of order 10% for plantation tropical tree species (Roque, 2004; Erskine et al., 2005). Similarly, site specific differences 5 have been noted when comparing the same species growing in different forests or site conditions (Gonzalez and Fisher, 1998; Woodcock et al., 2000; Muller-Landau, 2004; Parolin and Ferreira, 2004; Roque, 2004; Nogueira et al., 2005; Nogueira et al., 2007). Lower D_w is often associated with early and secondary succession vegetation (Saldarriaga 1989; Woodcock et al., 2000) and higher growth rates (Wiemann and Williamson, 10 Erskine et al., 2005; Nogueira et al., 2005; King et al., 2005, 2006).

We thus hypothesised that D_x is a “plastic” trait that reflects both phylogenetic heritage of trees and overall site conditions. Therefore variations in D_x at the stand level across Amazonia should not only reflect differences in species composition, but differences in soil and climate conditions. If a given species can grow in different sites, D_x of 15 that species may converge to an “average” value of that of co-occurring species within each site. If this is true, species with enough phenotypic plasticity can occupy more and larger areas and become more common than species with small plasticity which will occupy restricted areas and become “rare” species at a large-scale but common in just one site.

20 2 Methods

2.1 Study sites

Eighty forest plots from across the Amazon basin were sampled normally, at the end 25 of the rainy season of the region between January 2001 and December 2005. Additionally, two plots were sampled in Paracou, French Guiana in September 2007 (see details below). These plots form part of the RAINFOR project (www.rainfor.org; Malhi et al., 2002) and span local, regional and Basin-wide environmental gradients. Many

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of the plots have been described in detail elsewhere (Baker et al., 2004a, b; Malhi et al., 2002; Phillips et al., 2004; Vinceti, 2003) and Appendix A lists all the plots visited, including those not previously described and in some cases with updated information.

2.2 Sampling of plant material

5 Normally, around 20 trees greater than 10 cm dbh (diameter at breast height i.e. at 1.3 m from the base of the tree) were chosen in each plot. On some occasions, for instance when plots were unusually heterogeneous, as a consequence of topographic variations and/or shape (i.e. 1000×10 m) more trees were sampled (for example Bogi plots in Ecuador). For two of the Caxiúana plots (Central Pará, Brazil) we had sampled
10 in two consecutive years (2002 and 2003) and since there was no statistical difference in D_x for the two years, we merged the data in just one set. When a plot was clearly composed of different defined landscapes, and each landscape was considered as an individual plot with often less than 20 trees sampled within it (e.g. Jacaranda Plots, Km 34 Manaus, Brazil).

15 Trees were not chosen completely at random but taking into account two factors. First, often there was a selection of three to six contrasting areas (slopes, valleys, gaps, flood, etc) where these were present, a professional tree climber then choose a “climbable tree” within the generally identified areas. Naturally, this “climbable tree” varied from climber to climber according to the technique employed and overall climbing
20 skills. Nevertheless a general rule was that the “climbable tree” was >10 cm dbh with at least some neighbouring trees reachable by a clipper pole. From the climbed tree, the climber sampled one branch from each of the upper, (exposed to the sun), middle (semi light) and lower crown (shade), and one branch from the upper crown of the (usually three to five) surrounding trees.

25 The only difference for this sampling strategy was for two of the Guyaflux plots at Paracou, French Guiana where only lower branches from subcanopy trees were sampled. An analysis of over 200 trees comparing D_x of branches from the upper, middle

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



and lower crown of the trees using the techniques described in Lloyd et al. (2008b)² has, however, shown no statistically significant effect of height on D_x (data not shown). The subcanopy trees sampled at Paracou represented the distribution of the main families present in those plots with D_x showing the same distribution of frequencies found for D_w of 309 trees from Paracou (J. Beauchen, personal communication). In any case, because these measurements were made after the main analysis of this paper we have simply included these values to help illustrate regional patterns (Fig. 1).

2.3 Species identification

Details of the species identification from the permanent plots are described elsewhere (Baker et al., 2004b) and in this work we have used the new classification given by the 10 Angiosperm Phylogeny Group II (APG 2003, <http://www.mobot.org/MOBOT/Research/APweb/>), in which *Bombacaceae*, *Sterculiaceae*, and *Tiliaceae* are all included in the *Malvaceae*; *Papilionaceae*, *Caesalpiniaceae*, and *Mimosaceae* are included in the *Fabaceae*; *Cecropiaceae* in the *Urticaceae*; and *Flacourtiaceae* in the *Salicaceae*.

15 2.4 Xylem density determinations

A pair of stem segments of 0.05 to 0.1 m long and 0.01 to 0.02 m diameter were cut from each branch after harvesting and immediately placed in plastic bags to avoid desiccation and returned to the laboratory or field station. Normally within 12 h of sampling (but sometimes as long as 36 h later) the outer bark and phloem were removed from 20 one of the two sample stems and its fresh volume calculated from its stem length and the average diameter of the two equiaxial diameters at each end. When the pith was wider than 2 mm diameter the stem was cut into a small segment (0.02 to 0.05 m long)

²Lloyd, J., Patiño, S., Paiva, R. Q., Nardoto, G., Quesada, C. A., Santos, A. J. B., Baker, T. R., Brand, W. A., Hilke, I., Gielmann, H., Raessler, M., and Mercado, L. M.: Gradients in leaf physiological properties within Amazon forest canopies, Biogeosciences Discuss., in preparation, 2008b.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and the pith removed with a small screw driver or scalpel. When the pith was thinner than 2 mm it was not removed from the stem (as it was thus assumed to be of negligible mass) but its volume subtracted from the volume of the stem without bark. Pith volume was calculated by measuring the average diameter (two measurements of diameter at each end of the stem) and stem length. All stems were then dried at 70–90°C for three to four days (to constant mass) and weighed. Xylem density, D_x , was then determined as the dry mass divided by the green volume of the sample.

2.5 Statistical analysis

Basic statistics shown in Figs. (1), (2), and (3) were performed with Minitab 14 (Minitab Inc.). All Standard Major Axis Regression analysis (Fig. 5) were done using SMATR (Warton et al., 2006). Variance partitioning for Fig. 4 was accomplished by applying Residual Error Maximum Likelihood (REML) analysis (Gilmour et al., 1995) employing GENSTAT Discovery Edition. Mixed-effect modelling (Fig. 6) was carried out with the “Imne” package (Bates et al., 2007) and rank-based linear regression (Fig. 7) accomplished as in Terpstra and McKean (2005), both using the “R” statistical computing package (R Development Core Team, 2007). For the latter analysis, we applied the “high-breakpoint” option (Chang et al., 1999) to account for the possibility of “contaminated” data having been included in any of the D_w values assimilated from a wide range of sources into the RAINFOR “wood density” database.

In order to determine the extent to which D_x changes in a given species within the same plot and between plots, IPP, index of phenotypic plasticity (Valladares et al., 2000), was calculated as the absolute difference between the maximum value and the minimum value divided by the maximum value.

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



3 Results

We measured D_x of 1466 trees from 80 plots across the Amazon basin (Appendix A) <http://www.biogeosciences-discuss.net/5/2003/2008/bgd-5-2003-2008-supplement.pdf>.

Data for D_x followed a normal distribution with a mean value of 620 kg m^{-3} ; 5 normality test (StDev=0.13, $N=1466$, AD=1.82 $P<0.001$).

Of all the trees sampled, 92% (1352) had been identified to the family level, 86% (1258) to the genus level, and 69% (1006) to the species level. The trees sampled accounted for 58 families, representing 40% of the total number of families present in the neotropics (Mass and Westra, 1993) with 249 genera, and 503 species being sampled. 10 The most common families sampled were *Fabaceae*, *Sapotaceae*, *Moraceae*, *Lecythidaceae*, *Burseraceae*, *Myristicaceae*, *Lauraceae*, *Euphorbiaceae*, *Chrysobalanaceae*, *Annonaceae*; with the most common genera being *Eschweilera*, *Pouteria*, *Protium*, *Inga*, *Licania*, *Pseudolmedia*, *Virola*, *Pourouma*, *Miconia*. The most common species were *Eschweilera coriacea*, *Pseudolmedia laevis*, *Rinorea guianensis*, 15 *Tetragastris altissima*, *Minquartia guianensis*, *Pourouma guianensis*, and *Pseudolmedia macrophylla*. We had 20 undetermined *Protium* sp., 17 *Pouteria* sp., 14 *Inga* sp., 11 *Ocotea* sp., 10 *Eschweilera* sp.

3.1 Geographic variation

Arithmetic mean D_x for the 80 plots are shown in Fig. 1, which also shows our separation into 13 specified geographical regions and used for subsequent analysis. 20

From Fig. 1 a gradient of increasing D_x from south and north towards the Amazon River is apparent with high D_x being concentrated along the river itself. Plots located close to the Andes tend to have the lowest D_x . For example, all plots in Ecuador, some in North Perú, South Perú and Bolivia had a relatively low D_x compared to plots in 25 Colombia and North Perú which were all at a lower altitude and closer to the Amazon River (see Appendix A). Similar patterns of low D_w in forests close to the Andes have previously been reported (Chave et al., 2006; Baker et al., 2004b; ter Steege et al.,

BGD

5, 2003–2047, 2008

Amazonian xylem density variation

S. Patiño et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2006). Taking the basin as a whole (no division into regions), statistically significant differences existed between plot means ($P<0.001$) ranging from $800\pm50 \text{ kg m}^{-3}$ (\pm standard deviation) at the dry experiment plot at Caixuana (Projecto Secaflor), CAX-04, with the nearby control plot CAX-03 being the second highest at $780\pm120 \text{ kg m}^{-3}$.

5 These are both terra firme forests located in Central Pará, Brazil (CP-Brazil region). The lowest plot means were for TAM-03 a swamp forests in Tambopata, South Perú (S-Perú & AC-Brazil region) and JAS-05 a forest growing on recently deposited river sediments (fluvisol) in Jatun Sacha in the Ecuador region. Both these plots had a mean D_x of 470 kg m^{-3} . Data for all 80 plots are summarised in Appendix B.

10 Figure 2 gives means (\pm standard deviations) for all plots, grouped according to region, with regions being presented sequentially from top to bottom according to the overall mean D_x for the trees sampled within them. This shows that, although considerable plot-to-plot variation existed within regions (e.g. N. Perú and Colombia) large statistical differences between regions also existed ($P<0.001$). Of these, the highest

15 overall value was for Central Pará in Brazil ($760\pm130 \text{ kg m}^{-3}$, $N=144$) which had significantly higher D_x (Tukey Test) than the rest of the regions while Ecuador had the lowest overall values ($540\pm92 \text{ kg m}^{-3}$). Nevertheless, this region did not differ significantly from Acre, Mato Grosso (Brazil), Colombia and Bolivia. Within some regions:

20 PC-Brazil, PE-Brazil, N-Perú, PW-Brazil, Colombia, S-Perú, MT-Brazil, and Ecuador mean D_x of plots varied considerably (Appendix B), while in some regions Bolivia, AC-Brazil, NE-Venezuela, SW-Venezuela, plots were not significantly different from each other.

3.2 Taxonomic variation

In a similar manner to the Region/Plot analysis above, variation in D_x at the family and genera level is summarised in Fig. 3. Overall there were significant differences between the families sampled ($F=8.08$ $DF=57$ $P<0.001$). Families with D_x higher than the basin mean were *Olaceaceae*, *Celastraceae*, *Chrysobalanaceae*, *Humiriaceae*, *Ochnaceae*, *Linaceae*, *Scrophulariaceae*, *Myrtaceae*, and *Lecythidaceae* (Fig. 3). Families with

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

lower D_x were *Boraginaceae*, *Bixaceae*, *Sabiaceae*, *Lepidobotryaceae*, *Lacistemataceae*, *Rhamnaceae*, *Malvaceae*, *Annonaceae*, *Myristicaceae*, *Urticaceae*, *Vochysiaceae*, *Araliaceae*, *Dichapetalaceae*, *Bignoniaceae*, and *Euphorbiaceae*. The remaining families all contained genera characterised by both high and low D_x and include some of the most abundant families across the basin: *Fabaceae*, *Rubiaceae*, *Lauraceae*, *Sapotaceae*, *Apocynaceae* (Fig. 3). There were also significant differences between genera ($F=3.78$ $DF=249$ $P<0.0001$) with the highest density genera being *Ajouea*, *Callichlamys*, *Pithecellobium*, *Vatairea*, *Stachyarrhena*, *Dipteryx*, *Machaerium*. The genera with lower density were *Annona*, *Matisia*, *Tetrorchidium*, *Collophora*, *Onychopetalum*, *Hieronima*, *Luehea*.

3.3 Partialling out geographical and taxonomic differences

In order to apportion the total variance in the dataset observed (Searle et al., 2006) into geographical and taxonomic components, we used Residual Maximum Likelihood (REML) Analysis to fit a model according to

$$15 \quad D = \mu + r/p + f/g/s + \varepsilon \quad (1)$$

where μ represents the overall mean of the dataset (620 kg m^{-3}); effects of location are incorporated in the term r/p , which denotes that within each region (r) are nested more than one plot (p); genetic effects are represented by the term $f/g/s$, which denotes that within each family (f) are nested various genera (g), within which are nested several species (s); and (ε) represents the residual variance. All effects were taken as random variables, as we had sampled only a limited subset of plots within distinct but not comprehensive regions; also sampling a more or less random (and incomplete) selection of Amazon families, genera and species.

20 Results of this analysis are shown in Fig. 4. Taken together, the geographical parameters (region and plot) accounted for 33% of the total variation with 26% of this being attributable to between region variations (this effectively representing the average variation between plots in any one region). Taxonomic variability in D_x , was inferred as

Title Page	Abstract	Introduction
Conclusions	References	
Tables	Figures	
◀	▶	
◀	▶	
Back	Close	
Full Screen / Esc		
Printer-friendly Version		
Interactive Discussion		



27% of the total variability in the dataset, with species and family accounting for about 10% and genus per se accounting only for 6%. Overall, the proportion of the variance in the dataset that remained unexplained was 40%, this also being the “residual variance” reflecting within tree variation (only one branch per tree was sampled), tree-to-tree variation and also may reflect measurement errors. Note that in this analysis we only included species that were observed to occur in more than one site and included families and genera for which more than one species had been sampled. The analysis presented here for D_x differs from others (Chave et al., 2006; Baker et al., 2004b) in that we have not taken overall means for each species; but rather included intra-specific variation and the possibility of systematic plot-to-plot variations in our interpretation.

Figure 4 suggests that geographic location is as important, if not more important than taxonomic identity in determining the value of D_x observed for any given tree but with considerable variation accountable for by neither. This point is demonstrated further in Fig. 5, where we have taken the more widely abundant families, genera and species in our data set and plotted the values observed in all plots where they were sampled as a function of the average tree density of all other trees sampled in the same plot. Detailed results from this analysis (which also included a consideration of likely errors in both mean plot and mean family/genus/species values for that plot) are also given in Appendix C.

Figure 5A shows that nearly all families examined converged to a statistically significant trend for an increase in the average D_x observed with a slope close to 1:1 as the values observed by other families at that site also increased. Likewise the same trend was observed when different genera were examined (Fig. 5B) and, somewhat surprisingly, also for species (Fig. 5C). Notable exceptions did however exist for the families *Urticaceae* (Fig. 4A, panel M) the family containing the genus *Pourouma* which almost entirely consist of pioneer species (Withmore 1989) and *Myrtaceae* (Fig. 4A, panel U) which in our data set includes the genera *Eugenia* and *Myrcia*. Likewise at the genus level *Pourouma* and *Brosimum* also seemed to vary less than all the genera examined. The relationships for *Miconia* and *Trichilia* were also not highly significant (Appendix

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



C). Figure 5C shows the same trend for the species examined. Nearly half of the species examined showed significant relationships. Non significant relationships might be related to the “pioneer” character of the examined species i.e. *Pououma minor*, *Pououma guianensis* and perhaps to the reduced number of plots where they were collected and use for this analysis (Appendix C).

3.4 Geographical and taxonomic contributions to stand level differences

In order to evaluate the extent to which overall plot-to-plot variations might be accountable in terms of geographical versus taxonomic effects, we utilised estimates of the individual plot and species effects from Eq. (1) and compared them to direct stand level calculations. This was achieved by first estimating the average value for each species within each plot and then obtaining a weighted average value for D_x for that plot according to the observed abundance of each species within it, denoted here as $\langle D_x \rangle$. A similar calculation was done for the REML “species effects” which are plotted along with REML fixed plot effects (the r/p term from Eq. (1) as a function of $\langle D_x \rangle$ in Fig. 6. This analysis shows that by far the majority of the variation in $\langle D_x \rangle$ was accountable in terms of plot-to-plot differences, with the plot effects increasing almost linearly with $\langle D_x \rangle$ with a slope close to 1.0. By contrast the species (i.e. $f/g/s$) effects were more or less constant (and close to zero) for $\langle D_x \rangle > \text{ca. } 550 \text{ kg m}^{-3}$, although declining slightly thereafter. We treated our plot term as a fixed effect for the analysis in Fig. 6 (as opposed to a random effect in Fig. 4), as this permitted us to allow for different plots to have different intrinsic variances which were broadly consistent with difference of plot topographic and soil heterogeneity, also removing a slight bias in residuals which was present when treating the r/p term as random.

3.5 Phenotypic plasticity

In order to determine inter and intra-specific variation we compared the IPP of the same species collected several times within one plot and in many plots. The IPP of one

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



species collected in more than two plots (mean= 0.29 ± 0.12 , $N=86$) was significantly higher (mean= 0.14 ± 0.10 , $N=86$) than the variation of the same species collected more than twice within one plot ($DF=1$, $F=77.94$, $P<0.0001$).

BGD

5, 2003–2047, 2008

4 Discussion

- 5 Our results show that there is a large variation of branch xylem density across Amazonia and that considerable plasticity occurs for many species growing in different forests. This suggests that branch xylem density may not be a simple genetically inherited trait that is predictable on the basis of the knowledge of plant taxonomy alone, and that across basin patterns may not be only explained by patterns of species composition
10 and abundance as has been previously suggested for bole wood density (Enquist et al., 1999; Baker et al., 2004b; Chave et al., 2006). But, knowledge of the site conditions is also necessary. For example, irrespective of the genotypic level examined (Fig. 5C), D_x observed varied by as much as 400 kg m^{-3} across sites. Moreover, this variation was systematic with different trees sampled within a given family/genus/species tending to
15 have higher values of D_x along with other trees in the same plot (and vice versa).

Additional evidence for widespread plasticity comes from the REML variance partitioning of Fig. 4 in which the combined effects of Region/Plot are shown to have contributed to about the same proportion of the overall variation in the data set as did Family/Genus/Species. This result differs significantly from those for D_w (Baker et al.,
20 2004b).

Nevertheless, when our species level means for D_x are examined as a function of species mean D_w using an expanded database from that presented in (Baker et al., 2004b) then there is reasonably good relationship (Fig. 7a). And the average D_x for this study, 620 kg m^{-3} is also very similar to previous values reported for D_w for Amazonia. For example (Brown and Lugo, 1984) estimated 620 kg m^{-3} as the average
25 wood density of tropical America, (Chave et al., 2006) reported 650 kg m^{-3} for Central and South America together and (Baker et al., 2004b) estimated 620 kg m^{-3} as the

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



overall species-level mean for Amazonia.

As reflected in Fig. 7b, our results do, however, differ from the above authors in that much of the variation in D_w is attributed to genotype while in our case, variations in D_x does include site and genetic variations. This difference is understandable, as there
5 is a strong tendency of many species, genera and even families to be geographically confined to certain areas of the Basin (ter Steege et al., 2006). Thus, if there is equivalence between D_x and D_w what appears to be a genetic effect for the latter, may in fact be mostly a geographic (site and regional) effect attribute to variations in soils, climate and/or geological origin. In that respect it is only by studying replicated species growing
10 across a wide range of environments that we have been able to show the strong environmental influence on D_x (and by implication D_w). In essence the REML species effect in Fig. 7b represents the inferred value that each species would have were it to be found on some sort of “overall average site”.

Also worth noting is that, in contrast to the general trend, pioneer species (Whitmore
15 1989) -either short and long-lived such as within the genera *Pououma*, (*Urticaceae*) and *Miconia* (*Melastomataceae*) and some species belonging *Brossimum* (*Moraceae*), *Trichillia* (*Meliaceae*) and *Myrcia* (*Myrtaceae*) among others, often associated to secondary vegetation and/or late stages of forest succession (Banana and Tweheyo, 2001; Peña-Claros, 2003; Vieira et al., 2003; Poorter et al., 2006; Viera and Proctor, 2007;
20 Selaya and Anten, 2008)., showed little tendency to exhibit variation in D_x across the sites where they were found. This brings the question whether species showing little phenotypic plasticity and intermediate D_x values even when present in sites with relative low or high xylem density are more restricted to specific edaphic and microclimatic conditions that sustain colonisation and fast growth i.e. gaps with enough water supply
25 from the soil, light, and nutrients. Also, it is worth noticing that species such as *Pououma minor* and *P. guianensis*, which are generally considered low density-fast-growing species; were by far the species with the lowest densities in our study; xylem density varied from 410 to 690 kg m⁻³, being comparable to any slow-growth old forest species in our data set.

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



But why should such plasticity in D_x generally occurs? Most likely it is related to variation in resource availability and/or different site dependent soil physical characteristics and hydrological constraints to plant function. There is already abundant evidence of this from across a range of species, environments and controlled experiments (Sabatier et al., 1997; Laurance et al., 1999; Hacke et al., 2000; Vormisto et al., 2000; Pelissier et al., 2002; Baraloto and Goldberg, 2004; Kursar et al., 2005; Baraloto et al., 2007; Madelaine et al., 2007). Further evidence of the influence of site conditions on D_x of trees comes from our own data. For example, in a Mangrove forest in East Pará, Brazil (EP-Brazil, BRA-01, Appendix A). There, only two species were sampled (10 individuals per species) *Avicennia germinans* and *Rhizophora mangle* which mean D_x were $722 \pm 87 \text{ kg m}^{-3}$ and $723 \pm 99 \text{ kg m}^{-3}$ respectively. The two species are not phylogenetically related since they belong to two different families (*Scrophulariaceae* and *Rhizophoraceae*) and two different orders (*Lamiales* and *Malpighiales*). Nevertheless they converge to an almost identical D_x . An analysis of the most variable species in our data set *Protium paniculatum* (IPP=0.66), collected in two different plots in Caxiuana (Pará, Brazil): CAX-02 ($D_x=280 \text{ kg m}^{-3}$) and CAX-04 ($D_x=800 \text{ kg m}^{-3}$), and *Perebea tessmannii* (IPP=0.58) two samples collected in the same plot but 900 m apart in BOG-01 with 320 and 760 kg m^{-3} , in both cases low pH, high proportion of sand and low minerals concentrations were be associated with the trees exhibiting high D_x (Quesada et al., 2008³; Patiño et al., 2008a⁴). Hacke et al. (2000) have shown remarkable

³Quesada, C. A., Lloyd, J., Schwarz, M., Baker, T. R., Patiño, S., Czimczik, C., Schmerler, J., Hodnett, M., Arneth, A., Lloyd, G., Dezzeo, N., Gasior, R., Herrera, R., Hilke, I., Kuhlmann, I., Phillips, O., Raessler, M., Chaves, E., Cruz, O., Filho, J. M., Luizão, F. J., Pimentel, T., Santos, A. J. B., Almeida, S., Alvarez, E., Arroyo, L., Higuchi, N., Jimenez, E. M., Lezama, A. T., Neill, D. A., Paiva, R., Priante Filho, N., Silva, N., Silveira, M., Prieto, A., Rudas, A., and Vieira, I.: Chemical and physical properties of Amazonian forest soils in relation to their genesis, *Biogeosciences Discuss.*, in preparation, 2008.

⁴Patiño, S., Lloyd, J., Quesada, C. A., Paiva, R. Q., et al.: Factors influencing observed variation of branch xylem density across the Amazon basin, *Biogeosciences Discuss.*, in prepara-

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



influence of soil texture and porosity on the hydraulic properties of *Pinus taeda*.

In conjunction with concurrent measurements of associated foliar physiological characteristics undertaken on leaves from the branches with which D_x was determined in this study (Lloyd et al., 2008c⁵) as well as associated soil physical and chemical characteristics (Quesada et al., 2008⁶), further papers in this series consider the underlying factors contributing to the strong site effects in Fig. 6 (Patiño et al., 2008a⁴), along with the extent to which these site dependent variations in xylem density reflect a co-ordinated physiological response of different Amazonian tree species to variations in their growth environment (Patiño and Lloyd, 2008b⁷).

10 Appendix A

More precise coordinates will be available (Anderson and Malhi, 2008⁸). Abbreviations in regions are: AC=Acre, AM=Amazonas, MT=Mato Grosso, CP=Central Pará, EP=East Pará, WP=West Pará, N=North, S=South, NE=North East, SW=South
tion, 2008a.

⁵ Lloyd, J., Patiño, S., Paiva, R. Q., Quesada, C. A., Baker, T. R., and Mercado, L. M.: Basin-wide variations in foliar properties of Amazon forest trees, Biogeosciences Discuss., in preparation, 2008c.

⁶ Quesada, C. A., Lloyd, J., Schwarz, M., Baker, T. R., Patiño, S., Czimczik, C., Schmerler, J., Hodnett, M., Arneth, A., Lloyd, G., Dezzeo, N., Gasior, R., Herrera, R., Hilke, I., Kuhlmann, I., Phillips, O., Raessler, M., Chaves, E., Cruz, O., Filho, J. M., Luizão, F. J., Pimentel, T., Santos, A. J. B., Almeida, S., Alvarez, E., Arroyo, L., Higuchi, N., Jimenez, E. M., Lezama, A. T., Neill, D. A., Paiva, R., Priante Filho, N., Silva, N., Silveira, M., Prieto, A., Rudas, A., and Vieira I.: Chemical and physical properties of Amazonian forest soils in relation to their genesis, Biogeosciences Discuss., in preparation, 2008.

⁷ Patiño, S. and Lloyd, J.: Integration of branch xylem density variations into the tropical tree physiological spectrum, Biogeosciences Discuss., in preparation, 2008b.

⁸ Anderson, L. O. and Malhi, Y.: Landscape patterns of forest biophysical measurements in eastern and western Amazonia, Biogeosciences Discuss., in preparation, 2008.

BGD

5, 2003–2047, 2008

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



West,. ** not a permanent plot, samples were taken from trees around the Eddy covariance tower. Additional information of plots may be found in: (Malhi et al., 2002, 2003; Baker et al., 2004; Vinceti, 2003).

BGD

5, 2003–2047, 2008

Appendix B

- 5 Variation of D_x within regions. In the first column the number below the name of the country is the mean followed by the standard deviation in parenthesis. DF=degrees of freedom; F =statistical values, P =probability, N =number of samples, SE =standard error of mean. * means “significantly different”.

Appendix C

- 10 Regression analysis outputs for family (D1) genera (D2) and species (D3). See Figs. (5A), (5B) and (5C). Notice that for this Appendix units for D_x are given in g cm^{-3} .

Acknowledgements. We wish to thank M. Wagner for his help “cleaning up” our data set and for his useful comments that improved the quality of the present manuscript and N. Fyllas for advice on statistical analysis. We also thanks the students from the FTH module in UMR-

- 15 ECOFOG 2007: E. Alberny, S. Dutrey, J. Fournier, H. Richard for help in sampling the French Guiana plots and the VPN Corporation for making available Genstat Discovery Edition for SP when she was working in Colombia. We thank J. Stropp for help with the elaboration of the map shown in Fig. 1 and N. Rowe for comments that improved the last version of the present manuscript. Finally, we dedicate this work to the memory of our friend and colleague A. Santos.

20 References

- Asner, G. P. and Goldstein, G.: Correlating Stem Biomechanical Properties of Hawaiian Canopy Trees with Hurricane Wind Damagein, *Biotropica*, 29, 145–150, 1997.

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Baker, T. R., Phillips, O. L., Malhi, Y., Almeida, S., Arroyo, L., Di Fiore, A., Erwin, T., Higuchi, H., Killeen, T. J., Laurance, S. G., Laurance, W. F., Lewis, S. L., Monteagudo, A., Neill, D. A., Nuñez Vargas, P., Pitman, N. C. A., Silva, J. N. M., and Vásquez Martínes, R.: Increasing biomass in Amazonian forest plots, *Philos. Trans. R. Soc. London, Ser. B*, 359, 353–365, 2004a.

Baker, T. R., Phillips, O. L., Malhi, Y., Almeida, S., Arroyo, L., Di Fiore, A., Erwin, T., Killeen, T. J., Laurance, S. G., Laurance, W. F., Lewis, S. L., Lloyd, J., Monteagudo, A., Neill, D. A., Patiño, S., Pitman, N. C. A., Silva, J. N. M., and Vásquez Martínes, R.: Variation in wood density determines spatial patterns in Amazonian biomass, *Global Change Biol.*, 10, 1–18, 2004b.

Baltzer, J. L., Davies, S. J., Bunyavejchewin, S., and Noor, N. S. M.: The role of desiccation tolerance in determining tree species distributions along the Malay–Thai Peninsula, *Funct. Ecol.*, 22, 221–231, 2008.

Banana, A. Y. and Tweheyo, M.: The ecological changes of Echuya afromontane bamboo forest, Uganda, *Afr. J. Ecol.*, 39, 366–373, 2001.

Baraloto, C. and Goldberg, D. E.: Microhabitat associations and seedling bank dynamics in a neotropical forest, *Oecologia*, 141, 701–712, 2004.

Baraloto, C., Morneau, F., Bonal, D., Blanc, L., and Ferry, B.: Seasonal water stress tolerance and habitat associations within four neotropical tree genera, *Ecology*, 88, 478–489, 2007.

Bongers, F., L. Poorter, R. S. A. R. Van Rompaey, and M. P. E. Parren: Distribution of twelve moist forest canopy tree species in Liberia and Côte d'Ivoire: response curves to a climatic gradient, *J. Veg. Sci.*, 10, 371–382, 1999.

Borchert, R.: Responses of tropical trees to rainfall seasonality and its long-term changes, *Climate Change*, 39, 381–393, 1998.

Brown, S. and Lugo, A. E.: Biomass of tropical forests: a new estimate based on forest volumes, *Science*, 223, 1290–1293, 1984.

Bucci, S. J., Goldstein, G., Meinzer, F. C., Scholz, F. G., and Franco, A. C.: Functional convergence in hydraulic architecture and water relations of tropical savanna trees: from leaf to whole plant, *Tree Physiol.*, 24, 891–899, 2004.

Chave, J., Muller-Landau, H. C., Baker, T. R., Easdale, T. A., Ter Steege, H., and Webb, C. O.: Regional and phylogenetic variation of wood density across 2456 neotropical tree species, *Ecol. Appl.*, 16, 2356–2367, 2006.

Condit, R., Pitman, N., Leigh Jr., E. G., Chave, J., Terborgh, J., Foster, R. B., Nuñez-V., P.,

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Aguilar, S., Valencia, R., Villa, G., Muller-Landau, H. C., Losos, E., and Hubbell, S. P.: Beta-Diversity in Tropical Forest Trees, *Science*, 295, 666–669, 2002.
- Engelbrecht, B. M. J., Comita, L. S., Condit, R., Kursar, T. A., Tyree, M. T., Turner, B. L. and Hubbell, S. P.: Drought sensitivity shapes species distribution patterns in tropical forests, 5 *Nature*, 447, 80–83, 2007.
- Engelbrecht, B. M. J., Dalling, J. W., Pearson, T. R. H., Wolf, R. L., Gálvez, D. A., Koehler, T., Tyree, M. T., and Kursar, T. A.: Short dry spells in the wet season increase mortality of tropical pioneer seedlings, *Oecologia*, 148, 258–269, 2006.
- Engelbrecht, B. M. J., Kursar, T. A., and Tyree, M. T.: Drought effects on seedling survival in a 10 tropical moist forest, *Trees-Struct. Funct.*, 19, 312–321, 2005.
- Engelbrecht, B. M. J., Wright, B. W., and De Steven, D.: Survival and ecophysiology of tree seedlingsc during El Niño drought in a tropical moist forest in Panama, *J. Trop. Ecol.*, 18, 569–579, 2002.
- Enquist, B. J., West, G. B., Charnov, E. L., and Brown, J. H.: Allometric scaling of production 15 and life-history variation in vascular plants, *Nature*, 401, 907–911, 1999.
- Erskine, P. D., Lamb, D., and Borschmann, G.: Growth performance and management of a mixed rainforest tree plantation, *New Forest*, 29, 117–134, 2005.
- Gentry, A. H.: Tree species richness of upper Amazonian forests, *P. Natl. Acad. Sci. USA*, 85, 156–159, 1988.
- Gilmour, A. R., Thompson, R., and Cullis, B. R.: Average information REML, an efficient 20 algorithm for variance parameterisation in linear mixed models, *Biometrics*, 51, 1440–1450, 1995.
- Gonzalez, E. and Fisher, R. F.: Variation in selected wood Properties of *Vochysia guatemalensis* from four sites in Costa Rica, *Forest Sci.*, 44, 185–191, 1998.
- Hacke, U. G., Sperry, J. S., Ewers, B. E., Ellsworth, D. S., Schafer, K. V. R., and Oren, R.: 25 Influence of soil porosity on water use in *Pinus taeda*, *Oecologia*, 124, 495–505, 2000.
- Hacke, U. G., Sperry, J. S., Pockman, W. T., Davis, S. D., and McCulloch, K. A.: Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure, *Oecologia*, 126, 457–461, 2001.
- Harms, K. E., Condit, R., Hubbell, S. P., and Foster, R. B.: Habitat associations of trees and 30 shrubs in a 50-ha neotropical forest plot, *J. Ecol.*, 89, 947–959, 2001.
- Hubbell, S. P.: The Unified Neutral Theory of Biodiversity and Biogeography, Princeton, Princeton Univ. Press, 448 pp., 2001.

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Jacobsen, A. L., Agenbag, L., Esler, K. J., Pratt, R. B., Ewers, F. W., and Davis, S. D.: Xylem density, biomechanics and anatomical traits correlate with water stress in 17 evergreen shrub species of the Mediterranean-type climate region of South Africa, *J. Ecol.*, 95, 171–183, 2007.
- 5 Jacobsen, A. L., Pratt, R. B., Ewers, F. W., and Davis, S. D.: Cavitation resistance among 26 chaparral species of Southern California, *Ecol. Monogr.*, 77, 99–115, 2007.
- King, D. A., Davies, S. J., Nur Supardi, M. N., and Tan, S.: Tree growth is related to light interception and wood density in two mixed dipterocarp forest of Malaysia, *Funct. Ecol.*, 19, 445–453, 2005.
- 10 King, D. A., Davies, S. J., Tan, S., and Nur Supardi, M. N.: The role of wood density and stem support costs in the growth and mortality of tropical trees, *J. Ecol.*, 94, 670–680, 2006.
- Kursar, T. A., Engelbrecht, B. M. J., and Tyree, M. T.: A comparison of methods for determining soil water availability in two sites in Panama with similar rainfall but distinct tree communities, *J. Trop. Ecol.*, 21, 297–305, 2005.
- 15 Laurance, W. F., Oliveira, A. A., Laurance, S. G., Condit, R., Nascimento, H. E. M., Sanchez-Thorin, A. C., Lovejoy, T. E., Andrade, A., D'angelo, S., Ribeiro, J. E., and Dick, C. W.: Pervasive alteration of tree communities in undisturbed Amazonian forest, *Nature*, 428, 171–175, 2004.
- Laurance, W. F., Fearnside, P. M., Laurance, S. G., Delamonica, P., Lovejoy, T. E., Rankin-De
20 Merona, J. M., Chambers, J. Q., and Gascon, C.: Relationship between soils and Amazon forest biomass: a landscape-scale study, *Forest Ecol. Manag.*, 118, 127–138, 1999.
- Madelaine, C., Pelissier, R., Vincent, G., Molino, J. F., Sabatier, D., Prevost, M. F., and De Namur, C.: Mortality and recruitment in a lowland tropical rain forest of French Guiana: effects of soil type and species guild, *J. Trop. Ecol.*, 23, 277–287, 2007.
- 25 Malhi, Y. and Grace, J.: Tropical forests and atmospheric carbon dioxide, *Tree*, 15, 332–337, 2000.
- Malhi, Y., Phillips, O. L., Lloyd, J., Baker, T., Wright, J., Almeida, S., Arroyo, L., Frederiksen, T., Grace, J., Higuchi, N., Killeen, T. J., Laurance, W. F., Leaño, C., Lewis, S. L., Meir, P., Monteagudo, A., Neill, D. A., Núñez Vargas, P., Panfil, S. N., Patiño, S., Pitman, N. C. A., Quesada, C. A., Rudas-Ll., A., Salomão, R., Saleska, S., Silva, N., Silveria, M., Sombroek, W. G., Valencia, R., Vásquez Martínes, R., Vieira, I., and Vinceti, B.: An international network to monitor the structure, composition and dynamics of Amazonian forests (RAINFOR), *J. Veg. Sci.*, 13, 439–450, 2002.

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Mass, P. J. M. and Westra, L. Y. T.: Neotropical plant families. Koenigstein, Germany, Koeltz Scientific Books, 290 pp., 1993.
- Meinzer, F. C., Andrade, A., Goldstein, G., Holbrook, N. M., Cavelier, J., and Wright, S. J.: Partitioning of soil water among canopy trees in a seasonally dry tropical forest, *Oecologia*, 121, 293–301, 1999.
- 5 Meinzer, F. C.: Functional convergence in plant responses to the environment, *Oecologia*, 134, 1–11, 2003.
- Muller-Landau, H. C.: Interspecific and inter-site variation in wood specific gravity of tropical trees, *Biotropica*, 36, 20–32, 2004.
- 10 Niklas, J. K.: Mechanical properties of black locust (*Robinia pseudoacacia*) wood: correlations among elastic and rupture moduli, proportional limit, and tissue density and specific gravity, *Ann. Bot.* 79, 479–485, 1997.
- Nogueira, E. M., Fearnside, P. M., Nelson, B. W., and França, M. B.: Wood density in forests of Brazil's "arc of deforestation": Implications for biomass and flux of carbon from land-use change in Amazonia, *Forest Ecol. Manag.*, 248, 119–135, 2007.
- 15 Nogueira, E. M., Nelson, B. W., and Fearnside, P. M.: Wood density in dense forest in central Amazonia, Brazil, *Forest Ecol. Manag.*, 208, 261–286, 2005.
- Parolin, P. and Ferreira, L. V.: Are there differences in specific wood gravities between trees in varzea and igapo (Central Amazonia)?, *Ecotropica*, 4, 25–32, 2004.
- 20 Pelissier, R., Dray, S., and Sabatier, D.: Within-plot relationships between tree species occurrences and hydrological soil constraints: an example in French Guiana investigated through canonical correlation analysis, *Plant Ecol.*, 162, 143–156, 2002.
- Peña-Claros, M.: Changes in forest structure and species composition during secondary forest succession in the Bolivian Amazon, *Biotropica*, 35, 450–461, 2003.
- 25 Phillips, O. L., Baker, T. R., Arroyo, L., Higuchi, H., Killeen, T. J., Laurance, W. F., Lewis, S. L., Lloyd, J., Malhi, Y., Monteagudo, A., Neill, D. A., Nuñez Vargas, P., Silva, J. N. M., Terborgh, J., Vásquez Martínez, R., Alexiades, M., Almeida, S., Brown, S., Chave, J., Comiskey, J. A., Czimczik, C. I., Di Fiore, A., Erwin, T., Kuebler, C., Laurance, S. G., Nascimento, H. E. M., Olivier, J., Palacios, W., Patiño, S., Pitman, N. C. A., Quesada, C. A., Saldías, M., Torres Lezama, A., and Vinceti, B.: Pattern and process in Amazon tree turnover 1976–2001, *Philos. Trans. R. Soc. London, Ser. B*, 359, 381–407, 2004.
- 30 Phillips, O. L. and Gentry, A. H.: Increasing turnover through time in tropical forests, *Science*, 263, 954–958, 1994.

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Phillips, O. L., Vargas, P. N., Monteagudo, A. L., Cruz, A. P., Zans, M. E. C., Sanchez, W. G., Yli-Halla, M., and Rose, S.: Habitat association among Amazonian tree species: a landscape-scale approach, *J. Ecol.*, 91, 757–775, 2003.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar D., and the R Core team: Linear and Nonlinear Mixed Effects Models, R package version 3.1-86, 2007.
- Pitman, N. C. A., Terborgh, J. W., Silman, M. R., Nuñez Vargas, P., Neill, D. A., Cerón, C. E., Palacios, W. A., and Aulestia, M.: Dominance and distribution of tree species in upper Amazonian terra firme forest, *Ecology*, 82, 2101–2117, 2001.
- Poorter, H. and Arends, E. J. M. M.: Light environment and tree strategies in a Bolivian tropical moist forest: an evaluation of the light partitioning hypothesis, *Plant Ecol.*, 166, 295–306, 2003.
- Poorter, L., Bongers, L., and Bongers, F.: Architecture of 54 moist-forest tree species: traits, trade-offs, and functional groups laurent bongers, 1, 2 and frans bongers, *Ecology*, 87, 1289–1301, 2006.
- Pratt, R. B., Jacobsen, A. L., Ewers, F. W., and Davis, S. D.: Relationships among xylem transport, biomechanics and storage in stems and roots of nine Rhamnaceae species of the California chaparral, *New Phytol.*, 174, 787–798, 2007.
- R Development Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, ISBN 3-900051-07-0, <http://www.R-project.org>, 2007.
- Robert, J., Dalling, J. W., Harms, K. E., Yavitt, J. B., Stallard, R. F., Mirabello, M., Hubbell, S. P., Valencia, R., Navarrete, H., Vallejo, M., and Foster, R. B.: Soil nutrients influence spatial distributions of tropical tree species, *P. Natl. Acad. Sci. USA*, 104, 864–869, 2007.
- Roque, R. M.: Effect of management treatment and growing regions on wood properties of *Gmelina arborea* in Costa Rica, *New Forest*, 28, 325–330, 2004.
- Sabatier, D., Grimaldi, M., Prevost, M. F., Guillaume, J., Godron, M., Dosso, M., and Curni, P.: The influence of soil cover organization on the floristic and structural heterogeneity of a Guianan rain forest, *Plant. Ecol.*, 131, 81–108, 1997.
- Santiago, L. S., Kaoru, K., Joseph, W., and Mulkey, S.: Coordinated changes in photosynthesis, water relations and leaf nutritional traits of canopy trees along a precipitation gradient in lowland tropical forest, *Oecologia*, 139, 495–502, 2004.
- Scholz, F. G., Bucci, S. J., Goldstein, G., Meinzer, F. C., Franco, A. C., and Miralles-Wilhelm, F.: Biophysical properties and functional significance of stem water storage tissues in Neotropi-

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- cal savanna trees, *Plant. Cell. Environ.*, 30, 236–248, 2007.
- Searle, S. R., Casella, G., and Mcculloch, C. E.: *Variance Components*, USA, Wiley Interscience, 2006.
- Selway, N. G. and Anten, N. P. R.: Differences in biomass allocation, light interception and mechanical stability between lianas and trees in early secondary tropical forest, *Funct. Ecol.*, 5, 30–39, 2008.
- Soininen, J., Mcdonald, R., and Hillebrand, H.: The distance decay of similarity in ecological communities, *Ecography*, 30, 3–12, 2007.
- Stratton, L., Goldstein, G., and Meinzer, F. C.: Stem water storage capacity and efficiency of water transport: their functional significance in a Hawaiian dry forest, *Plant Cell Environ.*, 23, 10, 99–106, 2000.
- Taneda, H. and Tateno, M.: The criteria for biomass partitioning of the current shoot: water transport versus mechanical support, *Am. J. Bot.*, 91, 1949–1959, 2004.
- ter Steege, H. and Hammond, D. S.: Character convergence, diversity, and disturbance in 15 tropical rain forest in Guyana, *Ecology*, 82, 3197–3212, 2004.
- ter Steege, H., Pitman, N. C. A., Phillips, O. L., Chave, J., Sabatier, D., Duque, A., Molino, J. F., Prévost, M. F., Spchiger, R., Castellanos, H., Von Hildebrand, P., and Vásquez, R.: Continental-scale patterns of canopy tree composition and function across Amazonia, *Nature*, 443, 444–447, 2006.
- Terpstra, J. T. and McKean, J. W.: Rank Based analysis of linear models using R, *J. Stat. Soft.* 20 14, 7, 2005.
- Tuomisto, H., Ruokolainen, K., Kalliola, R., Linna, A., Danjoy, W., and Rodriguez, Z.: Dissecting Amazonian biodiversity, *Science*, 269, 63–66, 1995.
- Tuomisto, H., Ruokolainen, K., and Yli-Halla, M.: Dispersal, environment and floristic variation 25 of western Amazonia forests, *Science*, 299, 241–244, 2003.
- Tyree, M. T., Engelbrecht, B. M. J., Vargas, G., and Kursar, T. A.: Desiccation tolerance of five tropical seedlings in Panama. Relationship to a field assessment of drought performance, *Plant Physiol.*, 132, 1439–1447, 2003.
- Valladares, F., Wright, J., Lasso, E., Kitajima, K., and Pearcy, R. W.: Plastic phenotypic response to light of 16 congeneric shrubs from a Panamanian rainforest, *Ecology*, 81, 1925–30 1936, 2000.
- Veenendaal, E. M. and Swaine, M. D.: Limits to tree species distribution in lowland tropical rainforests. *Dynamics of Tropical Forest Communities*, 37th Symposium of the British Ecological

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Society, Blackwell Science, Oxford, 163–191, 1998.

Vieira, I. C. G. and Proctor, J.: Mechanisms of plant regeneration during succession after shifting cultivation in eastern Amazonia, *Plant Ecol.*, 192, 303–315, 2007.

Vieira, I. C. G., Silva De Almeida, A., Davidson, E. A., Stone, T. A., Reis De Carvalho, C. J., and Guerrero, J. B.: Classifying successional forests using Landsat spectral properties and ecological characteristics in eastern Amazônia, *Remote Sense Environ.*, 87, 470–481, 2003.

Vinceti, B.: Spatial and temporal analysis of Floristic composition and Dynamics in some low-land Amazonian forests, Ph.D. thesis, Institute of Ecology and Resources Management, Edinburgh University, Scotland, 229 pp., 2003.

Vormisto, J., Phillips, O. L., Ruokolainen, K., Tuomisto, H., and Vasquez, R.: A comparison of fine-scale distribution patterns of four plant groups in an Amazonian rainforest, *Ecography*, 23, 349–359, 2000.

Wagner, K. R., Ewers, F. W., and Davies, S. D.: Tradeoffs between hydraulic efficiency and mechanical strength in the stems of four co-occurring species of chaparral shrubs, *Oecologia*, 117, 53–62, 1998.

Warton, D. I., Wright, I. J., Falster, D. S., and Westoby, M.: Bivariate line-fitting methods for allometry, *Biol. Rev.*, 81, 259–291, 2006.

Whitmore, T. C.: Canopy Gaps and the Two Major Groups of Forest Trees, *Ecology*, 70, 536–538, 1989.

Wiemann, M. C. and Williamson, G. B.: Wood specific gravity gradients in tropical dry and montane rain forest trees, *Am. J. Bot.*, 76, 924–928, 1989.

Woodcock, D. W., Dos Santos, G., and Reynel, C.: Wood characteristics of Amazon forest types, *Iawa Journal*, 21, 277–292, 2000.

Woodward, F. I.: Climate and plant distribution, Cambridge: Cambridge University Press, 192 pp., 1987.

Yang, L. C., Lee, C. H., and Chiu, C. M.: Genetic variation of wood density in Luanta fir tested in central Taiwan, *Wood Fiber Sci.*, 33, 486–491, 2001.

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table A1. More precise coordinates will be available (Anderson and Malhi, 2008⁸). Abbreviations in regions are: AC=Acre, AM=Amazonas, MT=Mato Grosso, CP=Central Pará, EP=East Par, WP=West Par, N=North, S=South, NE=North East, SW=South West,. ** not a permanent plot, samples were taken from trees around the Eddy covariance tower. Additional information of plots may be found in: (Malhi et al., 2002; 2003; Baker et al., 2004; Vinceti, 2003).

Plot Name and Description	Region Code	Region	Plot Code	latitude	longitude	Altitude (m)	Mean T (°C)	Forest Type	Principal Investigator
Sinop	1	MT-Brazil	SIN-01	-11.41	-55.33	325	25.4	Terra firme	M. Silveria
Alta Floresta	1	MT-Brazil	ALF-01	-9.60	-55.94	255	25.6	Terra firme	M. Silveria
Los Fierros Bosque I	2	Bolivia	LFB-01	-14.56	-60.93	230	25.1	Terra firme	T. Killeen
Los Fierros Bosque II	2	Bolivia	LFB-02	-14.58	-60.83	225	25.1	Terra firme	T. Killeen
Huanchaca Dos, plot1	2	Bolivia	HCC-21	-14.56	-60.75	720	25.1	Gallery	L. Arollo
Huanchaca Dos, plot2	2	Bolivia	HCC-22	-14.57	-60.75	735	25.1	Gallery	L. Arollo
Las Londras, plot 1	2	Bolivia	LSL-01	-14.41	-61.14	170	25.9	Seasonally flooded	L. Arollo
Las Londras, plot 2	2	Bolivia	LSL-02	-14.41	-61.14	170	25.9	Seasonally flooded	L. Arollo
Chore 1	2	Bolivia	CHO-01	-14.39	-61.15	170	25.9	Liana forest	T. Killeen
Tambopata plot zero	3	S-Peru	TAM-01	-12.84	-69.29	205	25.1	Terra firme	O. Phillips and R. Vasquez
Tambopata plot one	3	S-Peru	TAM-02	-12.84	-69.29	210	25.1	Terra firme	O. Phillips and R. Vasquez
Tambopata plot two swamp	3	S-Peru	TAM-03	-12.84	-69.28	205	25.1	Swamp	O. Phillips and R. Vasquez
Tambopata plot two swamp edge clay	3	S-Peru	TAM-04	-12.84	-69.28	205	25.1	Terra firme	O. Phillips and R. Vasquez
Tambopata plot three	3	S-Peru	TAM-05	-12.83	-69.27	220	25.1	Terra firme	O. Phillips and R. Vasquez
Tambopata plot four (cerca rio)	3	S-Peru	TAM-06	-12.84	-69.30	200	25.1	Terra firme	O. Phillips and R. Vasquez
Tambopata plot six	3	S-Peru	TAM-07	-12.83	-69.26	225	25.1	Terra firme	O. Phillips and R. Vasquez
Cuzco Amazonico, CUZAM2E	3	S-Peru	CUZ-03	-12.50	-68.96	195	25.1	Terra firme	O. Phillips and R. Vasquez
Jurua, PAA-05	3	AC-Brazil	PAA-05	-8.88	-72.79	245	26.2	Terra firme	M. Silveira
RESEX Alto Juruá:	3	AC-Brazil	RES-02	-9.04	-72.27	275	25.9	Terra firme	M. Silveira
Seringal Restauração									
RESEX Chico Mendes:	3	AC-Brazil	RES-03	-10.82	-68.78	275	25.8	Terra firme	M. Silveira
Seringal Porongaba 1									
RESEX Chico Mendes:	3	AC-Brazil	RES-04	-10.80	-68.77	270	25.8	Terra firme	M. Silveira
Seringal Porongaba 2									
RESEX Chico Mendes:	3	AC-Brazil	RES-05	-10.57	-68.31	200	26.0	Terra firme	M. Silveira
Seringal Dois Irmãos 1									
RESEX Chico Mendes:	3	AC-Brazil	RES-06	-10.56	-68.30	210	26.0	Bamboo forest	M. Silveira
Seringal Dois Irmãos 2									

Table A1. Continued.

Plot Name and Description	Region Code	Region	Plot Code	latitude	longitude	Altitude (m)	Mean T (°C)	Forest Type	Principal Investigator
Alpahuayo A, poorly drained	4	N-Peru	ALP-11	-3.95	-73.43	125	26.5	Terra firme	O. Phillips and R. Vasquez
Alpahuayo A, well drained	4	N-Peru	ALP-12	-3.95	-73.43	125	26.5	Terra firme	O. Phillips and R. Vasquez
Alpahuayo B, sandy	4	N-Peru	ALP-21	-3.95	-73.43	125	26.5	Terra firme	O. Phillips and R. Vasquez
Alpahuayo B, clayed	4	N-Peru	ALP-22	-3.95	-73.43	115	26.4	Terra firme	O. Phillips and R. Vasquez
Alpahuayo C	4	N-Peru	ALP-30	-3.95	-73.43	125	26.4	Tall caatinga	O. Phillips and R. Vasquez
Sucursari A	4	N-Peru	SUC-01	-3.23	-72.90	110	26.4	Terra firme	O. Phillips and R. Vasquez
Sucursari B	4	N-Peru	SUC-02	-3.23	-72.90	110	26.4	Terra firme	O. Phillips and R. Vasquez
Sucursari C	4	N-Peru	SUC-03	-3.25	-72.93	110	26.4	Seasonally flooded	O. Phillips, A. Monteagudo
Sucursari D	4	N-Peru	SUC-04	-3.25	-72.89	160	26.4	Terra firme	O. Phillips, A. Monteagudo, T. Baker
Yanamono A	4	N-Peru	YAN-01	-3.43	-72.85	105	26.4	Terra firme	O. Phillips and R. Vasquez
Yanamono B	4	N-Peru	YAN-02	-3.43	-72.84	105	26.4	Terra firme	O. Phillips and R. Vasquez
Jenaro Herrera A- Clay rich high terrace	4	N-Peru	JEN-11	-4.88	73.63	130	26.8	Terra firme	T.R. Baker and O. Phillips
Jenaro Herrera B- sandy	4	N-Peru	JEN-12	-4.90	-73.63	130	26.8	Terra firme	T.R. Baker and O. Phillips
Sumaco	5	Ecuador	SUM-01	-1.75	-77.63	1200	—	Premontane forest	D. Neill
Jatun Sacha 2	5	Ecuador	JAS-02	-1.07	-77.60	435	23.3	Terra firme	D. Neill
Jatun Sacha 3	5	Ecuador	JAS-03	-1.07	-77.67	410	23.3	Terra firme	D. Neill
Jatun Sacha 4	5	Ecuador	JAS-04	-1.07	-77.67	430	23.3	Terra firme	D. Neill
Jatun Sacha 5	5	Ecuador	JAS-05	-1.07	-77.67	395	23.3	Terra firme	D. Neill
Bogi 1	5	Ecuador	BOG-01	-0.70	-76.48	270	26.0	Terra firme	N. Pitman, and T. DiFiore
Bogi 2	5	Ecuador	BOG-02	-0.70	-76.47	270	26.0	Terra firme	N. Pitman, and T. DiFiore
Tiputini 3	5	Ecuador	TIP-03	-0.64	-76.16	250	26.0	Seasonally flooded	N. Pitman
Tiputini 5	5	Ecuador	TIP-05	-0.64	-76.14	245	26.0	Terra firme	N. Pitman
Amacayacu: Lorena E	6	Colombia	LOR-01	-3.06	-69.99	95	25.9	Terra firme	A. Rudas and A. Prieto
Amacayacu: Lorena U	6	Colombia	LOR-02	-3.06	-69.99	95	25.9	Terra firme	A. Rudas and A. Prieto
Amacayacu: Agua Pudre E	6	Colombia	AGP-01	-3.72	-70.31	105	25.8	Terra firme	A. Rudas and A. Prieto
Amacayacu: Agua Pudre U	6	Colombia	AGP-02	-3.72	-70.30	110	25.8	Terra firme	A. Rudas and A. Prieto
EL Zafire: Varillal	6	Colombia	ZAR-01	-4.01	-69.91	130	25.6	Caatinga	M.C.Penuela and E. Alvarez
EL Zafire: Rebalse	6	Colombia	ZAR-02	-4.00	-69.90	120	25.6	Seasonally flooded	M.C.Penuela and E. Alvarez
EL Zafire: Terra Firme	6	Colombia	ZAR-03	-3.99	-69.90	135	25.6	Terra firme	M.C.Penuela and E. Alvarez
EL Zafire: Altura	6	Colombia	ZAR-04	-4.01	-69.90	120	25.6	Terra firme	M.C.Penuela and E. Alvarez
San Carlos Oxisol	7	SW-Venezuela	SCR-01	1.93	-67.02	120	26.0	Terra firme	R. Herrera

Amazonian xylem density variation

S. Patiño et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table A1. Continued.

Plot Name and Description	Region Code	Region	Plot Code	latitude	longitude	Altitude (m)	Mean T (°C)	Forest Type	Principal Investigator
San Carlos Tall Caatinga	7	SW-Venezuela	SCR-04	1.93	-67.04	120	26.0	Tall caatinga	R. Herrera
San Carlos Yévaro	7	SW-Venezuela	SCR-05	1.93	-67.04	120	26.0	Terra firme	R. Herrera
Rio Grande, plots DA1 (RIO-01) and DA2 (RIO-02)	8	NE-Venezuela	RIO-12	8.11	-61.69	270	24.9	Terra firme	A. Torres-Lezama
El Dorado, km 91, plots G1 (ELD-01) and G2 (ELD-02)	8	NE-Venezuela	ELD-12	6.10	-61.40	200	24.9	Terra firme	A. Torres-Lezama
El Dorado, km 98, plots G3 (ELD-03) and G4 (ELD-04)	8	NE-Venezuela	ELD-34	6.08	-61.41	360	24.9	Terra firme	A. Torres-Lezama
Manaus K34, plato	9	AM-Brazil	MAN-01	-2.61	-60.21	65	27.3	Terra firme	N. Higuchi
Manaus K34, vertiente	9	AM-Brazil	MAN-02	-2.61	-60.21	50	27.3	Terra firme	N. Higuchi
Manaus K34, campinarana	9	AM-Brazil	MAN-03	-2.60	-60.22	65	27.3	Tall caatinga	N. Higuchi
Manaus K34, baxio	9	AM-Brazil	MAN-04	-2.61	-60.22	45	27.3	Caatinga/swampy valley	N. Higuchi
Bionte 4: Manaus K 23	9	AM-Brazil	BNT-04	-2.63	-60.15	105	27.3	Terra firme	N. Higuchi
Manaus K14. Tower**	9	AM-Brazil	MAN-05	-2.59	-60.12	108	27.3	??	??
Tapajos, RP014, 1-4	10	WP-Brazil	TAP-01	-3.31	-54.94	187	26.5	Terra firme	N. Silva
Tapajos, RP014, 5-8	10	WP-Brazil	TAP-02	-3.31	-54.95	187	26.5	Terra firme	N. Silva
Tapajos, RP014, 9-12	10	WP-Brazil	TAP-03	-3.31	-54.94	187	26.5	Terra firme	N. Silva
Tapajos, LBA Tower, Transects 1, 2, 3 and 4	10	WP-Brazil	TAP-04	-2.85	-54.96	73	26.5	Terra firme	S. Saleska, Hammond-Pyle, Hytira, Wofsy,de Camargo, Vieira
Caxiuaná 1	11	CP-Brazil	CAX-01	-1.74	-51.46	40	25.6	Terra firme	S. Almeida
Caxiuaná 2	11	CP-Brazil	CAX-02	-1.74	-51.46	40	25.6	Terra firme	S. Almeida
Caxiuaná 3: A (Control drought experiment). Esecaflor	11	CP-Brazil	CAX-03	-1.73	-51.46	15	25.6	Terra firme	S. Almeida, A. L. da Costa, L de Sa, J. Grace, P. Meir and Y. Malhi
Caxiuaná 4: B (Drought experiment). Esecaflor	11	CP-Brazil	CAX-04	-1.73	-51.46	15	25.6	Terra firme	S. Almeida, A. L. da Costa, L de Sa, J. Grace, P. Meir and Y. Malhi
Caxiuaná 5: Eddy tower	11	CP-Brazil	CAX-05	-1.72	-51.46	15	25.6	Terra firme	S. Almeida, L de Sa, J. Grace, P. Meir and Y. Malhi
Jari 1	11	CP-Brazil	JRI-01	-0.89	-52.19	127	26.5	Terra firme	N. Silva
Braganca	12	EP-Brazil	BRA-01	-0.83	-46.64	10	25.8	??	A. L. da Costa and Y. Malhi
Mocambo 1	12	EP-Brazil	MBO-01	-1.45	-48.45	24	26.8	Terra firme	R. Salomao
Capitão Poço	12	EP-Brazil	CPP-01	-2.19	-47.33	66	25.9	Terra firme	I. Viera and E. Leal
Guyafux 7	13	Guiana	GFX-07			10	25.7	flooded	D. Bonal
Guyafux 9	13	Guiana	GFX-09			40	25.7	Terra firme	D. Bonal

Amazonian xylem density variation

S. Patiño et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table B1. D_x distribution across the Amazon basin. Plots and Regions are indicated in the right and left axes respectively. Horizontal lines represent the standard deviation. Vertical straight lines represent the Tukey test in which regions joined by the line are not significantly different while regions not included within a line are significant different.

BGD

5, 2003–2047, 2008

Region/Country	DF	F	P	Plot Code	N	Mean	SE Mean	StDev
CP-Brazil 0.757 (0.132)	5	4.32	0.001	CAX-02 *	15	0.669	0.052	0.203
				CAX-05	19	0.733	0.019	0.084
				CAX-01	20	0.740	0.037	0.166
				JRI-01	20	0.757	0.026	0.116
				CAX-03 *	38	0.788	0.018	0.112
				CAX-04 *	32	0.797	0.019	0.105
AM-Brazil 0.702 (0.082)	5	2.11	0.74	MAN-02 *	6	0.639	0.043	0.106
				MAN-04	10	0.675	0.023	0.072
				MAN-01	13	0.688	0.019	0.067
				MAN-05	20	0.694	0.021	0.095
				MAN-03	9	0.729	0.032	0.097
				BNT-04 *	21	0.736	0.011	0.051
EP-Brazil 0.668 (0.109)	3	4.58	0.014	MBO-01	18	0.627	0.030	0.126
				CPP-01	20	0.649	0.021	0.092
				BRA-01 *	20	0.723	0.020	0.091
				YAN-02 *	8	0.521	0.023	0.065
				YAN-01 *	17	0.570	0.024	0.101
				SUC-01 *	19	0.629	0.028	0.120
N-Peru 0.664 (0.117)	12	4.89	<0.001	ALP-12	9	0.644	0.021	0.064
				SUC-04	20	0.657	0.017	0.077
				JEN-11	19	0.659	0.032	0.138
				SUC-02	16	0.659	0.022	0.090
				ALP-11	10	0.672	0.036	0.114
				ALP-22	12	0.678	0.022	0.075
				SUC-03	18	0.694	0.030	0.125
				ALP-21	6	0.720	0.055	0.135
				JEN-12 *	20	0.746	0.026	0.115
				ALP-30 *	12	0.765	0.017	0.058
				TAP-04 *	33	0.627	0.024	0.136
				TAP-01	16	0.659	0.028	0.113
				TAP-03	20	0.673	0.014	0.060
				TAP-02 *	19	0.722	0.022	0.096
WP-Brazil 0.663 (0.114)	3	3.07	0.032	SCR-04	26	0.594	0.019	0.098
				SCR-05	34	0.596	0.017	0.101
				SCR-01	21	0.653	0.026	0.117

Amazonian xylem density variation

S. Patiño et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table B1. Continued.

BGD

5, 2003–2047, 2008

Region/Country	DF	F	P	Plot Code	N	Mean	SE Mean	StDev
S-Peru & AC-Brazil 0.589 (0.100)	13	2.90	0.001	TAM-03 *	6	0.468	0.040	0.098
				CUZ-03	23	0.573	0.015	0.070
				TAM-01	22	0.578	0.021	0.099
				TAM-04	15	0.588	0.025	0.095
				TAM-06	21	0.588	0.018	0.081
				TAM-02	19	0.625	0.026	0.114
				TAM-07	20	0.637	0.022	0.100
				TAM-05	20	0.642	0.020	0.090
				POR-01	19	0.545	0.023	0.100
				DOI-02	18	0.551	0.022	0.093
				POR-02	20	0.557	0.024	0.105
				RST-01	20	0.583	0.024	0.107
				DOI-01	18	0.613	0.018	0.075
				JUR-01	13	0.634	0.030	0.109
MT-Brazil 0.575 (0.093)	1	9.55	0.004	ALF-01 *	26	0.543	0.013	0.068
				SIN-01 *	17	0.625	0.026	0.105
NE-Venezuela 0.568 (0.125)	2	1.29	0.284	ELD-34	16	0.528	0.030	0.121
				RIQ-12	19	0.582	0.024	0.102
				ELD-12	16	0.593	0.037	0.149
Bolivia 0.561 (0.106)	6	0.74	0.62	LSL-02	16	0.530	0.021	0.085
				CHO-01	18	0.549	0.025	0.107
				HCC-22	21	0.550	0.025	0.114
				LFB-01	18	0.560	0.023	0.098
				LSL-01	14	0.569	0.037	0.140
				HCC-21	20	0.574	0.027	0.121
				LFB-02	16	0.601	0.017	0.068
Colombia 0.593 (0.103)	7	8.00	< 0.001	ZAR-02	20	0.572	0.019	0.084
				ZAR-03	18	0.612	0.018	0.078
				ZAR-04	20	0.616	0.018	0.082
				ZAR-01 *	20	0.712	0.020	0.088
				LOR-02	16	0.513	0.033	0.130
				AGP-02	20	0.545	0.019	0.087
				AGP-01	20	0.574	0.022	0.097
				LOR-01	17	0.582	0.020	0.081
Ecuador 0.535 (0.089)	8	2.32	0.021	JAS-05*	20	0.472	0.016	0.074
				SUM-01	18	0.510	0.016	0.069
				JAS-03	19	0.526	0.017	0.073
				JAS-02	21	0.531	0.019	0.086
				BOG-02	33	0.536	0.017	0.100
				TIP-03	20	0.550	0.012	0.054
				BOG-01	44	0.554	0.015	0.098
				JAS-04	22	0.559	0.018	0.084
				TIP-05	11	0.568	0.035	0.115



Amazonian xylem density variation

S. Patiño et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table C1. Regression analysis outputs for family (D1) genera (D2) and species (D3). See Figs. (5A), (5B) and (5C). Notice that for this Appendix units for D_x are given in g cm^{-3} .

Family	n	R ²	p	Slope	LowCI	UppCI	Interc	LowCI	UppCI	Ymean	Xmean
<i>Fabaceae</i>	64	0.679	<0.001	1.1629	1.0076	1.3421	-0.08505	-0.1903	0.02019	0.641	0.625
<i>Sapotaceae</i>	53	0.371	<0.001	1.1979	0.9602	1.4945	-0.10343	-0.27174	0.064871	0.645	0.625
<i>Moraceae</i>	48	0.263	<0.001	1.5312	1.19	1.9702	-0.34804	-0.58443	-0.11165	0.573	0.601
<i>Myristicaceae</i>	44	0.235	0.001	1.6258	1.2423	2.1279	-0.44166	-0.71386	-0.16946	0.548	0.609
<i>Lecythidaceae</i>	41	0.414	<0.001	1.2409	0.9709	1.5861	-0.10415	-0.29929	0.09099	0.677	0.629
<i>Burseraceae</i>	37	0.281	0.001	1.1498	0.8629	1.532	-0.1152	-0.32422	0.093814	0.597	0.619
<i>Lauraceae</i>	37	0.345	<0.001	1.7984	1.3671	2.3659	-0.52294	-0.83807	-0.20781	0.603	0.626
<i>Annonaceae</i>	33	0.539	<0.001	1.9531	1.5269	2.4984	-0.64874	-0.94373	-0.35376	0.528	0.603
<i>Euphorbiaceae</i>	30	0.411	<0.001	1.2956	0.9666	1.7365	-0.20745	-0.44246	0.027548	0.578	0.606
<i>Malvaceae</i>	27	0.457	<0.001	1.8604	1.3796	2.5087	-0.55923	-0.89383	-0.22463	0.536	0.589
<i>Meliaceae</i>	27	0.299	0.003	1.5244	1.0869	2.138	-0.28627	-0.596	0.02345	0.607	0.586
<i>Chrysobalanaceae</i>	26	0.477	<0.001	1.4788	1.0952	1.9967	-0.21232	-0.50491	0.080268	0.74	0.644
<i>Urticaceae</i>	20	0.019	0.557	0.9531	0.5942	1.5289	-0.01937	-0.29461	0.255877	0.538	0.585
<i>Olacaceae</i>	17	0.575	<0.001	1.3773	0.969	1.9575	-0.15213	-0.46956	0.165301	0.725	0.637
<i>Rubiaceae</i>	17	0.238	0.047	1.4992	0.9431	2.3832	-0.24574	-0.668	0.176529	0.626	0.581
<i>Violaceae</i>	16	0.255	0.046	1.4414	0.895	2.3215	-0.31222	-0.76893	0.144483	0.603	0.635
<i>Apocynaceae</i>	14	0.271	0.056	1.9349	1.1571	3.2355	-0.56654	-1.26576	0.092692	0.668	0.648
<i>Clusiaceae</i>	12	0.443	0.018	1.4535	0.8779	2.4063	-0.27101	-0.74962	0.207597	0.634	0.623
<i>Melastomataceae</i>	12	0.546	0.006	1.3648	0.8629	2.1588	-0.17818	-0.56802	0.211669	0.637	0.597
<i>Salicaceae</i>	12	0.324	0.054	1.2708	0.7324	2.205	-0.11586	-0.54314	0.311417	0.618	0.577
<i>Myrtaceae</i>	11	0.004	0.849	1.1165	0.5572	2.2374	-0.02775	-0.56646	0.510964	0.68	0.634
<i>Bignoniaceae</i>	10	0.568	0.012	1.2924	0.7736	2.1591	-0.21989	-0.64258	0.202799	0.563	0.606
<i>Sapindaceae</i>	10	0.199	0.196	1.0605	0.539	2.0867	-0.00513	-0.48015	0.469884	0.641	0.609
<i>Elaeocarpaceae</i>	9	0.471	0.041	1.254	0.6805	2.3108	-0.12868	-0.6266	0.369239	0.633	0.607

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table C1. Continued

Genera	n	R ²	p	Slope	LowCI	UppCI	Interc	LowCI	UppCI	Ymean	Xmean
<i>Eschweilera</i>	38	0.338	<0.001	1.208	0.9206	1.585	-0.08035	-0.2913	0.130594	0.68	0.629
<i>Pouteria</i>	33	0.296	0.001	1.0817	0.7992	1.4641	0.001807	-0.20788	0.211497	0.678	0.625
<i>Virola</i>	30	0.186	0.017	1.7252	1.2248	2.4299	-0.51423	-0.88692	-0.14154	0.542	0.612
<i>Inga</i>	29	0.457	<0.001	1.3303	0.9983	1.7727	-0.18135	-0.41348	0.050783	0.611	0.595
<i>Protium</i>	26	0.207	0.02	1.2025	0.8332	1.7355	-0.15964	-0.44904	0.129751	0.604	0.635
<i>Licania</i>	22	0.435	0.001	1.2808	0.9083	1.8062	-0.07048	-0.36435	0.223384	0.761	0.649
<i>Pououuma</i>	19	0.022	0.544	0.9597	0.5899	1.5612	-0.02545	-0.31228	0.261382	0.537	0.586
<i>Pseudolmedia</i>	18	0.241	0.039	1.2907	0.8257	2.0178	-0.17457	-0.52217	0.173022	0.575	0.581
<i>Brosimum</i>	16	0.002	0.858	1.8749	1.087	3.234	-0.59057	-1.25315	0.072006	0.559	0.613
<i>Ocotea</i>	15	0.201	0.094	1.0427	0.6243	1.7415	-0.07265	-0.42995	0.284653	0.589	0.634
<i>Tachigali</i>	15	0.653	<0.001	1.4829	1.0492	2.0957	-0.27776	-0.60405	0.048518	0.64	0.619
<i>Micropholis</i>	14	0.453	0.008	1.1959	0.7625	1.8757	-0.14913	-0.50019	0.201939	0.601	0.627
<i>Iryanthera</i>	13	0.362	0.03	1.0244	0.6165	1.7023	-0.06124	-0.39904	0.276568	0.572	0.618
<i>Guarea</i>	11	0.705	0.001	1.2925	0.8671	1.9267	-0.14834	-0.45503	0.158345	0.596	0.576
<i>Rinorea</i>	11	0.219	0.147	1.5798	0.8456	2.9515	-0.42908	-1.13096	0.272803	0.617	0.662
<i>Swartzia</i>	11	0.627	0.004	0.9748	0.6244	1.522	0.046443	-0.25213	0.345019	0.69	0.661
<i>Miconia</i>	10	0.207	0.187	2.068	1.054	4.0574	-0.58253	-1.46005	0.294994	0.623	0.583
<i>Minquartia</i>	10	0.721	0.002	1.5284	1.006	2.322	-0.27196	-0.71201	0.168102	0.743	0.664
<i>Tetragastris</i>	10	0.689	0.003	1.2043	0.7754	1.8703	-0.15214	-0.49909	0.194811	0.606	0.63
<i>Trichilia</i>	10	0.146	0.275	1.8155	0.9054	3.6406	-0.42628	-1.22306	0.370506	0.627	0.58
<i>Lecythis</i>	9	0.488	0.036	1.4459	0.7917	2.6408	-0.30197	-0.92802	0.324076	0.672	0.673
<i>Manilkara</i>	9	0.469	0.042	1.9199	1.0411	3.5408	-0.61887	-1.46185	0.22411	0.666	0.669
<i>Aspidosperma</i>	8	0.455	0.066	1.4701	0.7426	2.9102	-0.19256	-0.89258	0.507471	0.75	0.641

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Amazonian xylem density variation

S. Patiño et al.

Table C1. Continued.

Group	n	R ²	p	Slope	LowCI	UppCI	Interc	LowCI	UppCI	Ymean	Xmean
<i>Eschweilera coriacea</i>	14	0.461	0.008	0.9686	0.6197	1.514	0.09245	-0.18433	0.3692	0.687	0.614
<i>Pseudolmedia laevis</i>	12	0.334	0.049	1.8039	1.0434	3.1185	-0.47096	-1.0828	0.1409	0.589	0.588
<i>Minquartia guianensis</i>	9	0.678	0.006	1.5032	0.9229	2.4485	-0.25345	-0.77122	0.2643	0.76	0.674
<i>Virola-calophylla</i>	8	0.84	0.001	1.727	1.1702	2.5488	-0.5144	-0.9361	-0.093	0.537	0.609
<i>Licania-heteromorpha</i>	7	0.445	0.101	0.9109	0.4193	1.9789	0.21779	-0.30486	0.7404	0.825	0.667
<i>Rinorea-guianensis</i>	7	0.213	0.297	2.5887	1.0575	6.337	-1.18212	-3.03119	0.667	0.627	0.699
<i>Virola-pavonis</i>	7	0.441	0.104	1.363	0.6259	2.9681	-0.32976	-1.10846	0.4489	0.573	0.662
<i>Iryanthera-juruensis</i>	6	0.485	0.124	0.729	0.3027	1.7556	0.10677	-0.35901	0.5726	0.571	0.636
<i>Micropholis-guyanensis</i>	6	0.711	0.035	1.6374	0.821	3.2654	-0.44742	-1.24398	0.3491	0.617	0.65
<i>Pourouma-guianensis</i>	6	0.45	0.144	0.7968	0.3234	1.9634	0.05047	-0.42126	0.5222	0.507	0.573
<i>Pourouma-minor</i>	6	0.112	0.517	0.7733	0.2617	2.2851	0.10734	-0.46931	0.684	0.546	0.567
<i>Pseudolmedia-macrophylla</i>	6	0.69	0.041	1.8901	0.9279	3.85	-0.5422	-1.39521	0.3108	0.56	0.583
<i>Tetragastris-altissima</i>	6	0.169	0.418	1.7954	0.6238	5.1678	-0.49947	-1.8346	0.8357	0.554	0.587

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	






Fig. 1. Schematic representation of regions across the Amazon Basin. Each symbol represent one plot and the size of the symbol shows the arithmetic mean D_x (kg m^{-3}). Coordinates were changed to avoid overlapping of points in the map, correct coordinates are in Appendix A. Numbers in blue indicate the number of each Region in which we have grouped the sampled plots. Abbreviations in regions follow those in legend for Appendix A. Regions are: 1. MT-Brazil; 2. Bolivia; 3. S-Peru and AC-Brazil; 4. N-Peru; 5. Ecuador; 6. Colombia; 7. SE-Venezuela; 8. NE-Venezuela; 9. AM-Brazil; 10. WP-Brazil; 11. CP Brazil 12. Brazil-EP, 13. F-Guiana.

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. 2. D_x distribution across the Amazon basin. Plots and Regions are indicated in the right and left axes respectively. Horizontal lines represent the standard deviation. Vertical straight lines represent the Tukey test in which regions joined by the line are not significantly different while regions not included within a line are significant different.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Amazonian xylem density variation

S. Patiño et al.




Fig. 3.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Fig. 3. Variation of D_x (kg m^{-3}) within and between families (genera within families). Each dot represents the average D_x of genera. Left vertical axes represent genera, right vertical axes represent families and X-axis is the D_x . Grey and white shadows separate the families. Vertical dashed line represents the mean D_x of the basin. Horizontal lines represent the Standard Deviation. Families in the Fig. are sorted from high to low D_x from top-right (panel A) to left-bottom (Panel C). The three panels (A, B, and C) represent one continuous Fig., divided only for purpose of presentation. Abbreviation of the families are listed below D_x : OLA= *Olacaceae*, CEL= *Celastraceae*, CHR= *Chrysobalanaceae*, HUM= *Humiriaceae*, OCH= *Ochnaceae*, LIN= *Linaceae*, RHI= *Rhizophoraceae*, SCR= *Scrophulariaceae*, APO= *Apocynaceae*, MYRT= *Myrtaceae*, LEC= *Lecythidaceae*, SIM= *Simaroubaceae*, SAPO= *Sapotaceae*, SAPI= *Sapindaceae*, AQUI= *Aquifoliaceae*, STYR= *Styracaceae*, FAB= *Fabaceae*, ELA= *Elaeocarpaceae*, CLU= *Clusiaceae*, CAR= *Caryocaraceae*, VIO= *Violaceae*, VER= *Verbenaceae*, RUB= *Rubiaceae*, NYC= *Nyctaginaceae*, ANA= *Anacardiaceae*, RUT= *Rutaceae*, MELA= *Melastomataceae*, ICA= *Icacinaceae*, SAL= *Salicaceae*, COM= *Combretaceae*, LAU= *Lauraceae*, ROS= *Rosaceae*, MELI= *Meliaceae*, ULM= *Ulmaceae*, BUR= *Burseraceae*, PRO= *Proteaceae*, MALP= *Malpighiaceae*, OLEA= *Oleaceae*, MON= *Monimiaceae*, POL= *Polygonaceae*, MOR= *Moraceae*, EUP= *Euphorbiaceae*, BIG= *Bignoniaceae*, DIC= *Dichapetalaceae*, ARAL= *Araliaceae*, VOC= *Vochysiaceae*, URT= *Urticaceae*, MYRI= *Myristicaceae*, ANN= *Annonaceae*, MAL= *Malvaceae*, STA= *Staphyleaceae*, RHA= *Rhamnaceae*, LAC= *Lacistemataceae*, LEP= *Lepidobotryaceae*, SAB= *Sabiaceae*, BIX= *Bixaceae*, and BOR= *Boraginaceae*

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Amazonian xylem density variation

S. Patiño et al.




Fig. 4. Apportion of the total variance of D_x in the data set.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 5A. Relationships between mean D_x of plot (X-axis) and mean D_x of each family (**A**), genera (**B**), and species (**C**) within each plot. For each regression line a plot mean was calculated excluding the family, genera or species for which the analysis was done and plotted against the average of that family, genus or species. Families used in the analysis were collected at least in 6, genera 8 and species 6 plots. Regression lines in blue where not highly significant although follow the general trend. No regression lines in panels (**A**) M and U; (**B**) G and I; and (**C**) F, K, and M indicate that there were not significant relationships. Analysis were performed with SMATR. Mean of plot without species (D_x , kg m^{-3}).

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. 5B. Mean of plot without species (D_x , kg m^{-3}).

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Fig. 5C. Mean of plot without species (D_x , kg m^{-3}).

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Amazonian xylem density variation

S. Patiño et al.




Fig. 6. Plot level (abundance weighted) D_x , kg m^{-3} .

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)

- [Printer-friendly Version](#)
- [Interactive Discussion](#)






Fig. 7. The relationship between **(a)** observed species level values for xylem density (D_x) obtained in the current study and species level mean values for wood density D_w obtained from the RAINFOR database and **(b)** deduced species level effects on D_x from the REML analysis of Eq. (1) and mean values of D_w from the RAINFOR database.

Amazonian xylem density variation

S. Patiño et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

