



HAL
open science

Uptake and reaction of HOBr on frozen and dry NaCl/NaBr surfaces between 253 and 233K

J. W. Adams, N. S. Holmes, J. N. Crowley

► **To cite this version:**

J. W. Adams, N. S. Holmes, J. N. Crowley. Uptake and reaction of HOBr on frozen and dry NaCl/NaBr surfaces between 253 and 233K. Atmospheric Chemistry and Physics Discussions, 2002, 2 (1), pp.109-145. hal-00300822

HAL Id: hal-00300822

<https://hal.science/hal-00300822>

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Uptake and reaction
of HOBr**

J. W. Adams et al.

Uptake and reaction of HOBr on frozen and dry NaCl/NaBr surfaces between 253 and 233 K

J. W. Adams^{1,*}, N. S. Holmes¹, and J. N. Crowley¹

¹Max-Planck-Institut für Chemie, Division of Atmospheric Chemistry, Postfach 3060, 55020 Mainz, Germany

*Present address: Dept. of Chemistry, University of Cambridge, Lensfield Rd, Cambridge CB2 1EW, UK

Received: 20 December 2001 – Accepted: 4 January 2002 – Published: 22 January 2002

Correspondence to: J. N. Crowley (Crowley@mpch-mainz.mpg.de)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGS 2002

Abstract

The uptake and reaction of HOBr with frozen salt surfaces of variable NaCl / NaBr composition and temperature were investigated with a coated wall flow tube reactor coupled to a mass spectrometer for gas-phase analysis. HOBr is efficiently taken up onto the frozen surfaces at temperatures between 253 and 233 K where it reacts to form the di-halogens BrCl and Br₂, which are subsequently released into the gas-phase. The uptake coefficient for HOBr reacting with a frozen, mixed salt surface of similar composition to sea-spray was $\approx 10^{-2}$. The relative concentration of BrCl and Br₂ released to the gas-phase was found to be strongly dependent on the ratio of Cl⁻ to Br⁻ in the solution prior to freezing / drying. For a mixed salt surface of similar composition to sea-spray the major product at low conversion of surface reactants (i.e. Br⁻ and Cl⁻) was Br₂.

Variation of the pH of the NaCl / NaBr solution used to prepare the frozen surfaces was found to have no significant influence on the results. The observations are explained in terms of initial formation of BrCl in a surface reaction of HOBr with Cl⁻, and conversion of BrCl to Br₂ via reaction of surface Br⁻. Experiments on the uptake and reaction of BrCl with frozen NaCl / NaBr solutions served to confirm this hypothesis. The kinetics and products of the interactions of BrCl, Br₂ and Cl₂ with frozen salt surfaces were also investigated, and lower limits to the uptake coefficients of > 0.034 , > 0.025 and > 0.028 respectively, were obtained. The uptake and reaction of HOBr on dry salt surfaces was also investigated and the results closely resemble those obtained for frozen surfaces. During the course of this study the gas diffusion coefficients of HOBr in He and H₂O were also measured as (273 ± 1) Torr cm² s⁻¹ and (51 ± 1) Torr cm² s⁻¹, respectively, at 255 K. The implications of these results for modelling the chemistry of the Arctic boundary layer in springtime are discussed.

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

1. Introduction

Field measurements have conclusively shown that the episodic, catalytic destruction of ozone in the polar tropospheric boundary layer is strongly correlated with the presence of reactive bromine compounds (Barrie et al., 1988, 1994; Barrie and Platt, 1997).

5 Time resolved measurement of the concentrations of alkanes, alkenes and alkynes (Jobson et al., 1994; Ramacher et al., 1997, 1999; Rudolph et al., 1999; Boudries and Bottenheim, 2000), detection of photo-labile bromine compounds such as HOBr, BrCl and Br₂ (Impey et al., 1997, 1999; Foster et al., 2001), the direct measurement of the BrO radical in ground-based experiments (Hausmann and Platt, 1994; Miller et al., 1997; Tuckermann et al., 1997), and by satellite (Chance, 1998), measurement of active chlorine via chemical amplification (Perner et al., 1999) and isotopic studies of CO (Röckmann et al., 1999) provide the experimental evidence for the presence of reactive, gas-phase bromine and chlorine species and indicate their central role in ozone destruction in several arctic locations (Oltmans, 1981; Barrie et al., 1988; Oltmans et al., 1989; Anlauf et al., 1994; Solberg et al., 1996; Lorenzen-Schmidt et al., 1998). Measurements in the Antarctic indicate that bromine catalysed ozone loss can take place even in this relatively pristine environment (Kreher et al., 1997; Wessel et al., 1998).

The chemical mechanism(s) by which high concentrations of reactive bromine and chlorine compounds are sustained are not yet completely characterised although an important role for heterogeneous reactions of HOBr (Fan and Jacob, 1992; McConnell et al., 1992; Mozurkewich, 1995; Tang and McConnell, 1996) has been identified.

15 In a recent publication (Fickert et al., 1999) we were able to show that the interaction of HOBr with aqueous NaCl / NaBr solutions of similar composition to sea-spray was an efficient source of Br₂ (and BrCl) if the solution pH was less than ≈ 6 . The uptake coefficient (see below) for the interaction of HOBr with salt containing deliquescent aerosols has been shown to be large (Abbatt and Waschewsky, 1998). The initial step

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

is thought to be an acid catalysed reaction between HOBr and Cl^- :



Rapid, aqueous phase equilibria result in the conversion of BrCl to Br_2 ,



5 and the relative concentrations of Br_2 and BrCl released to the gas-phase is controlled by aqueous phase equilibrium reactions of Br_2Cl^- (Wang et al., 1994; Fickert et al., 1999). Only when the $[\text{Br}^-] / [\text{Cl}^-]$ ratio was very low compared to the composition of sea-water, was BrCl released in significant yield.

10 In the present study we investigate the efficiency of di-halogen release from the interaction of HOBr with frozen salt surfaces that mimic the arctic snow/ice pack. Supplementary experiments that examined the uptake and reaction of HOBr on “dry” surfaces and the uptake and reaction of molecular di-halogens with frozen surfaces were also carried out to gain insight into the parameters that control the rate and mechanism of the chemical interaction. Surprisingly, with few exceptions, the results from this work
15 represent the first measurements of the kinetics and mechanisms of the reactions of halogenated species on mixed $\text{Cl}^- / \text{Br}^-$ surfaces made from aqueous solutions that were of similar composition to atmospheric sea-spray /sea-water.

2. Experimental

2.1. Flow tube and mass spectrometer

20 The uptake and reaction of the halogen trace gases on frozen and dry surfaces was investigated using a coated wall flow tube (Fig. 1) combined with a differentially pumped quadrupole mass spectrometer (Holmes et al., 2001). The flow tube set up consists of a horizontally mounted Pyrex glass tube (length 300 mm, internal diameter 27.5 mm) with a thermostatted jacket to regulate the temperature. The surfaces of interest are

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Uptake and reaction
of HOBr**

J. W. Adams et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

coated onto the inner wall of a snug-fitting Pyrex insert ($l = 210$ mm, $id = 25.5$ mm), located in the centre of the flow tube. The experiments were conducted at total (flow tube) pressures of between 0.75 and 3 Torr He (1 Torr = 1.333 hPa), the temperature was varied between 233 and 253 K. The bulk flow rate of He through the flow tube was between 400 and 1000, standard $\text{cm}^3 \text{min}^{-1}$, resulting in linear velocities of between 300 and 1500cm s^{-1} . Reactive trace gases, diluted in He, were introduced into the flow tube via a movable injector, the translation of which enabled the contact time between trace gas and surface to be varied. When appropriate, the main gas flow, which entered the flow-tube through a sidearm of the reactor, was humidified to prevent evaporation of the frozen surface. Trace gases were analysed with a quadrupole mass spectrometer (QMS), equipped with an electron-impact ion source (Fickert et al., 1998). HOBr, BrCl and Br_2 were monitored at their parent ions ($m/z = 96, 116$ and 158 , respectively).

2.2. Surface preparation, chemicals

In order to prepare reproducible frozen surfaces the following methodology was adapted. The Pyrex insert was cleaned using a dilute HF solution before thorough rinsing with distilled water and the aqueous solution from which the frozen or dry surfaces were to be made. Following this, a few ml of the same solution was introduced between two Teflon constraining rings. By slowly turning the insert, a film of estimated thickness $\approx 100 \mu\text{m}$ was evenly distributed on the glass walls. The insert was then placed into the pre-cooled flow tube where a translucent, frozen film was formed within a few seconds. The insert could also be slowly turned during the freezing process. Once the flow tube was re-sealed the bulk gas flows, at the correct humidity to maintain the ice film in equilibrium with gas-phase, were established. Note that the use of pure chloride / bromide samples leads to a completely crystalline condensed phase, with no remaining liquid at temperatures below ≈ 251 K (Koop et al., 2000).

When dry films were required, the frozen film was exposed to dry He as the temperature was allowed to increase slowly to room temperature over a period of ≈ 30 min. In this way an even, unbroken, dry salt surface could be reproducibly obtained. In the

**Uptake and reaction
of HOBr**

J. W. Adams et al.

text we refer to these surfaces as “dry” although the relative humidity was also varied in these experiments. HOBr was prepared by the reaction of AgNO_3 with Br_2 in solution as described previously (Fickert et al., 1999), and was stored as an aqueous solution at $\approx -30^\circ\text{C}$. BrCl was prepared by allowing a $\text{Br}_2 / \text{Cl}_2$ mixture to come to equilibrium in a blacked glass storage bulb, and its mixing ratio determined by absorption spectroscopy. The response of the mass spectrometer was calibrated against known concentrations of Br_2 , Cl_2 and BrCl, and was determined over a range of flow-tube pressures, gas-phase water concentrations and instrument configurations. The mass spectrometer signals of HOBr were converted to concentrations as described previously (Fickert et al., 1999).

The aqueous bromide / chloride solutions were prepared from analysis grade NaCl and NaBr. For most experiments the solutions were 2 M NaCl and 3×10^{-3} M NaBr. The Cl^- to Br^- ratio (≈ 660) is thus the same as found in sea-water, and the concentrations similar to those expected to be present in sea-spray (Sander and Crutzen, 1996). In addition, solutions of varied Cl^- to Br^- ratio, and also pure Cl^- and Br^- solutions were used.

2.3. Determination of uptake coefficients

In the present experiments, the pseudo first-order rate coefficient for loss to the reactive surface k_w was calculated from the variation of signal with injector position according to:

$$[C]_{z_2} = [C]_{z_1} \cdot \exp\left(-k_w \cdot \frac{\Delta z}{v}\right) \quad (\text{i})$$

where $[C]_{z_{1,2}}$ is the trace gas concentrations at injector positions z_1 and z_2 , Δz represents the distance between the positions z_1 and z_2 and v is the gas flow velocity in the tube. The ratio $\Delta z/v$ thus defines the gas-surface contact time. The measured value of k_w has to be corrected (to give k_{cor}) to take into account radial concentration gradients caused by efficient interaction of the trace gas at the wall (Brown, 1978). This required

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

knowledge of the diffusion coefficients of HOBr in both He and H₂O, which were determined separately (see results). The uptake coefficient, γ , was then obtained directly from k_{cor} (Howard, 1979):

$$\gamma = \frac{2 \cdot r \cdot k_{\text{cor}}}{\omega} \quad (\text{ii})$$

5 where r is the internal radius of the coated glass insert, and ω is the mean molecular velocity of the trace gas, derived from the Boltzmann equation. Note that using this method, we make the assumption that the geometric surface area of the ice substrate is appropriate for calculating the uptake coefficient, though there is still some controversy regarding this (Hanson and Ravishankara, 1993; Keyser et al., 1993). Corrections
10 based on pore diffusion calculations have only been properly tested for ice surfaces that were formed by condensation from the gas-phase, which are considerably rougher than those formed from the relatively slow freezing of a liquid as practised here. We expect that our ice surfaces, which appear smooth and translucent to the eye, are significantly less porous than those grown by vapour deposition.

15 3. Results and discussion

3.1. Diffusion coefficients of HOBr in He and H₂O

Experiments to determine the diffusion coefficient of HOBr in water vapour, $D_{\text{H}_2\text{O}}^{\text{HOBr}}$, were carried out using a salt surface doped with HCl at 255 K. For this measurement, the He gas flow was accurately humidified, and the uptake of HOBr, limited by its
20 diffusion through both He and H₂O, was monitored. The diffusion coefficient of HOBr in He, $D_{\text{He}}^{\text{HOBr}}$, was also determined using the reaction with HCl on the surface of the glass insert. In these experiments, HCl was added to the bulk gas flow, and was present at a concentration in excess of HOBr by more than a factor of 10.

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

For diffusion-limited uptake onto the surface of a reactor with cylindrical geometry, the equation approximating the dependence of the uptake coefficient on the bath gas pressure is given by:

$$\frac{1}{\gamma} = \frac{1}{\alpha} + \frac{\omega \cdot r \cdot P_{H_2O}}{7.32 \cdot D_{H_2O}} + \frac{\omega \cdot r}{7.32 \cdot D_{He}} \cdot P_{He} \quad (\text{iii})$$

5 This assumes that reaction at the surface is rapid and not rate limiting. Doping the ice / glass surface with excess HCl ensures that this is the case (Abbatt, 1994).

Figure 2 (open circles) shows plots of $1/\gamma$ vs. He bath gas pressure for uptake onto a sea-salt substrate in the presence of excess HCl at 255 K. Two experiments were performed at each pressure, though the excellent reproducibility makes some data points difficult to see. During these experiments the water vapour pressure was held constant at ca. 0.8 Torr. The linear dependence of the inverse uptake coefficient on the pressure of the He bath gas enabled the diffusion coefficient to be extracted from the gradient (see Eq. iii above). The diffusion coefficient at 255 K was found to be $D_{He}^{HOBr} = (273 \pm 1) \text{ Torr cm}^2 \text{ s}^{-1}$, where the errors are precision only (2 σ). This result can be compared to the only previous experimental determination of $D_{He}^{HOBr} = (319 \pm 48) \text{ Torr cm}^2 \text{ s}^{-1}$ at 274 K. Assuming a $T^{3/2}$ dependence on temperature, this indicates a value of $(286 \pm 43) \text{ Torr cm}^2 \text{ s}^{-1}$ at 255 K, in good agreement with the present result.

A separate set of experiments (filled circles) was performed in which the diffusion limited uptake of HOBr onto frozen sea-salt surfaces and excess HCl in the presence of various pressures of added water vapour was studied. During these experiments the He pressure was held constant at 2 Torr. Each data point represents the average of at least three uptake experiments. The slope of this data set yields a diffusion coefficient of $D_{H_2O}^{HOBr} = (51 \pm 1) \text{ Torr cm}^2 \text{ s}^{-1}$ where the errors are precision only (2 σ). This is the first direct determination of this quantity. The data taken with the partially humidified sea-salt surface (open circles) was analysed to obtain the y-axis intercept of the data. Once the contribution from the diffusion of HOBr in water (see Eq. iii) was subtracted,

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGS 2002

an estimate of the mass accommodation coefficient for uptake of HOBr onto frozen, HCl doped, sea-salt surfaces can be calculated. A value of $\alpha \approx 0.15$ was obtained, which is comparable to the literature value of $\gamma = 0.25$, obtained for the uptake of HOBr onto ice doped with HCl (Abbatt, 1994).

5 3.2. HOBr uptake onto frozen salt surfaces: product formation

The uptake and reaction of HOBr on frozen salt substrates was examined at a variety of temperatures between 233 and 253 K, at various NaBr / NaCl concentration ratios and with variation of the initial HOBr concentration and total pressure. For each experiment a fresh substrate was used.

10 Figure 3a displays raw data obtained at 233 K with an initial (aqueous) composition of the salt of 2 M NaCl and 3×10^{-3} M NaBr. After a steady flow of HOBr at a concentration of $\approx 4 \times 10^9 \text{ cm}^{-3}$ has been established ($t = 0$ to $t = 400$ s, constant mass spectrometer signal at $m/z = 96$) the injector is withdrawn and the HOBr is exposed to a well defined area of frozen substrate. At this point, the HOBr signal drops precipitously, indicating
15 uptake onto the substrate, and the molecular ion of Br_2 is, within the time resolution of the measurement, instantaneously observed. The uptake of HOBr and the release of Br_2 proceeds at approximately the same rate until $t = 3400$ s, during which the HOBr is seen to be converted stoichiometrically to Br_2 . When the injector is re-inserted at $t \approx 3400$ s, and contact between HOBr and the substrate is stopped the HOBr signal
20 returns to its initial value, and Br_2 production ceases. We note that although the result shown is representative of several data sets, the exact shape of the HOBr uptake profile was somewhat variable from one experiment to the next, though the size of the initial uptake coefficient (directly after the injector was withdrawn) was reasonably constant. Despite efforts made to generate surfaces in a reproducible manner, this effect probably
25 reflects differences in surface characteristics such as the number of defect sites and porosity, which will influence the availability of reactive sites on the surface. As shown, there is no evidence for significant production of BrCl in this experiment. In a second experiment, at the same temperature (Fig. 3b), a fresh substrate was exposed to a

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Uptake and reaction
of HOBr**

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGS 2002

significantly higher concentration of HOBr ($8.5 \times 10^{10} \text{ cm}^{-3}$). At this concentration of HOBr the time dependence of its uptake is markedly different. Following an initially large uptake directly after the injector has been withdrawn ($t = 550 \text{ s}$) the concentration of HOBr increases rapidly until it almost reaches its original level. At the same time Br_2 is initially formed at high yield, but decreases in concentration as the uptake of HOBr slows down. In this experiment BrCl is also clearly formed, though there is a delay of $\approx 50 \text{ s}$ before its maximum concentration is reached. An examination of the HOBr, Br_2 and BrCl profiles at $600 \text{ s} < t < 850 \text{ s}$ shows that HOBr is being converted to Br_2 and BrCl in roughly equal amounts.


The change in the HOBr uptake profile as the HOBr concentration is varied indicates that, at higher concentrations, there is sufficient HOBr to modify the reactivity of the surface on the time scales of these experiments, i.e. reactive sites on the surface are depleted as the reaction progresses, and both the uptake of HOBr and the formation of products decrease with time. An important aspect of these results is the switch in products from only Br_2 at low HOBr to Br_2 and BrCl as the concentration of HOBr is increased.

In a further series of experiments, the dependence of the HOBr uptake and the formation of Br_2 and BrCl products was examined as a function of the initial ratio of Br^- to Cl^- in the aqueous solution. NaCl was always kept at 2 M, whilst NaBr was varied between 3×10^{-3} and 3×10^{-5} M. The data from an experiment with $\text{NaBr} = 3 \times 10^{-4}$ M carried out at 233 K is displayed in Fig. 4. When compared to the results of Fig. 3a, the decrease of the NaBr concentration from 3×10^{-3} to 3×10^{-4} M is seen to have a profound impact on the time dependence of the HOBr uptake and release of products, and also on the identity of the products. At the lower NaBr concentration of $= 3 \times 10^{-4}$ M the uptake of HOBr is characterised by a rapid decrease in γ from its initial value over a period of $\approx 50 \text{ s}$, followed by a phase in which the uptake decreases slowly over several hundred seconds. At the same time, the Br_2 formation curve is also “spiked” and a significant amount of BrCl is formed, again with a delay period before it is detected. The ratio of BrCl to Br_2 formed at short exposure times is clearly significantly higher

**Uptake and reaction
of HOBr**

J. W. Adams et al.

than in experiments with 3×10^{-3} M Br^- . For comparison, the results from experiments in which HOBr was reacted with pure NaCl and NaBr surfaces are shown in Fig. 5. Figure 5a shows the uptake of HOBr onto frozen chloride (initial concentration 2 M). The uptake was strongly time dependent in all experiments, irrespective of the initial HOBr concentration. At low HOBr concentrations ($< 1 \times 10^{10}$ cm $^{-3}$), the only product observed was BrCl, the release of which was not instantaneous, but took > 100 s to reach a maximum value. The yield of BrCl (per HOBr taken up) was constant at 0.5 ± 0.1 after this maximum was passed. At higher initial concentrations of HOBr, Br $_2$ O was often formed on the chloride ice surface, via the reaction.



Br $_2$ O was monitored at $m/z = 176$. The Br $_2$ O formation was found to be dependent on the temperature of the flow tube, with enhanced formation at low temperatures. By measuring the relative HOBr and Br $_2$ O signals at various temperatures it was possible to set up simultaneous equations and determine the sensitivity of the mass spectrometer to Br $_2$ O. When high concentrations of HOBr were present, the product (BrCl and Br $_2$ O) to reactant ratio was ≈ 1 initially, falling to 0.4 ± 0.1 at longer times. Figure 5b shows the uptake of HOBr onto frozen bromide (initial concentration 3×10^{-3} M). In this case instantaneously released Br $_2$ was the only product, with a constant yield of 0.9 ± 0.1 per HOBr taken up.

The above observations of HOBr uptake, Br $_2$ and BrCl formation, and the dependence of these on the initial HOBr concentration or the ratio of NaBr to NaCl are used to derive a qualitative description of the reactions taking place on the frozen substrate. In a first step we expect HOBr to react with either Cl $^-$ or Br $^-$ on the surface to form either BrCl or Br $_2$:



Due to its greater polarity, we expect BrCl to have a relatively long residence time on

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

the surface compared to Br_2 , thus, when sufficient Br^- is available on the surface for reaction, BrCl will be converted to Br_2 before it can leave the surface to be detected as a gas-phase species.



5 When the Br^- concentration is lowered, the rate of desorption of BrCl from the surface becomes comparable with the rate of its surface reaction with Br^- and some will be detected, with a delay period due to retention on the surface. When Br^- is low, BrCl may thus build up on the surface and block reactive sites for HOBr uptake, hence the decrease in HOBr uptake between $500 \text{ s} < t < 1500 \text{ s}$ in Fig. 4. The same mechanism
10 also explains the formation of both BrCl and Br_2 when high initial HOBr concentrations are employed. In this case, the surface Br^- is rapidly depleted by the $\text{BrCl} + \text{Br}^-$ reaction (Fig. 3b, $550 < t < 600 \text{ s}$), and further uptake of HOBr (which is reduced in magnitude due to BrCl occupation of reactive sites) can result in BrCl formation and release. Independent evidence for an efficient conversion of BrCl to Br^- on frozen
15 $\text{NaCl} / \text{NaBr}$ substrates was gained in a separate set of experiments that directly examined the uptake of BrCl (see below).

Until now the formation of Br_2 following reaction of HOBr with mixed salt surfaces has been discussed largely in terms of an indirect mechanism in via formation of BrCl reaction. This is due to the greater concentration of Cl^- compared to Br^- in the aqueous solutions used to generate the frozen surfaces, which favours Reaction (5) over
20 Reaction (4). The rate of each reaction will however depend on its rate coefficient and the surface concentration of the halide ions. In the aqueous phase the reaction of HOBr with Br^- is a factor of ≈ 3 larger than that for HOBr with Cl^- which alone cannot compensate the factor 660 in the relative concentration of Cl^- and Br^- in the bulk solutions. However, during the freezing process, which presumably starts at the surface
25 of the cold glass insert and proceeds outwards towards the gas-liquid interface, some separation of Br^- and Cl^- will be caused by the higher solubility of Br^- compared to Cl^- . This will result in a frozen film that has a higher surface $\text{Br}^- / \text{Cl}^-$ ratio compared

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Uptake and reaction
of HOBr**

J. W. Adams et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

to bulk. Observations of enhanced $\text{Br}^- / \text{Cl}^-$ concentration ratios at the surface have been made on NaBr / NaCl crystals grown from aqueous solutions (Zangmeister et al., 2001), and from a melt (Ghosal et al., 2000). In the present experiments, although a surface enhancement of the Br^- ion concentration probably occurs, we have no means to determine its extent, but note that even a surface enhancement of a factor of 10 in the $\text{Br}^- / \text{Cl}^-$ ratio would increase the contribution of Reaction (5) to just $\approx 5\%$ if the relative aqueous phase rate coefficients for these reactions are applicable.

The observations presented here appear to be completely consistent with the known aqueous-phase equilibrium chemistry involved in the uptake and reaction of HOBr on deliquescent sea-salt aerosols or solutions (Wang et al., 1994; Fickert et al., 1999), and it is tempting to suggest that the reactions are taking place in a liquid layer on the ice surface. However, an important aspect of the aqueous phase transformation of HOBr into Br_2 or BrCl is the requirement of low pH (Abbatt and Waschewsky, 1998; Fickert et al., 1999) as the reactions are acid catalysed (Wang et al., 1994). We have shown (Fickert et al., 1999) that the release of molecular halogens from aqueous solutions is suppressed at a pH of greater than ≈ 7 , and have therefore carried out a series of experiments in which the NaBr / NaCl solutions were adjusted in pH between 4 and 10 prior to freezing. It was found that the variation of pH had no influence on identity of the products or their yields per HOBr taken up to the surface. As the amount of liquid-phase available at the surface of a frozen sample will depend on the temperature, we also carried out experiments at various temperatures between 233 and 253 K. Again, no significant change in the uptake coefficient, time dependence or product distribution (i.e. the Br_2 to BrCl ratio) was observed over this temperature range. Inspection of the NaCl phase diagram (Koop et al., 2000) shows that at temperatures below 251 K (the temperature of the ice/liquid/ $\text{NaCl}\cdot 2\text{H}_2\text{O}$ eutectic), only ice and NaCl di-hydrate co-exist. The presence of bromide ions at concentrations of 10^{-3} M is not expected to modify this and, for the majority of our experiments, there is not expected to be an aqueous phase in equilibrium with the bulk material of the substrate. Even in those experiments where the temperature was above the eutectic for NaCl (i.e. at 253 K)

there was no obvious change in either the kinetics of the uptake process or the product distribution.

The efficient formation of products Br₂ and/or BrCl in the absence of acidity can be understood in terms of the dissociation of HOBr to H⁺ and OBr⁻ in an aqueous medium:



At low H⁺ concentrations, HOBr (pK_a ≈ 9) will exist mainly in the form of its ions, and is not available for reaction with Cl⁻ or Br⁻. Acidification of the solution pushes the equilibrium to the left of Reaction (7) to favour the existence of molecular HOBr which makes nucleophilic attack by halide ions possible. On the frozen surfaces the degree of dissociative ionisation of HOBr will be reduced if insufficient H₂O molecules are available, and acidification will not be necessary.

3.3. HOBr uptake onto frozen salt surfaces: uptake coefficients

As shown in, e.g. Fig. 3b, the uptake of HOBr onto a frozen, mixed salt surface is sometimes time dependent, with an initially large uptake coefficient that decreases with exposure time due to the deactivation of reactive surface sites. At low concentrations of HOBr and sufficiently high halide ion concentrations the uptake was independent of exposure time on the time scales of a typical experiment (e.g. Fig. 3a). In order to make meaningful comparison of uptake coefficients from different data sets, the uptake coefficient shortly after exposure, γ_{init} , is reported. The initial uptake coefficients listed have been corrected for radial concentrations gradients as described in the experimental section. Only those data where the criterion $k_{\text{cor}} \leq 4k_{\text{obs}}$ is fulfilled are reported/displayed as absolute values, otherwise a lower limit to γ_{init} is presented. One set of experiments was carried out to examine the effect of temperature on the initial uptake coefficient of HOBr to a mixed salt surface, whereby uptake coefficients were obtained at temperatures of 233, 243, 248 and 253 K, with the 2 M Cl / 3×10^{-3} M Br⁻ surfaces. The results are presented in Table 1, and plotted in Fig. 6 (solid circles).

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Uptake and reaction
of HOBr**

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGS 2002

The data points are averages from several experiments. Although the data are associated with some scatter (the errors bars represent reproducibility) there is a trend of increasing uptake coefficient with decreasing temperature. Also plotted with this data are results obtained using the same initial Cl^- concentration, but without bromide present (open circles). The initial uptake coefficients are, within error limits, indistinguishable from those with 3×10^{-3} M bromide, and clearly, at these concentrations, the presence of bromide on the surface has little influence on the magnitude of the HOBr uptake coefficient. This substantiates the arguments presented above that the initial step is the reaction of surface adsorbed HOBr with the dominant halide ion, Cl^- , rather than reaction with Br^- . Uptake coefficients were also obtained for the reaction of HOBr with a surface prepared from freezing NaBr solutions of either 3×10^{-3} or 0.1 M concentration.

3.4. BrCl , Br_2 and Cl_2 uptake onto frozen salt surfaces

The uptake of HOBr and release of Br_2 is described above in terms of formation of BrCl in a primary step, followed by reaction of BrCl on the surface with Br^- . This concept was tested in separate experiments by measuring the uptake of BrCl , Br_2 and also Cl_2 onto frozen salt surfaces of varying composition. Representative data from an experiment to examine the uptake of BrCl (plus Br_2 and Cl_2 impurity) on a mixed $\text{Br}^- / \text{Cl}^-$ surface are shown in Fig. 7, which shows that BrCl is taken up efficiently and that the increase in gas-phase Br_2 is approximately equal to the summed loss of BrCl and Cl_2 from the gas-phase. If the initial Cl^- to Br^- ratio is close to 660 as found in sea-water, BrCl will react on the surface to form Br_2 . A lower limit to the uptake coefficient of 0.034 was obtained at 233 K.

In a further set of experiments the uptake of Br_2 onto mixed salt surfaces was examined. Br_2 was found to react strongly with these surfaces in halogen inter-conversion reactions. Although BrCl is expected to be the primary product, it reacts with the surface bromide to form Br_2 . The uptake coefficient was determined to be > 0.025 at 233 K. On pure bromide at 243 K, Br_2 is both taken up to react with the surface ($\gamma > 0.01$) and

released to form a different Br_2 molecule. In this case, the use of isotopically labelled NaBr was used to discriminate between the reactant and product Br_2 .

The uptake of Cl_2 was examined on mixed salt surfaces at 233 K, and resulted in the stoichiometric formation of Br_2 , presumably via the intermediacy of BrCl . A lower limit to the uptake coefficients of > 0.028 was obtained.

3.5. HOBr uptake onto dry salt surfaces

Experiments to examine the uptake of HOBr onto dry surfaces were performed on both mixed $\text{Br}^- / \text{Cl}^-$ surfaces and pure Cl^- surfaces, and at various relative humidities. For the mixed salt at 243 K (see Fig. 8a), γ was found to be 2×10^{-2} and time independent.

Uptake onto the pure Cl^- surface (see Fig. 8b) behaved differently and was clearly time dependent, with an initial uptake coefficient of 6×10^{-2} that decreased to a steady value of $\approx 3 \times 10^{-3}$ after ≈ 500 s. BrCl was formed at high yield ($\approx 70\%$), but not stoichiometrically. Upon re-inserting the injector ($t \approx 2050$ s) some HOBr was released from the surface to the gas-phase, indicating the presence of unreactive, physisorption sites on the chemically aged surface, and explaining the observation that the yield of BrCl per HOBr lost from the gas-phase was not 100%. Similar to the uptake of HOBr onto frozen surfaces, the uptake onto dried and humidified surfaces resulted in the release of both Br_2 and trace amounts of BrCl , with Br_2 the major initial product released to the gas-phase, followed by BrCl release as the surface bromide was depleted. There was no change in the product distribution as the relative humidity was varied between 0 and 100%.

3.6. Comparison with literature

3.6.1. HOBr uptake to frozen and dry surfaces

The present results pertaining to the uptake and reaction of HOBr on frozen salt surfaces can be directly compared to the similar but less extensive studies of Kirchner et

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Uptake and reaction
of HOBr**

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGS 2002

al. (1997). Also using a flow-tube / mass spectrometer set-up, those authors found that the reaction of HOBr on a frozen, mixed NaCl / NaBr surface at 233 K resulted in the release to the gas-phase of both BrCl and Br₂ in roughly equal amounts. They determined an uptake coefficient for the process of $\approx 1 - 3 \times 10^{-3}$. As detailed in Sect. 2, the present work has clearly shown that the use of HOBr concentrations in excess of 10^{11} cm^{-3} in studies of the HOBr uptake to salt surfaces can strongly influence both the kinetic behaviour and the ratio of the Br₂ and BrCl products, and can also lead to the formation of Br₂O on the surface. The factor of 10 difference in the uptake coefficient and the vastly different product distribution seen by these authors can therefore be explained by their use of $\approx 10^{13} \text{ cm}^{-3}$ HOBr, i.e. a factor of 1000 greater than routinely used in the present study. As we have already shown, the use of high concentrations of HOBr leads to rapid depletion of surface bromide, and increases the ratio of BrCl to Br₂ observed in the gas-phase. At the same time, at such high HOBr concentrations, we expect a very rapid surface saturation that leads to a much lower uptake coefficient after an initially high value. Even at concentrations of $\approx 10^{11} \text{ cm}^{-3}$ this effect is apparent, and at 10^{13} cm^{-3} the initial high uptake coefficient, would not be observable. Following the initial large uptake, the subsequent low uptake is also expected to show some time dependence, and results in curvature in plots of HOBr signal versus injector position. Indeed, a close examination of the raw data presented by Kirchner et al. (1997) reveals pronounced curvature in several of the uptake profiles. The low uptake coefficient measured by these authors probably corresponds to uptake onto a surface that is severely depleted in available reactive sites.

There have been several studies of the uptake of HOBr onto pure ice surfaces, which reveal a number of common features (Abbatt, 1994; Allanic et al., 1997; Chu and Chu, 1999; Chaix et al., 2000) including a strong negative temperature dependence of the uptake coefficient between 175 and 239 K, and the formation of Br₂ and BrCl products when the surface is doped with HBr or HCl respectively. The uptake coefficients measured in these studies vary from $\gamma \approx 0.5$ at 175 K to $\gamma \approx 10^{-3}$ at ≈ 230 K on pure ice. The uptake coefficients on pure ice at 230 K are thus lower than those measured on

**Uptake and reaction
of HOBr**

J. W. Adams et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

the frozen salt substrates of the present study, which were $\gamma \approx 10^{-2}$ at 230 K. On the other hand, the presence of HCl on the surface increases the uptake coefficient up to $\gamma \approx 0.2$ at temperature close to 230 K, depending on the amount of HCl available on the surface for reaction (Abbatt, 1994; Chu and Chu, 1999). The observed formation of BrCl and Br₂ when Cl⁻ and Br⁻ are available at the surface is compatible with the present results, as is the observation of a fast reaction of BrCl with surface adsorbed HBr on ice surfaces at 180 – 210 K to form Br₂ (Allanic et al., 2000).

The interaction of HOBr with dry NaCl and KBr surfaces has been examined at room temperature (Mochida et al., 1998). In that study no data were obtained for mixed salt surfaces. For reaction with NaCl, the products observed were BrCl and Br₂, whereby the Br₂ was believed to arise from a self-reaction of HOBr on the surface, and decomposition of the subsequently formed Br₂O. The BrCl evolution into the gas-phase displayed an induction period as observed in the present work. Br₂ was observed as the sole gas-phase product of the HOBr + KBr interaction. Mochida et al. (1998) reported initial uptake coefficients of $\gamma = (6.5 \pm 2.5) \times 10^{-3}$ for NaCl and $\gamma = (0.18 \pm 0.04)$ for KBr.

3.6.2. BrCl, Cl₂ and Br₂ uptake to frozen and dry surfaces

There have been a number of studies of the interactions of the di-halogens BrCl, Cl₂ and Br₂ with both dry and frozen salt surfaces. Berko et al. (1991) examined qualitative aspects of the uptake and reaction of Cl₂ and BrCl with dry NaBr crystals. They found that whilst BrCl reacted with NaBr to form Br₂ directly, the uptake of Cl₂ resulted in initial formation and release of Br₂, followed by BrCl. This is consistent with the present results and mechanism for reaction of Cl₂ with mixed salt surfaces. BrCl is formed in the first step of the reaction in which Cl₂ reacts with NaBr, but is converted to Br₂ before it can leave the bromide-rich surface. Only after the surface has been sufficiently depleted in Br⁻ can BrCl escape to the gas-phase. Kinetic data on the reaction of Cl₂ with dry synthetic sea-salt have also been obtained, and an uptake coefficient of

**Uptake and reaction
of HOBr**

J. W. Adams et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

$\gamma \approx 2 \times 10^{-2}$ reported (Mochida et al., 1998). As in the present study, Br_2 (rather than BrCl) was observed as the dominant product. The reaction of Cl_2 with frozen, mixed salt surfaces of similar composition to those reported here was examined by Huff and Abbatt (2000). They found an efficient interaction at 248 K, but no evidence for reaction at 233 K ($\gamma < 0.002$), which the authors reported as surprising. This observation contrasts the present result in which the uptake coefficient was found to be greater than 0.028 at 233 K, and where Br_2 product was observed at high yield. The result of Huff and Abbatt (2000) at 248 K, whereby an uptake coefficient of $\gamma = 0.048$ was obtained is more in line with the present observations, as is the observation that Br_2 is the dominant product (98–99%). These authors also carried out a detailed study of the reaction of Cl_2 with surfaces prepared from solutions containing various amounts of Br^- (in the absence of Cl^-) and found a strong dependence of the uptake coefficient on this parameter at 233 K.

The same authors have also examined the reaction of BrCl with frozen, Br^- containing salt surfaces and also find an efficient conversion of BrCl to Br_2 and an uptake coefficient that depended on the Br^- concentration, with values of $\gamma = 0.018$ for $[\text{Br}^-] = 0.01$ M, and $\gamma = 0.042$ for $[\text{Br}^-] = 0.1$ M. Importantly, for reaction of both Cl_2 and BrCl , they also observed no dependence of the uptake coefficient on the pH of the solution used to prepare the surface, which is consistent with the present observations for HOBr.

4. Conclusions and implications for the chemistry of the Polar Boundary Layer

In the present study it has been shown that HOBr reacts efficiently with surfaces prepared by freezing synthetic salt solutions of similar $\text{Br}^- / \text{Cl}^-$ ratio and ionic strength to sea water. Use of very low HOBr concentrations has enabled us to show that previous results (Kirchner et al., 1997) on both the uptake coefficient and product yields were erroneous due to use of very high HOBr concentrations in the gas-phase, that modified the relative surface abundance of Br^- and Cl^- on the time scale of the measurements.

**Uptake and reaction
of HOBr**

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGS 2002

Various aspects of our observations are discussed below in the context of their implications for helping us understand the cause of low ozone events (LOEs) in both the Arctic and Antarctic boundary layers. In doing this we note that whereas our laboratory-made surfaces, prepared from pure chloride / bromide solutions are not perfect replicas of sea-ice, or sea-spray that has been deposited on the ice pack, they allow the variation of parameters (i.e. the Cl^- to Br^- ratio and the pH) that enable us to gain some insight into the chemical mechanisms of halogen activation and release.

Firstly we note that the uptake of HOBr to a fresh, frozen sea-salt surface does not initially lead to the release of BrCl, which is consistent with our measurements in the aqueous phase (Fickert et al., 1999). A significant depletion of surface Br^- is necessary before the slow rate of desorption of BrCl from the surface allows it to compete with its transformation to Br_2 . If halogen activation is initiated by gas-phase bromine chemistry that leads to HOBr formation, we would predict first bromine activation and later the release of chlorine. At first glance, this appears to disagree with the results of Ramacher et al. (1999) who, by making measurements of hydrocarbon depletion rates, were able to show that during each ozone depletion event the time integrated chlorine atom concentration increases earlier than the bromine. However, as discussed by these authors, the presence of high concentrations of Br atoms (available to react with hydrocarbons) is precluded by ozone mixing ratios in excess of ≈ 10 ppb, whereas Cl atom / hydrocarbon reactions are relatively unaffected by the presence of these ozone concentrations.

We also observe that, in stark contrast to reaction in the aqueous phase, the uptake of HOBr to Br^- / Cl^- containing ice surfaces leads to release of molecular di-halogens at high yield even at low H^+ concentrations. In the marine boundary layer at mid-latitudes the deposition of HOBr to the surface of the sea does not lead to release of di-halogens, but requires the acidification of aerosol sea-spray by uptake of gases such as SO_2 , HNO_3 (Sander and Crutzen, 1996) or methane sulphonic acid (Ayers et al., 1999) (MSA), and gives halogen activation a strong dependence on the concentration of airborne particles, and thus wind strength. In very clean environments, and in the

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGS 2002

absence of a significant source of CH_3SCH_3 (precursor of MSA), di-halogen activation by HOBr uptake is therefore inoperative. Our observation that the uptake of HOBr and the formation of Br_2 or BrCl is efficient on dry and frozen salt surfaces made from non-acidified solutions may help explain why low ozone events still occur in the clean air of the Antarctic even though mechanisms to acidify particles may not be available. A complicating issue here is caused by the presence of a liquid-phase which is believed to be associated with these frozen surfaces, and in which high concentrations of Cl^- and Br^- are expected to be present (Koop et al., 2000). The reaction of HOBr with Cl^- and or Br^- in liquid environments should require a pH which is lower than that of sea-water to result in halogen activation as long as sufficient liquid is present to cause dissolution and ionisation of HOBr. Our measurements of the uptake coefficient of HOBr on frozen sea-salt surfaces of $\gamma = 2 \times 10^{-2}$ enables comparison between, and linking of the heterogeneous and homogeneous gas-phase processes occurring in the boundary layer at the poles during a LOE to be made. If most of the heterogeneous chemistry involving salt surfaces is taking place at the snow pack rather than in suspended sea-salt aerosols (Tang and McConnell, 1996; Sander et al., 1997), values of γ of this magnitude imply that the rate of reaction of HOBr at the surface of the snow pack will be limited by its deposition velocity, which, following Huff and Abbatt (2000), and assuming a mixed layer of 100 m height, is $\approx 0.6 \text{ cm s}^{-1}$. Although there may be a weak source of Br atoms due to the photolysis of bromoform (Barrie et al., 1988), which may be sufficient to initiate bromine chemistry at polar sunrise, most photochemical activity will take place close to the snow pack, where Br_2 is released and photolysed (to form two Br atoms, Reaction 8) on time scales that may be short relative to times required for vertical mixing. The Br atoms can react with O_3 to form BrO .



**Uptake and reaction
of HOBr**

J. W. Adams et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

At normal levels of O_3 (≈ 40 ppb, Hausmann and Platt, 1994; Hopper et al., 1998), most Br atoms will react with O_3 , and the loss of one HOBr in a heterogeneous process (11) and the formation of 2 HOBr via Reaction sequence (9 and 10) implies that the photochemical / heterogeneous chemistry taking place at the snow pack is of a chain branching nature, and will result in efficient release of reactive halogen species as long as sufficient Br^- is present in the surface snow-pack and sufficient O_3 is present in the gas-phase so that formation of BrO is efficient compared to reaction with aldehydes. If the reservoir of Br^- is significantly depleted before all gas-phase O_3 has been consumed, then chlorine activation can be expected, as BrCl is released in preference to Br_2 . Fast surface reactions and the short lifetimes of the reactive bromine species with respect to photo-dissociation imply that mixing of O_3 rich air masses distant from the snow pack with O_3 poor air masses very close to the surface will be important in determining the occurrence and duration of LOEs. These considerations preclude a simple description of the heterogeneous and gas-phase reactions that initiate and maintain high concentrations of reactive halogens in the polar boundary layer, which can only be done with the use of models that contain both gas-phase and heterogeneous halogen chemistry (the objectives of this paper) and appropriate parameterisation of mixing and transport of air masses over the snow pack. The reader is referred to the recent modelling study by Michalowski et al. (2000) which shows that O_3 depletion is very sensitive to the concentration of HOBr, and the rate of its uptake to Br^- and Cl^- containing surfaces.

Acknowledgements. We extend our thanks to Thomas Koop for several helpful discussions regarding the freezing of halide solutions, and for making his data available prior to publication. We thank Rolf Sander for critically reading the manuscript. We acknowledge the European Union for financial support (ENV4-CT97-0394, HAMLET).

References

- Abbatt, J. P. D.: Heterogeneous reaction of HOBr with HBr and HCl on ice surfaces at 228 K, *Geophys. Res. Lett.*, 21, 665–668, 1994.
- Abbatt, J. P. D. and Waschewsky, G. C. G.: Heterogeneous interactions of HOBr, HNO₃, O₃ and NO₂ with deliquescent NaCl aerosols at room temperature, *J. Phys. Chem.*, 102, 3719–3725, 1998.
- Allanic, A., Oppliger, R., and Rossi, M. J.: Real-time kinetics of the uptake of HOBr and BrONO₂ on ice and in the presence of HCl in the temperature range 190–200 K, *J. Geophys. Res.*, 102, 23 529–23 541, 1997.
- Allanic, A., Oppliger, R., van den Bergh, H., and Rossi, M. J.: The heterogeneous kinetics of the reactions ClONO₂ + HX/ice (X = Br, I), BrONO₂ + HI/ice and the reactivity of the interhalogens BrCl, ICl and IBr with HX/ice in the temperature range 180 to 210 K, *Z. Phys. Chem.*, 214, 1479–1500, 2000.
- Anlauf, K. G., Mickle, R. E., and Trivett, N. B. A.: Measurement of ozone during polar sunrise experiment 1992, *J. Geophys. Res.*, 99, 24 345–24 353, 1994.
- Ayers, G. P., Gillett, R. W., Cainey, J. M., and Dick, A. L.: Chloride and bromide loss from sea-salt particles in Southern Ocean air, *J. Atmos. Chem.*, 33, 299–319, 1999.
- Barrie, L. and Platt, U.: Arctic tropospheric chemistry: An overview, *Tellus*, 49B, 450–454, 1997.
- Barrie, L. A., Bottenheim, J. W., and Hart, W. R.: Polar sunrise experiment 1992 (PSE 1992): Preface, *J. Geophys. Res.*, 99, 25 313–25 314, 1994.
- Barrie, L. A., Bottenheim, J. W., Schnell, R. C., Crutzen, P. J., and Rasmussen, R. A.: Ozone destruction and photochemical reactions at polar sunrise in the lower Arctic atmosphere, *Nature*, 334, 138–141, 1988.
- Berko, H. N., McCaslin, P. C., and Finlayson-Pitts, B. J.: Formation of gas-phase bromine compounds by reaction of solid NaBr with gaseous ClONO₂, Cl₂, and BrCl at 298 K, *J. Phys. Chem.*, 95, 6951–6958, 1991.
- Boudries, H. and Bottenheim, J. W.: Cl and Br atom concentrations during a surface boundary layer ozone depletion event in the Canadian high Arctic, *Geophys. Res. Lett.*, 27, 517–520, 2000.
- Brown, R. L.: Tubular flow reactors with first-order kinetics, *J. Res. Nat. Bur. Standards*, 83, 1–8, 1978.

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Uptake and reaction
of HOBr**

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGS 2002

Chaix, L., Allanic, A., and Rossi, M. J.: Heterogeneous chemistry of HOBr on different types of ice and on ice doped with HCl, HBr and HNO₃ at 175 <math>T/K < 215</math>, *J. Phys. Chem. A*, 104, 7268–7277, 2000.

Chance, K.: Analysis of BrO measurements from the global ozone monitoring experiment, *Geophys. Res. Lett.*, 25, 3335–3338, 1998.

Chu, L. and Chu, L. T.: Heterogeneous interaction and reaction of HOBr on ice films, *J. Phys. Chem. A*, 103, 8640–8649, 1999.

Fan, S.-M. and Jacob, D. J.: Surface ozone depletion in Arctic spring sustained by bromine reactions on aerosols, *Nature*, 359, 522–524, 1992.

Fickert, S., Adams, J. W., and Crowley, J. N.: Activation of Br₂ and BrCl via uptake of HOBr onto aqueous salt solutions, *J. Geophys. Res.*, 104, 23 719–23 727, 1999.

Fickert, S., Helleis, F., Adams, J. W., Moortgat, G. K., and Crowley, J. N.: Reactive uptake of ClNO₂ on aqueous bromide solutions, *J. Phys. Chem.*, 102, 10 689–10 696, 1998.

Foster, K. L., Plastridge, R. A., Bottenheim, J. W., Shepson, P. B., Finlayson-Pitts, B. J., and Spicer, C. W.: The role of Br₂ and BrCl in surface ozone destruction at polar sunrise, *Science*, 291, 471–474, 2001.

Ghosal, S., Shbeeb, A., and Hemminger, J. C.: Surface segregation of bromine in bromide doped NaCl: Implications for the seasonal variations in Arctic ozone, *Geophys. Res. Lett.*, 27, 1879–1882, 2000.

Hanson, D. R. and Ravishankara, A. R.: Response to “Comment on porosities of ice films used to simulate stratospheric cloud surfaces”, *J. Phys. Chem.*, 97, 2802–2803, 1993.

Hausmann, M. and Platt, U.: Spectroscopic measurements of bromine oxide and ozone in the high Arctic during polar sunrise experiment 1992, *J. Geophys. Res.*, 99, 25 399–25 414, 1994.

Holmes, N. S., Adams, J. W., and Crowley, J. N.: Uptake and reaction of HOI and IONO₂ on frozen and dry NaCl / NaBr surfaces and H₂SO₄, *Phys. Chem. Chem. Phys.*, 3, 1679–1687, 2001.

Hopper, J. F., Barrie, L. A., Silis, A., Hart, W., Gallant, A. J., and Dryfhout, H.: Ozone and meteorology during the 1994 polar sunrise experiment, *J. Geophys. Res.*, 103, 1481–1492, 1998.

Howard, C. J.: Kinetic measurements using flow tubes, *J. Phys. Chem.*, 83, 3–9, 1979.

Huff, A. K. and Abbatt, J. P. D.: Gas-Phase Br₂ production in heterogeneous reactions of Cl₂, HOCl and BrCl with halide-ice surfaces, *J. Phys. Chem.*, 104, 7284–7293, 2000.

**Uptake and reaction
of HOBr**

J. W. Adams et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

Impey, G. A., Mihele, C. M., Anlauf, K. G., Barrie, L. A., Hastie, D. R., and Shepson, P. B.: Measurements of photolyzable halogen compounds and bromine radicals during the polar sunrise experiments 1997, *J. Atmos. Chem.*, 34, 21–37, 1999.

Impey, G. A., Shepson, P. B., Hastie, D. R., Barrie, L.A., and Anlauf, K. G.: Measurement of photolyzable chlorine and bromine during the polar sunrise experiment 1995, *J. Geophys. Res.*, 102, 16 005–16 010, 1997.

Ingham, T., Bauer, D., Landgraf, J., and Crowley, J. N.: Ultraviolet-visible absorption cross sections of gaseous HOBr, *J. Phys. Chem.*, 102, 3293–3298, 1998.

Jobson, B. T., Niki, H., Yokouchi, Y., Bottenheim, Y., Hopper, F., and Leaitch, R.: Measurements of C2-C6 hydrocarbons during polar sunrise experiment 1992, *J. Geophys. Res.*, 99, 25 355–25 368, 1994.

Keyser, L. F., Leu, M.-T., and Moore, S. B.: Comment on porosities of ice films used to simulate stratospheric cloud surfaces, *J. Phys. Chem.*, 97, 2800–2801, 1993.

Kirchner, U., Benter, T., and Schindler, R. N.: Experimental verification of gas phase bromine enrichment in reactions of HOBr with sea salt doped ice surfaces, *Ber. Bunsen Ges. Phys. Chem.*, 101, 975–977, 1997.

Koop, T., Kapilashrami, A., Molina, L. T., and Molina, M. J.: Phase transitions of sea-salt/water mixtures at low temperatures: Implications for ozone chemistry in the polar marine boundary layer, *J. Geophys. Res.*, 105, 26 393–26 402, 2000.

Kreher, K., Johnson, P. V., Wood, S. W., Nardi, B., and Platt, U.: Ground-based measurements of tropospheric and stratospheric BrO at Arrival Heights, Antarctica, *Geophys. Res. Lett.*, 24, 3021–3024, 1997.

Lorenzen-Schmidt, H., Wessel, S., Unold, W., Solberg, S., Gernandt, H., Stordal, F., and Platt, U.: Ozone measurements in the European Arctic during the ARCTOC 1995 campaign, *Tellus*, 50B, 416–429, 1998.

McConnell, J. C., Henderson, G. S., Barrie, L., Bottenheim, J., Niki, H., Langford, C. H., and Tempelton, E. M. J.: Photochemical bromine production in Arctic boundary-layer ozone depletion, *Nature*, 355, 150–152, 1992.

Michalowski, B. A., Francisco, J. S., Li, S. M., Barrie, L. A., Bottenheim, J. W., and Shepson, P. B.: A computer model study of multiphase chemistry in the Arctic boundary layer during polar sunrise, *J. Geophys. Res.*, 105, 15 131–15 145, 2000.

Miller, H. L., Weaver, A., Sanders, R. W., and Arpag, K.: Measurements of arctic sunrise surface ozone depletion events at Kangerlussuaq, Greenland (67° N, 51° W), *Tellus*, 49B, 496–509,

1997.

Mochida, M., Akimoto, H., van den Bergh, H., and Rossi, M. J.: The heterogeneous kinetics of the uptake of HOBr on solid alkali halides at ambient temperatures, *J. Phys. Chem.*, 102, 4819–4828, 1998.

5 Mochida, M., Hirokawa, J., Kajii, Y., and Akimoto, H.: Heterogeneous reactions of Cl₂ with sea salts at ambient temperature: Implications for halogen exchange in the atmosphere, *Geophys. Res. Lett.*, 25, 3927–3930, 1998.

Mozurkewich, M.: Mechanisms for the release of halogens from sea-salt particles by free radical reactions, *J. Geophys. Res.*, 100, 14 199–14 207, 1995.

10 Oltmans, S. J.: Surface Ozone Measurements in Clean-Air, *J. Geophys. Res.*, 86, 1174–1180, 1981.

Oltmans, S. J., Schnell, R. C., Sheridan, P. J., Peterson, R. E., Li, S.-M., Winchester, J. W., Tans, P. P., Sturges, W. T., and Kahl, J. D.: Seasonal surface ozone and filterable bromine relationship in the high Arctic, *Atmos. Environ.*, 23, 2431–2441, 1989.

15 Perner, D., Arnold, T., Crowley, J., Klüpfel, T., Martinez, M., and Seuwen, R.: The measurement of active chlorine in the atmosphere by chemical amplification, *J. Atmos. Chem.*, 34, 9–20, 1999.

Ramacher, B., Rudolph, J., and Koppmann, R.: Hydrocarbon measurements in the spring arctic troposphere during the ARCTOC 95 campaign, *Tellus*, 49B, 466–485, 1997.

20 Ramacher, B., Rudolph, J., and Koppmann, R.: Hydrocarbon measurements during tropospheric ozone depletion events: Evidence for halogen atom chemistry, *J. Geophys. Res.*, 104, 3633–3653, 1999.

Röckmann, T., Brenninkmeijer, C. A. M., Crutzen, P. J., and Platt, U.: Short-term variations in the C¹³/C¹² ratio of CO as a measure of Cl activation during tropospheric ozone depletion events in the Arctic, *J. Geophys. Res.*, 104, 1691–1697, 1999.

25 Rudolph, J., Fu, B. R., Thompson, A., Anlauf, K., and Bottenheim, J.: Halogen atom concentrations in the Arctic troposphere derived from hydrocarbon measurements: Impact on the budget of formaldehyde, *Geophys. Res. Lett.*, 26, 2941–2944, 1999.

Sander, R. and Crutzen, P. J.: Model study indicating halogen activation and ozone destruction in polluted air masses transported to the sea, *J. Geophys. Res.*, 101, 9121–9138, 1996.

30 Sander, R., Vogt, R., Harris, G. W., and Crutzen, P. J.: Modeling the chemistry of ozone, halogen compounds, and hydrocarbons in the arctic troposphere during spring, *Tellus*, 49B, 522–532, 1997.

**Uptake and reaction
of HOBr**

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

**Uptake and reaction
of HOBr**

J. W. Adams et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

- Solberg, S., Schmidbauer, N., Semb, A., and Stordal, F.: Boundary-layer ozone depletion as seen in the Norwegian Arctic in spring, *J. Atmos. Chem.*, 23, 301–332, 1996.
- Tang, T. and McConnell, J. C.: Autocatalytic release of bromine from Arctic snow pack during polar sunrise, *Geophys. Res. Lett.*, 23, 2633–2636, 1996.
- 5 Tuckermann, M., Ackermann, R., Gölz, C., Lorenzen-Schmidt, H., Senne, T., Stutz, J., Trost, B., Unold, W., and Platt, U.: DOAS observations of halogen radical-catalyzed arctic boundary layer ozone destruction during the ARCTOC-campaigns 1995 and 1996 in Ny-Alesund, Spitsbergen, *Tellus*, 49B, 533–555, 1997.
- 10 Wang, T. X., Kelley, M. D., Cooper, J. N., Beckwith, C., and Margerum, D. W.: Equilibrium, kinetic and UV-spectral characteristics of aqueous bromine chloride, bromine and chlorine species, *Inorg. Chem.*, 33, 5872–5878, 1994.
- Wessel, S., Aoki, S., Winkler, P., Weller, R., Herber, A., Gernandt, H., and Schrems, O.: Tropospheric ozone depletion in polar regions. A comparison of observations in the Arctic and Antarctic, *Tellus*, 50B, 34–50, 1998.
- 15 Zangmeister, C. D., Turner, J. A., and Pemberton, J. E.: Segregation of NaBr in NaBr / NaCl crystals grown from aqueous solutions: Implications for sea salt surface chemistry, *Geophys. Res. Lett.*, 28, 995–998, 2001.

Uptake and reaction of HOBr

J. W. Adams et al.

Table 1. Summary of kinetic and product data obtained for HOBr uptake onto frozen salt surfaces

[Cl ⁻] / M	[Br ⁻] / M	<i>T</i> K	Major Products*	γ_{Init}	<i>n</i>
2	3×10^{-3}	248	Br ₂	0.026 ± 0.006	10
2	3×10^{-3}	233	Br ₂	0.032 ± 0.009	16
2	3×10^{-3}	253	Br ₂	0.012 ± 0.005	2
2	3×10^{-3}	243	Br ₂	0.017 ± 0.09	6
2	0	233	BrCl	0.043 ± 0.008	3
2	0	248	BrCl	0.016 ± 0.007	2
0	0.1	248	Br ₂	> 0.01	1
0	3×10^{-3}	248	Br ₂	0.036	1

*Product formed at short exposure times. The evolution of the product distribution as a function of reaction time is described in the text.

n = number of experiments.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

Table 2. Summary of kinetic and product data obtained for uptake of BrCl, Cl₂ and Br₂ onto frozen salt surfaces, and HOBr onto dry (and humidified)¹ salt surfaces

Trace gas	Surface ²	T K	Major Products ³	Y _{Init}
BrCl	Br ⁻ / Cl ⁻ (Frozen)	233	Br ₂	> 0.034
Br ₂	Br ⁻ / Cl ⁻ (Frozen)	233	Br ₂	> 0.025
Br ₂	Br ⁻ (Frozen)	243	Br ₂	> 0.01
Cl ₂	Br ⁻ / Cl ⁻ (Frozen)	233	Br ₂	> 0.028
HOBr	Br ⁻ / Cl ⁻ (Dry)	243	Br ₂	0.02
HOBr	Cl ⁻ (Dry)	233	BrCl	0.06

¹Relative humidity varied from 0 to 100%. ²Br⁻/Cl⁻ surfaces were made from 2 M Cl⁻ / 3 × 10⁻³ M Br⁻ solutions. The Cl⁻ surfaces were prepared from 2 M Cl⁻ solutions, the Br⁻ surfaces were prepared from 3 × 10⁻³ M solutions. ³Product formed at short exposure times. The evolution of the product distribution as a function of reaction time is described in the text.

Uptake and reaction of HOBr

J. W. Adams et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Print Version

Interactive Discussion

© EGS 2002

**Uptake and reaction
of HOBr**

J. W. Adams et al.




Fig. 1. Schematic Diagram of the experimental set-up. QMS = Quadrupole Mass Spectrometer.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

Uptake and reaction of HOBr

J. W. Adams et al.




Fig. 2. Plot of inverse uptake coefficient of HOBr vs. pressure of He or H_2O , for dry salt and HCl doped Pyrex surfaces at 253 K. Diffusion coefficients and accommodation coefficients may be extracted from the gradients and intercepts of these lines. (See Eq. iii in text)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

Uptake and reaction of HOBr

J. W. Adams et al.




Fig. 3. Uptake of HOBr and subsequent product formation, on a frozen mixed salt surface at 233 K. **(a)** $[\text{HOBr}]_{\text{init}} = 4 \times 10^9 \text{ molecules cm}^{-3}$, exposure of the surface to HOBr begins at 400 s when the injector is retracted. **(b)** $[\text{HOBr}]_{\text{init}} = 8.5 \times 10^{10} \text{ molecules cm}^{-3}$, exposure of the surface to HOBr begins at 550 s. In both cases, the Br_2 trace has been offset for clarity of presentation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

Uptake and reaction of HOBr

J. W. Adams et al.




Fig. 4. Uptake of HOBr onto a frozen (233 K) mixed salt surfaces made from an aqueous solution of composition 2 M NaCl, 3×10^{-4} NaBr. The Br₂ trace has been offset for clarity of presentation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

Uptake and reaction of HOBr

J. W. Adams et al.




Fig. 5. Uptake of HOBr onto frozen salt surfaces. **(a)** 2 M NaCl, $T = 233 \text{ K}$, **(b)** $3 \times 10^{-3} \text{ M}$ NaBr, 243 K. The Br_2 trace has been offset for clarity of presentation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

Uptake and reaction of HOBr

J. W. Adams et al.




Fig. 6. Temperature dependence of the initial uptake coefficient of HOBr onto frozen surfaces made from solutions containing 2 M NaCl with and without added NaBr. The error bars define the reproducibility of the result.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

**Uptake and reaction
of HOBr**

J. W. Adams et al.




Fig. 7. Uptake of BrCl and Cl₂ onto a mixed, frozen salt surface and formation of Br₂. $T = 233$ K, 2 M NaCl, 3×10^{-3} M NaBr. The Br₂ and Cl₂ traces have been offset for clarity of presentation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002

Uptake and reaction of HOBr

J. W. Adams et al.




Fig. 8. Uptake of HOBr onto dried salt surface at 233 K **(a)** 2 M NaCl, 3×10^{-3} M NaBr, **(b)** 2 M NaCl, 0 M NaBr.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Print Version](#)[Interactive Discussion](#)

© EGS 2002