



**HAL**  
open science

## Remote sensed and in situ constraints on processes affecting tropical tropospheric ozone

B. Sauvage, R. V. Martin, A. van Donkelaar, X. Liu, K. Chance, L. Jaeglé, P. I. Palmer, S. Wu, T.-M. Fu

► **To cite this version:**

B. Sauvage, R. V. Martin, A. van Donkelaar, X. Liu, K. Chance, et al.. Remote sensed and in situ constraints on processes affecting tropical tropospheric ozone. *Atmospheric Chemistry and Physics Discussions*, 2006, 6 (6), pp.11465-11520. hal-00302279

**HAL Id: hal-00302279**

**<https://hal.science/hal-00302279>**

Submitted on 18 Jun 2008

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Tropical tropospheric  
ozone**

B. Sauvage et al.

# Remote sensed and in situ constraints on processes affecting tropical tropospheric ozone

**B. Sauvage<sup>1</sup>, R. V. Martin<sup>1,2</sup>, A. van Donkelaar<sup>1</sup>, X. Liu<sup>2</sup>, K. Chance<sup>2</sup>, L. Jaeglé<sup>3</sup>, P. I. Palmer<sup>4</sup>, S. Wu<sup>5</sup>, and T.-M. Fu<sup>5</sup>**

<sup>1</sup>Department of Physics and Atmospheric Science, Dalhousie University, Halifax, Nova Scotia, Canada

<sup>2</sup>Atomic and Molecular Physics Division, Harvard-Smithsonian Center for Astrophysics, Cambridge, Massachusetts, USA

<sup>3</sup>Department of Atmospheric Sciences, University of Washington, Seattle, Washington, USA

<sup>4</sup>School of GeoSciences, University of Edinburgh, UK

<sup>5</sup>Department of Earth and Planetary Sciences and Division of Engineering and Applied Sciences, Harvard University, Cambridge, Massachusetts, USA

Received: 5 October 2006 – Accepted: 8 November 2006 – Published: 17 November 2006

Correspondence to: B. Sauvage (bsauvage@fizz.phys.dal.ca)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

## Abstract

We use a global chemical transport model (GEOS-Chem) to evaluate the consistency of satellite measurements of lightning flashes and ozone precursors with in situ measurements of tropical tropospheric ozone. The measurements are tropospheric O<sub>3</sub>, NO<sub>2</sub>, and HCHO columns from the GOME satellite instrument, lightning flashes from the OTD and LIS instruments, profiles of O<sub>3</sub>, CO, and relative humidity from the MOZAIC aircraft program, and profiles of O<sub>3</sub> from the SHADOZ ozonesonde network. We interpret these multiple data sources with our model to better understand what controls tropical tropospheric ozone. Tropical tropospheric ozone is mainly affected by lightning and convection in the upper troposphere and by surface emissions in the lower troposphere. Scaling the spatial distribution of lightning in the model to the observed flash counts improves the simulation of O<sub>3</sub> in the upper troposphere by 5–20 ppbv versus in situ observations and by 1–4 Dobson Units versus GOME retrievals of tropospheric O<sub>3</sub> columns. A lightning source strength of 5±2 Tg N/yr best represents in situ observations from aircraft and ozonesonde. Tropospheric NO<sub>2</sub> and HCHO columns from GOME are applied to provide top-down constraints on emission inventories of NO<sub>x</sub> (biomass burning and soils) and VOCs (biomass burning). The top-down biomass burning inventory is larger by a factor of 2 for HCHO and alkenes, and by 2.6 for NO<sub>x</sub> over northern equatorial Africa. These emissions increase lower tropospheric O<sub>3</sub> by 5–20 ppbv, improving the simulation versus aircraft observations, and by 4 Dobson Units versus GOME observations of tropospheric O<sub>3</sub> columns. Emission factors in the a posteriori inventory are more consistent with a recent compilation from in situ measurements. The ozone simulation using two different dynamical schemes (GEOS-3 and GEOS-4) is evaluated versus observations; GEOS-4 better represents O<sub>3</sub> observations by 5–15 ppbv due to enhanced convective detrainment in the upper troposphere. Heterogeneous uptake of HNO<sub>3</sub> on aerosols reduces simulated O<sub>3</sub> by 5–7 ppbv, reducing a model bias versus in situ observations over and downwind of deserts. Exclusion of HO<sub>2</sub> uptake on aerosols improves O<sub>3</sub> by 5 ppbv in biomass burning regions.

## Tropical tropospheric ozone

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 1 Introduction

Ozone ( $O_3$ ) in the tropical troposphere is a major component of atmospheric radiative forcing (de Forster et al., 1997; Lacis et al., 1990) and plays a key role in the global oxidizing power of the atmosphere (Logan et al., 1981). Indeed tropical regions present high ultraviolet radiation and humidity rates that promote hydroxyl (OH) creation through  $O_3$  photolysis (Thompson et al., 1992). Tropical tropospheric  $O_3$  production is limited by nitrogen oxides ( $NO_x = NO + NO_2$ ) emitted from biomass burning (Chatfield and Delany, 1990), biogenic sources, lightning, and fossil fuel combustion (Jacob et al., 1996). The motivation of the present manuscript is to better understand processes affecting tropical tropospheric  $O_3$ , using a global chemical and transport model constrained with satellite and in situ data.

Considerable uncertainty remains in the magnitude and distribution of tropical  $O_3$  precursor emissions, such as  $NO_x$  (Lee et al., 1997; Holland et al., 1999). Lightning produced  $NO_x$  (L- $NO_x$ ) are the most uncertain with recent estimates varying by an order magnitude from 1 to 13 Tg N/yr (Nesbitt et al., 2000; Price et al., 1997). Lightning  $NO_x$  emissions are largest over the Tropics, in the Inter Tropical Convergence Zone (ITCZ) area (Christian et al., 2003), and are directly emitted into the free troposphere where long lifetimes and efficient  $O_3$  production make the  $O_3$  burden very sensitive to those emissions (Martin et al., 2002a). Surface sources from biomass burning and soils are also highly uncertain (around 3–13 Tg N/yr and 4–21 Tg N/yr respectively, Holland et al., 1999). Biomass burning accounts for half of the global CO emissions (Andreae et al., 1993) and most recently soils have been highlighted to be an underestimated  $NO_x$  source (Jaeglé et al., 2004). Bottom-up estimates of these tropical emissions have been confounded by the lack of measurements in this remote region.

The goal of the present study is motivated by 2 objectives: 1/ use a global chemical transport model to evaluate the consistency of satellite measurements of lightning flash counts and  $O_3$  precursors with in situ measurements of tropospheric  $O_3$ , and 2/ interpret these multiple data sources with a global chemical transport model to bet-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ter understand what controls tropical tropospheric O<sub>3</sub>. Indeed evaluation of satellite data is limited over the tropics because of lack of in situ measurements especially for O<sub>3</sub> precursors. Understanding of tropical tropospheric O<sub>3</sub> is also limited by current uncertainties on anthropogenic and natural O<sub>3</sub> precursors sources, which can be readily inferred from satellite observations. Global measurements of nitrogen dioxide (NO<sub>2</sub>) atmospheric concentrations from space provide a top-down constraint on NO<sub>x</sub> emissions (Martin et al., 2003a; Jaeglé et al., 2005; Leue et al., 2001). Tropospheric NO<sub>2</sub> columns track surface NO<sub>x</sub> emissions on a regional scale since NO<sub>2</sub> is the dominant form of NO<sub>x</sub> in the boundary layer and the NO<sub>x</sub> lifetime against oxidation in the tropical boundary layer is several hours. Similarly, volatile organic compounds (VOC) emissions, critical for understanding radical chemistry in the troposphere, can be constrained by formaldehyde (HCHO) columns measured from space (Palmer et al., 2003). Indeed HCHO is a high-yield product of VOC oxidation with a lifetime of hours (Palmer et al., 2003). Interpretation of these two tropospheric column molecules is then fundamental for evaluation of a correct location and intensity of ground sources of O<sub>3</sub> precursors. In situ measurements from the Measurements of ozone and water vapor by in-service Airbus aircraft (MOZAIC) program (Marengo et al., 1998; Thouret et al., 2006) and the Southern Hemisphere Additional Ozonesondes SHADOZ network (Thompson et al., 2003a, b) provide vertical profile information that is unavailable from satellite. Few studies have used at the same time the different dataset available over the Tropics, through in situ measurements and satellite observations, to better understand tropical tropospheric O<sub>3</sub>. A global chemical transport model is a useful tool to relate measurements from these disparate sources.

We provide an overview of the data sets in Sect. 2. A complete description of the GEOS-Chem global chemical transport model is in Sect. 3.1. Then we introduce the standard simulation used in this study, based on improvements described in the same Sect. 3.2. These improvements enable a better understanding of factors controlling tropospheric tropical O<sub>3</sub>. In Sect. 4, we first evaluate the simulation and integration of satellite information with in situ data and satellite data; then we assess the dynamical

**Tropical tropospheric ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and chemical processes driving tropical tropospheric O<sub>3</sub>.

## 2 Presentation and overview of the data

The following measurements are used to improve and evaluate the GEOS-Chem chemical transport model.

### 5 2.1 In situ data. Aircraft and ozonesonde measurements

Since 1994, the MOZAIC airborne program provides regular measurements of ozone (the overall precision is  $\pm 2$  ppbv+2%) and water vapor at high spatial and temporal resolution (Marenco et al., 1998). Recent details are available at <http://mozaic.aero.obs-mip.fr>). Additional CO measurements are performed onboard the five instrumented aircraft (Nédélec et al., 2003) since the end of 2000 with an overall precision of  $\pm 5$  ppbv,  $\pm 5\%$ . Table 1 contains characteristics of the MOZAIC sites, with their locations shown in Fig. 1 in blue font. We use 19 of the 30 cities sampled by the MOZAIC program between 30° N–30° S, the most sampled ones, with 15 to 60 flights per month for a site. This corresponds to a total of 6750 flights over all regions.

15 We analyze the data in monthly average for the 1994–2005 period, except for West Africa where measurements began in 2001 (Savauge et al., 2005). For each site, we remove data within 15 km of a site, to avoid local pollution that is not representative of the broader region. This criterion removes the lowest 25–50 hPa.

20 The SHADOZ network complements the MOZAIC coverage as shown in Fig. 1 in black. It provides regular ozonesonde measurements (Thompson et al., 2003a,b), at different tropical stations, at least twice a month. Further details can be found on the SHADOZ Web site: <http://croc.gsfc.nasa.gov/shadoz/>. We use measurements over the 1998–2004 period.

25 For clarity and conciseness, we present a subset representative of the broader region indicated by the black rectangle in Fig. 1. We also examined other sites within

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

each region, but found similar features.

## 2.2 Space-based observations. The LIS, OTD and GOME instruments

The Optical Transient Detector (OTD) (Boccippio et al, 2000b) was launched in 1995 on the MicroLab-1 satellite. The OTD spatial resolution is 10 km over a field of view of 1300 km×1300 km. The OTD detects both intra-cloud (IC) and cloud-to-ground (CG) discharges during day and night conditions with a 40–65% detection efficiency. The Lightning Imaging sensor (LIS) was launched in 1997 aboard the Tropical Rainfall Measuring Mission (TRMM) Observatory into a nearly circular orbit inclined 35 degrees with an altitude of 350 km. It detects lightning with storm-scale resolution of 3–6 km (3 at nadir, 6 at limb) over 550×550 km. The system is enabled to detect weak lightning and achieve a 90% detection efficiency (Christian et al., 1989).

The Global Ozone Monitoring Experiment (GOME) (Burrows et al., 1999) instrument onboard the European Remote Sensing-2 satellite provided the capability for continuous global monitoring of O<sub>3</sub>, NO<sub>2</sub> and HCHO atmospheric columns through observation of solar backscatter over 1995–2003. GOME observes the atmosphere in the nadir view with a 40 km along track by 320 km across track. Global coverage is achieved every 3 days with an overpass time over the tropics between 10–11 local time (crossing the equator at 1030 local time). In this work we use GOME measurements for the year 2000.

We begin with tropospheric NO<sub>2</sub> line-of-sight (slant) columns retrieved from the GOME observations by Martin et al. (2002b) version 2 (Guerova et al., 2006), and HCHO slant columns retrieved by (Chance et al., 2000). Following Palmer et al. (2001) we calculate vertical columns by applying an air mass factor (AMF) algorithm to account for atmospheric scattering. The AMF is computed as the integral of the relative vertical distribution of the trace gas (shape factor), weighted by the altitude dependent scattering weights computed from the LIDORT radiative transfer model (Spurr et al., 2002). Coincident NO<sub>2</sub> and HCHO shape factors are from the standard GEOS-Chem simulation described in Sect. 3. The cloud correction uses local cloud information from

## Tropical tropospheric ozone

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Tropical tropospheric ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

GOME (Kurosu et al., 1999) as described in Martin et al. (2002b). The aerosol correction uses aerosol profiles from the GEOS-Chem model following Martin et al. (2003a). We exclude observations in which the fraction of backscattered intensity from clouds exceeds 50% of a GOME scene. The stratospheric NO<sub>2</sub> column is removed using observations over the central Pacific where there is little tropospheric NO<sub>2</sub>, and subtracting the corresponding column from the ensemble of GOME scenes for the appropriate latitude and month. The result is corrected for the small amount of tropospheric NO<sub>2</sub> over the Pacific. Variability in the stratospheric NO<sub>2</sub> columns is accounted for using assimilated stratospheric NO<sub>2</sub> columns from Boersma et al. (2004), a minor issue in the Tropics.

Martin et al. (2004) evaluated the GOME retrieval with airborne in situ measurements of NO<sub>2</sub> and HCHO over the Southeastern United States. Uncertainties include absolute errors of  $1 \times 10^{15}$  molecules cm<sup>-2</sup> for tropospheric NO<sub>2</sub> (Martin et al., 2002b) and  $4 \times 10^{15}$  molecules cm<sup>-2</sup> for HCHO (Chance et al., 2000) from the spectral fitting, the stratospheric NO<sub>2</sub> column and instrument artifacts. Other uncertainties arising from the AMF calculation include random and systematic contributions from surface reflectivity, clouds, aerosols, and the trace gas profile (Martin et al., 2003a; Boersma et al., 2004). The monthly mean uncertainty is  $\pm (5 \times 10^{14}$  molecules cm<sup>-2</sup> + 30%) for tropospheric NO<sub>2</sub> and a 30% (Millet et al., 2006) error on the HCHO column retrieval that increases in the presence of biomass burning aerosol (Fu et al., 2006<sup>1</sup>). van Noije et al. (2006) compared three difference retrievals of tropospheric NO<sub>2</sub> columns from GOME, and found the greatest degree of consistency in the tropics, well within the error estimates reported here.

For O<sub>3</sub>, we use version 2 of tropospheric O<sub>3</sub> columns retrieved by Liu et al. (2005). The retrieval uses an optimal estimation method (Rodgers, 2000). Tropospheric ozone

<sup>1</sup>Fu, T.-M., Jacob, D. J., Palmer, P. I., Chance, K., Wang, Y. X., Barletta, B., Blake, D. R., Stanton, J. C., and Pilling, M. J.: Space-based formaldehyde measurements as constraints on volatile organic compound emissions in east and south Asia, *J. Geophys. Res.*, submitted, 2006.



columns (TOC), the sum of tropospheric partial columns, are interpolated with the GEOS-Chem model tropopause used to divide the stratosphere and the troposphere. GOME retrievals and GEOS-Chem simulations are mapped onto a common regular grid.

### 3 General description of the GEOS-Chem model – original and standard versions

A global 3-D model of tropospheric chemistry provides a quantitative tool to assess the processes affecting tropospheric ozone. We use the GEOS-Chem chemical and transport model (Bey et al., 2001). In the following we first introduce the original model version (7-02-04 <http://www-as.harvard.edu/chemistry/trop/geos/index.html>). Then we describe the “standard” simulation, focusing on developments to improve the original simulation.

#### 3.1 Original version

The model is driven by assimilated meteorological data for 2000 from the Goddard Earth Observing System (GEOS-4) at the NASA Global Modeling and Assimilation Office (GMAO). The model version has 30 vertical sigma-levels (surface to 0.1 hPa), and a horizontal resolution of 1° latitude by 1.25° longitude, which can be degraded to 2° latitude by 2.5° longitude and 4° latitude by 5° longitude for computational expediency. We use the latter two resolutions in the study. The data have 6-hour temporal resolution (3-hour for surface variables and mixing depth). We present sensitivity simulations using GEOS-3 as discussed in Sect. 4.4.

The GEOS-Chem model includes a detailed simulation of tropospheric O<sub>3</sub>-NO<sub>x</sub>-hydrocarbon chemistry as well as of aerosols and their precursors, using 41 tracers, around 90 species, and 300 reactions. The model presently includes sulfate, nitrate, ammonium, black and organic carbon, mineral dust and sea salt (Park et al., 2004,

## Tropical tropospheric ozone

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Tropical tropospheric  
ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2005; Alexander et al., 2005; Fairlie et al., 2006<sup>2</sup>). The aerosol and gaseous simulations are coupled through formation of sulfate and nitrate,  $\text{HNO}_3(\text{g})/\text{NO}_3^-$  partitioning of total inorganic nitrate, heterogeneous chemistry on aerosols (Jacob, 2000; Evans et al., 2005), and aerosol effects on photolysis rates (Martin et al., 2003b). The model has been previously applied to interpret satellite observations of HCHO (Palmer et al., 2001, 2003, 2006; Shim et al., 2005; Millet et al., 2006),  $\text{NO}_2$  (Martin et al., 2002b, 2003a; Jaeglé et al., 2004, 2005; Guerova et al., 2006), and tropospheric  $\text{O}_3$  (Martin et al., 2002a; Chandra et al., 2002, 2003; Kim et al., 2005; Liu et al., 2006). However, none of these studies has examined all three species together.

Table 2 contains annual global  $\text{NO}_x$  emissions used in the model. Soil  $\text{NO}_x$  emissions are computed using a modified version of the algorithm of Yienger and Levy (1995) with the canopy reduction factors described in Wang et al. (1998). The biomass burning inventory is interannually varying and is based on satellite observations of fires as derived by Duncan et al. (2003). Emissions of lightning  $\text{NO}_x$  are linked to deep convection following the parameterization of Price et al. (1992) with vertical profiles from Pickering et al. (1998) as implemented by Wang et al. (1998).

### 3.2 Standard (improved) version

In the following section we present several developments which are necessary for accurate understanding and evaluation of the processes affecting tropical tropospheric  $\text{O}_3$  described in Sect. 4. These improvements deal with emissions and heterogeneous chemistry that are included in our standard simulation. GOME observations of  $\text{NO}_2$  and HCHO are applied to constrain surface emissions of  $\text{NO}_x$  and VOCs. Lightning flash counts are used to better represent its spatial distribution. Heterogeneous chemistry on aerosols is updated to reflect recent measurements.

<sup>2</sup>Fairlie, T. D., Jacob, D. J., and Park, R. J.: The impact of transpacific transport of mineral dust in the United States, *Atmos. Environ.*, submitted, 2006.

### 3.2.1 Soil NO<sub>x</sub> emissions

Strong signals from soil NO<sub>x</sub> emissions are apparent in satellite observations of tropospheric NO<sub>2</sub> columns (Bertram et al., 2005). We use the a posteriori NO<sub>x</sub> emission inventory derived from GOME observations of NO<sub>2</sub> columns by Jaeglé et al. (2005) for the year 2000. GOME tropospheric NO<sub>2</sub> column observations were related to surface NO<sub>x</sub> emissions via inverse modeling with GEOS-Chem model. They used the spatio-temporal distribution of remotely sensed fires and a priori inventory information on the locations of regions dominated by fuel combustion to partition among the different NO<sub>x</sub> sources. The resulting annual tropical emissions are 35% higher in the a posteriori inventory (Table 2) and account for 22% of tropical NO<sub>x</sub> emissions. During March-April-May (MAM) and June-July-August (JJA), emissions increase by a factor of 3 over tropical ecosystems of Africa, reflecting a better constraint on NO<sub>x</sub> emissions associated with the monsoon (Jaeglé et al., 2004). Emissions increase by 20% during the rainy season over South America, and the agricultural region of North India.

### 3.2.2 Biomass burning emissions of NO<sub>x</sub> and VOCs

We apply tropospheric NO<sub>2</sub> and HCHO columns retrieved from GOME to provide top-down constraints on regional biomass burning emissions of NO<sub>x</sub> and reactive VOCs. Richter et al. (2002) found a strong signal from biomass burning in the GOME NO<sub>2</sub> columns. Our inversion for biomass burning NO<sub>x</sub> is conducted after application of the a posteriori soil NO<sub>x</sub> inventory from Jaeglé et al. (2005). The NO<sub>x</sub> inversion accounts for the local NO<sub>2</sub>/NO ratio and the local NO<sub>x</sub> lifetime following Martin et al. (2003a). The inversion is applied here at regional scale in the form of NO<sub>x</sub> emission factors that should be applicable to simulations for other years.

Palmer et al. (2003) showed that HCHO columns over North America are closely related to isoprene emissions, and exploited that relationship to infer continental isoprene emissions from the GOME HCHO columns. Meyer-Arnek et al. (2005) found signals in the GOME HCHO columns from both biogenic and pyrogenic sources over

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Africa. Shim et al. (2005) extended the approach of Palmer et al. (2003) to infer global isoprene emissions, but found that large increases in biomass burning emissions were necessary to reconcile the GOME observations. More recently, Fu et al. (2006)<sup>1</sup> found over East and South East Asia a biomass burning source derived from GOME almost 5 times the estimate of a bottom-up emission inventory. We similarly find over the tropics an underestimate of more than a factor two in the GEOS-Chem HCHO columns during biomass burning. Neither scaling of the current GEIA isoprene emission inventory, nor application of the recently developed MEGAN inventory (Guenther et al., 2006), was able to account for the discrepancy without introducing biases outside of the biomass burning season. A recent compilation by Andreae (2005, personal communication) of in situ measurements of emission factors contain values that are higher than those used in GEOS-Chem for HCHO and alkenes. We tentatively attribute the regional difference between GOME and GEOS-Chem HCHO columns to biomass burning emissions of alkenes and HCHO, and calculate a tropical mean emission ratio for reactive VOCs emissions that is a factor of 2 larger for both species.

Figure 2 shows the seasonal NO<sub>x</sub> biomass burning emissions arising from the a priori (left) and top-down (right) inventories. Annual tropical NO<sub>x</sub> emissions are 30% higher in the top-down versus the a priori (Table 2). NO<sub>x</sub> emissions from Africa and eastern regions increase by 30%, whereas they decrease from South America by 30%. The largest absolute difference occurs in DJF over Northern Africa with top-down emissions of 0.96 Tg N /season compared to 0.41 Tg N /season, likely reflecting emission factors that were too low in the original simulation. There is also a 15% increase in emissions from Central/South Africa during JJA to 1.06 Tg N.

### 3.2.3 Lightning NO<sub>x</sub> emissions

We use space-based observations of lightning flash counts from the seasonally varying climatological OTD/LIS (Boccippio et al., 2000a, 2001) dataset (High Resolution Annual Climatology – HRAC – data) to constrain GEOS-Chem lightning flashes, by applying a

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

local seasonal rescaling factor,  $R$ :

$$R = \left( \frac{\text{Local}_{(\text{LIS}/\text{OTD})} \text{flashes}}{\text{Global}_{(\text{LIS}/\text{OTD})} \text{flashes}} \right)_{\text{season}} \bigg/ \left( \frac{\text{Local}_{(\text{GEOS}-\text{Chem})} \text{flashes}}{\text{Global}_{(\text{GEOS}-\text{Chem})} \text{flashes}} \right)_{\text{season}} \quad (1)$$

This approach is motivated by the seasonal latitudinal variation in tropical lightning activity that is not well represented by the GEOS fields. The scaling factor is applied to a 10-year average of the simulated and observed flashes, such that inter-annual variability of the lightning emissions is allowed. The climatology is a  $0.5^\circ \times 0.5^\circ$  gridded composite of total intra cloud – cloud to ground (IC+CG) lightning bulk production over 1995–2004. Lowpass temporal filtering of 110 days for the combined LIS/OTD is applied. Observations in the LIS/OTD v1.0 reanalysis have been corrected by the LIS Science Team for flash detection efficiency, applied as a function of sensor, viewing time, date of mission, and (for OTD) geographic location. For the entire dataset, these corrections correspond to average flash detection efficiencies of 47% (OTD) and 82% (LIS) (Boccippio et al., 2002; Christian et al., 2003). The adjustments derive from a combination of laboratory calibration, ground validation, and cross-normalization between OTD and LIS. The uncertainty in these corrections is  $\pm 10\%$ .

Figure 3 shows the seasonal average lightning  $\text{NO}_x$  emissions ( $\text{L-NO}_x$ ) during DJF and JJA, for the original (left), and standard (right) simulations. The LIS/OTD seasonal climatologies and the improved  $\text{L-NO}_x$  emissions in GEOS-Chem exhibit higher spatio-temporal correlations ( $r^2=0.97\text{--}0.98$ ) than in the original simulation ( $r^2=0.4\text{--}0.57$ ). Annual emissions are unchanged (Table 2). However substantial regional differences are inferred by the local rescaling. Emissions decrease over Africa by 16%, over South America by 42%, and increase from the Eastern tropics by 55% (mostly over Australia). During JJA, continental  $\text{L-NO}_x$  emissions decrease south of the ITCZ by 50% whereas they increase by 45% over North Africa. During DJF continental emissions

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

decrease in general by around 50%. Oceanic emissions increase by a factor of 2.9.

### 3.2.4 Heterogeneous chemistry

The original simulation used a reaction probability  $\gamma$  of  $\text{HO}_2$  on all aerosols equal to 0.2. Laboratory measurements by [Thornton et al. \(2005\)](#) demonstrated that  $\text{HO}_2$  uptake on aerosols is negligible at temperatures warmer than 270 K in the absence of Cu or Fe ions that would catalyze the reaction. Field measurements of biomass burning aerosol ([Yamasoe et al., 2000](#)) found insufficient Cu or Fe ions to catalyze that reaction. We exclude this reaction for biomass burning aerosols.

Following [Bauer et al. \(2004\)](#) we implement  $\text{HNO}_3$  uptake on mineral aerosols in the standard simulation using  $\gamma(\text{HNO}_3)=0.1$ . Laboratory experiments have shown  $\text{HNO}_3$  uptake on mineral dust is promoted by its alkalinity (Goodman et al., 2000; Grassian, 2000; Underwood et al., 2001; Michel et al., 2002; Hanisch and Crowley, 2003). Field measurements also support  $\text{HNO}_3$  uptake (Tabazadeh et al., 1998; Thakur et al., 1999). Rapid sedimentation of nitrate on mineral dust could reduce recycling of  $\text{NO}_x$  from  $\text{HNO}_3$ , and in turn  $\text{O}_3$ , with lower tropospheric  $\text{O}_3$  decreases of 8–30% over and downwind of deserts (Bian et al., 2003; Bauer et al., 2004; Umann et al., 2005; Liao et al., 2005).

There have been few comparisons with in situ measurements to evaluate these heterogeneous processes. In Sect. 4.3 we perform sensitivity studies to evaluate the uptake of  $\text{HO}_2$  on biomass burning aerosols and uptake of  $\text{HNO}_3$  on mineral dust.

## 4 Assessment of the dynamical and chemical processes affecting tropospheric tropical ozone

Of particular interest is 1/ the ability of the model to accurately simulate the distribution of tropospheric ozone and its precursors in order to 2/ accurately understand what controls tropical tropospheric ozone. We first give an overview of the distribution of

### Tropical tropospheric ozone

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

tropospheric ozone columns. We then discuss the processes affecting its distribution in the context of the MOZAIC and SHADOZ vertical O<sub>3</sub> profiles, as well as the GOME tropospheric NO<sub>2</sub> and HCHO columns.

Figure 4 shows seasonal TOC from GOME observations (left), our standard simulation (middle). We exclude retrievals with cloud fraction exceeding 0.7 of a GOME scene. The simulated and retrieved O<sub>3</sub> columns exhibit similar spatio-temporal variation over the Tropics (monthly  $r^2=0.91-0.98$ ; seasonal bias = 1.4–4.4 DU). Both show enhancements in the downwelling branches of the Hadley circulation, smaller values in the Tropics, and a zonal wave-one pattern, with maximum TOC between 40W-60E. The original and retrieved TOC are less consistent (monthly  $r^2=0.67-0.87$ ) although the tropical mean bias remains unchanged.

The right panels show large regional changes of 5 DU in the simulated O<sub>3</sub>. In the following sections we focus on the consequences of our developments on the comparison of the model versus observations.

## 4.1 Sensitivity to lightning

Here we discuss how the local rescaling of lightning flashes affects the comparison with O<sub>3</sub> observations. Then we discuss the sensitivity of the simulation to lightning intensity and to lightning vertical distribution.

### 4.1.1 Satellite constraint. Lightning rescaling

The local rescaling of lightning flashes to match OTD/LIS measurements yields substantial improvement in the modeled TOC as demonstrated below. We compare the original and standard simulations at MOZAIC and SHADOZ sites that exhibit the largest sensitivity to lightning. These sites are generally in subsidence regions downwind of lightning activity, allowing for O<sub>3</sub> production during transport. Figure 5 shows the seasonal O<sub>3</sub> vertical profiles for the in situ measurements (MOZAIC, SHADOZ, black lines); original (blue line) simulation, and the standard simulation (red line). Both simulations

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

are generally within one standard deviation of the in situ measurements. However improvements due to the lightning rescaling are apparent in the standard simulation in the middle and upper troposphere.

The first panel of Fig. 5 show continental sites with  $O_3$  concentrations of 40–50 ppbv throughout the year in the middle and upper troposphere, sustained by L- $NO_x$  emissions in the South American Convergence Zone (SACZ) or the ITCZ. The lightning rescaling reduces L- $NO_x$  emissions in South America (Fig. 3) decreasing in upper tropospheric  $O_3$  during DJF and MAM by 5–10 ppbv over Sao Paulo and by 10–15 ppbv over Caracas. The Middle East is under the influence of an anticyclonic circulation in the middle and upper troposphere (Hoskins and Rodwell, 1995) and of easterly flow through the Tropical Easterly jet in the upper troposphere, which brings lightning outflow during the Indian monsoon (Li et al., 2001), mainly during JJA as depicted by the easterly ozone flux (Fig. 4). Reductions in Indian L- $NO_x$  emissions improve the simulation at Dubai by 5–10 ppbv in JJA and SON. Bangkok is influenced by lightning mostly during the dry season from November to May when the circulation is convergent. Lightning rescaling improves the  $O_3$  simulation by 5–15 ppbv. Other continental sites exhibit less sensitivity due to their proximity to L- $NO_x$  emissions.

The effect of local lightning rescaling is also apparent in the TOC. Table 3 contains the TOC for the standard simulation, the in situ measurements, and the GOME retrievals. Lightning rescaling has a considerable effect on  $O_3$  over South America (Fig. 4, right panel) reducing the model bias versus the in situ measurements to within 2 DU over Caracas and within 4 DU over Sao Paulo, compared to more than 8 DU difference in the original simulation. The simulation is closer to in situ TOC than to GOME observations over both regions. Over the Middle East lightning rescaling improves the simulated TOC by 3–5 DU to within 2–5 DU. The remaining bias at Dubai arises from the  $O_3$  overestimate below 600 hPa (Fig. 5). GOME measurements are within 2 DU of the MOZAIC TOC except during DJF when there is a 5 DU underestimate that probably originates from the lower troposphere as noticed by Liu et al. (2006). Over South East Asia there is a positive bias of GEOS-Chem TOC compared to GOME, between 4 to

**Tropical tropospheric ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



7 DU. The lightning rescaling clearly yields better modeled TOC versus MOZAIC to within 1–4 DU, versus 7 DU for the original simulation.

The second panel of Fig. 5 show that lightning rescaling also yields improvements over oceanic sites. Lightning rescaling increases emissions over the South Pacific Convergence Zone (Fig. 3) especially in DJF and SON resulting in a 5–10 ppbv increase in  $O_3$  in the middle and upper troposphere (Samoa, Fig. 5). Over Reunion Island there is improvement in DJF due to a 7 ppbv increase in  $O_3$ . The Atlantic, Ascension and Natal depict similar  $O_3$  vertical profiles near the maximum of the zonal-wave one, with enhanced mid-upper tropospheric  $O_3$  throughout the year. Lightning is a significant source of this enhancement (Thompson et al., 2000; Martin et al., 2002a; Sauvage et al., 2006b<sup>3</sup>). The main improvements are in DJF and MAM with  $O_3$  increases of 7–10 ppbv from more lightning over Central Africa in the standard simulation (Fig. 3). During SON, both simulations in the middle troposphere and upper troposphere underestimate  $O_3$  by 10 to 20 ppbv, but  $O_3$  remains enhanced.

The TOC over oceans are generally consistent between the standard simulation, GOME and in situ measurements, within 5 DU everywhere. Over Ascension the standard simulation is closer than the original one to in situ measurements by 1–3 DU in DJF and MAM. However there is still an underestimate of 4 DU in SON. Better agreement is found over the Pacific and Indian Ocean, within 1–3 DU compared to in situ measurements, and within 2–5 DU versus GOME TOC.

Finally the seasonal cycle of the modeled TOC is reproduced for all sites, except over Caracas. The last line of Table 3 shows that for the TOC averaged for the tropical sites, the three datasets are within 2 DU bias and within  $1\sigma$  of the measurements. The seasonal cycle is well reproduced, with maximum in SON, minimum in MAM, as depicted by the southern hemispheric zonal-wave one pattern (Thompson et al., 2003b; Sauvage et al., 2006a).

<sup>3</sup>Sauvage, B., Martin, R. V., van Donkelaar, A., and Ziemke, J. R.: Quantification of the factors controlling tropical tropospheric ozone and the South Atlantic maximum, *J. Geophys. Res.*, under review, 2006b.

**Tropical tropospheric ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 4.1.2 Lightning intensity and distribution

### a. Sensitivity to intensity

In order to evaluate the lightning  $\text{NO}_x$  source of  $5 \text{ Tg N/yr}$ , we conduct sensitivity studies based on the standard simulation that vary the intensity over  $3$  to  $7 \text{ Tg N yr}^{-1}$ .

Figure 5 shows the sensitivity of the seasonal  $\text{O}_3$  vertical profiles to L- $\text{NO}_x$  intensity, using either  $3$  or  $7 \text{ Tg N/yr}$  (dashed green lines). Lower concentrations reflect the simulation with  $3 \text{ Tg N/yr}$ . In general  $\text{O}_3$  is perturbed throughout the entire troposphere by  $5$ – $10$  ppbv. The simulation remains nearly within one standard deviation of measurements. However  $3 \text{ Tg N/yr}$  is generally too low. In contrast  $7 \text{ Tg N/yr}$  is generally too high. The largest sensitivity to intensity is found over the Atlantic region where  $\text{O}_3$  concentrations change by  $10$ – $20$  ppbv. The simulation with  $7 \text{ Tg N/yr}$  reduces the model bias versus in situ measurements in SON at Ascension (Fig. 5), but creates a bias during other seasons, and at most other sites. Emissions of  $9.7 \text{ Tg N/yr}$  would be necessary to achieve in situ  $\text{O}_3$  concentrations in SON at Ascension. Another process is likely responsible for the bias.

In summary,  $5 \pm 2 \text{ Tg N yr}^{-1}$  represents the plausible range of lightning  $\text{NO}_x$  emissions. Outside of that range, simulated  $\text{O}_3$  becomes increasingly inconsistent with in situ measurements. This is obviously dependent of the accuracy of all surface sources. Martin et al. (2006)<sup>4</sup> found a similar magnitude of  $6 \pm 2 \text{ Tg N yr}^{-1}$  best agreed with space-based measurements of  $\text{NO}_2$ ,  $\text{O}_3$  and  $\text{HNO}_3$ .

### b. Sensitivity to distribution

The vertical distribution of lightning emissions is also important (Labrador et al.,

<sup>4</sup>Martin, R. V., Sauvage, B., Folkins, I., Sioris, C. E., Boone, C., Bernath, P., and Ziemke, J.: Space-based constraints on the production of nitric oxide by lightning, *J. Geophys. Res.*, in revision, 2006.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2005). Most previous studies assumed much higher NO<sub>x</sub> emissions per flash for cloud to ground (CG) flashes than intra-cloud (IC) flashes (Price et al., 1997; Pickering et al., 1998). However, recent studies provide evidence that the IC/CG ratio may be between 0.5–1.0 (DeCaria et al., 2000; Fehr et al., 2004). The implications have not yet been evaluated versus tropical in situ O<sub>3</sub> data. We explore the implications of increasing the IC/CG ratio to 0.75, instead of 0.1 in our standard simulation. The additional NO<sub>x</sub> from intra-cloud flashes is distributed within the cloud anvil.

The simulation using enhanced IC emissions is shown in Fig. 5 (solid green line). Generally, this lightning parameterization overestimates middle-upper tropospheric O<sub>3</sub>, but remains within one standard deviation of measurements. The effects vary with season and location, with for example negligible incidence at Caracas, a negative bias at Bangkok, and a large impact at Ascension. Over Ascension, O<sub>3</sub> concentrations are biased high in DJF and MAM by 10–15 ppbv, but the model bias in SON is eliminated suggesting a seasonal variation in the IC/CG ratio. In summary, a uniform increase in the IC/CG ratio is unsupported by the in situ O<sub>3</sub> profiles, but it could be higher for particular geographical regions.

## 4.2 Sensitivity to biomass burning and soils

In this section we address the following questions: What are the consequences of the modifications to surface emissions of NO<sub>x</sub> and VOCs on O<sub>3</sub> distributions? Do these changes improve the simulated tropospheric O<sub>3</sub> compared to in situ measurements?

### 4.2.1 Satellite constraint

Figure 6 shows seasonal average GOME (left) and GEOS-Chem (middle: standard; right: original) tropospheric columns of NO<sub>2</sub>, during 2000. The GOME and GEOS-Chem NO<sub>2</sub> standard columns are highly consistent over the Tropics during the 4 seasons. The coefficient of determination of the retrieved columns versus the standard simulation during the 4 seasons ( $r^2=0.86\text{--}0.91$ ,  $p<0.0001$ ) is considerably higher than

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

versus the original simulation, which are in the range of  $r^2=0.6-0.8$ . The annual mean absolute difference between the standard simulation and retrieved columns over the Tropics is  $0.2 \times 10^{15}$  molecules  $\text{cm}^{-2}$  compared with  $0.7 \times 10^{15}$  molecules  $\text{cm}^{-2}$  in the original simulation. The standard simulation better reproduces seasonal  $\text{NO}_2$  maxima observed by GOME. For instance over Northern Africa during DJF and MAM, top down biomass burning  $\text{NO}_x$  emissions enhance lower tropospheric  $\text{NO}_2$  concentrations by a factor of 2.6, reducing a regional model bias. Over Central Africa, the regional bias in JJA and SON is reduced, however a local bias remains during JJA reflecting the regional emission factor applied here. Over India  $\text{NO}_2$  tropospheric column are reduced by a factor 4 during the biomass burning season of MAM, better representing GOME columns. During May to July the a posteriori soil  $\text{NO}_x$  emission inventory better reproduces the  $\text{NO}_2$  column enhancement over the Sahel.

Figure 7 shows seasonal average GOME (left) and GEOS-Chem (middle standard; right original) tropospheric columns of HCHO during 2000. The spatio-temporal correlation is quite high with  $r^2=0.7-0.9$  compared with  $0.6-0.75$  respectively versus the original simulation. The mean absolute difference between GOME and the standard simulation is  $0.06 \times 10^{16}$  molecules  $\text{cm}^{-2}$ , versus  $0.2 \times 10^{16}$  molecules  $\text{cm}^{-2}$  with original simulation. Previous regional differences of more than a factor of 2 are reduced during the biomass burning season to 20% in the standard simulation over Northern Africa in DJF-MAM and to 35% over Central Africa and South America in JJA-SON. The remaining model biases likely reflect isoprene emissions.

#### 4.2.2 Evaluation with in situ data

Figure 8 shows  $\text{O}_3$  profiles at MOZAIC sites that have the greatest sensitivity to surface emissions. West equatorial (Lagos, Abidjan) and Central Africa (Brazzaville) sites exhibit  $\text{O}_3$  enhancements related to seasonal biomass burning fires (Fig. 8) driven by the lower tropospheric Harmattan and trade flow. The new CO measurements confirm the sensitivity of those sites to biomass burning (as shown in Fig. 9) and as noticed

## Tropical tropospheric ozone

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

by [Edwards et al. \(2003\)](#) in their analysis of CO retrieval from the MOPITT satellite instrument. During DJF, O<sub>3</sub> enhancements confined to the lower troposphere over West Africa come with the highest tropical CO concentrations measured by the MOZAIC program, with 200–500 ppbv at Lagos below 700 hPa (Fig. 9) and 200–400 ppbv below 500 hPa at Douala (not shown). High CO concentrations originating from biomass burning fires over Central Africa are measured near 600–700 hPa over Lagos and Douala during JJA. Trade winds allow this CO transport and the associated O<sub>3</sub> enhancement ([Sauvage et al., 2005](#)). Aghedo et al. (2006) found also high influence of biomass burning on surface O<sub>3</sub> near 1000 hPa. A persistent CO enhancement that may reflect local pollution is observed at Delhi, with more than 150 ppbv below 800 hPa. No CO measurements are performed south of the ITCZ.

As a result of the GOME constraints on surface emissions, the simulation better reproduces lower tropospheric O<sub>3</sub>. During DJF over Lagos and Abidjan, the intensity of the lower tropospheric O<sub>3</sub> enhancement is now well reproduced mostly because of the higher NO<sub>x</sub> emission factors that increase O<sub>3</sub> by 15–20 ppbv (+45%) compared to the original version. Five ppbv of the 15–20 ppbv increase are attributed to the additional biomass burning VOCs. Moreover, Brazzaville shows an O<sub>3</sub> enhancement in the lower troposphere through inter-hemispheric transport (+15/20 ppbv (+55%) compared to the original version).

During JJA over Brazzaville the intensity of the O<sub>3</sub> maximum is also better reproduced (+10 ppbv/+14%), as a consequence of both the higher NO<sub>x</sub> and VOC emissions. These emissions also yield a better reproduction of the O<sub>3</sub> enhancement at Lagos through inter hemispheric transport. The enhancement near 600–700 hPa is also increased by 7 ppbv due to the a posteriori soil NO<sub>x</sub> emissions.

There are improvements associated with biomass burning emissions over the South America Cerrado in SON, and over India in MAM. O<sub>3</sub> decreases in the lower troposphere of Bombay by around 5 to 7 ppbv (8–10%). However O<sub>3</sub> is still too high in the lower troposphere, perhaps reflecting a combination of local sea breeze, missing halogen chemistry (Dickerson et al., 1999; Stehr et al., 2002), or inefficient O<sub>3</sub> production in

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

urban areas (Dickerson et al., 2002) not resolved at the coarse resolution of the model.

The top-down emissions also affect the TOC (Table 3). Over the Gulf of Guinea (Lagos) the standard simulation is within 1 DU of MOZAIC versus 6 DU for the original simulation. Over Central Africa the standard simulation is within 2 DU versus MOZAIC during DJF compared to 6 DU in the original simulation. The seasonal cycle is well reproduced, with maximum during JJA and minimum during MAM over Central Africa, maximum during DJF and minimum during JJA over West Africa. Over Windhoek the bias is within 1–2 DU for all the seasons. Over India modeled TOC is within 1–5 DU of MOZAIC during all seasons except JJA, reflecting the lower tropospheric bias.

Comparisons between GOME and GEOS-Chem TOC also show substantial improvements. Most of the differences between GOME and the standard GEOS-Chem TOC are within 3 DU. The largest differences appear in the northern tropics, with a negative bias of 5 to 8 DU between GOME and GEOS-Chem. Table 3 shows that the GOME TOC underestimate MOZAIC in this region, perhaps reflecting the low sensitivity of GOME to lower tropospheric O<sub>3</sub>, especially in the presence of aerosols from biomass burning or mineral dust. A retrieval of tropospheric O<sub>3</sub> using the scan-angle method better captures lower tropospheric O<sub>3</sub> (Kim et al., 2005). Instrument sensitivity may also play a role over Central Africa during JJA, when GOME TOC biased by 10 DU compared to MOZAIC at Brazzaville.

#### 4.2.3 Biomass burning emission factors

We compare the standard simulation with a sensitivity simulation using a recent compilation of biomass burning emission factors (EF) from Andreae (Andreae and Merlet, 2001, personal communication, 2005), that were compiled from in situ measurements. The main differences versus the original simulation are a 23% lower NO<sub>x</sub> EF for savannas/grassland and a 15% higher NO<sub>x</sub> EF for tropical forest fires. The new EF for savanna and grassland also feature 100% higher values for HCHO and 200% for alkenes.

The green line in Fig. 8 shows the O<sub>3</sub> simulation using the in-situ-based emission

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

factors. The in-situ-based emission factors reduce the original bias versus O<sub>3</sub> over West Africa but still yield insufficient O<sub>3</sub> in contrast with the top-down emissions. Over Central and South Africa, in-situ-based emission factors increase O<sub>3</sub> by 10–15 ppbv in JJA with respect to the original simulation, 5 ppbv more than the standard simulation.

5 During DJF O<sub>3</sub> is 20 ppbv higher than the standard simulation. Over India during the biomass burning season, the new emission factors have no effect on the lower tropospheric O<sub>3</sub> distribution, in contrast with the space-based constraint, which decreases O<sub>3</sub> by 5 ppbv yielding a simulation more consistent with results to in situ measurements. The amount of biomass burned may be responsible for the bias in the original  
10 simulation as evident from a similar bias in CO (Heald et al., 2003).

In summary the recent compilation has similarities with the top-down emissions, but less successfully reproduces O<sub>3</sub> observations. We go on to infer regional NO<sub>x</sub> and VOC emission factors from the top-down inventory over Africa and the bottom-up estimate of biomass burned. The resulting emission factors for savanna/grassland  
15 fires are 2.9 gNO/kg over North Africa, 4.3 gNO/kg over Central/South Africa and 3.1 gNO/kg over the South American Cerrado. This leads to 3.4 gNO/kg mean for savanna/grassland, at the upper limit of the recommendation from Andreae (Andreae and Merlet, 2001, Andreae, personal communication, 2005) with 2.3±1.1 gNO/kg. For tropical forest fires the top-down EF are 2.3 gNO/kg over North Africa, 2.6 gNO/kg over  
20 Central/South Africa and 2.0 gNO/kg over South America leading to 2.3 gNO/kg mean versus 1.8±0.7 from Andreae (Andreae and Merlet, 2001, Andreae, personal communication, 2005).

The resulting emission factors for savanna/grassland fires are 0.96 g/kg for alkenes, and 0.7 g/kg for HCHO close to the recommendation by Andreae (Andreae and Merlet, 2001, Andreae, personal communication, 2005) with 1.1±0.6 g/kg and 0.7±0.4 g/kg for HCHO.  
25

---

**Tropical tropospheric ozone**B. Sauvage et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

#### 4.2.4 Sensitivity to biogenic emissions

We explore whether the HCHO bias in the original simulation could be related to isoprene emissions by conducting a sensitivity simulation using the recent Model Emissions of Gases and Aerosols from Nature (MEGAN) inventory (Guenther et al., 2006).

5 This inventory yields 600 Tg C/yr of isoprene emissions and has improved the HCHO simulation over the United States (Palmer et al., 2006). However the simulation using MEGAN (not shown) increases HCHO columns over the Amazonian and Equatorial African forest, for all seasons. The general effect is to introduce an overestimate of the tropospheric HCHO columns outside of the wet and biomass burning seasons. More-  
10 over the MEGAN inventory generally decreases  $O_3$  in the lower and middle troposphere by 3–5 ppbv, reducing agreement with in situ  $O_3$  data. In summary there is a higher consistency in the comparison of GOME vs. GEOS-Chem standard simulation than in the comparison of GOME vs. GEOS-Chem simulation using MEGAN, supporting the previous conclusion of an underestimate of biomass burning VOCs in the bottom-up  
15 emission inventory.

#### 4.3 Sensitivity to heterogeneous chemistry

Here we examine the implications of the heterogeneous chemistry updates described in Sect. 3.2.4, specifically the neglect of  $HO_2$  uptake on biomass burning aerosols, and the uptake of  $HNO_3$  on mineral dust. We also explore the effect of direct  $O_3$  destruction on mineral dust. This section provides a first overall evaluation of these processes,  
20 through comparison with in situ  $O_3$  measurements over a broad area.

The exclusion of  $HO_2$  uptake on biomass burning aerosols in our standard simulation systematically increases modeled  $O_3$  over biomass burning regions by 5–7 ppbv, improving the consistency with in situ measurements as shown in Fig. 8. Elsewhere  
25 no effect is found over the Tropics.

Figure 10 shows vertical profiles of  $O_3$  at locations and seasons in which  $HNO_3$  uptake had a large effect. As found by Bauer et al. (2004) heterogeneous uptake of  $HNO_3$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Tropical tropospheric ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

reduces  $O_3$  primarily over and downwind of deserts, i.e. northern Africa and South America, the Arabic peninsula, and India. We find that the reduction in  $O_3$  reduces model biases compared to in situ measurements. There is significant improvement over Dubai and Bombay during March to November, when  $O_3$  is reduced by 10–15%.

The simulated  $O_3$  column maximum over the Middle East is reduced by this process. Over Caracas, there is a 3–5 ppbv (10%)  $O_3$  decrease below 800 hPa from November to August, when there is a long range transport from the Sahara. Over Lagos  $HNO_3$  uptake reduces  $O_3$  by a maximum of 5%.

The uptake of  $HNO_3$  on mineral dust implemented here, using a reaction probability formulation for convenience, likely represents an upper limit. The particle alkalinity would likely be consumed during continued exposure to  $HNO_3$  and  $H_2SO_4$  and would be better represented in an equilibrium partitioning. Aerosol nitrate could photolyze to regenerate  $NO_x$  (Anastasio and Mc Gregor, 2001). Nonetheless, we find observational evidence in support of the reaction.

We also explored the effect of direct  $O_3$  destruction on mineral dust using  $\gamma(O_3)=10^{-5}$  as recommended from recently laboratory measurements by Hanisch and Crowley (2003). The effect of this reaction on  $O_3$  is smaller than that of  $HNO_3$  uptake as found by Bauer et al. (2004). However  $O_3$  uptake had a large negative role over Lagos during DJF in the lower troposphere, leading to a 15–20% reduction of the  $O_3$  biomass burning enhancement.

In summary,  $HNO_3$  uptake on mineral dust and the exclusion of  $HO_2$  uptake on biomass burning aerosols improves the simulation versus MOZAIC and SHADOZ sites. This is not the case for  $O_3$  uptake, which had no effect over the Middle East and India, and a negative effect over West Africa.

#### 4.4 Sensitivity to dynamics

Convective transport has considerable implications for upper tropospheric  $O_3$  (Lelieveld and Crutzen, 1999; Lawrence et al., 2003; Diab et al., 2004; Folkins and Martin, 2005; Rasch et al., 1997). The Goddard Earth Observing System data assimilation system

at the NASA Global Modeling and Assimilation office provides two different assimilated meteorological datasets, GEOS-3 and GEOS-4, for the year 2000. Three major differences between the two assimilations are the convective parameterization, the cloud optical depths, and cloud top heights. GEOS-3 uses the Relaxed Arakawa Schubert (Moorthi and Suarez, 1992) convective parameterization, and GEOS-4 uses the Zhang and McFarlane (Zhang and McFarlane, 1995) convective parameterization. As discussed by Folkins et al. (2006) the tropical cloud divergence is quite weak at all altitudes with GEOS-3, and is stronger in the upper troposphere with GEOS-4 model. Cloud optical depths are smaller in GEOS-4 than GEOS-3 leading to more active photochemistry (Liu et al., 2006). Cloud top heights are higher in GEOS-3 than GEOS-4 (Wu et al., 2006). We compare our standard simulation driven with GEOS-4 meteorological fields with one driven with GEOS-3 at the MOZAIC and SHADOZ sites for O<sub>3</sub> (Fig. 10); and also for CO and RH (Fig. 9). For clarity Fig. 10 contains sites and seasons that exhibited a high sensitivity to the dynamical scheme.

As shown in Fig. 10, the main differences in O<sub>3</sub> between the standard simulation using GEOS-4 and GEOS-3 are found in the middle and upper troposphere. GEOS-3 substantially overestimates O<sub>3</sub> compared to measurements, over all continental and oceanic regions, by 10–25 ppbv (15%–50%) with even higher overestimates over South America, the Middle East, and the Pacific. The main discrepancy above 400 hPa likely reflects an underestimate of convective detrainment which injects O<sub>3</sub> depleted air as shown by Folkins et al. (2006) with SHADOZ measurements. As a consequence, RH and CO modeled with GEOS-3 are generally more underestimated compared to MOZAIC at those levels (Fig. 9), than with the GEOS-4 standard simulation. These effects are apparent in a meridional average.

Figure 11 shows a meridional average (5° W–30° E), along MOZAIC flight altitudes, 200–300 hPa, during the monsoon season (JJA). MOZAIC data depict the ITCZ position over Africa (0–10° N) with depleted O<sub>3</sub> and enhanced RH and CO. GEOS-3 underestimates O<sub>3</sub> and overestimates RH in contrast with GEOS-4. However GEOS-4 overestimates the CO gradient versus the few CO measurements that are available.

**Tropical tropospheric ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Tropical tropospheric  
ozone**

B. Sauvage et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

There are few instances where MOZIC measurements are more consistent with GEOS-3 than with GEOS-4. For example during SON over Lagos, Ascension (Fig. 10), and Reunion, upper tropospheric  $O_3$  measurements are better reproduced with GEOS-3, than with GEOS-4. This bias at Ascension appears to reflect the low altitude of convective outflow in GEOS-4, as supported by the improvement in the simulation with enhanced intracloud lightning.

The two assimilation schemes also affect trace gases in the lower troposphere at some locations (Fig. 10, Lagos, Ascension, Bombay). The GEOS-3 simulation exhibits lower  $O_3$  concentrations than with GEOS-4 and in situ measurements. In contrast both GEOS-3 and GEOS-4 underestimate lower tropospheric CO at Lagos and Delhi (Fig. 9), but the simulation is within one standard deviation of the measurements and the CO seasonal cycle well reproduced. Both simulations are able to capture the lower tropospheric maximum in CO associated with the biomass burning season, in DJF and MAM over Lagos, and MAM over Delhi. CO is more sensitive than  $O_3$  to dynamics in the lower troposphere, reflecting the stronger gradients in CO. In the lower troposphere GEOS-4 CO is lower than GEOS-3, likely reflecting lower cloud optical depth that results in more active chemistry and more active convection that would transport CO from the lower troposphere.

The lower tropospheric CO underestimate with both GEOS-3 and GEOS-4, suggests an underestimate of CO emissions. We examine the possible implications in our  $O_3$  simulation by increasing CO biomass burning emissions by a factor 2. However the effect on  $O_3$  is negligible, increasing the  $O_3$  background by 3 ppbv.

More than a simple overview of two different meteorological datasets, this comparison clearly shows convection and clouds as major processes driving tropospheric  $O_3$ . These processes may be as important as the remaining uncertainties in chemical processes and emissions sources.

## 5 Conclusions

We used a global chemical transport model (GEOS-Chem) to evaluate the consistency of satellite measurements and to examine the processes affecting tropospheric O<sub>3</sub> over the Tropics. Space-based observations from the Global Ozone Monitoring Experiment (GOME), Optical Transient Detector (OTD) and Lightning Imaging sensor (LIS) instruments are used to constrain the model emissions necessary for an accurate estimation and understanding of processes affecting tropical tropospheric ozone. In-situ measurements from the Measurements of ozone and water vapor by in-service Airbus aircraft (MOZAIC) aircraft program and the Southern Hemisphere Additional Ozonesondes (SHADOZ) ozonesonde network, were subsequently used to evaluate the simulation.

Our standard simulation featured substantial modifications over the original simulation. A climatology of flash counts from the OTD and LIS instruments are used to improve the spatial distribution of lightning NO<sub>x</sub> emissions in the model. Tropospheric NO<sub>2</sub> and HCHO columns retrieved from GOME are applied to provide top-down constraints on emission inventories of NO<sub>x</sub> (biomass burning and soils) and VOCs (biomass burning). We remove HO<sub>2</sub> uptake on biomass burning aerosols, and implement HNO<sub>3</sub> uptake on mineral dust.

Upper tropospheric O<sub>3</sub> is highly sensitive to the spatial distribution of lightning NO<sub>x</sub> emissions. The lightning rescaling improves the simulation of middle and upper tropospheric O<sub>3</sub> for tropical sites, by 5–15 ppbv (10%–45%) versus in situ measurements from SHADOZ and MOZAIC. Biases in the simulation of tropospheric ozone columns are reduced by 1–6 DU versus GOME, MOZAIC and SHADOZ measurements. We evaluate lightning emissions in terms of intensity, by testing ±2 Tg N/yr around the 5 Tg N/yr used in the standard simulation; and in term of distribution by increasing the NO<sub>x</sub> emitted from intracloud lightning. A lightning source strength of 5±2 Tg N/yr best represents in situ observations from MOZAIC and SHADOZ. Increasing the ratio of intra-cloud (IC) to cloud-ground (CG) lightning NO emissions from 0.1 to 0.75 generally introduces an O<sub>3</sub> overestimate compared to in situ measurements. However,

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

substantial improvements are found at Ascension and Reunion during SON. A global mean increase in intra-cloud lightning  $\text{NO}_x$  is not supported by in situ  $\text{O}_3$  profiles. Prognostic determination of the IC/CG ratio could yield an improved simulation of tropical ozone.

5 The top-down constraints on  $\text{NO}_x$  emissions inferred from GOME  $\text{NO}_2$  columns increase biomass burning emissions, by a factor of 1.1 over Central Africa and by a factor of 2.6 over North Africa. The  $\text{NO}_x$  emission factor inferred from GOME  $\text{NO}_2$  columns over savanna/grassland is  $3.4 \text{ gNO/kg dm}$ , 40% higher than the recommendation by Andreae (Andreae and Merlet, 2001, personal communication, 2005) but within the  
10 given range. The GOME HCHO columns provide a measure of reactive VOC emissions. An increase in HCHO and alkenes emissions by a factor of 2 over biomass burning regions is necessary to reproduce GOME observations of HCHO columns. The top-down emissions increase the simulation of lower tropospheric ozone by 5–20 ppbv, improving the simulation versus MOZAIC in situ measurements, mainly over  
15 Africa where  $\text{O}_3$  is most sensitive to surface sources. The improvement in simulated  $\text{O}_3$  provides an indirect validation of the retrieved tropospheric  $\text{NO}_2$  and HCHO columns. The modeled TOC are within 1–3 DU of GOME, and within 1–4 DU compared to in situ measurements. The seasonal variations are well reproduced.

We evaluate the biogenic a posteriori  $\text{NO}_x$  emission inventory (Jaeglé et al., 2005)  
20 versus in situ  $\text{O}_3$  measurements. The largest influence appears over Africa and adjacent regions in MAM/JJA, with  $\text{O}_3$  increasing by 5–7 ppbv, and reducing a regional model bias.

We drive GEOS-Chem with two different assimilation schemes, GEOS-3 and GEOS-4, that feature different convective parameterizations and cloud fields. The two different  
25 dynamical schemes have considerable effect on the ozone simulation. GEOS-4 better represents  $\text{O}_3$  observations by 5–20 ppbv due to enhanced convective detrainment in the upper troposphere, compared to GEOS-3 which overestimates  $\text{O}_3$ . The role of enhanced convective outflow is particularly apparent in relative humidity and  $\text{O}_3$  in the upper troposphere across the ITCZ over Africa. The two assimilated fields most affect

---

**Tropical tropospheric ozone**B. Sauvage et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

carbon monoxide in the lower troposphere, and observations are better reproduced with GEOS-3 which has higher cloud optical depths.

Recent laboratory and field measurements provide evidence for uptake of  $\text{HNO}_3$  ( $\gamma_{\text{HNO}_3}=0.1$ ) on mineral dust, and the absence of  $\text{HO}_2$  uptake on biomass burning aerosols. We evaluate those processes with in situ measurements of  $\text{O}_3$ .  $\text{HNO}_3$  uptake reduces a regional model bias by 5–15% downwind of deserts. The neglect of  $\text{HO}_2$  uptake on biomass burning aerosols increases simulated  $\text{O}_3$  by 5 ppbv, improving our simulations versus in situ measurements in biomass burning regions. Direct uptake of  $\text{O}_3$  ( $\gamma_{\text{O}_3}=10^{-5}$ ) on mineral dust introduces a large model bias compared to MOZAIC  $\text{O}_3$  measurements over West Africa.

We have shown that satellite observations of lightning and  $\text{O}_3$  precursors improve substantially the simulation of tropical tropospheric  $\text{O}_3$  with a global chemical transport model due to better representation of emissions. The most prominent outstanding issues are related to lightning and convection. Future development of a prognostic parameterization of lightning that reproduces observed flash counts, should improve the accuracy of  $\text{O}_3$  simulations. In-situ measurements of trace gases in close proximity to deep convection in the Tropics would enable disentangling of issues related to lightning vertical profile and convective transport. Forthcoming high resolution space-based data, such as from Aura (Schoeberl et al., 2004), or GOME-2 and IASI should continue to provide additional insight into tropical tropospheric ozone.

*Acknowledgements.* The authors acknowledge MOZAIC funding agencies, the European Commission, CNRS (France), Forschungszentrum Jülich (Germany), Météo France, EADS (Airbus) and the airlines (Air France, Lufthansa, Austrian Airlines, and former Sabena who carry free of charge the MOZAIC equipment and perform the maintenance). We also acknowledge the European Agency, ESA/ERIN. SHADOZ is supported by NASA's Atmospheric Chemistry and Analysis Program (ACMAP), Aura Validation, and the TOMS and OMI projects. We thank D. Boccippio for providing the lightning flash counts used in this analysis. We thank M. O. Andreae for providing his biomass burning emission factors data base. This work was supported by the Atmospheric Composition Program of NASA's Earth-Sun System Division.

**Tropical tropospheric ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## References

- Aghebo, A. M., Schultz, M. G., and Rast, S.: The influence of African air pollution on regional and global tropospheric chemistry, *Atmos. Chem. Phys. Discuss.*, 6, 5797–5838, 2006.
- Alexander, B. J., Park, R. J., Jacob, D. J., Li, Q. B., Yantosca, R. M., Savarino, J., Lee, C. C. W., and Thiemens, M. H.: Sulfate formation in sea-salt aerosols: Constraints from oxygen isotopes, *J. Geophys. Res.*, 110, D10307, doi:10.1029/2004JD005659, 2005.
- Anastasio, C. and McGregor, K. G.: Chemistry of fog waters in California's Central Valley: 1. In situ photoformation of hydroxyl radical and singlet molecular oxygen, *Atm. Env.*, 35(6), 1079–1089, 2001. [11488](#)
- Andreae, M. O.: The influence of tropical biomass burning on climate and the atmospheric environment, in: *Biogeochemistry of Global Change: Radiatively Active Trace Gases*, edited by: Oremland, R. S., 113–150, Chapman and Hall, New York, 1993. [11467](#)
- Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cycles*, 15(4), 955–966, doi:10.1029/2000GB001382, 2001. [11485](#), [11486](#), [11492](#)
- Andreae, M. O., Artaxo, P., Fischer, H., et al.: Transport of biomass burning smoke to the upper troposphere by deep convection in the equatorial region, *Geophys. Res. Lett.*, 28(6), 951–954, 2001.
- Andreae, M. O., Rosenfeld, D., Artaxo, P., Costa, A. A., Frank, G. P., Longo, K. M., and Silva-Dias, M. A. F.: Smoking Rain Clouds over the Amazon, *Science*, 303, 1337–1342, 2004.
- Burrows, J. P., Weber, M., Buchwitz, M., et al.: The Global Ozone Monitoring Experiment (GOME): Mission concept and first scientific results, *J. Atmos. Sci.*, 56, 151–175, 1999. [11470](#)
- Bauer, S. E., Balkanski, Y., Schulz, M., Hauglustaine, D. A., and Dentener, F.: Global modeling of heterogeneous chemistry on mineral aerosol surfaces: Influence on tropospheric ozone chemistry and comparison to observations, *J. Geophys. Res.*, 109, D02304, doi:10.1029/2003JD003868, 2004. [11477](#), [11487](#), [11488](#)
- Bertram, T. H., Heckel, A., Richter, A., Burrows, J., and Cohen, R. C.: Satellite measurements of daily variations in soil NO<sub>x</sub> emissions, *Geophys. Res. Lett.*, 32, L24812, doi:10.1029/2005GL024640, 2005.
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and Schultz, M. G.: Global modeling of tropospheric Chemistry with

ACPD

6, 11465–11520, 2006

## Tropical tropospheric ozone

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

- assimilated meteorology: Model description and evaluation, *J. Geophys. Res.*, 106, 23 073–23 096, 2001. [11472](#)
- Bian, H. and Zender, C. S.: Mineral dust and global tropospheric chemistry: Relative roles of photolysis and heterogeneous uptake, *J. Geophys. Res.*, 108(D21), 4672, doi:10.1029/2002JD003143, 2003.
- Boccippio, D. J., Cummins, K. L., Christian, H. J., and Goodman, S. J.: Combined satellite and surface-based estimation of the intracloud cloud-to-ground lightning ratio over the continental United States, *C. Mon. Wea. Rev.*, 129, 108–122, 2001.
- Boccippio, D. J., Goodman, S. J., and Heckman, S.: Regional differences in tropical lightning distributions, *J. Appl. Met.*, 39, 2231–2248, 2000.
- Boccippio, D. J., Driscoll, K. T., Koshak, W. J., Blakeslee, R. J., Boeck, W. L., Mach, D. A., Buechler, D. E., Christian, H. J., and Goodman, S. J.: The Optical Transient Detector (OTD): Instrument characteristics and cross-sensor validation, *J. Atmos. Oc. Tech.*, 17, 441–458, 2000.
- Boccippio, D. J.: Lightning scaling laws revisited, *J. Atmos. Sci.*, 59, 1086–1104, 2002.
- Boersma, K. F., Eskes, H. J., and Brinksma, E. J.: Error analysis for tropospheric NO<sub>2</sub> retrieval from space, *J. Geophys. Res.*, 109, D04311, doi:10.1029/2003JD003962, 2004. [11471](#)
- Chatfield, R. B. and Delany, A. C.: Convection links biomass burning to increased tropical ozone: However, models will tend to overpredict O<sub>3</sub>, *J. Geophys. Res.*, 95, 18 473–18 488, 1990. [11467](#)
- Chance, K., Palmer, P. I., Spurr, R. J. D., Martin, R. V., Kurosu, T. P., and Jacob, D. J.: Satellite observations of formaldehyde over North America from GOME, *Geophys. Res. Lett.*, 27, 3461–3464, doi:10.1029/2000GL011857, 2000. [11470](#), [11471](#)
- Chandra, S., Ziemke, J. R., Bhartia, P. K., and Martin, R. V.: Tropical tropospheric ozone: Implications for biomass burning, *J. Geophys. Res.*, 107, doi:10.1029/2001JD000447, 2002.
- Chandra, S., Ziemke, J. R., and Martin, R. V.: Tropospheric ozone at tropical and middle latitudes derived from TOMS/MLS residual: Comparison with a global model, *J. Geophys. Res.*, 108, doi:10.1029/2002JD002912, 2003.
- Christian, H. J., Blakeslee, R. J., Boccippio, D. J., Boeck, W. L., Buechler, D. E., Driscoll, K. T., Goodman, S. J., Hall, J. M., Mach, D. M., and Stewart, M. F.: Global frequency and distribution of lightning as observed from space by the Optical Transient Detector, *J. Geophys. Res.*, 108, doi:10.1029/2002JD002347, 2003. [11467](#)
- Christian, H. J., Blakeslee, R. J., and Goodman, S. J.: The Detection of Lightning from Geosta-

**Tropical tropospheric  
ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- tionary Orbit, *J. Geophys. Res.*, 94, 13 329–13 337, 1989. [11470](#)
- de Forster, F., P. M., and Shine, K. P.: Radiative forcing and temperature trends from stratospheric ozone changes, *J. Geophys. Res.*, 102(D9), 10 841–10 856, 1997.
- DeCaria, A. J., Pickering, K. E., Stenchikov, G. L., Scala, J. R., Stith, J. L., Dye, J. E., Ridley, B. A., and Laroche, P.: A cloud-scale model study of lightning-generated  $NO_x$  in an individual thunderstorm during STERAO-A, *J. Geophys. Res.*, 105(D9), doi:10.1029/2000JD900033, 2000.
- Diab, R. D., Thompson, A. M., Mari, K., Ramsay, L., and Coetzee, G. J. R.: Tropospheric ozone climatology over Irene, South Africa, from 1990 to 1994 and 1998 to 2002, *J. Geophys. Res.*, 109, D20301, doi:10.1029/2004JD004793, 2004.
- Dickerson, R. R., Rhoads, K. P., Carsey, T. P., Oltmans, S. J., Burrows, J. P., and Crutzen, P. J.: Ozone in the remote marine boundary layer: A possible role for halogens, *J. Geophys. Res.*, 104(D17), 21 385–21 396, 1999.
- Dickerson, R. R., Andreae, M. O., Campos, T., Mayol-Bracero, O. L., Neusuess, C., and Streets, D. G.: Analysis of black carbon and carbon monoxide observed over the Indian Ocean: Implications for emissions and photochemistry, *J. Geophys. Res.*, 107(D19), 8017, doi:10.1029/2001JD000501., 2002.
- Duncan, B. N., Martin, R. V., Staudt, A. C., Yevich, R. M., and Logan, J. A.: Interannual and seasonal variability of biomass burning emissions constrained by remote-sensed observations, *J. Geophys. Res.*, 108, 4040, doi:10.1029/2002JD002378, 2003. [11473](#)
- Edwards, D. P., Lamarque, J. F., Attié, J. L., et al.: Tropospheric ozone over the Tropical Atlantic: a satellite perspective, *J. Geophys. Res.*, 108(D8), doi:10.1029/2002JD002927, 2003. [11484](#)
- Evans, M. J. and Jacob, D. J.: Impact of new laboratory studies of  $N_2O_5$  hydrolysis on global model budgets of tropospheric nitrogen oxides, ozone, and OH, *Geophys. Res. Lett.*, 32, L09813, doi:10.1029/2005GL022469, 2005.
- Fehr, T., Holler, H., and Huntrieser, H.: Model study on production and transport of lightning-produced  $NO_x$  in a EULINOX supercell storm, *J. Geophys. Res.*, 109, D09102, doi:10.1029/2003JD003935, 2004.
- Folkens, I. and Martin, R. V.: The vertical structure of tropical convection and its impact on the budgets of water vapor and ozone, *J. Atm. Sci.*, 62(5), 1560–1573, 2005.
- Folkens, I., Bernath, P., Boone, C., Eldering, A., Lesins, G., Martin, R. V., Sinnhuber, B.-M., and Walker, K.: Testing convective parameterizations with tropical measurements of  $HNO_3$ , CO,  $H_2O$ , and  $O_3$ : implications for the water vapor budget, *J. Geophys. Res.*, accepted, 2006.

**Tropical tropospheric  
ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, 2006. [11475](#), [11487](#)
- 5 Guerova, G., Bey, I., Attié, J.-L., Martin, R. V., Cui, J., and Sprenger, M.: Impact of transatlantic transport episodes on summertime ozone in Europe *Atmos. Chem. Phys.*, 6, 2057–2072, 2006. [11470](#)
- Goodman, A. L., Underwood, G. M., and Grassian, V. H.: A laboratory study of the heterogeneous resaction of nitric acid on calcium carbonate particles, *J. Geophys. Res.*, 105, D23, 29 053–29 064, 2000.
- 10 Grassian, V. H.: Heterogeneous uptake and reaction of nitrogen oxides and volatile organic compounds on the surface of atmospheric particles including oxides, carbonates, soot and mineral dust: Implications for the chemical balance of the troposphere, *Int. Rev. Phys. Chem.*, 20(20), 467–548, 2001.
- 15 Hanisch, F. and Crowley, J. N.: Ozone decomposition on Saharan dust: an experimental investigation, *Atmos. Chem. Phys.*, 3, 119–130, 2003. [11488](#)
- Hastenrath, S.: *Climate and circulations of the tropics*, D. Reidel, Dordrecht, 1985.
- Heald, C. L., Jacob, D. J., Fiore, A. M., et al.: Asian outflow and trans-Pacific transport of carbon monoxide and ozone pollution: An integrated satellite, aircraft, and model perspective, *J. Geophys. Res.*, 108(D24), doi:10.1029/2003JD003507, 2003. [11486](#)
- 20 Holland, E. A., Dentener, F. J., Braswell, B. H., and Sulzman, J. M.: Contemporary and pre-industrial global reactive nitrogen budgets, *Biogeochem.*, 46, 7–43, 1999 [11467](#)
- Hoskins, B. J. and Rodwell, M. J.: A model of the Asian summer monsoon. Part I: the global scale, *J. Atmos. Sci.*, 52(9), 1329–1340, 1995.
- 25 Jaeglé, L., Martin, R. V., Chance, K., Steinberger, L., Kurosui, T. P., Jacob, D. J., Modi, A. I., Yoboué, V., Sigha-Nkamdjou, L., and Galy-Lacaux, C.: Satellite mapping of rain-induced nitric oxide emissions from soils, *J. Geophys. Res.*, 109, D21310, doi:10.1029/2004JD004787, 2004. [11467](#), [11474](#)
- Jaeglé, L., Steinberger, L., Martin, R. V., and Chance, K.: Global partitioning of NO<sub>x</sub> sources using satellite observations: Relative roles of fossil fuel combustion, biomass burning and soil emissions, *Faraday Discuss.*, 130, 407–423, doi:10.1039/b502128f, 2005. [11474](#), [11492](#)
- 30 Jacob, D. J., Heikes, B. G., Fan, S.-M., et al.: Origin of ozone and NO<sub>x</sub> in the tropical troposphere: A photochemical analysis of aircraft observations over the South Atlantic basin, *J.*

---

**Tropical tropospheric  
ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Geophys. Res., 101(D19), 24 235–24 250, 1996. [11467](#)
- Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, *Atm. Env.*, 34, 2131–2159, 2000.
- Kim, J. H., Na, S., Newchurch, M. J., and Martin, R. V.: Tropical tropospheric ozone morphology and seasonality seen in satellite and in situ measurements and model calculations, *J. Geophys. Res.*, 110, D02303, doi:10.1029/2003JD004332, 2005. [11485](#)
- Kurosu, T. P., Chance, K., and Spurr, R. J. D.: CRAG-cloud retrieval algorithm for ESA's global ozone monitoring experiment, ESA WPP-161, 513–521, Eur. Space Res. and Tech Cent., Noordwijk, Netherlands, 1999. [11471](#)
- Labrador, L. J., von Kuhlmann, R., and Lawrence, M. G.: The effects of lightning-produced NO<sub>x</sub> and its vertical distribution on atmospheric chemistry: sensitivity simulations with MATCH-MPIC, *Atmos. Chem. Phys.*, 5, 1815–1834, 2005. [11481](#)
- Lacis, A. A., Wuebbles, D. J., and Logan, J. A.: Radiative forcing of climate by changes in the vertical distribution of ozone, *J. Geophys. Res.*, 95(D7), 9971–9981, 1990.
- Lawrence, M. G., von Kuhlmann, R., Salzmann, M., and Rasch, P. J.: The balance of effects of deep convective mixing on tropospheric ozone, *Geophys. Res. Lett.*, 30(18), doi:10.1029/2003GL017644, 2003.
- Lee, D. S., Kohler, I., Grobler, E., Rohrer, F., Sausen, R., Gallardo-Klenner, L., Olivier, J. H. J., Dentener, F. J., and Bouwman, A. F.: Estimation of global NO<sub>x</sub> emissions and their uncertainties, *Atm. Env.*, 31, 1735–1749, 1997.
- Leue, C., Wenig, M., Wagner, T., Klimm, O., Platt, U., and Jahne, B.: Quantitative analysis of NO<sub>x</sub> emissions from Global Ozone Monitoring Experiment satellite image sequences, *J. Geophys. Res.*, 106, 5493–5506, doi:10.1029/2000JD900572, 2001.
- Lelieveld, J. and Crutzen, P. J.: Role of deep cloud convection in the ozone budget of the troposphere, *Science*, 264, 1759–1761, 1994.
- Li, Q., Jacob, D. J., Logan, J. A., et al.: A tropospheric ozone maximum over the Middle East, *Geophys. Res. Lett.*, 28, 3235–3238, 2001.
- Liao, H. and Seinfeld, J. H.: Global impacts of gas-phase chemistry-aerosol interactions on direct radiative forcing by anthropogenic aerosols and ozone, *J. Geophys. Res.*, 110, D18208, doi:10.1029/2005JD005907, 2005.
- Liu, X., Chance, K., Sioris, C. E., Spurr, R. J. D., Kurosu, T. P., Martin, R. V., and Newchurch, M. J.: Ozone profile and tropospheric ozone retrievals from the Global Ozone Monitoring Experiment: Algorithm description and validation, *J. Geophys. Res.*, 110, D20307,

**Tropical tropospheric ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

doi:10.1029/2005JD006240, 2005. [11471](#)

Liu, X., Chance, K., Sioris, C. E., et al.: First directly retrieved global distribution of tropospheric column ozone from GOME: Comparison with the GEOS-CHEM model, *J. Geophys. Res.*, 111, D02308, doi:10.1029/2005JD006564, 2006. [11479](#), [11489](#)

5 Liu, H., Crawford, J. H., Pierce, R. B., et al.: Radiative effect of clouds on tropospheric chemistry in a global three-dimensional chemical transport model, *J. Geophys. Res.*, 111(D20), D20303, doi:10.1029/2005JD006403, 2006.

Logan, J. A., Prather, M. J., and Hierro, R. F.: Tropospheric chemistry – A Global perspective, *J. Geophys. Res.*, 86, 7210–7254, 1981. [11467](#)

10 Marengo, A., Thouret, V., Nédélec, P., et al.: Measurement of ozone and water vapor by Airbus in-service aircraft: The MOZAIC airborne program, An overview, *J. Geophys. Res.*, 103, 25 631–25 642, 1998. [11469](#)

Martin, R. V., Jacob, D. J., Logan, J. A., et al.: Interpretation of TOMS observations of tropical tropospheric ozone with a global model and in situ observations, *J. Geophys. Res.*, 107(D18), 4351, doi:10.1029/2001JD001480, 2002a. [11467](#)

15 Martin, R. V., Chance, K., Jacob, D. J., et al.: An improved retrieval of tropospheric nitrogen dioxide from GOME, *J. Geophys. Res.*, 107(D20), doi:10.1029/2001JD001027, 2002b. [11470](#), [11471](#)

20 Martin, R. V., Jacob, D. J., Chance, K., Kurosu, T. P., Palmer, P. I., and Evans, M. J.: Global inventory of nitrogen oxide emissions constrained by space-based observations of NO<sub>2</sub> columns, *J. Geophys. Res.*, 108(D17), 4537, doi:10.1029/2003JD003453, 2003a. [11471](#), [11474](#)

25 Martin, R. V., Jacob, D. J., Yantosca, R. M., Chin, M., and Ginoux, P.: Global and regional decreases in tropospheric oxidants from photochemical effects of aerosols, *J. Geophys. Res.*, 108(D3), 4097, doi:10.1029/2002JD002622, 2003b. [11473](#)

Martin, R. V., Parrish, D. D., Ryerson, T. B., Nicks Jr., D. K., Chance, K., Kurosu, T. P., Jacob, D. J., Sturges, E. D., Fried, A., and Wert, B. P.: Evaluation of GOME satellite measurements of tropospheric NO<sub>2</sub> and HCHO using regional data from aircraft campaigns in the southeastern United States, *J. Geophys. Res.*, 109, D24307, doi:10.1029/2004JD004869, 2004. [11471](#)

30 Meyer-Arneke, J., Ladstatter-Weissenmayer, A., Richter, A., Wittrock, F., and Burrows, J. P.: A study of the trace gas columns of O<sub>3</sub>, NO<sub>2</sub>, and HCHO over Africa in September 1997, *Faraday Discuss.*, 130, 387–405, 2005. [11474](#)

Michel, A. B., Usher, C. R., and Grassian, V. H.: Heterogeneous and catalytic uptake of ozone

**Tropical tropospheric ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Tropical tropospheric  
ozone**

B. Sauvage et al.

on mineral oxides and dusts: A Knudsen cell investigation, *Geophys. Res. Lett.*, 29(14), 1665, doi:10.1029/2002GL014896, 2002.

Millet, D. B., Jacob, D. J., Turquety, S., Hudman, R. C., Wu, S., Fried, A., Walega, J., Heikes, B. G., Blake, D. R., Singh, H. B., Anderson, B. E., and Clarke, A. D.: Formaldehyde distribution over North America: Implications for satellite retrievals of formaldehyde columns and isoprene emission, *J. Geophys. Res.*, in press, 2006. [11471](#)

Moorthi, S. and Suarez, M. J.: Relaxed Arakawa-Schubert – A parameterization of moist convection for general-circulation models *M. Weath. Rev.*, 120(6), 978–1002, 1992. [11489](#)

Nesbitt, S. W., Zhang, R., and Orville, R. E.: Seasonal and Global  $NO_x$  production by lightning estimated from the Optical Transient Detector (OTD), *Tellus*, 52B, 1206–1215, 2000.

Nédélec, P., Cammas, J.-P., Thouret, V., Athier, G., Cousin, J.-M., Legrand, C., Abonne, C., Lecoeur, F., Cayez, G., and Marizy, C.: An improved infrared carbon monoxide analyser for routine measurements aboard commercial airbus aircraft: Technical validation and first scientific results of the MOZAIC III programme, *Atmos. Chem. Phys.*, 3, 1551–1564, 2003. [11469](#)

Palmer, P. I., Jacob, D. J., Chance, K., Martin, R. V., Spurr, R. J. D., Kurosu, T. P., Bey, I., Yantosca, R., Fiore, A., and Li, Q.: Air mass factor formulation for spectroscopic measurements from satellites: Application to formaldehyde retrievals from the Global Ozone Monitoring Experiment, *J. Geophys. Res.*, 106(D13), 14 539–14 550, 2001. [11470](#)

Palmer, P. I., Jacob, D. J., Fiore, A. M., Martin, R. V., Chance, K., and Kurosu, T. P.: Mapping isoprene emissions over North America using formaldehyde column observations from space, *J. Geophys. Res.*, 108(D6), 4180, doi:10.1029/2002JD002153, 2003. [11468](#), [11474](#), [11475](#)

Palmer, P. I., Abbot, D. S., Fu, T.-M., et al.: Quantifying the seasonal and interannual variability of North American isoprene emissions using satellite observations of the formaldehyde column, *J. Geophys. Res.*, 111, D12315, doi:10.1029/2005JD006689, 2006. [11487](#)

Park, R. J., Jacob, D. J., Field, B. D., Yantosca, R. M., and Chin, M.: Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the United States: implications for policy, *J. Geophys. Res.*, 109, D15204, 10.1029/2003JD004473, 2004.

Park, R. J., Jacob, D. J., Palmer, P. I., et al.: Export efficiency of black carbon aerosol in continental outflow: global implications, *J. Geophys. Res.*, 110, D11205, doi:10.1029/2004JD005432, 2005.

Pickering, K. E., Wang, Y. S., Tao, W. K., Price, C., and Muller, J. F.: Vertical distributions of

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Tropical tropospheric  
ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

lightning  $\text{NO}_x$  for use in regional and global chemical transport models, *J. Geophys. Res.*, 103, 31 203–31 216, 1998. [11473](#)

Price, C. and Rind, D.: A simple lightning parameterization for calculating global lightning distributions, *J. Geophys. Res.*, 97, 9919–9933, 1992. [11473](#)

5 Price, C., Penner, J., and Prather, M.:  $\text{NO}_x$  from lightning (1). Global distribution based on lightning physics, *J. Geophys. Res.*, 102, 5929–5941, 1997.

Rasch, P. J., Mahowald, N. M., and Eaton, B. E.: Representations of transport, convection, and the hydrologic cycle in chemical transport models: Implications for the modeling of short-lived and soluble species, *J. Geophys. Res.*, 102(D23), 28 127–28 138, 1997.

10 Richter, A. and Burrows, J. P.: Tropospheric  $\text{NO}_2$  from GOME measurements, *Adv. Space Res.*, 29(D23), 1673–1683, 2002.

Rodgers, C. D.: Inverse methods for atmospheric soundings: Theory and practice, World Sci., Hackensack, N. J., 2000. [11471](#)

15 Sauvage, B., Thouret, V., Cammas, J.-P., Gueusi, F., Athier, G., and Nédélec, P.: Tropospheric ozone over Equatorial Africa: regional aspects from the MOZAIC data, *Atmos. Chem. Phys.*, 5, 311–335, 2005. [11469](#), [11484](#)

Sauvage, B., Thouret, V., Thompson, A. M., Witte, J. C., Cammas, J.-P., Nédélec, P., and Athier, G.: Tropical Atlantic “Ozone Paradox” and “Zonal Wave-one” from the In-situ MOZAIC and SHADOZ Data, *J. Geophys. Res.*, 111, D01301, doi:10.1029/2005JD006241, 2006a.

20 Shim, C., Wang, Y. H., Choi, Y., Palmer, P., Abbot, D. S., and Chance, K.: Constraining global isoprene emissions with GOME formaldehyde column measurements, *J. Geophys. Res.*, 110, doi:10.1029/2004JD005629, 2005.

Schoeberl, M. R., Douglass, A. R., Hilsenrath, E., Bhartia, P. K., Barnett, J., Gille, J., Beer, R., Gunson, M., Waters, J., Levelt, P. F., and deCola, P.: Earth Observing System missions benefit atmospheric research, *EOS*, 85, 177–184, 2004.

25 Spurr, R. J. D.: Simultaneous derivation of intensities and weighting functions in a general pseudo-spherical discrete ordinate radiative transfer treatment, *J. Quant. Spectrosc. Radiat. Transfer*, 75, 129–175, 2002. [11470](#)

30 Stehr, J. W., Ball, W. P., Dickerson, R. R., Doddridge, B. G., Piety, C. A., and Johnson, J. E.: Latitudinal gradients in  $\text{O}_3$  and  $\text{CO}$  during INDOEX 1999, *J. Geophys. Res.*, 107(D19), 8016, doi:10.1029/2001JD000446, 2002.

Tabazadeh, A., Jacobson, M. Z., Singh, H. B., Toon, O. B., Lin, J. S., Chatfield, R. B., Thakur, A. N., Talbot, R. W., and Dibb, J. E.: Nitric acid scavenging by mineral and biomass burning

- aerosols, *Geophys. Res. Lett.*, 25, 4185–4188, 1998.
- Thakur, A. N., Singh, H. B., Mariani, P., et al.: Distribution of reactive nitrogen species in the remote free troposphere: data and model comparisons, *Atm. Env.*, 99, 1403–1422, 1999.
- Thompson, A. M.: The oxidizing capacity of the earth's atmosphere: Probable past and future changes, *Science*, 256, 1157–1165, 1992. [11467](#)
- Thompson, A. M., Doddridge, B. G., Witte, J. C., Hudson, R. D., Luke, W. T., Johnson, J. E., Johnson, B. J., Oltmans, S. J., and Weller, R.: A tropical Atlantic paradox: Shipboard and satellite views of a tropospheric ozone maximum and wave-one in January-February 1999, *Geophys. Res. Lett.*, 27, 3317–3320, 2000.
- Thompson, A. M., Witte, J. C., Mc Peters, R. D., et al.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2000 tropical ozone climatology 2. Tropospheric variability and the zonal wave-one, *J. Geophys. Res.*, 108(D2), 8238, doi:10.1029/2001JD000967, 2003b. [11469](#)
- Thompson, A. M., Witte, J. C., Oltmans, S. J., et al.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2000 tropical ozone climatology 1. Comparison with Total Ozone Mapping Spectrometer (TOMS) and ground-based measurements, *J. Geophys. Res.*, 108(D2), 8241, doi:10.1029/2002JD002241, 2003a. [11469](#)
- Thouret, V., Cammas, J.-P., Sauvage, B., Athier, G., Zbinden, R., Nédélec, P., Simon, P., and Karcher, F.: Southern Hemisphere Tropopause referenced ozone climatology and inter-annual variability (1994–2003) from the MOZAIC programme, *Atmos. Chem. Phys.*, 6, 1033–1051, 2006.
- Thornton, J. and Abbatt, J. P. D.: Measurements of HO<sub>2</sub> uptake to aqueous aerosol: Mass accommodation coefficients and net reactive loss, *J. Geophys. Res.*, 110(D8), D08309, doi:10.1029/2004JD005402, 2005. [11477](#)
- Umann, B., Arnold, F., Schaal, C., Hanke, M., Uecker, J., Aufmhoff, H., Balkanski, Y., and Van Dingenen, R.: Interaction of mineral dust with gas phase nitric acid and sulfur dioxide during the MINATROC II field campaign: First estimate of the uptake coefficient  $\gamma$  HNO<sub>3</sub> from atmospheric data, *J. Geophys. Res.*, 110, D22306, doi:10.1029/2005JD005906, 2005.
- Underwood, G. M., Song, C. H., Phadnis, M., Carmichael, G. R., and Grassian, V. H.: Heterogeneous reactions of NO<sub>2</sub> and HNO<sub>3</sub> on oxides and mineral dust: A combined laboratory and modeling study, *J. Geophys. Res.*, 106(D16), 18055–18066, 2005.
- van Noije, T. P. C., Eskes, H. J., Dentener, F. J., et al.: Multi-model ensemble simulations of tropospheric NO<sub>2</sub> compared with GOME retrievals for the year 2000, *Atmos. Chem. Phys.*, 6

**Tropical tropospheric ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2943–2979, 2006.

Wang, Y., Jacob, D. J., and Logan, J. A.: Global simulation of tropospheric O<sub>3</sub>-NO<sub>x</sub>-hydrocarbon chemistry, 1 – Model formulation, *J. Geophys. Res.*, 103, 10 713–10 726, 1998. [11473](#)

Wang, K. Y. and Shallcross, D. E.: Modelling terrestrial biogenic isoprene fluxes and their potential impact on global chemical species using a coupled LSM-CTM model, *Atmos. Env.*, 34, 2909–2925, 2000.

Wu, S., Mickley, L. J., Jacob, D. J., Logan, J. A., and Yantosca, R. M.: Why are there large differences between models in global budgets of tropospheric ozone?, *J. Geophys. Res.*, accepted, 2006. [11489](#)

Yamasoe, M. A., Artaxo, P., Miguel, A. H., and Allen, A. G.: Chemical composition of aerosols particles from direct emissions of vegetation fires in the Amazon Basin: water-soluble species and trace elements *Atmos. Env.*, 34(10), 1641–1653, 2000. [11477](#)

Yienger, J. J. and Levy II, H.: Empirical model of global soil-biogenic NO<sub>x</sub> emissions, *J. Geophys. Res.*, 100(D6), 11 447–11 464, 1995. [11473](#)

Zhang, G. J. and McFarlane, N. A.: Sensitivity of climate simulations to the parameterization of cumulus in the Canadian Climate Centre General Circulation Model, *Atmosphere-Ocean*, 33, 407–446, 1995. [11489](#)

ACPD

6, 11465–11520, 2006

## Tropical tropospheric ozone

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU



**Table 1.** Characteristics of the MOZAIC and SHADOZ sites. Abbreviations are given in parenthesis. Number of CO measurements are given in parenthesis where available.

Site	Lon/Lat	Total number of O <sub>3</sub> and RH profiles	Region
Caracas (CAR)	67.0° W/10.5° N	651	northern South America
Cayenne (Cay)	52.3° W/4.9° N	175	northern South America
Bogota (Bog)	74.0° W/4.5° N	220	northern South America
San Cristobal (San C)	89.6° W/0.9° S	256	northern South America
Paramaribo (Par)	55.2° W/5.8° N	230	northern South America
Rio de Janeiro (Rio)	43.2° W/22.8° S	551	South America
Sao Paolo (SAO)	46.6° W/23.5° S	979	South America
Dakar (Dak)	17.4° W/14.5° N	89	north Africa
Lagos (LAG)	3.3° E/6.5° N	354 (139)	West Africa
Abidjan (Abi)	4.0° W/5.4° N	178	West Africa
Douala (Dou)	9.7° E/4.0° N	185	West Africa
Brazzaville (BRA)	15.3° E/4.2° S	114	Central Africa
Luanda (Lua)	13.2° E/8.5° S	48	Central Africa
Windhoek (Win)	17.4° E/22.4° S	138	South Africa
Johannesburg (Joh)	28.0° E/26.2° S	574	South Africa
Nairobi (Nai)	36.7° E/1.1° S	116	East Africa
Abu Dhabi (Abu)	54.6° E/24.4° N	215	Middle East
Dubai (DUB)	55.3° E/25.2° N	559 (89)	Middle East
Bombay (BOM)	72.8° E/19.0° N	145	India
Delhi (DEL)	77.3° E/28.5° N	678 (274)	India
Madras (Mad)	80.0° E/13.0° N	246	India
Bangkok (BAN)	100.5° E/13.9° N	659	Thailand
Natal (Nat)	35.3° W/5.4° S	253	Atlantic
Ascension (ASC)	14.4° W/7.9° S	305	Atlantic
Reunion Island (REU)	55.4° E/21.0° S	146	Indian Ocean
Kuala Lumpur (Kua)	112.6° E/-7.5° S	160	Pacific
Fiji (Fij)	178° E/18.0° S	229	Pacific
Samoa (SAM)	170.5° W/14.2° S	263	Pacific

**Tropical tropospheric ozone**

B. Sauvage et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Tropical tropospheric ozone

B. Sauvage et al.

**Table 2.** Annual  $\text{NO}_x$  emissions in the GEOS-Chem simulations for the year 2000. The tropical emissions are over  $20^\circ \text{S}$ – $20^\circ \text{N}$ .

Source	Original, Global/Tropics (Tg N/yr)	Standard, Global/Tropics (Tg N/yr)
Biomass Burning	5.9/4.0	7.0/5.0
Lightning	5.0/3.3	5.0/3.3
Soils	6.0/2.3	8.9/3.1
Anthropogenic	23.9/2.1	23.9/2.1
Biofuels	2.2/0.7	2.2/0.7
Aircraft	0.5/0.1	0.5/0.1

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Tropical tropospheric  
ozone

B. Sauvage et al.

**Table 3.** Seasonal tropospheric O<sub>3</sub> column (DU) from GEOS-Chem standard simulation (difference with original simulation is given in parenthesis)/MOZAIC or SHADOZ (standard deviation 1 $\sigma$  is in parenthesis)/and GOME. For MOZAIC we complete the column between the aircraft ceiling of 185 hPa and the tropopause with a fixed ozone mixing ratio of 70 ppbv.

GEOS-Chem/MOZAIC-SHADOZ/GOME TOC								
Regions	DJF		MAM		JJA	SON		
Caracas	26.6 (−4.2)/	22.2 (3.2) / 27	29.6 (−3.1)/	28.8 (4.5)/ 26.9	27.6 (−1.8)/	26.3 (4) / 27.9	25.3 (−2.5)/	25.2 (3.8)/ 25.8
Sao Paolo	32.7 (−0.5)/	29.4 (4.6)/ 35.4	29.4 (−4.1)/	24.7 (3.2)/ 32	30.1 (−0.2)/	29.4 (4.1)/ 29.5	37.4 (−0.6)/	34.7(5)/ 39
Dubai	40 (+0.1)/	38.8 (3.6)/ 34	43.9 (−1.7)/	41.6 (4.5)/ 44.5	50.1 (−2.7)/	45.2 (4.7)/ 43.5	40.9 (−2.2)/	36 (3.6)/ 35
Samoa	17.4 (+1.8)/	18.4 (3.2)/ 16.4	19 (+0.9)/	17.9 (5)/ 19.5	20.4 (+1)/	20.2 (4)/ 24.4	22.1 (+3.2)/	23 (5)/ 23
Reunion	32.6 (+2.2)/	32.4 (4.5)/ 30	31 (+1)/	32(6)/ 31.5	34.1 (+0.4)/	34.7 (5)/ 33	43.4 (+1.1)/	45.2 (5)/ 45
Ascension	36 (+3.2) /	35.4 (5)/ 32.8	31 (−1)/	30.9 (5)/ 30.5	38.2 (−0.6)/	40.6 (6.2)/ 34.5	40.7 (−0.9)/	44 (6.4)/ 37.7
Natal	34.3 (+2.5) /	34.2 (4.5)/ 33.5	27.2 (−1.2)/	26 (5)/ 29.4	33.8 (−0.4)/	36.2 (4.2)/ 32.73	37.3 (−0.7)/	41 (6)/ 37.4
Lagos	40.1 (+5.9)/	40.5 (3.9)/ 34	36 (+2.7)/	37.5 (2.2)/ 32.5	32.5 (+2.4)/	33 (3.2)/ 31	32 (−0.1)/	32.5 (2.6)/ 29.5
Brazzaville	34.5 (+4.8)/	36.7 (3.3)/ 29.5	31.5 (+2)/	32.5 (3.4)/ 29.4	48.8 (+6.3)/	49 (3.6)/ 37.5	40.3 (+1)/	43(2.8)/ 34.7
Windhoek	32.8 (+0.9)/	31.5 (2.6)/ 32	29.3 (−0.1)/	28.7 (3.1)/ 29.7	33.3 (+0)/	32.4 (3.2)/ 33.1	41.2 (+1.2)/	42.1 (3.4)/ 42.1
Bombay	40.7 (+0.4)/	40.1 (5)/ 33.3	41.6 (−2.5)/	40.3 (4) / 35.5	34.9 (−2.1)/	30.3 (5.4)/ 28.6	36.1 (+1.1)/	35.6 (4.1)/ 31.7
Average	31.8/	32.7(3.9)/30.7	31.7/	30.9(4.1)/31	34.8/	34.3(4.3)/32.3	36.0/	36.5(4.3)/33.9

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Tropical tropospheric  
ozone

B. Sauvage et al.



**Fig. 1.** MOZAIC (blue) and SHADOZ (black) sites used in this study. Capital letters refer to sites that represent the rectangular region. Abbreviations are defined in Table 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Tropical tropospheric  
ozone

B. Sauvage et al.




**Fig. 2.** Seasonal biomass burning emissions ( $10^9 \text{ molec N cm}^{-2} \text{ s}^{-1}$ ) for December-February (DJF) and June-August (JJA). The left panels represent emissions used in the original simulation. The right panels represent top-down emissions determined from GOME observations of tropospheric  $\text{NO}_2$  columns.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Tropical tropospheric  
ozone

B. Sauvage et al.




**Fig. 3.** Seasonal average lightning emissions ( $10^9$  molec  $\text{N}/\text{cm}^2/\text{s}$ ). The left panels show lightning emissions calculated from GEOS dynamics in the original version. The right panels show lightning emissions scaled to OTD/LIS measurements of flash rates as used in the standard (improved) simulation.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Tropical tropospheric  
ozone

B. Sauvage et al.



**Fig. 4.** Seasonal average GOME (left) and (middle) GEOS-Chem (convolved with GOME averaging kernels) tropospheric ozone columns for 2000. The right column represent difference between standard and original simulation of tropospheric ozone columns (TOC). The arrows in the middle column represent the horizontal ozone flux integrated from the surface to the tropopause.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Tropical tropospheric ozone**

B. Sauvage et al.





Fig. 5.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Tropical tropospheric  
ozone**

B. Sauvage et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Tropical tropospheric  
ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion

EGU

**Fig. 5.** Seasonal vertical profiles of  $O_3$  in ppbv. The plain black line indicates MOZAIC and SHADOZ measurements of  $O_3$ . Horizontal bars represent one standard deviation of measurements.  $O_3$  simulations are in blue (original) and red (standard). The solid green line indicates a simulation with enhanced intracloud  $NO_x$  emissions. The dashed green lines show simulation with L- $NO_x$  emissions of 3 TgN/yr and 7 TgN/yr.

Tropical tropospheric  
ozone

B. Sauvage et al.




**Fig. 6.** Seasonal averaged tropospheric  $\text{NO}_2$  columns ( $10^{15} \text{ molec cm}^{-2}$ ) during the year 2000. The left panels are for GOME, the middle for GEOS-Chem standard and the right for GEOS-Chem original. White areas indicate regions with persistent clouds.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Tropical tropospheric  
ozone

B. Sauvage et al.




**Fig. 7.** Seasonal averaged tropospheric HCHO columns ( $10^{16}$  molec  $\text{cm}^{-2}$ ). The left panels are for GOME, the middle are for GEOS-Chem standard and the right for GEOS-Chem original. White areas represent persistent clouds.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Tropical tropospheric  
ozone

B. Sauvage et al.




**Fig. 8.** Seasonal vertical profiles of  $O_3$  in ppbv. The plain black line indicates in situ  $O_3$ . Horizontal bars represent one standard deviation of measurements.  $O_3$  simulations are in blue (original), red (standard), and in green for in-situ-based on  $NO_x$  emission factors.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Tropical tropospheric  
ozone

B. Sauvage et al.



**Fig. 9.** Seasonal vertical profiles of relative humidity (RH) and CO. Black lines are for MOZIC RH, dashed-lines are for CO. GEOS-4 simulations are in solid lines, GEOS-3 in dashed lines, with blue for RH and red for CO.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Tropical tropospheric  
ozone

B. Sauvage et al.



Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Tropical tropospheric  
ozone**

B. Sauvage et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion

**Fig. 10.** Vertical profiles of  $O_3$  at MOZAIC and SHADOZ sites that exhibit a large sensitivity to either dynamics and heterogeneous chemistry. The red line is for the standard simulation (GEOS-4), the blue line for GEOS-3, and the green line for  $HNO_3$  uptake turned off.



Tropical tropospheric  
ozone

B. Sauvage et al.



**Fig. 11.** Meridional average (5° W–30° E) of CO (top left), RH (top right) and O<sub>3</sub> (bottom) at flight altitude (200–300 hPa) for MOZAIC (black line, squares), GEOS-4 standard (red line, circle) and GEOS-3 standard (blue line, triangle) simulations during JJA.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)