

Measurements and modelling of I2, IO, OIO, BrO and NO3 in the mid-latitude marine boundary layer

A. Saiz-Lopez, J. A. Shillito, H. Coe, J. M. C. Plane

▶ To cite this version:

A. Saiz-Lopez, J. A. Shillito, H. Coe, J. M. C. Plane. Measurements and modelling of I2, IO, OIO, BrO and NO3 in the mid-latitude marine boundary layer. Atmospheric Chemistry and Physics Discussions, 2005, 5 (5), pp.9731-9767. hal-00303944

HAL Id: hal-00303944 https://hal.science/hal-00303944

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Atmos. Chem. Phys. Discuss., 5, 9731–9767, 2005 www.atmos-chem-phys.org/acpd/5/9731/ SRef-ID: 1680-7375/acpd/2005-5-9731 European Geosciences Union



ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.

Measurements and modelling of I_2 , IO, OIO, BrO and NO₃ in the mid-latitude marine boundary layer

A. Saiz-Lopez¹, J. A. Shillito², H. Coe², and J. M. C. Plane¹

¹School of Environmental Sciences, University of East Anglia, Norwich, UK ²School of Earth, Atmospheric and Environmental Sciences, University of Manchester, Manchester, UK

Received: 26 August 2005 - Accepted: 4 September 2005 - Published: 10 October 2005

Correspondence to: J. M. C. Plane (j.plane@uea.ac.uk)

 $\ensuremath{\textcircled{O}}$ 2005 Author(s). This work is licensed under a Creative Commons License.

Title Page		
Abstract	Introduction	
Conclusions	References	
Tables	Figures	
- 19		
•	•	
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		

Abstract

Time series observations of molecular iodine (I₂), iodine oxides (IO, OIO), bromine oxide (BrO), and the nitrate radical (NO₃) in the mid-latitude coastal marine boundary layer (MBL) are reported. Measurements were made using a new long-path DOAS instrument during a summertime campaign at Mace Head on the west coast of Ireland. I₂ was detected using the $B^3\Pi(0^+_u) - X^1\Sigma_g^x$ electronic transition between 535 and 575 nm. The I₂ mixing ratio was found to vary from below the detection limit (~5 ppt) up to a nighttime maximum of 93 ppt. Along with I₂, observations of IO, OIO and NO₃ were also made during the night. Surprisingly, IO and OIO were detected at mixing ratios up to 2.5 and 10.8 ppt, respectively. A model is employed to show that the reaction between I₂ and NO₃ is the likely nighttime source of these radicals. The BrO mixing ratio varied from below the detection limit at night (~1 ppt) to a maximum of 6 ppt in the first hours after sunrise. A model shows that this diurnal behaviour can be explained by halogen recyling in sea-salt aerosol building up photolabile precursors of atomic Br

¹⁵ during the preceding night. In the same campaign a zenith sky DOAS was employed to determine the column density variation of NO_3 as a function of solar zenith angle (SZA) during sunrise, from which vertical profiles of NO_3 through the troposphere were obtained. On several occasions a positive gradient of NO_3 was observed over the first 2 km, possibly due to dimethyl sulphide (DMS) removing NO_3 at the ocean surface.

20 1. Introduction

Interest in atmospheric halogen chemistry has grown enormously in the past three decades, originally because of its active role in the destruction of stratospheric (Molina and Rowland, 1974; Wofsy et al., 1975; Farman et al., 1985) and polar tropospheric (Barrie et al., 1988; Finlayson-Pitts et al., 1990; McConnell et al., 1992; Hausmann and Platt, 1994; Hoppinger and Platt, 2002) ozone. However, halogen radicals could play a

Platt, 1994; Honninger and Platt, 2002) ozone. However, halogen radicals could play a key role in a number of other important tropospheric processes, including NO_x (NO and 5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment



 NO_2) and HO_x (OH and HO_2) chemistry, the oxidation of a range of organic molecules, and in the formation of ultra-fine particles. These impacts are discussed in greater detail below. In the mid-latitude MBL, BrO has recently been detected both by multi-axis DOAS (Leser et al., 2003) and long-path DOAS (Saiz-Lopez et al., 2004a). The source

- ⁵ of bromine is almost certainly the release of species such as Br₂, BrCl and IBr from sea-salt aerosol, following the uptake from the gas phase, and subsequent aqueous-phase reactions, of hypohalous acids (HOX, where X=Br, I) (Vogt et al., 1996). In semi-polluted areas with higher NO_x, the uptake of di-nitrogen pentoxide (N₂O₅), formed at night by the recombination of the NO₃ with NO₂, also leads to the release of BrNO₂
- ¹⁰ (Behnke et al., 1994). These heterogeneous mechanisms are supported by the observation that the bromide ions in sea-salt aerosols are substantially depleted (Ayers et al., 1999; Gabriel et al., 2002). Once released into the gas phase, these bromine-containing compounds will photolyse rapidly during the day, and the resulting Br atoms will react fast with O_3 to form BrO.
- Recent measurements of IO and OIO (Alicke et al., 1999; Allan et al., 2000a, 2001; Saiz-Lopez and Plane, 2004) have confirmed the importance of iodine chemistry in the MBL. The source of atmospheric iodine is the evasion of biogenic iodocarbons such as CH₃I, CH₂I₂, CH₂IBr and CH₂ICI from both the open ocean and coastal areas (Carpenter et al., 1999). Additionally, the detection of I₂ during NAMBLEX indicates that this
 molecule is probably the major source of atmospheric iodine in certain coastal loca-
- tions (Saiz-Lopez and Plane, 2004). Also, I_2 would account for the surprising nighttime observations of IO and OIO, since the rate constant between NO₃ and I_2 to yield INO₃ and I is relatively fast (Chambers et al., 1992). The atomic I produced in this way would rapidly form IO by reaction with O₃. A number of studies have proposed iodine as a
- ²⁵ significant catalyst in tropospheric O₃ depletion (Jenkin et al., 1985; Davis et al., 1996; McFiggans et al., 2000). BrO and IO also compete effectively with the OH radical in the oxidation of DMS (Saiz-Lopez et al., 2004a; Glasow and Crutzen, 2004), and a very recent study has shown that the HO₂/OH ratio in the coastal MBL is significantly altered by IO (Bloss et al., 2005). Another potentially significant aspect of the iodine

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment



chemistry is the condensation of iodine oxide vapours, leading to the formation of new ultra-fine particles in the daytime marine atmosphere (O'Dowd et al., 2002; McFiggans et al., 2004; Saiz-Lopez et al., 2005).

Field observations of the NO₃ radical over the past 25 years have demonstrated that

⁵ it plays a major role in the chemistry of the nighttime troposphere (Platt et al., 1980, 1990; Heintz et al., 1996; Allan et al., 2000b, 2002a; Brown et al., 2003, 2004). NO₃ reacts rapidly with DMS in the MBL (Butkovskaya and LeBras, 1994). The radical also recombines with NO₂ to establish an equilibrium with N₂O₅. The subsequent heterogeneous uptake of N₂O₅ on aqueous sea-salt aerosol both removes NO_x and activates halogens (see above).

In this paper we report results from a 5 week campaign of boundary layer observations of I_2 , IO, OIO, BrO and NO₃ made by long-path differential optical absorption spectroscopy (DOAS), and measurements of the vertical concentration profiles of NO₃ by zenith sky DOAS. Box models are then used to investigate the nighttime chemistry of iodine, and the diurnal behaviour of BrO.

2. Experimental

15

2.1. Long-path DOAS instrument

The DOAS measurements were performed at the Mace Head Atmospheric Research Station (MHARS) on the west coast of Ireland (53°20′ N, 9°54′ W), during the NAM-

- BLEX (North Atlantic Marine Boundary Layer EXperiment) campaign in July/August 2002. The long-path DOAS instrument was housed in an observatory on the foreshore at the MHARS. It consists of a Newtonian telescope containing the xenon arc lamp source, and the transmitting and receiving optics. The light beam was folded back to the transmitter by a retro-reflector situated 4.2 km away on Croaghnakeela Island (see
- Fig. 2 in Saiz-Lopez et al., 2005), providing an optical absorption path of 8.4 km, about 5–20 m above sea level. Differential absorption spectra were collected by averaging

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.



EGU

for 30 min when measuring I₂, IO, OIO and BrO, or 20 min when detecting NO₃. More details on the instrument design and spectral deconvolution procedures are provided elsewhere (Plane and Saiz-Lopez, 2005). The following reference spectra were taken from the literature and adapted to the spectrometer instrument function: I₂ (Saiz-Lopez et al., 2004b), IO (Harwood et al., 1997), OIO (P. Spietz, University of Bremen, personal communication, 2005), BrO (Wahner et al., 1988) and NO₃ (Yokelson et al., 1994).

2.2. Zenith sky DOAS instrument

This instrument consists of a 0.8° field-of-view telescope, fixed in a zenith-viewing geometry and coupled to a spectrometer by means of a fibre optic cable. Spectra were recorded with a resolution of 0.5 nm in the $\lambda = 640-700 \text{ nm}$ region for detection of NO₃. A detailed description of the instrument is given in (Allan et al., 2002a). Spectra were acquired every 1–2 min (depending on scattered light levels) for approximately 2 h through sunrise (SZA 95°). The NO₃ column abundances were obtained using the same de-convolution method as for the long-path DOAS measurements, with a reference spectrum recorded after sunrise when the NO₃ mixing ratio is negligible due to the rapid photodissociation of the radical (Allan et al., 2002a). The vertical concentration profile was then determined using an optimal estimation algorithm described below.

3. Results and discussion

- 3.1. Long-path DOAS measurements of NO₃
- Observations of NO₃ were performed during 16 nights under different meteorological conditions. Figure 1 summarises the NO₃ time series set during the campaign. The maximum value reached during the observation period was 25 ppt, with averages of ~3 and 13 ppt for clean marine and semi-polluted air masses. These values are consistent with previously reported observations at the same location, under a similar range

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

Title	Title Page		
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
١٩	►I		
•	►		
Back	Close		
Full Sci	Full Screen / Esc		
Print	Print Version		
Interactive	Interactive Discussion		

of meteorological conditions (Allan et al., 2000b). Comparable nighttime mixing ratio profiles have been reported in other marine areas (Heintz et al., 1996; Allan et al., 1999; Brown et al., 2004; Vrekoussis et al., 2004). The significant variations in NO_3 levels during the period of measurements can be explained by considering the back

- ⁵ trajectories of the air masses arriving in the MBL at Mace Head. Figures 2 and 3 show two different types of air masses and the associated NO₃ mixing ratios. The solid and broken grey bars correspond to sunset and sunrise, respectively. Typically, NO₃ mixing ratios below 5 ppt were associated with air masses from the Atlantic Ocean sector, Fig. 2a and b. By contrast, NO₃ mixing ratios above 10 ppt coincided with semi-polluted
- ¹⁰ air masses from the UK or the European continent Fig. 3a and b. Note that although the predicted back trajectories on 3 September have their origin in the north Atlantic, the air mass has traversed over parts of the United Kingdom and the European continent before reaching Mace Head.

NO₃ is formed by the oxidation of NO₂ by O₃, with a rate constant $k(NO_2 + O_3) = 1.2 \times 10^{-13} \exp(-2450/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Sander et al., 2003). In the immediate postdusk hours, once photolysis ceases and before NO₃ builds up to a chemical steady state, assuming other losses are negligible, the rate of increase of NO₃ should just be the rate of production, $R_{NO_3} = k(NO_2 + O_3)$ [O₃][NO₂]. Taking the observed average NO₂ and O₃ mixing ratios of 450 ppt and 22 ppb, respectively, and an average temperature of 288 K, then $R_{NO_3} = 22 \text{ ppt h}^{-1}$ is obtained. This rate is in good accord with the observations (see Fig. 3a). We have shown previously that under clean marine air conditions at Mace Head during summer, DMS controls the removal of NO₃, whereas in semi-polluted air masses NO₃ removal is dominated by the indirect loss of N₂O₅ (Allan et al., 2000b).

²⁵ During NAMBLEX the broad-band cavity ring-down spectroscopy (BBCRDS) technique was also employed to make in-situ measurements of the NO₃ radical. An intercomparison of the local and long-path observations can be found in Bitter et al. (2005)¹.

¹Bitter, M., Ball, S. M., Povey, I. M., Jones, R. L., Saiz-Lopez, A., and Plane, J. M. C.:

ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment



A photochemical box model of the nighttime NO₃ chemistry (Allan et al., 2000b) was used to simulate the production of N₂O₅ from its equilibrium with NO₃. The model was run under typical westerly clean marine conditions, for which an average NO₃ mixing ratio of 3 ppt was observed. For marine air masses, the N₂O₅ accommodation coefficient in the model, γ N₂O₅, was set at 0.03 (Behnke et al., 1997; Allan et al., 2000b). Under these clean marine conditions, with an aerosol volumetric surface area of 10⁻⁷ cm² cm⁻³, the modelled N₂O₅ mixing ratio throughout the night is ~3 ppt. This N₂O₅ mixing ratio will be used in a bromine chemistry model to assess the contribution of NO₃ chemistry to halogen activation on sea-salt particles (Sect. 3.6).

¹⁰ 3.2. Zenith sky observations of NO₃ vertical profiles

25

NO₃ vertical column measurements were carried out at sunrise during six days, as shown in Fig. 4. The left-hand panels (Fig. 4a) show the expected dependence of NO₃ column on SZA: as the solar terminator sweeps down through the atmosphere, more and more of the overlying NO₃ is removed by photolysis. Note that the errors associated with individual column density measurements are quite variable. These uncertainties arise mainly from the visibility conditions, and the temperature dependence of the NO₃ absorption cross-section (Allan et al., 2002a). The literature cross-section employed is that of Yokelson et al. (1994) at 280 K, a temperature which can be considered representative of the lower troposphere where the contribution to the NO₃ total absorption column should be strongest. In Fig. 4a it can be seen that the NO₃ persisted after dawn on several occasions. This effect is most likely due to the thermal

- decomposition of N₂O₅ competing with photolysis to sustain the NO₃ mixing ratio past sunrise (Smith and Solomon, 1990; Smith et al., 1993; Allan et al., 2002a; Coe et al., 2002).
 - Measurements of NO_3 , N_2O_5 , OIO, I_2 , water vapour and aerosol optical depth by broadband cavity ringdown spectroscopy during the NAMBLEX campaign, Atmos. Chem. Phys. Discuss., in preparation, 2005.

The conversion from column abundance to vertical mixing ratio profiles was made

ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment



using the optimal estimation method (OEM) (Rodgers, 1976, 1990). Briefly, the forward model used is described by the expression $\mathbf{y} = \mathbf{K}\mathbf{x}$, where knowing the state of the system or weighing matrix function (**K**) and the vertical mixing ratio profile (**x**), the column density time series (**y**) can be predicted. However, since **x** is not known the forward model is inverted to give the backward model $\mathbf{x} = \mathbf{K}^{-1}\mathbf{y}$. The inversion process makes use of 'a priori' information, i.e. the likely NO₃ vertical profile, to allow NO₃ vertical distribution information to be retrieved. **K** comprises photolysis of NO₃ as a function of height, SZA and time (Coe et al., 2002).

In Fig. 4a the predicted column abundance (solid red line) using the forward model is compared with the measured, as a function of SZA. Figure 4b illustrates the corresponding retrieved NO_3 vertical concentration profiles. These profiles exhibit strong gradients, consistent with other observations (Allan et al., 2002a). On all six days, the highest NO_3 concentrations are in the mid- to low troposphere. There are probably three reasons for this. First, higher temperatures and NO_2 concentrations will cause

- ¹⁵ NO₃ to be produced more rapidly. Second, the thermal decomposition of N_2O_5 back to NO₃ will be faster. Third, DMS evading from the ocean will remove NO₃ at the base of the MBL, creating a positive gradient of NO₃ in the MBL. This probably explains the observations in clean marine conditions on 3, 7 and 10 August. On the night of 2 August (when the site was influenced by semi-polluted continental air masses) long-path
- DOAS and zenith sky measurements of NO₃ were performed simultaneously. In Fig. 4b (top panel) the boundary layer measurement of ~10 ppt just before dawn is comparable to the 7 ppt retrieved for the lowest box in the vertical profile, which is centred at 2 km. On 9 and 11 August, back trajectories show boundary layer air arriving after traversing north-south along the west coast of Ireland, which probably accounts for the relatively high NO₃ concentrations in the lowest box.
- $_{25}$ nign NO₃ concentrations in the lowest bo
 - 3.3. lodine species (I₂, IO, OIO)

5

Figure 5 illustrates the time series of I_2 , IO and OIO during NAMBLEX. The broken line in each of the panels shows the average DOAS detection limit for the particular

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment



species during the campaign. Measurements of I₂ were carried out during 18 days. As we have demonstrated in an accompanying paper (Saiz-Lopez et al., 2005), the photolysis lifetime of I_2 is so short (~15 s) that during daytime I_2 is very inhomogeneously distributed along the DOAS light path. Thus the daytime mixing ratios of I_2 (and IO)

5 in Fig. 5 should be treated with caution – they are useful as relative rather than absolute measurements. The maximum I₂ mixing ratio observed during the campaign was 93 ppt. This was observed at night, when the average I_2 mixing ratio was considerably higher, as expected in the absence of photolysis.

Figure 5 shows that the I₂ mixing ratio is characterised by sharp peaks lasting for

- 30 to 45 min. These peaks always coincided with low tide (Saiz-Lopez and Plane, 10 2004). Indeed, a simple plot of I₂ mixing ratio versus tidal height exhibits a clear anticorrelation, as shown in Fig. 6. The source of I₂ at this coastal environment is almost certainly macro-algae (e.g. Laminaria), which accumulate iodine compounds well over the concentration of such chemical species in seawater, leading to direct injection of
- I₂ when the plant is exposed to the atmosphere at low tide (Truesdale et al., 1995; 15 Kupper et al., 1998; McFiggans et al., 2004). The role of I_2 in the production of new particles in coastal areas has been recently proposed by Saiz-Lopez and Plane (2004) and McFiggans et al. (2004), and explored in detail using an iodine chemistry model (Saiz-Lopez et al., 2005). These studies show that I_2 is the major contributor to the particle bursts observed at Mace Head.

20

In contrast to the state of the tide, the I_2 mixing ratio did not correlate significantly with local wind direction. This is not surprising since the prevailing wind directions tended to be westerly or easterly, and macroalgae were exposed at both ends of the light path (see the map in Saiz-Lopez et al., 2005). Nevertheless, one potentially interesting detail is that I₂ was observed at night in westerly winds, irrespective of the state of 25 the tide. This may indicate that I_2 is also produced over the open ocean, as originally proposed by Garland and Curtis (1981).

Figure 5 also shows the complete time series of IO during NAMBLEX, totalling 15 days of measurements. The maximum IO mixing ratio during the campaign was \sim 7 ppt.

ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment



measured during daytime. This is in accord with previously reported IO observations at Mace Head (Alicke et al., 1999; Allan et al., 2000a). Note that this mixing ratio is probably the average of a rather inhomogeneous distribution along the DOAS path (Saiz-Lopez et al., 2005). During daytime, IO exhibited a clear anti-correlation with tidal
⁵ height and solar irradiation, as shown in the mesh plot in Fig. 7. This is good evidence for its photochemical production from coastal emissions of I₂ (Saiz-Lopez and Plane, 2004). IO was only measured during two nights of NAMBLEX, when mixing ratios up to ~2.5 ppt were observed.

OIO measurements were made simultaneously with I₂ in the 535–575 nm wavelength region. During the day, OIO was not observed above the daytime detection limit of ~4 ppt. The radical was only observed at night, with a maximum mixing ratio of 10.8 ppt. In fact, on this occasion the OIO peaked after local midnight, and about 2 h after a night-time peak of I₂ at low tide (Saiz-Lopez and Plane, 2004). The observation that OIO is only detectable at night is consistent with previous measurements made by us at Cape Grim (Tasmania), another remote marine location (Allan et al., 2001).

There appear to be two possible explanations for the absence of OIO during daytime. First, it could photolyse following absorption in the strong bands between about 470 and 610 nm. Whilst certain experimental evidence led us to conclude this was the case (Ashworth et al., 2002), we have recently shown that absorption leads to interconversion into the highly vibrationally-excited ground state rather than dissociation to I+O₂, so the quantum yield for photolysis is probably much less than 10% at 562 nm (D. M. Joseph, S. H. Ashworth, and J. M. C. Plane, University of East Anglia, personal communication). Nevertheless, a small probability of photolysis, integrated over the absorption bands, could still account for the removal of OIO during daytime (Saiz-Lopez

et al., 2005). A second possibility is that OIO is removed during daytime by reaction with radicals that have marked diurnal cycles with daytime maxima, such as OH, NO and IO. Based on our current experimental and theoretical work, we believe that the most likely of these is IO, as discussed in our recent modelling study (Saiz-Lopez et al., 2005).

ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment



3.4. Nighttime iodine chemistry

In order to understand the source of IO and OIO during the night in the MBL, we have modified our existing iodine chemistry model (McFiggans et al., 2000; Saiz-Lopez et al., 2005). A plausible mechanism for the formation of atomic iodine at night is the reaction of I₂+NO₃→INO₃+I, which has a rate constant at 295K of 1.5×10⁻¹² cm³ molecule⁻¹ s⁻¹ (Chambers et al., 1992). The resulting I atom would form IO in the atmosphere, and a simple calculation showed that this reaction might be fast enough to account for the nighttime observations of IO (Saiz-Lopez and Plane, 2004). The other product is INO₃. Assuming an accommodation coefficient, γINO₃, of
 0.05 and a clean MBL aerosol volumetric surface area of 10⁻⁷ cm² cm⁻³, the loss rate

- due to uptake on aerosols is $\sim 2.3 \times 10^{-5} \text{ s}^{-1}$. An upper limit to the rate of thermal dissociation at 285 K has been determined to be $4.5 \times 10^{-4} \text{ s}^{-1}$ (Allan and Plane, 2002b). Hence, at night in the MBL there will be a competition between uptake on aerosols and thermal decomposition of INO₃.
- ¹⁵ We also consider in the model the thermal stabilities of I₂O₂ and I₂O₄, dimers that form from the self reactions of IO and OIO, respectively. Ab initio quantum calculations, using a level of theory described previously (Allan and Plane, 2002b), show that the binding energy of the OIO dimer is only 72 kJ mol⁻¹, while that of the IO dimer (in the form IOIO) is about 86 kJ mol⁻¹. Using these binding energies, vibrational frequen-²⁰ cies and rotational constants as input for a Rice-Ramsberger-Kassel-Markus (RRKM)
- ²⁰ Cles and rotational constants as input for a Rice-Ramsberger-Kassel-Markus (RRKM) calculation, we predict that the rates of dissociation of these dimer molecules will be fast at a typical mid-latitude MBL nighttime temperature of 285 K and a pressure of 1 atmosphere. I₂O₄ will dissociate back to OIO+OIO at a rate of ~7×10³ s⁻¹. IOIO will dissociate to OIO+I and IO+IO at rates of ~12 and 4 s⁻¹, respectively. The atmospheric implication of these results will be a delay in the rate of disappearance of IO and OIO in
- the nighttime MBL. These dissociation rates, and the rate constant for the association of the OIO dimer (predicted from RRKM theory to be $\sim 5 \times 10^{-11}$ cm³ molecule⁻¹ s⁻¹ at 1 atm and 285 K), are included in the model.

ACPD

5, 9731–9767, 2005

Measurements and modelling of trace gases in a coastal marine environment



When running the model, the source of I_2 was constrained to produce the average nighttime I₂ peak of about 40 ppt that was observed during nocturnal low tide. The rate of I₂ emission in the model followed a Gaussian distribution, peaking at low tide, with a standard width of 22 min (Saiz-Lopez et al., 2005). The NO₃ mixing ratio was fixed at 10 ppt. Figure 8a shows the time- profiles of I_2 , IO and OIO. The O₃ profile is also shown, although the predicted O_3 loss rate is negligible (3×10⁻² ppb h⁻¹). In the

absence of photolysis, the I2 peak decays much more slowly than during the day, in accord with our observations (Saiz-Lopez and Plane, 2004).

In this model run the IO and OIO mixing ratios peak at 2 ppt and 1.7 ppt, respectively, coincident with low tide. Hence, there are two things that differ from the nighttime observations. First, the OIO/IO ratio predicted by the model is 0.9, whereas the measurements indicate that this ratio is ~4. Second, the observations show that the OIO peak occurs after I_2 reaches its maximum at low tide. Therefore, to reconcile the measurements with the model results, another route for converting IO to OIO, in addition to the self reaction of IO, is required. Although the reaction $IO+O_3 \rightarrow OIO+O_2$ is exothermic 15 by about 150 kJ mol⁻¹ (Misra and Marshall, 1998), a very small upper limit to the rate

constant of 10⁻¹⁵ cm³ molecule⁻¹ s⁻¹ has been determined (Atkinson et al., 2000), so that this reaction would be too slow. Another candidate is the reaction:

 $IO + NO_3 \rightarrow OIO + NO_2$

5

Although this reaction does not appear to have been studied, it is also exothermic 20 (by about 50 kJ mol⁻¹), and NO₃ is in general a more aggressive reactant than O₃: for instance, it reacts about 10^5 times faster with I_2 .

When Reaction (1) is included in the model, the rate constant needs to be larger than 7×10^{-12} cm³ molecule⁻¹ s⁻¹ in order to increase the modelled OIO/IO ratio to the observed ratio of 4. This model run is shown in Fig. 8b; note that the OIO peak now oc-25 curs after the maxima of I₂ and IO, as observed. Although this model does not include vertical mixing or consider inhomogeneity along the DOAS light path when comparing with the observations, it does include the uptake of OIO (and IO) onto a background

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.



(1)

aerosol loading typical of Mace Head (Saiz-Lopez et al., 2005). We therefore consider the level of agreement with the observations to be encouraging. A study of Reaction (1) appears to be a priority to develop further understanding of nighttime iodine chemistry.

3.5. BrO observations

- ⁵ BrO was measured over 6 days during NAMBLEX (Saiz-Lopez et al., 2004a). A maximum mixing ratio of 6 ppt was observed, coinciding with westerly and relatively high speed winds (up to 11 m s⁻¹), whereas a mean daytime mixing ratio of 2.3 ppt was observed. The diurnal profile was characterised by a short-lived pulse after dawn. Figure 9 shows an average mixing ratio time-profile from three days of observations (3,
- ¹⁰ 4 and 10 August) when BrO was observed from dawn onwards. The sunrise time at Mace Head for this time of the year was around 04:50 UT, with a 43 min twilight period defined as the period when the SZA is between 96° and 90°. Twilight needs to be considered because, as we show below, some bromine atom precursors are very photolabile. After the BrO pulse, a rapid decrease in its mixing ratio was observed, followed by a partial recovery before noon. Figure 9 indicates that a second small maximum many also he means the late of the second s
- maximum may also be present in the late afternoon before sunset, after which BrO decreased below the detection limit of the DOAS instrument (~1 ppt).

It is unlikely that the post-dawn BrO pulse results from the photolysis of organic bromine species, because of their long photolytic lifetimes in the lower atmosphere

- (Carpenter et al., 1999). One relatively short-lived alkyl bromide is CH₂IBr; however, its photolytic lifetime is still about 1 h, and its mixing ratio in the MBL at Mace Head is ~0.1 ppt (Carpenter et al., 1999). In any case, halocarbon emissions at Mace Head are driven by tidal height (Carpenter et al., 1999), and there is no correlation with tidal height evident in the BrO measurements.
- Hence, the most likely BrO precursors are inorganic bromine species such as Br₂, IBr, BrCl and BrNO₂, which have built up after heterogeneous processing through seasalt aerosol during the preceding night. At sunrise they are rapidly photolysed to yield Br atoms which react with O₃ to form BrO. An interesting problem is to determine which

ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment



of these potential precursors is the main contributor to the post-sunrise BrO pulse. In Fig. 10a the absorption cross-sections of these four bromine precursors are plotted against wavelength, using data from Atkinson et al. (2000). It can be seen that IBr has the largest cross-section, peaking around 500 nm and extending throughout the visible

- ⁵ region of the spectrum. IBr will therefore photolyse most readily around dawn. In contrast, BrNO₂ has an absorption cross-section nearly an order of magnitude smaller, peaking at 300 nm. The rates of photodissociation (*J*) of BrNO₂, Br₂, IBr and BrCl are shown in Fig. 10b as a function of time of day, for the location of Mace Head in early August. The *J* values are calculated, assuming a quantum yield of unity, by an
 ¹⁰ explicit two-stream radiation code (Thompson, 1984), where the incident actinic flux
- (photons cm⁻² s⁻¹ nm⁻¹) at the Earth's surface is obtained after attenuation through 50 atmospheric layers of 1 km height. Immediately after sunrise, J_{IBr} is twice as fast as J_{Br_2} and up to an order of magnitude faster than J_{BrNO_2} . Indeed, if BrNO₂ (and to a lesser extent BrCl) were the major sources of atomic Br, then the BrO pulse would be delayed in appearance and also less sharp.

Laboratory studies of the uptake of N_2O_5 on a saline surface mimicking sea-salt aerosols, which are ~660 times more concentrated in Cl⁻ than Br⁻, have been reported by Behnke et al. (1994):

$$N_2 O_{5(q)} \rightarrow N_2 O_{5(aq)} \tag{2}$$

²⁰
$$N_2O_{5(aq)} + X^- \rightarrow NO_3^- + XNO_2$$

25

where X is Cl or Br. The authors found that ~10% of the released product was $BrNO_2$, with 73% $CINO_2$ and 17% Br_2 . Assuming this product yield for $BrNO_2$ and using the NO_3 chemistry model described in Sect. 3.1, we calculate that for clean marine conditions ([NO_3] and [N_2O_5] ~3 ppt) the cycling through sea-salt aerosol during the night would yield a BrNO2 mixing ratio of ≤ 0.2 ppt before sunrise.

When photolysis ceases after sunset, the uptake of HOBr or $BrNO_3$ on sea-salt aerosol will be their major removal pathways. These species will be rapidly recycled

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.



(3)

through heterogeneous reactions to gas-phase Br_2 and BrCI, depending on the availability of CI^- and Br^- (Finlayson-Pitts, 2003):

 $HOBr + Cl^{-} + H^{+} \rightarrow BrCl + H_{2}O$ (4)

 $BrCl + Br^{-} \leftrightarrow Br_{2}Cl^{-}$

 $5 \quad Br_2Cl^- \leftrightarrow Br_2 + Cl^-$

If the Br^- concentration is sufficient (i.e., in freshly generated sea-salt aerosol), BrCl will be converted into Br_2 via Reactions (5) and (6). In aged sea-salt aerosol that is depleted in Br^- , BrCl becomes the major product (Fickert et al., 1999; Liu and Margerum, 2001; Adams et al., 2002).

- ¹⁰ Although the I⁻ content in sea-salt is negligible compared to CI⁻ and Br⁻, a similar mechanism to Reaction (4) can proceed following uptake of species such as HOI and INO₃, leading to the release of ICI and IBr to the gas phase (Vogt et al., 1999; Mc-Figgans et al., 2000). From laboratory experiments, the main di-halogen gas-phase product is IBr from freshly generated NaCl/NaBr surfaces with sufficient Br⁻ content.
- ¹⁵ Release of ICI was only observed after depletion of Br⁻ (Holmes et al., 2001). The corresponding Henry's law constants (mol l⁻¹ atm⁻¹) for the different species are: BrNO₂ 0.3, Br₂ 1.8, BrCl 0.59 (Frenzel et al., 1998) and 110 and 24 for ICI and IBr, respectively (Wagman et al., 1982).

Considering the above photodissociation rates and the mechanisms for recycling halogens through sea salt, it seems that Br₂ and IBr should be the primary sources of the large post-sunrise BrO pulses observed at Mace Head. However, from our DOAS observations there is no evidence of a post-sunrise pulse of IO (apart from when low tide coincides with sunrise). Putting this together with the fact that the solubility constant of IBr is an order of magnitude greater than that of Br₂, we infer that Br₂ is probably the dominant source of Br at sunrise in this location. **ACPD**

5, 9731-9767, 2005

(5)

(6)

Measurements and modelling of trace gases in a coastal marine environment

Title Page			
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
•	•		
Back	Close		
Full Scr	Full Screen / Esc		
Print	Print Version		
Interactive Discussion			

3.6. Modelling the diurnal behaviour of BrO

We now describe a model of bromine chemistry, containing the gas-phase reactions, photochemistry, and heterogeneous uptake processes listed in Table 1. The photolysis rates were calculated off-line using a two-stream radiation scheme. The model

⁵ is solved using a variable step-size fourth-order Runge-Kutta integrator (Press et al., 1986). All bromine-containing species and O₃ are allowed to vary. The mixing ratios of the following species are held constant at values typical of clean MBL conditions: [NO₂]=30 ppt; [NO]=5 and 0 ppt at day- and nighttime, respectively; [DMS]=100 ppt; [CH₂IBr]=0.1 ppt. The mixing ratios of OH and HO₂ are constrained in the model to follow the diurnal profiles of these radicals measured during NAMBLEX, with noon maxima of 0.1 and 5 ppt, respectively (Bloss et al., 2005).

The heterogeneous reaction of HOBr with sea salt and subsequent liberation of Br_2 and BrCl to the gas phase has been observed to occur rapidly (Abbatt and Waschewsky, 1998; Fickert et al., 1999; Adams et al., 2002). These experiments were

¹⁵ conducted on surfaces with a Cl⁻/Br⁻ ratio of ~660. Considering the aqueous phase reaction rates and equilibrium constants for Reactions (4–6) (Liu and Margerum, 2001; Finlayson-Pitts, 2003), a simple calculation shows that it would take a few minutes for HOBr, once taken up on a sea-salt aerosol surface, to be released as Br₂ and, to a lesser extent BrCl, depending upon the Br⁻ content and acidity of the aerosol (Fickert et al., 1999). Hence, the rate-limiting step of this heterogeneous process will be the uptake of HOBr on the aerosol surface. In the model runs shown here we consider the uptake of HOBr to generate 75% Br₂ and 25% BrCl.

Likewise, aqueous phase reactions involving $BrNO_3$ could also lead to the release of Br_2 and BrCl via hydrolysis or reaction with Cl^- and Br^- (Sander et al., 1999):

$BrNO_3 + H_2O \rightarrow HOBr + HNO_3$	
--	--

 $BrNO_3 + Cl^- \rightarrow BrCl + NO_3^-$ (8)

$$BrNO_3 + Br^- \rightarrow Br_2 + NO_3^-$$

ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.



(7)

(9)

Reaction of BrNO₃ on aerosol surfaces via Reactions (8) and (9) can proceed without aerosol acidity, and hence allow halogen activation on alkaline aerosols (Sander et al., 1999). Figure 11 shows the result of a model simulation for clean marine conditions, where the sea-salt surface area is set to 10^{-7} cm² cm⁻³. The model is initialised with ⁵ Br₂ and BrCl mixing ratios of 4.5 and 1.5 ppt, respectively. A rapid pulse of BrO up to 4.5 ppt is generated as soon as photolysis starts. In order to match the observed pulse of BrO (Fig. 9) the rate of post-sunrise Br atom production needed is $\sim 8 \times 10^3$ molecule $cm^{-3} s^{-1}$. As Br₂ and BrCl are photolysed and their mixing ratio rapidly decreases, BrO reaches a minimum followed by a slow recovery throughout the day leading to a second maximum in the late afternoon. BrO then disappears as photolysis ceases after 10 sunset. Similar diurnal behaviour has been predicted in previous studies (Sander et al., 1999; Glasow et al., 2002). The modelled BrO time-profile in Fig. 11 is comparable to that observed (Fig. 9), with midday/afternoon mixing ratios around 1.5-2.5 ppt (Saiz-Lopez et al., 2004). Note that this model does not incorporate BrO mixing within the boundary layer. The BrO increase towards the late afternoon is due to the recycling of

15 HOBr and BrNO₃ through sea-salt aerosol.

After sunset, large amounts of BrNO₃ build up as photolysis ceases, and this becomes the major bromine species. In this model run BrNO₃ has a loss rate due to uptake on aerosols of $\sim 1 \times 10^{-5} \text{ s}^{-1}$ if $\gamma = 0.02$ (note that the uptake of BrNO₃ on deliquesced sea salt aerosols does not appear to have been measured). The daytime

- 20 average photolysis rate of BrNO₃ is $\sim 8.5 \times 10^{-4} \text{ s}^{-1}$, so this will probably be the dominant loss process, competing with thermal decomposition which has a measured rate of 9×10^{-6} s⁻¹ at 290 K (Orlando and Tyndall, 1996). The BrNO₃ time-profile in Fig. 11 exhibits two distinct peaks: after the sunrise BrO pulse, and in the early evening when
- photolysis slows down. The BrNO₃ minimum around noon is caused by photolysis, 25 which is sharply peaked to noon because the molecule absorbs almost entirely in the near-UV. After sunset the remaining BrNO₃ is slowly taken up on aerosols, and this is most likely the major route to the overnight build-up of gas-phase Br₂.

HOBr follows the diurnal behaviour of HO₂, with a maximum at midday. HOBr re-

ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment



moval during daytime is dominated by photodissociation, as J_{HOBr} is about 40 times greater than heterogeneous loss to form Br₂ and BrCl. Here a γ of 0.05 for HOBr is used, in accord with a lower limit of $\gamma > 10^{-2}$ measured on aqueous salt solutions (Fickert et al., 1999; Adams et al., 2002). Note that Abbatt and Waschewsky (1998) reported $\gamma > 0.2$ for deliquescent NaCl aerosols within an acidity pH range 0.3–7. HOBr uptake and production of gas-phase bromine-containing is dependant on aerosol acidity, and decreases with decreasing acidity (Fickert et al., 1999). Here we are assuming a scenario of freshly generated sea-salt aerosols with pH = 8, the same as ocean water (Glasow et al., 2002, and references therein). It must be noted that the pro-

- ¹⁰ gressive acidification of aerosols by uptake of species such as HNO₃, H₂SO₄ and SO₂ is not treated in the model. Considering an average volume concentration of aerosol (diameter = 0.3–47 μ m) measured during NAMBLEX of 400 μ m³ cm⁻³ (G. McFiggans, personal communication, 2005) and a Br⁻ content in sea salt of 8×10⁻³ mol l⁻¹ we calculate a Br⁻ depletion from sea-salt of 22% over the 1-day model run.
- After sunset, the main sink for HOBr is heterogeneous uptake, contributing to the nighttime production of gas-phase bromine atom precursors: Fig. 11 shows the Br₂ build up once photolysis ceases leading to a maximum before sunrise, which will be responsible of the second post-dawn BrO pulse. Note that O₃ is depleted through the day at a rate of approximately 0.06 ppb h⁻¹, mainly through the cycle involving
 BrO+HO₂. Of course, this is a zero-dimensional model and does not include mixing with O₃-rich air from aloft.

4. Conclusions

The techniques of long-path DOAS and zenith sky spectroscopy have been employed to acquire a comprehensive set of measurements of I_2 , IO, OIO, BrO and NO₃ in a coastal mid-latitude location. The time series of the different species over a month of observations are summarised in this paper. The observed daytime mixing ratios of the halogen oxides will affect a number of atmospheric processes such as ozone de-

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

Title Page			
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
	►I		
•	►		
Back	Close		
Full Scr	Full Screen / Esc		
Print Version			
Interactive Discussion			

pletion, DMS oxidation capacity, new particle formation and regulation of the HO₂/OH and NO₂/NO ratios. Model simulations show that the reaction between NO₃ and I₂ probably accounts for the significant nighttime levels of iodine oxides found at Mace Head. The observed pulses of BrO at sunrise, and subsequent diurnal behaviour of the radical, are explained with a model that incorporates the processing of the major daytime bromine reservoirs, BrNO₃ and HOBr, through sea-salt aerosol. However, the NAMBLEX data-set on BrO is quite small and more field observations are needed in order to establish this diurnal behaviour. Finally, the combination of boundary layer and vertical profile measurements of NO₃ indicates that this radical often exists at higher concentration in the upper MBL and lower free troposphere, probably reflecting the

concentration in the upper MBL and lower free troposphere, proba efficient removal of NO₃ by DMS close to the ocean surface.

Acknowledgements. This work was supported by the Natural Environment Research Council of the UK. The authors wish to thank D. Heard (University of Leeds) for leading the NAMBLEX campaign and B. Allan for helpful discussions.

15 References

20

Abbatt, J. P. D. and Waschewsky, G. C. G.: Heterogeneous interactions of HOBr, HNO₃, O₃ and NO₂ with deliquescent NaCl aerosols at room temperature, J. Phys. Chem. A, 102, 3719–3725, 1998. 9746, 9748

Adams, J. W., Holmes, N. S., and Crowley, J. N.: Uptake and reaction of HOBr on frozen and

dry NaCl/NaBr surfaces between 253 and 233 K, Atmos. Chem. Phys., 2, 79–91, 2002, SRef-ID: 1680-7324/acp/2002-2-79. 9745, 9746, 9748, 9756

Alicke, B., Hebestreit, K., Stutz, J., and Platt, U.: Iodine oxide in the marine boundary layer, Nature, 397, 572–573, 1999. 9733, 9740

Allan, B. J. and Plane, J. M. C.: A study of the recombination of IO and NO₂ and the stability

of INO₃: implications for the atmospheric chemistry of iodine, J. Phys. Chem. A, 106, 8634– 8641, 2002b. 9741

Allan, B. J., Carslaw, N., Coe, H., Burgess, R. A., and Plane, J. M. C.: Observations of the nitrate radical in the marine boundary layer, J. Atmos. Chem., 33, 129–154, 1999. 9736

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.



EGU

- Allan, B. J., McFiggans, G., Plane, J. M. C., and Coe, H.: Observations of iodine monoxide in the remote marine boundary layer, J. Geophys. Res.-Atmos, 105, 14363–14369, 2000a. 9733, 9740
- Allan, B. J., McFiggans, G., Plane, J. M. C., and McFadyen, G. G.: The nitrate radical in the
- remote marine boundary layer, J. Geophys. Res.-Atmos, 105, 24191–24204, 2000b. 9734, 9736, 9737
 - Allan, B. J., Plane, J. M. C., and McFiggans, G.: Observations of OIO in the remote marine boundary layer, Geophys. Res. Lett., 28, 1945–1948, 2001. 9733, 9740
 - Allan, B. J., Plane, J. M. C., Coe, H., and Shillito, J. A.: Observations of NO₃ concentration pro-
- files in the troposphere, J. Geophys. Res., 107, art. no. 4588, doi:10.1029/2002JD002112, 2002a. 9734, 9735, 9737, 9738
 - Ashworth, S. H., Allan, B. J., and Plane, J. M. C.: High resolution spectroscopy of the OIO radical: Implications for the ozone-depleting potential of iodine, Geophys. Res. Lett., 29, art. no. 1456, doi:10.1029/2001GL013851, 2002. 9740
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson Jr., R. F., Hynes, R. G., Jenkin, M. E., Kerr, J. A., Rossi, M. J., and Troe, J.: Summary of evaluated kinetic and photochemical data for atmospheric chemistry, IUPAC, J. Phys. Chem. Ref. Data, 29, 167– 266, http://www.iupac-kinetic.ch.cam.ac.uk/, 2000. 9742, 9744, 9756

Ayers, G. P., Gillet, R. W., Cainey, J. M., and Dick, A. L.: Chloride and bromide loss froma sea-salt particles in Southern Ocean air, J. Atmos. Chem., 33, 299–319, 1999. 9733

Barrie, L. A., Bottenheim, J. W., Schnell, R. C., Crutzen, P. J., and Rasmussen, R. A.: Ozone destruction and photochemical-reactions at polar sunrise in the lower Arctic atmosphere, Nature, 334, 138–141, 1988. 9732

20

- Behnke, W., Scheer, V., and Zetzsch, C.: Production of BrNO₂, Br₂ and CINO₂ from the reaction between sea spray aerosol and N₂O₅, J. Aero. Sci., 25, S277–S278, 1994. 9733, 9744
- ²⁵ between sea spray aerosol and N₂O₅, J. Aero. Sci., 25, S277–S278, 1994. 9733, 9744
 Behnke, W., George, C., Scheer, V., and Zetzsch, C.: Production and decay of CINO₂ from the reaction of gaseous N₂O₅ with NaCl solution: bulk and aerosol experiments, J. Geophys. Res.-Atmos, 102, 3795–3804, 1997. 9737

Bloss, W. J., Lee, J. D., Johnson, G. P., Sommariva, R., Heard, D. E., Saiz-Lopez, A., Plane,

J. M. C., McFiggans, G., Coe, H., Flynn, M., Williams, P., Rickard, A. R., and Fleming, Z.: Impact of halogen monoxide chemistry upon boundary layer OH and HO₂ concentrations at a coastal site, Geophys. Res. Lett., 32, art. no. L06814, doi:10.1029/2004GL022084, 2005. 9733, 9746 5, 9731–9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.

Title Page		
Abstract	Introduction	
Conclusions	References	
Tables	Figures	
•	•	
	Close	
Back	Close	
Back Full Scr	Close een / Esc	
Back Full Scr	Close een / Esc	
Back Full Scree Print V	Close een / Esc /ersion	

EGU

- Brown, S. S., Stark, H., Ryerson, T. B., Williams, E. J., Nicks, D. K., Trainer, M., Fehsenfeld, F. C., and Ravishankara, A. R.: Nitrogen oxides in the nocturnal boundary layer: Simultaneous in situ measurements of NO₃, N₂O₅, NO₂, NO, O₃, J. Geophys. Res.-Atmos, 108, art. no. 4299, doi:10.1029/2002JD002917, 2003. 9734
- ⁵ Brown, S. S., Dibb, J. E., Stark, H., Aldener, M., Vozella, M., Whitlow, S., Williams, E. J., Lerner, B. M., Jakoubek, R., Middlebrook, A. M., DeGouw, J. A., Warneke, C., Goldan, P. D., Kuster, W. C., Angevine, W. M., Sueper, D. T., Quinn, P. K., Bates, T. S., Meagher, J. F., Fehsenfeld, F. C., and Ravishankara, A. R.: Nighttime removal of NO_x in the summer marine boundary layer, Geophys. Res. Lett., 31, art. no. L07108, doi:10.1029/2004GL019412, 2004. 9734, 9736
 - Butkovskaya, N. I. and LeBras, G.: Mechanism of the NO₃ + DMS reaction by discharge flow mass- spectrometry, J. Phys. Chem. A, 98, 2582–2591, 1994. 9734
 - Carpenter, L. J., Sturges, W. T., Penkett, S. A., Liss, P. S., Alicke, B., Hebestreit, K., and Platt, U.: Short-lived alkyl iodides and bromides at Mace Head, Ireland: Links to biogenic sources
- and halogen oxide production, J. Geophys. Res.-Atmos, 104, 1679–1689, 1999. 9733, 9743
 Chambers, R. M., Heard, A. C., and Wayne, R. P.: Inorganic gas-phase reactions of the nitrate radical I₂+NO₃ and I+NO₃, J. Phys. Chem. A, 96, 3321–3331, 1992. 9733, 9741
 - Coe, H., Allan, B. J., and Plane, J. M. C.: Retrieval of vertical profiles of NO₃ from zenith sky measurements using an optimal estimation method, J. Geophys. Res.-Atmos, 107, art. no. 4587, doi:10.1029/2002JD002111, 2002. 9737, 9738
 - Davis, D. J., Crawford, J., Liu, S., McKeen, S., Bandy, A., Thorton, D., Rowland, F., and Blake,
 D.: Potential impact of iodine on tropospheric levels of Ozone and other critical oxidants, J.
 Geophys. Res.-Atmos, 101, 2135–2147, 1996. 9733
 - Deiber, G., George, C., Calve, S. L., Schweitzer, F., and Mirabel, P.: Uptake study of CINO₂ and BrNO₂ by halide containing droplets, Atmos. Chem. Phys., 4, 1291–1299, 2004,

SRef-ID: 1680-7324/acp/2004-4-1291 9756

20

25

- Farman, J. C., Gardiner, B. G., and Shanklin, J. D.: Large losses of total ozone in Antarctica reveal seasonal CIO_x/NO_x interaction, Nature, 315, 207–210, 1985. 9732
- Fickert, S., Adams, J. W., and Crowley, J. N.: Activation of Br₂ and BrCl via uptake of HOBr onto
- ³⁰ aqueous salt solutions, J. Geophys. Res.-Atmos, 104, 23719–23727, 1999. 9745, 9746, 9748, 9756
 - Finlayson-Pitts, B. J.: The tropospheric chemistry of sea salt: a molecular-level view of the chemistry of NaCl and NaBr, Chem. Rev., 103, 4801–4822, 2003. 9745, 9746

ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

Title Page			
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
•	►		
Back	Close		
Full Scr	Full Screen / Esc		
Print Version			
Interactive Discussion			

- Finlayson-Pitts, B. J., Livingston, F. E., and Berko, H. N.: Ozone destruction and bromine photochemistry at ground-level in the Arctic spring, Nature, 343, 622–625, 1990. 9732
- Frenzel, A., Scheer, V., Sikorski, R., George, C., Behnke, W., and Zetzsch, C.: Heterogeneous interconversion reactions of BrNO₂, CINO₂, Br₂ and Cl₂, J. Phys. Chem. A, 102, 1329–1337,
- ⁵ 1998. 9745

10

25

30

- Gabriel, R., von Glasow, R., Sander, R., Andreae, M. O., and Crutzen, P. J.: Bromide content of sea-salt aerosols collected over the Indian Ocean during INDOEX 1999, J. Geophys. Res.-Atmos, 107, 8032, doi:10.1029/2001JD001133, 2002. 9733
- Garland, J. A. and Curtis, H.: Emission of iodine from the sea surface in the presence of ozone, J. Geophys. Res.-Atmos, 86, 3183–3186, 1981. 9739
- Glasow, R. V. and Crutzen, P. J.: Model study of multiphase DMS oxidation with a focus on halogens, Atmos. Chem. Phys., 4, 589–608, 2004,

SRef-ID: 1680-7324/acp/2004-4-589. 9733

Glasow, R. V., Sander, R., Bott, A., and Crutzen, P. J.: Modelling halogen chemistry

- ¹⁵ in the marine boundary layer 1. Cloud-free MBL, J. Geophys. Res.-Atmos, 107, 4341, doi:10.1029/2001JD000942, 2002. 9747, 9748
 - Harwood, M. H., Burkholder, J. B., Hunter, M., Fox, R. W., and Ravishankara, A. R.: Absorption cross sections and self-reaction kinetics of the IO radical, J. Phys. Chem. A, 101, 853–863, 1997. 9735
- Hausmann, M. and Platt, U.: Spectroscopic measurement of bromine oxide and ozone in the high Arctic during Polar Sunrise Experiment 1992, J. Geophys. Res.-Atmos, 99, 25399– 25413, 1994. 9732
 - Heintz, F., Platt, U., Flentje, H., and Dubois, R.: Long-term observation of nitrate radicals at the Tor station, Kap Arkona (Rugen), J. Geophys. Res.-Atmos, 101, 22891–22910, 1996. 9734, 9736
 - Holmes, N. S., Adams, J. W., and Crowley, J. N.: Uptake and reaction of HOI and IONO₂ on frozen and dry NaCl/NaBr surfaces and H₂SO₄, Phys. Chem. Chem. Phys., 3, 1679–1687, 2001. 9745
 - Honninger, G. and Platt, U.: Observations of BrO and its vertical distribution during surface ozone depletion at Alert, Atmos. Environ., 36, 2481–2489, 2002. 9732
 - Jenkin, M. E., Cox, R. A., and Candeland, D. E.: Photochemical aspects of tropospheric iodine behavior, J. Atmos. Chem., 2, 359–375, 1985. 9733

Kupper, F. C., Schweigert, N., Gall, E. A., Legendre, J. M., Vilter, H., and Kloareg, B.: Iodine

ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.

Title Page		
Abstract	Introduction	
Conclusions	References	
Tables	Figures	
	P 1	
•	•	
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		

EGU

uptake in Laminariales involves extracellular, haloperoxidase-mediated oxidation of iodide, Planta, 207, 163–171, 1998. 9739

- Leser, H., Honninger, G., and Platt, U.: MAX-DOAS measurements of BrO and NO₂ in the marine boundary layer, Geophys. Res. Lett., 30, doi:10.1029/2002GL015811, 2003. 9733
- Liu, Q. and Margerum, D. W.: Equilibrium and kinetics of bromine chloride hydrolysis, Environ. Sci. Tech., 35, 1127–1133, 2001. 9745, 9746
 - McConnell, J. C., Henderson, G. S., Barrie, L., Bottenheim, J., Niki, H., Langford, C. H., and Templeton, E. M. J.: Photochemical bromine production implicated in Arctic boundary-layer ozone depletion, Nature, 355, 150–152, 1992. 9732
- McFiggans, G., Plane, J. M. C., Allan, B. J., Carpenter, L. J., Coe, H., and O'Dowd, C.: A modeling study of iodine chemistry in the marine boundary layer, J. Geophys. Res.-Atmos, 105, 14371–14385, 2000. 9733, 9741, 9745
 - McFiggans, G., Coe, H., Burgess, R., Allan, J., Cubison, M., Alfarra, M. R., Saunders, R., Saiz-Lopez, A., Plane, J. M. C., Wevill, D. J., Carpenter, L. J., Rickard, A. R., and Monks,
- P. S.: Direct evidence for coastal iodine particles from Laminaria macroalgae linkage to emissions of molecular iodine, Atmos. Chem. Phys., 4, 701–713, 2004, SRef-ID: 1680-7324/acp/2004-4-701. 9734, 9739
 - Misra, A. and Marshall, P.: Computational investigations of iodine oxides, J. Phys. Chem. A, 102, 9056–9060, 1998. 9742
- Molina, M. J. and Rowland, F. S.: Stratospheric sink for chlorofluoromethanes chlorine atomcatalysed destruction of ozone, Nature, 249, 810–812, 1974. 9732
 - O'Dowd, C. D., Jimenez, J. L., Bahreini, R., Flagan, R. C., Seinfeld, J. H., Hameri, K., Pirjola, L., Kulmala, M., Jennings, S. G., and Hoffmann, T.: Marine aerosol formation from biogenic iodine emissions, Nature, 417, 632–636, 2002. 9734
- Orlando, J. J. and Tyndall, G. S.: Rate coefficients for the thermal decomposition of BrONO₂ and the heat of formation of BrONO₂, J. Phys. Chem. A, 100, 19398–19405, 1996. 9747, 9756

Plane, J. M. C. and Saiz-Lopez, A.: UV-visible Differential Optical Absorption Spectroscopy (DOAS), in: Analytical Techniques for Atmospheric Measurement; edited by: Heard, D. E.,

- ³⁰ Blackwell Publishing, Oxford, in press, 2005. 9735
 - Platt, U., Perner, D., Harris, G. W., Winer, A. M., and Pitts, J. N.: Detection of NO₃ in the polluted troposphere by differential optical absorption, Geophys. Res. Lett., 7, 89–92, 1980. 9734
 Platt, U., LeBras, G., Poulet, G., Burrows, J. P., and Moortgat, G.: Peroxy radicals from night-

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

Title Page		
Abstract	Introduction	
Conclusions	References	
Tables	Figures	
•	•	
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		

time reaction of NO₃ with organic compounds, Nature, 348, 147–149, 1990. 9734 Press, W. H., Flannery, B. P., Teukolsky, S. A., and Vetterling, W. T.: Numerical recipes: The art of scientific computing, Cambridge University Press, Cambridge, 1986. 9746 Rodgers, C. D.: Retrieval of atmospheric temperature and composition from remote measure-

- ments of thermal-radiation, Rev. Geophys., 14, 609–624, 1976. 9738
 Bedgere, C. D.: Characterization and error analysis of profiles retrieved from remote soundir
 - Rodgers, C. D.: Characterization and error analysis of profiles retrieved from remote sounding measurements, J. Geophys. Res.-Atmos, 95, 5587–5595, 1990. 9738
 - Saiz-Lopez, A. and Plane, J. M. C.: Novel iodine chemistry in the marine boundary layer, Geophys. Res. Lett., 31, art. no. L04112, doi:10.1029/2003GL019215, 2004. 9733, 9739, 9740, 9741, 9742
- Saiz-Lopez, A., Plane, J. M. C., and Shillito, J. A.: Bromine oxide in the mid-latitude marine boundary layer, Geophys. Res. Lett., 31, art. no. L03111, doi:10.1029/2003GL018956, 2004a. 9733, 9743

Saiz-Lopez, A., Saunders, R. W., Joseph, M., and Plane, J. M. C.: Absolute absorption cross-

section and photolysis rate of I₂, Atmos. Chem. Phys., 4, 1443–1450, 2004b, SRef-ID: 1680-7324/acp/2004-4-1443. 9735

10

20

Saiz-Lopez, A., Plane, J. M. C., McFiggans, G., Williams, P. I., Ball, S. M., Bitter, M., Jones, R. L., Hongwei, C., and Hoffmann, T.: Modelling molecular iodine emissions in a coastal marine environment: the link to new particle formation, Atmos. Chem. Phys. Discuss., 5, 5405–5439, 2005.

SRef-ID: 1680-7375/acpd/2005-5-5405. 9734, 9739, 9740, 9741, 9742, 9743

- Sander, R., Rudich, Y., von Glasow, R., and Crutzen, P. J.: The role of BrNO₃ in marine tropospheric chemistry: A model study, Geophys. Res. Lett., 26, 2857–2860, 1999. 9746, 9747
- ²⁵ Sander, S. P., Friedl, R. R., Golden, D. M., Kurylo, M. J., Huie, R. E., Orkin, V. L., Moortgat, G. K., Ravishankara, A. R., Kolb, C. E., Molina, M., and Finlayson-Pitts, B. J.: Chemical kinetics and photochemical data for use in stratospheric modeling: Evaluation 14, Tech. rep., Jet Propulsion Laboratory, Pasadena, California, USA, http://jpldataeval.jpl.nasa.gov/, 2003. 9736, 9756
- ³⁰ Schweitzer, F., Mirabel, P., and George, C.: Multiphase chemistry of N₂O₅, CINO₂, BrNO₂, J. Phys. Chem. A, 102, 3942–3952, 1998. 9756
 - Smith, J. P. and Solomon, S.: Atmospheric NO₃. 3. Sunrise disappearance and the stratospheric profile, J. Geophys. Res.-Atmos, 95, 13819–13827, 1990. 9737

5, 9731–9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

Title Page		
Abstract	Introduction	
Conclusions	References	
Tables	Figures	
I	۶I	
•	•	
Back	Close	
Full Screen / Esc		
Print Version		
Interactive Discussion		

EGU

- Thompson, A. M.: The effect of clouds on photolysis rates and ozone formation in the unpolluted troposphere, J. Geophys. Res.-Atmos, 89, 1341–1349, 1984. 9744
- Truesdale, V. W., Luther, G. W., and Canosamas, C.: Molecular iodine reduction in seawater – an improved rate equation considering organic compounds, Mar. Chem., 48, 2, 143–150, 1995. 9739
- Vogt, R., Crutzen, P. J., and Sander, R.: A mechanism for halogen release from sea-salt aerosol in the remote marine boundary layer, Nature, 383, 327–330, 1996. 9733
- Vogt, R., Sander, R., Glasow, R. V., and Crutzen, P. J.: Iodine chemistry and its role in halogen activation and ozone loss in the marine boundary layer: A model study, J. Atmos. Chem., 32, 375–395, 1999. 9745

Vrekoussis, M., Kanakidou, M., Mihalopoulos, N., Crutzen, P. J., Lelieveld, J., Perner, D.,

¹⁵ Berresheim, H., and Baboukas, E.: Role of the NO₃ radicals in oxidation processes in the eastern Mediterranean troposphere during the MINOS campaign, Atmos. Chem. Phys., 4, 169–182, 2004,

SRef-ID: 1680-7324/acp/2004-4-169. 9736

Wagman, D. D., Evans, W. H., Parker, V. B., Schumm, R. H., Halow, I., Bailey, S. M., Churney,

- K. L., and Nuttall, R. L.: The NBS tables of chemical thermodynamic properties; Selected values for inorganic and C₁ and C₂ organic substances in SI units, J. Phys. Chem. Ref. Data, 11, suppl. 2, 1982. 9745
 - Wahner, A., Ravishankara, A. R., Sander, S. P., and Friedl, R. R.: Absorption cross-section of BrO between 312 and 385 nm at 298 and 223 K, Chem. Phys. Lett., 152, 507–512, 1988. 9735
 - Wofsy, S. C., McElroy, M. B., and Yung, Y. L.: The chemistry of atmospheric bromine, Geophys. Res. Lett., 2, 215–218, 1975. 9732
 - Yokelson, R. J., Burkholder, J. B., Fox, R. W., Talukdar, R. K., and Ravishankara, A. R.: Temperature-Dependence of the NO₃ absorption spectrum, J. Phys. Chem. A, 98, 13144–
- ³⁰ 13 150, 1994. 9735, 9737

5

10

25

ACPD

5, 9731–9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

Title Page		
Abstract	tract	
Conclusions	References	
Tables	Figures	
	►	
Back	Close	
Back Full Scre	Close en / Esc	
Back Full Scree Print V	Close en / Esc ersion	

No	Reaction	Rate constant	Remarks ^b
(R1)	$Br + O_3 \rightarrow BrO + O_2$	$1.7 \times 10^{-11} e^{(-800/T)}$	1
(R2)	HBr + $OH \rightarrow Br + H_2O$	1.1×10 ⁻¹¹	1
(R3)	Br + HO ₂ \rightarrow HBr + O ₂	$1.5 \times 10^{-11} e^{(-600/T)}$	1
(R4)	$Br + NO_2 + M \rightarrow BrNO_2$	$k_o = 4.2 \times 10^{-31} \times (T/300)^{-2.4}$	1
		$k_{\infty} = 2.7 \times 10^{-11} \times (T/300)^{-0}$	
(R5)	$BrO + NO_2 + M \rightarrow BrNO_3$	$k_o = 5.2 \times 10^{-31} \times (T/300)^{-3.2}$	1
		$k_{\infty} = 6.9 \times 10^{-12} \times (T/300)^{-2.9}$	
(R6)	BrO + HO ₂ \rightarrow HOBr + O ₂	$3.4 \times 10^{-12} e^{(540/T)}$	1
(R7)	$BrO + NO \rightarrow Br + NO_2$	8.8×10 ⁻¹² e ^(260/T)	1
(R8)	$BrO + CH_3SCH_3 \rightarrow CH_3SOCH_3 + Br$	$1.5 \times 10^{-14} e^{(850/T)}$	1
(R9)	$BrO + BrO \rightarrow 2Br + O_2$	$2.4 \times 10^{-12} e^{(40/T)}$	1
(R10)	$BrO + BrO \rightarrow Br_2 + O_2$	2.8×10 ⁻¹⁴ e ^(860/T)	1
(R11)	$BrNO_3 \rightarrow BrO + NO_2$	2.8×10 ¹³ e ^{-(12360/T)}	2
(R12)	$BrO + hv \rightarrow Br + O$	0.05	3
(R13)	$Br_2 + h\nu \rightarrow 2Br$	0.0315	3
(R14)	$IBr + hv \rightarrow Br + I$	0.067	3
(R15)	$BrCl + hv \rightarrow Br + Cl$	9.8×10 ⁻³	3
(R16)	$BrNO_2 + hv \rightarrow Br + NO_2$	5.8×10 ⁻³	3
(R17)	$BrNO_3 + hv \rightarrow BrO + NO_2 (\Phi=0.71)$	1.4×10 ⁻³	3
	\rightarrow Br + NO ₃ (Φ = 0.29)		
(R18)	$HOBr + hv \rightarrow Br + OH$	1.8×10 ⁻³	3
(R19)	$CH_2IBr + hv \rightarrow CH_2 + Br + I$	3.8×10 ⁻⁴	3
(R20)	Uptake coefficient of HOBr	0.05	4
(R21)	Uptake coefficient of HBr	0.03	5
(R22)	Uptake coefficient of BrNO ₃	0.02	6

Table 1. Bromine chemistry scheme used in the photochemical box model^a.

^a Units: unimolecular reactions, s⁻¹; photolysis rate constants, s⁻¹; bimolecular reactions, cm³ molecule⁻¹ s⁻¹; termolecular reactions, cm⁶ molecule⁻² s⁻¹, calculated using the formalism of Sander et al. (2003), where $k = ((k_o [M]/(k_{\infty})) \times F_c^n)$, $F_c = 0.6$ and $n = (1 + (\log_{10}(k_o [M]/k_{\infty}))^2)^{-1}$.

^b Remarks: 1, Sander et al. (2003); 2, from Orlando and Tyndall (1996), applicable for atmospheric pressure; 3, absorption cross-sections taken from Atkinson et al. (2000); 4, from Fickert et al. (1999); Adams et al. (2002); 5, from Schweitzer et al. (1998); 6, from latest IUPAC dataset 2005, estimated from Deiber et al. (2004).

Title Page Introduction Abstract Conclusions References **Tables** Figures ∎∎ ► Back Close Full Screen / Esc **Print Version** Interactive Discussion EGU

ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.





Fig. 1. NO₃ time series as measured by the LP-DOAS instrument during NAMBLEX. The mixing ratios are plotted together with the 2σ uncertainties.





ACPD 5, 9731–9767, 2005 Measurements and modelling of trace gases in a coastal

Marine environment



EGU





5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment









Fig. 4. (a) Zenith-sky spectroscopy measurements of the NO_3 column abundance as a function of SZA during six days of observations at Mace Head. The solid red line indicates the profile predicted by the OEM forward model. **(b)** The corresponding retrieved vertical concentration profile of NO_3 . 9760

EGU





5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment





Fig. 6. Plot of I_2 mixing ratio versus tidal height at Mace Head.

ACPD

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment



5, 9731–9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.



Fig. 7. Mesh plot showing the observed daytime IO as a function of tidal height and solar irradiance.



5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.



Fig. 8. (a) Nighttime run of the iodine model, with low tide 60 min after the start. **(b)** Model run with the same initial conditions as (a) but including the reaction $IO+NO_3 \rightarrow OIO+NO_2$.

a)

50

40

30

20

10

0 -

I₂ / ppt

6 5

4

3

2

1

0 + 0

50

IO, OIO / ppt

b)



03

ю

100

010

150

Time / minutes

200

250

25.1

25.0

- 24.9 qdd / [©]O

24.8

24.7

300





5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.



EGU



5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.



Fig. 10. (a) Absorption cross-sections of Br_2 , BrCl, IBr and $BrNO_2$ as a function of wavelength. **(b)** Diurnal variation of the corresponding rates of photolysis, calculated assuming a quantum yield of unity for clear sky conditions at Mace Head in August.

5, 9731-9767, 2005

Measurements and modelling of trace gases in a coastal marine environment

A. Saiz-Lopez et al.





Fig. 11. Model simulation of the diurnal profiles of the main inorganic bromine species and O_3 , for low NO_x conditions in the MBL.