



HAL
open science

On oriented arc-coloring of subcubic graphs

Alexandre Pinlou

► **To cite this version:**

Alexandre Pinlou. On oriented arc-coloring of subcubic graphs. The Electronic Journal of Combinatorics, 2006, 13, pp.131. hal-00307094

HAL Id: hal-00307094

<https://hal.science/hal-00307094>

Submitted on 30 Jul 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On oriented arc-coloring of subcubic graphs

Alexandre Pinlou

Alexandre.Pinlou@labri.fr

LaBRI, Université Bordeaux I,
351, Cours de la Libération,
33405 Talence, France

January 17, 2006

Abstract. A homomorphism from an oriented graph G to an oriented graph H is a mapping φ from the set of vertices of G to the set of vertices of H such that $\overrightarrow{\varphi(u)\varphi(v)}$ is an arc in H whenever \overrightarrow{uv} is an arc in G . The oriented chromatic index of an oriented graph G is the minimum number of vertices in an oriented graph H such that there exists a homomorphism from the line digraph $LD(G)$ of G to H (Recall that $LD(G)$ is given by $V(LD(G)) = A(G)$ and $\overrightarrow{ab} \in A(LD(G))$ whenever $a = \overrightarrow{uv}$ and $b = \overrightarrow{vw}$). We prove that every oriented subcubic graph has oriented chromatic index at most 7 and construct a subcubic graph with oriented chromatic index 6.

AMS Subject Classification: 05C15.

Keywords: Graph coloring, oriented graph coloring, arc-coloring, cubic graphs.

1 Introduction

We consider finite simple *oriented graphs*, that is digraphs with no opposite arcs. For an oriented graph G , we denote by $V(G)$ its set of vertices and by $A(G)$ its set of arcs.

In [2], Courcelle introduced the notion of vertex-coloring of oriented graphs as follows: an *oriented k -vertex-coloring* of an oriented graph G is a mapping φ from $V(G)$ to a set of k colors such that (i) $\varphi(u) \neq \varphi(v)$ whenever \overrightarrow{uv} is an arc in G , and (ii) $\varphi(u) \neq \varphi(x)$ whenever \overrightarrow{uv} and \overrightarrow{wx} are two arcs in G with $\varphi(v) = \varphi(w)$. The *oriented chromatic number* of an oriented graph G , denoted by $\chi_o(G)$, is defined as the smallest k such that G admits an oriented k -vertex-coloring.

Let H and H' be two oriented graphs. A *homomorphism* from H to H' is a mapping φ from $V(H)$ to $V(H')$ that preserves the arcs: $\overrightarrow{\varphi(u)\varphi(v)} \in A(H')$ whenever $\overrightarrow{uv} \in A(H)$. An oriented k -vertex-coloring of G can be equivalently defined as a homomorphism φ from G to H , where H is an oriented graph of order k . The existence of such a homomorphism from G to H is denoted by $G \rightarrow H$. The graph H will be called *color-graph* and its vertices will be called *colors*, and we will say that G is

H -colorable. The oriented chromatic number can be then equivalently defined as the smallest order of an oriented graph H such that $G \rightarrow H$.

Oriented vertex-colorings have been studied by several authors in the last past years (see e.g. [1, 3, 5] or [7] for an overview).

One can define *oriented arc-colorings* of oriented graphs in a natural way by saying that, as in the undirected case, an oriented arc-coloring of an oriented graph G is an oriented vertex-coloring of the line digraph $LD(G)$ of G (Recall that $LD(G)$ is given by $V(LD(G)) = A(G)$ and $\vec{ab} \in A(LD(G))$ whenever $a = \vec{uv}$ and $b = \vec{vw}$). We will say that an oriented graph G is *H -arc-colorable* if there exists a homomorphism ϕ from $LD(G)$ to H and ϕ is then an *H -arc-coloring* or simply an *arc-coloring* of G . Therefore, an oriented arc-coloring ϕ of G must satisfy (i) $\phi(\vec{uv}) \neq \phi(\vec{vw})$ whenever \vec{uv} and \vec{vw} are two consecutive arcs in G , and (ii) $\phi(\vec{vw}) \neq \phi(\vec{xy})$ whenever $\vec{uv}, \vec{vw}, \vec{xy}, \vec{yz} \in A(G)$ with $\phi(\vec{uv}) = \phi(\vec{yz})$. The *oriented chromatic index* of G , denoted by $\chi'_o(G)$, is defined as the smallest order of an oriented graph H such that $LD(G) \rightarrow H$.

The notion of oriented chromatic index can be extended to undirected graphs as follows. The oriented chromatic index $\chi'_o(G)$ of an undirected graph G is the maximum of the oriented chromatic indexes taken over all the orientations of G (an orientation of an undirected graph G is obtained by giving one of the two possible orientations to every edge of G).

In this paper, we are interested in oriented arc-coloring of subcubic graphs, that is graphs with maximum degree at most 3.

Oriented vertex-coloring of subcubic graphs has been first studied in [4] where it was proved that every oriented subcubic graph admits an oriented 16-vertex-coloring. In 1996, Sopena and Vignal improved this result:

Theorem 1 [6] *Every oriented subcubic graph admits an oriented 11-vertex-coloring.*

It is not difficult to see that every oriented graph having an oriented k -vertex-coloring admits a k -arc-coloring (from a k -vertex-coloring f , we obtain a k -arc-coloring g by setting $g(\vec{uv}) = f(u)$ for every arc \vec{uv}). Therefore, every oriented subcubic graph admits an oriented 11-arc-coloring.

We improve this bound and prove the following

Theorem 2 *Every oriented subcubic graph admits a 7-arc-coloring.*

More precisely, we shall show that every oriented subcubic graph admits a homomorphism to QR_7 , a tournament on 7 vertices described in section 3.

Note that Sopena conjectured that every oriented connected subcubic graph admits an oriented 7-vertex-coloring [4].

This paper is organised as follows. In the next section, we introduce the main definitions and notation. In section 3, we described the tournament QR_7 and give some properties of this graph. Finally, Section 4 is dedicated to the proof of Theorem 2.

2 Definitions and notation

In the rest of the paper, oriented graphs will be simply called *graphs*. For a graph G and a vertex v of G , we denote by $d_G^-(v)$ the indegree of v , by $d_G^+(v)$ its outdegree and by $d_G(v)$ its degree. A vertex of degree k (resp. at most k , at least k) will be called a k -vertex (resp. $\leq k$ -vertex, $\geq k$ -vertex). A *source*

vertex (or simply a *source*) is a vertex v with $d^-(v) = 0$ and a *sink vertex* (or simply a *sink*) is a vertex v with $d^+(v) = 0$. A source (resp. sink) of degree k will be called a k -source (resp. a k -sink).

We denote by $N_G^+(v)$, $N_G^-(v)$ and $N_G(v)$ respectively the set of successors of v , the set of predecessors of v and the set of neighbors of v in G . The *maximum degree* and *minimum degree* of a graph G are respectively denoted by $\Delta(G)$ and $\delta(G)$.

We denote by \overrightarrow{uv} the arc from u to v or simply uv whenever its orientation is not relevant (therefore $uv = \overrightarrow{uv}$ or $uv = \overleftarrow{uv}$).

For a graph G and a vertex v of $V(G)$, we denote by $G \setminus v$ the graph obtained from G by removing v together with the set of its incident arcs; similarly, for an arc a of $A(G)$, $G \setminus a$ denotes the graph obtained from G by removing a . These two notions are extended to sets in a standard way: for a set of vertices V' , $G \setminus V'$ denotes the graph obtained from G by successively removing all vertices of V' and their incident arcs, and for a set of arcs A' , $G \setminus A'$ denotes the graph obtained from G by removing all arcs of A' .

The drawing conventions for a configuration are the following: a vertex whose neighbors are totally specified will be black (i.e. vertex of fixed degree), whereas a vertex whose neighbors are partially specified will be white. Moreover, an edge will represent an arc with any of its two possible orientations.

3 Some properties of the tournament QR_7

For a prime $p \equiv 3 \pmod{4}$, the Paley tournament QR_p is defined as the oriented graph whose vertices are the integers modulo p and such that \overrightarrow{uv} is an arc if and only if $v - u$ is a non-zero quadratic residue of p .

For instance, let us consider the tournament QR_7 with $V(QR_7) = \{0, 1, \dots, 6\}$ and $\overrightarrow{uv} \in A(QR_7)$ whenever $v - u \equiv r \pmod{7}$ for $r \in \{1, 2, 4\}$.

This graph has the two following useful properties [1]:

(P_1) Every vertex of QR_7 has three successors and three predecessors.

(P_2) For every two distinct vertices u and v , there exists four vertices w_1, w_2, w_3 and w_4 such that:

- $\overrightarrow{uw_1} \in A(QR_7)$ and $\overleftarrow{vw_1} \in A(QR_7)$;
- $\overrightarrow{uw_2} \in A(QR_7)$ and $\overleftarrow{vw_2} \in A(QR_7)$;
- $\overleftarrow{w_3u} \in A(QR_7)$ and $\overleftarrow{w_3v} \in A(QR_7)$;
- $\overleftarrow{w_4u} \in A(QR_7)$ and $\overleftarrow{vw_4} \in A(QR_7)$.

4 Proof of Theorem 2

Let G be an oriented subcubic graph and C be a cycle in G (C is a subgraph of G). A vertex u of C is a *transitive vertex of C* if $d_C^+(u) = d_C^-(u) = 1$ (therefore $2 \leq d_G(u) \leq 3$).

A cycle C in G is a *special cycle* if and only if:

- (1) every non-transitive vertex of C is a 2-source or a 2-sink in G ;
- (2) C has either exactly 1 transitive vertex or exactly 2 transitive vertices, and in this case, both transitive vertices have the same orientation on C .




Figure 1: Two special cycles

Figure 1 shows two special cycles; the first one has exactly 1 transitive vertex while the second has exactly 2 transitive vertices oriented in the same direction. Vertices s_i , s'_j and t_k are respectively the sinks, sources, and transitive vertices of the special cycles.

Remark 3 Every 2-source (resp. 2-sink) in a special cycle C is necessarily adjacent to a 2-sink (resp. 2-source). This directly follows from the fact that C does not contain two transitive vertices oriented in opposite direction.

We shall denote by $SS_G(C)$ the set of 2-sources and 2-sinks of the cycle C in G .




Figure 2: Graphs with a special cycle

Remark 4 Note that a special cycle may only be connected to the rest of the graph by its transitive vertices (see Figure 2 for an example).

A QR_7 -arc-coloring f of an oriented subcubic graph G is *good* if and only if :

- for every 2-source u , $|C_f^+(u)| = 1$,
- for every 2-sink v , $|C_f^-(v)| = 1$.

Note that if a subcubic graph G admits a good QR_7 -arc-coloring, then for every 2-vertex v of G , $|C_f^+(v)| \leq 1$ and $|C_f^-(v)| \leq 1$.

We first prove the following:

Theorem 5 *Every oriented subcubic graph with no special cycle admits a good QR_7 -arc-coloring.*

We define a partial order \prec on the set of all graphs. Let $n_2(G)$ be the number of ≥ 2 -vertices of G . For any two graphs G_1 and G_2 , $G_1 \prec G_2$ if and only if at least one of the following conditions holds:

- G_1 is a proper subgraph of G_2 ;
- $n_2(G_1) < n_2(G_2)$.

Note that this partial order is well-defined, since if G_1 is a proper subgraph of G_2 , then $n_2(G_1) \leq n_2(G_2)$. The partial order \prec is thus a partial linear extension of the subgraph poset.

In the rest of this section, let H be a counter-example to Theorem 5 which is minimal with respect to \prec .

We shall show in the following lemmas that H does not contain some configurations.

In all the proofs which follow, we shall proceed similarly. We suppose that H contains some configurations and, for each of them, we consider a reduction H' of H with no special cycle such that $H' \prec H$. Therefore, due to the minimality of H , there exists a good QR_7 -arc-coloring f of H' . The coloring f is a partial good QR_7 -arc-coloring of H , that is an arc-coloring of some subset S of $A(H)$ and we show how to extend it to a good QR_7 -arc-coloring of H . This proves that H cannot contain such configurations.

We will extensively use the following proposition:

Proposition 6 *Let \vec{G} be an oriented graph which admits a good QR_7 -arc-coloring. Let \overleftarrow{G} be the graph obtained from \vec{G} by giving to every arc its opposite direction. Then, \overleftarrow{G} admits a good QR_7 -arc-coloring.*

Proof : Let f be a good QR_7 -arc-coloring of \vec{G} . Consider the coloring $f' : V(QR_7) \rightarrow A(\overleftarrow{G})$ defined by $f'(\overleftarrow{uv}) = 6 - f(\overrightarrow{vu})$.

It is easy to see that for every arc $\overrightarrow{uv} \in A(QR_7)$, we have $\overrightarrow{xy} \in A(QR_7)$ for $x = 6 - v$ and $y = 6 - u$. Moreover, the two incident arcs to a 2-source (or a 2-sink) will get the same color by f' since they got the same color by f . \square

Therefore, when considering oriented good QR_7 -arc-coloring of an oriented graph G , we may assume that one arc in G has a given orientation.

The following remark will be extensively used in the following lemmas :

Remark 7 Let G be a graph with no special cycle and $A \subseteq A(G)$ be an arc set. If the graph $G' = G \setminus A$ contains a special cycle C , then at least one of the vertices incident to A is a 2-source or a 2-sink in G' and belongs to $V(C)$, since otherwise C would be a special cycle in G .

Lemma 8 *The graph H is connected.*

Proof : Suppose that $H = H_1 \uplus H_2$ (disjoint union). We have $H_1 \prec H$ and $H_2 \prec H$. The graphs H_1 and H_2 contain no special cycle and then, by minimality of H , H_1 and H_2 admits good QR_7 -arc-colorings f_1 and f_2 respectively that can easily be extended to a good QR_7 -arc-coloring $f = f_1 \cup f_2$ of H . \square

Lemma 9 *The graph H contains no 3-source and no 3-sink.*

Proof : By Proposition 6, we just have to consider the 3-source case. Let u be a 3-source in H and H' be the graph obtained from H by splitting u into three 1-vertices u_1, u_2, u_3 . We have $H' \prec H$ since $n_2(H') = n_2(H) - 1$. Any good QR_7 -arc-coloring of H' is clearly a good QR_7 -arc-coloring of H . \square

Lemma 10 *The graph H contains no 1-vertex.*

Proof : Let u_1 be a 1-vertex in H , v be its neighbor and $N_H(v) = \{u_i, 1 \leq i \leq d_H(v)\}$. By Proposition 6, we may assume $\overrightarrow{u_1 v} \in A(H)$. We consider three subcases.

1. $d_H(v) = 1$.

By Lemma 8, $H = \overrightarrow{u_1 v}$ and obviously, H admits a good QR_7 -arc-coloring.

2. $d_H(v) = 2$.

Let $H' = H \setminus u_1$; we have $H' \prec H$ and H' contains no special cycle by remark 7. By minimality of H , H' admits a good QR_7 -arc-coloring f that can easily be extended to H : if v is a 2-sink, we set $f(\overrightarrow{u_1 v}) = f(\overrightarrow{u_2 v})$; otherwise, we have three available colors for $f(\overrightarrow{u_1 v})$ by Property (P_1)

3. $d_H(v) = 3$.

Let $H' = H \setminus u_1$; we have $H' \prec H$.

If H' contains no special cycle then, by minimality of H , H' admits a good QR_7 -arc-coloring f such that $|C_f^+(v)| \leq 1$. The coloring f can then be extended to H since we have three available colors to set $f(\overrightarrow{u_1 v})$ by property (P_1) .

If H' contains a special cycle C , $v \in C$ and v is a 2-source in H' by Remark 7 and Lemma 9. We may assume w.l.o.g. that u_2 is a 2-sink by Remark 3. Let $N_H(u_2) = \{v, x\}$ and $H'' = H \setminus \{\overrightarrow{v u_2}, \overrightarrow{u_1 v}\}$. We have $H'' \prec H$ and H'' contains no special cycle by Remark 7. By minimality of H , H'' admits a good QR_7 -arc-coloring f that can be extended to H : we set $f(\overrightarrow{v u_2}) = f(\overrightarrow{x u_2})$, and we have at least one available color for $f(\overrightarrow{u_1 v})$ by Property (P_2) . \square

Recall that a *bridge* in a graph G is an edge whose removal increases the number of components of G .

Lemma 11 *The graph H contains no bridge.*

Proof : Suppose that H contains a bridge uv . Let $H \setminus uv = H_1 \uplus H_2$. For $i = 1, 2$, consider $H'_i = H_i + uv$. By Lemma 10, uv is not a dangling arc in H . Moreover $H'_i \prec H$ for $i = 1, 2$. Clearly, the graphs H'_1 and H'_2 have no special cycle and therefore, by minimality of H , they admit good QR_7 -arc-colorings f_1 and f_2 respectively. By cyclically permuting the colors of f_2 if necessary, we may assume that $f_1(uv) = f_2(uv)$. The mapping $f = f_1 \cup f_2$ is then clearly a good QR_7 -arc-coloring of H . \square




Figure 3: Configurations of Lemma 14

Lemma 12 *The graph H contains no 2-sink adjacent to a 2-source.*

Proof : Suppose that H contains a 2-sink v adjacent to a 2-source w . Let $N(v) = \{u, w\}$ and $N(w) = \{v, x\}$. Since H contains no special cycle, u and x are distinct vertices and $\overrightarrow{wx} \notin A(H)$.

Let H' be the graph obtained from $H \setminus \{v, w\}$ by adding \overrightarrow{ux} (if it did not already belong to $A(H)$). We have $H' \prec H$ since $n_2(H') \leq n_2(H) - 2$. Since the vertices u and x are neither 3-sources nor 3-sinks in H by Lemma 9, they are neither 2-sources nor 2-sinks in H' and therefore, by Remark 7, H' contains no special cycle. Hence, by minimality of H , H' admits a good QR_7 -arc-coloring f' that can be extended to H by setting $f(\overrightarrow{uv}) = f(\overrightarrow{wv}) = f(\overrightarrow{wx}) = f(\overrightarrow{ux})$. \square

Lemma 13 *Every 2-source (resp. 2-sink) of H is adjacent to a vertex v with $d^+(v) = 2$ (resp. $d^-(v) = 2$).*

Proof : Suppose that H contains a 2-source u adjacent to two vertices v and w such that $d^+(v) \neq 2$ and $d^+(w) \neq 2$ (by Proposition 6, it is enough to consider this case). Let $H' = H \setminus u$; by hypothesis and by Lemmas 9 and 12, the vertices v and w are such that $d_{H'}^+(v) = d_{H'}^-(v) = d_{H'}^+(w) = d_{H'}^-(w) = 1$. Therefore, the graph H' contains no special cycle by Remark 7. By minimality of H , H' admits a good QR_7 -arc-coloring f that can be extended to H in such a way that $f(\overrightarrow{uv_1}) = f(\overrightarrow{uv_2})$ thanks to Property (P_2) . \square

Recall that we denote by $SS_G(C)$ the set of 2-sources and 2-sinks of the cycle C in G .

Lemma 14 *Let u be a vertex of H and $H' = H \setminus u$. Then H' does not contain a special cycle C with $|N_H(u) \cap SS_{H'}(C)| = 1$.*

Proof : Let $v_1 \in N(u)$ and w.l.o.g., suppose that $H' = H \setminus u$ contains a special cycle C such that $N_H(u) \cap SS_{H'}(C) = \{v_1\}$; by Remark 7, v_1 is a 2-source or a 2-sink in H' and by Proposition 6 we may assume w.l.o.g. that v_1 is a 2-source.

By Remark 3, v_1 is adjacent to a 2-sink v_2 . By Lemma 12, the only pair of adjacent 2-source and 2-sink in H' is v_1, v_2 . Therefore, we have $3 \leq |C| \leq 4$. Let $V(C) = \{v_1, v_2, v_3, v_4\}$ and $v_3 = v_4$ if $|C| = 3$. Moreover v_3 and v_4 are necessarily two transitives vertices of C . Furthermore, we have $\overrightarrow{v_3v_4} \in A(H)$ by Lemma 13 and $\overrightarrow{uv_1} \in A(H)$ by Lemma 9. Then, we have only two possible configurations, depicted in Figure 3.

- If $|C| = 3$ (see Figure 3(a)), consider $H'_1 = H \setminus \overrightarrow{v_1 v_2}$. This graph contains no special cycle by Remark 7 and we have $H'_1 \prec H$. By minimality of H , H'_1 admits a good QR_7 -arc-coloring f that can be extended to H : we first erase $f(\overrightarrow{v_1 v_3})$; then, we can set $f(\overrightarrow{v_1 v_2}) = f(\overrightarrow{v_3 v_2})$ thanks to Property (P_2) and then we have one available color for $f(\overrightarrow{v_1 v_3})$ by Property (P_2) since $f(\overrightarrow{u v_1}) \neq f(\overrightarrow{v_3 v_2})$.
- If $|C| = 4$ (see Figure 3(b)), consider the graph $H'_2 = H \setminus v_2$. We have $H'_2 \prec H$.
 - If H'_2 contains no special cycle, by minimality of H , H'_2 admits a good QR_7 -arc-coloring f that can be extended to H in such a way that $f(\overrightarrow{v_3 v_2}) = f(\overrightarrow{v_1 v_2})$ thanks to Property (P_2) since $f(\overrightarrow{v_4 v_3}) = f(\overrightarrow{y v_3})$.
 - Suppose now that H'_2 contains a special cycle C' . By Remark 7, v_3 belong to C' and by Remark 3, y is a 2-sink. By Lemma 12, the only pair of adjacent 2-source and 2-sink in H' is v_3, y , and therefore $|C'|$ is a special cycle of length 3 or 4. Suppose first that $\{u, v_1, v_4, v_3, y\} \subseteq V(C')$; we thus have $u = y$, that is a contradiction since by hypothesis $N_H(u) \cap SS_{H'}(C) = \{v_1\} \neq \{v_1, v_3\}$. Therefore $V(C') = \{y, v_3, v_4, z\}$, and then $\overrightarrow{z v_4} \in A(H)$. If $|C'| = 3$, we have $y = z$ and in this case, the graph H contains a bridge $\overrightarrow{u v_1}$ that is forbidden by Lemma 11. Therefore, we have $|C'| = 4$ and z is a transitive vertex of C' . Consider in this case the graph $H'_3 = H \setminus v_4$. This graph contains no special cycle since the vertices v_1 and v_3 are two transitive 2-vertices oriented in opposite directions. We have $H'_3 \prec H$ and therefore, by minimality of H , there exists a good QR_7 -arc-coloring f of H'_3 such that $C_f^-(v_1) = \{c_1\}$, $C_f^-(v_2) = \{c_2\}$ and $C_f^+(y) = C_f^+(z) = \{c_3\}$. The mapping f can be extended to H as follows: we can set $f(\overrightarrow{v_4 v_3}) = c_4 \notin \{c_1, c_3\}$ thanks to Property (P_1) . Then, by Property (P_2) , we have one available color for $f(\overrightarrow{v_1 v_4})$ since $c_1 \neq c_4$ and one available color for $f(\overrightarrow{z v_4})$ since $c_3 \neq c_4$.

□

Lemma 15 *The graph H does not contain two adjacent 2-vertices.*

Proof : Suppose that H contains two adjacent 2-vertices v and w . Let $N(v) = \{u, w\}$ and $N(w) = \{v, x\}$ and $H' = H \setminus v$. By Lemma Remark 7 and 14, H contains no special cycle. We have $H' \prec H$ and by minimality of H , H' admits a good QR_7 -arc-coloring f .

We shall consider two cases depending on the orientation of the arcs incident to v and w (by Proposition 6, we may assume that $\overrightarrow{u v} \in A(H)$).

1. v is a 2-sink and w is a transitive vertex.

By Lemma 12, u is not a 2-source in H . We have $|C_f^-(u)| \leq 1$ and then, we can set $f(\overrightarrow{u v}) = f(\overrightarrow{w v})$ thanks to Property (P_2) .

2. v and w are transitive vertices.

By the previous case, u is not a 2-source. We have $|C_f^-(u)| \leq 1$. Thanks to Property (P_1) , we can set $f(\overrightarrow{u v}) \neq f(\overrightarrow{w x})$ and finally, we have one available color for $f(\overrightarrow{v w})$ by Property (P_2) since $f(\overrightarrow{u v}) \neq f(\overrightarrow{w x})$.

□




Figure 4: Configurations of Lemma 16

Lemma 16 *The graph H contains no 2-vertex.*

Proof : Suppose that H contains a 2-vertex u and let $N(u) = \{u_1, u_2\}$. The vertices u_1 and u_2 are 3-vertices by Lemma 15. By Proposition 6, we may assume w.l.o.g. that $\overrightarrow{uu_1} \in A(H)$. Let $H'_1 = H \setminus u$; we have $H'_1 \prec H$.

If H'_1 contains no special cycle, then by minimality of H , H'_1 admits a good QR_7 -arc-coloring f of H'_1 that can be extended to H as follows. If u is a 2-source, we can set $f(\overrightarrow{uu_1}) = f(\overrightarrow{uu_2})$ thanks to Property (P_2) since $|C_f^+(u_1)| \leq 1$ and $|C_f^+(u_2)| \leq 1$. If u is a transitive vertex, we can set $f(\overrightarrow{uu_1}) \notin C_f^-(u_2)$ thanks to Property (P_1) and then we have one available color for $f(\overrightarrow{u_2u})$ by Property (P_2) .

Suppose now that H'_1 contains a special cycle C . By Lemma 14, u_1 and u_2 belongs to C and at least one of them is a 2-source or a 2-sink.

Suppose first that u_1 is a 2-source in H'_1 and u_2 is neither a 2-source nor a 2-sink in H'_1 . Then, since H contains no adjacent 2-vertices by Lemma 15, we have only three possible configurations depicted in Figures 4(a), 4(b) and 4(c).

Clearly, the configuration of Figure 4(a) admits a good QR_7 -arc-coloring. The white vertex of the configuration of Figure 4(b) is a 3-vertex by Lemma 15, but in this case, the graph contains a bridge, that is forbidden by Lemma 11. The white vertex of the configuration of Figure 4(c) is of degree two by Lemma 11 and this configuration clearly admits a good QR_7 -arc-coloring.

Therefore, u_1 and u_2 are either 2-sources or 2-sinks in H'_1 . In this case, since H contains no adjacent 2-vertices by Lemma 15, we have only three possible configurations depicted in Figure 4(d), 4(e) and 4(f).

- Figure 4(d): by Lemma 9, we have $\overrightarrow{u_2u}, \overrightarrow{uu_1} \in A(H)$. Consider the graph $H'_2 = H \setminus \overrightarrow{u_1u_2}$; H'_2 contains no special cycle. Since $H'_2 \prec H$, by minimality of H , H'_2 admits a good QR_7 -arc-coloring f that can be extended to H thanks to Property (P_2) since $f(\overrightarrow{u_2u}) \neq f(\overrightarrow{uu_1})$.
- Figure 4(e): by Lemma 9, we have $\overrightarrow{u_2u}, \overrightarrow{uu_1} \in A(H)$. By Lemma 15, u_4 is a 3-vertex. If $d^-(u_4) = 2$, this configuration is forbidden by Lemma 13. If $d^+(u_4) = 2$, this configuration is also forbidden by Lemma 13.

- Figure 4(f): by Lemma 9, we have $\overrightarrow{uu_1}, \overrightarrow{uu_2} \in A(H)$. Therefore, by Lemma 13, $d^-(u_4) = 2$. Consider $H'_4 = H \setminus \overrightarrow{u_1u_3}$; clearly, H'_4 contains no special cycle. By minimality of H , H'_4 admits a good QR_7 -arc-coloring that can be extended to H as follows. We first erase $f(\overrightarrow{u_2u_4})$ and $f(\overrightarrow{u_4u_3})$; then, thanks to Property (P_2) , we can set $f(\overrightarrow{u_1u_3}) = f(\overrightarrow{u_4u_3})$. Finally, since $f(\overrightarrow{uu_2}) \neq f(u_4u_3)$, we can extend f to a good QR_7 -arc-coloring of H thanks to Property (P_2) .

□




Figure 5: Configurations of Lemma 17

Lemma 17 *The graph H contains no 3-vertex.*

Proof : By Lemmas 10 and 16, H is a 3-regular graph. Let u be a vertex of H with neighbors u_1, u_2 and u_3 . By Lemma 9, u is neither a 3-source nor a 3-sink and therefore, by Proposition 6, we may assume w.l.o.g. that $d^+(u) \geq d^-(u)$. Let $\overrightarrow{u_1u}, \overrightarrow{uu_2}, \overrightarrow{uu_3} \in A(H)$.

If $H'_1 = H \setminus u$ contains no special cycle, by minimality of H , H'_1 admits a good QR_7 -arc-coloring f that can be extended to H as follows. We can set $f(\overrightarrow{u_1u}) \notin C_f^+(u_2) \cup C_f^+(u_3)$ thanks to Property (P_1) . Then, thanks to Property (P_2) , we can extend f to a good QR_7 -arc-coloring of H .

Suppose now that H'_1 contains a special cycle C . The graph H'_1 contains three 2-vertices. Since a special cycle consists in k pairs of 2-sources and 2-sinks, C contains only one pair of adjacent 2-source and 2-sink (w.l.o.g. u_1 and u_2 respectively). Therefore, we have only four possible configurations depicted in Figure 5.

Clearly, the configuration of Figure 5(a) admits a good QR_7 -arc-coloring. The white vertex of the configuration of Figure 5(c) is a 2-vertex by Lemma 11 and it is easy to check that there exists a good QR_7 -arc-coloring of this graph. Consider now the configurations of Figures 5(b) and 5(d) and let $H'_2 = H \setminus \overrightarrow{u_1u_2}$. We have $H'_2 \prec H$ and clearly, H'_2 contains no special cycle. Therefore, by minimality of H , H'_2 admits a good QR_7 -arc-coloring f that can be extended to H thanks to Property (P_2) since for any orientation of H , $C_f^-(u_1) \cap C_f^+(u_2) = \emptyset$. □

Proof of Theorem 2 :

By Lemmas 10, 16 and 17, a minimal counter-example to Theorem 5 does not exist.

We now say that a QR_7 -arc-coloring f of an oriented subcubic graph G is *quasi-good* if and only if for every 2-source u , $|C_f^+(u)| = 1$.

Note that if a subcubic graph admits a quasi-good QR_7 -arc-coloring f , we have $|C_f^+(v)| \leq 1$ for every ≤ 2 -vertex v of G .

We shall then prove Theorem 2 by showing that every subcubic graph admits a quasi-good QR_7 -arc-coloring.

Let H be a minimal counter-example to Theorem 2.

If H contains no special cycle, by Theorem 5, H admits a good QR_7 -arc-coloring which is a quasi-good QR_7 -arc-coloring.

Suppose now that H contains at least one special cycle. By definition, a special cycle contains at least one 2-source. We inductively define a sequence of graphs H_0, H_1, \dots, H_n for $n \geq 0$, and a sequence of vertices u_0, u_1, \dots, u_{n-1} such that:

- $H_0 = H$;
- H_i contains a special cycle, and thus a 2-source u_i for $0 \leq i < n$;
- $H_{i+1} = H_i \setminus u_i$ for $0 \leq i < n$;
- H_n has no special cycle.

By Theorem 5, H_n admits a good QR_7 -arc-coloring, and therefore a quasi-good QR_7 -arc-coloring. Suppose that H_{i+1} admits a quasi-good QR_7 -arc-coloring f_{i+1} for $1 \leq i < n$; we claim that we can extend f_{i+1} to a quasi-good QR_7 -arc-coloring f_i of H_i as follows. To see that, let v_i and w_i be the two neighbors of u_i which are ≤ 2 -vertices in H_{i+1} . Therefore, we have $|C_{f_{i+1}}^+(v_i)| \leq 1$ and $|C_{f_{i+1}}^+(w_i)| \leq 1$ and thanks to Property (P_2) , we can set $f_i(\overrightarrow{u_i v_i}) = f_i(\overrightarrow{u_i w_i})$.

Therefore, any quasi-good QR_7 -arc-coloring of H_n can be extended to $H_0 = H$, that is a contradiction. A minimal counter-example to Theorem 2 does not exist, that completes the proof. \square




Figure 6: Cubic graph G with $\chi'_o(G) = 6$

Currently, we cannot provide an oriented subcubic graph with oriented chromatic index 7. However, the oriented cubic graph G depicted in Figure 6 has oriented chromatic index 6.

Suppose we want to color G with five colors 1, 2, 3, 4, 5. Necessarily the colors of \overrightarrow{vw} , \overrightarrow{xy} and \overrightarrow{zu} are pairwise distinct and we may assume w.l.o.g. that $f(\overrightarrow{vw}) = 1$, $f(\overrightarrow{xy}) = 2$ and $f(\overrightarrow{zu}) = 3$. Clearly, each of the colors 4 and 5 will appear at most once on \overrightarrow{uv} , \overrightarrow{wx} and \overrightarrow{yz} . Therefore, w.l.o.g. we may assume that $f(\overrightarrow{yz}) = 1$, which implies w.l.o.g. that we must set $f(\overrightarrow{ux}) = 4$. Thus, we must set $f(\overrightarrow{yv}) = 5$, and then we have no remaining color to color $f(\overrightarrow{wz})$.

Therefore, we have the following:

Proposition 18 *Let \mathcal{C} be the class of subcubic graphs. Then $6 \leq \chi'_o(\mathcal{C}) \leq 7$.*

References

- [1] O. V. Borodin, A. V. Kostochka, J. Nešetřil, A. Raspaud, and E. Sopena. On the maximum average degree and the oriented chromatic number of a graph. *Discrete Math.*, 206:77–89, 1999.
- [2] B. Courcelle. The monadic second order-logic of graphs VI : on several representations of graphs by relational structures. *Discrete Appl. Math.*, 54:117–149, 1994.
- [3] A. V. Kostochka, E. Sopena, and X. Zhu. Acyclic and oriented chromatic numbers of graphs. *J. Graph Theory*, 24:331–340, 1997.
- [4] E. Sopena. The chromatic number of oriented graphs. *J. Graph Theory*, 25:191–205, 1997.
- [5] E. Sopena. Oriented graph coloring. *Discrete Math.*, 229(1-3):359–369, 2001.
- [6] E. Sopena and L. Vignal. A note on the chromatic number of graphs with maximum degree three. Technical Report 1125-96, LaBRI, Université Bordeaux 1, 1996.
- [7] D. R. Wood. Acyclic, star and oriented colourings of graph subdivisions. *Discrete Mathematics and Theoretical Computer Science*, 7(1):37–50, 2005.