



HAL
open science

Some metrics on Teichmüller spaces of surfaces of infinite type

Lixin Liu, Athanase Papadopoulos

► **To cite this version:**

Lixin Liu, Athanase Papadopoulos. Some metrics on Teichmüller spaces of surfaces of infinite type. Transactions of the American Mathematical Society, 2011, 363 (8), pp.4109-4134. 10.1090/S0002-9947-2011-05090-7 . hal-00309330v2

HAL Id: hal-00309330

<https://hal.science/hal-00309330v2>

Submitted on 22 Jul 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SOME METRICS ON TEICHMÜLLER SPACES OF SURFACES OF INFINITE TYPE

LIXIN LIU AND ATHANASE PAPADOPOULOS

ABSTRACT. Unlike the case of surfaces of topologically finite type, there are several different Teichmüller spaces that are associated to a surface of topological infinite type. These Teichmüller spaces first depend (set-theoretically) on whether we work in the hyperbolic category or in the conformal category. They also depend, given the choice of a point of view (hyperbolic or conformal), on the choice of a distance function on Teichmüller space. Examples of distance functions that appear naturally in the hyperbolic setting are the length spectrum distance and the bi-Lipschitz distance, and there are other useful distance functions. The Teichmüller spaces also depend on the choice of a basepoint. The aim of this paper is to present some examples, results and questions on the Teichmüller theory of surfaces of infinite topological type that do not appear in the setting the Teichmüller theory of surfaces of finite type. In particular, we point out relations and differences between the various Teichmüller spaces associated to a given surface of topological infinite type.

AMS Mathematics Subject Classification: 32G15 ; 30F30 ; 30F60.

Keywords: Teichmüller space, infinite-type surface, Teichmüller metric, quasiconformal metric, length spectrum metric, bi-Lipschitz metric.

Acknowledgement: L. Liu was partially supported by the NSF of China.

1. INTRODUCTION

Let X be a connected oriented surface of infinite topological type. Unlike the case of a surface of finite type, there are several Teichmüller spaces associated to X . Such a Teichmüller space can be defined as a space of equivalence classes of marked hyperbolic surfaces homeomorphic to X , the marking being a homeomorphism between a base hyperbolic structure on X and a hyperbolic surface homeomorphic to X . To define a distance function (and then, a topology) on Teichmüller space, we ask that this marking be either length spectrum bounded or a bi-Lipschitz homeomorphism. Depending on such a choice, the Teichmüller spaces that we obtain are setwise different. Taking a different point of view, a Teichmüller space associated to X can also be defined as a space of equivalence classes of marked Riemann surfaces homeomorphic to X , the marking being here a quasiconformal homeomorphism between a base Riemann surface structure on X and a Riemann surface homeomorphic to X . We shall be more precise below. Unlike the case of surfaces of finite type, the various definitions of Teichmüller spaces are not equivalent and the resulting Teichmüller space strongly depends on the point of view (hyperbolic or conformal), on the distance function we choose on the set of equivalence classes of marked surfaces, and on the choice of a base surface. The aim of this paper is to review some of the Teichmüller spaces associated to X , precisely when X has infinite type, and to point out some differences, some relations and some analogies between these various Teichmüller spaces.

The plan of this paper is as follows.

Date: July 22, 2009.

In Section 2, we study the length spectrum Teichmüller spaces associated to a surface of infinite type. This study involves the consideration of the length spectrum distance between hyperbolic surfaces.

In Section 3, we study the bi-Lipschitz Teichmüller spaces associated to a hyperbolic surface of infinite type, and this involves the consideration of the bi-Lipschitz distance between hyperbolic surfaces.

In Section 4, we consider the quasiconformal point of view. We define the quasiconformal Teichmüller space of a Riemann surface of infinite type and the quasiconformal metric on this space. This space is related to the bi-Lipschitz Teichmüller space. We give a sufficient condition under which the three Teichmüller spaces coincide as sets.

Each section contains basic definitions and results, and some open questions.

We now fix some notation which will be used throughout the paper.

We consider a connected orientable surface X of genus $g \geq 0$, with $p \geq 0$ punctures and $n \geq 0$ boundary components, and we assume that $g + n + p$ is countably infinite. We note right away that even in the case of surfaces with no punctures and no boundary components, two surfaces of countably infinite genus are not necessarily homeomorphic. (For instance, the surfaces represented in Figures 1, 2 and 3 below are pairwise non-homeomorphic.) This contrasts with the case of surfaces of finite type in which the triple (genus, number of punctures, number of boundary components) determines the surface up to homeomorphism. The classification of noncompact surfaces has a long history, and for this subject we refer the reader to the work of Kerékjártó [15], to the later paper by Richards [26], and to recent update by Prishlyak and Mischenko [25]. In particular, there exist invariants for surfaces of infinite type. (We note that these invariants are, unlike those of surfaces of finite type, not discrete.)

Using Zorn's Lemma, on any topological surface X as above, we can find a countable family of disjoint simple closed curves C_i , $i = 1, 2, \dots$ such that each connected component of $S - \cup_{i=1}^{\infty} C_i$ is homeomorphic to a pair of pants, that is, a sphere with three holes. Here, a hole is either a puncture (that is, a point removed from the surface) or an open disk removed. In the latter case the surface has a boundary component at the hole.

All the hyperbolic structures that we consider on X will be complete, such that each boundary component of this surface a closed geodesic and each puncture of this surface has a neighborhood isometric to a cusp, that is, to the quotient of a region of the form $\{(x, y) \mid x > a\}$ in the upper half-space model of the hyperbolic plane \mathbb{H}^2 , with $a > 0$, by the transformation $z \mapsto z + 1$. The neighborhood of a cusp is biholomorphically equivalent to a punctured disk in \mathbb{C} .

For future reference, we reformulate our requirements as follows:

(\star) The surface X admits a pants decomposition \mathcal{P} with a countable number of pair of pants. The hyperbolic structures that we consider on X are all metrically complete, and they have the property that if we replace each boundary component of a pair of pants in the decomposition \mathcal{P} by the closed geodesic in its homotopy class, then each pair of pants becomes a sphere with three holes, a hole being either a boundary component which is a closed geodesic (and this can hold when the hole is either a boundary component of X or a closed curve in the pants decomposition \mathcal{P}) or a cusp (and this holds when the hole is a puncture of X). In the case where the surface is just a Riemann surface (i.e. not necessarily equipped with a hyperbolic metric), then, we assume that the unique complete hyperbolic metric on $X \setminus \partial X$ that uniformizes the complex structure on that surface, after cutting the flares if such cylindrical ends exist, satisfies the above conditions.

We point out right away that the completeness of a hyperbolic surface satisfying Property (\star) is not a redundant property. This contrasts with the case of hyperbolic surfaces of finite type with geodesic boundary, where a surface is complete if and only if the neighborhood of each puncture is a cusp. Examples of non-complete infinite type hyperbolic surfaces with no punctures and no boundary components have been given by Basmajian in [4].

It is easy (although not completely trivial) to prove that any hyperbolic surface satisfying (\star) has infinite diameter.

A large part of the classical quasiconformal theory of Teichmüller space (as started by Teichmüller, and developed by Ahlfors and Bers) is valid for surfaces of infinite type. In contrast, Thurston's surface theory does not easily extend to surfaces of infinite type. For instance, on any surface of infinite type, we can find an infinite sequence of homotopy classes of simple closed curves which are disjoint, pairwise nonhomotopic and nonhomotopic to a point, and such a sequence does not converge in any reasonable sense to a homotopy class of measured foliations.

We shall consider ordered pairs (f, H) , where H is a hyperbolic metric on a surface S homeomorphic to X , and $f : X \rightarrow S$ a homeomorphism. Such a pair (f, H) is called a *marked hyperbolic surface* (in general, with a base surface X being understood). The homeomorphism f is called the *marking* of H , or of S . We shall also consider a hyperbolic metric on X as a marked hyperbolic surface, taking as marking the identity map. Finally, we shall also talk about markings $f : H_0 \rightarrow H$ between hyperbolic surfaces, when considering the hyperbolic surface H_0 on S as a basepoint in our space.

A *simple closed curve* on X is a one-dimensional submanifold of X homeomorphic to a circle. We shall call a curve on X *essential* if it is a simple closed curve which is neither homotopic to a point nor to a puncture (but it can be homotopic to a boundary component).

We let $\mathcal{S} = \mathcal{S}(X)$ be the set of homotopy classes of essential curves on X .

The *intersection number* $i(\alpha, \beta)$ of two elements α and β in \mathcal{S} is the minimum number of intersection points of two closed curves representing the classes α and β .

Given a hyperbolic structure H on X and given a homotopy class α of essential curves on X , we denote by $l_H(\alpha)$ the length of the closed H -geodesic on X in the class α .

Finally, we note that the Teichmüller spaces that we consider in this paper are, in the classical terminology, "reduced Teichmüller spaces". We shall recall the definitions below.

We thank Daniele Alessandrini and the referee for carefully reading the original manuscript and for their very useful remarks.

2. THE LENGTH SPECTRUM TEICHMÜLLER SPACES

Throughout this section, H_0 is a fixed hyperbolic metric on the surface X called the base hyperbolic structure. With this, a marking $f : X \rightarrow S$ (where S is a hyperbolic surface) becomes a homeomorphism between hyperbolic surfaces. Given a homeomorphism f between two surfaces S and S' equipped with hyperbolic metrics H and H' respectively, we say that f is *length-spectrum bounded* if the following holds:

$$(1) \quad K(f) = \sup_{\alpha \in \mathcal{S}(H)} \left\{ \frac{l_{H'}(f(\alpha))}{l_H(\alpha)}, \frac{l_H(\alpha)}{l_{H'}(f(\alpha))} \right\} < \infty.$$

We shall call the quantity $K(f)$ that appears in (1) the *length-spectrum constant* of f . Note that this quantity only depends on the homotopy class of f .

We consider the collection of marked hyperbolic structures (f, H) relative to the base structure H_0 and where the marking $f : H_0 \rightarrow H$ is length-spectrum bounded. Given two such marked hyperbolic structures (f, H) and (f', H') , we write $(f, H) \sim (f', H')$ if there exists an isometry (or, equivalently, a length spectrum preserving homeomorphism, see Proposition 2.2 below) $f'' : H \rightarrow H'$ which is homotopic to $f' \circ f^{-1}$. We note that all the homotopies of a surface that we consider in this paper preserve the punctures and preserve setwise the boundary components of the surface at all times. The relation \sim is an equivalence relation on the set of length-spectrum bounded marked hyperbolic surfaces (f, H) , based at (X, H_0) .

Definition 2.1. The *length-spectrum Teichmüller space* $\mathcal{T}_{ls}(H_0)$ is the space of \sim -equivalence classes $[f, H]$ of length-spectrum bounded marked hyperbolic surfaces (f, H) . The *basepoint* of this Teichmüller space is the equivalence class $[\text{Id}, H_0]$.

We note that the fact that we do not ask our homotopies to preserve pointwise the boundary of the surface corresponds to working with what is usually called the *reduced* Teichmüller space, instead of Teichmüller space. (In the latter case, the homotopies that define the equivalence relation are required to induce the identity map on each boundary component.) There is a substantial difference between the two theories. For instance, in the quasiconformal setting that we consider in Section 4 below, the Teichmüller space of the unit disk in \mathbb{C} is, in the non-reduced theory, infinite-dimensional (and it is called the *universal Teichmüller space*), whereas in the reduced theory, this space is reduced to a point. Of course, for surfaces that do not have boundary components, the reduced and non-reduced Teichmüller spaces coincide. We also note that in conformal geometry, dealing with boundary points of a Riemann surface generally involves working with the Poincaré metric and its boundary theory, that is, using hyperbolic geometry.

Since all the Teichmüller spaces that we use in this paper are reduced, we shall use, for simplicity, the terminology *Teichmüller space* instead of *reduced Teichmüller space*.

The topology of $\mathcal{T}_{ls}(H_0)$ is induced by the *length-spectrum* metric d_{ls} , defined by taking the distance $d_{ls}([f, H], [f', H'])$ between two points in $\mathcal{T}_{ls}(H_0)$ represented by two marked hyperbolic surfaces (f, H) and (f', H') to be

$$d_{ls}([f, H], [f', H']) = \frac{1}{2} \log K(f' \circ f^{-1}).$$

(It may be useful to recall here that the length-spectrum constants of length-spectrum bounded homeomorphism only depends on the homotopy class of such a homeomorphism.)

The fact that the function d_{ls} satisfies the properties of a metric is straightforward, except perhaps for the separation axiom, which is a consequence of the following:

Proposition 2.2. *Let H and H' be two hyperbolic metrics on a surface X and suppose that $f : (X, H) \rightarrow (X, H')$ is a homeomorphism whose length-spectrum constant $K(f)$ equals 1. Then, f is isotopic to an isometry.*

Proof. From the hypothesis, it follows that for each α in $\mathcal{S}(X)$ we have $l_H(\alpha) = l_{H'}(f(\alpha))$. We show that f is isotopic to an isometry. Consider a sequence K_i , $i = 0, 1, \dots$ of finite type subsurfaces with boundary satisfying $K_i \subset K_{i+1}$ for all $i \geq 0$ and $\cup_{i=0}^{\infty} K_i = X$. For each $i \geq 0$, let $L_i \subset X$ (respectively $M_i \subset X$) be the hyperbolic subsurface of X isotopic to K_i and having geodesic boundary, with respect to the hyperbolic metric H (respectively H'). Then, for each $i \geq 0$, f is homotopic to a map $f'_i : X \rightarrow X$ which sends homeomorphically L_i to M_i . From a well known result for surfaces of finite type, since f'_i preserves the length spectrum

and since each L_i is of finite type, the restriction of f'_i to each L_i is isotopic to an isometry $f''_i : L_i \rightarrow M_i$. Now an isometry between two surfaces of finite type is unique in its isotopy class. Thus, for each $i \geq 1$, the restriction of f'_i to L_{i-1} is equal to f''_{i-1} . By taking the union of the maps f''_i , we obtain an isometry from (X, H) to (X, H') . \square

The length function $\alpha \mapsto l_H(\alpha)$ on $\mathcal{S}(X)$ associated to a hyperbolic structure H on X induces a length function $\alpha \mapsto l_x(\alpha)$ associated to an element x of the Teichmüller space $\mathcal{T}_{ls}(H_0)$. The first observation is the following:

Proposition 2.3. *For any base hyperbolic metric H_0 on X and for any α in \mathcal{S} , the map $x \mapsto \log l_x(\alpha)$ defined on $\mathcal{T}_{ls}(H_0)$ is 2-Lipschitz.*

Proof. Let x and x' be two points in $\mathcal{T}_{ls}(H_0)$, represented by marked surfaces (f, H) and (f', H') respectively. Then, for any α in \mathcal{S} and for any length-spectrum bounded homeomorphism $f'' : H \rightarrow H'$ with associated length-spectrum constant $K(f'')$, we have

$$\begin{aligned} |\log l_x(\alpha) - \log l_{x'}(\alpha)| &= \left| \log \frac{l_{H'}(\alpha)}{l_H(\alpha)} \right| \\ &\leq \max \left\{ \log \frac{l_{H'}(\alpha)}{l_H(\alpha)}, \log \frac{l_H(\alpha)}{l_{H'}(\alpha)} \right\} \\ &\leq \log K(f''). \end{aligned}$$

Taking the infimum over all length-spectrum bounded homeomorphisms $f'' : H \rightarrow H'$ in the homotopy class of $f' \circ f^{-1}$, we obtain

$$|\log l_x(\alpha) - \log l_{x'}(\alpha)| \leq 2d_{ls}(x, x').$$

\square

The next observation is the existence of pairs of marked hyperbolic structures on X which are not related by any length-spectrum bounded homeomorphism. This will show that two Teichmüller spaces $\mathcal{T}_{ls}(H_0)$ and $\mathcal{T}_{ls}(H_1)$ based at two different hyperbolic structures H_0 and H_1 on X are in general different as sets.




FIGURE 1.

Example 2.4. Consider the surface drawn in Figure 1, where C_1, C_2, \dots are the curves in the homotopy classes represented. We equip this surface with two hyperbolic metrics H_0 and H_1 such that for all $i = 1, 2, \dots$, $l_{H_0}(C_i) = 1$ and $l_{H_1}(C_i) = 1/i$. The metrics H_0 and H_1 have the further property that any closed ball of radius 1 on the surface is contained in a finite number of pairs of pants (of the given decomposition), and therefore it is compact. Thus, by the theorem of Hopf-Rinow, the two metrics are complete. It is clear that the hyperbolic structures H_0 and H_1 are not length-spectrum equivalent, and therefore $H_1 \notin \mathcal{T}_{ls}(H_0)$. In particular, $\mathcal{T}_{ls}(H_0) \neq \mathcal{T}_{ls}(H_1)$.

Example 2.5. As a different example of the same phenomenon, we take X to be the surface represented in Figure 2 and we let T be the step-1 right-translation suggested in this figure, transforming C_i into C_{i+1} for every i in \mathbb{Z} . Consider a hyperbolic structure H_0 on X that is translation-invariant. In other words, we suppose that all the two-holed tori that are in the complement of the curves C_i are isometric, and that the twist parameters about these curves are all equal. For each $i = 1, 2, \dots$, let τ_i be the i -th power of the positive Dehn twist along C_i . Let $f = \tau_1 \circ \tau_2 \circ \tau_3 \circ \dots$ be the infinite composition of the τ_i 's and let H_1 be the hyperbolic structure $f(H_0)$. By using the Hopf-Rinow theorem as in Example 2.4 above, we can see that the metrics H_0 and H_1 are complete. Let D_0 be an essential simple closed curve on X satisfying $i(C_0, D_0) > 0$ and $i(C_i, D_0) = 0$ for all $i \neq 0$. For each i in \mathbb{Z} , let $D_i = T^i(D_0)$. Then, we have

$$\frac{l_{H_1}(D_i)}{l_{H_0}(D_i)} \rightarrow \infty$$

as $i \rightarrow \infty$. Therefore, H_0 and H_1 are not related by a length-spectrum bounded homeomorphism. Hence, $H_1 \notin \mathcal{J}_{ls}(H_0)$, and again we have $\mathcal{J}_{ls}(H_0) \neq \mathcal{J}_{ls}(H_1)$.

More generally, we have the following:

Proposition 2.6. *Given a hyperbolic structure H_0 on an infinite type surface X , there exists a hyperbolic structure H_1 on X such that H_0 and H_1 are not length-spectrum equivalent. In particular, $\mathcal{J}_{ls}(H_0) \neq \mathcal{J}_{ls}(H_1)$.*

Proof. Consider an infinite collection C_i , $i = 1, 2, \dots$ of disjoint pairwise non-homotopic essential curves on X . Up to taking a subsequence of (C_i) , we can find, for each $i = 1, 2, \dots$, a curve D_i satisfying $i(C_i, D_i) > 0$ and $i(C_i, D_j) = 0$ for all $j \neq i$. For each $i = 1, 2, \dots$, we take $\tau_i : X \rightarrow X$ to be a power of a positive Dehn twist along C_i satisfying $l_{H_0}(\tau_i(D_i)) > i$. Let f be the infinite composition $f = \tau_1 \circ \tau_2 \circ \dots$. Then,

$$\frac{l_{H_0}(f(D_i))}{l_{H_0}(D_i)} \rightarrow \infty$$

as $i \rightarrow \infty$. This shows that the hyperbolic structure $f(H_0)$ is not related to H_0 by a length-spectrum bounded homeomorphism. \square

We denote by $\text{Hom}(X)$ the group of self-homeomorphisms of X , and by $\text{Hom}_{ls}(H_0)$ the subgroup of $\text{Hom}(X)$ consisting of self-homeomorphisms that are length-spectrum bounded with respect to the metric H_0 . We observe that in general $\text{Hom}_{ls}(H_0) \neq \text{Hom}(X)$. This can be seen in the following example.

Example 2.7. Let X be the topological surface represented in Figure 3. We consider a homeomorphism f obtained by applying a $2\pi/3$ -counterclockwise rotation suggested by the picture. For $i = 1, 2, \dots$, the curves C_i , C'_i and C''_i are in the homotopy classes represented in this figure. We choose a hyperbolic metric H_0 on X such that for all $i = 1, 2, \dots$, $l_{H_0}(C_i) = 1/i$, $l_{H_0}(C'_i) = i$ and $l_{H_0}(C''_i) = i^2$. By using, as in the previous examples, the theorem of Hopf-Rinow, we can insure that




FIGURE 2.

this metric is complete. It is clear that f is not homotopic to a length-spectrum bounded homeomorphism.

More generally, we have the following.

Proposition 2.8. *For any hyperbolic structure H_0 on a surface X of infinite type, we have $\text{Hom}_{l_s}(H_0) \subsetneq \text{Hom}(X)$.*

Proof. The homeomorphism of $f : X \rightarrow X$ constructed in the proof of Proposition 2.6 is not in $\text{Hom}_{l_s}(H_0)$. \square

We let $\text{MCG}(X)$ denote the quotient of the group $\text{Hom}(X)$ by the normal subgroup consisting of orientation-preserving homeomorphisms that are homotopic to the identity. (Recall that our homotopies preserve setwise the boundary components, at each time.) The group $\text{MCG}(X)$ is the *mapping class group* of X .

Definition 2.9 (The length-spectrum mapping class group). The *length-spectrum mapping class group* of X relative to the hyperbolic metric H_0 , which we denote by $\text{MCG}_{l_s}(H_0)$, is the subgroup of $\text{MCG}(X)$ consisting of homotopy classes of homeomorphisms that are length-spectrum bounded with respect to the metric H_0 .

It follows from Proposition 2.8 that for any hyperbolic structure H_0 on a surface X of infinite type, we have $\text{MCG}_{l_s}(H_0) \subsetneq \text{MCG}(X)$.

The group $\text{MCG}_{l_s}(H_0)$ acts on the Teichmüller space $\mathcal{T}_{l_s}(H_0)$ by the quotient of the action of homeomorphisms on marked hyperbolic surfaces by right-composition by the inverse. More precisely, for each marked hyperbolic surface (f, H) and for each homeomorphism $g : (X, H_0) \rightarrow (X, H)$ which is length-spectrum bounded, we set

$$(2) \quad g \cdot (f, H) = (f \circ g^{-1}, H).$$

This action preserves the metric d_{l_s} on $\mathcal{T}_{l_s}(H_0)$.

The following is a natural question:

Question 2.10. Given any hyperbolic metric on an infinite-type surface, is the natural map from $\text{MCG}_{l_s}(H_0)$ into the isometry group $\text{Isom}(\mathcal{T}_{l_s}(H_0), d_{l_s})$ of the




FIGURE 3.

metric space $(\mathcal{T}_{l_s}(H_0), d_{l_s})$ an isomorphism? Is it an isomorphism onto an index-2 subgroup? (The reason for talking about an index-two subgroup is that one may have to extend the image of $\text{MCG}_{l_s}(H_0)$ in $\text{Isom}(\mathcal{T}_{l_s}(H_0), d_{l_s})$ by the homotopy class of an orientation-reversing length-spectrum bounded homeomorphism of (X, H_0) , if such a homeomorphism exists.)

Remark 2.11. The study of the image of $\text{MCG}_{l_s}(H_0)$ in the isometry group $\text{Isom}(\mathcal{T}_{l_s}(H_0), d_{l_s})$ can also be asked in the case of finite-type surfaces, and it does not seem to have been settled. We recall that in the case of a finite-type hyperbolic surface S , we have $\text{MCG}_{l_s}(S) \simeq \text{MCG}(S)$, and that a famous result by Royden, completed by Earle and Kra, says that the natural homomorphism from the extended mapping-class group of S (which is an order-two extension of $\text{MCG}(S)$) into the isometry group of $\mathcal{T}(S)$ equipped with the Teichmüller metric is an isomorphism, except for some finite set of surfaces of small genus and small number of boundary components (see [27] and [7]). For surfaces of finite type, Li and Liu (see [17] and [18]) proved that the length-spectrum metric on Teichmüller space defines the same topology as the Teichmüller metric, but of course this result is not sufficient to prove that the result of Royden and Earle-Kra also applies to the length-spectrum metric.

We denote by $\text{MCG}_f(X)$ the subgroup of $\text{MCG}(X)$ consisting of the homotopy classes of homeomorphisms supported on a subsurface of X of finite topological type.

Proposition 2.12. *We have a natural inclusion $\text{MCG}_f(X) \subset \text{MCG}_{l_s}(H_0)$.*

Proof. Since every element of the mapping class group of a surface of finite topological type is a finite composition of Dehn twists, it suffices to show that a Dehn twist about a simple closed curve in X is length spectrum bounded.

Let τ_β be a Dehn twist about a simple closed curve β and let α be an arbitrary element of $\mathcal{S}(X)$.

If $i(\alpha, \beta) = 0$, then $\alpha = \tau_\beta(\alpha)$ and $l_{H_0}(\alpha) = l_{H_0}(\tau_\beta(\alpha))$.

If $i(\alpha, \beta) \neq 0$, then, by the Collar Lemma (cf. [3]), there exists a constant t_0 , depending only on the hyperbolic length $l_{H_0}(\beta)$ such that the cylinder in X defined as a $t_0/2$ -neighborhood of the closed geodesic β is embedded in X . Since α traverses this cylinder $i(\alpha, \beta)$ times, we have, $l_{H_0}(\alpha) \geq i(\alpha, \beta)t_0$. Furthermore, from the definition of a Dehn twist, we have

$$l_{H_0}(\tau_\beta(\alpha)) \leq l_{H_0}(\alpha) + i(\alpha, \beta)l_{H_0}(\beta).$$

This gives

$$\frac{l_{H_0}(\tau_\beta(\alpha))}{l_{H_0}(\alpha)} \leq 1 + \frac{l_{H_0}(\beta)}{t_0}.$$

Similarly, we have

$$\frac{l_{H_0}(\alpha)}{l_{H_0}(\tau_\beta(\alpha))} \leq 1 + \frac{l_{H_0}(\beta)}{t_0}.$$

This implies that

$$\log \sup_{\alpha \in \mathcal{S}(X)} \left\{ \frac{l_{H_0}(\tau_\beta(\alpha))}{l_{\phi(H_0)}(\alpha)}, \frac{l_{\phi(H_0)}(\alpha)}{l_{H_0}(\tau_\beta(\alpha))} \right\} < \infty.$$

Thus, τ_β is length spectrum bounded. □

Note that the inclusion in Proposition 2.12 is in general strict.

Proposition 2.13. *Let H be a hyperbolic metric on X such that there exists a sequence α_n , $n = 1, 2, \dots$ of elements in $\mathcal{S}(X)$ represented by disjoint simple closed curves satisfying $l_H(\alpha_n) \rightarrow 0$ as $n \rightarrow \infty$. For every $n = 1, 2, \dots$, let τ_n be the positive Dehn twist about α_n . Then,*

$$d_{ls}(\tau_n(H), H) \rightarrow 0.$$

Proof. Note that $l_{\tau_n(H)}(\alpha) = l_H(\tau_n^{-1}(\alpha))$. We need to prove that

$$(3) \quad \log \sup_{\alpha \in \mathcal{S}(X)} \left\{ \frac{l_H(\tau_n^{-1}(\alpha))}{l_H(\alpha)}, \frac{l_H(\alpha)}{l_H(\tau_n^{-1}(\alpha))} \right\} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Let α be an element of $\mathcal{S}(X)$ and let n be a positive integer.

If $i(\alpha, \alpha_n) = 0$, then $\alpha = \tau_n^{-1}(\alpha)$ and $l_H(\tau_n^{-1}(\alpha)) = l_H(\alpha)$.

Assume that $i(\alpha, \alpha_n) \neq 0$ and let $\varepsilon_n = l_H(\alpha_n)$. By a version of the Collar Lemma [3], there exists $B_\alpha > 0$ such that

$$l_H(\alpha) = i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha.$$

From the definition of a Dehn twist, we have

$$\begin{aligned} i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha - i(\alpha, \alpha_n) \varepsilon_n &\leq l_H(\tau_n^{-1}(\alpha)) \\ &\leq i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha + i(\alpha, \alpha_n) \varepsilon_n. \end{aligned}$$

Then we have

$$\begin{aligned} \frac{l_H(\tau_n^{-1}(\alpha))}{l_H(\alpha)} &\leq \frac{i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha + i(\alpha, \alpha_n) \varepsilon_n}{i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha} \\ &= 1 + \frac{i(\alpha, \alpha_n) \varepsilon_n}{i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha} \\ &\leq 1 + \frac{i(\alpha, \alpha_n) \varepsilon_n}{i(\alpha, \alpha_n) |\log \varepsilon_n|} \\ &\leq 1 + \frac{\varepsilon_n}{|\log \varepsilon_n|}. \end{aligned}$$

In the same way, we can show that

$$\frac{l_H(\alpha)}{l_H(\tau_n^{-1}(\alpha))} \leq 1 + \frac{\varepsilon_n}{|\log \varepsilon_n|}.$$

This proves that

$$\sup_{\alpha \in \mathcal{S}(X)} \left\{ \frac{l_H(\tau_n^{-1}(\alpha))}{l_H(\alpha)}, \frac{l_H(\alpha)}{l_H(\tau_n^{-1}(\alpha))} \right\} \rightarrow 1 \text{ as } n \rightarrow \infty,$$

which implies (3). \square

We shall say that the action of a group on a topological space is *discrete* if the orbit of any point is discrete.

As a consequence of Proposition 2.13, we have the following.

Corollary 2.14. *There exist hyperbolic structures H such that the action of the group $\text{MCG}_{ls}(H)$ on the Teichmüller space $\mathcal{T}_{ls}(H)$ equipped with the metric d_{ls} is not discrete*

Corollary 2.15. *A necessary condition for the action of the group $\text{MCG}_{ls}(H)$ on the Teichmüller space $\mathcal{T}_{ls}(H)$ equipped with the metric d_{ls} to be discrete is that there exists a constant $\epsilon > 0$, such that for any element α in $\mathcal{S}(X)$, $l_H(\alpha) \geq \epsilon$.*

Question 2.16. Find necessary and sufficient conditions on a hyperbolic structure H on a surface of infinite type such that the action of the group $\text{MCG}_{l_s}(H)$ on the Teichmüller space $\mathcal{T}_{l_s}(H)$ equipped with the metric d_{l_s} is discrete. (It is known that for surfaces of finite analytical type, the action is always discrete.)

Remark 2.17. It is known that for surfaces of infinite analytical type, the action of the quasiconformal mapping class group $\text{MCG}_{qc}(H)$ on the quasiconformal Teichmüller space $\mathcal{T}_{qc}(H_0)$ is in general not discrete (see Section 4 below for the definitions, and [22], [12] and [13] for work on this question).

Proposition 2.6 implies that for every hyperbolic metric H_0 on an infinite-type surface, we have $\text{MCG}_{l_s}(H_0) \neq \text{MCG}(X)$.

Example 2.18 (A noncompact closed ball). Let X be the topological surface represented in Figure 2. We let $f : X \rightarrow X$ be the homeomorphism defined as the step-one translation to the right suggested by this picture, and we equip X with a hyperbolic metric H which is invariant by the homeomorphism f . In particular, the homotopy classes of curves C_n satisfy $C_n = f^n(C_0)$ for every $n \in \mathbb{Z}$.

Let $x \in \mathcal{T}_{l_s}(H)$ denote the equivalence class of the marked hyperbolic surface (Id, H) . For each $n \in \mathbb{Z}$, let τ_n denote the positive Dehn twist along C_n and let $x_n \in \mathcal{T}_{l_s}(H)$ denote the equivalence class of the marked surface (τ_n, H) .

From the translation-invariance of H , the distance $d_{l_s}(x, x_n)$ does not depend on n . Thus, all the points x_n are contained in a geometric ball B centered at x and of a fixed radius.

We claim that there exists a constant $K > 0$ such that

$$(4) \quad \forall m \neq n \in \mathbb{Z}, \quad d_{l_s}(x_m, x_n) \geq K.$$

Indeed, let α_0 be the homotopy class of an essential curve satisfying $i(\alpha_0, C_0) \neq 0$ and $i(\alpha_0, C_n) = 0$ for all $n \neq 0$ and for all m in \mathbb{Z} , let α_m denote the homotopy class $f^m(\alpha_0)$. Then, for all integers $m \neq n$, the value

$$\max \left\{ \log \frac{l_{H_m}(\alpha_m)}{l_{H_n}(\alpha_m)}, \log \frac{l_{H_n}(\alpha_m)}{l_{H_m}(\alpha_m)} \right\}$$

is independent of m and n . Calling this constant K , we have (4).

Property (4) implies that the sequence x_n does not have a convergent subsequence. Therefore the closed ball B is not compact.

We recall that a metric space is called *proper* if every closed ball is compact. From the preceding example, there exist hyperbolic structures H such that the Teichmüller space $\mathcal{T}_{l_s}(H)$ is not proper. We shall study properness in more generality in Proposition 2.20 below. For that, we shall use the following.




FIGURE 4.

Let F be a compact sphere with four holes, equipped with a hyperbolic structure H with geodesic boundary and suppose that F is decomposed into a union of two hyperbolic pairs of pants, P_1 and P_2 . (See Figure 4 in which β is a boundary component of P_1 and of P_2 .) We assume that there exists a positive constant M such that each boundary component C of either of the two pairs of pants P_1 and P_2 , including the curve β itself, satisfies

$$\frac{1}{M} \leq l_H(C) \leq M.$$

We let α be a simple closed geodesic on F defined as follows. We first take, in each of the pairs of pants P_1 and P_2 the shortest geodesic segment joining β to itself and not homotopic to a point relative to the curve β (this shortest geodesic segment is called the *seam* of the pair of pants). Then, we join together the endpoints of these two seams by using geodesic segments contained in β , producing a closed curve in F . Then, the curve α is the simple closed geodesic homotopic to that curve. (There are two possibilities for joining the endpoints of the two seams, namely by turning “to the left” or “to the right” along β , and making such a choice will not affect the result we prove in the next lemma.)

Lemma 2.19. *With the above notation, there exists a constant $K > 1$ that depends only on M such that if τ denotes the positive Dehn twist along β , we have*

$$\max \left(\frac{l_H(\tau^{-1}(\alpha))}{l_H(\alpha)}, \frac{l_H(\alpha)}{l_H(\tau^{-1}(\alpha))}, \frac{l_H(\tau(\alpha))}{l_H(\alpha)}, \frac{l_H(\alpha)}{l_H(\tau(\alpha))} \right) > K.$$

Proof. The isometry type of the sphere with four holes F is determined by the values of six parameters, five of them being the lengths of the geodesic boundary components of the pairs of pants P_1 and P_2 , of which two are equal, and the sixth value being the twist parameter along the curve β . The values $l_H(\tau(\alpha))$ and $l_H(\tau^{-1}(\alpha))$ are continuous functions of these parameters. Furthermore, for any element α in $\mathcal{S}(F)$, at least one of the two values $|l_H(\tau(\alpha)) - l_H(\alpha)|$ and $|l_H(\tau^{-1}(\alpha)) - l_H(\alpha)|$ is nonzero. This can be deduced from the fact that the length function $\theta \mapsto l_H(\tau_\theta(\alpha))$ under the twist deformation along the curve β is strictly convex (see [10] p. 130).

Therefore, for each $\alpha \in \mathcal{S}$, the value

$$\max \left(\frac{l_H(\tau^{-1}(\alpha))}{l_H(\alpha)}, \frac{l_H(\alpha)}{l_H(\tau^{-1}(\alpha))}, \frac{l_H(\tau(\alpha))}{l_H(\alpha)}, \frac{l_H(\alpha)}{l_H(\tau(\alpha))} \right)$$

is strictly positive. Since in this maximum, each time a number occurs, then its inverse also occurs, the value of the maximum is > 1 . This value varies continuously in terms of the six parameters referred to above. Under the hypotheses considered, the six parameters vary in a compact set. This shows the existence of the constant K . \square

We now consider a surface X , equipped with a hyperbolic structure H and a pair of pants decomposition \mathcal{P} satisfying Property (\star) stated in the introduction, and such that if C_i , $i = 1, 2, \dots$ is the collection of simple closed curves that are boundaries of pairs of pants in the collection \mathcal{P} , then,

$$(5) \quad \exists M > 0, \forall i = 1, 2, \dots, \frac{1}{M} \leq l_H(C_i) \leq M.$$

Proposition 2.20. *Let H be a hyperbolic metric on an infinite type surface X satisfying Property (5). Then, the Teichmüller space $\mathcal{T}_{ls}(H)$ is not proper.*

Proof. We can find a collection F_i , $i = 1, 2, \dots$ of four-holed spheres with geodesic boundary embedded in X and satisfying the following.

- (1) for each $i = 1, 2, \dots$, the four-holed sphere F_i is the union of two pairs of pants in the collection \mathcal{P} ;
- (2) for all $i \neq j$, the intersection of the interiors of F_i and F_j is empty.

In particular, in the interior of each four-holed sphere F_i , there is a curve β which belongs to the collection $\{C_i\}$. For each four-holed sphere F_i , we denote by τ_i the positive Dehn twist along the curve β in its interior. We let $x_i \in \mathcal{T}_{ls}(H)$ be the point representing (τ_i, H) and we let x be the point representing (Id, H) . By the arguments used in the proof of Proposition 2.12, there exists a constant $R > 0$ such that we have

$$\forall i = 1, 2, \dots, d_{ls}(x, x_i) \leq R.$$

Thus, the sequence (x_i) is contained in the closed ball B of radius R and centered at x . From Lemma 2.19, we deduce that there exists a constant K such that for each $i \neq j$, $d_{ls}(x_i, x_j) \geq K$. This shows that the sequence x_i does not have any Cauchy subsequence. This implies that the closed ball B is not compact, which proves that $\mathcal{T}_{ls}(H)$ is not proper. \square

It should be true that for any hyperbolic structure H on an infinite type surface, $\mathcal{T}_{ls}(H)$ is not proper, and this should follow from the fact that this space is infinite-dimensional.

We also ask the following.

Question 2.21. If H and H' are arbitrary hyperbolic metrics on surfaces on an infinite-type surface X , are the two Teichmüller spaces $\mathcal{T}_{ls}(H)$ and $\mathcal{T}_{ls}(H')$ locally bi-Lipschitz equivalent ?

In this regard, we note that in [11], A. Fletcher proved that the answer to Question 2.21 is yes if instead of the length-spectrum metric, one takes the Teichmüller metric (see Section 4 below). As a matter of fact, Fletcher proved that the Teichmüller metric on any Teichmüller space of an infinite-type Riemann surface is locally bi-Lipschitz equivalent to the Banach space l^∞ of bounded sequences with the supremum norm.

We end this section by a discussion of the completeness of the length spectrum metric.

In the paper [28] (Corollary 1.1), H. Shiga showed that for surfaces of infinite type, the topology defined by the length spectrum metric is in general not complete, but that under condition (5), this metric is complete. A natural question then is the following:

Question 2.22. Give a necessary and sufficient condition on a hyperbolic metric H on a surface of infinite type under which the Teichmüller space $\mathcal{T}_{ls}(H)$ is complete.

Note that for surfaces of finite type, the completeness of the length spectrum metric does not follow automatically from the completeness of the Teichmüller metric, even though the topologies induced by the length spectrum and the Teichmüller metrics are the same (see [18]). But it is true that the length spectrum metric in the case of surfaces of finite type is complete. For completeness, we now provide a proof of this fact.

We shall discuss the Teichmüller metric for Teichmüller spaces of surfaces of infinite type in Section 4 below, but we assume here that the reader is familiar with the Teichmüller metric of surfaces of finite type.

Let S be a Riemann surface of conformally finite type (g, n) , where g is the genus and n the number of punctures of S . For any $\epsilon > 0$, let $\mathcal{T}^\epsilon(S)$ be the subset of $\mathcal{T}(S)$ consisting of equivalence classes of marked hyperbolic surfaces in which the length of any simple closed curve which is not homotopic to a puncture is at least ϵ . We call $\mathcal{T}^\epsilon(S)$ the *thick part* of $\mathcal{T}(S)$. We recall that for surfaces of finite type, all

the Teichmüller spaces coincide from the set-theoretic point of view, and therefore we do not distinguish between the spaces \mathcal{T}_{l_s} and \mathcal{T}_{q_c} . We denote by d_{l_s} and d_{q_c} respectively the length-spectrum and the Teichmüller metrics on $\mathcal{T}(S)$.

We shall use the following result, proved in [20].

Lemma 2.23. *There exists a constant $N(g, n, \epsilon)$, depending only on g, n, ϵ , such that for any x and y in $\mathcal{T}^\epsilon(S)$, we have*

$$d_{l_s}(x, y) \leq d_{q_c}(x, y) \leq 4d_{l_s}(x, y) + N(g, n, \epsilon).$$

We have the following.

Theorem 2.24. *Let S be a conformally finite type hyperbolic surface and let $x_n, n = 0, 1, \dots$ be a sequence of elements in $\mathcal{T}(S)$. Then*

$$\lim_{n \rightarrow \infty} d_{q_c}(x_n, x_0) = \infty \iff \lim_{n \rightarrow \infty} d_{l_s}(x_n, x_0) = \infty.$$

Proof. We first recall that a result of Wolpert (see [1]) says that given any K -quasiconformal map $f : S_0 \rightarrow S$, then, for any hyperbolic metrics H_0 and H on S_0 and S respectively and for any element α in $\mathcal{S}(S_0)$, we have

$$(6) \quad \frac{1}{K} \leq \frac{l_H(f(\alpha))}{l_{H_0}(\alpha)} \leq K.$$

This implies that if $\lim_{n \rightarrow \infty} d_{l_s}(x_n, x_0) = \infty$, then $\lim_{n \rightarrow \infty} d_{q_c}(x_n, x_0) = \infty$.

Now suppose that $\lim_{n \rightarrow \infty} d_{q_c}(x_n, x_0) = \infty$. Consider first the following two cases:

(1) the sequence (x_n) leaves any thick part $\mathcal{T}^\epsilon(S)$. In this case it is clear that $\lim_{n \rightarrow \infty} d_{l_s}(x_n, x_0) = \infty$.

(2) the sequence (x_n) stays in some thick part $\mathcal{T}^\epsilon(S)$ for some $\epsilon > 0$. In this case, from Lemma 2.23 we have $\lim_{n \rightarrow \infty} d_{l_s}(x_n, x_0) = \infty$.

Now we discuss the general case. Assume that $d_{l_s}(x_n, x_0)$ does not go to ∞ as $n \rightarrow \infty$. Then there exists subsequence $x_{n_k}, n_k = 1, 2, \dots$ of (x_n) such that the set $\{d_{l_s}(x_{n_k}, x_0), n_k = 1, \dots\}$ is bounded. From the above discussion, if for some K_0 , a subsequence $x_{n_k}, n_k > K_0$ leaves any thick part of $\mathcal{T}^\epsilon(S)$, we get a contradiction. Otherwise, there exists a subsequence of $x_{n_k}, n_k = 1, 2, \dots$ which stays in some thick part of $\mathcal{T}^\epsilon(S)$, which also leads a contradiction. \square

Theorem 2.25. *Let S be a conformally finite type Riemann surface. Then $\mathcal{T}_{l_s}(S)$ equipped with the length spectrum metric is complete.*

Proof. We need to prove that for any Cauchy sequence $x_n, n = 1, \dots$ in $(\mathcal{T}(S), d_{l_s})$, there exists $x_0 \in \mathcal{T}(S)$, such that $\lim_{n \rightarrow \infty} d_{l_s}(x_n, x_0) = 0$.

The sequence (x_n) is bounded in the metric d_{l_s} . From Theorem 2.24, it follows that (x_n) is also bounded in the metric d_{q_c} . Since d_{q_c} is complete and since the Teichmüller space considered is finite-dimensional, there exists a subsequence (x_{n_k}) of (x_n) and $x_0 \in \mathcal{T}(S)$ such that $\lim_{n_k \rightarrow \infty} d_{q_c}(x_{n_k}, x_0) = 0$. From the topological equivalence of d_{q_c} and d_{l_s} on $\mathcal{T}(S)$ [18], it follows that

$$\lim_{n_k \rightarrow \infty} d_{l_s}(x_{n_k}, x_0) = 0.$$

As $\{x_n\}_{n=1}^\infty$ is a Cauchy sequence in the metric d_{l_s} , $\lim_{n \rightarrow \infty} d_{l_s}(x_n, x_0) = 0$. \square

From [28] and [20], Theorem 2.25 is not true for certain Teichmüller spaces of infinite type surfaces.

3. THE BI-LIPSCHITZ TEICHMÜLLER SPACES

We say that a homeomorphism $f : (X_1, d_1) \rightarrow (X_2, d_2)$ between metric spaces is *bi-Lipschitz* if there exists a real number $K \geq 1$ satisfying

$$(7) \quad \frac{1}{K}d_1(x, y) \leq d_2(f(x), f(y)) \leq Kd_1(x, y).$$

The real number K in (7) is called a *bi-Lipschitz constant of f* . Two metric spaces are said to be *bi-Lipschitz equivalent* if there exists a bi-Lipschitz homeomorphism between them.

Definition 3.1. Let H_0 be a hyperbolic metric on the surface X . The *bi-Lipschitz Teichmüller space of H_0* , denoted by $\mathcal{T}_{bL}(H_0)$, is the set of equivalence classes $[f, H]$ of pairs (f, H) where H is a hyperbolic metric on a surface homeomorphic to X and where $f : H_0 \rightarrow H$ (the *marking of H*) is a bi-Lipschitz homeomorphism, and where two such pairs (f, H) and (f', H') are considered equivalent if there exists an isometry $f'' : H \rightarrow H'$ homotopic to $f' \circ f^{-1}$.

The topology of $\mathcal{T}_{bL}(H_0)$ is the one induced by the *bi-Lipschitz metric d_{bL}* , defined by

$$d_{bL}([f, H], [f', H']) = \frac{1}{2} \log \inf\{K\},$$

where the infimum is taken over all bi-Lipschitz constants K of bi-Lipschitz homeomorphisms $f'' : H \rightarrow H'$ homotopic to $f' \circ f^{-1}$, with (f, H) and (f', H') being two hyperbolic surfaces representing the two points $[f, H]$ and $[f', H']$ in $\mathcal{T}_{bL}(H_0)$.

To see that this defines indeed a metric on $\mathcal{T}_{bL}(H_0)$, we note that if we take a sequence of bi-Lipschitz maps homotopic to $f' \circ f^{-1}$ whose bi-Lipschitz constants tend to 1, then there exists an isometry homotopic to $f' \circ f^{-1}$. One easy way to see this is to use the following remarks that show that under these hypotheses, the length-spectrum constant for the map $f' \circ f^{-1}$ must be 1, hence, by Proposition 2.2, there exists an isometry homotopic to $f' \circ f^{-1}$.

If $f : H_0 \rightarrow H$ is a bi-Lipschitz homeomorphism with constant K , then, from the definition of the length of a curve, we have, for every element α in \mathcal{S}_0 ,

$$\frac{1}{K} \leq \frac{l_H(f(\alpha))}{l_{H_0}(\alpha)} \leq K.$$

It follows that a bi-Lipschitz homeomorphism f between two hyperbolic surfaces is also length-spectrum bounded, with length-spectrum constant bounded by any bi-Lipschitz constant of f . This implies that we have a natural inclusion

$$(8) \quad \mathcal{T}_{bL}(H_0) \hookrightarrow \mathcal{T}_{ls}(H_0)$$

and that for any x and y of $\mathcal{T}_{bL}(H_0)$, we have

$$(9) \quad d_{ls}(x, y) \leq d_{bL}(x, y).$$

In particular, the inclusion map in (8) is continuous. It will follow from the discussion in the next section that this inclusion is generally strict.

Using the inclusion map in (8), the same example as in 2.4 shows that there exist pairs of marked hyperbolic structures on X which are not bi-Lipschitz equivalent. Therefore the Teichmüller spaces $\mathcal{T}_{bL}(H_0)$ and $\mathcal{T}_{bL}(H_1)$ based at two different hyperbolic structures H_0 and H_1 on X are in general setwise different.

Question 3.2. Characterize the hyperbolic surfaces H_0 on an infinite-type hyperbolic surface such that the inclusion in (8) is strict.

Definition 3.3. The *bi-Lipschitz mapping class group* of the surface X relatively to the base hyperbolic metric H_0 , denoted by $\text{MCG}_{bL}(H_0)$, is the subgroup of

$\text{MCG}(X)$ consisting of homotopy classes of self-homeomorphisms of X that are bi-Lipschitz with respect to the metric H_0 .

From (9), we have an inclusion

$$\text{MCG}_{bL}(H_0) \subset \text{MCG}_{ls}(H_0).$$

Since we know that $\text{MCG}_{ls}(H_0) \subsetneq \text{MCG}(X)$ (Proposition 2.8), we have $\text{MCG}_{bl}(H_0) \subsetneq \text{MCG}(X)$.

The group $\text{MCG}_{bL}(H_0)$ acts naturally on the Teichmüller space $\mathcal{T}_{bL}(H_0)$, by right-composition with the inverse at the level of markings (see Formula (2) above), and this action preserves the metric d_{bL} .

Denoting as before $\text{MCG}_f(X)$ the subgroup of $\text{MCG}(X)$ consisting of the homotopy classes of homeomorphisms which are supported on a subsurface of finite type of X , we have the following.

Proposition 3.4. *There is a natural inclusion $\text{MCG}_f(X) \subset \text{MCG}_{bl}(H_0)$.*

Proof. As in the proof of Proposition 2.12, we use the fact that any element of the mapping class group of a surface of finite topological type is the product of a finite number of Dehn twist elements. A Dehn twist homeomorphism along a closed curve can be represented by a bi-Lipschitz homeomorphism supported on a compact annular neighborhood of that curve. The composition of these bi-Lipschitz homeomorphisms is a bi-Lipschitz homeomorphism of X . □

4. THE QUASICONFORMAL TEICHMÜLLER SPACE OF A SURFACE OF INFINITE TYPE

The Riemann surfaces, that is, the complex structures, that we shall consider will often be induced by hyperbolic metrics, and for this reason we shall use a metric definition of a quasiconformal map. We recall that if $f : S \rightarrow S'$ is a homeomorphism between two surfaces S and S' equipped with Riemannian metrics, and if f is differentiable at a point x of S , then the differential of f at x is an \mathbb{R} -linear map from the tangent space $T_x S$ of S at x to the tangent space $T_{f(x)} S'$ of S' at $f(x)$, and therefore, such a linear map sends circles in $T_x S$ centered at the origin to ellipses in $T_{f(x)} S'$ centered at the origin. The *quasiconformal dilatation* $K(f)_x$ of f at x is then equal to the *eccentricity* of an image ellipse, that is, the ratio of the large axis to the small axis of this ellipse. This quantity is independent of the choice of the circle in $T_x S$ centered at the origin. Now if $f : S \rightarrow S'$ is a homeomorphism, then the *quasiconformal constant* of f at a point x in S where f is not necessarily differentiable is defined as follows. For $r > 0$, we consider the quantities

$$L_f(x, r) = \sup\{d_{S'}(f(x), f(y)) \mid d_S(x, y) = r\}$$

and

$$l_f(x, r) = \inf\{d_{S'}(f(x), f(y)) \mid d_S(x, y) = r\}.$$

The *quasiconformal dilatation* $K(f)_x$ of f at x is the element of $\mathbb{R} \cup \{\infty\}$ defined as

$$K(f)_x = \limsup_{r \rightarrow 0} \frac{L_f(x, r)}{l_f(x, r)}.$$

In the case where f is differentiable at x , the two definitions coincide.

The quasiconformal dilatation of f at x depends only on the conformal classes of the Riemannian metrics on S and S' . Therefore we can talk about the quasiconformal constant of a homeomorphism between Riemann surfaces, by choosing Riemannian metrics in the given conformal classes and using the above definition.

The *maximal dilatation* $K(f)$ of a homeomorphism $f : S \rightarrow S'$ between two Riemann surfaces S and S' , also called the *quasiconformal dilatation*, or, in short, the *dilatation* of f , is defined as the supremum of the quasiconformal dilatation $K(f)_x$ of f over all points x of S . The homeomorphism f is said to be *quasiconformal* if its dilatation is finite.

Definition 4.1. Consider a Riemann surface structure S_0 on X . Its *quasiconformal Teichmüller space*, $\mathcal{T}_{qc}(S_0)$, is defined as the set of equivalence classes $[f, S]$ of pairs (f, S) , where S is a Riemann surface homeomorphic to X and $f : S_0 \rightarrow S$ a quasiconformal homeomorphism (called the marking of S), and where two pairs (f, S) and (f', S') are considered to be equivalent (and are said to be *conformally equivalent*) if there exists a conformal homeomorphism $f'' : S \rightarrow S'$ homotopic to $f' \circ f^{-1}$.

The space $\mathcal{T}_{qc}(S_0)$ is equipped with the Teichmüller metric, defined as follows. Given two elements $[f, S]$ and $[f', S']$ of $\mathcal{T}_{qc}(S_0)$ represented by marked conformal surfaces (f, S) and (f', S') , their *quasiconformal distance* (also called the *Teichmüller distance*, because it was introduced by Teichmüller) is defined as

$$(10) \quad d_{qc}([f, S], [f', S']) = \frac{1}{2} \log \inf \{K(f'')\}$$

where the infimum is taken over maximal dilatations $K(f'')$ of homeomorphisms $f'' : S \rightarrow S'$ homotopic to $f' \circ f^{-1}$.

The equivalence class of the marked Riemann surface (Id, S_0) is the *basepoint* of $\mathcal{T}_{qc}(S_0)$.

It is important to note that the definition of the Teichmüller distance given in (10) is the same as in the case of surfaces of finite conformal type, but that for surfaces of infinite type, choosing different basepoints may lead to different Teichmüller spaces. (But choosing a different basepoint by just applying a quasiconformal map leads to the same Teichmüller space, up to biholomorphism.) Any homeomorphism between two Riemann surfaces of finite conformal type is homotopic to a differentiable map, which is automatically quasiconformal, but there exist homeomorphisms between surfaces of infinite conformal type that are not homotopic to quasiconformal maps. We shall discuss this fact in more detail below.

We refer to Nag [24] for an exposition of the quasiconformal theory of infinite-dimensional Teichmüller spaces. In particular, it is known that the quasiconformal metric is complete.

The quasiconformal dilatation and the bi-Lipschitz constant of a homeomorphism are related to each other. We shall see precise relations between these two quantities (see Theorem 4.3 below).

By the uniformization theorem, given any conformal structure S_0 on X , there is a hyperbolic structure H_0 on X whose underlying conformal structure is S_0 . We shall also denote the space $\mathcal{T}_{qc}(S_0)$ by $\mathcal{T}_{qc}(H_0)$, meaning that we consider the conformal structure represented by the hyperbolic structure H_0 .

From Wolpert's inequality (6), there is a natural inclusion map

$$(11) \quad \mathcal{T}_{qc}(H_0) \hookrightarrow \mathcal{T}_{ls}(H_0).$$

This inclusion map is continuous since Wolpert's result (6) also implies that for any two elements x and y in $\mathcal{T}_{qc}(S_0)$, we have

$$(12) \quad d_{ls}(x, y) \leq d_{qc}(x, y).$$

Proposition 4.2. *In general, we have $\mathcal{T}_{ls}(H) \neq \mathcal{T}_{qc}(H)$.*

Proof. It suffices to construct a hyperbolic structure H on X and a self-homeomorphism of (X, H) which is not quasiconformal and which is length-spectrum bounded. This is done in Proposition 4.7 below. \square

Concerning the comparison between the quasiconformal distance and the bi-Lipschitz distance, there is the following.

Theorem 4.3 (Thurston [29] p. 268). *For every hyperbolic structure H , we have the set-theoretic equality*

$$(13) \quad \mathcal{T}_{qc}(H) = \mathcal{T}_{bL}(H)$$

and there exists a constant C such that for every x and y in $\mathcal{T}_{qc}(H)$, we have

$$(14) \quad d_{qc}(x, y) \leq d_{bL}(x, y) \leq Cd_{qc}(x, y).$$

It may be worth recalling that we are considering, as in Thurston's setting, complete hyperbolic structures.

The equality between the Teichmüller spaces in (13) follows from the inequalities between distances in (14). The left hand side inequality in (14) follows from the fact that the quasiconformal dilatation is the supremum of an infinitesimal distortion at each point, and the bi-Lipschitz constant measures distortion of distances at all scales. The existence of a constant C for which the right hand side inequality is satisfied is non-trivial, and it can be proved using the Douady-Earle quasiconformal barycentric extension theory in [6] (see [29] p. 268).

Corollary 4.4. *For any hyperbolic structure H on a surface X , we have the set-theoretic equality $MCG_{qc}(H) = MCG_{bL}(H)$.*

Douady and Earle gave in [6] a proof of the fact that any quasiconformal Teichmüller space $\mathcal{T}_{qc}(H_0)$ is contractible (see [6] Theorem 3, where this result is also attributed to Tukia). We ask the following:

Question 4.5. Are the Teichmüller spaces $\mathcal{T}_{ls}(H_0)$ contractible? Are they even connected?

Again, using the inclusion map in (11) and taking the conformal structures S_0 and S_1 underlying the two hyperbolic structures H_0 and H_1 of Example 2.4, we obtain an element $H_1 \notin \mathcal{T}_{qc}(H_0)$. This gives an example of two Teichmüller spaces $\mathcal{T}_{qc}(S_0)$ and $\mathcal{T}_{qc}(S_1)$ whose basepoints are two homeomorphic Riemann surfaces S_0 and S_1 , such that $\mathcal{T}_{qc}(S_0) \neq \mathcal{T}_{qc}(S_1)$ setwise.

Definition 4.6. Let S be a Riemann surface structure on the topological surface X . The *quasiconformal mapping class group* of S , denoted by $MCG_{qc}(S)$, is the subgroup of $MCG(X)$ consisting of homotopy classes of homeomorphisms that are quasiconformal with respect the structure S .

The group $MCG_{qc}(S)$ naturally acts on the Teichmüller space $\mathcal{T}_{qc}(S)$, by right-composition with the inverse at the level of markings (see the action described in (2)), and this action preserves the metric d_{qc} .

The homomorphism of the group $MCG_{qc}(S)$ into the group of isometries of $\mathcal{T}_{qc}(S)$ is injective (see [9] and [8]).

A theorem whose most general version is due to V. Markovic (see [21]) asserts that any isometry of the space $\mathcal{T}_{qc}(S)$ is induced by an element of $MCG_{qc}(S)$. Special cases of this result had been obtained by several people, including Earle and Kra [7], Lakic [16] and Matsuzaki [23]. The earliest version of this theorem, in the case of finite-type surfaces, is due to Royden [27].

Proposition 4.7. *Let H be a hyperbolic structure on X such that there exists a sequence of homotopy classes of disjoint essential closed curves on X whose length tends to 0. Then, $MCG_{qc}(H) \subsetneq MCG_{ls}(H)$.*

Proof. We produce an element $T \in \text{MCG}_{ls}(H)$ with $T \notin \text{MCG}_{qc}(H)$.

Without loss of generality, we can assume that there exists a sequence $\alpha_n, n = 1, 2, \dots$, of homotopy classes of disjoint essential closed curves on X whose lengths satisfy $l_H(\alpha_n) = \varepsilon_n$ with $e^{-(n+1)^2} < \varepsilon_n < e^{-n^2}$.

Let

$$t_n = \left\lceil \frac{\log |\log \varepsilon_n|}{\varepsilon_n} \right\rceil + 1, \quad n = 1, 2, \dots$$

($\lceil x \rceil$ denoting the integral part of the real number x).

For each $n = 1, 2, \dots$, let τ_n be the t_n -th power of the positive Dehn twist about α_n . We take all the positive Dehn twists to be supported on disjoint annuli, and we define $T \in \text{MCG}(X)$ to be the infinite composition $T = \tau_1 \circ \tau_2 \circ \dots$. From [20], we have that T is not in $\text{MCG}_{qc}(X)$.

Now we show that $T \in \text{MCG}_{ls}(X)$. We need to show that

$$\log \sup_{\alpha \in \mathcal{S}(X)} \left\{ \frac{l_H(f(\alpha))}{l_H(\alpha)}, \frac{l_H(\alpha)}{l_H(f(\alpha))} \right\} < \infty.$$

The proof parallels the proof of Proposition 2.13.

Let α be an arbitrary homotopy class of essential curves in X .

If $i(\alpha, \alpha_n) = 0, n = 1, 2, \dots$, then $\alpha = T(\alpha)$ and $l_H(T(\alpha)) = l_H(\alpha)$.

Suppose that $i(\alpha, \alpha_n) \neq 0$ for some n . Without loss of generality, we can assume that $i(\alpha, \alpha_n) \neq 0$, for all $n = 1, 2, \dots, N$. By the Collar Lemma, on any hyperbolic surface, any closed geodesic whose length ε is sufficiently small has an embedded collar neighborhood of width $|\log \varepsilon|$. Thus, we can write

$$l_H(\alpha) = \sum_{n=1}^N i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha$$

where $B_\alpha > 0$ is a constant that depends on the closed curve α .

From the definition of a Dehn twist, we have

$$\begin{aligned} & \sum_{n=1}^N i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha - \sum_{n=1}^N i(\alpha, \alpha_n) t_n \varepsilon_n \leq l_H(T(\alpha)) \\ & \leq \sum_{n=1}^N i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha + \sum_{n=1}^N i(\alpha, \alpha_n) t_n \varepsilon_n. \end{aligned}$$

Therefore we have

$$\begin{aligned} \frac{l_H(T(\alpha))}{l_H(\alpha)} & \leq \frac{\sum_{n=1}^N i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha + \sum_{n=1}^N i(\alpha, \alpha_n) t_n \varepsilon_n}{\sum_{n=1}^N i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha} \\ & = 1 + \frac{\sum_{n=1}^N i(\alpha, \alpha_n) t_n \varepsilon_n}{\sum_{n=1}^N i(\alpha, \alpha_n) |\log \varepsilon_n| + B_\alpha} \\ & \leq 1 + \frac{\sum_{n=1}^N i(\alpha, \alpha_n) \log |\log \varepsilon_n|}{\sum_{n=1}^N i(\alpha, \alpha_n) |\log \varepsilon_n|} \\ & \leq 1 + \sum_{n=1}^N \frac{\log |\log \varepsilon_n|}{|\log \varepsilon_n|} \\ & \leq 1 + 2 \sum_{n=1}^N \frac{\log(n+1)}{n^2} \\ & \leq 1 + 2 \sum_{n=1}^\infty \frac{\log(n+1)}{n^2} < \infty \end{aligned}$$

which is bounded independently of α and n . In the same way, we can prove that $\frac{l_H(\alpha)}{l_H(T(\alpha))}$ is also bounded independently of α and n . This implies that $T \in \text{MCG}_{ls}(H)$. □

Question 4.8. Proposition 4.7 gives a sufficient condition under which the inclusion in (11) is strict. The question is to find sufficient and necessary conditions for the same fact.

Recall that $\text{MCG}_f(X)$ denotes the group of homotopy classes of homeomorphisms of X which are supported on subsurfaces of finite analytical type.

Proposition 4.9. *For any conformal structure S on X , we have a natural inclusion $\text{MCG}_f(X) \subset \text{MCG}_{qc}(S)$.*

Proof. Any homeomorphism of a subsurface of X of finite analytical type is homotopic to a diffeomorphism, and therefore to a quasiconformal homeomorphism. This quasiconformal homeomorphism can be extended to a quasiconformal homeomorphism of X with the same quasiconformal dilatation constant. \square

It was shown in [20] that the following is a necessary condition for the equality $\mathcal{T}_{qc}(H_0) = \mathcal{T}_{ls}(H_0) = \mathcal{T}_{bL}(H_0)$: there exists a positive constant ϵ , such that for any element α in $\mathcal{S}(X)$, $l_{H_0}(\alpha) \geq \epsilon$.

Let C_i ($i = 1, 2, \dots$) be the collection of the simple closed curves that are boundaries of pairs of pants in the pants decomposition \mathcal{P} of the surface X . We recall that Condition (5) above, for a hyperbolic structure H on X , states the following:

$$\exists M > 0, \forall i = 1, 2, \dots, \frac{1}{M} \leq l_H(C_i) \leq M.$$

Theorem 4.10. *Suppose that the base hyperbolic metric H satisfies Condition (5) and let $x_n, n = 0, 1, \dots$, be a sequence of points in $\mathcal{T}_{qc}(H)$. Then the following are equivalent:*

- (a) $\lim_{n \rightarrow \infty} d_{qc}(x_n, x_0) = 0$;
- (b) $\lim_{n \rightarrow \infty} d_{ls}(x_n, x_0) = 0$;
- (c) $\lim_{n \rightarrow \infty} d_{bL}(x_n, x_0) = 0$.

Proof. From (14), we have (a) \Rightarrow (b). From Theorem 4.3, we have (a) \iff (c). Finally, from Shiga's result in [28] (proof of Theorem 1.2), we have (b) \Rightarrow (a). \square

We consider for a while surfaces of finite type.

The following follows from Lemma 2.23.

Lemma 4.11. *Let S be a finite type Riemann surface of genus g with n punctures and c boundary components, and let $\epsilon > 0$. Suppose that there exists $n_0 > 0$ such that $g + n + c \leq n_0$. For any x and y in $\mathcal{T}^\epsilon(S)$, there exists a constant $N(n_0, \epsilon)$, depending only on n_0 and ϵ , such that*

$$d_{ls}(x, y) \leq d_{qc}(x, y) \leq 4d_{ls}(x, y) + N(n_0, \epsilon).$$

Proof. For $c = 0$, this lemma is a reformulation of Lemma 2.23. For $c \neq 0$, the result follows by taking doubles of surfaces along geodesic boundary components. \square

Let S be a Riemann surface equipped with a hyperbolic metric and let α be a simple closed geodesic on S . We recall that a time- t Fenchel-Nielsen deformation of S with respect to α is a Riemann surface S_t obtained by cutting S along α and regluing the cut off borders after twisting by a hyperbolic distance t along α . If S is a marked surface, then S_t is also equipped with a natural marking. We refer to [14] for background material.

The following two lemmas are inspired from Lemma 3.1 in [28], and a corresponding lemma in [20].

Lemma 4.12. *Let S be a hyperbolic surface and let α be a simple closed curve on S which is not homotopic to a boundary component of S . Let S_t be the time- t Fenchel-Nielsen deformation of S about α , let $\tau_\alpha^t : S \rightarrow S_t$ be a quasiconformal homeomorphism and let $K(t)$ be its dilatation. Then $\lim_{t \rightarrow \infty} K(t) = \infty$.*

Proof. Let ρ be the hyperbolic distance on S . For $s > 0$, let

$$W_\alpha = \{p \in S \mid \rho(p, \alpha) < s\}.$$

The positive constant s is chosen to be so small that W_α is a tubular neighborhood of α in S . We construct a quasiconformal homeomorphism g_t from S to S_t which represents the positive twisting along α by t . This map is best described by looking to a lift to the universal covers of the two surfaces S and S_t .

We identify the universal covers of S and S_t with the upper-half model of the hyperbolic plane, \mathbb{H}^2 . We denote by $\pi : \mathbb{H}^2 \rightarrow S$ the covering map. We consider connected components of $\pi^{-1}(W_\alpha)$ and of $\pi^{-1}(\alpha)$ in \mathbb{H}^2 . Without loss of generality, we may assume that the connected component of $\pi^{-1}(\alpha)$ is the positive half of the imaginary axis. Let \tilde{W}_α be the connected component of $\pi^{-1}(W_\alpha)$ containing the positive half of the imaginary axis. For a suitable θ with $0 < \theta < \frac{\pi}{2}$, we can express \tilde{W}_α as

$$\tilde{W}_\alpha = \{z \in \mathbb{H}^2 \mid \frac{\pi}{2} - \theta < \arg z < \frac{\pi}{2} + \theta\}.$$

Let ω_t be the lift of a quasiconformal mapping in the homotopy class of g_t , normalized by $\omega_t(0) = 0, \omega_t(i) = i$ and $\omega_t(\infty) = \infty$. It is well known that ω_t can be extended to a homeomorphism of $\overline{\mathbb{H}^2}$ and that the boundary mapping $\omega_t|_{\mathbb{R}}$ depends only on the homotopy class of g_t .

Finally, we let δ be the geodesic in \mathbb{H}^2 which (as a set) is defined as

$$\delta = \{z \in \mathbb{H}^2, |z| = 1.\}$$

Let z_1 and z_2 , ($\Re z_1 < 0 < \Re z_2$) be the points of $\delta \cap \tilde{W}_\alpha$. From the fact that g_t is the positive time- t Fenchel-Nielsen deformation, we have $\omega_t(z_1) = z_1, \omega_t(z_2) = e^t z_2$, and $\omega_t(\delta \cap \tilde{W}_\alpha)$ is an arc connecting z_1 and $e^t z_2$ in the component \tilde{W}_α (see Figure 5). There is a similar description of the image by ω_t of a subarc of δ in each component of $\pi^{-1}(W_\alpha)$. Let $z_1 = -\sin \theta + i \cos \theta$ and $z_2 = \sin \theta + i \cos \theta$. Then,

$$\omega_t(z_2) = e^t(\sin \theta + i \cos \theta)$$

and

$$\Re \omega_t(z_2) - \Re z_2 = (e^t - 1) \sin \theta.$$




FIGURE 5.

Now we study the action of the quasiconformal map ω_t on the line at infinity \mathbb{R} . From the above construction we have

$$\omega_t(1) - 1 > \Re \omega_t(z_2) - \Re z_2 = (e^t - 1) \sin \theta$$

and

$$-1 < \omega_t(-1) < 0.$$

Summing up, we have

$$(15) \quad -1 < \omega_t(-1) < 0 < 1 + (e^t - 1) \sin \theta < \omega_t(1).$$

We use the following definition of the cross ratio of quadruples in \mathbb{H}^2 :

$$[a, b, c, d] = \frac{(a-b)(c-d)}{(a-d)(c-b)}.$$

This definition extends to a notion of cross ratio defined for quadruples of points at infinity, and we have

$$[-1, 0, 1, \infty] = -1.$$

We shall use the geometric definition of a quasiconformal mapping, that involves the ratios of moduli of quadrilaterals, and we refer to Ahlfors' book [2] for this definition.

We can talk about the conformal modulus of ideal quadrilaterals (quadrilaterals having their vertices at infinity), and we have the following formula:

$$(16) \quad \text{Mod}(H(-1, 0, 1, \infty)) = 1.$$

We set

$$\nu_t = [\omega_t(-1), \omega_t(0), \omega_t(1), \omega_t(\infty)] = [\omega_t(-1), 0, \omega_t(1), \infty] = \frac{\omega_t(-1)}{\omega_t(1)}.$$

Then

$$|\nu_t| = \left| \frac{\omega_t(-1)}{\omega_t(1)} \right| \leq \frac{1}{1 + (e^t - 1) \sin \theta} \rightarrow 0, \quad t \rightarrow \infty.$$

Therefore, we have the following.

$$(17) \quad \text{Mod}(H(\omega_t(-1), \omega_t(0), \omega_t(1), \omega_t(\infty))) \rightarrow 0, \quad \text{as } t \rightarrow \infty.$$

By the geometric definition of quasiconformal mappings, we get

$$(18) \quad \frac{1}{K(\omega_t)} \leq \frac{\text{Mod}(H(\omega_t(-1), \omega_t(0), \omega_t(1), \omega_t(\infty)))}{\text{Mod}(H(-1, 0, 1, \infty))}.$$

Summing up, (16), (17) and (18) give

$$(19) \quad K(\omega_t) \rightarrow \infty, \quad n \rightarrow \infty.$$

Since any quasiconformal mapping in the homotopy class of ω_t induces the same boundary map as ω_t , its quasiconformal dilatation $K(t)$ tends to infinity as $t \rightarrow \infty$. \square

Lemma 4.13. *Let S be a Riemann surface and let α_n be a sequence of simple closed curves on S which are not homotopic to a boundary component of S . Given an unbounded sequence t_n of real numbers, let $\tau_{\alpha_n}^{t_n} : S \rightarrow S_{t_n}$ be the time- t_n Fenchel-Nielsen deformation about α_n , and let $K(t_n)$ be the dilatation of $\tau_{\alpha_n}^{t_n}$. Then $\lim_{t_n \rightarrow \infty} K(t_n) = \infty$.*

Lemma 4.13 can be proved in the same way as Lemma 4.12.

Theorem 4.14. *Suppose that the base hyperbolic metric H_0 satisfies Condition (5). Then, $\mathcal{J}_{qc}(H_0) = \mathcal{J}_{Is}(H_0) = \mathcal{J}_{bL}(H_0)$.*

Proof. It is easy to see that if H_0 satisfies Condition (5), then every element of $\mathcal{J}_{Is}(H_0)$ satisfies that condition (but of course, not necessarily with the same constant M). Since the hyperbolic length of the image of a homotopy class α of closed curves by a K -Lipschitz homeomorphism is bounded from above by K times the hyperbolic length of α , it follows that if H_0 satisfies Condition (5), then every element of $\mathcal{J}_{bL}(H_0)$ satisfies that condition. Finally, using Wolpert's inequality (6), it is also easy to show that if H_0 satisfies Condition (5), then every element of

$\mathcal{T}_{qc}(H_0)$ satisfies the same condition. (One can also use the set-theoretic equality $\mathcal{T}_{bL}(H_0) = \mathcal{T}_{qc}(H_0)$ and their inclusion in $\mathcal{T}_{ls}(H_0)$.)

From (8), we have $\mathcal{T}_{bL}(H_0) \subset \mathcal{T}_{ls}(H_0)$. From (11), we have $\mathcal{T}_{qc}(H_0) \subset \mathcal{T}_{ls}(H_0)$. From Theorem 4.3, we have $\mathcal{T}_{bL}(H_0) = \mathcal{T}_{qc}(H_0)$. Therefore, it remains to show that $\mathcal{T}_{ls}(H_0) \subset \mathcal{T}_{qc}(H_0)$. We now prove this by showing that any two hyperbolic structures at bounded length spectrum distance are at bounded quasiconformal distance.

We consider two decompositions of X by surfaces with disjoint interiors, $X = \bigcup_{n=1}^{\infty} X_n$ and $X = \bigcup_{n=1}^{\infty} X'_n$, satisfying the following properties:

- For each $n = 1, 2, \dots$, each of the surfaces X_n and X'_n is of finite type, and the sum of its genus, number of punctures and number of boundary components is bounded by an integer n_0 that is independent of n .
- If $S_i, i = 1, 2, \dots$ and $S'_i, i = 1, 2, \dots$ are the decomposition curves of the surface decompositions $X = \bigcup_{n=1}^{\infty} X_n$ and $X = \bigcup_{n=1}^{\infty} X'_n$ respectively, then $\{S_i\}_{i=1}^{\infty} \subset \{C_i\}_{i=1}^{\infty}$ and $\{S'_i\}_{i=1}^{\infty} \subset \{C_i\}_{i=1}^{\infty}$, where $C_i, i = 1, 2, \dots$ are the curves in the pants decomposition \mathcal{P} of the surface X satisfying condition (5).
- We have $\{\bigcup_{i=1}^{\infty} S_i\} \cap \{\bigcup_{i=1}^{\infty} S'_i\} = \emptyset$.

Let $x_1 = [f_1, H_1]$ and $x_2 = [f_2, H_2]$ be two elements of the Teichmüller space $\mathcal{T}_{ls}(H_0)$ and let $M_0 = d_{ls}(x_1, x_2)$

For $i = 0, 1, 2$ and for each $n = 1, 2, \dots$, let H_i^n , be the restriction of the hyperbolic structure H_i , on X_n , let $f_i^n : H_0^n \rightarrow H_i^n$, be the restriction of f_i to H_0^n and let $x_i^n = [f_i^n, H_i^n]$.

For any $i = 1, 2$ and for each $n = 1, 2, \dots$, we have $x_i^n \in \mathcal{T}^{\epsilon}(H_0^n)$, where ϵ only depends on the constant M that appears in Condition (5). From Lemma 4.11, we can write, for any $n = 1, 2, \dots$,

$$d_{ls}(x_1^n, x_2^n) \leq d_{qc}(x_1^n, x_2^n) \leq 4d_{ls}(x_1^n, x_2^n) + N(n_0, \epsilon) \leq 4M_0 + N(n_0, \epsilon).$$

It remains to show that the amounts of twist along the curves $S_i, i = 1, \dots$, are bounded.

For an arbitrary integer $p \geq 1$, we can find an integer n_p such that $S_p \subset X'_{n_p}$. For $i = 0, 1, 2$, let H'_{i_p} , be the restriction of H_i to X'_{n_p} and let $f'_{i_p} : H'_{0_p} \rightarrow H'_{i_p}$, $i = 0, 1, 2$, be the map obtained by restriction of f . Let $x'_{i_p} = [f'_{i_p}, H'_{i_p}]$, $i = 0, 1, 2$. Then, by the same arguments that we used above, we have

$$(20) \quad d_{qc}(x'_{1_p}, x'_{2_p}) \leq 4M_0 + N(n_0, \epsilon).$$

From Lemmas 4.13 and 4.12, the amount of twist about $C_{i_0}, i = 0, 1, 2$ is bounded by a constant B , depending only on M, M', n_0 , since otherwise, if this amount of twist had no bounds, Lemma 4.12 would imply that the dilatation of the corresponding quasiconformal mapping tends to infinity. This contradicts Inequality (20).

Since for an arbitrary integer $p \geq 1$ $d_{qc}(x'_{1_p}, x'_{2_p}) \leq 4M_0 + N(n_0, \epsilon)$, and since the twists about the boundary curves have a common upper bound, we have $d_{qc}(x_1, x_2) < \infty$. \square

Theorem 4.15. *Suppose that the base hyperbolic metric H_0 satisfies Condition (5). Let $x_n, n = 0, 1, \dots$, be a sequence of marked Riemann surface relative to the base hyperbolic structure H_0 . Then the following are equivalent:*

- $\lim_{n \rightarrow \infty} d_{qc}(x_n, x_0) = \infty$,
- $\lim_{n \rightarrow \infty} d_{ls}(x_n, x_0) = \infty$,
- $\lim_{n \rightarrow \infty} d_{bL}(x_n, x_0) = \infty$.

Proof. From (14) we know that (a) is equivalent to (c). From (9) we know that (a) implies (b). By quoting Shiga's result as we did in the proof of Theorem 4.10, we have (b) implies (a). \square

Combining Theorems 4.10 and 2.25, we obtain

Theorem 4.16. *Suppose that the base hyperbolic metric H satisfies Condition (5). Then $\mathcal{T}_{qc}(H)$, $\mathcal{T}_{ls}(H)$ and $\mathcal{T}_{bL}(H)$ are all complete.*

Corollary 4.17. *If the base hyperbolic metric H satisfies Condition (5), then $MCG_{qc}(H) = MCG_{ls}(H)$.*

REFERENCES

- [1] W. Abikoff, The real analytic theory of Teichmüller space, Lecture Notes in Mathematics 820, Springer-Verlag (1980).
- [2] L. V. Ahlfors, Lectures on quasiconformal maps, Van Nostrand-Reinhold, Princeton, New Jersey, 1966. New edition: AMS University Lecture Series, vol. 38, 2006.
- [3] P. Buser, Geometry and Spectra of Compact Riemann Surfaces, Birkhäuser (1992).
- [4] A. Basmajian, Hyperbolic structures for surfaces of infinite type, Trans. Amer. Math. Soc. Vol. 336, No. 1 (1993) 421–444.
- [5] M. Gromov, Structures métriques pour les variétés riemanniennes, Textes Mathématiques, 1. Paris: Cedec/Fernand Nathan, (1981).
- [6] A. Douady & C. J. Earle, Conformally natural extensions of homeomorphisms of the circle, Acta Math. 157 (1986) 23–48.
- [7] C. Earle & I. Kra, On holomorphic mappings between Teichmüller spaces, *Contributions to Analysis*, Academic Press, New York (1974), 107–124.
- [8] C. Earle, F. Gardiner & N. Lakic, Teichmüller spaces with asymptotic conformal equivalence, Preprint, IHES, 1995.
- [9] A. Epstein, Effectiveness of Teichmüller modular groups, In the Tradition of Ahlfors and Bers, Contemporary Math. 256, AMS, 2000, 69–74.
- [10] A. Fathi, F. Laudenbach & V. Poénaru, *Travaux de Thurston sur les surfaces*, Astérisque 66–67 (1979).
- [11] A. Fletcher, Local rigidity of Infinite dimensional Teichmüller spaces, *J. London Math. Soc.*, **74** (1) (2006), 26–40.
- [12] E. Fujikawa, Limit sets and regions of discontinuity of Teichmüller modular groups, Proc. Amer. Math. Soc. 132 (2004), 117–126.
- [13] E. Fujikawa, H. Shiga & M. Taniguchi, On the action of the mapping class group for Riemann surfaces of infinite type, J. Math. Soc. Japan 56, No. 4, 1069–1086 (2004).
- [14] Y. Iwayoshi & M. Taniguchi, An introduction to Teichmüller spaces, Springer-Verlag, Berlin - Heidelberg - New York, 1992.
- [15] B. V. Kerékjártó, Vorlesungen über Topologie. I, Verlag von Julius Springer, Berlin, 1923.
- [16] N. Lakic, An isometry theorem for quadratic differentials on Riemann surfaces of finite genus, *Trans. Amer. Math. Soc.*, **349**, no.7 (1997), 2951–2967.
- [17] Z. Li, Teichmüller metric and length spectrum of Riemann surfaces, Sci. China Ser. A 3, 1986, 802–810.
- [18] L. Liu, On the length spectrums of non-compact Riemann surfaces, Ann. Acad. Sci. Fenn. Math. 24, 1999, 11–22.
- [19] L. Liu, On the metrics of length spectrum in Teichmüller space, Chinese J. Cont. math. 22 (1) (2001), 23–34.
- [20] L. Liu, Z. Sun and H. Wei, Topological equivalence of metrics in Teichmüller space, Ann. Acad. Sci. Fenn. Math. 33, No. 1, 159–170 (2008).
- [21] V. Markovic, Biholomorphic maps between Teichmüller spaces, *Duke Math. J.*, **120**, no. 2 (2003), 405–431.
- [22] K. Matsuzaki, The infinite product of Dehn twists acting on infinite dimensional Teichmüller spaces, Kodai Math. J. 26 (2003), 279–287.
- [23] K. Matsuzaki, Inclusion relations between the Bers embeddings of Teichmüller spaces, *Israel J. Math.*, **140** (2004), 113–123.
- [24] S. Nag, The complex analytic theory of Teichmüller spaces, John Wiley, Canadian Mathematical Society Series of Monographs and Advanced Texts, 1988.
- [25] A. O. Prishlyak & K. I. Mischenko, Classification of noncompact surfaces with boundary, Methods of Functional Analysis and Topology Vol. 13 (2007), no. 1, pp. 62–66

- [26] I. Richards, On the classification of noncompact surfaces, *Trans. Amer. Math. Soc.* 106 (1963), 159–269.
- [27] H. Royden, Automorphisms and isometries of Teichmüller space, in: *Advances of the Theory of Riemann Surfaces*, *Ann. Math. Stud.*, **66** (1971), 369–384.
- [28] H. Shiga, On a distance defined by the length spectrum on Teichmüller space, *Ann. Acad. Sci. Fenn. Math.* 28, No. 2, 315–326 (2003).
- [29] W. P. Thurston, *Three-dimensional geometry and topology*, Vol. 1, Princeton mathematical series **35**, Princeton University Press, 1997.

L. LIU, DEPARTMENT OF MATHEMATICS, ZHONGSHAN UNIVERSITY, 510275, GUANGZHOU, P. R. CHINA

E-mail address: `mcsllx@mail.sysu.edu.cn`

A. PAPADOPOULOS, INSTITUT DE RECHERCHE MATHÉMATIQUE AVANCÉE, UNIVERSITÉ DE STRASBOURG AND CNRS, 7 RUE RENÉ DESCARTES, 67084 STRASBOURG CEDEX - FRANCE

E-mail address: `papadopoulos@math.u-strasbg.fr`