



HAL
open science

Solar Cycle variations of f_oF_2 from IGY to 1990

N. K. Sethi, M. K. Goel, K. K. Mahajan

► **To cite this version:**

N. K. Sethi, M. K. Goel, K. K. Mahajan. Solar Cycle variations of f_oF_2 from IGY to 1990. *Annales Geophysicae*, 2002, 20 (10), pp.1677-1685. hal-00317187

HAL Id: hal-00317187

<https://hal.science/hal-00317187v1>

Submitted on 18 Jun 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Solar Cycle variations of f_oF2 from IGY to 1990

N. K. Sethi, M. K. Goel, and K. K. Mahajan

Radio and Atmospheric Sciences Division, National Physical

Received: 19 November 2001 – Revised: 27 March 2002 – Accepted: 24 April 2002

Abstract. Noontime monthly median values of F2-layer critical frequency f_oF2 (m) for some ionospheric stations representing low- and mid-latitudes are examined for their dependence on solar activity for the years 1957 (IGY) to 1990. This is the period for which ionospheric data in digital form is available in two CD-ROMs at the World Data Center, Boulder. It is observed that at mid-latitudes, f_oF2 (m) shows nearly a linear relationship with R12 (the 12-month running average of the Zurich sunspot number), though this relation is nonlinear for low-latitudes. These results indicate some departures from the existing information often used in theoretical and applied areas of space research.

Key words. Ionosphere (equatorial ionosphere; mid-latitude ionosphere; modelling and forecasting)

1 Introduction

Electron concentration in the F2-region of the ionosphere is primarily due to ionization of the neutral atmosphere by the solar UV radiations. These radiations are now known to show very definitive solar cycle variations. Consequently, electron concentrations and thus, the critical frequency of the F2-region (f_oF2) is also expected to reflect these variations. Although there were no solar UV measurements during the early years of ionospheric research, sunspots data for several decades were available and solar cycle changes in f_oF2 were detected in the very beginning of ionospheric research (see Mitra, 1952 for early works). In fact, excellent correlations between the sunspot number and the monthly mean f_oF2 were reported and a detailed analysis of ionosonde data for several stations by Jones and Gallet (1962, 1965) and later by Rush et al. (1983, 1984) helped in generating global maps of f_oF2 as a function of sunspot number and other geophysical parameters. These maps have since been used by international organizations like CCIR and URSI as predictive tools for HF propagation. An important feature of these predictive

models is that f_oF2 saturates or increases very slowly at all stations for R12 (12-month running average of sunspot number) more than 150 units. This saturation, however, is not expected from theory, since there is no evidence that solar UV flux saturates at high solar activity. On the other hand, Huang (1960) in a detailed analysis of f_oF2 data for the period 1954 to 1958 from stations in the eastern sector concluded that f_oF2 saturation is subject to diurnal and geophysical variations. The f_oF2 -R12 plots published by him show that f_oF2 does not always saturate at all stations for R12=150. Noontime f_oF2 -R12 plots for several stations, with geomagnetic latitudes varying from 48° S to 83° N for the period 1954–1964 published by Rao and Rao (1969) do not show much evidence of f_oF2 saturation for R12 above 150 units for all the stations.

However, all the above studies were based on a limited data set, involving either one solar cycle or a part of it. Ionospheric data from several stations are now available in two CD-ROMs, covering the period from 1957, the International Geophysical Year (IGY), to 1990. In the present paper, we have studied the f_oF2 (m) data obtained from these CD-ROMs for a few stations, covering low- to mid-latitudes, namely Maui (20.8° N, 203.5° E), Chungli (24.9° N, 121.2° E), Poitiers (46.6° N, 0.3° E), Slough (51.5° N, 359.4° E) and Moscow (55.5° N, 37.3° E). We have also used the data for the Indian station Kodaikanal (10.2° N, 77.5° E). Results of our analysis of 30 years of ionospheric data for the above mentioned stations indicate some departures from the existing information on the noontime R12- f_oF2 relationship, often used in theoretical and applied areas of space research. This paper presents an improvement on the existing information.

2 Database

We have used the Ionospheric database available on the two CD-ROMS obtained from the World Data Center, Boulder and the Ionospheric data bulletins published by the National




Fig. 1. Shows the plots of observed monthly median noon *foF2* values against *R12* during summer. The regression fits are shown as solid line. The left panels show low latitude stations, while the mid-latitude stations are shown on the right panel of the figure.




Fig. 2. Same as Fig. 1 but for winter.




Fig. 3. Same as Fig. 1 but for equinox.




Fig. 4. Shows the variation of regressions fits with $R12$ along with the CCIR and IRI models for summer. Low- and mid-latitude stations are shown on left and right panels of the figure.




Fig. 5. Same as Fig. 4 but for winter.




Fig. 6. Same as Fig. 4 but for equinox.

Physical Laboratory (NPL), New Delhi, India. These bulletins contain ionospheric data for Indian stations. In the present paper, we have selected six stations covering locations from low- to mid-latitudes for the period 1957–1990. The quality of data for these locations is known to be excellent, since these data have undergone some stringent tests to bring these to a “highest possible level of correctness before being archived and made available to users” (NGDC, 1994), and the data for most of the stations exist on a continuous basis for more than 30 years. The noontime monthly median $foF2$ (m) values have been examined with respect to their dependence on conventional solar index R12 (12-month running average of sunspot number). We have selected the months of June, April and January as representatives of summer, equinox and winter seasons respectively.

3 Analysis and results

In order to determine the relationships between the $foF2$ (m) and R12, a regression analysis was carried out for each station, and it was observed that the relationship is nearly linear at mid-latitudes. For low-latitudes, a second degree fit gives a better correlation. This is demonstrated in Fig. 1 to 3, where the variation of $foF2$ (m) with R12 is shown for summer, winter and equinox respectively. These figures contain both the observed data and regression fits. It can be observed from Fig. 1–3 (right panels) that at mid-latitudes, $foF2$ (m) varies more or less linearly with R12, right up to the highest level of solar activity recorded so far. However, during summer, the variation of $foF2$ (m) with R12 is nonlinear. At low-latitudes (left panels), however, $foF2$ (m) shows more or less a quadratic variations with R12 up to around 150 during all the seasons. After R12 of 150 it is observed that during winter and equinox, $foF2$ (m) either shows saturation or a decrease. However, in summer, the increase of $foF2$ (m) with R12 is faster at lower levels of solar activity and very slow at higher levels.

Since the linear relationship seen by us between $foF2$ (m) and R12 was observed only at mid-latitudes, we examined $foF2$ (m) data for a few more mid-latitude stations (Washington (38.7° N, 252.1° E), Boulder (40.0° N, 254.7° E), Rostov (47.2° N, 39.7° E)). Here too similar behaviour was observed.

4 Comparison with empirical models

International Reference Ionosphere (IRI) is the most widely used empirical model for upper atmospheric studies and is being updated and improved off and on following the annual IRI Workshops. This model used the CCIR coefficients up to its 1990 version (Bilitza, 1990). This version, as expected, shows $foF2$ (m) saturation for R12 above 150 units. In the 1995 version, the solar index R12 was replaced by the ionospheric index IG12 (Bilitza, 1997) and it shows $foF2$ (m) saturation for R12 above 200. The latest IRI model (Bilitza, 2001) again uses the ionospheric index IG12 in place

of solar index R12. Figures 4–6 show the comparisons of regression curves with the CCIR and IRI models at all the six stations for summer, winter and equinox respectively. It can be observed from these figures that during all the seasons, for low-latitudes (left panels), the agreement between the observed fit and the models is fairly good at low solar activity, with somewhat lower values for models. During high solar activity, it can be noticed that in the observed fit, there is a tendency of either saturation or fall in $foF2$ (m) after R12 of 150, in agreement with CCIR, whereas IRI shows saturation beyond R12 of 200. Further, it can be seen that during equinox, the observed fit shows lower values of $foF2$ (m) at high solar activity as compared to model values. For mid-latitudes, as seen in Figs. 4–6 (right panels), the agreement between the IRI and the observed fit is found to be fairly good for summer and winter months, during the low solar activity periods. This agreement becomes excellent during equinox, right up to R12 of 200. However, CCIR shows saturation for R12 above 150 units. During high solar activity, as shown in the right panels of Figs. 4–6, $foF2$ (m) values of the fit are higher than those obtained from the IRI, especially during winter and summer months. It is to be noted that both the CCIR and IRI show the saturation effect in $foF2$ (CCIR beyond R12 of 150 and IRI beyond R12 of 200). It seem to us that the saturation effect may not occur at all at mid-latitudes, even beyond R12 > 200. This can be inferred from the figures if the observed values are extrapolated for R12 beyond 200. However, some stations like Slough and Moscow might be the exceptions during summer.

5 Conclusion

Detailed analyses of a long series of solar and ionospheric observations indicate that $foF2$ (m) at mid-latitudes increases linearly with R12 right up to the highest level of solar activity so far. However, at low-latitudes, the observed $foF2$ (m) values show a tendency to saturate, and in some cases even fall beyond a certain level of solar activity during all the seasons. Comparative studies with the CCIR and IRI show, in general, a fairly good agreement with the observed $foF2$ during low solar activity period. However, major discrepancies exist during high solar activity period, at both the low- and mid-latitudes, especially for equinox and winter months respectively. The observations do not provide much evidence of $foF2$ saturation at mid-latitudes.

Acknowledgements. We are thankful to World Data Center for supplying us with two CD-ROMs containing ionospheric data. K. K. Mahajan is thankful to the Council of Scientific and Industrial Research (CSIR) for the award of Emeritus Scientist position during the progress of this work.

Topical Editor M. Lester thanks two referees for their help in evaluating this paper.

References

- Bilitza, D.: (Ed) NSSDC 90–22, World Data Center A, Rockets and Satellites Greenbelt, Md, USA, November, 1990.
- Bilitza, D.: International Reference Ionosphere – Status 1995/96, *Adv. Space Res.*, 20, 9, 1751–1754, 1997.
- Bilitza, D.: International Reference Ionosphere 2000, *Radio Science*, 36, 2, 261–275, 2001.
- Huang, C.-M.: Some abnormalities in the variations of F2-layer critical frequency during the period of high solar activity of solar cycle 8–19, *J. Geophys. Res.*, 65, 897–906, 1960.
- Jones, W. B. and Gallet, R. M.: The representation of diurnal and geographic variations of ionospheric data by numerical methods, *Telecomm. J.*, 29, 129–147, 1962.
- Jones, W. B. and Gallet, R. M.: The representation of diurnal and geographic variations of ionospheric data by numerical methods, *Telecomm. J.*, 32, 18–28, 1965.
- Mitra, S. K.: *The Upper Atmosphere*, 2nd edition, The Asiatic Society, Calcutta, 1952.
- NGDC: National Geophysical Data Center, NOAA, Mail Code E / GC2, 325 Broadway, Boulder, Colorado, 80303 USA., 1994.
- Rao, M. S. V. G. and Rao, R. S.: The hysteresis variation in F2-layer parameters, *J. Atmos. Terr. Phys.*, 31, 1119–1125, 1969.
- Rush, C. M., PoKempner, M., Anderson, D. N., Stewart, F. G., and Perry, J.: Improving ionospheric maps using theoretically derived values of f_oF_2 , *Radio Science*, 18, 95–107, 1983.
- Rush, C. M., PoKempner, M., Anderson, D. N., Perry, J., Stewart, F. G., and Reasoner, R.: Maps of f_oF_2 derived from observations and theoretical data., *Radio Science*, 19, 1083–1097, 1984.