



HAL
open science

On the Schrodinger equation outside strictly convex obstacles

Oana Ivanovici

► **To cite this version:**

| Oana Ivanovici. On the Schrodinger equation outside strictly convex obstacles. 2008. hal-00319146v2

HAL Id: hal-00319146

<https://hal.science/hal-00319146v2>

Preprint submitted on 18 Dec 2008 (v2), last revised 4 Sep 2009 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the Schrödinger equation outside strictly convex obstacles

Oana Ivanovici *

Universite Paris-Sud, Orsay,
Mathematiques, Bat. 430, 91405 Orsay Cedex, France
oana.ivanovici@math.u-psud.fr

Abstract

We prove sharp Strichartz estimates for the semi-classical Schrödinger equation on a compact Riemannian manifold with smooth, strictly geodesically concave boundary. We deduce classical Strichartz estimates for the Schrödinger equation outside a strictly convex obstacle, local existence for the H^1 -critical (quintic) Schrödinger equation and scattering for the sub-critical Schrödinger equation in $3D$.

1 Introduction

Let (M, g) be a Riemannian manifold of dimension $n \geq 2$. Strichartz estimates are a family of dispersive estimates on solutions $u(x, t) : M \times [-T, T] \rightarrow \mathbb{C}$ to the Schrödinger equation

$$i\partial_t u + \Delta_g u = 0, \quad u(x, 0) = u_0(x), \quad (1.1)$$

where Δ_g denotes the Laplace-Beltrami operator on (M, g) . In their most general form, local Strichartz estimates state that

$$\|u\|_{L^q([-T, T], L^r(M))} \leq C \|u_0\|_{H^s(M)}, \quad (1.2)$$

where $H^s(M)$ denotes the Sobolev space over M and $2 \leq q, r \leq \infty$ satisfy $(q, r, n) \neq (2, \infty, 2)$ (for the case $q = 2$ see [26]) and are given by the scaling admissibility condition

$$\frac{2}{q} + \frac{n}{r} = \frac{n}{2}. \quad (1.3)$$

*The author was supported by grant A.N.R.-07-BLAN-0250

In \mathbb{R}^n and for $g_{ij} = \delta_{ij}$, Strichartz estimates in the context of the wave and Schrödinger equations have a long history, beginning with Strichartz pioneering work [38], where he proved the particular case $q = r$ for the wave and (classical) Schrödinger equations. This was later generalized to mixed $L_t^q L_x^r$ norms by Ginibre and Velo [17] for Schrödinger equations, where (q, r) is sharp admissible and $q > 2$; the wave estimates were obtained independently by Ginibre-Velo [19] and Lindblad-Sogge [28], following earlier work by Kapitanski [24]. The remaining endpoints for both equations were finally settled by Keel and Tao [26]. In that case $s = 0$ and $T = \infty$; (see also Kato [25], Cazenave-Weissler [11]). Estimates for the flat 2-torus were shown by Bourgain [4] to hold for $q = r = 4$ and any $s > 0$.

In the variable coefficients case, even without boundaries, the situation is much more complicated: we simply recall the pioneering work of Staffilani and Tataru [37], dealing with compact, non trapping perturbations of the flat metric, and recent work of Bouclet and Tzvetkov [3] which considerably weakens the decay of the perturbation (retaining the non-trapping character at spatial infinity). On compact manifolds without boundaries, Burq, Gerard and Tzvetkov [8] established Strichartz estimates with $s = 1/p$, hence with a loss of derivatives when compared to the case of flat geometries. Recently, Blair, Smith and Sogge [2] improved on the current results for compact (M, g) where either $\partial M \neq \emptyset$, or $\partial M = \emptyset$ and g Lipschitz, by showing that Strichartz estimates hold with a loss of $s = 4/3p$ derivatives. This appears to be the natural analog of the estimates of [8] for the general boundaryless case.

In this paper we prove that Strichartz estimates for the semi-classical Schrödinger equation also hold on compact Riemannian manifolds with smooth, strictly geodesically concave boundaries. By the last condition we understand that the second fundamental form on the boundary of the manifold is strictly positive definite.

Assumptions 1.1. Let (M, g) be a smooth n -dimensional compact Riemannian manifold with C^∞ boundary. We shall assume that $n \geq 2$ and that ∂M is strictly geodesically concave throughout. Let Δ_g be the Laplace-Beltrami operator associated to g on M , acting on $L^2(M)$, with domain $H^2(M) \cap H_0^1(M)$. We assume that we can globally write

$$\Delta_g = \sum_{j,k=2}^n g^{j,k}(x) \partial_j \partial_k + \sum_{j=1}^n a_j(x) \partial_j, \quad (1.4)$$

where the coefficients belong to a bounded set of C^∞ and the principal part is uniformly elliptic.

Let $0 < \alpha_0 \leq 1/2$, $2 \leq \beta_0$, $\Psi \in C_0^\infty(\mathbb{R} \setminus \{0\})$ be compactly supported in the interval (α_0, β_0) . We introduce the operator $\Psi(-h^2 \Delta_g)$ using the Dynkin-Helffer-Sjöstrand formula

[15] and refer to [31], [15] or [23] for a complete overview of its properties (see also [8] for compact manifolds without boundaries).

Definition 1.2. Given $\Psi \in C_0^\infty(\mathbb{R})$ we have

$$\Psi(-h^2\Delta_g) = -\frac{1}{\pi} \int_{\mathbb{C}} \bar{\partial} \tilde{\Psi}(z) (z + h^2\Delta_g)^{-1} dL(z),$$

where $dL(z)$ denotes the Lebesgue measure on \mathbb{C} and $\tilde{\Psi}$ is an almost analytic extension of Ψ , e.g., with $\langle z \rangle = (1 + |z|^2)^{1/2}$, $N \geq 0$,

$$\tilde{\Psi}(z) = \left(\sum_{m=0}^N \partial^m \Psi(\operatorname{Re} z) (i \operatorname{Im} z)^m / m! \right) \tau(\operatorname{Im} z / \langle \operatorname{Re} z \rangle),$$

where τ is a non-negative C^∞ function such that $\tau(s) = 1$ if $|s| \leq 1$ and $\tau(s) = 0$ if $|s| \geq 2$.

Our main result reads as follows:

Theorem 1.3. *Under the Assumptions 1.1, given (q, r) satisfying the scaling condition (1.3), $q > 2$ and $T > 0$ sufficiently small, there exists a constant $C = C(T) > 0$ such that the solution $v(x, t)$ of the semi-classical Schrödinger equation on $M \times \mathbb{R}$ with Dirichlet boundary condition*

$$\begin{cases} ih\partial_t v + h^2\Delta_g v = 0 & \text{on } M \times \mathbb{R}, \\ v(x, 0) = \Psi(-h^2\Delta_g)v_0(x), \\ v|_{\partial M} = 0 \end{cases} \quad (1.5)$$

satisfies

$$\|v\|_{L^q((-T, T), L^r(M))} \leq Ch^{-\frac{n}{2}(\frac{1}{2} - \frac{1}{r})} \|\Psi(-h^2\Delta_g)v_0\|_{L^2(M)}. \quad (1.6)$$

Remark 1.4. An example of compact manifold with smooth, strictly concave boundary is given by the Sinai billiard (i.e. the complementary of a strictly convex obstacle on a cube of \mathbb{R}^n with periodic boundary conditions).

We deduce from Theorem 1.3 and [23, Thm.1.1] (see also Lemma 3.3), as in [8], the following Strichartz estimates with derivative loss:

Corollary 1.5. *Under the Assumptions 1.1, given (q, r) satisfying the scaling condition (1.3), $q > 2$ and I any finite time interval, there exists a constant $C = C(I) > 0$ such that the solution $u(x, t)$ of the (classical) Schrödinger equation on $M \times \mathbb{R}$ with Dirichlet boundary condition*

$$\begin{cases} i\partial_t u + \Delta_g u = 0, & \text{on } M \times \mathbb{R}, \\ u(x, 0) = u_0(x), \quad u|_{\partial M} = 0 \end{cases} \quad (1.7)$$

satisfies

$$\|u\|_{L^q(I, L^r(M))} \leq C(I) \|u_0\|_{H^{\frac{1}{q}}(M)}. \quad (1.8)$$

The proof of Theorem 1.3 is based on the finite speed of propagation of the semi-classical flow (see Lebeau [27]) and the energy conservation which allow us to use the arguments of Smith and Sogge [34] for the wave equation: using the Melrose and Taylor parametrix for the stationary wave (see [29], [30] or Zworski [40]) we obtain, by Fourier transform in time a parametrix for the Schrödinger operator near a "glancing" point. Since in the elliptic and hyperbolic regions the solution of (1.9) will clearly satisfy the same Strichartz estimates as on a manifold without boundary, we need to restrict our attention only on the glancing region.

As an application of Theorem 1.3 we prove classical, global Strichartz estimates for the Schrödinger equation outside a strictly convex domain in \mathbb{R}^n .

Assumptions 1.6. Let $\Omega = \mathbb{R}^n \setminus \Theta$, where Θ is a compact with smooth boundary. We assume that $n \geq 2$ and that $\partial\Omega$ is strictly geodesically concave throughout. Let $\Delta_D = \sum_{j=1}^n \partial_j^2$ denote the Dirichlet Laplace operator (with constant coefficients) on Ω .

Theorem 1.7. *Under the Assumptions 1.6, given (q, r) satisfying the scaling condition (1.3), $q > 2$ and $u_0 \in L^2(\Omega)$, there exists a constant $C > 0$ such that the solution $u(x, t)$ of the Schrödinger equation on $\Omega \times \mathbb{R}$ with Dirichlet boundary condition*

$$\begin{cases} i\partial_t u + \Delta_D u = 0, & \text{on } \Omega \times \mathbb{R}, \\ u(x, 0) = u_0(x), \\ u|_{\partial\Omega} = 0 \end{cases} \quad (1.9)$$

satisfies

$$\|u\|_{L^q(\mathbb{R}, L^r(\Omega))} \leq C \|u_0\|_{L^2(\Omega)}. \quad (1.10)$$

The proof of Theorem 1.7 combines several arguments: firstly, we perform a time rescaling, first used by Lebeau [27] in the context of control theory, which transforms the equation into a semi-classical problem for which we can use the local in time semi-classical Strichartz estimates proved in Theorem 1.3. Secondly, we adapt a result of Burq [5] which provides Strichartz estimates without loss for a non-trapping problem, with a metric that equals the identity outside a compact set. The proof relies on a local smoothing effect for the free evolution $\exp(it\Delta_D)$, first observed in the case of the flat space in the works of Constantin and Saut [14], Doi [16], Burq, Gérard and Tzvetkov [7] in the non-trapping case. Following a strategy suggested by Staffilani and Tataru [37], we prove that away from the obstacle the free evolution enjoys the Strichartz estimates exactly as for the free space.

We give two applications of Theorem 1.7 : the first one is a local existence result for the quintic Schrödinger equation in $3D$, while the second one is a scattering result for the subcritical (sub-quintic) Schrödinger equation in $3D$ domains.

Theorem 1.8. (*Local existence for the quintic Schrödinger equation*) Let Ω be a three dimensional Riemannian manifold satisfying the Assumptions 1.6. Let $T > 0$ and $u_0 \in H_0^1(\Omega)$. Then there exists a unique solution $u \in C([0, T], H_0^1(\Omega)) \cap L^5((0, T], W^{1,30/11}(\Omega))$ of the quintic nonlinear equation

$$i\partial_t u + \Delta_D u = \pm |u|^4 u \text{ on } \Omega \times \mathbb{R}, \quad u|_{t=0} = u_0 \text{ on } \Omega, \quad u|_{\partial\Omega} = 0. \quad (1.11)$$

Moreover, for any $T > 0$, the flow $u_0 \rightarrow u$ is Lipschitz continuous from any bounded set of $H_0^1(\Omega)$ to $C([-T, T], H_0^1(\Omega))$. If the initial data u_0 has sufficiently small H^1 norm, then the solution is global in time.

Theorem 1.9. (*Scattering for subcritical Schrödinger equation*) Let Ω be a three dimensional Riemannian manifold satisfying the Assumptions 1.6. Let $3 \leq p < 5$ and $u_0 \in H_0^1(\Omega)$. Then the global in time solution of the defocusing Schrödinger equation

$$i\partial_t u + \Delta_D u = |u|^{p-1} u, \quad u|_{t=0} = u_0 \text{ on } \Omega, \quad u|_{\partial\Omega} = 0 \quad (1.12)$$

scatters in $H_0^1(\Omega)$. If $p = 5$ and the gradient ∇u_0 of the initial data has sufficiently small L^2 norm, then the global solution of the critical Schrödinger equation scatters in $H_0^1(\Omega)$.

Results for the Cauchy problem associated to the critical wave equation outside a strictly convex obstacle were obtained by Smith and Sogge [34]. Their result was a consequence of the fact that the Strichartz estimates for the Euclidian wave equation also hold on Riemannian manifolds with smooth, compact and strictly concave boundaries.

In [9], Burq, Lebeau and Planchon proved that the defocusing quintic wave equation with Dirichlet boundary conditions is globally wellposed on $H^1(M) \times L^2(M)$ for any smooth, compact domain $M \subset \mathbb{R}^3$. Their proof relies on L^p estimates for the spectral projector obtained by Smith and Sogge [35]. A similar result for the defocusing critical wave equation with Neumann boundary conditions was obtained in [10].

In the case of Schrödinger equation in $\mathbb{R}^3 \times \mathbb{R}_t$, Colliander, Keel, Staffilani, Takaoka and Tao [13] established global well-posedness and scattering for energy-class solutions to the quintic defocusing Schrödinger equation (1.11), which is energy-critical. When the domain is the complementary of an obstacle in \mathbb{R}^3 , non-trapping but not convex, the counterexamples constructed in [22] for the wave equation suggest that losses are likely to occur in the Strichartz estimates for the Schrödinger equation too. In this case Burq, Gerard and Tzvetkov [7] proved global existence for subcubic defocusing nonlinearities and Anton [1] for the cubic case. Recently, Planchon and Vega [32] improved the local well-posedness theory

to H^1 -subcritical (subquintic) nonlinearities for $n = 3$. Theorem 1.9 is proved in [32] in the case of the exterior of a star-shaped domain for the particular case $p = 3$, using

$$\|e^{it\Delta_D}u_0\|_{L^4_{t,x}} \lesssim \|u_0\|_{\dot{H}^{1/4}},$$

but since this estimate fails to provide control of $L^4_t L^\infty_x$ one has to use local smoothing estimates close to the boundary, and Strichartz estimates for the usual Laplacian on \mathbb{R}^3 away from it, the sub-criticality with respect to H^1 compensating the weakness of the local smoothing estimate. Here we give a simpler proof on the exterior of a strictly convex obstacle and for every $3 < p < 5$ using the Strichartz estimates (1.10).

Acknowledgements

The author would like to thank Nicolas Burq for many helpful and stimulating discussions and Fabrice Planchon for having suggested to her the applications. She would also like to thank Michael Taylor for having sent her the manuscript "Boundary problems for the wave equations with grazing and gliding rays" and the referees for helpful comments and suggestions on an early version of this paper which greatly improved the presentation.

2 Estimates for semi-classical Schrödinger equation in a compact domain with strictly concave boundary

In this section we prove Theorem 1.3. In what follows Assumptions 1.1 are supposed to hold. We may assume that the metric g is extended smoothly across the boundary, so that M is a geodesically concave subset of a complete, compact Riemannian manifold \tilde{M} . By the free semi-classical Schrödinger equation we mean the semi-classical Schrödinger equation on \tilde{M} , where the data v_0 has been extended to \tilde{M} by an extension operator preserving the Sobolev spaces. By a broken geodesic in M we mean a geodesic that is allowed to reflect off ∂M according to the reflection law for the metric g .

2.1 Restriction in a small neighborhood of the boundary. Elliptic and hyperbolic regions

We consider $\delta > 0$ a small positive number and for $T > 0$ small enough we set

$$S(\delta, T) := \{(x, t) \in M \times [-T, T] \mid \text{dist}(x, \partial M) < \delta\}.$$

On the complement of $S(\delta, T)$ in $M \times [-T, T]$, the solution $v(x, t)$ equals the solution of the semi-classical Schrödinger equation on \mathbb{R}^n for which Strichartz estimates are known by the work of Staffillani and Tataru [37], thus it suffices to establish Strichartz estimates for the norm of v over $S(\delta, T)$.

We show that in order to prove Theorem 1.3 it will be sufficient to consider only data v_0 supported outside a small neighborhood of the boundary. Recall that in [27] Lebeau proved that if Ψ is supported in an interval $[\alpha_0, \beta_0]$ and if $\varphi \in C_0^\infty(\mathbb{R})$ is equal to 1 near the interval $[-\beta_0, -\alpha_0]$, then for t in a bounded set (and for $D_t = \frac{1}{i}\partial_t$) one has

$$\forall N \geq 1, \quad \exists C_N > 0 \quad |(1 - \varphi)(hD_t) \exp(ih\Delta_g)\Psi(-h^2\Delta_g)v_0| \leq C_N h^N. \quad (2.1)$$

For δ and T sufficiently small, let $\chi(x, t) \in C_0^\infty$ be compactly supported and be equal to 1 on $S(\delta, T)$. Let $t_0 > 0$ be such that $T = t_0/4$ and let $A \in C^\infty(\mathbb{R}^n)$, $A = 0$ near ∂M , $A = 1$ outside a neighborhood of the boundary be such that every broken bicharacteristic γ starting at $t = 0$ from the support of $\chi(x, t)$ and for $-\tau \in [\alpha_0, \beta_0]$ satisfies

$$\text{dist}(\gamma(t), \text{supp}(1 - A)) > 0, \quad \forall t \in [-2t_0, -t_0]. \quad (2.2)$$

Let $\psi \in C^\infty(\mathbb{R})$, $\psi(t) = 0$ for $t \leq -2t_0$, $\psi(t) = 1$ for $t > -t_0$ and set

$$w(x, t) = \psi(t) \exp(ih\Delta_g)\Psi(-h^2\Delta_g)v_0.$$

Then w satisfies

$$\begin{cases} ih\partial_t w + h^2\Delta_g w = ih\psi'(t)e^{ih\Delta_g}\Psi(-h^2\Delta_g)v_0, \\ w|_{\partial\Omega \times \mathbb{R}} = 0, \quad w|_{t \leq -2t_0} = 0, \end{cases}$$

and writing Duhamel formula we have

$$w(x, t) = \int_{-2t_0}^t e^{i(t-s)h\Delta_g} \psi'(s) e^{ish\Delta_g} \Psi(-h^2\Delta_g) v_0 ds.$$

Notice that $w(x, t) = v(x, t)$ if $t \geq -t_0$, hence for $t \in [-t_0, T]$ we can write

$$v(x, t) = \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_g} \psi'(s) e^{ish\Delta_g} \Psi(-h^2\Delta_g) v_0 ds. \quad (2.3)$$

In particular, for $t \in [-T, T]$, $T = t_0/4$, $v(x, t) = w(x, t)$ is given by (2.3). We want to estimate the $L_t^q L_x^r$ norms of $v(x, t)$ for (x, t) on $S(\delta, T)$ where $v = \chi v$. Let

$$v_Q(x, t) = \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_g} \psi'(s) Q(x) e^{ish\Delta_g} \Psi(-h^2\Delta_g) v_0 ds, \quad Q \in \{A, 1 - A\},$$

then $v = v_A + v_{1-A}$, where v_{1-A} solves

$$\begin{cases} ih\partial_t v_{1-A} + h^2\Delta_g v_{1-A} = ih\psi'(t)(1-A)e^{ith\Delta_g}\Psi(-h^2\Delta_g)v_0, \\ v_{1-A}|_{\partial M \times \mathbb{R}} = 0, \quad v_{1-A}|_{t < -2t_0} = 0. \end{cases}$$

We apply Proposition 4.8 from the Appendix with $Q = 1 - A$, $\tilde{\psi} = \psi'$ to deduce that if $\rho_0 \in WF_b(v_{1-A})$ then the broken bicharacteristic starting from ρ_0 must intersect $WF_b((1-A)v) \cap \{t \in [-2t_0, -t_0]\}$. Since we are interested in estimating the norm of v on $S(\delta, T)$ it is enough to consider only $\rho_0 \in WF_b(\chi v_{1-A})$. Thus, if γ is a broken bicharacteristic starting at $t = 0$ from ρ_0 , $-\tau \in [\alpha_0, \beta_0]$, then Proposition 4.8 implies that for some $t \in [-2t_0, -t_0]$, $\gamma(t)$ must intersect $WF_b((1-A)v)$. On the other hand from (2.2) this implies (see Definition 4.2) that for every $\sigma \geq 0$

$$\forall N \geq 0 \quad \exists C_N > 0 \quad \|\chi v_{1-A}\|_{H^\sigma(M \times \mathbb{R})} \leq C_N h^N. \quad (2.4)$$

We are thus reduced to estimating $v(x, t)$ for initial data supported outside a small neighborhood of the boundary. Indeed, suppose that the estimates (1.6) hold true for any initial data compactly supported where $A \neq 0$. It follows from (2.3), (2.4) that

$$\begin{aligned} \|\chi v_A\|_{L^q((-T, T), L^r(M))} &\leq \|\psi'(s)A(x)e^{ish\Delta_g}\Psi(-h^2\Delta_g)v_0\|_{L^1(s \in (-2t_0, -t_0), L^2(M))} \lesssim \\ &\left(\int_{-2t_0}^{-t_0} |\Psi'(s)| ds \right) \|\Psi(-h^2\Delta_g)v_0\|_{L^2(M)} = \|\Psi(-h^2\Delta_g)v_0\|_{L^2(M)}, \end{aligned} \quad (2.5)$$

where we used the fact that the semi-classical Schrödinger flow $\exp(ihs\Delta_g)\Psi(-h^2\Delta_g)$, which maps data at time 0 to data at time s , is an isomorphism on $H^\sigma(M)$ for every $\sigma \geq 0$.

Remark 2.1. Notice that when dealing with the wave equation, since the speed of propagation is exact, one can take $\psi(t) = 1_{\{t \geq -t_0\}}$ for some small $t_0 \geq 0$ and reduce the problem to proving Strichartz estimates for the flow $\exp(ih(t_0 + \cdot)\Delta_g)\Psi(-h^2\Delta_g)$ and initial data compactly supported outside a small neighborhood of ∂M . This was precisely the strategy followed by Smith and Sogge [34].

Let Δ_0 denote the Laplacian on \tilde{M} coming from the extension of the metric g smoothly across the boundary ∂M . We let \mathcal{M} denote the outgoing solution to the Dirichlet problem for the semiclassical Schrödinger operator on $M \times \mathbb{R}$. Thus, if g is a function on $\partial M \times \mathbb{R}$ which vanishes for $t \leq -2t_0$, then $\mathcal{M}g$ is the solution on $M \times \mathbb{R}$ to

$$\begin{cases} ih\partial_t \mathcal{M}g + h^2\Delta_g \mathcal{M}g = 0, \\ \mathcal{M}g|_{\partial M \times \mathbb{R}} = g. \end{cases} \quad (2.6)$$

Then, for $t \in [-t_0, T]$ and data f supported outside a small neighborhood of the boundary and localized at frequency $1/h$, i.e. such that $f = \Psi(-h^2\Delta_g)f$, we have

$$\begin{aligned}\chi v_A(x, t) &= \chi \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_g} \psi'(s) A(x) e^{ish\Delta_g} f ds = \\ &= \chi \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_0} \psi'(s) A(x) e^{ish\Delta_0} f ds - \mathcal{M} \left(\chi \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_0} \psi'(s) A(x) e^{ish\Delta_0} f ds \Big|_{\partial M \times \mathbb{R}} \right).\end{aligned}$$

The cotangent bundle of $\partial M \times \mathbb{R}$ is divided into three disjoint sets: the hyperbolic and elliptic regions where the Dirichlet problem is respectively hyperbolic and elliptic, and the glancing region which is the boundary between the two.

Let local coordinates be chosen such that $M = \{(x', x_n) | x_n > 0\}$ and $\Delta_g = \partial_{x_n}^2 - r(x, D_{x'})$. A point $(x', t, \eta', \tau) \in T^*(\partial M \times \mathbb{R})$ is classified as one of three distinct types. It is said to be *hyperbolic* if $-\tau + r(x', 0, \eta') \geq c > 0$ for some $c > 0$, so that there are two distinct nonzero real solutions η_n to $\tau - r(x', 0, \eta') = \eta_n^2$. These two solutions yield two distinct bicharacteristics, one of which enters M as t increases (the *incoming ray*) and one which exits M as t increases (the *outgoing ray*). The point is *elliptic* if $-\tau + r(x', 0, \eta') \leq -c < 0$, so there are no real solutions η_n to $\tau - r(x', 0, \eta') = \eta_n^2$. In the remaining case $-\tau + r(x', 0, \eta') \in [-c, c]$, $c > 0$ small enough, there is a unique solution which yields a glancing ray, and the point is said to be a *glancing point*. We decompose the identity operator into

$$\text{Id}(x, t) = \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}((x'-y')\eta' + (t-s)\tau)} (\chi_h + \chi_e + \chi_{gl})(y', \eta', \tau) d\eta' d\tau,$$

where at (y', η', τ) we have

$$\chi_h := \mathbf{1}_{\{-\tau + r(y', 0, \eta') \geq c\}}, \quad \chi_e := \mathbf{1}_{\{-\tau + r(y', 0, \eta') \leq -c\}}, \quad \chi_{gl} := \mathbf{1}_{\{-\tau + r(y', 0, \eta') \in [-c, c]\}},$$

for some $c > 0$ sufficiently small. The corresponding operators with symbols χ_h, χ_e , denoted Π_h, Π_e , respectively, are pseudo-differential cutoffs essentially supported inside the hyperbolic and elliptic regions, while the operator with symbol χ_{gl} , denoted Π_{gl} , is essentially supported in a small set around the glancing region. Thus, on $S(\delta, T)$ we can write χv_A as the sum of four terms

$$\begin{aligned}\chi \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_g} \psi'(s) A(x) e^{ish\Delta_g} f ds &= \chi \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_0} \psi'(s) A(x) e^{ish\Delta_0} f ds - \\ &- \sum_{\Pi \in \{\Pi_e, \Pi_h, \Pi_{gl}\}} \mathcal{M} \Pi \left(\chi \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_0} \psi'(s) A(x) e^{ish\Delta_0} f ds \Big|_{\partial M \times \mathbb{R}} \right).\end{aligned}\tag{2.7}$$

Remark 2.2. For the first term in the right hand side, $\chi \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_0} \psi'(s) A(x) e^{ish\Delta_0} f ds$, the desired estimates follow as in the boundaryless case by the results of Staffilani and Tataru [37] (since we considered the extension of the metric g across the boundary to be smooth).

2.1.1 Elliptic region

Using the compactness argument of the proof of Proposition 4.7 from the Appendix, together with the inclusion (which follows from Proposition 4.3 in the Appendix)

$$WF_b(\chi \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_0} \psi'(s) A(x) e^{ish\Delta_0} f ds |_{\partial M \times \mathbb{R}}) \subset \mathcal{H} \cup \mathcal{G},$$

where \mathcal{H} and \mathcal{G} denote the hyperbolic and the glancing regions, respectively, it follows that the elliptic part satisfies for all $\sigma \geq 0$

$$\mathcal{M}\Pi_e(\chi \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_0} \psi'(s) A(x) e^{ish\Delta_0} f ds |_{\partial M \times \mathbb{R}}) = O(h^\infty) \|f\|_{H^\sigma(M)}.$$

For the definition and properties of the b -wave front set see Appendix.

2.1.2 Hyperbolic region

If local coordinates are chosen such that $M = \{(x', x_n) | x_n > 0\}$, on the essential support of Π_h the forward Dirichlet problem can be solved locally, modulo smoothing kernels, on an open set in $\tilde{M} \times \mathbb{R}$ around ∂M . Precisely, near a hyperbolic point, the solution v to (1.5) can be decomposed modulo smoothing operators into an incoming part v_- and an outgoing part v_+ where

$$v_\pm(x, t) = \frac{1}{(2\pi h)^d} \int e^{\frac{i}{h}\varphi_\pm(x, t, \xi)} \sigma_\pm(x, t, \xi, h) d\xi,$$

where the phases φ_\pm satisfy the eikonal equations

$$\begin{cases} \partial_s \varphi_{\pm+} < d\varphi_\pm, d\varphi_\pm >_g = 0, \\ \varphi_+ |_{\partial M} = \varphi_- |_{\partial M}, \quad \partial_{x_n} \varphi_+ |_{\partial M} = -\partial_{x_n} \varphi_- |_{\partial M}, \end{cases}$$

where $\langle \cdot, \cdot \rangle_g$ denotes the inner product induced by the metric g . The symbols are asymptotic expansions in h and write $\sigma_\pm(\cdot, h) = \sum_{k \geq 0} h^k \sigma_{\pm, k}$, where σ_0 solves the linear transport equation

$$\partial_s \sigma_{\pm, 0} + (\Delta_g \varphi_\pm) \sigma_{\pm, 0} + \langle d\varphi_\pm, d\sigma_{\pm, 0} \rangle_g = 0,$$

while for $k \geq 1$, $\sigma_{\pm, k}$ satisfies the non-homogeneous transport equations

$$\partial_s \sigma_{\pm, k} + (\Delta_g \varphi_\pm) \sigma_{\pm, k} + \langle d\varphi_\pm, d\sigma_{\pm, k} \rangle_g = i \Delta_g \sigma_{\pm, k-1}.$$

A direct computation shows that

$$\left\| \sum_{\pm} v_\pm \right\|_{H^\sigma(M \times \mathbb{R})}^2 \simeq \sum_{\pm} \|v_\pm\|_{H^\sigma(M \times \mathbb{R})}^2 \simeq \|v\|_{H^\sigma(M \times \mathbb{R})}^2 \simeq \|v\|_{L^\infty(\mathbb{R})H^\sigma(M)}^2.$$

Each component v_{\pm} is a solution of linear Schrödinger equation (without boundary) and consequently satisfies the usual Strichartz estimates (see Burq, Gérard and Tzvetkov [8]).

Note that $\sum_{\pm} v_{\pm}$ contains the contribution from

$$\mathcal{M}\Pi_h(\chi \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_0} \psi'(s) A(x) e^{ish\Delta_0} \Psi(-h^2 \Delta_g) v_0 ds |_{\partial M \times \mathbb{R}})$$

and a contribution from $\chi \int_{-2t_0}^{-t_0} e^{i(t-s)h\Delta_0} \psi'(s) A(x) e^{ish\Delta_0} \Psi(-h^2 \Delta_g) v_0 ds$.

2.2 Glancing region

Near a diffractive point we use the Melrose and Taylor construction for the wave equation in order to write, following Zworski [40], the solution to the wave equation as a finite sum of pseudo-differential cutoffs, each essentially supported in a suitably small neighborhood of a glancing ray. Using the Fourier transform in time we obtain a parametrix for the semi-classical Schrödinger equation (1.5) microllocally near a glancing direction and modulo smoothing operators.

2.2.1 Preliminaries. Parametrix for the wave equation near the glancing region

We start by recalling the results by Melrose and Taylor [29], [30], Zworski [40, Prop.4.1] for the wave equation near the glancing region. Let w solve the (semi-classical) wave equation on M with Dirichlet boundary conditions

$$\begin{cases} h^2 D_t^2 w + h^2 \Delta_g w = 0, & M \times \mathbb{R}, \quad w|_{\partial M \times \mathbb{R}} = 0, \\ w(x, 0) = f(x), \quad D_t w(x, 0) = g(x), \end{cases} \quad (2.8)$$

where f, g are compactly supported in M and localized at spacial frequency $1/h$, and where $D_t = \frac{1}{i} \partial_t$.

Proposition 2.3. *Near a glancing direction the solution to (2.8) writes, modulo smoothing operators*

$$w(x, t) = \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^n} e^{\frac{i}{h}(\theta(x, \xi) + it\xi_1)} \left[a(x, \xi/h) \left(A_-(\zeta(x, \xi/h)) - A_+(\zeta(x, \xi/h)) \frac{A_-(\zeta_0(\xi/h))}{A_+(\zeta_0(\xi/h))} \right) + \right. \\ \left. b(x, \xi/h) \left(A'_-(\zeta(x, \xi/h)) - A'_+(\zeta(x, \xi/h)) \frac{A_-(\zeta_0(\xi/h))}{A_+(\zeta_0(\xi/h))} \right) \right] \times \widehat{K}(f, g) \left(\frac{\xi}{h} \right) d\xi, \quad (2.9)$$

where the symbols a, b and the phases θ, ζ have the following properties: a and b are symbols of type $(1, 0)$ and order $1/6$ and $-1/6$, respectively, both of which are supported in a small

conic neighborhood of the ξ_1 axis and where K is a classical Fourier integral operator of order 0 in f and order -1 in g , compactly supported on both sides. The phases θ and ζ are real, smooth and homogeneous of degree 1 and $2/3$, respectively. If we denote Ai the Airy function then A_{\pm} are defined by $A_{\pm}(z) = Ai(e^{\mp 2\pi i/3} z)$.

Remark 2.4. If local coordinates are chosen so that Ω is given by $x_n > 0$, the phase functions θ, ζ satisfy the eikonal equations

$$\begin{cases} \xi_1^2 - \langle d\theta, d\theta \rangle_g + \zeta \langle d\zeta, d\zeta \rangle_g = 0, \\ \langle d\theta, d\zeta \rangle_g = 0, \\ \zeta(x', 0, \xi) = \zeta_0(\xi) = -\xi_1^{-1/3} \xi_n, \end{cases} \quad (2.10)$$

in the region $\zeta \leq 0$. Here $x' = (x_1, \dots, x_{n-1})$ and $\langle \cdot, \cdot \rangle_g$ denotes the inner product given by the metric g . The phases also satisfy the eikonal equations (2.10) to infinite order at $x_n = 0$ in the region $\zeta > 0$.

Remark 2.5. Notice that one can think of $A_-(\zeta)$ as the incoming contribution and of $A_+(\zeta) \frac{A_-(\zeta_0)}{A_+(\zeta_0)}$ as the outgoing one. The A_{\pm} terms are not temperate and one really exploits the cancellation

$$A_-(\zeta) - A_+(\zeta) \frac{A_-(\zeta_0)}{A_+(\zeta_0)} = e^{i\pi/3} (Ai(\zeta) - A_+(\zeta) \frac{Ai(\zeta_0)}{A_+(\zeta_0)}),$$

since $Ai(\zeta) = e^{i\pi/3} A_+(\zeta) + e^{-i\pi/3} A_-(\zeta)$. The term $Ai(\zeta)$ gives us the direct term (bicharacteristic not hitting the boundary), while the oscillatory one corresponds to the billiard ball map shift corresponding to reflection.

2.2.2 Parametrix for the solution to the semi-classical Schrödinger equation near the glancing region

Let now $v(x, t)$ be the solution of the semi-classical Schrödinger equation (1.5) where the initial data $v_0 \in L^2(M)$ is spectrally localized at spatial frequency $1/h$, i.e. $v_0(x) = \Psi(-h^2 \Delta_g) v_0(x)$. From the discussion at the beginning of this section we see that it will be enough to consider v_0 compactly supported outside some small neighborhood of ∂M . Under this assumption $\Psi(-h^2 \Delta_g) v_0$ is a well-defined pseudo-differential operator for which the results of Burq, Gérard and Tzvetkov [8, Section 2] apply.

Let $(e_{\lambda}(x))_{\lambda \geq 0}$ be the eigenbasis of $L^2(M)$ consisting in eigenfunctions of $-\Delta_g$ associated to the eigenvalues (λ^2) , so that $-\Delta_g e_{\lambda} = \lambda^2 e_{\lambda}$. We write

$$\Psi(-h^2 \Delta_g) v_0(x) \simeq \sum_{h^2 \lambda^2 \in [\alpha_0, \beta_0]} \Psi(h^2 \lambda^2) v_{\lambda} e_{\lambda}(x), \quad (2.11)$$

and hence

$$e^{ith\Delta_g}\Psi(-h^2\Delta_g)v_0(x) \simeq \sum_{h^2\lambda^2 \in [\alpha_0, \beta_0]} \Psi(h^2\lambda^2)e^{-ith\lambda^2}v_\lambda e_\lambda(x). \quad (2.12)$$

If δ denotes the Dirac function, then the Fourier transform of $v(x, t)$ writes

$$\hat{v}(x, \frac{\tau}{h}) \simeq h \sum_{h^2\lambda^2 \in [\alpha_0, \beta_0]} \Psi(h^2\lambda^2)\delta_{\{-\tau=h^2\lambda^2\}}v_\lambda e_\lambda(x). \quad (2.13)$$

For $t \in \mathbb{R}$ we can define (since \hat{v} has compact support away from 0)

$$\begin{aligned} w(x, t) &:= \frac{1}{2\pi h} \int_0^\infty e^{\frac{it\sigma}{h}} \hat{v}(x, -\frac{\sigma^2}{h}) d\sigma = -\frac{1}{4\pi h} \int_{-\infty}^0 e^{\frac{it\sqrt{-\tau}}{h}} \frac{1}{\sqrt{-\tau}} \hat{v}(x, \frac{\tau}{h}) d\tau \\ &\simeq -\frac{1}{2} \sum_{h^2\lambda^2 \in [\alpha_0, \beta_0]} \Psi(h^2\lambda^2) \left(\frac{1}{2\pi} \int_{-\infty}^0 e^{\frac{it\sqrt{-\tau}}{h}} \frac{1}{\sqrt{-\tau}} \delta_{\{-\tau=h^2\lambda^2\}} d\tau \right) v_\lambda e_\lambda(x) \\ &\simeq -\frac{1}{2} \sum_{h^2\lambda^2 \in [\alpha_0, \beta_0]} \frac{1}{h\lambda} \Psi(h^2\lambda^2) e^{it\lambda} v_\lambda e_\lambda(x). \end{aligned} \quad (2.14)$$

Then $w(x, t)$ solves the wave equation

$$\begin{cases} h^2 D_t^2 w + h^2 \Delta_g w = 0, & \text{on } M \times \mathbb{R}, & w|_{\partial M \times \mathbb{R}} = 0, \\ w(x, 0) = f_h(x), & D_t w(x, 0) = g_h(x), \end{cases} \quad (2.15)$$

where the initial data are given by

$$f_h(x) \simeq -\frac{1}{2} \sum_{h^2\lambda^2 \in [\alpha_0, \beta_0]} \frac{1}{h\lambda} \Psi(h^2\lambda^2) v_\lambda e_\lambda(x), \quad (2.16)$$

$$g_h(x) \simeq -\frac{1}{2h} \sum_{h^2\lambda^2 \in [\alpha_0, \beta_0]} \Psi(h^2\lambda^2) v_\lambda e_\lambda(x) \simeq -\frac{1}{2h} \Psi(-h^2\Delta_g) v_0(x). \quad (2.17)$$

From (2.16), (2.17) it follows that

$$h \|g_h\|_{L^2(M)} \simeq \|f_h\|_{L^2(M)} \simeq \|\Psi(-h^2\Delta_g)v_0\|_{L^2(M)}. \quad (2.18)$$

Indeed, in order to prove (2.18) notice that w defined by (2.14) satisfies in fact

$$(D_t - \sqrt{-\Delta_g})w = 0$$

and (since Δ_g and D_t commute) we have

$$f_h = w|_{t=0} = [(\sqrt{-\Delta_g})^{-1} D_t w]|_{t=0} = (\sqrt{-\Delta_g})^{-1} (D_t w|_{t=0}) = (\sqrt{-\Delta_g})^{-1} g_h.$$

Due to the spectral localization and since $g_h = -\frac{1}{2h}\Psi(-h^2\Delta_g)v_0$ we deduce (2.18).

By the L^2 continuity of the (classical) Fourier integral operator K introduced in Proposition 2.3 we deduce

$$\|K(f_h, g_h)\|_{L^2(M)} \leq C(\|f_h\|_{L^2(M)} + h\|g_h\|_{L^2(M)}) \simeq \|\Psi(-h^2\Delta_g)v_0\|_{L^2(M)}. \quad (2.19)$$

The solution $v(x, t)$ of (1.5) writes

$$\begin{aligned} v(x, t) &= \frac{1}{2\pi h} \int e^{-\frac{it\sigma^2}{h}} (-2\sigma) \hat{v}(x, -\frac{\sigma^2}{h}) d\sigma = \\ &= \frac{1}{2\pi h} \int_0^\infty e^{-i\frac{t\sigma^2}{h}} (-2\sigma) \int_{s \in \mathbb{R}} e^{-\frac{is\sigma}{h}} w(x, s) ds d\sigma. \end{aligned} \quad (2.20)$$

The next step is to use Proposition 2.8 in order to obtain a representation of $v(x, t)$ near the glancing region: notice that the glancing part of the stationary wave $\hat{w}(x, \frac{\sigma}{h})$ is given by

$$\begin{aligned} 1_{\{\sigma^2+r(x',0,\eta') \in [-c,c]\}} \hat{w}(x, \frac{\sigma}{h}) &= 1_{\{\sigma^2+r(x',0,\eta') \in [-c,c]\}} \hat{v}(x, -\frac{\sigma^2}{h}) = \\ &= 1_{\{-\tau+r(x',0,\eta') \in [-c,c]\}} \hat{v}(x, \frac{\tau}{h}), \end{aligned} \quad (2.21)$$

where $c > 0$ is sufficiently small. The equality in (2.21) follows from (2.14) and from the fact that \hat{v} is essentially supported for the second variable in the interval $[-\beta_0, -\alpha_0]$. Consequently we can apply Proposition 2.8 and determine a representation for v near the glancing region (for the Schrödinger equation) as follows

$$\begin{aligned} v(x, t) &= \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^n} e^{\frac{i}{h}(\theta(x,\xi) - t\xi_1^2)} 2\xi_1 \left[a(x, \xi/h) \left(Ai(\zeta(x, \xi/h)) - A_+(\zeta(x, \xi/h)) \frac{Ai(\zeta_0(\xi/h))}{A_+(\zeta_0(\xi/h))} \right) + \right. \\ &\quad \left. b(x, \xi/h) \left(Ai'(\zeta(x, \xi/h)) - A'_+(\zeta(x, \xi/h)) \frac{Ai(\zeta_0(\xi/h))}{A_+(\zeta_0(\xi/h))} \right) \right] \widehat{K}(f_h, g_h) \left(\frac{\xi}{h} \right) d\xi, \end{aligned} \quad (2.22)$$

where a , b and K are those defined in Proposition 2.3 and f_h, g_h are given by (2.16), (2.17). The initial data f_h, g_h are both supported, like v_0 , away from ∂M , and consequently their $\dot{H}^\sigma(M)$ norms for $\alpha < n/2$ will be comparable to the norms of the non-homogeneous Sobolev space $H^\sigma(\mathbb{R}^n)$, so we shall henceforth work with the latter norms on the data f_h, g_h .

Remark 2.6. Notice that it is enough to prove semi-classical Strichartz estimates only for the "outgoing" piece corresponding to the oscillatory term $A_+(\zeta) \frac{Ai(\zeta_0)}{A_+(\zeta_0)}$, since the direct solution has already been dealt with (see the remark following (2.7)).

We deduce from (2.19), (2.22) that in order to finish the proof of Theorem 1.3 we need only to show that the operator A_h defined, for f supported away from ∂M and spectrally localized at frequency $1/h$, i.e. such that $f = \Psi(-h^2\Delta_g)f$, by

$$A_h f(x, t) = \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^n} 2\xi_1 (a(x, \xi/h)A_+(\zeta(x, \xi/h)) + b(x, \xi/h)A'_+(\zeta(x, \xi/h))) \times \quad (2.23)$$

$$\times e^{\frac{i}{h}(\theta(x, \xi) - t\xi_1^2)} \frac{Ai(\zeta_0(\xi/h))}{A_+(\zeta_0(\xi/h))} \widehat{f}\left(\frac{\xi}{h}\right) d\xi,$$

satisfies

$$\|A_h f\|_{L^q((0, T], L^r(\mathbb{R}^n))} \leq Ch^{-n/2(1/2-1/r)} \|f\|_{L^2(\mathbb{R}^n)}. \quad (2.24)$$

Remark 2.7. We introduce a cut-off function $\chi_1 \in C_0^\infty(\mathbb{R}^n)$ equal to 1 on the support of f and equal to 0 near ∂M . Since χ_1 is supported away from the boundary it follows from [8, Prop.2.1] (which applies here in its adjoint form) that $\Psi(-h^2\Delta_g)\chi_1 f$ is a pseudo-differential operator and writes (in a patch of local coordinates)

$$\Psi(-h^2\Delta_g)\chi_1 f = d(x, hD_x)\chi_2 f + O_{L^2(M)}(h^\infty), \quad (2.25)$$

where $\chi_2 \in C_0^\infty(\mathbb{R}^n)$ is equal to 1 on the support of χ_1 and where $d(x, D_x)$ is defined for x in the suitable coordinate patch using the usual pseudo-differential quantization rule,

$$d(x, D_x)f(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{ix\xi} d(x, \xi) \widehat{f}(\xi) d\xi, \quad d \in C_0^\infty,$$

with symbol d compactly supported for $|\xi|_g^2 := \langle \xi, \xi \rangle_g \in [\alpha_0, \beta_0]$, which follows by the condition of the support of Ψ . Since the principal part of the Laplace operator Δ_g is uniformly elliptic, we can introduce a smooth radial function $\psi \in C_0^\infty([\frac{1}{\delta}\alpha_0^{1/2}, \delta\beta_0^{1/2}])$ for some $\delta \geq 1$ such that $\psi(|\xi|)d = d$ everywhere. In what follows we shall prove (2.24) where, instead of f we shall write $\psi(|\xi|)f$, keeping in mind that f is supported away from the boundary and localized at spatial frequency $1/h$.

The proof of Theorem 1.3 will be completed once we prove (2.24). In order to do that, we split the operator A_h into two parts: a main term and a diffractive term. To this end, let $\chi(s)$ be a smooth function satisfying

$$\text{supp}\chi \subset (-\infty, -1], \quad \text{supp}(1 - \chi) \subset [-2, \infty).$$

We write this operator as a sum $A_h = M_h + D_h$, by decomposing

$$A_+(\zeta) = (\chi A_+)(\zeta) + ((1 - \chi)A_+)(\zeta),$$

and letting the "main term" be defined for f like in Remark 2.7 by

$$M_h f(x, t) = \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^n} 2\xi_1 (a(x, \xi/h)(\chi A_+) (\zeta(x, \xi/h)) + b(x, \xi/h)(\chi A'_+) (\zeta(x, \xi/h))) \times \\ \times e^{\frac{i}{h}(\theta(x, \xi) - t\xi_1^2)} \frac{Ai(\zeta_0(\xi/h))}{A_+(\zeta_0(\xi/h))} \psi(|\xi|) \hat{f}\left(\frac{\xi}{h}\right) d\xi.$$

The "diffractive term" is then defined for f like before by

$$D_h f(x, t) = \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^n} 2\xi_1 (a(x, \xi/h)((1-\chi)A_+) (\zeta(x, \xi/h)) + b(x, \xi/h)((1-\chi)A'_+) (\zeta(x, \xi/h))) \times \\ \times e^{\frac{i}{h}(\theta(x, \xi) - t\xi_1^2)} \frac{Ai(\zeta_0(\xi/h))}{A_+(\zeta_0(\xi/h))} \psi(|\xi|) \hat{f}\left(\frac{\xi}{h}\right) d\xi.$$

We analyze these operators separately, following the ideas of [34]:

2.2.3 The main term M_h

To estimate the "main term" M_h we first use the fact that

$$\left| \frac{Ai(s)}{A_+(s)} \right| \leq 2, \quad s \in \mathbb{R}. \quad (2.26)$$

Consequently, since the term $\frac{Ai(\zeta_0)}{A_+(\zeta_0)}$ acts like a multiplier and so does ξ_1 which is localized in the interval $[\alpha_0, \beta_0]$ (this follows from (2.1)), the estimates for M_h will follow from showing that the operator

$$f \rightarrow \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^n} (a(x, \xi/h)(\chi A_+) (\zeta(x, \xi/h)) + b(x, \xi/h)(\chi A'_+) (\zeta(x, \xi/h))) \times \\ \times e^{\frac{i}{h}(\theta(x, \xi) - t\xi_1^2)} \psi(|\xi|) \hat{f}\left(\frac{\xi}{h}\right) d\xi \quad (2.27)$$

satisfies the same bounds like in (2.24) for f spectrally localized at frequency $1/h$. Following [40, Lemma 4.1], we write χA_+ and $(\chi A_+)'$ in terms of their Fourier transform to express the phase function of this operator,

$$\phi(t, x, \xi) = -t\xi_1^2 + \theta(x, \xi) - \frac{2}{3}(-\zeta)^{3/2}(x, \xi) \quad (2.28)$$

which satisfies the eikonal equation (2.10). We denote its symbol $c_m(x, \xi/h)$, $c_m(x, \xi) \in \mathcal{S}_{2/3, 1/3}^0(\mathbb{R}^n \times \mathbb{R}^n)$ and we also denote the operator defined in (2.27) by W_h^m , thus

$$W_h^m f(x, t) = \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^n} e^{\frac{i}{h}\phi(t, x, \xi)} c_m(x, \xi/h) \psi(|\xi|) \hat{f}\left(\frac{\xi}{h}\right) d\xi.$$

Proposition 2.8. *Let (q, r) be an admissible pair with $q > 2$, let $T > 0$ be sufficiently small and for $f = d(x, D_x)\chi_2 f + O_{L^2(\Omega)}(h^\infty)$ like in Remark 2.7 let*

$$W_h f(x, t) := \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}\phi(t, x, \xi)} c_m(x, \xi/h) \psi(|\xi|) \hat{f}\left(\frac{\xi}{h}\right) d\xi.$$

Then the following estimates hold

$$\|W_h f\|_{L^q((0, T], L^r(\mathbb{R}^n))} \leq Ch^{-n/2(1/2-1/r)} \|f\|_{L^2(\mathbb{R}^n)}. \quad (2.29)$$

In the rest of this section we prove Proposition 2.8. The first step in the proof is a TT* argument. Explicitly,

$$\widehat{W_h^*(F)}\left(\frac{\xi}{h}\right) = \int e^{-\frac{i}{h}\phi(s, y, \xi)} F(y, s) \overline{c_m(y, \xi/h)} dy ds,$$

and if we set

$$(T_h F)(x, t) = (W_h W_h^* F)(x, t) = \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}(\phi(t, x, \xi) - \phi(s, y, \xi))} c_m(x, \xi/h) \overline{c_m(y, \xi/h)} \psi^2(|\xi|) F(y, s) d\xi ds dy,$$

then inequality (2.29) is equivalent to

$$\|T_h F\|_{L^q((0, T], L^r(\mathbb{R}^n))} \leq Ch^{-n(1/2-1/r)} \|F\|_{L^{q'}((0, T], L^{r'}(\mathbb{R}^n))}, \quad (2.30)$$

where (q', r') satisfies $1/q + 1/q' = 1$, $1/r + 1/r' = 1$. To see, for instance, that (2.30) implies (2.29), notice that the dual version of (2.29) is

$$\|W_h^* F\|_{L^2(\mathbb{R}^n)} \leq Ch^{-n/2(1/2-1/r)} \|F\|_{L^{q'}((0, T], L^{r'}(\mathbb{R}^n))},$$

and we have

$$\|W_h^* F\|_{L^2(\mathbb{R}^n)}^2 = \int W_h W_h^* F \bar{F} dt dx \leq \|T_h F\|_{L^q((0, T], L^r(\mathbb{R}^n))} \|F\|_{L^{q'}((0, T], L^{r'}(\mathbb{R}^n))}.$$

Therefore we only need to prove (2.30). Since the symbols are of type $(2/3, 1/3)$ and not of type $(1, 0)$, before starting the proof of (2.30) for the operator T_h we need to make a further decomposition: let $\rho \in C_0^\infty(\mathbb{R})$ satisfying $\rho(s) = 1$ near 0, $\rho(s) = 0$ if $|s| \geq 1$. Let

$$T_h F = T_h^f F + T_h^s F,$$

where

$$T_h^s F(x, t) = \int K_h^s(t, x, s, y) F(y, s) ds dy, \quad (2.31)$$

$$K_h^s(t, x, s, y) = \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}(\phi(t, x, \xi) - \phi(s, y, \xi))} (1 - \rho(h^{-1/3}|t - s|)) c_m(x, \xi/h) \overline{c_m(y, \xi/h)} \psi^2(|\xi|) d\xi, \quad (2.32)$$

while

$$T_h^f F(x, t) = \int K_h^f(t, x, s, y) F(y, s) ds dy, \quad (2.33)$$

$$K_h^f(t, x, s, y) = \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}(\phi(t, x, \xi) - \phi(s, y, \xi))} \rho(h^{-1/3}|t - s|) c_m(x, \xi/h) \overline{c_m(y, \xi/h)} \psi^2(|\xi|) d\xi. \quad (2.34)$$

Remark 2.9. The two pieces will be handled differently. The kernel of T_h^f is supported in a suitable small set and it will be estimate by "freezing" the coefficients. To estimate T_h^s we shall use the stationary phase method for type (1, 0) symbols. For type (2/3, 1/3) symbols, these stationary phase arguments break down if $|t - s|$ is smaller than $h^{1/3}$, which motivates the decomposition. We use here the same arguments as in [34].

- The "stationary phase admissible" term T_h^s

Proposition 2.10. *There is a constant $1 < C_0 < \infty$ such that the kernel K_h^s of T_h^s satisfies*

$$|K_h^s(t, x, s, y)| \leq C_N h^N \quad \forall N, \quad \text{if } \frac{|t - s|}{|x - y|} \notin [C_0^{-1}, C_0]. \quad (2.35)$$

Moreover, there is a function $\xi_c(t, x, s, y)$ which is smooth in the variables (t, s) , uniformly over (x, y) , so that if $C_0^{-1} \leq \frac{|t - s|}{|x - y|} \leq C_0$, then

$$|K_h^s(t, x, s, y)| \lesssim h^{-n} \left(1 + \frac{|t - s|}{h}\right)^{-n/2}, \quad \text{for } |t - s| \geq h^{1/3}. \quad (2.36)$$

Proof. We shall use stationary phase lemma to evaluate the kernel K_h^s of T_h^s . The critical points occur when $|t - s| \simeq |x - y|$. For some constant C_0 and for $|\xi| \in \text{supp}\psi$, ξ_1 in a small neighborhood of 1, we have

$$|\nabla_\xi(\phi(t, x, \xi) - \phi(s, y, \xi))| \simeq |t - s| + |x - y| \geq h^{1/3}, \quad \text{if } \frac{|t - s|}{|x - y|} \notin [C_0^{-1}, C_0].$$

Since $c \in S_{2/3, 1/3}^0$, an integration by parts leads to (2.35). If $|t - s| \simeq |x - y|$ we introduce a cutoff function $\kappa\left(\frac{|x - y|}{|t - s|}\right)$, $\kappa \in C_0^\infty(\mathbb{R} \setminus \{0\})$. The phase function can be written as

$$\phi(t, x, \xi) - \phi(s, y, \xi) = (t - s)\Theta(t, x, s, y, \xi) \quad \text{for } |t - s| \simeq |x - y| \geq h^{1/3}.$$

We want to apply the stationary phase method with parameter $|t - s|/h \geq h^{-2/3} \gg 1$ to estimate K_h^s . For x, y, t, s fixed we must show that the critical points of Θ are non-degenerate.

Lemma 2.11. *If T is sufficiently small then the phase function $\Theta(t, x, s, y, \xi)$ admits a unique, non-degenerate critical point ξ_c . Moreover, for $0 \leq t, s \leq T$, the function $\xi_c(t, x, s, y)$ solving $\nabla_\xi \Theta(t, x, s, y, \xi_c) = 0$ is smooth in t and s , with uniform bounds on derivatives as x and y vary and we have*

$$|\partial_{t,s}^\alpha \partial_{x,y}^\gamma \xi_c(t, x, s, y)| \leq C_{\alpha,\gamma} h^{-|\alpha|/3} \quad \text{if } |x - y| \geq h^{1/3}. \quad (2.37)$$

Proof. The phase $\Theta(t, x, s, y, \xi)$ writes

$$\begin{aligned} \Theta(t, x, s, y, \xi) &= \xi_1^2 + \frac{1}{(t-s)} (\phi(0, x, \xi) - \phi(0, y, \xi)) = \\ &= \xi_1^2 + \frac{1}{(t-s)} \sum_{j=1}^n (x_j - y_j) \partial_{x_j} \phi(0, z_{x,y}, \xi), \end{aligned}$$

for some $z_{x,y}$ close to x, y (if T is sufficiently small then $|t - s| \simeq |x - y|$ is small), and using the eikonal equations (2.10) we can write

$$\Theta(t, x, s, y, \xi) = \langle \nabla_x \phi, \nabla_x \phi \rangle_g(0, z_{x,y}, \xi) - \frac{1}{(t-s)} \sum_{j=1}^n (x_j - y_j) \partial_{x_j} \phi(0, z_{x,y}, \xi).$$

Let us write $\langle \nabla_x \phi, \nabla_x \phi \rangle_g = \sum_{j,k} g^{j,k} \partial_{x_j} \phi \partial_{x_k} \phi$ and compute explicitly $\nabla_\xi \Theta$. For each $l \in \{1, \dots, n\}$ we have

$$\partial_{\xi_l} \Theta(t, x, s, y, \xi) = \sum_{j=1}^n \partial_{\xi_l, x_j}^2 \phi(0, z_{x,y}, \xi) \left(2 \sum_{k=1}^n g^{j,k}(z_{x,y}) \partial_{x_k} \phi(0, z_{x,y}, \xi) - \frac{(x_j - y_j)}{(t-s)} \right), \quad (2.38)$$

thus

$$\nabla_\xi \Theta(t, x, s, y, \xi) = \nabla_{\xi,x}^2 \phi(0, z_{x,y}, \xi) \cdot \begin{pmatrix} 2 \sum_k g^{1,k}(z_{x,y}) \partial_{x_k} \phi(0, z_{x,y}, \xi) - \frac{x_1 - y_1}{(t-s)} \\ \vdots \\ 2 \sum_k g^{n,k}(z_{x,y}) \partial_{x_k} \phi(0, z_{x,y}, \xi) - \frac{x_n - y_n}{(t-s)} \end{pmatrix}, \quad (2.39)$$

where $\nabla_{\xi,x}^2 \phi = (\partial_{\xi_l, x_j}^2 \phi)_{l,j \in \{1, \dots, n\}}$ is the matrix $n \times n$ whose elements are the second derivatives of ϕ with respect to ξ and x . We need the following:

Lemma 2.12. *If T is sufficiently small, $|x - y| \simeq |t - s| \leq 2T$ then*

$$\det(\nabla_{\xi, x}^2 \phi)(0, z_{x, y}, \xi) \neq 0. \quad (2.40)$$

In what follows we complete the proof of Lemma 2.11 and then we prove (2.40).

A critical point for Θ satisfies $\nabla_{\xi} \Theta(t, x, s, y, \xi) = 0$ and from (2.39) and (2.40) this translates into

$$\left((g^{j, k}(z_{x, y}))_{j, k} \right) (\nabla_x \phi)^t(0, z_{x, y}, \xi) = \frac{(x - y)}{(t - s)}. \quad (2.41)$$

Since $(g^{j, k})_{j, k}$ is invertible we determine $\nabla_x \phi(0, z_{x, y}, \xi)$, and by (2.40) we can apply the implicit function's theorem to obtain (for T small enough) a critical point $\xi_c = \xi_c(t, x, s, y)$ for Θ . In order to show that ξ_c is non-degenerate we compute

$$\begin{aligned} \partial_{\xi_q} \partial_{\xi_l} \Theta(t, x, s, y, \xi) &= \sum_{j=1}^n \partial_{\xi_q, \xi_l, x_j}^3 \phi(0, z_{x, y}, \xi) \left(2 \sum_{k=1}^n g^{j, k}(z_{x, y}) \partial_{x_k} \phi(0, z_{x, y}, \xi) - \frac{(x_j - y_j)}{(t - s)} \right) \\ &\quad + 2 \sum_{j=1}^n \partial_{\xi_l, x_j}^2 \phi(0, z_{x, y}, \xi) \left(\sum_{k=1}^n g^{j, k}(z_{x, y}) \partial_{\xi_q, x_k}^2 \phi(0, z_{x, y}, \xi) \right), \end{aligned} \quad (2.42)$$

consequently at the critical point $\xi = \xi_c$ the hessian matrix $\nabla_{\xi, \xi}^2 \Theta$ is given by

$$\nabla_{\xi, \xi}^2 \Theta(t, x, s, y, \xi_c) = 2(\nabla_{\xi, x}^2 \phi)(g^{ij}(z_{x, y}))_{i, j} (\nabla_{\xi, x}^2 \phi)|_{(0, z_{x, y}, \xi_c)},$$

consequently for T small enough the critical point ξ_c is non-degenerate by (2.40).

It remains to prove (2.40). To deduce this we need to use the precise form of the initial phase function in (2.27). Writing again χA_+ and $(\chi A_+)'$ in terms of their Fourier transform, we express the phase function of (2.27) in the form

$$-t\xi_1^2 + \theta(x, \xi) + \sigma\xi_1^{-2/3}\zeta(x, \xi) + \sigma^3/3\xi_1^2. \quad (2.43)$$

Notice that at $t = 0$ this phase is homogeneous of degree 1 in ξ and the proof of (2.40) reduces to checking that the Jacobian J of the mapping

$$(\xi, \sigma) \rightarrow (\nabla_x(\theta(x, \xi) + \sigma\zeta(x, \xi)), \zeta(x, \xi) + \sigma^2)$$

does not vanish at the critical point of the phase (2.43). At this (critical) point $\sigma = \zeta(x, \xi) = 0$ and $\nabla_{x'} \zeta(x, \xi) = 0$. Since $\partial_{x_n} \zeta(x, \xi) \neq 0$ and $\partial_{\xi_n} \zeta(x, \xi) \neq 0$ there, the result follows by the nonvanishing of $|\nabla_{x'} \nabla_{\xi'} \theta(x, \xi)|$. In fact we have

$$\det \begin{pmatrix} \nabla_{x'} \nabla_{\xi'} \theta & \nabla_{\xi'} \partial_{x_n} \theta & \nabla_{\xi'} \zeta \\ \partial_{\xi_n} \nabla_{x'} \theta & \partial_{\xi_n} \partial_{x_n} \theta & \partial_{\xi_n} \zeta \\ \nabla_{x'} \zeta & \partial_{x_n} \zeta & 2\sigma \end{pmatrix} \Big|_{\sigma^2 = -\zeta} \neq 0.$$

This geometric property can be restated in an equivalent form by saying that at $t = 0$, ϕ satisfies (2.40) if $|x - y| \leq 2T$ and T sufficiently small. \square

On the support of κ it follows that the kernel K_h^s writes

$$K_h^s(t, x, s, y) = \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}|t-s|\Theta(t,x,s,y,\xi)} \psi^2(|\xi|) (1 - \rho(h^{-1/3}|t-s|)) c_m(x, \xi/h) \overline{c_m(y, \xi/h)} d\xi, \quad (2.44)$$

where, if $\omega = |t - s|/h$ and $\xi_1 \simeq 1$, the symbol satisfies

$$|\partial_{t,s}^\alpha \partial_\omega^k \sigma_h(t, x, s, y, \omega\xi/|t-s|)| \leq C_{\alpha,k} h^{-|\alpha|/3} (|t-s|^{3/2}/h)^{-2k/3},$$

where we set

$$\sigma_h(t, x, s, y, \omega\xi/|t-s|) = (1 - \rho(h^{-1/3}|t-s|)) c_m(x, \omega\xi/|t-s|) \overline{c_m(y, \omega\xi/|t-s|)}.$$

Indeed, since $c_m \in S_{2/3,1/3}^0$, for $\alpha = 0$ one has

$$|\partial_\omega^k \sigma_h| \leq |\xi| |t-s|^{-k} |(\partial_\xi^k c)(t, x, \omega\xi/|t-s|)| \leq C_{0,k} |t-s|^{-k} (\omega/|t-s|)^{-2k/3} = C_{0,k} |t-s|^{-k} h^{2k/3}.$$

We conclude using the next lemma with $\omega = \frac{|t-s|}{h}$ and $\delta = |t-s|^{3/2} \geq h^{1/2} \gg h$.

Lemma 2.13. *Suppose that $\Theta(z, \xi) \in C^\infty(\mathbb{R}^{2(n+1)} \times \mathbb{R}^n)$ is real, $\nabla_\xi \Theta(z, \xi_c(z)) = 0$, $\nabla_\xi \Theta(z, \xi) \neq 0$ if $\xi \neq \xi_c(z)$, and*

$$|\det \nabla_{\xi\xi}^2 \Theta| \geq c_0 > 0, \text{ if } |\xi| \leq 1.$$

Suppose also that

$$|\partial_z^\alpha \partial_\xi^\beta \Theta(z, \xi)| \leq C_{\alpha,\beta} h^{-|\alpha|/3}, \quad \forall \alpha, \beta.$$

In addition, suppose that the symbol $\sigma_h(z, \xi, \omega)$ vanishes when $|\xi| \geq 1$ and satisfies

$$|\partial_z^\alpha \partial_\xi^\gamma \partial_\omega^k \sigma_h(z, \xi, \omega)| \leq C_{k,\alpha,\gamma} h^{-(|\alpha|+|\gamma|)/3} (\delta/h)^{-2k/3}, \quad \forall k, \alpha, \gamma,$$

where on the support of σ_h we have $\omega \geq h^{-2/3}$ and $\delta > 0$. Then we can write

$$\int_{\mathbb{R}^n} e^{i\omega\Theta(z,\xi)} \sigma_h(z, \xi, \omega) d\xi = \omega^{-n/2} e^{i\omega\Theta(z,\xi_c(z))} b_h(z, \omega),$$

where b_h satisfies

$$|\partial_\omega^k \partial_z^\alpha b_h(z, \omega)| \leq C_{k,\alpha} h^{-|\alpha|/3} (\delta/h)^{-2k/3}$$

and where each of the constants depend only on c_0 and the size of finitely many of the constants $C_{\alpha,\beta}$ and $C_{k,\alpha,\gamma}$ above. In particular, the constants are uniform in δ if $1 \geq \delta \geq h$.

This Lemma is used in [34, Lemma 2.6] and also in the thesis of Grieser [20] and it follows easily from the proof of standard stationary phase lemma (see [36, pag. 45]). Proposition 2.10 is thus proved. \square

For each t, s , let $T_h^s(t, s)$ be the "frozen" operator defined by

$$T_h^s(t, s)g(x) = \int K_h^s(t, x, s, y)g(y)dy.$$

From Proposition 2.10 we deduce

$$\|T_h^s(t, s)g\|_{L^\infty(\mathbb{R}^n)} \leq C \max(h^{-n}, (h|t-s|)^{-n/2})\|g\|_{L^1(\mathbb{R}^n)}. \quad (2.45)$$

We need the following

Lemma 2.14. *For t, s fixed the frozen operator $T_h^s(t, s)$ is bounded on $L^2(\mathbb{R}^n)$,*

$$\|T_h^s(t, s)g\|_{L^2(\mathbb{R}^n)} \leq C\|g\|_{L^2(\mathbb{R}^n)}. \quad (2.46)$$

Proof. Explicitly we have

$$T_h^s(t, s)g(x) = (1 - \rho(h^{-1/3}|t-s|)) \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}(\phi(t,x,\xi) - \phi(s,y,\xi))} c_m(x, \xi/h) \overline{c_m(y, \xi/h)} \psi^2(|\xi|) g(y) d\xi dy$$

and (since $0 \leq \rho(h^{-1/3}|t-s|) \leq 1$) it is enough to prove that the frozen operator $W_h(\cdot, t)$ is bounded on $L^2(\mathbb{R}^n)$. This follows from $\|A_h f(\cdot, t)\|_{L^2(\mathbb{R}^n)} \simeq \|f\|_{L^2(\mathbb{R}^n)}$ together with the fact that the $L^2(\mathbb{R}^n)$ norms of $A_h f$, $M_h f$ and thus $W_h f$ are equivalent. \square

Interpolation between (2.45) and (2.46) with weights $1-2/r$ and $2/r$ respectively yields

$$\|T_h^s(t, s)g\|_{L^r(\mathbb{R}^n)} \leq Ch^{-n(1-2/r)} \left(1 + \frac{|t-s|}{h}\right)^{-n(1/2-1/r)} \|g\|_{L^{r'}(\mathbb{R}^n)} \quad (2.47)$$

and hence

$$\|T_h^s F\|_{L^q(0,T), L^r(\mathbb{R}^n)} \leq Ch^{-n/2(1-2/r)} \left\| \int_{1 \ll \frac{|t-s|}{h}}^T |t-s|^{-n/2(1-2/r)} \|F(\cdot, s)\|_{L^{r'}(\mathbb{R}^n)} ds \right\|_{L^{q'}((0,T))}.$$

Since $n(\frac{1}{2} - \frac{1}{r}) = \frac{2}{q} < 1$ the application $|t|^{-2/q} : L^{q'} \rightarrow L^q$ is bounded and by Hardy-Littlewood-Sobolev inequality we deduce

$$\|T_h^s F\|_{L^q((0,T), L^r(\mathbb{R}^n))} \leq Ch^{-n(1/2-1/r)} \|F\|_{L^{q'}((0,T), L^{r'}(\mathbb{R}^n))}. \quad (2.48)$$

- The "frozen" term T_h^f

To estimate T_h^f it suffices to obtain bounds for its kernel K_h^f with both the variables (t, x) and (s, y) restricted to lie in a cube of \mathbb{R}^{n+1} of sidelength comparable to $h^{1/3}$. Let us decompose S_T into disjoint cubes $Q = Q_x \times Q_t$ of sidelength $h^{1/3}$. We then have

$$\|T_h^f F\|_{L^q([0,T], L^r(\mathbb{R}^n))}^q = \int_0^T \left(\sum_{Q=Q_x \times Q_t} \|\chi_Q T_h^f F\|_{L^r(Q_x)}^r \right)^{q/r} dt = \sum_Q \|\chi_Q T_h^f F\|_{L^q([0,T], L^r(\mathbb{R}^n))}^q,$$

where by χ_Q we denoted the characteristic function of the cube Q . In fact, by the definition, the integral kernel $K_h^f(t, x, s, y)$ of T_h^f vanishes if $|t-s| \geq h^{1/3}$. If $|t-s| \leq h^{1/3}$ and $|x-y| \geq C_0 h^{1/3}$, then the phase

$$\phi(t, x, \xi) - \phi(s, y, \xi)$$

has no critical points with respect to ξ_1 (on the support of ψ), so that

$$|K_h^f(t, x, s, y)| \leq C_N h^N \quad \forall N, \text{ if } |x-y| \geq C_0 h^{1/3}.$$

It therefore suffices to estimate $\|\chi_Q T_h^f \chi_{Q^*} F\|_{L^q([0,T], L^r(\mathbb{R}^n))}$, where Q^* is the dilate of Q by some fixed factor independent of h . Since $q > 2 > q'$, $r \geq 2 \geq r'$, where q', r' are such that $1/q + 1/q' = 1$, $1/r + 1/r' = 1$, then we shall obtain

$$\sum_Q \|\chi_Q T_h^f \chi_{Q^*} F\|_{L^q([0,T], L^r(\mathbb{R}^n))}^q \leq C_1 \sum_Q \|\chi_{Q^*} F\|_{L^{q'}([0,T], L^{r'}(\mathbb{R}^n))}^q \leq C_2 \|F\|_{L^{q'}([0,T], L^{r'}(\mathbb{R}^n))}^q. \quad (2.49)$$

In order to prove (2.49) we shall use the following:

Proposition 2.15. *Let $b(\xi) \in L^\infty(\mathbb{R}^n)$ be elliptic near $\xi_1 \simeq 1$, $b_h(\xi) := b(\xi/h)$, then for $h \ll |t-s| \leq h^{1/3}$, $h \ll |x-y| \leq h^{1/3}$ the operator defined by*

$$B_h f(x, t) = \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}\phi(t, x, \xi)} \psi(|\xi|) b_h(\xi) \hat{f}\left(\frac{\xi}{h}\right) d\xi \quad (2.50)$$

satisfies

$$\|B_h f\|_{L^q((0,T], L^r(\mathbb{R}^n))} \leq C h^{-n/2(1/2-1/r)} \|f\|_{L^2(\mathbb{R}^n)}. \quad (2.51)$$

Proof. We use again the TT* argument. Since $b(\xi)$ acts as an L^2 multiplier we can apply the stationary phase theorem in the integral

$$\int e^{\frac{i}{h}(\phi(t, x, \xi) - \phi(s, y, \xi))} \psi(|\xi|) d\xi$$

in order to obtain

$$\|B_h B_h^* F\|_{L^q((0,T], L^r(\mathbb{R}^n))} \lesssim h^{-n(1/2-1/r)} \|F\|_{L^{q'}((0,T], L^{r'}(\mathbb{R}^n))}.$$

Notice that we haven't used the special properties of the phase function at $t = 0$. \square

Let now Q be a fixed cube in \mathbb{R}^{n+1} of sidelength $h^{1/3}$. Let

$$b_h(t, x, s, y, \xi) = \rho(h^{-1/3}|t - s|)c_m(x, \xi/h)\overline{c_m(y, \xi/h)},$$

and write

$$\begin{aligned} b_h(t, x, s, y, \xi) &= b_h(0, 0, s, y, \xi) + \int_0^t \partial_t b_h(r, 0, s, y, \xi) dr + \dots \\ &+ \dots + \int_0^t \dots \int_0^{x_n} \partial_t \dots \partial_{x_n} b_h(r, z_1, \dots, z_n, s, y, \xi) dr dz. \end{aligned}$$

If the symbol c is independent of t, x then the estimates (2.29) follow from Proposition 2.15. We use this, for instance, to deduce

$$\begin{aligned} &\|\chi_Q T_h^f \chi_{Q^*} F\|_{L^q((0,T], L^r(\mathbb{R}^n))} \leq \\ &Ch^{-n/2(1/2-1/r)} (\| \int \int e^{\frac{i}{h}(x\xi - \phi(s,y,\xi))} \psi(|\xi|) b_h(0, 0, s, y, \xi) F(y, s) d\xi ds dy \|_{L^2(\mathbb{R})} + \dots \\ &\dots + \| \int_0^{h^{1/3}} \dots \int_0^{h^{1/3}} \int \int e^{\frac{i}{h}(x\xi - \phi(s,y,\xi))} \partial_t \dots \partial_{x_n} \psi(|\xi|) b_h(r, z, s, y, \xi) F(y, s) d\xi ds dy \|_{L^2(\mathbb{R}^n)} dr dz. \end{aligned}$$

Each derivative of $b_h(t, x, s, y, \xi)$ loses a factor of $h^{-1/3}$, but this is compensated by the integral over (r, z) , so that it suffices to establish uniform estimates for fixed (r, z) . By duality, we have to establish the estimate

$$\| \int \int e^{\frac{i}{h}\phi(s,y,\xi)} \psi(|\xi|) b_h(0, 0, s, y, \xi) \hat{f}\left(\frac{\xi}{h}\right) d\xi \|_{L^q((0,T], L^r(\mathbb{R}^n))} \leq C \|f\|_{L^2(\mathbb{R}^n)},$$

which follows by using the same argument of freezing the variables (s, y) together with Proposition 2.15.

2.2.4 The diffractive term D_h

In order to estimate the diffractive term we shall proceed again like in [34, Sect.2].

Lemma 2.16. *For $x_n \geq 0$ and for ξ in a small conic neighborhood of the positive ξ_1 axis, the symbol q of S_h can be written in the form*

$$\begin{aligned} q(x, \xi) &:= (a(x, \xi)((1 - \chi)A_+)(\zeta(x, \xi)) + b(x, \xi)((1 - \chi)A_+)'(\zeta(x, \xi))) \frac{Ai(\zeta_0(\xi))}{A_+(\zeta_0(\xi))} \\ &= p(x, \xi, \zeta(x, \xi)), \end{aligned}$$

where, for some $c > 0$

$$|\partial_\xi^\alpha \partial_\zeta^j \partial_{x'}^\beta \partial_{x_n}^k p(x, \xi, \zeta(x, \xi))| \leq C_{\alpha,j,\beta,k} \xi_1^{1/6 - |\alpha| + 2k/3} e^{-cx_n^{3/2} \xi_1 - |\zeta|^{3/2}/2}.$$

Proof. Since

$$|\partial_\zeta^k((1-\chi)A_+)(\zeta)| \leq C_{k,\epsilon} e^{(2/3+\epsilon)|\zeta|^{3/2}}, \quad \forall \epsilon > 0,$$

and the symbols a and b belong to $S_{1,0}^{1/6}$, the above fact will follow by showing that in the region $\zeta(x, \xi) \geq -2$,

$$\frac{Ai}{A_+}(\zeta_0(\xi)) = \tilde{p}(x, \xi', \zeta(x, \xi)),$$

where if $\xi' = (\xi_1, \dots, \xi_{n-1})$

$$|\partial_{\xi'}^\alpha \partial_\zeta^j \partial_{x'}^\beta \partial_{x_n}^k \tilde{p}(x, \xi', \zeta)| \leq C_{\alpha,j,\beta,k,\epsilon} \xi_1^{-|\alpha|+2k/3} e^{-cx_n^{3/2} \xi_1 - (4/3-\epsilon)|\zeta|^{3/2}}. \quad (2.52)$$

At $x_n = 0$, one has $\zeta = \zeta_0$, $\partial_{x_n} \zeta < 0$. It follows that for some $c > 0$

$$\zeta_0(x, \xi) \geq \zeta(x, \xi) + cx_n \xi_1^{2/3}.$$

By the asymptotic behavior of the Airy function we have, in the region $\zeta(x, \xi) \geq -2$

$$\left| \left(\frac{Ai}{A_+} \right)^{(k)}(\zeta_0) \right| \leq C_{k,\epsilon} e^{-cx_n^{3/2} \xi_1 - (4/3-\epsilon)|\zeta(x,\xi)|^{3/2}}. \quad (2.53)$$

We introduce a new variable $\tau(x, \xi) = \xi_1^{1/3} \zeta(x, \xi)$. At $x_n = 0$ one has $\tau = -\xi_n$, so that we can write $\xi_n = \sigma(x, \xi', \tau)$, where σ is homogeneous of degree 1 in (ξ', τ) . We set

$$\tilde{p}(x, \xi', \zeta) = \frac{Ai}{A_+}(-\xi_1^{-1/3} \sigma(x, \xi', \xi_1^{1/3} \zeta)).$$

The estimates (2.52) will follow by showing that

$$|\partial_{\xi'}^\alpha \partial_\tau^j \partial_{x'}^\beta \partial_{x_n}^k \frac{Ai}{A_+}(-\xi_1^{-1/3} \sigma(x, \xi', \tau))| \leq C_{\alpha,j,\beta,k,\epsilon} \xi_1^{-|\alpha|-j+2k/3} e^{-cx_n^{3/2} \xi_1 - (4/3-\epsilon)|\tau|^{3/2} \xi_1^{-1/2}}. \quad (2.54)$$

For $k = 0$, the estimates (2.54) follow from (2.53), together with the fact that

$$|\partial_{\xi'}^\alpha \partial_\tau^j \partial_{x'}^\beta \frac{Ai}{A_+}(-\xi_1^{-1/3} \sigma(x, \xi', \tau))| \leq C_{\alpha,\beta,j} (x_n \xi_1^{2/3} + \xi_1^{-1/3} |\tau|) \xi_1^{-|\alpha|-j},$$

which, in turn, holds by homogeneity, together with the fact that $\sigma(x, \xi', \tau) = 0$ if $x_n = \tau = 0$. If $k > 0$, the estimate (2.54) follows by observing that the effect of differentiating in x_n is similar to multiplying by a symbol of order $2/3$. This concludes the proof of Lemma 2.16. \square

We now write the Schwartz kernel of the diffractive term D_h in the form

$$\int e^{i(\theta(x,\xi) - ht\xi_1^2)} \psi(h|\xi|) q(x, \xi) d\xi$$

$$= \int e^{i(\theta(x,\xi) - ht\xi_1^2 + s\xi_1^{-2/3}\zeta(x,\xi) + s^3/3\xi_1^2 - \langle y, \xi \rangle)} \psi(h|\xi|) a(x, \xi, s\xi_1^{-2/3}) ds d\xi,$$

where

$$|\partial_\xi^\alpha \partial_s^j \partial_{x'}^\beta \partial_{x_n}^k a(x, \xi, s\xi_1^{-2/3})| \leq C_{\alpha,j,\beta,k,N} \xi_1^{-1/2 - |\alpha| - 2j/3 + 2k/3} e^{-cx_n^{3/2}\xi_1} \langle \xi^{-2/3} s \rangle^{-N}, \quad \forall N.$$

From now on we proceed as for the main term and we reduce the problem to considering the operator

$$W_h^d f(x, t) = \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}\phi(t,x,\xi)} c_d(x, \xi/h) \psi(|\xi|) \hat{f}\left(\frac{\xi}{h}\right) d\xi,$$

where now $x_n^j \partial_{x_n}^k c_d(x, \xi) \in S_{2/3, 1/3}^{2(j-k)/3}(\mathbb{R}_{x',t}^n \times \mathbb{R}_\xi^n)$ uniformly over x_n . Using the freezing arguments behind the proof of the estimates for T_h^f and Minkovski inequality we have

$$\begin{aligned} \|W_h^d f\|_{L^q((0,T], L^r(\mathbb{R}^n))} &\leq \left\| \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}\phi(t,x,\xi)} c_d(x', 0, \xi/h) \psi(|\xi|) \hat{f}\left(\frac{\xi}{h}\right) d\xi \right\|_{L^q((0,T], L^r(\mathbb{R}^n))} + \\ &h^{-2/3} \int_0^{h^{2/3}} \left\| \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}\phi(t,x,\xi)} (h^{2/3} \partial_{x_n} c_d(x', r, \xi/h)) \psi(|\xi|) \hat{f}\left(\frac{\xi}{h}\right) d\xi \right\|_{L^q((0,T], L^r(\mathbb{R}^n))} dr + \\ &h^{2/3} \int_{r>h^{2/3}} \frac{dr}{r^2} \left\| \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}\phi(t,x,\xi)} (h^{-2/3} r^2 \partial_{x_n} c_d(x', r, \xi/h)) \psi(|\xi|) \hat{f}\left(\frac{\xi}{h}\right) d\xi \right\|_{L^q((0,T], L^r(\mathbb{R}^{n-1} \times \{x_n > r\}))}. \end{aligned}$$

Since $c_d(x', 0, \xi)$ and $h^{2/3}(1 + h^{-4/3}r^2)\partial_{x_n} c_d(x', r, \xi)$ are symbols of order 0 and type $(2/3, 1/3)$ with uniform estimates over r , the estimates for the diffractive term also follow from Proposition 2.8. Indeed, the term in the second line loses a factor $h^{-2/3}$, but this is compensated by the integral over $r \leq h^{2/3}$. The term in the third line can be bounded by above by

$$\begin{aligned} &\left\| \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}\phi(t,x,\xi)} (h^{-2/3} r^2 \partial_{x_n} c_d(x', r, \xi/h)) \psi(|\xi|) \hat{f}\left(\frac{\xi}{h}\right) d\xi \right\|_{L^q((0,T], L^r(\mathbb{R}^n))} \times h^{2/3} \int_{r>h^{2/3}} \frac{dr}{r^2} \\ &\leq \left\| \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}\phi(t,x,\xi)} (h^{-2/3} r^2 \partial_{x_n} c_d(x', r, \xi/h)) \psi(|\xi|) \hat{f}\left(\frac{\xi}{h}\right) d\xi \right\|_{L^q((0,T], L^r(\mathbb{R}^n))}. \end{aligned}$$

We conclude using the same arguments as in the proof of Proposition 2.8, where now W_h is replaced by operators with symbols $c_d(x', 0, \xi)$, $h^{2/3}\partial_{x_n} c_d(x', r, \xi)$ and $h^{-2/3}r^2\partial_{x_n} c_d(x', r, \xi)$ respectively.

3 Strichartz estimates for the classical Schrödinger equation outside a strictly convex obstacle in \mathbb{R}^n

In this section we prove Theorem 1.7 under the Assumptions 1.6. In what follows we shall work with the Laplace operator with constant coefficients $\Delta_D = \sum_{j=1}^n \partial_j^2$ acting on $L^2(\Omega)$

to avoid technicalities, where Ω is the exterior in \mathbb{R}^n of a strictly convex domain. In the proof of Theorem 1.7 we distinguish two main steps: we start by performing a time rescaling which transforms the equation (1.9) into a semi-classical problem: due to the finite speed of propagation (proved by Lebeau [27]), we can use the (local) semi-classical result of Theorem 1.3 together with the smoothing effect (by Staffilani and Tataru [37] and Burq [5]) to obtain classical Strichartz estimates near the boundary. Outside a fixed neighborhood of $\partial\Omega$ we use a method suggested by Staffilani and Tataru [37] which consists in considering the Schrödinger flow as a solution of a problem in the whole space \mathbb{R}^n , for which the Strichartz estimates are known.

Following [5], we split $u(x, t) = e^{it\Delta_D}u_0(x)$ as a sum of two terms $u = \chi u + (1 - \chi)u$, where $\chi \in C_0^\infty(\mathbb{R}^n)$ equals 1 in a neighborhood of $\partial\Omega$.

- Study of $v = \chi u$. We have

$$i\partial_t v - \Delta_D v = [\Delta_D, \chi]u, \quad v|_{t=0} = \chi u_0. \quad (3.1)$$

Let $\Psi \in C_0^\infty(\mathbb{R} \setminus 0)$ and let $\varphi \in C_0^\infty((-1, 2))$ equal to 1 on $[0, 1]$. Let $v_h = \Psi(-h^2\Delta_D)v$ and for $l \in \mathbb{Z}$, $v_{h,l} = \varphi(t/h - l)v_h$, solution to

$$\begin{cases} i\partial_t v_{h,l} - \Delta_D v_{h,l} = \varphi(t/h - l)\Psi(-h^2\Delta_D)[\Delta_D, \chi]u + i\frac{\varphi'(t/h-l)}{h}\Psi(-h^2\Delta_D)\chi u, \\ v_{h,l}|_{t < hl-h} = 0, \quad v_{h,l}|_{t > hl+2h} = 0. \end{cases} \quad (3.2)$$

If we denote $\Psi(-h^2\Delta_D)V_{h,l}$ the right-hand side of (3.2), the Duhamel formula writes

$$v_{h,l}(x, t) = \int_{hl-l}^t e^{i(t-s)\Delta_D}\Psi(-h^2\Delta_D)V_{h,l}(x, s)ds.$$

Using Minkovski inequality we obtain

$$\|v_{h,l}\|_{L^q([hl-h, hl+2h], L^r(\Omega))} \leq C \int_{[hl-h, hl+2h]} \|e^{i(t-s)\Delta_D}\Psi(-h^2\Delta_D)V_{h,l}(s)\|_{L^q([hl-h, hl+2h], L^r(\Omega))} ds. \quad (3.3)$$

In Appendix we deduce from Theorem 1.3 the following

Proposition 3.1. *We have*

$$\begin{aligned} & \|e^{i(t-s)\Delta_D}\Psi(-h^2\Delta_D)V_{h,l}(s)\|_{L^q([hl-h, hl+2h], L^r(\Omega))} \leq \\ & C\|\Psi(-h^2\Delta_D)V_{h,l}(s)\|_{L^2(\Omega)} + O(h^\infty)\|\Psi(-h^2\Delta_D)V_{h,l}(s)\|_{L^2(\Omega)}. \end{aligned} \quad (3.4)$$

We finish the proof of Theorem 1.7 in this first case. It follows from (3.3) and (3.4) that we have, modulo $O(h^\infty)$

$$\begin{aligned} \|v_{h,l}\|_{L^q([hl-h,hl+2h],L^r(\Omega))} &\leq C \int_{[hl-h,hl+2h]} \|\Psi(-h^2\Delta_D)V_{h,l}(s)\|_{L^2(\Omega)} ds \quad (3.5) \\ &\leq Ch^{1/2}\|\Psi(-h^2\Delta_D)V_{h,l}\|_{L^2([hl-h,hl+2h],L^2(\Omega))}, \end{aligned}$$

and since $q \geq 2$ it follows that, modulo $O(h^\infty)$

$$\begin{aligned} \|v_h\|_{L^q(\mathbb{R},L^r(\Omega))}^q &\leq Ch^{q/2} \sum_{l=-\infty}^{\infty} \|\Psi(-h^2\Delta_D)V_{h,l}(s)\|_{L^2([hl-h,hl+2h],L^2(\Omega))}^q \quad (3.6) \\ &\leq Ch^{q/2} \left(\sum_{l=-\infty}^{\infty} \|\Psi(-h^2\Delta_D)V_{h,l}\|_{L^2([hl-h,hl+2h],L^2(\Omega))}^2 \right)^{q/2}. \end{aligned}$$

Thanks to the spectral cutoff Ψ , we obtain

$$\begin{aligned} \|\Psi(-h^2\Delta_D)V_{h,l}\|_{L^2([hl-h,hl+2h],L^2(\Omega))} &= \quad (3.7) \\ \|\varphi(t/h-l)\Psi(-h^2\Delta_D)[\Delta_D,\chi]u + i\frac{\varphi'(t/h-l)}{h}\Psi(-h^2\Delta_D)\chi u\|_{L^2([hl-h,hl+2h],L^2(\Omega))} &\leq \\ Ch^{-1/2}\|\varphi(t/h-l)\Psi(-h^2\Delta_D)[\Delta_D,\chi]u\|_{L^2([hl-h,hl+2h],H^{-1/2}(\Omega))} + \\ Ch^{1/2}\|\frac{\varphi'(t/h-l)}{h}\Psi(-h^2\Delta_D)\chi u\|_{L^2([hl-h,hl+2h],H^{1/2}(\Omega))}. \end{aligned}$$

If $\tilde{\varphi} \in C_0^\infty(\mathbb{R})$ and $\tilde{\chi} \in C_0^\infty(\Omega)$ are chosen such that they are equal to 1 on the supports of φ, χ , we get, modulo $O(h^\infty)$

$$\|\Psi(-h^2\Delta_D)V_{h,l}\|_{L^2([hl-h,hl+2h],L^2(\Omega))} \leq Ch^{-1/2}\|\tilde{\varphi}(t/h-l)\tilde{\psi}(-h^2\Delta_D)\tilde{\chi}u\|_{L^2([hl-h,hl+2h],H^{1/2}(\Omega))}, \quad (3.8)$$

from which we deduce, using also (3.6)

$$\|v_h\|_{L^q(\mathbb{R},L^r(\Omega))} \leq C\|\psi(-h^2\Delta_D)\tilde{\chi}u\|_{L^2(\mathbb{R},H^{1/2}(\Omega))}. \quad (3.9)$$

We also need the next result

Proposition 3.2. (*[7, Prop.2.7]*) *Assume that $\Omega = \mathbb{R}^n \setminus \mathcal{O}$, where $\mathcal{O} \neq \emptyset$ is a compact non-trapping obstacle. Then for $T > 0$, for every $\tilde{\chi} \in C_0^\infty(\mathbb{R}^n)$, $n \geq 2$, one has*

$$\|\tilde{\chi}u\|_{L^2([-T,T],H^{s+1/2}(\Omega))} \leq C\|u_0\|_{H^s(\Omega)},$$

where $s \in [0, 1]$ and $u(x, t) = e^{it\Delta_D}u_0(x)$. Moreover, the constant C do not depend on T , i.e. the estimates are global in time.

In order to finish the proof in this case it suffices to use the next lemma proved in [23]:

Lemma 3.3. (see [23, Thm.1.1]) Let $\Psi_0 \in C_0^\infty(\mathbb{R})$, $\Psi \in C_0^\infty((1/2, 2))$ satisfy

$$\Psi_0(\lambda) + \sum_{j \geq 1} \Psi(2^{-2j}\lambda) = 1, \quad \forall \lambda \in \mathbb{R}.$$

Then for all $r \in [2, \infty)$ we have

$$\|f\|_{L^r(\Omega)} \leq C_r \left(\|\Psi_0(-\Delta_D)u_0\|_{L^r(\Omega)} + \left(\sum_{j=1}^{\infty} \|\Psi(-2^{-2j}\Delta_D)f\|_{L^r(\Omega)}^2 \right)^{1/2} \right). \quad (3.10)$$

Applying Lemma 3.3 to $f = \chi u(\cdot, t)$ and taking the L^q norm in time yields

$$\|\chi u\|_{L^q(\mathbb{R}, L^r(\Omega))} \leq \left\| \left(\|\Psi_0(-\Delta_D)u_0\|_{L^r(\Omega)} + \left(\sum_{j \geq 1} \|e^{it\Delta_D} \Psi(-2^{-2j}\Delta_D)u_0\|_{L^r(\Omega)}^2 \right)^{1/2} \right) \right\|_{L^q(\mathbb{R})} \quad (3.11)$$

which, by Minkowski inequality, leads to

$$\|\chi u\|_{L^q(\mathbb{R}, L^r(\Omega))} \leq C \|u_0\|_{L^2(\Omega)}. \quad (3.12)$$

- Study of $w = (1 - \chi)u$. We follow here the method suggested by Staffilani and Tataru [37]. We have

$$i\partial_t w - \Delta_D w = [\Delta_D, \chi]u, \quad w|_{t=0} = (1 - \chi)u_0. \quad (3.13)$$

Since on the support of w the metric coincides with the flat metric $(\delta_{i,j})$, we can replace in (3.13) Δ_D by the Euclidian Laplace operator Δ (on \mathbb{R}^n) and we find

$$w(x, t) = e^{it\Delta}(1 - \chi)u_0 - i \int_0^t e^{i(t-s)\Delta} [\Delta_D, \chi]u(s) ds. \quad (3.14)$$

The contribution of the first term can be easily evaluated. For the second term we shall use the next lemma, due to Christ and Kiselev [12]:

Lemma 3.4. (Christ and Kiselev) Consider a bounded operator

$$T : L^{q'}(\mathbb{R}, B_1) \rightarrow L^q(\mathbb{R}, B_2)$$

given by a locally integrable kernel $K(t, s)$ with values in bounded operators from B_1 to B_2 , where B_1 and B_2 are Banach spaces. Suppose that $q' < q$. Then the operator

$$\tilde{T}f(t) = \int_{s < t} K(t, s)f(s) ds$$

is bounded from $L^{q'}(\mathbb{R}, B_1)$ to $L^q(\mathbb{R}, B_2)$ and

$$\|\tilde{T}\|_{L^{q'}(\mathbb{R}, B_1) \rightarrow L^q(\mathbb{R}, B_2)} \leq C(1 - 2^{-(1/q - 1/q')})^{-1} \|T\|_{L^{q'}(\mathbb{R}, B_1) \rightarrow L^q(\mathbb{R}, B_2)}.$$

This lemma allows (since $q > 0$) to replace the study of the second term in the right hand side of (3.14) by that of

$$W(x, t) = \int_0^\infty e^{i(t-s)\Delta} [\Delta_D, \chi] u(s) ds = U_0 U_0^*(x, t),$$

where $U_0 = e^{it\Delta}$ is bounded from $L^2(\mathbb{R}^n)$ to $L^q(\mathbb{R}, L^r(\mathbb{R}^n))$ and U_0^* is bounded from $L^2(\mathbb{R}, H_{comp}^{-1/2})$ to $L^2(\mathbb{R}^n)$, which gives the estimates for W and thus for w .

3.1 Applications

In this section we sketch the proofs of Theorem 1.8 and Theorem 1.9.

We start with Theorem 1.8. From Theorem 1.7 we have an estimate of the linear flow of the Schrödinger equation

$$\|e^{-it\Delta_D} u_0\|_{L^5(\mathbb{R}, L^{30/11}(\Omega))} \leq C \|u_0\|_{L^2(\Omega)}. \quad (3.15)$$

One may shift regularity by 1 and obtain

$$\|e^{-it\Delta_D} u_0\|_{L^5(\mathbb{R}, W^{1,30/11}(\Omega))} \leq C \|u_0\|_{H_0^1(\Omega)}. \quad (3.16)$$

Hence for small $T > 0$ the left hand side in (3.15), (3.16) will be small; for such T let $X_T := L^5((0, T], W^{1,30/11}(\Omega))$. One may then set up the usual fixed point argument in X_T , as if $u \in X_T$ then $u^5 \in L^1([0, T], H^1(\Omega))$.

Let us proceed with Theorem 1.9. From the work of Planchon and Vega [32], one has a global in time control on the solution u , at the level of $\dot{H}^{\frac{1}{4}}$ regularity:

$$u \in L^4((0, +\infty), L^4(\Omega)).$$

By interpolation with either mass or energy conservation, combined with the local existence theory, one may bootstrap this global in time control into

$$u \in L^{p-1}((0, +\infty), L^\infty(\Omega)),$$

from which scattering in $H_0^1(\Omega)$ follows immediately.

4 Appendix

4.1 Finite speed of propagation for the semi-classical equation

In this section we recall several properties of the semi-classical Schrödinger flow (for further discussions and proofs we refer the reader to [27]). Let M be a compact manifold with smooth boundary ∂M .

Definition 4.1. We say that a symbol $q(y, \eta) \in S_{\rho, \delta}^m$ is of type (ρ, δ) and of order m if

$$\forall \alpha, \beta \quad \exists C_{\alpha, \beta} > 0 \quad |\partial_y^\beta \partial_\eta^\alpha q(y, \eta)| \leq C_{\alpha, \beta} (1 + |\eta|)^{m - \rho|\alpha| + \delta|\beta|}.$$

For $q \in S_{1,0}^m$ we let $Op_h(q) = Q(y, hD, h)$ be the h -pseudo differential operator defined by

$$Op_h(q)f(y) = \frac{1}{(2\pi h)^n} \int e^{\frac{i}{h}(y-\tilde{y})\eta} q(y, \eta, h) f(\tilde{y}) d\tilde{y}.$$

We set $y = (x, t) \in M$ and denote $\eta = (\xi, \tau)$ the dual variable of y . Near a point $x_0 \in \partial M$ we can choose a system of local coordinates such that M is given by $M = \{x = (x', x_n) | x_n > 0\}$. We define the tangential operators

$$Op_{h, \text{tang}}(q)f(y) = \frac{1}{(2\pi h)^{n-1}} \int e^{\frac{i}{h}(y'-\tilde{y}')\eta'} q(y, \eta', h) f(\tilde{x}', x_n, \tilde{t}) d\tilde{y}' d\eta',$$

where $y = (x', x_n, t)$, $y' = (x', t)$, $\tilde{y}' = (\tilde{x}', \tilde{t})$, $\eta = (\xi', \xi_n, \tau)$, $\eta' = (\xi', \tau)$ and where the symbol $q(y, \eta', h) \in S_{1,0, \text{tang}}^m$ i.e. such that

$$\forall \alpha, \beta \quad \exists C_{\alpha, \beta} > 0 \quad |\partial_y^\alpha \partial_{\eta'}^\beta q(y, \eta', h)| \leq C_{\alpha, \beta} (1 + |\eta'|)^{m - |\beta|}.$$

In what follows we let (M, g) be a compact Riemannian manifold with strictly concave boundary satisfying the Assumptions 1.1. Let also $v_0 \in L^2(M)$ be compactly supported outside a small neighborhood of the boundary, $\Psi \in C_0^\infty((\alpha_0, \beta_0))$ and let $v(x, t) = e^{iht\Delta_g} \Psi(-h^2\Delta_g)v_0$ denote the linear semi-classical Schrödinger flow with initial data at time $t = 0$ equal to $\Psi(-h^2\Delta_g)v_0$ and such that $\|\Psi(-h^2\Delta_g)v_0\|_{L^2(M)} \lesssim 1$.

Let $\pi : T^*(\bar{M} \times \mathbb{R}) \rightarrow T^*(\partial M \times \mathbb{R}) \cup T^*(M \times \mathbb{R})$ be the canonical projection defined, for $y = (x, t)$, $\eta = (\xi, \tau)$ by

$$\pi|_{T^*(M \times \mathbb{R})} = Id, \quad \pi(y, \eta) = (y, \eta|_{T^*(\partial M \times \mathbb{R})}), \quad \text{for } (y, \eta) \in T^*(\bar{M} \times \mathbb{R})|_{\partial M \times \mathbb{R}}.$$

We introduce the characteristic set

$$\Sigma_b := \pi\{(y, \eta) | \eta = (\xi, \tau), \tau + |\xi|_g^2 = 0, -\beta_0 \leq \tau \leq -\alpha_0\},$$

where $|\xi|_g^2 = \langle \xi, \xi \rangle_g =: \xi_n^2 + r(x, \xi')$ denotes the inner product given by the metric g and where, due to the strict concavity of the boundary we have $\partial_{x_n} r(x', 0, \eta') < 0$.

Definition 4.2. We say that a point $\rho_0 = (y_0, \eta_0) \in T_b^*(\partial M \times \mathbb{R}) := T^*(\partial M \times \mathbb{R}) \cup T^*(M \times \mathbb{R})$ doesn't belong to the b -wave front set $WF_b(v)$ of v if there exists a h -pseudo-differential operator of symbol $q(y, \eta, h)$ (respectively $q(y, \eta', h)$ if $\rho_0 \in T^*(\partial M \times \mathbb{R})$) with compact support in (y, η) , elliptic at ρ_0 , and a smooth function $\phi \in C_0^\infty$ equal to 1 near y_0 , such that for every $\sigma \geq 0$ the following holds

$$\forall N \geq 0 \quad \exists C_N > 0 \quad \|Op_h(q)\phi v\|_{H^\sigma(M \times \mathbb{R})} \leq C_N h^N.$$

We shall write $\rho_0 \notin WF_b(v)$.

Proposition 4.3. (Elliptic regularity [27, Thm.3.1]) Let $q(y, \eta)$ a symbol such that $q = 0$ on Σ_b . Then for every $\sigma \geq 0$ we have

$$\forall N \geq 0 \quad \exists C_N > 0 \quad \|Op_h(q)v\|_{H^\sigma(M)} \leq C_N h^N.$$

Proposition 4.3 is proved by Lebeau [27] for eigenfunctions of the Laplace operator, but the same arguments apply in this setting. From Proposition 4.3 and [27, Sections 2,3] it follows:

Corollary 4.4. There exists a constant $D > 0$ such that

$$WF_b(v) \subset \Sigma_b \cap \{-\tau \in [\alpha_0, \beta_0], |\xi|_g \leq D\}.$$

Corollary 4.5. ([27, Chp.3]) Let $\varphi \in C_0^\infty(\mathbb{R})$ be equal to 1 near the interval $[-\beta_0, -\alpha_0]$. Then for t in any bounded interval I one has

$$\forall N \geq 1, \quad \exists C_N > 0 \quad |(1 - \varphi)(hD_t)v| \leq C_N h^N, \quad \forall t \in I. \quad (4.1)$$

Corollary 4.6. (Elliptic regularity at " ∞ ") Let $\vartheta \in C_0^\infty(\mathbb{R}^n)$ be equal to 1 on $\{|\xi|_g \leq D\}$, then

$$\forall N \geq 1, \quad \exists C_N > 0 \quad |(1 - \vartheta)(hD_x)v| \leq C_N h^N. \quad (4.2)$$

Proposition 4.7. Let $y_0 \notin pr_y(WF_b(v))$, where by pr_y we mean the projection on the variable $y = (x, t)$. Then there exists $\phi \in C_0^\infty$, $\phi = 1$ near y_0 such that for every $\sigma \geq 0$ we have

$$\forall N \geq 0 \quad \exists C_N > 0 \quad \|\phi v\|_{H^\sigma(M)} \leq C_N h^N.$$

Proof. Let φ, ϑ be the functions defined in Corollaries 4.5, 4.6. Then, using again Proposition 4.3 we have

$$v(x, t) = \varphi(hD_t)\vartheta(hD_x)v + O(h^\infty). \quad (4.3)$$

Let now $y_0 = (x_0, t_0) \notin pr_y(WF_b(v))$. It follows that for every $\eta \neq 0$, $(y_0, \eta) \notin WF_b(v)$ and in particular for every $\eta_0 \in \text{supp}(\vartheta) \times \text{supp}(\varphi)$ there exists a symbols $q_0(y, \eta, h)$ with compact support in (y, η) near (y_0, η_0) and elliptic at (y_0, η_0) , and there exists $\phi_0 \in C_0^\infty$ equal to 1 in a neighborhood U_0 of y_0 such that for every $\sigma \geq 0$

$$\forall N \geq 0 \quad \exists C_N > 0 \quad \|Op_h(q_0)\phi v\|_{H^\sigma(M)} \leq C_N h^N.$$

Eventually shrinking U_0 , suppose that q_0 is elliptic on $U_0 \times W_0$ where W_0 is an open neighborhood of η_0 . Then it follows that on U_0 , for every $\sigma \geq 0$

$$\forall N \geq 0 \quad \exists C_N > 0 \quad \|\phi v\|_{H^\sigma(U_0)} \leq C_N h^N.$$

Since the set $\text{supp}(\vartheta) \times \text{supp}(\varphi)$ is compact there exist η^α , $\alpha \in \{1, \dots, N\}$ for some fixed $N \geq 1$ and for each η^α there exist symbols q_α elliptic on some neighborhoods $U_\alpha \times W_\alpha$ of (y_0, η^α) and smooth functions $\phi_\alpha \in C_0^\infty$ equal to 1 on the neighborhoods U_α of y_0 , such that $\text{supp}(\vartheta) \times \text{supp}(\varphi) \subset \cup_{j=1}^N W_\alpha$. Let $\phi \in C_0^\infty$ be equal to 1 in an open neighborhood of y_0 strictly included in the intersection $\cap_{\alpha=1}^N U_\alpha$ (which has nonempty interior) and supported in the compact set $\cap_{\alpha=1}^N \text{supp}(\phi_\alpha)$. Considering a partition of unity associated to $(U_\alpha \times W_\alpha)_\alpha$ and using (4.3) we deduce that ϕ satisfies Proposition 4.7. \square

Proposition 4.8. [6, Lemma B.7] *Let $v(x, t) = e^{ith\Delta_g}\Psi(-h^2\Delta_g)v_0$ like before, $v_0 \in L^2(M)$ and let Q be a h -pseudo-differential operator of order 0, $t_0 > 0$ and $\tilde{\psi} \in C_0^\infty((-2t_0, -t_0))$. Let w denote the solution to*

$$\begin{cases} (ih\partial_t + h^2\Delta_g)w = ih\tilde{\psi}(t)Q(v), & \text{on } M \times \mathbb{R}, \\ w|_{\partial M} = 0, \quad w|_{t < -2t_0} = 0. \end{cases} \quad (4.4)$$

If $\rho_0 \in WF_b(w)$ then the broken bicharacteristic starting from ρ_0 has a nonempty intersection with $WF_b(v) \cap \{t \in \text{supp}(\tilde{\psi})\}$.

4.2 Proof of Proposition 3.1

After the change of variables $t = hs$ we are reduced to prove that

$$\|e^{ish\Delta_D}\Psi(-h^2\Delta_D)\chi V_0\|_{L^q(s \in I)L^r(\Omega)} \leq \quad (4.5)$$

$$Ch^{-\frac{n}{2}(\frac{1}{2} - \frac{1}{r})}\|\Psi(-h^2\Delta_D)\tilde{\chi}V_0\|_{L^2(\Omega)} + O(h^\infty)\|\Psi(-h^2\Delta_D)\tilde{\chi}V_0\|_{L^2(\Omega)},$$

where I is a fixed time interval, say $I = [-1, 2]$, $\chi, \tilde{\chi} \in C_0^\infty$, $\chi = 1$ near $\partial\Omega$, $\tilde{\chi} = 1$ on the support of χ and $C = C(I)$ is independent of h .

Lemma 4.9. *Let $\phi \in C^\infty(\mathbb{R})$ with Fourier transform $\hat{\phi}$ compactly supported in the interval $[-4, 4]$. Then the estimate (4.5) holds true if $\Psi(-h^2\Delta_D)$ is replaced by $\phi(h\sqrt{-\Delta_D})$, say*

$$\begin{aligned} & \|e^{ish\Delta_D}\phi(h\sqrt{-\Delta_D})\chi V_0\|_{L^q(s \in I)L^r(\Omega)} \leq \\ & Ch^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{r})}\|\phi(h\sqrt{-\Delta_D})\tilde{\chi}V_0\|_{L^2(\Omega)} + O(h^\infty)\|\phi(h\sqrt{-\Delta_D})\tilde{\chi}V_0\|_{L^2(\Omega)}, \end{aligned} \quad (4.6)$$

Proof. Recall from the Assumption 1.6 that Ω is the complementary in \mathbb{R}^n of a strictly convex obstacle Θ with smooth boundary. Let $Q \subset \mathbb{R}^n$ be an open cube such that $\bar{\Theta} \subset Q$ and define the manifold M as the complementary of Θ in the compact manifold obtained by imposing periodic boundary conditions on ∂Q . Notice that, defined in this way, M coincides with the Sinai billiard and it is a compact manifold with strictly concave boundary $\partial\Omega$. Define

$$\Delta_M := \sum_{j=1}^n \partial_j^2 \quad \text{the Laplace operator on } M.$$

We write

$$\phi(h\sqrt{-\Delta_D})\chi V_0 = \frac{1}{2\pi} \int_{-4}^4 e^{i\lambda h\sqrt{-\Delta_D}} \chi V_0 \hat{\phi}(\lambda) d\lambda.$$

By the finite speed of propagation of the wave flow, on a time of size $h\lambda$ the support of $e^{i\lambda h\sqrt{-\Delta_D}}\chi$ remains for $\lambda \in [-4, 4]$ in a neighborhood of the boundary $\partial\Omega$ where the coefficients of $-\Delta_D$ equal the coefficients of $-\Delta_M$,

$$-\Delta_D \chi V_0 = -\Delta_M \chi V_0,$$

consequently we have $\tilde{\chi}\phi(h\sqrt{-\Delta_D})\chi V_0 = \tilde{\chi}\phi(h\sqrt{-\Delta_M})\chi V_0$ and by the same argument

$$\begin{aligned} e^{ish\Delta_D}\phi(h\sqrt{-\Delta_D})\chi V_0 &= \tilde{\chi}e^{ish\Delta_D}\phi(h\sqrt{-\Delta_D})\chi V_0 + O(h^\infty) = \\ \tilde{\chi}e^{ish\Delta_M}\phi(h\sqrt{-\Delta_M})\chi V_0 + O(h^\infty) &= e^{ish\Delta_M}\phi(h\sqrt{-\Delta_M})\chi V_0 + O(h^\infty), \quad \forall s \in [-4, 4], \end{aligned} \quad (4.7)$$

where we recall that $\tilde{\chi} \in C_0^\infty$ is equal to 1 near the support of χ .

In order to prove (4.6) we apply Theorem 1.3 to the semi-classical $e^{ish\Delta_M}\phi(h\sqrt{-\Delta_M})\chi V_0$ on the compact manifold M with strictly convex boundary $\partial\Omega$ together with the next lemma:

Lemma 4.10. *Under the Assumptions 1.1 and for any bounded interval of time I , the estimates*

$$\|e^{ish\Delta_M}\Psi(-h^2\Delta_M)\chi V_0\|_{L^q(s \in I)L^r(M)} \leq C(I)h^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{r})}\|\Psi(-h^2\Delta_M)\tilde{\chi}V_0\|_{L^2(M)} \quad (4.8)$$

hold true for any $\Psi \in C_0^\infty(\mathbb{R} \setminus \{0\})$ if and only if the estimates

$$\|e^{ish\Delta_M}\phi(h\sqrt{-\Delta_M})\chi V_0\|_{L^q(s \in I)L^r(M)} \leq C(I)h^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{r})}\|\phi(h\sqrt{-\Delta_M})\tilde{\chi}V_0\|_{L^2(M)} \quad (4.9)$$

hold true for any $\phi \in C^\infty$ with $\hat{\phi} \in C_0^\infty(\mathbb{R})$.

In order to prove the last lemma one uses exactly the same arguments like in the end of proof of Proposition 3.1 below.

□

End of the proof of Proposition 3.1

Let now $\psi \in C_0^\infty(\mathbb{R}_+ \setminus \{0\})$ such that $\Psi(\tau^2) = \psi(\tau)$ and write ψ as a sum

$$\psi(\tau) = \sum_{k \in \mathbb{Z}} \phi_k(\tau), \quad \|\partial^\alpha \phi_k\|_{L^\infty} \leq C_{k,\alpha,N} (1 + 2^{-k})^{-N}, \quad \forall N \geq 0, \quad \forall \alpha \in \mathbb{N} \quad (4.10)$$

and where the Fourier transform of each ϕ_k is compactly supported in

$$[-4, 4] \quad \text{if } k = 0; \quad [2^k, 2^{k+2}] \quad \text{if } k \geq 1; \quad [-2^{k+2}, -2^k] \quad \text{if } k \leq -1. \quad (4.11)$$

The contribution of ϕ_0 in (4.5) has been dealt with in Lemma 4.9. In order to estimate the contribution for the remaining ϕ_k , $|k| \geq 1$, the idea is to rescale the time variable and to use the (proof of) Lemma (4.9) to estimate the norms of the rescaled operators: using the fast decay of the Fourier transforms $\hat{\phi}_k$ we obtain a contribution $O(2^{-k\infty})$ for every $|k| \geq 1$.

Lemma 4.11. *Let $|k| \geq 1$ and $\phi_k \in \mathcal{S}(\mathbb{R})$ given by (4.11). Then*

$$\|e^{ish\Delta_D} \phi_k(h\sqrt{-\Delta_D}) \tilde{\chi} V_0\|_{L^q(s \in I) L^r(\Omega)} \leq O(2^{-k\infty}) \|\phi_k(h\sqrt{-\Delta_D}) \tilde{\chi} V_0\|_{L^2(\Omega)}. \quad (4.12)$$

Proof. Let $V_k := e^{ish\Delta_D} \phi_k(h\sqrt{-\Delta_D}) \tilde{\chi} V_0$ solve the semi-classical Schrödinger equation

$$\begin{cases} ih \partial_s V_k + h^2 \Delta_D V_k = -e^{ish\Delta_D} \phi_k(h\sqrt{-\Delta_D}) [h^2 \Delta_D, \tilde{\chi}] V_0, \\ V_k|_{s=0} = \phi_k(h\sqrt{-\Delta_D}) \tilde{\chi} V_0. \end{cases} \quad (4.13)$$

For $|k| \geq 1$ we introduce $W_k(x, 2^{|k|}s) = V_k(x, s)$ defined on an interval of time of size $\simeq 2^{|k|}$ and we set $h_k = h2^{-|k|}$. Then W_k solves

$$\begin{cases} ih_k \partial_s W_k + h_k^2 \Delta_D W_k = -e^{ish\Delta_D} \phi_k((h\sqrt{-\Delta_D}) [h_k^2 \Delta_D, \tilde{\chi}] V_0, \\ W_k|_{s=0} = \phi_k(h\sqrt{-\Delta_D}) \tilde{\chi} V_0. \end{cases} \quad (4.14)$$

Inspired from [8], we write the Duhamel formula to obtain Strichartz estimates (without losses) for W_k on intervals of time $[m, m+1]$, $m \in [-2^{|k|}, \dots, 2^{|k|}] \cap \mathbb{Z}$

$$\|W_k\|_{L^q(s \in [m, m+1]) L^r(\Omega)} \lesssim \|\phi_k(h\sqrt{-\Delta_D}) \tilde{\chi} V_0\|_{L^2(\Omega)}, \quad (4.15)$$

where (q, r) is an admissible pair in dimension n , $q > 2$. Summing up the q -th powers of those norms for all $-2^{|k|} \leq m \leq 2^{|k|}$ we obtain Strichartz estimates for W_k on $I_k = [-2^{|k|}, 2^{|k|+1}]$,

$$\|W_k\|_{L^q(s \in I_k) L^r(\Omega)} \lesssim 2^{k/q} \|\phi_k(h\sqrt{-\Delta_D}) \tilde{\chi} V_0\|_{L^2(\Omega)}. \quad (4.16)$$

On the other hand, if $|k| \geq 1$ then performing integrations by parts in the integral

$$\widehat{\phi}_k(\lambda) = \int e^{-i\tau\lambda} \phi_k(\tau) d\tau, \quad \phi_k \in \mathcal{S}(\mathbb{R})$$

and using (4.10) we find that for every $N \geq 1$ there exists $C_N > 0$ such that $|\widehat{\phi}_k| \leq C_N 2^{-kN}$. This remark, together with (4.16), yields (4.12). \square

We conclude the proof of Proposition 3.1 writing the function Ψ in the form (4.10) and noticing that the only nontrivial contribution in (4.5) comes from the term corresponding to ϕ_0 . Since ϕ_0 is compactly supported, it remains to apply Lemma 4.9.

References

- [1] R. Anton, Global existence for defocusing cubic NLS and Gross-Pitaevskii equations in exterior domains, *Journal de Mathématiques Pures et Appliquées*, to appear
- [2] M. Blair, H. Smith, C. Sogge, On Strichartz estimates for the Schrödinger operators in compact manifolds with boundary, *Proceedings AMS*, 136:247-256, 2008
- [3] J.M. Bouclet, N. Tzvetkov, On global Strichartz estimates for non trapping metrics, *Journal of Functional Analysis*, 254, 6:1661-1682, 2008
- [4] J. Bourgain, Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations, I. Schrödinger equations, *Geom. Funct. Anal.* 3, no.2:107–156, 2003.
- [5] N. Burq, Estimations de Strichartz pour des perturbations à longue portée de l'opérateur de Schrödinger, *Séminaire X-EDP 2002*
- [6] N. Burq, Contrôle de l'équation des plaques en présence d'obstacles strictement convexes, *Mémoires de la S.M.F.* 55, 1993
- [7] N. Burq, P. Gérard, N. Tzvetkov, On nonlinear Schrödinger equations in exterior domains, *Ann.I.H.Poincaré*, 21:295-318, 2004
- [8] N. Burq, P. Gérard, N. Tzvetkov, Strichartz inequality and the nonlinear Schrödinger equation on compact manifolds, *Am. J. Math.* 126, 569-605, 2004
- [9] N. Burq, G. Lebeau, F. Planchon, Global existence for energy critical waves in $3D$ domains, *J.Amer.Math.Soc.* 21:831-845, 2008

- [10] N. Burq, F. Planchon, Global existence for energy critical waves in 3D domains: Neumann boundary conditions, Amer.J.of Math, to appear
- [11] T. Cazenave, F. Weissler, The Cauchy problem for the critical Schrödinger equation in H^s , Nonlinear Anal. 14:807-836, 1990
- [12] M. Christ, A. Kiselev, Maximal functions associated to filtrations, J.Funct.Anal. 179, 2:409-425, 2001
- [13] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, T. Tao, Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in \mathbb{R}^3 , arXiv:math/0402129
- [14] P. Constantin, J.C. Saut, Local smoothing properties of Schrödinger equations, Indiana Univ.Math.J.38:791-810, 1989
- [15] E.B. Davies, Heat kernels and spectral theory, Cambridge Univ. Press, 1989
- [16] S.I. Doi, Smoothing effects of Schrödinger equations, second ed., Textos de Métodos Matemáticos 26, 1996
- [17] J. Ginibre, G. Velo, The global Cauchy problem for the nonlinear Schrödinger equation, Ann.I.H.Poincaré, Anal.Non Linéaire 2:309-327, 1985
- [18] J. Ginibre, G. Velo, Smoothing properties and retarded estimates for some dispersive evolution equations , Comm.Math.Phys. 144:163-188, 1992
- [19] J. Ginibre, G. Velo, Generalized Strichartz inequalities for the wave equation, J.Funct.Anal.133:50-68, 1995
- [20] D. Grieser, L^p bounds for eigenfunctions and spectral projections of the Laplacian near concave boundaries, Thesis, UCLA, 1992
- [21] L. Hörmander, The Analysis of Linear Partial Diff. Operators, Springer-Verlag 1985
- [22] O. Ivanovici, Counterexamples to Strichartz estimates for the wave equation in domains, arXiv:0805.2901
- [23] O. Ivanovici, F. Planchon, A poor man's square function estimate on domains, arXiv:0812.2733
- [24] L. Kapitanski, Some generalizations of the Strichartz-Brenner inequality, Leningrad Math.J.1:693-676, 1990

- [25] T. Kato, On nonlinear Schrödinger equations, *Ann.I.H.P., Phys.Théor.*, 46:13-129, 1987
- [26] M. Keel, T. Tao, Endpoints Strichartz estimates, *Amer.J.Math.*, 120:955-980,1998
- [27] G. Lebeau, Contrôle de l'équation de Schrödinger, *J.Math.Pures Appl.* 71:267-291, 1992
- [28] H. Lindblad, C. Sogge, On existence and scattering with minimal regularity for semi-linear wave equations, *J.Funct.Anal.*130:357-426, 1995
- [29] R. Melrose, M. Taylor, Near peak scattering and the corrected Kirchoff approximation for a convex obstacle, *Adv.Math.*55:242-315, 1985
- [30] R. Melrose, M. Taylor, The radiation pattern of a diffractive wave near the shadow boundary, *Comm.Partial Diff.Eq.* 11:599-672, 1985
- [31] F. Nier, A variational formulation of Schrödinger-Poisson systems in dimension $d \leq 3$, *Comm.Part.Diff.Equations* 18, 7-8:1125-1147, 1993
- [32] F. Planchon, L. Vega, Bilinear virial identities and applications, arXiv:0712.4076, *Ann.Sci.E.N.S.* to appear
- [33] H.F. Smith, C.D. Sogge, L^p regularity for the wave equation with strictly convex obstacles, *Duke Mathematical Journal*, vol.73, 1:97-153, 1994
- [34] H.F. Smith, C.D. Sogge, On the critical semilinear wave equation outside convex obstacles, *J.A.M.S.*, vol.8, 4:897-916, 1995
- [35] H. Smith, C.D. Sogge, On the L^p norm of spectral clusters for compact manifolds with boundary, *Acta Mathematica*, 2006, arXiv:math.AP/0605682
- [36] C.D. Sogge, *Fourier integrals in classical analysis*, Cambridge Univ.Press, Cambridge and New York, 1993
- [37] G. Staffilani, D. Tataru, Strichartz estimates for a Schrödinger operator with nonsmooth coefficients, *Comm.Part.Diff.Eq.* 27, vol.5-6:1337-1372, 2002
- [38] R.S. Strichartz, Restriction of Fourier transforms to quadratic surfaces and decay of solutions of wave equation, *Duke.Math.J.*, 44:705-714, 1977
- [39] H. Triebel, *Theory of function spaces*, volume 78 of *Monographs in Mathematics*, Birkhäuser Verlag, Basel, 1983
- [40] M. Zworski, High frequency scattering by a convex obstacle, *Duke Math.J.* 61:545-634, 1990