



HAL
open science

On a nonlocal Zakharov equation

Brigitte Bidégaray

► **To cite this version:**

Brigitte Bidégaray. On a nonlocal Zakharov equation. *Nonlinear Evolution Equations and Dynamical Systems (NEEDS'92)*, 1992, Dubna, Russia. pp.348-355. ⟨hal-00319997⟩

HAL Id: hal-00319997

<https://hal.science/hal-00319997v1>

Submitted on 9 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

On a nonlocal Zakharov equation

B. Bidégaray

*CMLA, Ecole Normale Supérieure de Cachan, 61 av. du Pdt Wilson
94235 Cachan CEDEX, France.*

July 11th 1992

Abstract

This article deals with the Cauchy problem for a nonlocal Zakharov equation. We will first recall the physical motivation (due to V.E. Zakharov) of this system. Then we will study the local Cauchy problem for certain initial data, and will identify the limit of the solutions when the ion velocity tends to infinity.

1 Origin of the nonlocal Zakharov system

The Physical theory which follows has been developed by V.E. Zakharov^{5,6} to describe the Langmuir Oscillations in Plasma Physics.

We assume that:

1. the plasma is sufficiently uniform,
2. the magnetic field is sufficiently weak,
3. the nonlinearity level is not too high,
4. there are no transverse high frequency electromagnetic waves.

We consider the following equations that model the phenomena:

Linearized hydrodynamical equations

$$\frac{\partial}{\partial t} \delta n_e + \operatorname{div}(n_0 + \delta n) \vec{V}_e = 0,$$

$$\frac{\partial}{\partial t} \delta \vec{V}_e + \frac{3V_{Te}^2}{n} \vec{\nabla} \delta n = \frac{e}{m_e} \vec{E}.$$

Maxwell's equation

$$\frac{1}{c^2} \frac{\partial^2 \vec{E}}{\partial t^2} + \operatorname{curl} \operatorname{curl} \vec{E} + \frac{4\pi e}{c^2} (n_0 + \delta n) \frac{\partial}{\partial t} \delta \vec{V}_e = 0.$$

Vlasov's equation

$$\frac{\partial f_i}{\partial t} + (V \cdot \vec{\nabla}) f_i - n \frac{e}{m_i} \vec{\nabla} \varphi_{el} \frac{\partial f_i}{\partial V} = 0,$$

$$\delta n_i = \frac{en_0}{T_e}(\varphi_{el} - \varphi).$$

We moreover set

$$\vec{E} = \frac{1}{2} \left(\vec{E} \exp(-i\omega_{pl}t) + \vec{E}^* \exp(i\omega_{pl}t) \right),$$

and

$$\vec{E} = \vec{\nabla}\psi.$$

Eq. 1-7 imply that

$$\Delta(2i\omega_{pl}\psi_t + 3V_{Te}^2 \Delta\psi) = \omega_{pl}^2 \operatorname{div}\left(\frac{\delta n}{n_0} \nabla\psi\right).$$

Two different hypothesis can be made:

First hypothesis

The nonlinear phenomena have such a long period that the ions have enough time to reach the Boltzmann distribution law in a low frequency field:

$$\frac{\delta n}{n_0} = -e \frac{\varphi_{el}}{T_i}.$$

After some computations and a change of scale, Eq. 8 and 9 lead to

$$\Delta(i\psi_t + \Delta\psi) = \operatorname{div}(|\nabla\psi|^2 \nabla\psi).$$

This equation has been widely studied by T. Colin^{1,2}.

Second hypothesis

The ions do not have the time to reach this distribution:

$$\left(\frac{\partial^2}{\partial t^2} + 2\gamma_s \frac{\partial}{\partial t} - c_s^2 \Delta \right) \delta n = \frac{1}{16\pi m_i} \Delta |E|^2.$$

We will suppose that the damping rate is zero, i.e. we neglect $2\gamma_s \frac{\partial}{\partial t}$.

After a change of scale, Eq. 11 and 8 become

$$\begin{cases} \Delta(i\psi_t + \Delta\psi) = \operatorname{div}(n\nabla\psi), \\ \frac{1}{c^2} n_{tt} - \Delta n = \Delta(|\nabla\psi|^2). \end{cases} \quad (1)$$

In what follows, we will not give any complete proof. They may be found in an article which is to appear⁷.

2 Existence and uniqueness for the Cauchy Problem

We first consider the system 12 with $c = 1$:

$$\begin{cases} \Delta(i\psi_t + \Delta\psi) = \operatorname{div}(n\nabla\psi), \\ n_{tt} - \Delta n = \Delta(|\nabla\psi|^2). \end{cases} \quad (2)$$

We set $B = \nabla(-\Delta)^{-1}\nabla$. and $\varphi = \nabla\psi$. Eq. 13 is equivalent to

$$\begin{cases} i\varphi_t + \Delta\varphi = -B(n\varphi), \\ n_{tt} - \Delta n = \Delta(|\varphi|^2). \end{cases} \quad (3)$$

B is homogeneous of order 0 in the Fourier variables. Thanks to Calderón-Zygmund's theorem we have the following result:

For all $1 < p < \infty$, there exists $C_{s,p}$ such that

$$\|Bf\|_{\#2} \leq C_{s,p} \|f\|_{\#2}.$$

2.1 Theorem for the Cauchy Problem

Theorem 1 *Let us consider the problem on R^N , $N = 1, 2, 3$.*

$$\begin{cases} i\dot{\varphi} + \Delta\varphi = \nabla\Delta^{-1}\nabla\cdot(n\varphi), \\ \ddot{n} - \Delta n = \Delta|\varphi|^2, \\ n(x, 0) = n_0(x), \\ \partial_t n(x, 0) = n_1(x), \\ \varphi(x, 0) = \varphi_0(x), \end{cases} \quad (4)$$

with $n_0 \in H^1$, $n_1 \in L^2$ and $\varphi_0 \in H^2$.

Then there exists a time $T > 0$ depending only on $\|n_0\|_{\#2}$, $\|n_1\|_{\#2}$, $\|\varphi_0\|_{\#2}$ and N and a unique solution $(\varphi(t), n(t))$ to Eq. 16 which satisfies

$$\begin{cases} \varphi(t) \in \mathcal{C}^0([0, T]; H^2) \cap \mathcal{C}^1([0, T]; L^2), \\ \varphi(t) \in W^{1,8/N}(0, T; L^4), \\ n(t) \in \mathcal{C}^0([0, T]; H^1) \cap \mathcal{C}^1([0, T]; L^2) \cap \mathcal{C}^2([0, T]; H^{-1}). \end{cases} \quad (5)$$

Thanks to Eq. 15, we have been inspired for the idea of the proof by the work of Ozawa and Tsutsumi³. We first make a change of variables (in order not to lose regularity at each step). We then use a fixed point method to find existence and uniqueness in a subset of the original functional space. To conclude, we prove uniqueness in the whole space and return to the initial variables.

2.2 Setting of the fixed point method

Setting $F = \partial_t \varphi$, we formally get

$$\begin{cases} iF_t + \Delta F = -B(\partial_t n(\varphi_0 + \int_0^t F ds) + nF), \\ n_{tt} - \Delta n = \Delta(|\varphi|^2), \\ \varphi = (-\Delta + 1)^{-1} \{iF + B(n(\varphi_0 + \int_0^t F ds)) + (\varphi_0 + \int_0^t F ds)\}, \end{cases}$$

with $F(0) = F_0$, $n(0) = n_0$, $\partial_t n(0) = n_1$.

We work in R^N with $1 \leq N \leq 3$. We use the following functional space:

$$X = [L^\infty(I; L^2) \cap L^8/N(I; L^4)] \oplus [L^\infty(I; H^1) \cap W^{1,\infty}(I; L^2)],$$

where $I = [0, T]$.

We set $N = (N_1, N_2)$ with

$$\begin{cases} N_1[F, n](t) = U(t)F_0 + i \int_0^t U(t-s) \{B(\partial_s n(\varphi_0 + \int_0^s F d\tau) + nF)\} ds, \\ N_2[F, n](t) = \cos(\omega t)n_0 + \omega^{-1} \sin(\omega t)n_1 + \int_0^t \omega^{-1} \sin(\omega(t-s)) \Delta |\varphi(s)|^2 ds, \\ (-\Delta + 1)\varphi = \{iF + B(n(\varphi_0 + \int_0^t F ds)) + (\varphi_0 + \int_0^t F ds)\}, \end{cases}$$

where $U(t)$ is the group generated by the Schrödinger operator and ω denotes $\sqrt{-\Delta}$ (multiplication by $|\xi|$ in Fourier variables).

We set

$$a = \max\{\|\varphi_0\|_{\#2}, \|\varphi_0\|_{\#2}, \|\Delta\varphi_0 + B(n_0\varphi_0)\|_{\#2}, \|n_0\|_{\#2} + \|n_1\|_{\#2}\},$$

$$Y = \{(F(t), n(t)) \in X / \begin{aligned} &\|F\|_{\#2} \leq 2a, \|F\|_{\#2} \leq 2\delta a, \\ &\|n\|_{\#2} \leq 2, \left\| \frac{dn}{dt} \right\|_{\#2} \leq 2a \}. \end{aligned} \quad (6)$$

The fixed point method will consist in solving

$$\begin{cases} N1[F, n] = F, \\ N2[F, n] = n. \end{cases}$$

$N: Y \rightarrow Y$ is a contraction if T is sufficiently small, thus we may find a fixed point.

2.3 Return to the initial problem

The fixed point method yields:

$$\begin{cases} F(t) \in \left[\bigcap_{j=0}^1 \mathcal{C}^j([0, T]; H^{-2j}) \right] \cap L^{8/N}(0, T; L^4), \\ n(t) \in \bigcap_{j=0}^2 \mathcal{C}^j([0, T]; H^{1-j}), \\ \varphi(t) \in \mathcal{C}([0, T]; H^2). \end{cases}$$

This regularity enables us to return to the initial variables; we get the desired regularity and two conservation laws:

$$\int |\varphi(t)|^2 = \int |\varphi_0|^2,$$

$$\int (|\nabla\varphi(t)|^2 + n(t)|\varphi(t)|^2 + \frac{1}{2}(\nabla\Phi(t))^2 + \frac{1}{2}n^2(t)) = \int (|\nabla\varphi_0|^2 + n_0|\varphi_0|^2 + \frac{1}{2}(\nabla\Phi_0)^2 + \frac{1}{2}n_0^2),$$

where $-\Delta\Phi = \frac{\partial n}{\partial t}$.

3 Limit as c tends to ∞

We now consider the problem

$$\begin{cases} \frac{1}{c^2} n_{tt} - \Delta(n + |E|^2) = 0, \\ iE_t + \Delta E + B(nE) = 0. \end{cases} \quad (7)$$

The result obtained in Section 2 still holds but T depends on c .

We shall work with variables in H^s with $s > \left\lceil \frac{k}{2} \right\rceil + 3$.

There is no need to prove uniqueness since this has been already done in some larger space.

The formal limit of the solution is the couple $(-|\tilde{E}|^2, \tilde{E})$ solution of

$$i\tilde{E}_t + \Delta\tilde{E} - B(|\tilde{E}|^2\tilde{E}) = 0.$$

3.1 The Theorem for the Limit

Theorem 2 *When c tends to ∞ ,*

$$n^c + |E^c|^2 \rightarrow 0 \text{ in } \mathcal{C}^0([0, T] \times R^k),$$

$$\nabla(n^c + |E^c|^2) \rightarrow 0 \text{ in } \mathcal{C}^0([0, T]; H^{s-2}),$$

$$E^c \rightarrow \tilde{E} \text{ in } \mathcal{C}^1([0, T] \times R^k) \cap \mathcal{C}([0, T]; \mathcal{C}^2),$$

where \tilde{E} is the unique solution of

$$i\tilde{E}_t + \Delta\tilde{E} - B(|\tilde{E}|^2\tilde{E}) = 0.$$

This time we adapt the method of Schochet and Weinstein⁴. We first carry out a transformation of the system into a dispersive perturbation of a symmetric hyperbolic one. Then we prove the existence of a regular solution for a time independent of c and pass to the limit when c tends to ∞ .

3.2 Transformation of the system

We set

$$\begin{cases} V = -\frac{1}{c}\Delta^{-1}\nabla n_t, \\ Q = n + |E|^2, \end{cases} \quad (8)$$

$$\sqrt{2}E = F + iG \text{ et } \sqrt{2}\nabla E = H + iL,$$

and

$$U = {}^t(Q, V, F, G, H, L),$$

then the system may be rewritten as

$$U_t + \sum_{j=1}^k \{R(A^j(U)U_{x_j}) + cC^j U_{x_j}\} + S(\tilde{B}(U)U) = K\Delta U.$$

R and S are Calderón–Zygmund’s operators.

\tilde{B} is a nonlocal operator.

K is an antisymmetric matrix.

C^j and $A^j(U)$ are symmetric matrices.

These properties will be very useful to make estimation in Sobolev spaces and to use the classical theory of hyperbolic equations.

3.3 Existence of a regular solution for a time independent of c

To this end, we use the following iteration scheme:

$$U^0(x, t) = U_0(x),$$

$$\frac{\partial U^{p+1}}{\partial t} + \sum_{j=1}^k \{R(A^j(U^p)U_{x_j}^{p+1}) + cC^j U_{x_j}^{p+1}\} + S(\tilde{B}(U^p)U^{p+1}) = K\Delta U^{p+1},$$

$$U^{p+1}(x, 0) = U_0(x).$$

One can show the following estimates.

- $\forall p \geq 0 \quad \| \| U^p \| \|_{s,T} \leq \delta,$
- $\forall p \geq 0 \quad \| \| U^{p+1} - U^p \| \|_{0,T} \leq C \| \| U^p - U^{p-1} \| \|_{0,T},$
with $C < 1$.

- Then
$$\begin{aligned} U^p &\rightarrow U \text{ in } L^\infty(0, T; L^2), \\ U^p &\text{ is bounded in } L^\infty(0, T; H^s). \end{aligned} \tag{9}$$

Then $U \in \mathcal{C}([0, T]; \mathcal{C}^1)$ and the solution is a classical one.

Moreover we have
$$\begin{aligned} U &\in Lip([0, T]; H^{s-2}), \\ U &\in \mathcal{C}([0, T]; H^s) \cap \mathcal{C}^1([0, T]; H^{s-2}). \end{aligned} \tag{10}$$

The time T is independent of c (because \mathcal{C}^j is symmetric).

To gain some regularity we use the theory of commutators.

There is no problem in returning to the initial variables and we get

Theorem 3 *Let $s \geq \left\lceil \frac{k}{2} \right\rceil + 3$. For $n_0 \in H^s$, $n_1 \in H^{s-1}$ and $E_0 \in H^{s+1}$, there exists a unique solution to system 27 endowed with the following initial data:*

$$\begin{cases} n(0, x) = n_0(x), \\ \partial_t n(0, x) = n_1(x), \\ E(0, x) = E_0(x), \end{cases} \tag{11}$$

on a time interval $[0, T]$, T not depending on c but only on $\|n_0\|_{\#2}$, $\|n_1\|_{\#2}$ and $\|E_0\|_{\#2}$. Moreover, for all $t \in [0, T]$ we have the estimate

$$\|E^c\|_{\#2} + \|E_t^c\|_{\#2} + \|n^c\|_{\#2} + \frac{1}{c} \|n_t^c\|_{\#2} + \frac{1}{c^2} \|n_{tt}^c\|_{\#2} \leq Cst$$

Thanks to the two conservation laws Eq. 25 and 26 we can, as for the classical Zakharov equation ($B = -I$), show that the solutions are global in time in the 1-dimensional case and also in the 2-dimensional case when the initial data are sufficiently small.

ACKNOWLEDGEMENTS I want to thank J.C. Saut for suggesting me this subject and his encouragements and T. Colin for helpful discussions.

1. T. Colin, Sur une équation de Schrödinger non linéaire et non locale intervenant en physique des plasmas. *C.R.A.S. Paris*, **314**, série I, (1992) 449-453.
2. T. Colin, On a nonlocal, nonlinear Schrödinger equation occurring in plasma Physics. *Publication CMLA*, ENS Cachan, **9208** (1992).
3. T. Ozawa, Y. Tsutsumi, Existence and Smoothing Effect of Solutions for the Zakharov Equations. 1990.
4. S.H. Schochet, M.I. Weinstein, The Nonlinear Schrödinger Limit of the Zakharov Equations Governing Langmuir Turbulence. *Commun. Math. Phys.*, **106**, (1986) 569-580.
5. V.E. Zakharov, Collapse and Self-Focusing of Langmuir Waves. in *Basic Plasma Physics II*, ed. A.A. Galeev and R.N. Sudan, (North Holland, 1984).
6. V.E. Zakharov, Collapse of Langmuir Waves. *Soviet Physics JETP*, **Vol. 35, 5** (1972) 908-914.
7. B. Bidégaray, On a nonlocal Zakharov equation, to appear