

# Hypergeometric Solutions of Systems of Linear Difference Equations

Moulay Barkatou

`moulay.barkatou@unilim.fr`

XLIM, Université de Limoges-CNRS  
FRANCE.

June 24, 2008

## Introduction

- ▶ This talk is based on a joint work with Mark van Hoeij (MvH) from FSU.
- ▶ We present a **direct** algorithm for computing hypergeometric solutions of systems of linear difference equations.
- ▶ “*direct algorithm*” means “*without first reducing the system to a single scalar equation by the use of a cyclic vector*”
- ▶ Motivation: finding  $d$ 'th order ( $d \geq 2$ ) right-hand factors of a given linear difference operator with polynomial coefficients.

## The differential case

- ▶ Beke (1894) gave two algorithms for factoring linear differential operators.
- ▶ Given a differential operator  $L$  of order  $n$ , the first algorithm computes the exponential solutions of  $L$ , or equivalently, the first order right-hand factors of  $L$ .
- ▶ The second algorithm reduces the problem of finding  $d$ 'th order right-hand factors of  $L$  to computing 1'st order right-hand factors of the so-called associated equations, which have order  $\binom{n}{d}$ .
- ▶ This was improved by M. Bronstein (1994).

- ▶ Bronstein's approach has several efficiency advantages:
  - (i) Instead of computing exponential solutions of a number of associated equations  $L_1(y_1) = 0$ ,  $L_2(y_2) = 0$ ,  $\dots$  one now computes the exponential solutions of just one associated system  $Y' = AY$ .
  - (ii) Consequently: a combinatorial problem disappears, namely the problem of deciding which exponential solution of each  $L_i$  should be combined with each other.
  - (iii) The most important advantage is: The coefficients of the associated system are much smaller than the coefficients of the associated equations.

- ▶ A consequence of advantage (iii) is that solving this one system  $Y' = AY$  potentially takes less time, or even much less time, than solving even one of the associated equations  $L_i$ , let alone solving all of them.
- ▶ We would not have this advantage if we used the cyclic vector approach to solve  $Y' = AY$ , because the resulting operator would be an associated equation.
- ▶ Thus, to benefit fully from the advantages of Bronstein's approach, we need to be able to directly compute the exponential solutions of  $Y' = AY$  without first reducing this system to an operator by the use of a cyclic vector.
- ▶ Algorithms for directly computing the exponential solutions are implemented in ISOLDE (a Maple package developed by E. Pfluegel and Bark.) .

## The difference case

- ▶ First order right-hand factors of a difference operator  $L$  correspond to so-called hypergeometric solutions.
- ▶ Algorithms for computing hypergeometric solutions of difference operators are given in [Petkovšek (1992), Cluzeau-van Hoeij (2006)].
- ▶ For computing higher order factors, Bronstein's approach works for the difference case as well in much the same way.
- ▶ Again, one reduces the computation of a  $d$ 'th order factor of  $L$  to solving (hypergeometric solutions) of an associated difference system

$$Y(x+1) = M(x)Y(x).$$

## Main Contributions

- ▶ We show that Petkovšek's algorithm for hypergeometric solutions of operators can be extended to computing hypergeometric solutions of systems.
- ▶ As in the scalar case we need to compute first the local data at infinity (the integer slopes of the Newton polygon and the corresponding polynomials)
- ▶ In the operator case the local data can be easily obtained from the coefficients of the operator.
- ▶ However, in the case of systems, we need a reduction algorithm [Bark. 1989] to determine the local data.
- ▶ Algorithms for computing polynomial and rational solutions of difference systems are given in [Abramov&Bark. (1998), Barkatou (1999)].

## Petkovšek's algorithm

Let  $\mathbf{C}$  be a field of characteristic zero.

An operator

$$L = a_n \tau^n + \dots + a_0 \tau^0 \in \mathbf{C}(x)[\tau]$$

acts as follows:

If  $u = u(x)$  is a function then

$$L(u) = a_n u(x+n) + \dots + a_1 u(x+1) + a_0 u(x)$$

If

$$L(u) = 0 \quad \text{and} \quad \frac{u(x+1)}{u(x)} \in \mathbf{C}(x)$$

then  $u$  is called a **hypergeometric solution** and corresponds to a **right-hand factor**  $\tau - r$  of  $L$  where

$$r = \frac{u(x+1)}{u(x)}.$$



A hypergeometric solution  $u$  of  $L = a_n\tau^n + \dots + a_0\tau^0$  can always be written as

$$u(x) = c^x P(x) \prod_i \Gamma(x - \alpha_i)^{e_i}$$

for some

$$c \in \mathbf{C}^*, \quad P(x) \in \mathbf{C}[x], \quad \alpha_i \in \mathbf{C}, \quad e_i \in \mathbb{Z}.$$

Write

$$A = \prod_{e_i > 0} (x - \alpha_i)^{e_i} \quad \text{and} \quad B = \prod_{e_i < 0} (x - \alpha_i)^{-e_i}$$

so that

$$u(x) = c^x P(x) \text{Sol}\left(\tau - \frac{A}{B}\right).$$

$$L = a_n(x)\tau^n + \cdots + a_0(x)\tau^0.$$

- ▶ Petkovšek (1992) gave a criterion for  $A, B \in \mathbf{C}[x]$ , namely:

$A$  divides  $a_0(x)$

$B$  divides  $a_n(x - n + 1)$

- ▶ This leaves only a finite (but exponential) number of potential  $A, B$  in the hypergeometric solutions:

$$\begin{aligned} u(x) &= c^x P(x) \text{Sol}\left(\tau - \frac{A}{B}\right) \\ &= c^x P(x) \prod_i \Gamma(x - \alpha_i)^{e_i} \end{aligned}$$

► Petkovšek's algorithm (1992):

For all possible combinations of:

- a monic factor  $A \in \mathbf{C}[x]$  of  $a_0$ ,
- a monic factor  $B$  of  $a_n(x - n + 1)$ ,
- and  $c$  in some finite list

compute a recurrence  $L_{A,B,c}$  that has  $P(x)$  as a solution, and solve it to find hypergeometric solutions:

$$c^x P(x) \text{Sol}\left(\tau - \frac{A}{B}\right)$$

► Another algorithm was given by Cluzeau and v.Hoeij (2006).

## Petkovšek's algorithm extended to systems

- ▶ The system to solve looks like

$$\begin{pmatrix} y_1(x+1) \\ \vdots \\ y_n(x+1) \end{pmatrix} = M \begin{pmatrix} y_1(x) \\ \vdots \\ y_n(x) \end{pmatrix}$$

where  $M$  is a  $n$  by  $n$  invertible matrix over  $\mathbf{C}(x)$ .

- ▶ The solution we search for is hypergeometric, meaning that it can be written as

$$\begin{pmatrix} y_1(x) \\ \vdots \\ y_n(x) \end{pmatrix} = c^x \begin{pmatrix} P_1(x) \\ \vdots \\ P_n(x) \end{pmatrix} \text{Sol}\left(\tau - \frac{A}{B}\right)$$

where  $A, B, P_1, \dots, P_n \in \mathbf{C}[x]$  and  $c \in \mathbf{C}^*$ .

- ▶ The goal: Find a solution

$$Y = c^x \begin{pmatrix} P_1(x) \\ \vdots \\ P_n(x) \end{pmatrix} \text{Sol}\left(\tau - \frac{A}{B}\right)$$

of the system  $\tau(Y) = MY$ .

- ▶ Once you have  $A, B$  and  $c$ , then you can compute  $P_1, \dots, P_n$  using Abramov-Barkatou algorithm for polynomial solutions of difference systems (1998).

## How to find $A, B$ ?

$$Y = c^x \begin{pmatrix} P_1(x) \\ \vdots \\ P_n(x) \end{pmatrix} \text{Sol}\left(\tau - \frac{A}{B}\right)$$

We may assume that the gcd of  $P_1, \dots, P_n$  is 1.

Now look at

$$\tau(Y) = \frac{A}{B} c^{x+1} \begin{pmatrix} P_1(x+1) \\ \vdots \\ P_n(x+1) \end{pmatrix} \text{Sol}\left(\tau - \frac{A}{B}\right)$$

- ▶ The denominator  $B$  in  $\tau(Y)$  does not appear in  $Y$ , and since  $\tau(Y) = MY$  it follows that the denominator  $B$  must divide the denominator of  $M$ .
- ▶ A similar argument shows that  $A$  divides the denominator of  $M^{-1}$ .

So one can generalize Petkovšek's algorithm to systems:

- ▶  $B$  has to divide denominator( $M$ ),
- ▶ and  $A$  has to divide denominator( $M^{-1}$ ).

## How to find $c$ ?

- ▶ If the system  $\tau Y = MY$  has a hypergeometric solution

$$c^x \text{Sol}\left(\tau - \frac{A}{B}\right) P(x), \quad P(x) \in \mathbf{C}[x]^n$$

then it has a formal solution of the form

$$Y(x) = \Gamma(x)^s c^x x^d F(x)$$

where

$$(s, c, d) \in \mathbb{Z} \times \mathbf{C}^* \times \mathbf{C}$$

$F(x)$  is an  $n$ -dimensional vector with entries in  $\mathbf{C}[[x^{-1}]]$ :

$$F(x) = (F_0 + x^{-1}F_1 + \dots) \quad F_0 \neq 0.$$



In fact, if we write  $A, B$  as

$$A = x^\ell + \alpha_{\ell-1}x^{\ell-1} + \dots + \alpha_0$$

$$B = x^m + \beta_{m-1}x^{m-1} + \dots + \beta_0$$

then

$$s = \ell - m$$

and

$$d = \alpha_{\ell-1} - \beta_{m-1} + \deg P.$$

## Local Hypergeometric Solutions

Consider a difference system

$$\tau Y = MY.$$

We call *local hypergeometric solution* (lhs, in short) any nonzero formal solution of the form

$$Y(x) = \Gamma(x)^s c^x x^d F(x)$$

where  $s \in \mathbb{Z}$ ,  $c \in \mathbf{C}^*$ ,  $d \in \mathbf{C}$  and where  $F(x)$  is an  $n$ -dimensional vector with entries in  $\mathbf{C}[[x^{-1}]]$ :

$$F(x) = (F_0 + x^{-1}F_1 + \dots) \quad F_0 \neq 0.$$

If a formal solution of the form

$$Y(x) = \Gamma(x)^s c^x x^d F(x), \quad F(x) \in \mathbf{C}[[x]]^n.$$

exists then

- ▶  $s$  is a slope of the  $\tau$ -Newton polygon of the system  $\tau Y = MY$ ,
- ▶  $c$  is a root of the Newton polynomial associated with  $s$
- ▶ and  $d$  is then a root of the indicial equation associated with the slope zero in the  $\Delta$ -Newton polygon of the system

$$\Delta Z(x) := Z(x+1) - Z(x) = (x^{-s} c^{-1} M(x) - I)Z(x).$$

$$Y(x) = \Gamma(x)^s c^x x^d F(x), \quad F(x) \in \mathbf{C}[[x]]^n.$$

- ▶ The pairs  $(s, c)$  are invariants under *gauge transformations*:

$$Y(x) \mapsto T(x)Y(x) \quad T \in \mathrm{GL}(n, \mathbf{C}((x^{-1}))).$$

- ▶ However, the  $d$ 's may change into  $d + \nu$ , for some  $\nu \in \mathbb{Z}$ .

- ▶ We are interested in the computation of all the triplets  $(s, c, d) \in \mathbb{Z} \times \mathbf{C}^* \times \mathbf{C}$  that appear in the lhs of our system.
- ▶ In general, the triplets  $(s, c, d)$  cannot be obtained directly from the matrix system  $M(x)$ .
- ▶ However, we show that all the possible pairs  $(s, c)$  can be obtained from a *super-irreducible form* of the matrix  $\hat{M} := M - I_n$ .
- ▶ Once a pair  $(s, c)$  is obtained, the corresponding  $d$ 's can be computed from a super-irreducible form of the matrix

$$(x^{-s}c^{-1}M(x) - I_n).$$

## Super-irreducible Forms

The notion of super-reduced forms of linear difference systems had been developed in [Bark 1989].

### Notation

- ▶  $\Delta = \tau - 1$  denotes the difference operator defined by

$$\Delta f(x) = f(x + 1) - f(x)$$

for any function  $f$  of  $x$ .

- ▶ Any difference system

$$\tau Y = MY,$$

can be rewritten as

$$\Delta Y = \hat{M}Y$$

where

$$\hat{M} := M - I_n.$$

## Gauge transformations

Let  $T(x)$  be an invertible matrix.

The gauge transformation

$$Y = TZ$$

yields the following *equivalent* system

$$\tau Z = NZ \quad N = T_\tau[M] := \tau T^{-1}MT$$

and in the  $\Delta$ -form

$$\Delta Z = \hat{N}Z \quad \hat{N} = T_\Delta[\hat{M}] := \tau T^{-1}\hat{M}T - \tau T^{-1}\Delta T.$$

## Super-irreducible Forms

- ▶ Consider a difference system of the form :

$$\Delta Y(x) = A(x)Y(x),$$

with

$$A(x) = x^q \sum_{\nu=0}^{\infty} x^{-\nu} A_{\nu}, \quad q \in \mathbb{Z}, \quad A_0 \neq 0.$$

- ▶ For  $k = 1, \dots, q + 1$ , put

$$m_k(A) = \begin{cases} 0 & \text{if } q \leq -1 \\ q + 1 + \frac{n_0}{n} + \frac{n_1}{n^2} + \dots + \frac{n_{k-1}}{n^k} & \text{if } q > -1 \end{cases}$$

where  $n_i$  is the number of rows of  $A$  with valuation  $-q + i$ .



- ▶ Define

$$\mu_k(A) = \min_T m_k(T_\Delta[A]).$$

- ▶ The matrix  $A$  is said to be  $k$ -irreducible if

$$m_k(A) = \mu_k(A).$$

Otherwise  $A$  is called  $k$ -reducible.

- ▶  $A$  is  $k$ -reducible  $\iff$  there exists  $T$  s.t.  
 $m_k(T_\Delta[A]) < m_k(A)$ .
- ▶ The matrix  $A$  is said to be super-irreducible, if it is  $k$ -irreducible for every  $k$ , or equivalently if

$$m_{q+1}(A) = \mu_{q+1}(A).$$

## A Criterion for $k$ -reducibility

- ▶ For  $1 \leq k \leq q + 1$  define

$$\theta_k(\lambda) = x^{-r_k(A)} \det(x^{k-q}A(x) - \lambda I_n)|_{x=\infty}$$

where

$$r_k(A) = kn_0 + (k - 1)n_1 + \cdots + n_{k-1}.$$

- ▶ One can check that  $\theta_k(\lambda)$  belongs to  $\mathbf{C}[\lambda]$ .

**Theorem**[Bark 1989]

The matrix  $A$  is  $k$ -irreducible, if and only if the polynomials

$$\theta_1(\lambda), \theta_2(\lambda), \dots, \theta_k(\lambda)$$

do not vanish identically in  $\lambda$ .

**Theorem**[Bark 1989]

Given a difference system

$$\Delta Y = AY$$

one can construct a nonsingular matrix  $T$ , which is **polynomial** in  $x$ , such that the change of variable

$$Z = TY$$

leads to an equivalent system

$$\Delta Z = BZ$$

which is super-irreducible.

- ▶ This algorithm is implemented in Maple. Given a difference system it computes an equivalent super-reduced form as well as the corresponding polynomials  $\theta_j(\lambda)$ .

## How to compute the pairs $(s, c)$

We look for lhs of the form

$$Y(x) = \Gamma(x)^s c^x x^d (F_0 + F_1 x^{-1} + O(x^{-1})) \quad F_0 \neq 0$$

Write our system  $\tau Y = MY$  in the  $\Delta$ -form :

$$\Delta Y(x) = \widehat{M}(x) Y(x),$$

with

$$\widehat{M}(x) = x^q \sum_{\nu=0}^{\infty} x^{-\nu} \widehat{M}_{\nu} \quad \widehat{M}_0 \neq 0$$

We can always suppose that  $q \geq 0$ .

The possible values of  $s$  are then such that:  $0 \leq s \leq q$ .

We can suppose that our system is super-irreducible and denote by

$$\theta_1(\lambda), \theta_2(\lambda), \dots, \theta_{q+1}(\lambda)$$

the corresponding polynomials.

Let  $0 \leq s \leq q$  be fixed.

Rewrite the system as

$$\mathcal{L}(Y) := D(x)\Delta Y(x) - N(x)Y(x) = 0.$$

with

$$D(x) = \text{diag}(x^{\nu_1}, \dots, x^{\nu_n}) \quad \text{and} \quad N(x) = D(x)\widehat{M}(x).$$

where

$$\nu_i = \min(v(\widehat{M}_{i,\cdot}), -s)$$

$\widehat{M}_{i,\cdot}$  denoting the  $i$ th row of the matrix  $\widehat{M}$ .

The matrices  $D$  and  $N$  can be expanded as:

$$D(x) = x^{-s} (D_0 + O(x^{-1})) \quad \text{and} \quad N(x) = N_0 + O(x^{-1}).$$

Compute

$$\mathcal{L}(Y) = \Gamma(x)^s c^x x^d (((c - \varepsilon_s)D_0 - N_0)F_0 + O(x^{-1}))$$

where  $\varepsilon_s = 1$ , when  $s = 0$  and  $\varepsilon_s = 0$ , otherwise.

If the system has a local hypergeometric solution

$$Y(x) = \Gamma(x)^s c^x x^d (F_0 + F_1 x^{-1} + O(x^{-2})) \quad F_0 \neq 0$$

then one must have

$$((c - \varepsilon_s)D_0 - N_0)F_0 = 0.$$

- ▶ If a nonzero local hypergeometric solution

$$Y(x) = \Gamma(x)^s c^x x^d F(x)$$

exists then  $c$  must annihilate the polynomial

$$E_s(\lambda) := \det(N_0 - (\lambda - \varepsilon_s)D_0).$$

### Proposition

$$E_s(\lambda) = \theta_{q-s}(\lambda - \varepsilon_s)$$

where  $\varepsilon_s = 1$ , when  $s = 0$  and  $\varepsilon_s = 0$ , otherwise.

## Algorithm $\Delta$ -HS

**Input:** A field  $\mathbf{C}$  of characteristic 0, a variable  $x$ , and an  $n$  by  $n$  matrix  $M$  with entries in  $\mathbf{C}(x)$ .

**Output:** A basis  $B$  of the hypergeometric solutions that are defined over  $\mathbf{C}$ .

- ▶ Let  $d_M$  resp.  $d_{M^{-1}}$  be the denominator of  $M$  resp.  $M^{-1}$ .
- ▶ Compute the local data of  $\tau(Y) = MY$ .  
This produces a list  $L$  consisting of the triplets  $(s, c, d) \in \mathbb{Z} \times \mathbf{C}^* \times \mathbf{C}$  that occur in the local hypergeometric solutions of  $\tau(Y) = MY$ .
- ▶ Form the *AB-list*.  
This produces a set  $S = \{[A_1, B_1, s_1, c_1, d_1], \dots, [A_l, B_l, s_l, c_l, d_l]\}$  of pairs  $(A, B)$  of monic factors  $A|d_{M^{-1}}$  and  $B|d_M$  with  $A, B \in K[x]$  that match the local data in  $L$ .



## Algorithm $\Delta$ -HS





- ▶ Let  $B = \emptyset$ .
- ▶ For all  $i$  from 1 to  $l$  do
  - ▶  $N_i := B_i / (c_i A_i) \cdot M$ .
  - ▶ Compute a basis  $P_1, \dots, P_{m_i} \in K[x]^n$  of polynomial solutions for  $\tau(Y) = N_i Y$ .





If  $m_i > 0$  then for  $j = 1, \dots, m_i$  do:

- ▶ Let  $u = \text{Sol}(\tau - c_i \frac{A_i}{B_i}) P_j$ .
    - ▶  $B := B \cup \{u\}$ .
- ▶ Return  $B$  and stop.

- ▶ The algorithm we implemented in Maple includes efficiency improvements obtained by adapting ideas from [Cluzeau&Hoeij 2006] to the case of systems.
- ▶ We use modular computations (namely,  $p$ -curvature test) to reduce the number of pairs  $(A, B)$  to check.
- ▶ We also use a method to partially desingularize systems.
- ▶ Examples show that this provides significant efficiency benefits.



-  S. A. Abramov, M.A Barkatou (1998): Rational solutions of first order linear difference systems *Proc. of ISSAC'98*, 124–131.
-  Barkatou M.A. (1989): Contribution à l'étude des équations différentielles et aux différences dans le champ complexe, thèse soutenue le 6 juin 1989 à l'institut national polytechnique de Grenoble (France).
-  M. A. Barkatou and G. Chen (1996) Computing the exponential part of a formal fundamental matrix solution of a linear difference system, *J. Difference Eq. Appl.*, to appear.
-  M. A. Barkatou and G. Chen (1997): Some formal invariants of linear difference systems , *Pub. IRMA Lille*, Vol. 44, N° 2, 1997.
-  M. A. Barkatou, E. Pflügel (1998): On the Equivalence Problem of Linear Differential Systems and its Application for Factoring Completely Reducible Systems, *Proc. of ISSAC'98*, ACM Press, 268–275

-  E. Beke (1894): Die Irreduzibilität der homogenen linearen differentialgleichungen. In *Math. Ann.*, **45**, 278-294.
-  M. Bronstein M (1994): An improved algorithm for factoring linear ordinary differential operators. Proceedings ISSAC 94, Oxford, U.K., ACM Press, p. 336-340.
-  Hilali A. , Wazner A. (1987): Formes super-irréductibles des systèmes différentiels linéaires. *Num. Math.***50**, 429-449.
-  G.D. Birkhoff (1911): General Theory of irregular linear difference equations, *Trans. A.M.S.* 12, pp. 243-284.
-  G. D. Birkhoff (1930): Formal Theory of irregular linear difference equations, *Acta Math.* 54, pp. 205-246.
-  G. D. Birkhoff, W.J. Trjitzinsky (1933): Analytic Theory of singular difference equations, *Acta Math.* 60, pp. 1-89.
-  C. Praagman (1983): The formal classification of linear difference operators, *Ind. Math.* **45**, 249-261.



H. L. Turrittin: The formal theory of systems of irregular homogeneous linear difference equations, *Bollett della Società Matematica Italiana* **5** (1960), 255-264.



M. van Hoeij (1998): Rational solutions of linear difference equations *Proc. of ISSAC'98*, 120–123.



M. Petkovšek, H. Wilf, D. Zeilberger (1996): *A = B*, A K Peters, Wellesley, Massachusetts.



M. van der Put, M. F. Singer (1997): Galois Theory of Difference Equations, V. 1666 in *Lecture Notes in Mathematics*, Springer-Verlag