



HAL
open science

Markov operators in q -Fourier analysis

Lazhar Dhaouadi

► **To cite this version:**

| Lazhar Dhaouadi. Markov operators in q -Fourier analysis. 2008. hal-00322168v1

HAL Id: hal-00322168

<https://hal.science/hal-00322168v1>

Preprint submitted on 16 Sep 2008 (v1), last revised 31 Jan 2013 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Markov operators in q -Fourier analysis

Lazhar Dhaouadi*

Abstract

Essential result in q -Fourier analysis proved by a new way and a new set of markov operators acting on the $\mathcal{L}_{q,2,v}$ space of square q -integrable function are characterized.

1 Introduction

In the mathematical literature one finds much article which treats the theory of q -Fourier analysis associated with the q -Hankel transform. This theory finds its origins in an important article of H.T. Koornwinder and R.F. Swart-touw [5]. The two authors proved that the q -Bessel functions of Hahn-Exton satisfy an orthogonality relation, which it gives a rigorous proof. They noticed that the q -Hankel transform would be an involutive isometric operator of the $\mathcal{L}_{q,2,v}$ space. Our goal is to give a detailed proof of this important assertion. It should be noticed that in [1] we provide the essential results of q -Fourier analysis in particular that the q -Hankel transform is prolonged on the $\mathcal{L}_{q,2,v}$ space like an isometric involutive operator. But we used the positivity of the q -Bessel translation operator to lead to this results. However, this properties is not ensured for any q in the interval $]0, 1[$, from where the interest of this paper. Since, we will prove some essential results of q -Fourier analysis without recourse to the positivity of the q -Bessel translation operator, which are

- Inversion Formula in the $\mathcal{L}_{q,p,v}$ spaces with $p \geq 1$.
- Plancherel Formula in the $\mathcal{L}_{q,p,v} \cap \mathcal{L}_{q,1,v}$ spaces with $p > 2$.
- Plancherel Formula in the $\mathcal{L}_{q,2,v}$ spaces.
- Product formula for the q -Hankel transform .

Note that in a recent paper [2] we prove that the positivity of the q -Bessel translation operator is ensured in all point of the interval $]0, 1[$ when

*Ecole Préparatoire d'Ingénieur, Bizerte, Tunisia. E-mail lazhardhaouadi@yahoo.fr

$v \geq 0$. In this article we also try to show in a clear way the part which the positivity of the q -Bessel translation operator plays in q -Fourier analysis. In particular, when we tries to introduce the concept of Markov operators of the $\mathcal{L}_{q,2,v}$ space. An interesting examples of such operators are the q -Bessel translation operator and the q -heat semigroup.

2 Preliminaries

In the following we will always assume $0 < q < 1$ and $v > -1$. We denote by

$$\mathbb{R}_q = \{\pm q^n, n \in \mathbb{Z}\}, \quad \mathbb{R}_q^+ = \{q^n, n \in \mathbb{Z}\}.$$

For more information on the q -series theory the reader can see the reference [3,4,6] and the reference [1,5] about the q -bessel Fourier analysis. Also for details of the proof of the following result in this section can be fond in [1,5].

Definition 1 *The q -Bessel operator is defined as follows*

$$\Delta_{q,v} f(x) = \frac{1}{x^2} [f(q^{-1}x) - (1 + q^{2v})f(x) + q^{2v}f(qx)].$$

Definition 2 *The eigenfunction of $\Delta_{q,v}$ associated with the eigenvalue $-\lambda^2$ is the function $x \mapsto j_v(\lambda x, q^2)$, where $j_v(\cdot, q^2)$ is the normalized q -Bessel function defined by*

$$j_v(x, q^2) = \sum_{n=0}^{\infty} (-1)^n \frac{q^{n(n+1)}}{(q^{2v+2}, q^2)_n (q^2, q^2)_n} x^{2n}.$$

Definition 3 *The q -Jackson integral of a function f defined on \mathbb{R}_q is*

$$\int_0^{\infty} f(t) d_q t = (1 - q) \sum_{n \in \mathbb{Z}} q^n f(q^n).$$

Definition 4 *We denote by $\mathcal{L}_{q,p,v}$ the space of even functions f defined on \mathbb{R}_q such that*

$$\|f\|_{q,p,v} = \left[\int_0^{\infty} |f(x)|^p x^{2v+1} d_q x \right]^{1/p} < \infty.$$

Definition 5 We denote by $\mathcal{C}_{q,0}$ the space of even functions defined on \mathbb{R}_q tending to 0 as $x \rightarrow \pm\infty$ and continuous at 0 equipped with the topology of uniform convergence. The space $\mathcal{C}_{q,0}$ is complete with respect to the norm

$$\|f\|_{q,\infty} = \sup_{x \in \mathbb{R}_q} |f(x)|.$$

Proposition 1 The normalized q -Bessel function $j_v(\cdot, q^2)$ satisfies the orthogonality relation

$$c_{q,v}^2 \int_0^\infty j_v(xt, q^2) j_v(yt, q^2) t^{2v+1} d_q t = \delta_q(x, y), \quad \forall x, y \in \mathbb{R}_q.$$

where

$$\delta_q(x, y) = \begin{cases} 0 & \text{if } x \neq y \\ \frac{1}{(1-q)x^{2(v+1)}} & \text{if } x = y \end{cases}$$

and

$$c_{q,v} = \frac{1}{1-q} \frac{(q^{2v+2}, q^2)_\infty}{(q^2, q^2)_\infty}.$$

Remark 1 Let f be an even function defined on \mathbb{R}_q then

$$\int_0^\infty f(y) \delta_q(x, y) y^{2v+1} d_q y = f(x).$$

Proposition 2 The normalized q -Bessel function $j_v(\cdot, q^2)$ satisfies

$$|j_v(q^n, q^2)| \leq \frac{(-q^2; q^2)_\infty (-q^{2v+2}; q^2)_\infty}{(q^{2v+2}; q^2)_\infty} \begin{cases} 1 & \text{if } n \geq 0 \\ q^{n^2 - (2v+1)n} & \text{if } n < 0 \end{cases}.$$

Definition 6 The q -Bessel Fourier transform $\mathcal{F}_{q,v}$ is defined by

$$\mathcal{F}_{q,v} f(x) = c_{q,v} \int_0^\infty f(t) j_v(xt, q^2) t^{2v+1} d_q t, \quad \forall x \in \mathbb{R}_q.$$

Proposition 3 Let $f \in \mathcal{L}_{q,1,v}$ then $\mathcal{F}_{q,v} f$ exists and $\mathcal{F}_{q,v} f \in \mathcal{C}_{q,0}$.

3 Inversion formula in the $\mathcal{L}_{q,p,v}$ space

Theorem 1 Let f be a function in the $\mathcal{L}_{q,p,v}$ space where $p \geq 1$ then

$$\mathcal{F}_{q,v}^2 f = f.$$

Proof. If $f \in \mathcal{L}_{q,p,v}$ then $\mathcal{F}_{q,v}f$ exist, and we write

$$\begin{aligned}
\mathcal{F}_{q,v}^2 f(x) &= c_{q,v} \int_0^\infty \mathcal{F}_{q,v}f(t) j_v(xt, q^2) t^{2v+1} d_q t \\
&= \int_0^\infty f(y) \left[c_{q,v}^2 \int_0^\infty j_v(xt, q^2) j_v(yt, q^2) t^{2v+1} d_q t \right] y^{2v+1} d_q y \\
&= \int_0^\infty f(y) \delta_q(x, y) y^{2v+1} d_q y \\
&= f(x).
\end{aligned}$$

We can permute the two integrals: in fact if $p > 1$ then we use the Hölder's inequality

$$\begin{aligned}
&\int_0^\infty |f(y)| \left[\int_0^\infty |j_v(xt, q^2) j_v(yt, q^2)| t^{2v+1} d_q t \right] y^{2v+1} d_q y \\
&\leq \left(\int_0^\infty |f(y)|^p y^{2v+1} d_q y \right)^{1/p} \times \left(\int_0^\infty \sigma(y)^{\bar{p}} y^{2v+1} d_q y \right)^{1/\bar{p}}.
\end{aligned}$$

The numbers p and \bar{p} above are conjugates and

$$\sigma(y) = \int_0^\infty |j_v(xt, q^2) j_v(yt, q^2)| t^{2v+1} d_q t,$$

then

$$\begin{aligned}
&\int_0^\infty \sigma(y)^{\bar{p}} y^{2v+1} d_q y \\
&= \int_0^1 \sigma(y)^{\bar{p}} y^{2v+1} d_q y + \int_1^\infty \sigma(y)^{\bar{p}} y^{2v+1} d_q y.
\end{aligned}$$

Not that

$$\begin{aligned}
\int_0^1 \sigma(y)^{\bar{p}} y^{2v+1} d_q y &\leq \|j_v(\cdot, q^2)\|_{q,\infty}^{\bar{p}} \int_0^1 \left\{ \int_0^\infty |j_v(xt, q^2)| t^{2v+1} d_q t \right\}^{\bar{p}} y^{2v+1} d_q y \\
&\leq \|j_v(\cdot, q^2)\|_{q,\infty}^{\bar{p}} \|j_v(\cdot, q^2)\|_{q,1,v}^{\bar{p}} x^{-2(v+1)\bar{p}} \left[\int_0^1 y^{2v+1} d_q y \right] < \infty,
\end{aligned}$$

and

$$\begin{aligned}
&\int_1^\infty \sigma(y)^{\bar{p}} y^{2v+1} d_q y \\
&\leq \|j_v(\cdot, q^2)\|_{q,\infty}^{\bar{p}} \|j_v(\cdot, q^2)\|_{q,1,v}^{\bar{p}} \int_1^\infty \frac{y^{2v+1}}{y^{2(v+1)\bar{p}}} d_q y \\
&\leq \|j_v(\cdot, q^2)\|_{q,\infty}^{\bar{p}} \|j_v(\cdot, q^2)\|_{q,1,v}^{\bar{p}} \int_1^\infty \frac{1}{y^{2(v+1)(\bar{p}-1)+1}} d_q y < \infty.
\end{aligned}$$

If $p = 1$ then

$$\begin{aligned} & \int_0^\infty |f(y)| \left\{ \int_0^\infty |j_v(xt, q^2) j_v(yt, q^2)| t^{2v+1} d_q t \right\} y^{2v+1} d_q y \\ & \leq \|f\|_{q,1,v} \|j_v(\cdot, q^2)\|_{q,\infty} \|j_v(\cdot, q^2)\|_{q,1,v} \times \frac{1}{x^{2(v+1)}}, \end{aligned}$$

this finish the proof. ■

4 Plancherel Formula in the $\mathcal{L}_{q,1,v} \cap \mathcal{L}_{q,p,v}$ space

Theorem 2 Let f be a function in the $\mathcal{L}_{q,1,v} \cap \mathcal{L}_{q,p,v}$ space, where $p > 2$ then

$$\|\mathcal{F}_{q,v} f\|_{q,2,v} = \|f\|_{q,2,v}$$

Proof. Let $f \in \mathcal{L}_{q,1,v} \cap \mathcal{L}_{q,p,v}$ then by Theorem 1 we see that

$$\mathcal{F}_{q,v}^2 f = f,$$

this implies

$$\begin{aligned} \int_0^\infty \mathcal{F}_{q,v} f(x)^2 x^{2v+1} d_q x &= \int_0^\infty \mathcal{F}_{q,v} f(x) \left\{ c_{q,v} \int_0^\infty f(t) j_v(xt, q^2) t^{2v+1} d_q t \right\} x^{2v+1} d_q x \\ &= \int_0^\infty f(t) \left\{ c_{q,v} \int_0^\infty \mathcal{F}_{q,v} f(x) j_v(xt, q^2) x^{2v+1} d_q x \right\} t^{2v+1} d_q t \\ &= \int_0^\infty f(t)^2 t^{2v+1} d_q t. \end{aligned}$$

We can permute the two integrals because

$$\begin{aligned} & \int_0^\infty |f(t)| \left\{ c_{q,v} \int_0^\infty |\mathcal{F}_{q,v} f(x)| |j_v(xt, q^2)| x^{2v+1} d_q x \right\} t^{2v+1} d_q t \\ & \leq \left(\int_0^\infty |f(t)|^p t^{2v+1} d_q t \right)^{1/p} \times \left(\int_0^\infty |\phi(t)|^{\bar{p}} t^{2v+1} d_q t \right)^{1/\bar{p}}, \end{aligned}$$

where

$$\phi(t) = c_{q,v} \int_0^\infty |\mathcal{F}_{q,v} f(x)| |j_v(xt, q^2)| x^{2v+1} d_q x,$$

then

$$\begin{aligned}
|\mathcal{F}_{q,v}f(x)| &\leq c_{q,v} \int_0^\infty |f(y)| |j_v(xy, q^2)| y^{2v+1} d_q y \\
&\leq c_{q,v} \left(\int_0^\infty |f(y)|^p y^{2v+1} d_q y \right)^{1/p} \times \left(\int_0^\infty |j_v(xy, q^2)|^{\bar{p}} y^{2v+1} d_q y \right)^{1/\bar{p}} \\
&\leq c_{q,v} \left(\int_0^\infty |f(y)|^p y^{2v+1} d_q y \right)^{1/p} \times \left(\int_0^\infty |j_v(y, q^2)|^{\bar{p}} y^{2v+1} d_q y \right)^{1/\bar{p}} x^{-2(v+1)/\bar{p}} \\
&\leq c_{q,v} \|f\|_{q,p,v} \|j_v(\cdot, q^2)\|_{q,\bar{p},v} x^{-2(v+1)/\bar{p}}.
\end{aligned}$$

This gives

$$\begin{aligned}
\phi(t) &\leq c_{q,v}^2 \|f\|_{q,p,v} \|j_v(\cdot, q^2)\|_{q,\bar{p},v} \int_0^\infty |j_v(xt, q^2)| x^{(2v+1)-2(v+1)/\bar{p}} d_q x \\
&\leq c_{q,v}^2 \|f\|_{q,p,v} \|j_v(\cdot, q^2)\|_{q,\bar{p},v} \left[\int_0^\infty |j_v(x, q^2)| x^{2(v+1)/p-1} d_q x \right] t^{-2(v+1)/p} \\
&\leq C_1 t^{-2(v+1)/p},
\end{aligned}$$

and

$$\begin{aligned}
\phi(t) &= c_{q,v} \int_0^\infty |\mathcal{F}_{q,v}f(x)| |j_v(xt, q^2)| x^{2v+1} d_q x \\
&= \left[c_{q,v} \int_0^\infty |\mathcal{F}_{q,v}f(x/t)| |j_v(x, q^2)| x^{2v+1} d_q x \right] t^{-2(v+1)} \\
&\leq c_{q,v} \|\mathcal{F}_{q,v}f\|_{q,\infty} \times \|j_v(\cdot, q^2)\|_{q,1,v} \times t^{-2(v+1)} \\
&\leq C_2 t^{-2(v+1)}.
\end{aligned}$$

Hence,

$$\begin{aligned}
\int_0^\infty |\phi(t)|^{\bar{p}} t^{2v+1} d_q t &= \int_0^1 |\phi(t)|^{\bar{p}} t^{2v+1} d_q t + \int_1^\infty |\phi(t)|^{\bar{p}} t^{2v+1} d_q t \\
&\leq C_1 \int_0^1 t^{-2(v+1)\bar{p}/p} t^{2v+1} d_q t + C_2 \int_1^\infty t^{-2(v+1)\bar{p}} t^{2v+1} d_q t < \infty.
\end{aligned}$$

In fact

$$\begin{cases} -1 < -2(v+1)\frac{\bar{p}}{p} + 2v+1 \\ -2(v+1)\bar{p} + 2v+1 < -1 \end{cases} \Leftrightarrow \begin{cases} 0 < -2(v+1)(\bar{p}-2) \\ -2(v+1)(\bar{p}-1) < 0 \end{cases} \Leftrightarrow 1 < \bar{p} < 2 \Leftrightarrow p > 2,$$

this finish the proof. ■

5 Plancherel Formula in the $\mathcal{L}_{q,2,v}$ space

Theorem 3 *Let f be a function in the $\mathcal{L}_{q,2,v}$ space then*

$$\|\mathcal{F}_{q,v}f\|_{q,2,v} = \|f\|_{q,2,v}.$$

Proof. We introduce the function ψ_x as follows

$$\psi_x(t) = c_{q,v} j_v(tx, q^2)$$

The inner product \langle, \rangle in the Hilbert space $\mathcal{L}_{q,2,v}$ is defined by

$$f, g \in \mathcal{L}_{q,2,v} \Rightarrow \langle f, g \rangle = \int_0^\infty f(t)g(t)t^{2v+1}d_qt.$$

By Proposition 1 we write

$$x \neq y \Rightarrow \langle \psi_x, \psi_y \rangle = 0$$

$$\|\psi_x\|_{q,2,v}^2 = \frac{1}{1-q} x^{-2(v+1)}.$$

We have

$$\mathcal{F}_{q,v}f(x) = \langle f, \psi_x \rangle,$$

and by Theorem 1

$$f \in \mathcal{L}_{q,2,v} \Rightarrow \mathcal{F}_{q,v}^2 f = f,$$

then

$$\langle f, \psi_x \rangle = 0, \forall x \in \mathbb{R}_q^+ \Rightarrow \mathcal{F}_{q,v}f = 0 \Rightarrow f = 0.$$

Hence, $\{\psi_x, x \in \mathbb{R}_q^+\}$ form an orthogonal basis of the Hilbert space $\mathcal{L}_{q,2,v}$ and we have

$$\overline{\{\psi_x, x \in \mathbb{R}_q^+\}} = \mathcal{L}_{q,2,v} \quad (1)$$

Now

$$f \in \mathcal{L}_{q,2,v} \Rightarrow f = \sum_{x \in \mathbb{R}_q^+} \frac{1}{\|\psi_x\|_{q,2,v}^2} \langle f, \psi_x \rangle \psi_x,$$

and then

$$\|f\|_{q,2,v}^2 = \sum_{x \in \mathbb{R}_q^+} \frac{1}{\|\psi_x\|_{q,2,v}^2} \langle f, \psi_x \rangle^2 = (1-q) \sum_{x \in \mathbb{R}_q^+} x^{2(v+1)} \mathcal{F}_{q,v}f(x)^2 = \|\mathcal{F}_{q,v}f\|_{q,2,v}^2,$$

this finish the proof. ■

6 q -translation operator and q -convolution product

Definition 7 The q -translation operator is given as follows (see [1])

$$T_{q,x}^v f(y) = c_{q,v} \int_0^\infty \mathcal{F}_{q,v} f(t) j_v(yt, q^2) j_v(xt, q^2) t^{2v+1} d_q t.$$

Proposition 4 If $f \in \mathcal{L}_{q,1,v}$ then $T_{q,x}^v f$ exists and

$$T_{q,x}^v f(y) = \int_0^\infty f(z) D_v(x, y, z) z^{2v+1} d_q z,$$

where

$$D_v(x, y, z) = c_{q,v}^2 \int_0^\infty j_v(xs, q^2) j_v(ys, q^2) j_v(zs, q^2) s^{2v+1} d_q s.$$

Proof. In fact for all $x \in \mathbb{R}_q$

$$\begin{aligned} |T_{q,x}^v f(y)| &\leq \int_0^\infty |\mathcal{F}_{q,v} f(t)| |j_v(yt, q^2)| |j_v(xt, q^2)| t^{2v+1} d_q t \\ &\leq \|\mathcal{F}_{q,v} f\|_{q,\infty} \|j_v(\cdot, q^2)\|_{q,\infty} \|j_v(\cdot, q^2)\|_{q,1,v} \frac{1}{x^{2(v+1)}} < \infty. \end{aligned}$$

On the other hand

$$\begin{aligned} T_{q,x}^v f(y) &= c_{q,v} \int_0^\infty \mathcal{F}_{q,v} f(t) j_v(yt, q^2) j_v(xt, q^2) t^{2v+1} d_q t \\ &= c_{q,v} \int_0^\infty \left[c_{q,v} \int_0^\infty f(z) j_v(zt, q^2) z^{2v+1} d_q z \right] j_v(yt, q^2) j_v(xt, q^2) t^{2v+1} d_q t \\ &= \int_0^\infty f(z) D_v(x, y, z) z^{2v+1} d_q z. \end{aligned}$$

We can permute the two integrals because

$$\begin{aligned} &\int_0^\infty \left[\int_0^\infty |f(z)| |j_v(zt, q^2)| z^{2v+1} d_q z \right] |j_v(yt, q^2)| |j_v(xt, q^2)| t^{2v+1} d_q t \\ &\leq \|f\|_{q,1,v} \|j_v(\cdot, q^2)\|_{q,\infty}^2 \|j_v(\cdot, q^2)\|_{q,1,v} \times \frac{1}{x^{2(v+1)}} < \infty, \end{aligned}$$

which finish the proof. ■

Definition 8 The q -convolution product is defined by

$$f *_q g(x) = c_{q,v} \int_0^\infty T_{q,x}^v f(y) g(y) y^{2v+1} d_q y.$$

Proposition 5 Let $f, g \in \mathcal{L}_{q,1,v}$ then $f *_q g$ exists and we have

$$f *_q g = g *_q f.$$

Proof. Let $f, g \in \mathcal{L}_{q,1,v}$ then

$$\begin{aligned} f *_q g(x) &= c_{q,v} \int_0^\infty T_{q,x}^v f(y) g(y) y^{2v+1} d_q y \\ &= c_{q,v} \int_0^\infty \left[c_{q,v} \int_0^\infty \mathcal{F}_{q,v} f(t) j_v(yt, q^2) j_v(xt, q^2) t^{2v+1} d_q t \right] g(y) y^{2v+1} d_q y \\ &= c_{q,v} \int_0^\infty \mathcal{F}_{q,v} f(t) \left[c_{q,v} \int_0^\infty g(y) j_v(yt, q^2) y^{2v+1} d_q y \right] j_v(xt, q^2) t^{2v+1} d_q t \\ &= c_{q,v} \int_0^\infty \mathcal{F}_{q,v} f(t) \mathcal{F}_{q,v} g(t) j_v(xt, q^2) t^{2v+1} d_q t \\ &= g *_q f(x), \end{aligned}$$

The exchange of the signs integrals is valid sense because

$$\begin{aligned} &\int_0^\infty \int_0^\infty |\mathcal{F}_{q,v} f(t)| \times |j_v(yt, q^2)| \times |j_v(xt, q^2)| \times |g(y)| y^{2v+1} t^{2v+1} d_q t d_q y \\ &\leq \|\mathcal{F}_{q,v} f\|_{q,\infty} \|j_v(\cdot, q^2)\|_{q,\infty} \|g\|_{q,1,v} \|j_v(\cdot, q^2)\|_{q,1,v} \times x^{-(v+1)}, \end{aligned}$$

which prove the result. ■

Proposition 6 Let $f \in \mathcal{L}_{q,1,v}$ and $g \in \mathcal{L}_{q,2,v}$ then

$$\mathcal{F}_{q,v}(f *_q g) = \mathcal{F}_{q,v}(f) \times \mathcal{F}_{q,v}(g).$$

Proof. In fact we have

$$\mathcal{F}_{q,v} : \mathcal{L}_{q,2,v} \rightarrow \mathcal{L}_{q,2,v}$$

and

$$\mathcal{F}_{q,v} : \mathcal{L}_{q,1,v} \rightarrow \mathcal{C}_{q,0}.$$

If $f \in \mathcal{L}_{q,1,v}$ and $g \in \mathcal{L}_{q,2,v}$ then

$$\omega = \mathcal{F}_{q,v} f \times \mathcal{F}_{q,v} g \in \mathcal{L}_{q,2,v}$$

On the other hand

$$\mathcal{F}_{q,v}\omega(x) = c_{q,v} \int_0^\infty \mathcal{F}_{q,v}f(t)\mathcal{F}_{q,v}g(t)j_v(xt, q^2)t^{2v+1}d_qt.$$

From the inversion formula in Theorem 1 we have

$$\mathcal{F}_{q,v}(\mathcal{F}_{q,v}\omega) = \mathcal{F}_{q,v}f \times \mathcal{F}_{q,v}g.$$

Now we write

$$\begin{aligned} \mathcal{F}_{q,v}\omega(x) &= c_{q,v} \int_0^\infty \mathcal{F}_{q,v}f(t)\mathcal{F}_{q,v}g(t)j_v(xt, q^2)t^{2v+1}d_qt \\ &= c_{q,v} \int_0^\infty \mathcal{F}_{q,v}f(t) \left[c_{q,v} \int_0^\infty g(y)j_v(ty, q^2)y^{2v+1}d_qy \right] j_v(xt, q^2)t^{2v+1}d_qt \\ &= c_{q,v} \int_0^\infty g(y) \left[c_{q,v} \int_0^\infty \mathcal{F}_{q,v}f(t)j_v(ty, q^2)j_v(xt, q^2)t^{2v+1}d_qt \right] y^{2v+1}d_qy \\ &= c_{q,v} \int_0^\infty g(y)T_{q,x}^v f(y)y^{2v+1}d_qy \\ &= f *_q g(x). \end{aligned}$$

This implies that

$$\mathcal{F}_{q,v}(f *_q g) = \mathcal{F}_{q,v}f \times \mathcal{F}_{q,v}g.$$

The exchange of the signs integrals is valid sense, indeed let

$$\begin{aligned} \phi(y) &= \int_0^\infty |\mathcal{F}_{q,v}f(t)| \times |j_v(xt, q^2)| \times |j_v(yt, q^2)| t^{2v+1}d_qt \\ &= \int_0^\infty \varrho(t) \times |j_v(yt, q^2)| t^{2v+1}d_qt \end{aligned}$$

where

$$\varrho(t) = |\mathcal{F}_{q,v}f(t)| \times |j_v(xt, q^2)| \Rightarrow \varrho \in \mathcal{L}_{q,1,v}$$

then

$$\lim_{y \downarrow 0} \phi(y) < \infty,$$

and

$$\begin{aligned} \phi(y) &= \int_0^\infty \varrho(t) \times |j_v(yt, q^2)| t^{2v+1}d_qt \\ &= \left[\int_0^\infty \varrho\left(\frac{t}{y}\right) \times |j_v(t, q^2)| t^{2v+1}d_qt \right] y^{-2(v+1)} \\ &\leq \left[\|\mathcal{F}_{q,v}f\|_{q,\infty} \|j_v(\cdot, q^2)\|_{q,\infty} \|j_v(\cdot, q^2)\|_{q,1,\infty} \right] y^{-2(v+1)}, \end{aligned}$$

which prove that $\phi \in \mathcal{L}_{q,2,v}$. This leads to the result. ■

7 Positivity of the q -translation operator

Definition 9 The operator $T_{q,x}^v$ is said positive if $T_{q,x}^v f \geq 0$ when $f \geq 0$ for all $x \in \mathbb{R}_q$. We denote by Q_v the domain of positivity of $T_{q,x}^v$

$$Q_v = \{q \in]0, 1[, \quad T_{q,x}^v \text{ is positive}\}.$$

In a recent paper [2] the authors prove that if $v \geq 0$ then $Q_v =]0, 1[$. In the following we assume that $q \in Q_v$.

Proposition 7 The operator $T_{q,x}^v$ is positive if and only if

$$D_v(x, y, z) \geq 0, \quad \forall x, y, z \in \mathbb{R}_q.$$

Proof. If we take $f(z) = \delta_a(z)$ then

$$T_{q,x}^v f(y) = (1 - q)a^{2(v+1)} D_v(x, y, a),$$

which prove the result. ■

Proposition 8 If $f \in \mathcal{L}_{q,1,v}$ then

$$\int_0^\infty T_{q,x}^v f(y) y^{2v+1} d_q y = \int_0^\infty f(y) y^{2v+1} d_q y.$$

Proof. In fact

$$\begin{aligned} \int_0^\infty T_{q,x}^v f(y) y^{2v+1} d_q y &= \int_0^\infty \left[\int_0^\infty f(z) D_v(x, y, z) z^{2v+1} d_q z \right] y^{2v+1} d_q y \\ &= \int_0^\infty \left[\int_0^\infty D_v(x, y, z) y^{2v+1} d_q y \right] f(z) z^{2v+1} d_q z. \end{aligned}$$

On the other hand

$$D_v(x, y, z) = \mathcal{F}_{q,v} \phi(y)$$

where

$$\phi(t) = c_{q,v} j_v(tx, q^2) j_v(tz, q^2),$$

then

$$\begin{aligned} \int_0^\infty D_v(x, y, z) y^{2v+1} d_q y &= \int_0^\infty \mathcal{F}_{q,v} \phi(y) y^{2v+1} d_q y \\ &= \frac{1}{c_{q,v}} \mathcal{F}_{q,v}^2 \phi(0) = \frac{1}{c_{q,v}} \phi(0) = 1. \end{aligned}$$

This leads to the result. ■

Proposition 9 Let $f, g \in \mathcal{L}_{q,1,v}$ then

$$\mathcal{F}_{q,v}(f *_q g) = \mathcal{F}_{q,v}(f) \times \mathcal{F}_{q,v}(g).$$

Proof. In fact

$$\begin{aligned} & \mathcal{F}_{q,v}(f *_q g)(t) \\ &= c_{q,v} \int_0^\infty f *_q g(x) j_v(xt, q^2) x^{2v+1} d_q x \\ &= c_{q,v} \int_0^\infty \left[\int_0^\infty T_{q,x}^v f(y) g(y) y^{2v+1} d_q y \right] j_v(xt, q^2) x^{2v+1} d_q x \\ &= c_{q,v} \int_0^\infty \left(\int_0^\infty g(y) \left[\int_0^\infty f(z) D_v(x, y, z) z^{2v+1} d_q z \right] y^{2v+1} d_q y \right) j_v(xt, q^2) x^{2v+1} d_q x \\ &= c_{q,v} \int_0^\infty f(z) \left(\int_0^\infty g(y) \left[\int_0^\infty D_v(x, y, z) j_v(xt, q^2) x^{2v+1} d_q x \right] y^{2v+1} d_q y \right) z^{2v+1} d_q z \\ &= c_{q,v}^2 \int_0^\infty f(z) \left(\int_0^\infty g(y) j_v(yt, q^2) j_v(xt, q^2) y^{2v+1} d_q y \right) z^{2v+1} d_q z \\ &= \mathcal{F}_{q,v}(f)(t) \times \mathcal{F}_{q,v}(g)(t). \end{aligned}$$

One can see that

$$\int_0^\infty D_v(x, y, z) j_v(xt, q^2) x^{2v+1} d_q x = c_{q,v} j_v(zt, q^2) j_v(yt, q^2),$$

and

$$\int_0^\infty D_v(x, y, z) x^{2v+1} d_q x = 1.$$

We can permute the two signs integrals because

$$\begin{aligned} & \int_0^\infty \left(\int_0^\infty |g(y)| \left[\int_0^\infty |f(z)| |D_v(x, y, z)| z^{2v+1} d_q z \right] y^{2v+1} d_q y \right) |j_v(xt, q^2)| x^{2v+1} d_q x \\ & \leq \|j_v(\cdot, q^2)\|_{q,\infty} \int_0^\infty |f(z)| \left(\int_0^\infty |g(y)| \left[\int_0^\infty |D_v(x, y, z)| x^{2v+1} d_q x \right] y^{2v+1} d_q y \right) z^{2v+1} d_q z \\ & \leq \|j_v(\cdot, q^2)\|_{q,\infty} \int_0^\infty |f(z)| \left(\int_0^\infty |g(y)| \left[\int_0^\infty D_v(x, y, z) x^{2v+1} d_q x \right] y^{2v+1} d_q y \right) z^{2v+1} d_q z \\ & \leq \|j_v(\cdot, q^2)\|_{q,\infty} \|f\|_{q,1,v} \|g\|_{q,1,v} < \infty. \end{aligned}$$

Note that if $q \in Q_v$ then $|D_v(x, y, z)| = D_v(x, y, z)$. ■

8 Markov operators on the $\mathcal{L}_{q,2,v}$ space

Definition 10 A linear bonded mapping $T : \mathcal{L}_{q,2,v} \rightarrow \mathcal{L}_{q,2,v}$ is called a Markov operator if

- T is positive, that is $f \geq 0$ implies $Tf \geq 0$.
- The constant function $f = 1$ is a fixed point of T .
- For all function $f, g \in \mathcal{L}_{q,2,v}$ we have

$$\langle Tf, g \rangle = \langle f, Tg \rangle.$$

Lemma 1 Let μ be a probability measure on \mathbb{R}_q^+ . If f is a positive function in $\mathcal{L}_{q,v,1}$ and g is convex on \mathbb{R}^+ then

$$g\left(\int_0^\infty f d\mu(x)\right) \leq \int_0^\infty g \circ f(x) d\mu(x).$$

In particular

$$\left[\int_0^\infty f(z) D_v(x, y, z) z^{2v+1} d_q z\right]^2 \leq \int_0^\infty f(z)^2 D_v(x, y, z) z^{2v+1} d_q z.$$

Proof. There exists $c \in \mathbb{R}$ such that for all x, y in a convex set of \mathbb{R}^+ it holds

$$g(x) \geq g(y) + c(x - y).$$

Let $t = \int_0^\infty f(x) d\mu(x)$. Then $t \in \mathbb{R}^+$. Now let $x \in \mathbb{R}_q^+$ then

$$g(f(x)) \geq g(t) + c(f(x) - t).$$

Integrating both sides and using the special value of t gives

$$\int_0^\infty g(f(x)) d\mu(x) \geq \int_0^\infty [g(t) + c(f(x) - t)] d\mu(x) = g(t).$$

In particular for $g(x) = x^2$ and $d\mu(z) = D_v(x, y, z) z^{2v+1} d_q z$ which is a probability measure on \mathbb{R}_q^+ the result follows immediately. ■

Proposition 10 $T_{q,x}^v$ is a Markov operator of $\mathcal{L}_{q,2,v}$.

Proof. We will prove that $T_{q,x}^v$ sends $\mathcal{L}_{q,2,v}$ to $\mathcal{L}_{q,2,v}$. Let $f \in \mathcal{L}_{q,2,v}$ then

$$\begin{aligned} [T_{q,x}^v f(y)]^2 &= \left[\int_0^\infty f(z) D_v(x, y, z) z^{2v+1} d_q z \right]^2 \\ &\leq \int_0^\infty f(z)^2 D_v(x, y, z) z^{2v+1} d_q z = T_{q,x}^v f^2(y). \end{aligned}$$

$$\int_0^\infty [T_{q,x}^v f(y)]^2 y^{2v+1} d_q y \leq \int_0^\infty T_{q,x}^v f^2(y) y^{2v+1} d_q y.$$

$$f \in \mathcal{L}_{q,2,v} \Rightarrow f^2 \in \mathcal{L}_{q,1,v}$$

Hence

$$\int_0^\infty T_{q,x}^v f^2(y) y^{2v+1} d_q y = \int_0^\infty f^2(y) y^{2v+1} d_q y$$

which gives

$$\|T_{q,x}^v f\|_{q,2,v} \leq \|f\|_{q,2,v}.$$

The operator $T_{q,x}^v$ is said to be a contraction of the $\mathcal{L}_{q,2,v}$ space. Now we will prove that $T_{q,x}^v$ is symmetric. Let $f, g \in \mathcal{L}_{q,2,v}$ then

$$|\langle T_{q,x}^v f, g \rangle| = \int_0^\infty |T_{q,x}^v f(y)| |g(y)| y^{2v+1} d_q y \leq \|T_{q,x}^v f\|_{q,2,v} \|g\|_{q,2,v} \leq \|f\|_{q,2,v} \|g\|_{q,2,v}$$

$$\begin{aligned} \langle T_{q,x}^v f, g \rangle &= \int_0^\infty T_{q,x}^v f(y) g(y) y^{2v+1} d_q y = \int_0^\infty \left[\int_0^\infty f(z) D_v(x, y, z) z^{2v+1} d_q z \right] g(y) y^{2v+1} d_q y \\ &= \int_0^\infty \left[\int_0^\infty g(z) D_v(x, y, z) z^{2v+1} d_q z \right] f(y) y^{2v+1} d_q y \\ &= \langle f, T_{q,x}^v g \rangle. \end{aligned}$$

The exchange of the signs integrals is valid sense

$$\begin{aligned} &\int_0^\infty \left[\int_0^\infty |g(z)| D_v(x, y, z) z^{2v+1} d_q z \right] |f(y)| y^{2v+1} d_q y \\ &\leq \left(\int_0^\infty \left[\int_0^\infty |g(z)| D_v(x, y, z) z^{2v+1} d_q z \right]^2 y^{2v+1} d_q y \right)^{1/2} \left(\int_0^\infty |f(y)|^2 y^{2v+1} d_q y \right)^{1/2} \\ &\leq \left(\int_0^\infty \left[\int_0^\infty |g(z)|^2 D_v(x, y, z) z^{2v+1} d_q z \right] y^{2v+1} d_q y \right)^{1/2} \|f\|_{q,2,v} = \|g\|_{q,2,v} \|f\|_{q,2,v}. \end{aligned}$$

On the other hand

$$T_{q,x}^v 1 = \int_0^\infty D_v(x, y, z) z^{2v+1} d_q z = 1.$$

Also

$$f \geq 0 \Rightarrow T_{q,x}^v f \geq 0.$$

Not that if f is bonded then

$$\begin{aligned} |T_{q,x}^v f(x)| &\leq \int_0^\infty |f(z)| |D_v(x, y, z)| z^{2v+1} d_q z \\ &\leq \|f\|_{q,\infty} \left[\int_0^\infty D_v(x, y, z) z^{2v+1} d_q z \right] = \|f\|_{q,\infty}. \end{aligned}$$

which finish the proof. ■

Theorem 4 Let $c_{q,v} \varrho(t) t^{2v+1} d_q t$ be a probability measure on \mathbb{R}_q^+ . Then

$$K : \mathcal{L}_{q,2,v} \rightarrow \mathcal{L}_{q,2,v}, \quad f \mapsto f *_q \varrho$$

is a Markov operator.

Proof. From Proposition 6 we see that K send $\mathcal{L}_{q,2,v}$ to $\mathcal{L}_{q,2,v}$. On the other hand

$$Kf(x) = f *_q \varrho(x) = c_{q,v} \int_0^\infty T_{q,x} f(y) \varrho(y) y^{2v+1} d_q y,$$

which implies that $K1 = 1$ and if $f \geq 0$ then $Kf \geq 0$. Given two functions $f, g \in \mathcal{L}_{q,2,v}$ we write

$$\begin{aligned} \langle Kf, g \rangle &= \int_0^\infty Kf(x) g(x) x^{2v+1} d_q x \\ &= \int_0^\infty \left[c_{q,v} \int_0^\infty T_{q,x} f(y) \varrho(y) y^{2v+1} d_q y \right] g(x) x^{2v+1} d_q x \\ &= \int_0^\infty \left[c_{q,v} \int_0^\infty T_{q,y} f(x) g(x) x^{2v+1} d_q x \right] \varrho(y) y^{2v+1} d_q y \\ &= \int_0^\infty \left[c_{q,v} \int_0^\infty f(x) T_{q,y} g(x) x^{2v+1} d_q x \right] \varrho(y) y^{2v+1} d_q y \\ &= \int_0^\infty \left[c_{q,v} \int_0^\infty T_{q,x} g(y) \varrho(y) y^{2v+1} d_q y \right] f(x) x^{2v+1} d_q x \\ &= \langle f, Kg \rangle. \end{aligned}$$

The exchange of the signs integral is valid sense

$$\begin{aligned} &\int_0^\infty \left[\int_0^\infty |T_{q,y} f(x)| |g(x)| x^{2v+1} d_q x \right] \varrho(y) y^{2v+1} d_q y \\ &\leq \|f\|_{q,2,v} \|g\|_{q,2,v} \|\varrho\|_{q,1,v}. \end{aligned}$$

Not that if f is bonded then

$$|Kf(x)| \leq \left[c_{q,v} \int_0^\infty \varrho(y) y^{2v+1} d_q y \right] \|f\|_{q,\infty} = \|f\|_{q,\infty}$$

which finish the proof. ■

Proposition 11 *The eigenfunctions of the operator $T_{q,x}^v$ are in the following form*

$$f_n(t) = \frac{\psi_{q^n}(t)}{\|\psi_{q^n}\|_{q,2,v}}, \quad \forall n \in \mathbb{Z}$$

associated to the eigenvalues

$$\lambda_n = \frac{f_n(x)}{f_n(0)}, \quad \forall n \in \mathbb{Z}.$$

Proof. From (1), the sequence $\{f_n\}_{n \in \mathbb{Z}}$ forme a normal orthogonal basis of the Hilbert space $\mathcal{L}_{q,2,v}$. On the other hand

$$T_{q,x}^v j_v(q^n y, q^2) = j_v(q^n x, q^2) j_v(q^n y, q^2),$$

then

$$T_{q,x}^v f_n(y) = \frac{f_n(x)}{f_n(0)} f_n(y),$$

which leads to the result. ■

Remark 2 *Note that the kernel of $T_{q,x}^v$ can be written as follows*

$$D_v(x, y, z) = \sum_{n \in \mathbb{Z}} \frac{f_n(x) f_n(y) f_n(z)}{f_n(0)} \geq 0,$$

then the sequence $\{f_n\}$ satisfies the hypergroup property at the point 0 with measure $x^{2v+1} d_q x$.

Proposition 12 *A linear mapping*

$$K : \mathcal{L}_{q,2,v} \rightarrow \mathcal{L}_{q,2,v}, \quad f_n \mapsto c_n f_n$$

is a Markov operator if and only if

$$c_n = c_{q,v} \int_0^\infty \frac{f_n(y)}{f_n(0)} \varrho(y) y^{2v+1} d_q y$$

where $c_{q,v} \varrho(y) y^{2v+1} d_q y$ is a probability measure on \mathbb{R}_q^+ .

Proof. In fact

$$\begin{aligned} f_n *_q \varrho(x) &= c_{q,v} \int_0^\infty T_{q,x} f_n(y) \varrho(y) y^{2v+1} d_q y \\ &= \left[c_{q,v} \int_0^\infty \frac{f_n(y)}{f_n(0)} \varrho(y) y^{2v+1} d_q y \right] f_n(x) = c_n f_n(x), \end{aligned}$$

Then

$$K(f) = f *_q \varrho, \quad \forall f \in \mathcal{L}_{q,2,v}.$$

Theorem 4 leads to the result. ■

9 The q -heat semigroup

Definition 11 *The q -exponential function is defined by*

$$e(z, q) = \sum_{n=0}^{\infty} \frac{z^n}{(q, q)_n} = \frac{1}{(z; q)_\infty}, \quad |z| < 1.$$

Lemma 2 *The unique solution of the following q -difference equation*

$$\begin{cases} -x^2 \psi(t) = (1 - q^2) D_{q^2, t} \psi(t) \\ \psi(0) = c(x) \end{cases}$$

is in the following form

$$\psi(t) = c(x) e(-tx^2, q^2).$$

Lemma 3 *Let $f \in \mathcal{L}_{q,2,v}$ such that $\Delta_{q,v} f \in \mathcal{L}_{q,2,v}$ then*

$$\mathcal{F}_{q,v} [\Delta_{q,v} f](x) = -x^2 \mathcal{F}_{q,v} f(x).$$

Proof. Let $g = \mathcal{F}_{q,v} f$. From the inversion formula we obtain

$$f = \mathcal{F}_{q,v}^2 f = \mathcal{F}_{q,v} g.$$

Then

$$\Delta_{q,v} f(x) = \mathcal{F}_{q,v} \left[y \mapsto -y^2 g(y) \right](x)$$

Again, we use the inversion formula we obtain the result. ■

Definition 12 The q -heat semigroup is introduced in [1] as follows:

$$P_{q,t}^v f(x) = G^v(., t, q^2) *_q f(x), \quad t > 0$$

where $G^v(., t, q^2)$ is the q -Gauss kernel

$$G^v(x, t, q^2) = \mathcal{F}_{q,v} \left[y \mapsto e(-ty^2, q^2) \right] (x) = A(t) e \left(-\frac{q^{-2v}}{t} x^2, q^2 \right),$$

and

$$A(t) = \frac{(-q^{2v+2}t, -q^{-2v}/t; q^2)_\infty}{(-t, -q^2/t; q^2)_\infty}.$$

Note that $A(t^2) = t^{-2(v+1)} A(1)$.

Proposition 13 The unique solution of the q -heat equation

$$\begin{cases} \Delta_{q,v} u(x, t) = (1 - q^2) D_{q^2, t} u(x, t), & \forall x \in \mathbb{R}_q, \quad \forall t > 0 \\ u(x, 0) = f(x), & f \in \mathcal{L}_{q,2,v} \\ x \mapsto u(x, t) \in \mathcal{L}_{q,2,v}, & \forall t > 0. \end{cases}$$

is in the following form

$$u(x, t) = P_{q,t}^v f(x).$$

Proof. Let

$$\psi(t) = \mathcal{F}_{q,v} \left[u(., t) \right] (x).$$

From the following equation

$$\Delta_{q,v} u(x, t) = (1 - q^2) D_{q^2, t} u(x, t)$$

we see that

$$\Delta_{q,v} u(., t) \in \mathcal{L}_{q,2,v},$$

then from Lemma 3 we have

$$\mathcal{F}_{q,v} \left[\Delta_{q,v} u(., t) \right] (x) = -x^2 \mathcal{F}_{q,v} \left[u(., t) \right] (x)$$

and

$$\mathcal{F}_{q,v} \left[(1 - q^2) D_{q^2, t} u(., t) \right] (x) = (1 - q^2) D_{q^2, t} \mathcal{F}_{q,v} \left[u(., t) \right] (x),$$

which prove that ψ satisfies

$$(1 - q^2) D_{q^2, t} \psi(t) = -x^2 \psi(t), \quad \psi(0) = \mathcal{F}_{q,v} f(x).$$

From Lemma 1 and Proposition 5 we see that

$$\psi(t) = e(-tx^2, q^2) \mathcal{F}_{q,v} f(x) = \mathcal{F}_{q,v} \left[G^v(\cdot, t, q^2) \right] (x) \times \mathcal{F}_{q,v} f(x) = \mathcal{F}_{q,v} \left[G^v(\cdot, t, q^2) *_q f \right] (x).$$

From Theorem 1 we obtain

$$u(x, t) = G^v(\cdot, t, q^2) *_q f(x).$$

This finish the proof. ■

Proposition 14 *The q -heat semigroup $P_{q,t}^v$ is a Markov operator of $\mathcal{L}_{q,2,v}$.*

Proof. In [1] it is proved that

$$\|G^v(\cdot, t, q^2)\|_{q,1,v} = \frac{1}{c_{q,v}},$$

then

$$y \mapsto c_{q,v} G^v(y, t, q^2) y^{2v+1} d_q y$$

is a probability measure on \mathbb{R}_q^+ . Theorem 4 leads to the result. ■

References

- [1] L. Dhaouadi, A. Fitouhi and J. El Kamel, Inequalities in q -Fourier Analysis, Journal of Inequalities in Pure and Applied Mathematics, Volume 7, Issue 5, Article 171, 2006.
- [2] L. Dhaouadi, A. Fitouhi, On the Grafts addition theorem for Hahn Exton q -Bessel function, arXiv:0707.1745v2 [math.CA].
- [3] G. Gasper and M. Rahman, Basic hypergeometric series, Encyclopedia of mathematics and its applications 35, Cambridge university press, 1990.
- [4] F. H. Jackson, On a q -Definite Integrals, Quarterly Journal of Pure and Application Mathematics 41, 1910, 193-203.
- [5] T. H. Koornwinder and R. F. Swarttouw, On q -Analogues of the Hankel and Fourier Transform, Trans. A. M. S. 1992, 333, 445-461.
- [6] R. F. Swarttouw, The Hahn-Exton q -Bessel functions PhD Thesis The Technical University of Delft (1992).