



HAL
open science

Multiplicity results for the assigned Gauss curvature problem in \mathbb{R}^2

Jean Dolbeault, Maria J. Esteban, Gabriella Tarantello

► **To cite this version:**

Jean Dolbeault, Maria J. Esteban, Gabriella Tarantello. Multiplicity results for the assigned Gauss curvature problem in \mathbb{R}^2 . *Nonlinear Analysis: Theory, Methods and Applications*, 2009, 70 (8), pp.2870-2881. hal-00323700

HAL Id: hal-00323700

<https://hal.science/hal-00323700v1>

Submitted on 22 Sep 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Multiplicity results for the assigned Gauss curvature problem in \mathbb{R}^2

Jean Dolbeault^a Maria J. Esteban^a Gabriella Tarantello^b

^a*Ceremade (UMR CNRS no. 7534), Université Paris-Dauphine, Place de Lattre de Tassigny, 75775 Paris Cedex 16, France.*

^b*Dipartimento di Matematica. Università di Roma "Tor Vergata", Via della Ricerca Scientifica, 00133 Roma, Italy.*

Abstract

To study the problem of the assigned Gauss curvature with conical singularities on Riemannian manifolds, we consider the Liouville equation with a single Dirac measure on the two-dimensional sphere. By a stereographic projection, we reduce the problem to a Liouville equation on the euclidean plane. We prove new multiplicity results for bounded radial solutions, which improve on earlier results of C.-S. Lin and his collaborators. Based on numerical computations, we also present various conjectures on the number of unbounded solutions. Using symmetries, some multiplicity results for non radial solutions are also stated.

Key words: Gauss curvature, conical singularities, Riemannian manifolds, Liouville equation, Onsager equation, self-dual gauge field vortices, stereographic projection, Emden-Fowler transformation, blow-up, radial symmetry, uniqueness, multiplicity
MSC (2000): 35J60; 34L30, 53C21, 58J05, 58J70

1 Introduction

In recent years much attention has been devoted to the study of mean field equations of Liouville type on Riemann surfaces and in the presence of singular sources. Such an interest has originated from various areas of mathematics

Email addresses: dolbeaul@ceremade.dauphine.fr (Jean Dolbeault),
esteban@ceremade.dauphine.fr (Maria J. Esteban),
tarantel@mat.uniroma2.it (Gabriella Tarantello).

URLs: www.ceremade.dauphine.fr/~dolbeaul (Jean Dolbeault),
www.ceremade.dauphine.fr/~esteban (Maria J. Esteban).

and physics, starting with the *assigned Gauss curvature* problem which can be reduced to analyze the solution set of

$$\Delta u + K(x) e^{2u} = 0 \quad \text{in } \mathbb{R}^2 \quad (1)$$

for a given function K defined in \mathbb{R}^2 . If u is a solution to (1), then the metric $g = e^{2u} |dx|^2$ is conformal to the flat metric $|dx|^2$ and such that K is the Gaussian curvature of the new metric g . Equation (1) also appears in the analysis of gravitating systems, in the statistical mechanics description of the vorticity in fluid mechanics (see [2,3,4,20,21]) and has been studied more recently in the context of self-dual gauge field vortices (see [16,35,32]).

The solution set of equation (1) depends very much on the properties of K . When K is negative, uniqueness results are always available, while for total positive curvature K , either uniqueness or multiplicity of solutions holds, depending on K . See [24] for various examples. In this paper we will focus on particular cases of positive functions K .

We are also interested in the problem of the *assigned Gauss curvature with conical singularities* (see [34,33,27,14,10]). For a given Riemann surface (M, g) , we aim at determining the range of the parameters $\lambda, \rho \in \mathbb{R}$ such that

$$\Delta_g u + \lambda \left(\frac{e^{2u}}{\int_M e^{2u} d\sigma_g} - \frac{1}{|M|} \right) - 2\pi\rho \left(\delta_P - \frac{1}{|M|} \right) = f, \quad (2)$$

is solvable on M , where Δ_g is the Laplace-Beltrami operator, $d\sigma_g$ is the volume element corresponding to the metric g , $f \in C(M)$ with $\int_M f d\sigma_g = 0$ and δ_P is the Dirac measure with singularity at $P \in M$. In case M has a non-empty boundary, both Dirichlet or Neumann boundary conditions on ∂M are of interest for the applications. One could also consider sums of Dirac measures located at several source points.

In the applications, the simplest situations correspond to the 2-sphere $M = \mathbb{S}^2$ and the flat 2-torus $M = \mathbb{C}/(\xi_1\mathbb{Z} + \xi_2\mathbb{Z})$, with periodic cell domain generated by ξ_1 and ξ_2 . Recall that the 2-sphere, with the standard metric induced by Lebesgue's measure in \mathbb{R}^3 , has already played a special role in the assigned curvature problem (see [17,19,18]), while the torus is important since many vortex-like configurations naturally develop into periodic lattices.

For the sphere, $\rho = 0$ corresponds to a particular case of the so-called Onsager vortex problem (see [24,25]). In fact, for closed surfaces and when there is no singularity (*i.e.* $\partial M = \emptyset$ and $\rho = 0$), the solvability of (2) is quite well understood in terms of the topological properties of M . Starting with the work of Y. Y. Li in [23], subsequently completed by C.-C. Chen and C.-S. Lin in [5,6], we know that, when $\rho = 0$, the solutions of (2) are uniformly

bounded for any fixed $\lambda \in \mathbb{R} \setminus 4\pi\mathbb{N}$, and the Leray-Schauder degree d_λ of the corresponding Fredholm operator can be explicitly computed: for $\lambda \in (4\pi(m-1), 4\pi m)$, $m \in \mathbb{N}^*$,

$$d_\lambda = 1 \text{ if } m = 1, \quad d_\lambda = \frac{(-\chi(M) + 1) \cdots (-\chi(M) + m - 1)}{(m - 1)!} \text{ if } m \geq 2,$$

where $\chi(M)$ is the Euler characteristics of M (see [6,28]). Actually, for the flat 2-torus we have that $d_\lambda = 1$ also when $\lambda \in 4\pi\mathbb{N}$, and so (2) has a solution for every $\lambda \in \mathbb{R}$, if $\rho = 0$ (see [6]). For the standard 2-sphere, we have that $d_\lambda = 0$ for all $\lambda > 8\pi$. But, by a more precise topological argument (see [13]), it can be shown that in this case equation (2) admits a solution for any $\lambda \in \mathbb{R} \setminus 4\pi\mathbb{N}$, if $\rho = 0$, and also some multiplicity results can be proved.

The situation is much more complex in presence of a Dirac measure. In fact, an expression for the Leray-Schauder degree, given by $d_\lambda = -\chi(M) + 2$, is available only when $\rho \geq 1$ and $\lambda \in (4\pi, 8\pi)$ (see [7]). Hence, for \mathbb{S}^2 , such a degree formula yields no information about the solvability of (2), and indeed this issue turns out to be very delicate (see [34,33]). A similarly delicate situation occurs for the flat 2-torus when $\lambda = 4\pi$ and $\rho = 2$. Consider for instance the equation

$$\Delta_g u + 4\pi \left(\frac{e^{2u}}{\int_M e^{2u} d\sigma_g} - \delta_P \right) = 0 \quad \text{in } M = \mathbb{C}/(\xi_1\mathbb{Z} + \xi_2\mathbb{Z}),$$

with $P = 0$. Then, C.-S. Lin and C. L. Wang have shown in [26] that there is no solution in the case of a rectangular lattice (*i.e.* $\xi_1 = a$, $\xi_2 = ib$, $a, b > 0$), while there is a solution for a rhombus lattice (*i.e.* $\xi_1 = a$, $\xi_2 = a e^{i\pi/3}$, $a > 0$).

As usual, the cause for such surprising existence or nonexistence situations, there is a lack of compactness for the solution set. It can be explained by looking at the singular Liouville equation

$$\Delta u + e^{2u} = 2\pi\rho \delta_{z_0} \quad \text{in } \mathbb{R}^2. \quad (3)$$

The solutions of (3) are described using complex notations (with $\mathbb{R}^2 \simeq \mathbb{C}$), by means of the Liouville formula:

$$u(z) = \frac{1}{2} \log \left(\frac{4|f'(z)|^2}{(1 + |f(z)|^2)^2} \right),$$

where f is a meromorphic function. To account for the singularity at z_0 , we must require that it corresponds to a pole of f , or a zero of f' , or a branch point of f , with the appropriate order. In particular, for all solutions u of (3)

that satisfy $e^{2u} \in L^1(\mathbb{R}^2)$, there holds:

$$\int_{\mathbb{R}^2} e^{2u} dx = 4\pi(1 + \rho),$$

(see [8,9,29]). This explains the role of 4π for the regular problem, $\rho = 0$.

One of the motivations of this paper is to connect the solvability of (2) to some weighted Liouville-type equations in \mathbb{R}^2 which generalize (3). To see how this class of equations could arise, let us first focus on problem (2) over the square 2-torus. In this case we know that a solution exists if we replace the Dirac measure by a smooth function (see [6]). Let $\Omega = (-1, 1)^2 \subset \mathbb{R}^2$ and suppose that the source point P coincides with the origin $0 \in \Omega$. For given $\lambda > 0$ and $\rho > -1$, denote by u_ε the solution of

$$\begin{cases} \Delta u_\varepsilon + \lambda \left(\frac{e^{2u_\varepsilon}}{\int_{\Omega} e^{2u_\varepsilon} dx} - \frac{1}{|\Omega|} \right) = 2\pi\rho \left(\frac{\varepsilon^2}{\pi(\varepsilon^2 + |x|^2)^2} - \frac{c_\varepsilon}{|\Omega|} \right) & \text{in } \Omega, \\ u_\varepsilon \text{ doubly periodic on } \partial\Omega, \end{cases} \quad (4)$$

where $c_\varepsilon := \int_{\Omega} \frac{\varepsilon^2}{\pi(\varepsilon^2 + |x|^2)^2} dx$ converges to 1 as $\varepsilon \rightarrow 0$. By doubly periodic on $\partial\Omega$, we mean $u(x + 2e) = u(x)$ for any $x \in \partial\Omega$, with $e = (1, 0)$ or $e = (0, 1)$. To prove the existence of a solution of (2), it is natural to investigate under which conditions on λ and ρ we can pass to the limit in (4), along with a subsequence of $\varepsilon \rightarrow 0$. In other words, whenever possible, we need to establish *a priori* estimates for u_ε in suitable norms. To this end, denote by $u_{\varepsilon,0}$ the unique solution to the problem

$$\begin{cases} \Delta u = 2\pi\rho \left(\frac{\varepsilon^2}{\pi(\varepsilon^2 + |x|^2)^2} - \frac{c_\varepsilon}{|\Omega|} \right) & \text{in } \Omega, \\ u \text{ doubly periodic on } \partial\Omega, \quad \int_{\Omega} u dx = 0, \end{cases}$$

which takes the form $u_{\varepsilon,0}(x) = \frac{\rho}{2} \log(\varepsilon^2 + |x|^2) + \psi_\varepsilon(x)$, for some suitable function ψ_ε , which is uniformly bounded in $C^{2,\alpha}$ -norm, with respect to $\varepsilon > 0$. Then $u_\varepsilon = v_\varepsilon + u_{\varepsilon,0}$ is a solution of (4) if and only if v_ε satisfies:

$$\begin{cases} \Delta v_\varepsilon + \lambda \left(\frac{e^{2(u_{\varepsilon,0} + v_\varepsilon)}}{\int_{\Omega} e^{2(u_{\varepsilon,0} + v_\varepsilon)} dx} - \frac{1}{|\Omega|} \right) = 0 & \text{in } \Omega, \\ v_\varepsilon \text{ doubly periodic on } \partial\Omega. \end{cases}$$

The function $e^{u_{\varepsilon,0}}$ is bounded from above and from below away from zero in $C_{\text{loc}}^0(\Omega \setminus \{0\})$, uniformly in $\varepsilon \in (0, 1)$. Therefore, by well known estimates based on blow-up analysis (see, *e.g.*, [23,5,6,1,31]), we know that for $\lambda \notin 4\pi\mathbb{N}$, then u_ε is bounded uniformly in $C_{\text{loc}}^{2,\alpha}(\Omega \setminus \{0\})$ with respect to $\varepsilon > 0$.

Next, we have to investigate what may happen in a neighborhood of the origin $0 \in \Omega$. To this purpose, define $w_\varepsilon := v_\varepsilon + \frac{1}{2} \log \lambda - \frac{1}{2} \log (\int_\Omega e^{2u_\varepsilon} dx)$ and $W_\varepsilon := e^{2\psi_\varepsilon}$ in $B_r(0) \subset \Omega$, for $r > 0$ small enough. There holds:

$$\begin{cases} -\Delta w_\varepsilon = (\varepsilon^2 + |x|^2)^\rho W_\varepsilon(x) e^{2w_\varepsilon} - \frac{\lambda}{|\Omega|} & \text{in } B_r(0), \\ \int_{B_r(0)} (\varepsilon^2 + |x|^2)^\rho W_\varepsilon(x) e^{2w_\varepsilon} dx = \lambda \frac{\int_{B_r(0)} e^{2u_\varepsilon} dx}{\int_\Omega e^{2u_\varepsilon} dx} \leq \lambda. \end{cases}$$

If w_ε was not uniformly bounded in $B_r(0)$, then we could find a sequence $(\varepsilon_n)_{n \in \mathbb{N}}$ with $\lim_{n \rightarrow \infty} \varepsilon_n = 0$, and a sequence $(x_n)_{n \in \mathbb{N}}$ of points in $B_r(0)$ with $\lim_{n \rightarrow \infty} x_n = 0$ such that

$$w_{\varepsilon_n}(x_n) = \max_{B_r(0)} w_{\varepsilon_n} \longrightarrow +\infty \quad \text{as } n \rightarrow \infty. \quad (5)$$

For any $n \in \mathbb{N}$, define $s_n := \max \{ \varepsilon_n, |x_n|, \exp(-\frac{w_{\varepsilon_n}(x_n)}{2(1+\rho)}) \}$ and observe that $\lim_{n \rightarrow \infty} s_n = 0$. Let $R_n := W_{\varepsilon_n}(x_n + s_n x)$, $B_n := B_{r/s_n}(0)$. For n large, $U_n(x) := w_{\varepsilon_n}(x_n + s_n x) + 2(1+\rho) \log s_n$ satisfies

$$\begin{cases} -\Delta U_n = \left(\left| \frac{\varepsilon_n}{s_n} \right|^2 + \left| \frac{x_n}{s_n} + x \right|^2 \right)^\rho R_n e^{2U_n} + o(1) & \text{in } B_n, \\ U_n(0) = w_{\varepsilon_n}(x_n) + 2(1+\rho) \log s_n, \\ \int_{B_n} \left(\left| \frac{\varepsilon_n}{s_n} \right|^2 + \left| \frac{x_n}{s_n} + x \right|^2 \right)^\rho R_n e^{2U_n} dx \leq \lambda. \end{cases}$$

By definition of s_n , we know that $\limsup_{n \rightarrow \infty} s_n \exp(\frac{w_{\varepsilon_n}(x_n)}{2(1+\rho)}) \geq 1$. We do not know whether this limit is finite or not. If

$$\limsup_{n \rightarrow \infty} s_n \exp\left(\frac{w_{\varepsilon_n}(x_n)}{2(1+\rho)}\right) < \infty, \quad (6)$$

using Harnack's estimates, we can determine a subsequence along which

$$U_n \rightarrow U_\infty \text{ in } C_{\text{loc}}^{2,\alpha}(\mathbb{R}^2), \quad \frac{\varepsilon_n}{s_n} \rightarrow \varepsilon_\infty \in [0, 1], \quad \frac{x_n}{s_n} \rightarrow x_\infty \in B(0, 1) \subset \mathbb{R}^2$$

and $W_{\varepsilon_n}(x_n + s_n x)$ pointwise converges to a positive constant $W_\infty > 0$. In addition, U_∞ satisfies:

$$\begin{cases} -\Delta U_\infty = (\varepsilon_\infty^2 + |x_\infty + x|^2)^\rho W_\infty e^{2U_\infty} & \text{in } \mathbb{R}^2, \\ U_\infty(0) = \max_{\mathbb{R}^2} U_\infty \geq 0, \\ \int_{\mathbb{R}^2} (\varepsilon_\infty^2 + |x_\infty + x|^2)^\rho W_\infty e^{2U_\infty} dx \leq \lambda. \end{cases}$$

If $\varepsilon_\infty = 0$, then by well known classification results (see [9,29]), we know that

$$W_\infty \int_{\mathbb{R}^2} |x_\infty + x|^{2\rho} e^{2U_\infty} dx = 4\pi(1 + \rho).$$

So, we could rule out the occurrence of (5) in this case by restricting the problem to $\lambda < 4\pi(1 + \rho)$.

Hence, assume that $\varepsilon_\infty > 0$. Then by adding a constant to U , scaling and translating we arrive at the limiting problem:

$$\begin{cases} -\Delta U = (1 + |x|^2)^\rho e^{2U} & \text{in } \mathbb{R}^2, & \text{(a)} \\ \int_{\mathbb{R}^2} (1 + |x|^2)^\rho e^{2U} dx \leq \lambda. & \text{(b)} \end{cases} \quad (7)$$

Now, with $\varepsilon_\infty > 0$, it is much harder to identify the range of values assumed by $\int_{\mathbb{R}^2} (1 + |x|^2)^\rho e^{2U} dx$ for all solutions of (7a). It is no longer a singleton, but a whole interval contained in $(2\pi(1 + \rho), 4\pi(1 + \rho))$ and containing $(4\pi \max(1, \rho), 4\pi(1 + \rho))$, whose explicit range is still under investigation, even when we restrict the problem to radially symmetric solutions of (7a), see [11,24,25]. In this paper, we are going to identify necessary conditions on λ that allow us to rule out the occurrence of (5) under Condition (6). In other words, if, for some λ_0 , there is no solution of (7) for any $\lambda \leq \lambda_0$, then (2) with $M = \Omega$ would have a solution, provided Condition (6) holds.

On the other hand, consider on \mathbb{R}^2 the solutions of

$$\begin{cases} -\Delta u = (1 + |x|^2)^N e^{2u} & \text{in } \mathbb{R}^2, \\ \int_{\mathbb{R}^2} (1 + |x|^2)^N e^{2u} dx = \lambda. \end{cases} \quad (8)$$

This problem is equivalent to (2). Let $\Sigma : \mathbb{S}^2 \rightarrow \mathbb{R}^2$ be the stereographic projection with respect to the north pole, $\mathbf{N} := (0, 0, 1)$. To any solution u of (8), we can associate

$$v(y) := u(x) - \frac{1}{2} \log \left(\frac{\lambda}{(1 + |x|^2)^{N+2}} \right) - \log 2, \quad (9)$$

for $x = \Sigma(y)$ and any $y \in \mathbb{S}^2$. The function v solves

$$\Delta_g v + \lambda \frac{e^{2v}}{\int_{\mathbb{S}^2} e^{2v} d\sigma_g} = \frac{2\pi(N+2)}{|\mathbb{S}^2|} \quad \text{on } \mathbb{S}^2 \setminus \{\mathbf{N}\}. \quad (10)$$

Moreover, if u is a locally bounded solution of (8), according to [11], $u(x) \sim -\frac{\lambda}{2\pi} \log |x|$ as $|x| \rightarrow \infty$, which shows that

$$\lim_{y \in \mathbb{S}^2, y \rightarrow \mathbf{N}} \frac{v(y)}{\log |y - \mathbf{N}|} = \frac{\lambda}{2\pi} - (N+2),$$

Here, by $|y - \mathbf{N}|$, we denote the euclidean distance from y to \mathbf{N} in $\mathbb{R}^3 \supset \mathbb{S}^2$. As a consequence,

$$\Delta_g v = \left(\lambda - 2\pi(N+2) \right) \delta_{\mathbf{N}} + O(1)$$

as $y \rightarrow \mathbf{N}$. Hence v is a solution of (2) in S^2 with $\rho = \frac{\lambda}{2\pi} - (N+2)$ and $f = 0$. Here the parameter ρ is not the same as in (7). Notice that v given by (9) is such that $\int_{\mathbb{S}^2} e^{2v} d\sigma_g = 1$. As a special case, if $\lambda = 2\pi(N+2)$, we find that v is a bounded solution of (2) with $\rho = 0$. Viceversa, if v is a solution of (2) with $f = 0$, then

$$u = v \circ \Sigma^{-1} - \frac{1}{2} \log \left(\int_{\mathbb{S}^2} e^{2v} d\sigma_g \right) + \log 2 + \frac{1}{2} \log \left(\frac{\lambda}{(1 + |x|^2)^{N+2}} \right)$$

is a solution of (8). So, existence and multiplicity of solutions to (8) provide us with existence and multiplicity of solutions to (2) on the 2-sphere.

From now on, we shall focus on the study of (8), for all positive values of N . Our main result establishes the existence of many bounded, radial solutions.

Theorem 1 *For all $k \geq 2$ and $N > k(k+1) - 2$, there are at least $2(k-2) + 2$ distinct radial solutions of (8) with $\lambda = 2\pi(N+2)$, one of them being the function $u_N^*(r) := \frac{1}{2} \log \left(\frac{2(N+2)}{(1+r^2)^{N+2}} \right)$ with $r = |x|$.*

To conclude this introduction let us mention that in many papers devoted to other applications than assigned Gauss curvature problems, the convention is to consider equation (2) with e^u instead of e^{2u} . Necessary adaptations are straightforward (replace u by $\frac{1}{2}(u - \log 2)$) and therefore are left to the interested reader.

2 Known results

In this section, we collect known existence results and some of the properties of the bounded solutions to

$$\Delta u + (1 + |x|^2)^N e^{2u} = 0, \quad x \in \mathbb{R}^2, \quad \text{and} \quad \int_{\mathbb{R}^2} (1 + |x|^2)^N e^{2u} dx < \infty, \quad (11)$$

where $N > 0$ is a real parameter which in the applications usually enters as an integer value. Several papers, mostly by C.-S. Lin *et al.* [11,24,25], have dealt with this class of equations. Almost all results are concerned with the set of radially symmetric solutions of (11), which can be parametrized by a parameter $a \in \mathbb{R}$ as follows:

$$\begin{cases} u_a'' + \frac{u_a'}{r} + (1 + r^2)^N e^{2u_a} = 0 & \text{in } (0, +\infty), \\ u_a(0) = a, \quad u_a'(0) = 0, \quad \int_0^\infty (1 + r^2)^N e^{2u_a} r dr < +\infty. \end{cases} \quad (12)$$

It is proven in [11,24,25] that, for every $a \in \mathbb{R}$, there exists a unique solution u_a of (12), which moreover satisfies

$$\lim_{r \rightarrow \infty} (u_a(r) + \alpha(a) \log r) = \beta(a), \quad (13)$$

where $a \mapsto \alpha(a)$ and $a \mapsto \beta(a)$ are two $C^1(\mathbb{R}, \mathbb{R})$ functions defined by:

$$\alpha(a) = \int_0^\infty (1 + r^2)^N e^{2u_a} r dr \quad \text{and} \quad \beta(a) = \int_0^\infty (1 + r^2)^N e^{2u_a} r \log r dr.$$

Moreover it was proved in [11] (also see [25]) that, for any $N > 0$,

$$\lim_{a \rightarrow -\infty} \alpha(a) = 2(N + 1) \quad \text{and} \quad \lim_{a \rightarrow +\infty} \alpha(a) = 2 \min\{1, N\}. \quad (14)$$

Pohozaev's identity applied to (11) shows that $\alpha(a) \in (2, 2(N+1))$. For integrability reasons, we also know that $\alpha(a) > N+1$, and so

$$\max\{2, N+1\} < \alpha(a) < 2(N+1) \quad \forall a \in \mathbb{R}.$$

Our problem is to find the solutions of (12) corresponding to a given α , that is

$$\begin{cases} u'' + \frac{u'}{r} + (1+r^2)^N e^{2u} = 0 & \text{in } (0, +\infty), \\ u'(0) = 0, \quad \int_0^\infty (1+r^2)^N e^{2u} r dr = \alpha. \end{cases} \quad (15)$$

In [24,25], C.-S. Lin investigates the uniqueness issue for problem (15) by identifying the values of α for which there is a unique a such that $\alpha(a) = \alpha$. He proves in [25] that uniqueness holds for $\alpha \in (2N, 2(N+1))$ if $N > 1$ and for all $\alpha \in (2, 2(N+1))$ if $N \leq 1$. On the other hand, it is easy to verify that for all N , the function

$$u_N^*(r) := \frac{1}{2} \log \left(\frac{2(N+2)}{(1+r^2)^{N+2}} \right) \quad (16)$$

is a solution to (12) for $a = a_N^* := \frac{1}{2} \log(2(N+2))$, and satisfies $\alpha(a_N^*) = N+2$. Since $N+2 < 2N < 2(N+1)$ for all $N > 2$, by continuity of $a \mapsto \alpha(a)$, it appears that there exists at least two different values of a such that $\alpha(a) = \alpha$, for any $\alpha \in (\min_{a \in \mathbb{R}} \alpha(a), 2N)$. In other words, for those values of α there exists at least two radially symmetric solutions of (11) satisfying:

$$\int_{\mathbb{R}^2} (1+|x|^2)^N e^{2u} dx = 2\pi \alpha.$$

Moreover for $N=2$, $\alpha(a_2^*) = 4 = 2N$ and $\alpha'(a_2^*) < 0$, so the above multiplicity results also holds true for $N=2$. Summarizing, we can state the following result.

Theorem 2 [11,24,25] *Let N be any positive real number.*

- (i) *If $N \leq 1$, then the curve $a \mapsto \alpha(a)$ is monotone decreasing. Moreover, there exists a radially symmetric solution u of (15) if and only if $\alpha \in (2, 2(N+1))$, and such a solution is unique.*
- (ii) *If $N > 1$, then for all $\alpha \in (2N, 2(N+1))$, there exists a unique $a \in \mathbb{R}$ such that $\alpha(a) = \alpha$. In other words, for such α , problem (15) is satisfied by a unique radial solution.*

(iii) If $N \geq 2$, then $\min_{a \in \mathbb{R}} \alpha(a) < 2N$, and for all $\alpha \in (\min_{a \in \mathbb{R}} \alpha(a), 2N)$, there exists at least two radial solutions of (15).

Remark. Concerning part (iii) of Theorem 2, by a closer inspection of the results of [24], actually we know that for $N \geq 2$, problem (15) is satisfied by a unique radial solution also when $\alpha = 2N$.

An important tool in the proof of the above uniqueness results is the study of the linearized problem

$$\begin{cases} \varphi_a'' + \frac{\varphi_a'}{r} + 2(1+r^2)^N e^{2u_a} \varphi_a = 0, & r \in (0, +\infty), \\ \varphi_a(0) = 1, \quad \varphi_a'(0) = 0, \end{cases} \quad (17)$$

and in particular the number of zeroes of the function φ_a when α is in the range $(2N, 2(N+1))$. The number of critical points of $a \mapsto \alpha(a)$ is also connected with the number of zeroes of φ_a in the range $(\min_{a \in \mathbb{R}} \alpha(a), 2N)$. It is indeed easy to prove that as r goes to $+\infty$,

$$\varphi_a(r) \sim -\alpha'(a) \log r + b'(a) + o(1), \quad (18)$$

and hence, that φ_a is a bounded function if and only if $a \in \mathbb{R}$ is a critical point of the function α . As a special case, for all N , if $\min_{a \in \mathbb{R}} \alpha(a)$ is achieved for some finite \underline{a} , then $\varphi_{\underline{a}}$ is bounded.

2.1 Non radially symmetric solutions

By (9), to any solution u_a of (12), we can associate a function v_a on \mathbb{S}^2 , such that $\int_{\mathbb{S}^2} e^{2v} d\sigma_g = 1$, which solves (10) for $\lambda = 2\pi\alpha(a)$. At level $\alpha = N+2$, v_a is a bounded solution of (2) (with $f = 0$, $\rho = 0$), which is axially symmetric with respect to the unit vector $(0, 0, 1)$ pointing towards the north pole \mathbf{N} of \mathbb{S}^2 . Since $v_N^* = v_{a_N^*}$ is the unique constant solution of (2), if we know the existence of more than one solution at level $\alpha = N+2$, then there is an axially symmetric solution of (10) which is not constant, and that can be thus rotated in order to be axially symmetric with respect to any vector $\mathbf{e} \in \mathbb{S}^2 \setminus \{\mathbf{N}, \mathbf{S}\}$. Let us denote by $v_{\mathbf{e}}$ such a solution. Applying (9) to $v_{\mathbf{e}}$, we find a solution $u_{\mathbf{e}}$ of (11) which is not radially symmetric. If at level $\lambda = 2\pi(N+2)$ we find k solutions of (11) different from u_N^* , then we get k punctured spheres of non radially symmetric solutions of (11). Details can be found in [25].

3 Multiplicity results for radially symmetric solutions

For given N , critical levels of the curve $a \mapsto \alpha(a)$ determine the multiplicity of the radial solutions at a given level. The number of zeros of the solutions of the linearized problem can change only at critical points of α , see below Section 3.2. In the special case $\alpha = N + 2$, a bifurcation argument provides us with a very precise multiplicity result, which is our main result, see Section 3.3.

3.1 A preliminary result

Let $(c_k^N)_{k=1}^{n-1}$ be the ordered sequence of all critical values, counted with multiplicity, of the curve $a \mapsto \alpha(a)$, $c_0^N := \inf_{a \in \mathbb{R}} \alpha(a)$ and $c_n^N = 2(N + 1)$. Denote by $(a_k^N)_{k=1}^{n-1}$ a sequence of critical points corresponding to $(c_k^N)_{k=1}^{n-1}$ and, for any $k = 1, 2, \dots, n - 1$, let $\epsilon_k^N := +2$ if a_k^N is a local minimum, $\epsilon_k^N := -2$ if a_k^N is a local maximum, and $\epsilon_k^N := 0$ otherwise. Also let $\epsilon_0^N := 2$ if $\inf_{a \in \mathbb{R}} \alpha(a)$ is not achieved, and 0 otherwise. Let $\chi_N(\alpha) := 1$ if $\alpha > 2N$ and 0 otherwise. The next proposition links these values with the number of solutions of (15).

Proposition 3 *Let N be any positive real number. With the above notations, for any $\alpha > 0$, Equation (15) has exactly $\sum_{j=0}^k \epsilon_j^N - \chi(\alpha)$ solutions such that $\alpha(a) = \alpha$ if $\alpha \in (c_k^N, c_{k+1}^N) \cap (\mathbb{R} \setminus \{2N\})$, for any $k = 0, 1, \dots, n - 1$.*

The proof is straightforward and left to the reader. We shall now focus on the study of the critical points of the curve $a \mapsto \alpha(a)$. Our main tool is the linearization of (12).

3.2 Study of the linearized problem

In order to study the multiplicity of radial solutions for (8), it is convenient to perform the Emden-Fowler transformation in the linearized equation (17):

$$t = \log r, \quad w_a(t) := \varphi_a(r).$$

The equation in (17) is then transformed into

$$w_a''(t) + 2e^{2t}(1 + e^{2t})^N e^{2ua(e^t)} w_a(t) = 0, \quad t \in (-\infty, +\infty). \quad (19)$$

When $a = a_N^*$, the equation for $w_N^* := w_{a_N^*}$ reads

$$w_N^{*''}(t) + \frac{(N + 2)}{2(\cosh t)^2} w_N^*(t) = 0, \quad t \in (-\infty, +\infty). \quad (20)$$

With one more change of variables, $w(t) = \psi(s)$, $s = \tanh t$, we find Legendre's equation:

$$\frac{d}{ds} \left((1-s^2) \frac{d\psi}{ds} \right) + \frac{N+2}{2} \psi = 0$$

which defines the Legendre polynomial of order $k \in \mathbb{N}^*$ if $N+2 = k(k+1)$. Notice that the composition of the two above changes of variables amounts to write $s = \frac{r^2-1}{r^2+1}$. A very precise spectral analysis made in [22] shows that the above equation has bounded solutions if and only if there is a positive integer k such that $1+2k = \sqrt{1+4(N+2)}$, that is, if and only if

$$\mathfrak{N}(N) := \frac{-1 + \sqrt{1+4(N+2)}}{2}$$

is a positive integer. This is solved by $N = N_k := k(k+1) - 2$, $k \in \mathbb{N}^*$. Actually, we are interested only in $k \geq 2$ since we only deal with $N > 0$. As a consequence, a_N^* is a critical point of α if and only if $\mathfrak{N}(N)$ is a positive integer: $N = 4, 10, 18, \dots$

For $N = N_k > 0$, we know explicitly the solutions to (20). They are the Legendre polynomials, namely, with $s = \tanh t$,

$$w_{N_k}^*(t) \equiv P_k(s) \text{ for all integer } k \geq 2. \quad (21)$$

Lemma 4 [22] *Take $N \geq 1$. Then, there exist bounded solutions of (20) if and only if $\mathfrak{N}(N)$ is a positive integer. In such a case, $\varphi_{a_N^*}$ has exactly $\mathfrak{N}(N)$ zeroes in the interval $(-\infty, +\infty)$.*

For all $a \in \mathbb{R}$, φ_a has at least two zeroes in the interval $(0, +\infty)$. This observation is a key step in the uniqueness proofs of [24,25]. Zeroes of φ_a will also play an important role in multiplicity results. The next observation is a standard result for linear ordinary differential equations.

Lemma 5 *For any $N > 0$, $a_0 > 0$ and $R > 0$, if φ_{a_0} has k zeroes in $(0, R)$ and $\varphi_{a_0}(R) \neq 0$, then there exists an $\varepsilon > 0$ such that φ_a also has exactly k zeroes in $(0, R)$ for any $a \in (a_0 - \varepsilon, a_0 + \varepsilon)$.*

Proof. If the result were wrong, then we could find some $r \in (0, R)$ such that $\varphi_{a_0}(r) = \varphi'_{a_0}(r) = 0$ and so we would get $\varphi_{a_0} \equiv 0$. \square

Corollary 6 *For any $N \in [N_k, N_{k+1})$, $k \in \mathbb{N}$, $k \geq 1$, solutions to (20) have exactly $k+1$ zeroes in the interval $(-\infty, +\infty)$.*

Proof. If we normalize all functions w_N^* so that $\lim_{t \rightarrow -\infty} (w_N^*(t), w_N^{*\prime}(t)) = (1, 0)$, then by continuity in N the number of zeroes changes if and only if w_N^* is bounded, *i.e.* if $\mathfrak{N}(N)$ is a positive integer, by Lemma 4. At $N_1 = 0$, we have: $w_0^*(t) = \tanh(t)$. A careful analysis shows that, as a function of N , the number of zeroes is continuous from the right and increasing. \square

Lemma 7 *Take $N \geq 1$ and consider $a_1, a_2 \in \mathbb{R}$ such that $\alpha'(a_1) = \alpha'(a_2) = 0$ and $\alpha'(a) \neq 0$ if $a \in (a_1, a_2)$. Then, for all $a \in (a_1, a_2)$, the functions φ_a have the same number of zeroes.*

Proof. The proof is based on the same arguments as the proof of Corollary 6. One has just to replace the continuity in N by the continuity in a . \square

Lemma 5 means that when a varies, zeroes may appear or disappear only at infinity. For a given a , the sign of the function

$$J_N(a) := \int_0^{+\infty} (1+r^2)^N e^{2u_a} \varphi_a^3 r dr \quad (22)$$

governs the dynamics of the zeroes of φ_a at infinity as follows. Denote by $r(a) := \max\{r > 0 : \varphi_a(r) = 0\}$ the largest zero of φ_a .

Lemma 8 *Let $\bar{a} > 0$ be such that, for $\zeta = \pm 1$, $\lim_{a \rightarrow \bar{a}, \zeta(a-\bar{a}) > 0} r(a) = \infty$. Then there exists $\varepsilon > 0$ such that, on $(\bar{a} - \varepsilon, \bar{a})$ if $\zeta = -1$, on $(\bar{a}, \bar{a} + \varepsilon)$ if $\zeta = +1$,*

$$\frac{dr}{da}(a) = -\frac{4}{r(a) |\varphi_a'(r(a))|^2} \int_0^{r(a)} (1+r^2)^N e^{2u_a} \varphi_a^3 r dr$$

and $\frac{dr}{da} J_N(a) < 0$ if $J_N(\bar{a}) \neq 0$.

Proof. First, we choose $\varepsilon > 0$ small enough so that $\int_0^{r(a)} (1+r^2)^N e^{2u_a} \varphi_a^3 r dr$ and $J_N(a)$ have the same sign, if $J_N(\bar{a}) \neq 0$. Next, we take $b > 0$, small. Multiplying the equation satisfied by φ_a and φ_{a+b} by $r \varphi_{a+b}$ and $r \varphi_a$ respectively, and integrating by parts in the interval $(0, r(a))$, we get

$$2 \int_0^{r(a)} (1+r^2)^N (e^{2u_a} - e^{2u_{a+b}}) \varphi_a \varphi_{a+b} r dr = -r(a) \varphi_a'(r(a)) \varphi_{a+b}(r(a)).$$

By definition of φ_a and using the uniform continuity properties of the functions u_a and φ_a on $(0, r(a))$, we obtain: $\|u_{a+b} - u_a - b \varphi_a\|_{L^\infty(0, r(a))} = o(b)$ as $b \rightarrow 0$,

$(1 - e^{2b\varphi_a}) \sim -2b\varphi_a$. Since $\varphi_{a+b}(r(a)) = \varphi'_a(r(a))(r(a+b)) - r(a) + o(b)$, we get, as $b \rightarrow 0$,

$$-4 \int_0^{r(a)} (1+r^2)^N e^{2u_a} \varphi_a^3 r dr = r(a) |\varphi'_a(r(a))|^2 \frac{r(a+b) - r(a)}{b} + o(b).$$

□

Notice that, with the notations of Lemma 8, $\varphi_{\tilde{a}}$ is a bounded function. As a consequence, we have the following result.

Corollary 9 *Let \tilde{a} be a critical point of α . There exists $\varepsilon > 0$, small enough, such that the following properties hold.*

- (i) *If $J_N(\tilde{a}) > 0$ and if, for any $a \in (\tilde{a} - \varepsilon, \tilde{a})$, all functions φ_a are unbounded and have k zeroes in $(0, +\infty)$, then $\varphi_{\tilde{a}}$ is bounded and has k zeroes, and for any $a \in (\tilde{a}, \tilde{a} + \varepsilon)$, φ_a is unbounded and has either k or $k + 1$ zeroes in $(0, +\infty)$.*
- (ii) *If $J_N(\tilde{a}) < 0$ and if, for any $a \in (\tilde{a} - \varepsilon, \tilde{a})$, all functions φ_a are unbounded and have k zeroes in $(0, +\infty)$, then $\varphi_{\tilde{a}}$ is bounded and has either k or $k - 1$ zeroes, and for any $a \in (\tilde{a}, \tilde{a} + \varepsilon)$, φ_a has the same number of zeroes as $\varphi_{\tilde{a}}$.*

As already seen in Proposition 3, if \tilde{a} is a local extremum of α , the number of zeroes changes when a goes through \tilde{a} , since the sign of φ_a at infinity also changes, by (18). Otherwise, if \tilde{a} is an inflection point, the number of zeroes is constant when a passes through \tilde{a} . This explains the ambiguity in the previous result.

Actually for the particular case $a = a_N^*$, $N = N_k$ we can exactly compute the value of J_N as follows:

Proposition 10 *Let us define $j(k) := J_N(a_{N_k}^*)$ for any integer $k \geq 2$. Then, $j(k) = 0$ if k is odd, and $j(k) > 0$ if k is even.*

Proof. By using (21), we can easily compute

$$j(k) = \frac{1}{2}k(k+1) \int_{-1}^1 P_k(s)^3 ds. \quad (23)$$

Now, when k is odd, P_k is also odd and so, $j(k) = 0$. On the contrary, Gaunt's formula, see [15, Identity (14), page 195] shows that $j(k) > 0$ if k is even. □

Remark. If $N = N_k$, k even, a_N^* is a local minimum for the function α . Indeed, this follows from the fact that we know that $J_N(a_N^*) > 0$ and the number of zeroes of P_k . From this we infer that the function w_N^* is positive at infinity for N close to N_k with $N < N_k$, and negative at infinity for N close to N_k with $N > N_k$. Hence, α is decreasing to the left of a_N^* and increasing afterwards.

3.3 A multiplicity result at level $\alpha = N + 2$

This section is devoted to the proof of Theorem 1, that is a multiplicity result for the solutions of problem (15) at level $\alpha = N + 2$, which also helps to illustrate Theorem 1.10 in [11]. As seen in the introduction this amounts to study the number of bounded solutions to (2) with $\rho = 0$ (without singularities). We show that when $\lambda = 2\pi\alpha$ and $\alpha = N + 2$ becomes large, there are more and more bounded solutions to (2).

Let u be a radial solution of (8) with $\alpha = N + 2$. We may reformulate this problem in terms of $f := u - u_N^* \in \mathcal{D}^{1,2}(\mathbb{R}^2)$ as a solution to:

$$\Delta f + \frac{\mu}{(1 + |x|^2)^2} (e^{2f} - 1) = 0 \text{ in } \mathbb{R}^2, \quad \int_{\mathbb{R}^2} \frac{e^{2f}}{(1 + |x|^2)^2} dx = \pi, \quad (24)$$

with $\mu = 2(N + 2)$. Solutions of (24) are bounded by (13). Moreover, (24) is trivially invariant under the Kelvin transformation:

Lemma 11 *If f is a solution of (24), then the function $x \mapsto f\left(\frac{x}{|x|^2}\right)$ is also a solution of (24).*

This lemma allows us to characterize many branches of solutions of (24).

Theorem 12 *The function $f \equiv 0$ is a trivial solution of (24) for any $\mu > 0$. For any $k \geq 2$, there are two continuous half-branches, \mathcal{C}_k^+ and \mathcal{C}_k^- , of solutions (μ, f) of (24) bifurcating from the branch of trivial solutions at, and only at, $(\mu_k = 2k(k + 1), 0)$. Solutions in \mathcal{C}_k^\pm are such that $\pm f(0) > 0$.*

Away from the trivial solutions, all branches are disjoint, unbounded and characterized by the number of zeroes. In \mathcal{C}_k^\pm , the solutions of (24) have exactly k zeroes. If k is odd, the branch \mathcal{C}_k^\mp is the image of \mathcal{C}_k^\pm by the Kelvin transform. If k is even, the half-branches \mathcal{C}_k^\pm are invariant under the Kelvin transform. Finally, \mathcal{C}_k^\pm for $k \geq 3$ and \mathcal{C}_2^- are locally bounded in μ .

We divide the proof in three steps.

Step 1: Existence of unbounded branches of solutions.

We use a bifurcation method to study the set of radial solutions of (24) with the bifurcation parameter μ . Since branches may be multi-valued in terms of N , we will reparametrize them with a parameter s . Classical results apply for instance in $\mathbb{R}^+ \times (\mathcal{D}^{1,2} \cap C^2)$, see [12,30].

By Lemma 4, there is local bifurcation from the trivial line $\{(\mu, 0)\}$ at the points $(\mu_k := 2(N_k + 2), 0)$, and there is no other bifurcation point in this trivial branch. By the properties of the Legendre polynomials P_k , if we denote by \mathcal{C}_k^\pm the two continuous half-branches of non-trivial solutions that meet at $(\mu_k, 0)$, one easily proves that for any $(\mu, f) \in \mathcal{C}_k^\pm$ in a neighborhood of $(\mu_k, 0)$, $f(r)$ has exactly k zeroes in the interval $(0, +\infty)$.

Actually, on \mathcal{C}_k^\pm , the number of zeroes of the solutions is constant, namely equal to k . For instance, let us prove it for \mathcal{C}_k^+ by smoothly parametrizing the branch as follows:

$$\mu(s) \text{ with } \mu(0) = 2(N_k + 2), \text{ and } f = f_s \text{ with } f_0 \equiv 0, \quad s \in \mathbb{R}. \quad (25)$$

Let $\Lambda_k := \{s \in (0, \infty) : f_s \text{ admits exactly } k \text{ zeroes}\}$. Clearly Λ_k is not empty. Since a solution $f \not\equiv 0$ of (24) cannot vanish at a point together with its derivative, the smoothness of the map $s \mapsto f_s$ ensures that Λ_k is open. We check that Λ_k is also closed. Indeed, for $(s_n)_{n \in \mathbb{N}} \in \Lambda_k^\mathbb{N}$ with $\lim_{n \rightarrow \infty} s_n = s_\infty \in (0, \infty)$, we see that the zeroes of $f_n = f_{s_n}$ cannot collide at a point $r_0 \in \mathbb{R}^+$ since there, $\lim_{n \rightarrow \infty} f_n = f_{s_\infty} \not\equiv 0$ would vanish together with its derivatives. So, for a zero to appear or disappear, this would require the existence of a sequence $(r_n)_{n \in \mathbb{N}}$ with $f_n(r_n) = 0$ and $\lim_{n \rightarrow \infty} r_n = \infty$. But then, $\tilde{f}(r) = f_{s_\infty}(1/r)$, which is still a solution of (24) by Lemma 11, would vanish at $r = 0$ together with its derivatives, and this is again impossible. In conclusion, $\Lambda_k = (0, +\infty)$.

Consequently, non-trivial branches with different k cannot intersect or join two different points of bifurcation in the trivial branch. For any $k \geq 2$, the half-branches \mathcal{C}_k^\pm are therefore unbounded and we can distinguish them as follows: if $(\mu, f) \in \mathcal{C}_k^+$, resp. $(\mu, f) \in \mathcal{C}_k^-$, then $f(0) > 0$, resp. $f(0) < 0$.

Step 2: Symmetry under Kelvin transform

Branches of solutions of (24) have an interesting symmetry property. If k is odd, the solutions in the branches \mathcal{C}_k^\pm have an odd number of zeroes in $(0, +\infty)$ and so, they cannot be invariant under the Kelvin transform, because they take values of different sign at 0 and near $+\infty$. Since $\mu_k = 2(N_k + 2)$ is a simple bifurcation point, the only possibility is that the branches \mathcal{C}_k^\pm transform into

each other through the Kelvin transform. Otherwise, there would be at least four half-branches bifurcating from $(\mu_k, 0)$, which is impossible.

If k is even, the solutions of \mathcal{C}_k^\pm have an even number of zeroes. So, they take values of the same sign at 0 and near $+\infty$. If they were not invariant under the Kelvin transform, we would find two new branches, $\tilde{\mathcal{C}}_k^\pm$, bifurcating from $(\mu_k, 0)$, which is again impossible.

Step 3: Asymptotic behaviour of the branches

Non trivial branches of radial solutions are contained in the region $\{\mu > 8\}$, that is $N > 2$ (see Theorem 2 and the remark immediately afterwards for $N = 2$). Furthermore, with the notations of Theorem 2, by (14), there exists a unique $a(N) \in \mathbb{R}$ such that $\alpha(a) > 2N$ for all $a < a(N)$. For $N > 2$, since $N + 2 < 2N < 2(N + 1)$, if $\alpha(a) = N + 2$, then $a > a(N)$. Hence $f(0) = u(0) - u_N^*(0) > a(N) - u_N^*(0)$ with $u = u_a$ given by (15).

As a consequence, the branches \mathcal{C}_k^- are locally bounded for $\mu \in [8, +\infty)$ for any $k \geq 2$. By Step 2, \mathcal{C}_3^+ is also locally bounded for $\mu \in [8, +\infty)$. Since non trivial branches do not intersect, \mathcal{C}_k^\pm , $k \geq 3$, are all locally bounded for $\mu \in [8, +\infty)$. \square

As a simple consequence of Theorem 12, we have the following corollary, which is the counterpart of Theorem 1 written for (24).

Corollary 1 *For all $k \geq 2$, for all $\mu > \mu_k = 2k(k+1)$, there at least $2(k-2)+2$ distinct radial solutions of (24), one of them being the zero solution.*

The bifurcation diagram obtained for equation (24) (see Fig. 1, left) is easily transformed into a bifurcation diagram for the solutions of (8) with $\lambda = 2\pi(N + 2)$ (see Fig. 1, right) through the transformation $u = f + u_N^*$. In the case of equation (8), branches bifurcate from the set of trivial solutions $\mathcal{C} := \{(N, \frac{1}{2} \log(2(N + 2)))\}$, in the representation $(N, a = u(0))$.

Based on numerical evidence (see Fig. 1), it is reasonable to conjecture that, in contrast, the branch $\mathcal{C}_{k=2}^+$ admits a vertical asymptote in the sense that as $s \rightarrow +\infty$, then N converges to 2, which is the only admissible value by (14). So for $(\mu(s), f_s) \in \mathcal{C}_2^+$, f_s should develop a concentration phenomenon at the origin, and as $s \rightarrow +\infty$, we should have: $a \rightarrow +\infty$, $N \rightarrow 2_+$ and $\frac{\mu(s)}{(1+|x|^2)^2} e^{2f_s} \rightharpoonup 8\pi \delta_{z=0}$, weakly in the sense of measures.




Fig. 1. Bifurcation diagram in the representation $(N, f(0))$ for equation (24) (left) and (N, a) for equation (8) with $\lambda = 2\pi(N+2)$ (right). Non trivial branches bifurcate from $N_k = 4, 10, 18, 28, \dots$

3.4 Non radially symmetric solutions

As already described in Section 2.1, to any solution $u \neq u_N^*$ of (11) such that $\int_{\mathbb{R}^2} (1 + |x|^2)^N e^{2u} dx = 2\pi(N+2)$, we can associate a punctured sphere of non radially symmetric solutions of (11), u_e with $e \in \mathbb{S}^2 \setminus \{\mathbf{N}, \mathbf{S}\}$, satisfying also $\int_{\mathbb{R}^2} (1 + |x|^2)^N e^{2u_e} dx = 2\pi(N+2)$ for all $e \in \mathbb{S}^2 \setminus \{\mathbf{N}, \mathbf{S}\}$. And so, for $N > N_k$, there are at least $2(k-2) + 1$ punctured spheres of non radially symmetric solutions to (11) at level $\lambda = 2\pi(N+2)$.

4 Further results, numerical observations and conjectures

In the study of (15), multiplicity results for general values of α are difficult to deduce from Proposition 3 since they require a detailed analysis of the nature of each critical point: maximum, minimum, and even more in the case of an inflection point, as well as precise estimates of the corresponding critical value. Numerically, Proposition 3 gives straightforward results, which can be observed directly from the plots of the curve $a \mapsto \alpha(a)$ for various values of N , see Figs. 2, 3.

Let us give some details. We consider the solution u_a of (12) parametrized by $a = u_a(0)$. Recall that for a given $N > 1$, the curve $a \mapsto \alpha(a) = \int_0^\infty (1 + r^2)^N e^{2u_a} r dr$ is such that $\lim_{a \rightarrow -\infty} \alpha(a) = 2(N+1)$, $\lim_{a \rightarrow \infty} \alpha(a) = 2N$, and for N large enough, its range is an interval $[\alpha_N, 2(N+1))$ if α_N is achieved, or $(2N, 2(N+1))$ otherwise. If $N > 2$, $\alpha_N := \min_{a \in \mathbb{R}} \alpha(a) \leq \alpha(a_N^*) = N+2 < 2N$, where $a_N^* := \frac{1}{2} \log(2(N+2))$, which provides a multiplicity result for (15) in the range $\alpha \in (\alpha_N, 2N) \supset (N+2, 2N)$.

As a function of $N > 0$, we observe that $\alpha_N = \inf_{a \in \mathbb{R}} \alpha(a) < 2N$ if and only if $N > N_0$, where N_0 is numerically found of the order of 1.27 ± 0.02 , although its exact value is not easy to determine, see Fig. 4, left. In the range $(N_0, 20)$, we observe that α_N is achieved by the first critical point, see Fig. 4, right.




Fig. 2. Curves $a \mapsto \alpha(a)$ for various values of N : $N = 25$ (left) and $N = 1, 2, 3, \dots, 12$ (right). The point $(a_N^*, N + 2)$ corresponding to the explicit solution (16) is represented by a gray dot.




Fig. 3. Curves $a \mapsto \alpha(a)$ for $N = N_k$, $k = 2, 3, 4, 5$ (left) and $a \mapsto \alpha(a) - 2N$ for $N = 1, 3, 5, \dots, 19$. The function $N \mapsto \alpha_N - 2N$ is monotone decreasing.




Fig. 4. Critical points (left) and critical values (divided by $4N$, right) of α , as a function of N .

For any $N > N_0$, we observe that there are at least two radial solutions of (15) for any $\alpha \in (\alpha_N, 2N)$. By the observations of Section 3.4, it follows that there should be at least a radial solution and a whole punctured sphere of non radially symmetric solutions at level $\alpha = N + 2$. This supports a conjecture by C.-S. Lin in [25]. Actually, we can state the following result, which slightly improves on Theorem 2, (iii), and rigorously defines N_0 .

Proposition 13 *There exists $N_0 \in (1, 2)$ such that, for all $N \in (N_0, \infty)$, $\alpha_N = \min_{a \in \mathbb{R}} \alpha(a) < 2N$, and for all $\alpha \in (\alpha_N, 2N)$, there exists at least two solutions of (15).*

Proof. The proof relies on the continuity of the curve $a \mapsto \alpha(a)$ with respect

to N and the fact that $\alpha'(a_2^*) < 0$, thus proving that $\alpha_N < 2N$ also for $N < 2$, close enough to 2. \square

It seems that as $N \rightarrow N_0$, $N > N_0$, α_N is achieved by a unique $\tilde{a}_N \rightarrow \infty$, so we may conjecture that (15) has multiple solutions for $\alpha \in (\alpha_N, 2N)$ if and only if $N \in (N_0, \infty)$, for some $N_0 \in (1, 2)$; for $N \in (0, N_0)$, the solution is unique, whenever it exists. A possible way to tackle such a conjecture could be to show that for $N > 1$, the function $N \mapsto \alpha_N - 2N$ is monotonically nonincreasing in N , as it appears to be the case in our numerical study, see Fig. 4, right, and to exploit the fact that $\tilde{a}_N \rightarrow \infty$ as $N \rightarrow N_0$, $N > N_0$, see Fig. 4, left.

On the basis of our numerical results, we may also conjecture that for $N_0 < N < 10$ and $\alpha \in (\alpha_N, 2N)$, there exist exactly two radially symmetric solutions of (8). This conjecture is supported by the bifurcation analysis of Section 3.3 concerning the specific value $\alpha = N + 2 \in (\alpha_N, 2N)$ for $N > 2$ and $N \neq 4$. Note that for $N = 4$, $\alpha_N = N + 2$ should hold. As N increases, the curves $a \mapsto \alpha(a)$ appear to have more and more critical points. Thus, for suitable values of α , the number of solutions increases as N increases. We have already checked this fact in Theorem 1 for $\lambda = 2\pi\alpha$, $\alpha = N + 2$, but apparently it also holds for other values of α .

The last observations are concerned with the function $a \mapsto J_N(a)$ defined by (22). It seems that such a function always takes positive values on $(-\infty, c(N))$ and negative values on $(c(N), \infty)$, for some $c(N) > 0$. See Fig. 5, left. We may formulate this as the following conjecture: *There exists a function $N \mapsto c(N)$ on $(0, +\infty)$ such that $J_N(a) = 0$ if and only if $a = c(N)$ and $J_N(a) > 0$ if and only if $a < c(N)$.* See Fig. 5, right.




Fig. 5. *The function $a \mapsto J_N(a)$ for $N = 1, 3, 5, \dots, 11$ (left) and the curve $N \mapsto c(N)$, where, at N fixed, $c(N)$ is the first positive zero of $a \mapsto J_N(a)$; the dotted line corresponds to $N \mapsto \frac{1}{2} \log(2(N+2))$. These two curves are tangent at $N = 10 = N_3$ and $N = 28 = N_5$ (right).*

Quite interesting is the comparison of a_N^* with $c(N)$. By Proposition 10, we know that $j(k) := J_N(a_{N_k}^*) = 0$ if k is odd and $j(k) > 0$ if k is even. Recall that $\alpha'(a_N^*) = 0$ if and only if $N = N_k$. We observe numerically, and conjecture,

that: For any $N > 2$, $a_N^* \leq c(N)$, with equality if and only if $N = N_{2l+1}$ for some $l \geq 1$. This is observed numerically with a very high accuracy for $k = 3, 5, 7, 9, 11$, see Figs. 5, right, and also 6, left.

Summarizing the results of Section 3.2, we have shown that, in the interval $(c(N), \infty) \ni a$, the number of nodes of φ_a given by (17) increases as N grows each time a new critical point of α appears. This needs to be interpreted in terms of Morse index, which is still an open question.

To investigate whether a critical point of α is a local minimum, we may look at the functional

$$K_N(a) := \int_0^{+\infty} (1+r^2)^N e^{2u_a} (\psi_a + 2\varphi_a^2) r dr$$

where ψ_a solves the ordinary differential equation

$$\begin{cases} \psi_a'' + \frac{\psi_a'}{r} + 2(1+r^2)^N e^{2u_a} (\psi_a + 2\varphi_a^2) = 0, & r \in (0, +\infty), \\ \psi_a(0) = 0, & \psi_a'(0) = 0. \end{cases}$$

We have indeed $\alpha''(a) = 2K_N(a)$. No simple criterion for the positivity of $K_N(a)$ is known, but our numerical results at level $\alpha = N + 2$, see Fig. 6, right, combine very well with the results of Theorem 1 and the bifurcation diagrams shown in Fig. 1.




Fig. 6. Left: the curve $N \mapsto J_N(a_N^*)$ is nonnegative and achieves its minimum value, 0, (resp. local maxima) for $N = N_{2l+1}$, $l \geq 1$ (resp. $N = N_{2l}$). Right: the curve $N \mapsto K_N(a_N^*)$ changes sign, but is always nonnegative when $\alpha'(N) = 0$. When $N = N_{2l}$, $l \geq 1$, $K_N(a_N^*)$ is positive.

Acknowledgments. This work has been partially supported by the *Fondation Sciences Mathématiques de Paris*, by the project *IFO* of the French National Research Agency (ANR) and by the italian P.R.I.N. project *Variational Methods and Non Linear Differential Equation*, Italy. The third author wishes also to express her grat-

itude to Ceremade for the warm and kind hospitality during her visits. Figures have been computed and plotted with Mathematica™.

© 2008 by the authors. This paper may be reproduced, in its entirety, for non-commercial purposes.

References

- [1] D. BARTOLUCCI AND G. TARANTELLA, *Liouville type equations with singular data and their applications to periodic multivortices for the electroweak theory*, Comm. Math. Phys., 229 (2002), pp. 3–47.
- [2] E. CAGLIOTI, P.-L. LIONS, C. MARCHIORO, AND M. PULVIRENTI, *A special class of stationary flows for two-dimensional Euler equations: a statistical mechanics description*, Comm. Math. Phys., 143 (1992), pp. 501–525.
- [3] ———, *A special class of stationary flows for two-dimensional Euler equations: a statistical mechanics description. II*, Comm. Math. Phys., 174 (1995), pp. 229–260.
- [4] S. CHANILLO AND M. KIESSLING, *Rotational symmetry of solutions of some nonlinear problems in statistical mechanics and in geometry*, Comm. Math. Phys., 160 (1994), pp. 217–238.
- [5] C.-C. CHEN AND C.-S. LIN, *Sharp estimates for solutions of multi-bubbles in compact Riemann surfaces*, Comm. Pure Appl. Math., 55 (2002), pp. 728–771.
- [6] ———, *Topological degree for a mean field equation on Riemann surfaces*, Comm. Pure Appl. Math., 56 (2003), pp. 1667–1727.
- [7] C.-C. CHEN, C.-S. LIN, AND G. WANG, *Concentration phenomena of two-vortex solutions in a Chern-Simons model*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 3 (2004), pp. 367–397.
- [8] W. X. CHEN AND C. LI, *Classification of solutions of some nonlinear elliptic equations*, Duke Math. J., 63 (1991), pp. 615–622.
- [9] ———, *Qualitative properties of solutions to some nonlinear elliptic equations in \mathbb{R}^2* , Duke Math. J., 71 (1993), pp. 427–439.
- [10] ———, *What kinds of singular surfaces can admit constant curvature ?*, Duke Math. J., 78 (1995), pp. 437–451.
- [11] K.-S. CHENG AND C.-S. LIN, *On the conformal Gaussian curvature equation in \mathbb{R}^2* , J. Differential Equations, 146 (1998), pp. 226–250.
- [12] M. G. CRANDALL AND P. H. RABINOWITZ, *Bifurcation from simple eigenvalues*, J. Functional Analysis, 8 (1971), pp. 321–340.

- [13] Z. DJADLI, *Existence result for the mean field problem on Riemann surfaces of all genres*, Commun. Contemp. Math., 10 (2008), pp. 205–220.
- [14] A. EREMENKO, *Metrics of positive curvature with conic singularities on the sphere*, Proc. Amer. Math. Soc., 132 (2004), pp. 3349–3355 (electronic).
- [15] J. A. GAUNT, *The triplets of helium. Appendix*, Phil. Trans. R. Soc. London (A), 228 (1929), pp. 192–196.
- [16] A. JAFFE AND C. TAUBES, *Vortices and monopoles*, vol. 2 of Progress in Physics, Birkhäuser Boston, Mass., 1980. Structure of static gauge theories.
- [17] J. L. KAZDAN, *Prescribing the curvature of a Riemannian manifold*, vol. 57 of CBMS Regional Conference Series in Mathematics, Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1985.
- [18] J. L. KAZDAN AND F. W. WARNER, *Curvature functions for open 2-manifolds*, Ann. of Math. (2), 99 (1974), pp. 203–219.
- [19] ———, *Existence and conformal deformation of metrics with prescribed Gaussian and scalar curvatures*, Ann. of Math. (2), 101 (1975), pp. 317–331.
- [20] M. K.-H. KIESSLING, *Statistical mechanics of classical particles with logarithmic interactions*, Comm. Pure Appl. Math., 46 (1993), pp. 27–56.
- [21] ———, *Statistical mechanics approach to some problems in conformal geometry*, Phys. A, 279 (2000), pp. 353–368. Statistical mechanics: from rigorous results to applications.
- [22] L. LANDAU AND E. LIFSCHITZ, *Physique théorique. Tome III: Mécanique quantique. Théorie non relativiste (French)*, Deuxième édition. Translated from russian by E. Gloukhian. Éditions Mir, Moscow, 1967.
- [23] Y. Y. LI, *Harnack type inequality: the method of moving planes*, Comm. Math. Phys., 200 (1999), pp. 421–444.
- [24] C.-S. LIN, *Uniqueness of conformal metrics with prescribed total curvature in \mathbb{R}^2* , Calc. Var. Partial Differential Equations, 10 (2000), pp. 291–319.
- [25] ———, *Uniqueness of solutions to the mean field equations for the spherical Onsager vortex*, Arch. Ration. Mech. Anal., 153 (2000), pp. 153–176.
- [26] C.-S. LIN AND C. L. WANG, *Elliptic functions, Green functions and mean field equations on torus*. Preprint, 2006.
- [27] F. LUO AND G. TIAN, *Liouville equation and spherical convex polytopes*, Proc. Amer. Math. Soc., 116 (1992), pp. 1119–1129.
- [28] A. MALCHIODI, *Morse theory and a scalar mean field equation on compact surfaces*. Preprint, 2007.
- [29] J. PRAJAPAT AND G. TARANTELLA, *On a class of elliptic problems in \mathbb{R}^2 : symmetry and uniqueness results*, Proc. Roy. Soc. Edinburgh Sect. A, 131 (2001), pp. 967–985.

- [30] P. H. RABINOWITZ, *Some aspects of nonlinear eigenvalue problems*, Rocky Mountain J. Math., 3 (1973), pp. 161–202. Rocky Mountain Consortium Symposium on Nonlinear Eigenvalue Problems (Santa Fe, N.M., 1971).
- [31] G. TARANTELLA, *Analytical aspects of Liouville-type equations with singular sources*, in Stationary partial differential equations. Vol. I, Handb. Differ. Equ., North-Holland, Amsterdam, 2004, pp. 491–592.
- [32] ———, *Selfdual gauge field vortices: An analytical approach*, Progress in Nonlinear Differential Equations and their Applications, 72, Birkhäuser Boston Inc., Boston, MA, 2008.
- [33] M. TROYANOV, *Metrics of constant curvature on a sphere with two conical singularities*, in Differential geometry (Peñíscola, 1988), vol. 1410 of Lecture Notes in Math., Springer, Berlin, 1989, pp. 296–306.
- [34] ———, *Prescribing curvature on compact surfaces with conical singularities*, Trans. Amer. Math. Soc., 324 (1991), pp. 793–821.
- [35] Y. YANG, *Solitons in field theory and nonlinear analysis*, Springer Monographs in Mathematics, Springer-Verlag New York, 2001.