



HAL
open science

Powers of sequences and convergence of ergodic averages

Nikos Frantzikinakis, Michael Johnson, Emmanuel Lesigne, Mate Wierdl

► **To cite this version:**

Nikos Frantzikinakis, Michael Johnson, Emmanuel Lesigne, Mate Wierdl. Powers of sequences and convergence of ergodic averages. 2008. hal-00327604v1

HAL Id: hal-00327604

<https://hal.science/hal-00327604v1>

Preprint submitted on 8 Oct 2008 (v1), last revised 29 Jun 2009 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POWERS OF SEQUENCES AND CONVERGENCE OF ERGODIC AVERAGES

N. FRANTZIKINAKIS, M. JOHNSON, E. LESIGNE, AND M. WIERDL

ABSTRACT. A sequence (s_n) of integers is good for the mean ergodic theorem if for each invertible measure preserving system (X, \mathcal{B}, μ, T) and any bounded measurable function f , the averages $\frac{1}{N} \sum_{n=1}^N f(T^{s_n} x)$ converge in the L^2 norm. We construct a sequence (s_n) that is good for the mean ergodic theorem, but the sequence (s_n^2) is not. Furthermore, we show that for any set of bad exponents B , there is a sequence (s_n) where (s_n^k) is good for the mean ergodic theorem exactly when k is not in B . We then extend this result to multiple ergodic averages of the form $\frac{1}{N} \sum_{n=1}^N f_1(T^{s_n} x) f_2(T^{2s_n} x) \dots f_\ell(T^{\ell s_n} x)$. We also prove a similar result for pointwise convergence of single ergodic averages.

CONTENTS

1.	Introduction	1
2.	Single mean convergence	5
3.	Multiple mean convergence	10
4.	Pointwise convergence	12
5.	Appendix	17
	References	23

1. INTRODUCTION

1.1. Main result. It is well known that for any fixed positive integer k the sequence $1^k, 2^k, 3^k \dots$ is good for the mean ergodic theorem. This means that for every measure preserving system and function f in $L^2(\mu)$, the averages

$$(1.1) \quad \frac{1}{N} \sum_{n=1}^N f(T^{n^k} x)$$

converge in L^2 norm as $N \rightarrow \infty$. Using the spectral theorem for unitary operators, this is equivalent to the convergence of the averages

$$(1.2) \quad \frac{1}{N} \sum_{n=1}^N e^{2\pi i n^k \alpha}$$

as $N \rightarrow \infty$ for any real number α .

Date: October 8, 2008.

2000 Mathematics Subject Classification. Primary: 37A30; Secondary: 28D05.

Key words and phrases. ergodic theorems, ergodic averages, multiple ergodic averages.

The first and last authors were supported by NSF grants DMS-0701027 and DMS-0801316.

An illustrative question for our paper is the following. Is there a sequence $s_1, s_2, s_3 \dots$ of positive integers so that the averages

$$(1.3) \quad \frac{1}{N} \sum_{n=1}^N e^{2\pi i s_n \alpha}$$

converge for any real number α , but for some α , the averages

$$(1.4) \quad \frac{1}{N} \sum_{n=1}^N e^{2\pi i s_n^2 \alpha}$$

do not converge as $N \rightarrow \infty$? In other words, we ask if there is a sequence $s_1, s_2, s_3 \dots$ of positive integers which is good for the mean ergodic theorem, but the sequence of squares $s_1^2, s_2^2, s_3^2 \dots$ of the sequence is not good for the mean ergodic theorem?

Similarly, we can ask: is there a sequence $s_1, s_2, s_3 \dots$ of positive integers which is *not* good for the mean ergodic theorem, but the sequence of squares $s_1^2, s_2^2, s_3^2 \dots$ of the sequence is good for the mean ergodic theorem?

Perhaps surprisingly, the answer to *both* questions is *yes*, indicating that the convergence properties of positive powers of a sequence are independent of those of the original sequence. In fact, in this paper we prove the following result showing the total independence of powers of a sequence for the mean ergodic theorem.

Theorem A. *Let B be an arbitrary set of positive integers. Then there exists an increasing sequence $s_1, s_2, s_3 \dots$ of positive integers such that*

- *The sequence $s_1^g, s_2^g, s_3^g \dots$ is good for the mean ergodic theorem for any “good” exponent $g \in \mathbb{N} \setminus B$.*
- *The sequence $s_1^b, s_2^b, s_3^b \dots$ is not good for the mean ergodic theorem for any “bad” exponent $b \in B$.*

Using the spectral theorem for unitary operators, we get the following equivalent formulation of our theorem:

Theorem A’. *Let B be an arbitrary set of positive integers. Then there exists an increasing sequence $s_1, s_2, s_3 \dots$ of positive integers such that*

- *For $g \in \mathbb{N} \setminus B$, the averages*

$$(1.5) \quad \frac{1}{N} \sum_{n=1}^N e^{2\pi i s_n^g \alpha}$$

converge as $N \rightarrow \infty$ for any real number α .

- *For $b \in B$, there exists a real number α such that the averages*

$$(1.6) \quad \frac{1}{N} \sum_{n=1}^N e^{2\pi i s_n^b \alpha}$$

do not converge as $N \rightarrow \infty$.

Similar results related to issues of recurrence were proved in [FrLW2]. The original motivation to search for results which express the independence of powers of a sequence for various properties, comes from the papers of Deshouillers, Erdős, Sárközy ([DES]) and Deshouillers, Fouvry ([DFo]). In these papers, the authors prove results analogous to ours but for bases of the positive integers.

In our paper, we generalize Theorem A to multiple ergodic averages and prove a version for pointwise convergence of single ergodic averages. We state these generalizations in the next subsection, where we also give precise definitions of the concepts used throughout the paper.

1.2. Definitions and generalizations. All along the article we will use the word *system* or the term *measure preserving system*, to designate a quadruple (X, \mathcal{B}, μ, T) , where T is an *invertible* measure preserving transformation of a probability space (X, \mathcal{B}, μ) . By insisting that our system is invertible, we make sure that T^k is well defined for negative integers k .

We give a strengthening of Theorem A that deals with problems related to convergence of multiple ergodic averages.

Definition 1.1. Let ℓ be a positive integer, (s_n) be a sequence of integers, and (X, \mathcal{B}, μ, T) be a system.

We say that the sequence of integers (s_n) is *good for ℓ -convergence for the system* (X, \mathcal{B}, μ, T) if for any bounded, measurable functions f_1, f_2, \dots, f_ℓ , the averages

$$\frac{1}{N} \sum_{n=1}^N f_1(T^{s_n}x) \cdot f_2(T^{2s_n}x) \cdot \dots \cdot f_\ell(T^{\ell s_n}x)$$

converge in L^2 norm as $N \rightarrow \infty$.

We say the sequence (s_n) is *universally good for ℓ -convergence* if it is good for ℓ -convergence for any system (X, \mathcal{B}, μ, T) . We often abbreviate this by saying that “ (s_n) is good for ℓ -convergence” and refer to the case $\ell = 1$ as *single mean convergence*.

We refer the reader to [RW] for examples of sequences that are good for single mean convergence. Examples of sequences that are good for ℓ -convergence for every $\ell \in \mathbb{N}$, include $s_n = n$, shown by Host and Kra ([HKr1]) and later by Ziegler ([Z]), as well as $s_n = p(n)$ where $p(n)$ is an integer polynomial, as shown by Host and Kra ([HKr2]).

Theorem 1.2. *Let B be an arbitrary set of positive integers and $\ell \in \mathbb{N}$. Then there exists an increasing sequence (s_n) of positive integers such that*

- *For every $g \in \mathbb{N} \setminus B$, the sequence (s_n^g) is good for ℓ -convergence.*
- *For every $b \in B$, the sequence (s_n^b) is not good for ℓ -convergence.*

We also give a strengthening of Theorem A that deals with problems related to pointwise convergence of ergodic averages.

Definition 1.3. Let (s_n) be a sequence of integers, and let (X, \mathcal{B}, μ, T) be a system.

We say that the sequence (s_n) is *good for the pointwise ergodic theorem for the system* (X, \mathcal{B}, μ, T) if for any $f \in L^2$, the averages

$$\frac{1}{N} \sum_{n=1}^N f(T^{s_n}x)$$

converge as $N \rightarrow \infty$ for almost every x .

We say the sequence (s_n) is *universally good for the pointwise ergodic theorem* if it is good for the pointwise ergodic theorem for any system (X, \mathcal{B}, μ, T) . We often abbreviate this by saying that “ (s_n) is good for the pointwise ergodic theorem.”

Notice that any sequence that is good for the pointwise ergodic theorem is also good for the mean ergodic theorem. Various examples of sequences that are good for the pointwise ergodic theorem are known (see for example [RW]). A particular example that will be used later, is the case of a sequence $(p(n))$ where p is any polynomial with integer coefficients. This case was treated by Bourgain in [Bou].

Theorem 1.4. *Let B be an arbitrary set of positive integers. Then there exists an increasing sequence (s_n) of positive integers such that*

- *For every $g \in \mathbb{N} \setminus B$, the sequence (s_n^g) is good for the pointwise ergodic theorem.*
- *For every $b \in B$, the sequence (s_n^b) is not good for the mean ergodic theorem.*

1.3. Further remarks and conjectures. If for a given $\ell \in \mathbb{N}$, the sequence (s_n) is good for ℓ -convergence, then, as Theorem 1.2 shows, we cannot assert in general that any of its powers (s_n^k) , where $k \geq 2$, is good for ℓ -convergence. In direct contrast with this, we expect the following result to be true:

Conjecture 1. *Suppose that the sequence of integers (s_n) is good for ℓ -convergence for every $\ell \in \mathbb{N}$. Then for every $k \in \mathbb{N}$, the sequence (s_n^k) is good for ℓ -convergence for every $\ell \in \mathbb{N}$.*

To support this conjecture, let us mention that if the sequence (s_n) is good for 2-convergence then the sequence (s_n^2) is good for single mean convergence (see Lemma 3.1 below). In fact, Conjecture 1 would be true if the following more general statement holds:

Conjecture 2. *Suppose that the sequence of integers (s_n) is good for mean $(k\ell)$ -convergence. Then the sequence (s_n^k) is good for ℓ -convergence.*

It is easy to verify this conjecture for $\ell = 1$. The reason for that, is because the spectral theorem for unitary operators gives convenient necessary and sufficient conditions for single mean convergence. We currently do not have such a convenient characterization for ℓ -convergence when $\ell \geq 2$. The following conjecture would fill this gap if true:

Conjecture 3. *Let (s_n) be a sequence of integers. The following three statements are equivalent:*

- *The sequence (s_n) is good for ℓ -convergence.*
- *The sequence (s_n) is good for ℓ -convergence for every ℓ -step nilsystem.¹*
- *For every ℓ -step nilmanifold $X = G/\Gamma$, every $a \in G$, and $f \in C(X)$, the sequence $(\frac{1}{N} \sum_{n=1}^N f(a^{s_n} \Gamma))$ converges as $N \rightarrow +\infty$.*

We are mainly interested in knowing if the third (or second) condition implies the first. In the case that the set $S = \{s_n, n \in \mathbb{N}\}$ has positive upper density, this implication follows immediately from the nilsequence decomposition result of Bergelson, Host, and Kra in [BeHKr].

For general sequences (s_n) , using the spectral theorem for unitary operators, we can easily verify Conjecture 3 for $\ell = 1$. It is possible to see that if Conjecture 3 is true then Conjecture 2 is also true (and hence Conjecture 1).

¹If G is an ℓ -step nilpotent Lie group and Γ is a discrete cocompact subgroup, then the homogeneous space $X = G/\Gamma$ is called an ℓ -step nilmanifold. If $T_a(g\Gamma) = (ag)\Gamma$ for some $a \in G$, \mathcal{G}/Γ is the Borel σ -algebra of X , and m is the Haar measure on X , then the system $(X, \mathcal{G}/\Gamma, m, T_a)$ is called an ℓ -step nilsystem.

Notation: The following notation will be used throughout the article: $Tf = f \circ T$, $e(t) = e^{2\pi it}$.

Acknowledgment. The authors wish to thank the Mathematical Sciences Research Institute for providing partial support to complete work on the present article during its special semester program on Ergodic Theory and Additive Combinatorics.

2. SINGLE MEAN CONVERGENCE

In this section we shall prove Theorem A. This will help us illustrate the main ideas behind the more complicated arguments we will use in the proof of Theorem 1.2 and of Theorem 1.4 (which both extend Theorem A).

We shall prove our theorem in its equivalent formulation of Theorem A'.

Let B be a fixed set of positive integers (possibly empty), and let α be a fixed irrational number. It is sufficient to find a sequence (s_n) satisfying the following two conditions

(s1) For every $b \in B$, the sequence $\left(\frac{1}{N} \sum_{n=1}^N e(s_n^b \alpha)\right)$ diverges.

(s2) For every $g \in \mathbb{N} \setminus B$ and every $\beta \in \mathbb{R}$, the sequence $\left(\frac{1}{N} \sum_{n=1}^N e(s_n^g \beta)\right)$ converges.

The rest of this section will be devoted to the construction of a sequence (s_n) that satisfies conditions (s1) and (s2).

2.1. Definition of the sequence (s_n) . We denote

$$(2.1) \quad I_+ = \left\{x \in \mathbb{T} \mid \cos(2\pi x) \geq \sqrt{2}/2\right\} \quad \text{and} \quad I_- = \left\{x \in \mathbb{T} \mid \cos(2\pi x) \leq -\sqrt{2}/2\right\}.$$

These are intervals of length $1/4$ on the torus. The sequence (s_n) consists of the elements of a set S , taken in increasing order, that is defined as follows:

$$(2.2) \quad S = \bigcup_{j \geq 1} \left\{n \in \mathbb{N} \mid [2^{2j-1} \leq n < 2^{2j}, n^{b_j} \alpha \in I_+] \text{ or } [2^{2j} \leq n < 2^{2j+1}, n^{b_j} \alpha \in I_-]\right\}$$

for some appropriately chosen sequence (b_j) of elements of B . We will construct a sequence (b_j) so that:

- Every element of B appears infinitely often in the sequence (b_j) . This easily implies that condition (s1) holds.
- The first appearance of elements of B in the sequence (b_j) happens late enough (with respect to j) to guarantee that certain equidistribution properties are satisfied. All unspecified elements b_j will be set to be equal to some fixed element of B . This will enable us to verify condition (s2).

Let us now state explicitly the properties that the sequence (b_j) is going to satisfy:

- (b1) We have $b_j \in B$, and all elements of B appear infinitely often in the sequence (b_j) .
- (b2) We have

$$\lim_{j \rightarrow \infty} \sup_{N > 2^{2j-1}} \left| \frac{1}{N} \sum_{n=1}^N \mathbb{1}_{I_{\pm}}(n^{b_j} \alpha) - \frac{1}{4} \right| = 0.$$

(b3) For every nonzero $l \in \mathbb{Z}$, $\beta \in \mathbb{R}$, and $g \in G$, we have

$$\lim_{j \rightarrow \infty} \sup_{N \geq 2^{2^j-1}} \left| \frac{1}{N} \sum_{n=1}^N e(ln^{b_j} \alpha + n^g \beta) \right| = 0.$$

In the next subsection we construct such a sequence (b_j) .

2.2. Construction of the sequence (b_j) . The following lemma will be essential for our construction:

Lemma 2.1. *Let $b, g \in \mathbb{N}$ with $b > g$ and $\alpha \in \mathbb{R} \setminus \mathbb{Q}$. Then*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mathbb{1}_{I_{\pm}}(n^b \alpha) = \frac{1}{4}, \text{ and}$$

$$\lim_{N \rightarrow \infty} \sup_{\beta \in \mathbb{R}} \left| \frac{1}{N} \sum_{n=1}^N e(n^b \alpha + n^g \beta) \right| = 0.$$

Proof. The first part follows from Weyl's equidistribution theorem. The second part is a direct consequence of van der Corput's classical inequality. Applying it g times, we obtain an estimate of the trigonometric sums in which the number β does not appear anymore. \square

Note that, for each fixed b only finitely many g 's are involved in the lemma. Therefore, the convergence is also uniform in g .

Proposition 2.2. *There exists a sequence (b_j) that satisfies conditions (b1), (b2), and (b3) of Section 2.1.*

Proof. Using Lemma 2.1, we get that for every $b \in \mathbb{N}$, nonzero $l \in \mathbb{Z}$, and $\varepsilon > 0$, there exists $J = J(b, l, \varepsilon) \geq 1$ that satisfies

$$(2.3) \quad \sup_{N \geq 2^{2^J-1}} \left| \frac{1}{N} \sum_{n=1}^N \mathbb{1}_{I_{\pm}}(n^b \alpha) - \frac{1}{4} \right| \leq \varepsilon;$$

and such that

$$(2.4) \quad \sup_{N \geq 2^{2^J-1}} \sup_{\beta \in \mathbb{R}} \sup_{g \in \mathbb{N}, g < b} \left| \frac{1}{N} \sum_{n=1}^N e(ln^b \alpha + n^g \beta) \right| \leq \varepsilon.$$

Furthermore, we can assume that $J = J(b, l, \varepsilon)$ is increasing with respect to the variables b, l , and decreasing with respect to the variable ε .

We write $B = \{a_t, t \in \mathbb{N}\}$ where $a_1 < a_2 < \dots$. We construct the sequence (b_j) using the following guidelines: (i) every integer a_t appears infinitely often in the range of (b_j) , (ii) the first appearance of a_t in (b_j) happens at a time j greater than $J(a_t, t, 1/t)$, and (iii) all values of b_j that are left unspecified are set to be equal to a_1 . Notice that condition (ii) guarantees that (2.3) and (2.4) hold for $b = b_j$ and $J = j$.

The explicit construction goes as follows:

Define $J_1 = J(a_1, 1, 1)$ and $b_{J_1} = a_1$.

Define $J_2 = \max\{J(a_2, 2, 1/2), J_1 + 2\}$ and $b_{J_2} = a_2, b_{J_2+1} = a_1$.

By induction, define $J_t = \max\{J(a_t, t, 1/t), J_{t-1} + t\}$

and $b_{J_t} = a_t, b_{J_t+1} = a_{t-1}, \dots, b_{J_t+t-1} = a_1$.

We claim that this sequence (b_j) has the advertised properties. First, we note that every integer a_t appears infinitely many times in the range of (b_j) , so condition (b1) holds. Now let $\beta \in \mathbb{R}$, $g \in G$, and $l \in \mathbb{N}$. Choose j large enough so that $J_t \leq j < J_{t+1}$ for some $t \geq l$. Then, by construction, we have $b_j \in \{a_1, a_2, \dots, a_t\}$. If $N \geq 2^{2^{j-1}}$ then $N \geq 2^{2^{J_t-1}}$, and $J_t \geq J(a_k, l, 1/t)$ for all k between 1 and t . It follows from (2.3) that

$$\sup_{N \geq 2^{2^{j-1}}} \left| \frac{1}{N} \sum_{n=1}^N \mathbb{1}_{I_{\pm}}(n^{b_j} \alpha) - \frac{1}{4} \right| \leq \frac{1}{t},$$

and so condition (b2) holds. Furthermore, it follows from (2.4) that for every b_j greater than g we have

$$(2.5) \quad \sup_{N \geq 2^{2^{j-1}}} \left| \frac{1}{N} \sum_{n=1}^N e(ln^{b_j} \alpha + n^g \beta) \right| \leq \frac{1}{t}.$$

It remains to deal with those b_j that are less than g . Since there are only finitely many values of the sequence (b_j) that are less than g , by Weyl's equidistribution theorem we have

$$\lim_{N \rightarrow \infty} \sup_{j \in \mathbb{N}, b_j < g} \left| \frac{1}{N} \sum_{n=1}^N e(ln^{b_j} \alpha + n^g \beta) \right| = 0.$$

Combining this with (2.5) easily implies that condition (b3) also holds, completing the proof. \square

2.3. The sequence (s_n) satisfies conditions (s1) and (s2). The goal of this section is complete the proof of Theorem A' by proving the following proposition which will

Proposition 2.3. *Suppose that the sequence (b_j) satisfies conditions (b1), (b2), and (b3) of Section 2.2. Then the sequence (s_n) defined by (2.2) satisfies conditions (s1) and (s2).*

First we show that the set S in (2.2) has positive density.

Lemma 2.4. *Suppose that the sequence (b_j) satisfies condition (b2) of Section 2.1. Then the set S in (2.2) has density 1/4.*

Proof. This is an easy consequence of Lemma 5.3 in the Appendix. \square

Since S has positive density, condition (s1) is equivalent to

$$(2.6) \quad \text{For every } b \in B, \text{ the sequence } \left(\frac{1}{N} \sum_{n=1}^N \mathbb{1}_S(n) e(n^b \alpha) \right) \text{ diverges,}$$

and condition (s2) is equivalent to

$$(2.7) \quad \text{For every } g \in \mathbb{N} \setminus B \text{ and all } \beta \in \mathbb{R}, \left(\frac{1}{N} \sum_{n=1}^N \mathbb{1}_S(n) e(n^g \beta) \right) \text{ converges.}$$

We first show that the conditions imposed on the sequence (b_j) guarantee that condition (2.6) (and as a result condition (s1)) is satisfied by the set S .

Proposition 2.5. *Suppose that the sequence (b_j) satisfies conditions (b1) and (b2) of Section 2.1. Then the sequence (s_n) defined by (2.2) satisfies condition (s1).*

Proof. Fix $b \in B$. By condition (b1), there are arbitrarily large values of j for which $b_j = b$. We fix such a j . We have

$$\begin{aligned} & \frac{1}{2^{2j}} \sum_{n=1}^{2^{2j}} \mathbb{1}_S(n) \cos(2\pi n^b \alpha) - \frac{1}{2^{2j+1}} \sum_{n=1}^{2^{2j+1}} \mathbb{1}_S(n) \cos(2\pi n^b \alpha) = \\ & \frac{1}{2^{2j+1}} \sum_{n=1}^{2^{2j}} \mathbb{1}_S(n) \cos(2\pi n^b \alpha) - \frac{1}{2^{2j+1}} \sum_{n=2^{2j}+1}^{2^{2j+1}} \mathbb{1}_S(n) \cos(2\pi n^b \alpha) \geq \\ & \frac{1}{2^{2j+1}} \sum_{n=1}^{2^{2j-1}} \mathbb{1}_S(n)(-1) + \frac{1}{2^{2j+1}} \sum_{n=2^{2j-1}+1}^{2^{2j}} \mathbb{1}_S(n)(\sqrt{2}/2) - \frac{1}{2^{2j+1}} \sum_{n=2^{2j}+1}^{2^{2j+1}} \mathbb{1}_S(n)(-\sqrt{2}/2). \end{aligned}$$

Using condition (b2) and Lemma 2.4, we see that, for large j , the last quantity is near $-\frac{1}{16} + \frac{\sqrt{2}}{32} + \frac{\sqrt{2}}{16}$, which is positive. This shows that (2.6) holds. Since the set S has positive density (Lemma 2.4), condition (s1) is also satisfied. \square

Next we show that the conditions imposed on the sequence (b_j) guarantee that condition (2.7) is satisfied. We first need a couple of lemmas.

Lemma 2.6. *Let l, m be two nonzero integers, $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, and suppose that the sequence (b_j) satisfies condition (b3) of Section 2.1. Let us define a sequence (e_n) by*

$$e_n = \begin{cases} e(ln^{b_j} \alpha), & n \in [2^{2j-1}, 2^{2j}); \\ e(mn^{b_j} \alpha), & n \in [2^{2j}, 2^{2j+1}). \end{cases}$$

Then for every $\beta \in \mathbb{R}$ and $g \in \mathbb{N} \setminus B$, we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N e_n e(n^g \beta) = 0.$$

Proof. Fix $\beta \in \mathbb{R}$ and $g \in \mathbb{N} \setminus B$. By condition (b3) we have for every nonzero integer k that

$$\lim_{j \rightarrow \infty} \sup_{N > 2^{2j-1}} \left| \frac{1}{N} \sum_{n=1}^N e(kn^{b_j} \alpha + n^g \beta) \right| = 0.$$

Applying this for $k = l$, and $k = m$, and using Lemma 5.3 in the Appendix we easily get the advertised result. \square

Given a sequence (b_j) of positive integers and functions $\phi, \psi: \mathbb{T} \rightarrow \mathbb{C}$, define the sequence

$$(2.8) \quad f_n(\phi, \psi) = \begin{cases} \phi(n^{b_j} \alpha), & n \in [2^{2j-1}, 2^{2j}); \\ \psi(n^{b_j} \alpha), & n \in [2^{2j}, 2^{2j+1}). \end{cases}$$

Lemma 2.7. *Let ϕ_0, ψ_0 be two Riemann integrable functions on the torus with zero integral, $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, and suppose that the sequence (b_j) satisfies condition (b3) of Section 2.1. Then for every $\beta \in \mathbb{R}$ and $g \in \mathbb{N} \setminus B$, we have*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f_n(\phi_0, \psi_0) e(n^g \beta) = 0.$$

Proof. By Lemma 2.6, the result is true when ϕ and ψ are trigonometric polynomials. To complete the proof, we use the following standard approximation argument.

Let ϕ_0, ψ_0 be Riemann integrable functions. It suffices to show that for every $\varepsilon > 0$ there exist trigonometric polynomials ϕ and ψ with zero integral such that

$$(2.9) \quad \limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N |f_n(\phi_0, \psi_0) - f_n(\phi, \psi)| \leq \varepsilon.$$

We can approximate in $L^1(\mathbb{T})$ the function ϕ_0 by a trigonometric polynomial. After composing with a translation this trigonometric polynomial will approximate the function ψ_0 as well. So there exist two trigonometric polynomials ϕ and ψ , with zero integral, such that $\int_{\mathbb{T}} |\phi - \phi_0| = \int_{\mathbb{T}} |\psi - \psi_0| = \theta \leq \varepsilon$. Notice that

$$(2.10) \quad |f_n(\phi_0, \psi_0) - f_n(\phi, \psi)| = f_n(|\phi - \phi_0|, |\psi - \psi_0|).$$

So in order to establish (2.9) it suffices to show that

$$(2.11) \quad \limsup_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f_n(|\phi - \phi_0|, |\psi - \psi_0|) \leq \varepsilon.$$

Let $\Phi_0 = |\phi - \phi_0|$ and $\Psi_0 = |\psi - \psi_0|$. Since both Φ_0 and Ψ_0 are Riemann integrable and have integral θ , the following holds: For every $\delta > 0$ it is possible to find four continuous functions $\phi_1, \phi_2, \psi_1, \psi_2$, with zero integral, such that

$$\phi_1 + \theta - \delta \leq \Phi_0 \leq \phi_2 + \theta + \delta \quad \text{and} \quad \psi_1 + \theta - \delta \leq \Psi_0 \leq \psi_2 + \theta + \delta.$$

It follows that

$$(2.12) \quad f_n(\phi_1, \psi_1) + \theta - \delta \leq f_n(\Phi_0, \Psi_0) \leq f_n(\phi_2, \psi_2) + \theta + \delta.$$

Moreover, since

$$(2.13) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f_n(\phi, \psi) = 0$$

for every trigonometric polynomials ϕ and ψ with zero integral, by uniform approximation, this remains true if ϕ and ψ are continuous functions on the torus, with zero integral. Thus, we deduce from (2.12) and (2.13) (applied to $\phi = \phi_i, \psi = \psi_i$ for $i = 1, 2$), and the fact that δ was arbitrary, that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f_n(\Phi_0, \Psi_0) = \theta \leq \varepsilon.$$

Hence, (2.11) is established, completing the proof. □

Proposition 2.8. *Suppose that the sequence (b_j) satisfies conditions (b2) and (b3) of Section 2.1. Then the sequence (s_n) defined by (2.2) satisfies condition (s2).*

Proof. We apply Lemma 2.7 to the functions $\phi = \mathbb{1}_{I_+} - 1/4$ and $\psi = \mathbb{1}_{I_-} - 1/4$. We get for every $\beta \in \mathbb{R}$ and $g \in \mathbb{N} \setminus B$ that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N (\mathbb{1}_S(n) - 1/4)e(n^g \beta) = 0.$$

Hence, condition (2.7) is satisfied. Since the set S has positive density (Lemma 2.4), condition (s2) is also satisfied. □

Combining Propositions 2.5 and 2.8, we have Proposition 2.3.

3. MULTIPLE MEAN CONVERGENCE

In this section we shall prove Theorem 1.2. The argument is similar to the one used to prove Theorem A, but there are some extra complications since our analysis relies on some deeper multiple convergence results. To avoid repetition, we do not give details of proofs that can be easily extracted using arguments of the previous section.

3.1. A reduction. We need one preliminary result that was proved in [FrLW1] in the special case where the polynomial p is a monomial. A very similar argument gives the following more general result:

Lemma 3.1. *Suppose that the sequence (s_n) is good for ℓ -convergence. Then for every polynomial $p \in \mathbb{R}[t]$ with $\deg p \leq \ell$ the sequence $(p(s_n))$ has an asymptotic distribution mod 1.*

Example 1. Let us illustrate how one proves Lemma 3.1 in the case where $\ell = 2$. Suppose that the sequence (s_n) is good for 2-convergence. Let $p(t) = 2\alpha t^2 + \beta t + \gamma$ for some $\alpha, \beta, \gamma \in \mathbb{R}$. We define the transformation $R: \mathbb{T}^2 \rightarrow \mathbb{T}^2$ by

$$R(t_1, t_2) = (t_1 + \alpha, t_2 + 2t_1 + \alpha, t_3 + \beta),$$

and for $k \in \mathbb{Z}$ the functions

$$f_1(t_1, t_2, t_3) = e(k(-2t_2 + t_3)), \quad f_2(t_1, t_2, t_3) = e(kt_2).$$

An easy computation shows that

$$R^n(t_1, t_2) = (t_1 + n\alpha, t_2 + 2nt_1 + n^2\alpha, t_3 + n\beta).$$

As a consequence, the averages

$$\frac{1}{N} \sum_{n=1}^N R^{s_n} f_1 \cdot R^{2s_n} f_2 = e(k(t_3 - t_2)) \cdot \frac{1}{N} \sum_{n=1}^N e(k(2s_n^2\alpha + s_n\beta))$$

converge as $N \rightarrow \infty$. It follows that the sequence $(p(s_n))$ has an asymptotic distribution mod 1.

Using the previous lemma, it is easy to deduce Theorem 1.2 from the following result that we will set to prove:

Proposition 3.2. *Let B be an arbitrary set of positive integers, $\ell \in \mathbb{N}$, and $\alpha \in \mathbb{R} \setminus \mathbb{Q}$. Then there exists an increasing sequence (s_n) of integers such that*

- (s1) *For every $b \in B$, the sequence $\left(\frac{1}{N} \sum_{n=1}^N e((s_n^{\ell b} + s_n^b)\alpha)\right)$ diverges.*
- (s2) *For every $g \in \mathbb{N} \setminus B$, system (X, \mathcal{B}, μ, T) , and functions $f_1, \dots, f_\ell \in L^\infty(\mu)$, the sequence $\left(\frac{1}{N} \sum_{n=1}^N T^{s_n^g} f_1 \cdot T^{2s_n^g} f_2 \cdot \dots \cdot T^{\ell s_n^g} f_\ell\right)$ converges in $L^2(\mu)$.*

The rest of this section will be devoted to the construction of a sequence (s_n) that satisfies conditions (s1) and (s2).

3.2. Definition of the sequence (s_n) . Let α be any irrational number and I_+ and I_- be the intervals defined in (2.1). The sequence (s_n) consists of the elements of a set S , taken in increasing order, that is defined as follows:

$$(3.1) \quad S = \bigcup_{j \geq 1} \left\{ n \in \mathbb{N} \mid [2^{2j-1} \leq n < 2^{2j}, (n^{\ell b_j} + n^{b_j})\alpha \in I_+] \text{ or } [2^{2j} \leq n < 2^{2j+1}, (n^{\ell b_j} + n^{b_j})\alpha \in I_-] \right\}$$

where the sequence (b_j) satisfies the following conditions

- (b1) We have $b_j \in B$, and all elements of B appear infinitely often in the sequence (b_j) .
- (b2) We have

$$\lim_{j \rightarrow \infty} \sup_{N > 2^{2j-1}} \left| \frac{1}{N} \sum_{n=1}^N \mathbb{1}_{I_{\pm}}(n^{b_j} \alpha) - \frac{1}{4} \right| = 0.$$

- (b3) For every system (X, \mathcal{B}, μ, T) , functions $f_1, \dots, f_{\ell} \in L^{\infty}(\mu)$, and nonzero $k \in \mathbb{Z}$, we have

$$\lim_{j \rightarrow \infty} \sup_{N \geq 2^{2j-1}} \left\| \frac{1}{N} \sum_{n=1}^N e(k(n^{\ell b_j} + n^{b_j})\alpha) \cdot T^{n^g} f_1 \cdot T^{2n^g} f_2 \cdot \dots \cdot T^{\ell n^g} f_{\ell} \right\|_{L^2(\mu)} = 0.$$

In the next subsection we construct such a sequence (b_j) .

3.3. Construction of the sequence (b_j) . We need two preliminary results. The first was proved in [FrLW2] using deep machinery from ergodic theory:

Lemma 3.3. *Let (X, \mathcal{B}, μ, T) be a system, $f_1, \dots, f_{\ell} \in L^{\infty}(\mu)$ and $\alpha \in \mathbb{R} \setminus \mathbb{Q}$. If b, g are distinct positive integers then*

$$(3.2) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N e((n^{\ell b} + n^b)\alpha) \cdot T^{n^g} f_1 \cdot T^{2n^g} f_2 \cdot \dots \cdot T^{\ell n^g} f_{\ell} = 0$$

where the convergence takes place in $L^2(\mu)$.

The second is the following result:

Lemma 3.4. *Let $\ell, g \in \mathbb{N}$. Then there exists $d(\ell, g) \in \mathbb{N}$ such that for every $d \geq d(\ell, g)$ and $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, we have*

$$\lim_{N \rightarrow \infty} \sup_{S_1} \left\| \frac{1}{N} \sum_{n=1}^N e((n^{\ell d} + n^d)\alpha) \cdot T^{n^g} f_1 \cdot T^{2n^g} f_2 \cdot \dots \cdot T^{\ell n^g} f_{\ell} \right\|_{L^2(\mu)} = 0,$$

where S_1 is the collection of all systems (X, \mathcal{B}, μ, T) and functions f_1, \dots, f_{ℓ} with $\|f_i\|_{L^{\infty}(\mu)} \leq 1$ for $i = 1, \dots, \ell$.

Proof. The main idea is to apply van der Corput's lemma several times in order to get an upper bound for the expression

$$(3.3) \quad \left\| \frac{1}{N} \sum_{n=1}^N e((n^{\ell d} + n^d)\alpha) \cdot T^{n^g} f_1 \cdot T^{2n^g} f_2 \cdot \dots \cdot T^{\ell n^g} f_{\ell} \right\|_{L^2(\mu)}^{2^{\ell d-1}}$$

that does not depend on the transformation T or the functions f_1, \dots, f_{ℓ} . Such an estimate can be obtained using a rather standard argument, very much along the

lines of the polynomial exhaustion technique introduced by Bergelson in [Bel]. To do this we will use Lemma 5.6 and Proposition 5.7 in the Appendix. We get that for d large enough (depending on ℓ and g only), the quantity in (3.3) is bounded by a constant multiple of

$$(3.4) \quad \frac{1}{H_1 \cdots H_{\ell d-1}} \sum_{1 \leq h_i \leq H_i} \left| \frac{1}{N} \sum_{n=1}^N e(\Delta_{h_1, \dots, h_{\ell d-1}}(n^{\ell d} + n^d)\alpha) \right| + o_{N, H_i, H_i < N}(1),$$

where $\Delta_h(a_n) = a_{n+h} - a_n$, $\Delta_{h_1, \dots, h_r}(a_n) = \Delta_{h_1} \Delta_{h_2} \cdots \Delta_{h_r}(a_n)$, and $o_{N, H_i, H_i < N}(1)$ denotes a quantity that goes to zero as $N, H_i \rightarrow \infty$ in a way that $H_i/N \rightarrow 0$. Notice that the sequence $\Delta_{h_1, \dots, h_{\ell d-1}}(n^{\ell d} + n^d)$ is linear in n . Since α is irrational, letting $N \rightarrow +\infty$ and then $H_i \rightarrow +\infty$, we get that the quantity (3.4) converges to 0. This completes the proof. \square

Using that $\frac{1}{N} \sum_{n=1}^N \mathbb{1}_{I_{\pm}}((n^{\ell b} + n^b)\alpha) \rightarrow \frac{1}{4}$ for $b \in \mathbb{N}$, and Lemmas 3.3 and 3.4, the next result is proved in a similar fashion as Proposition 2.2.

Proposition 3.5. *There exists a sequence (b_j) that satisfies conditions (b1), (b2), and (b3) of Section 3.2.*

3.4. The sequence (s_n) satisfies (s1) and (s2). The next result is proved in essentially the same way as Proposition 2.3 and allows us to immediately deduce Theorem 1.2.

Proposition 3.6. *Suppose that the sequence (b_j) satisfies conditions (b1), (b2), and (b3) of Section 3.2. Then the sequence (s_n) defined by (3.1) satisfies conditions (s1) and (s2) of Proposition 3.2.*

4. POINTWISE CONVERGENCE

We shall prove Theorem 1.4. The argument is similar to the one used to prove Theorem A. However, extra complications arise since, as is typical for pointwise results, we need to establish quantitative estimates for some trigonometric sums.

Throughout this section we will assume that the irrational α is badly approximable, that means, there exists a $c > 0$ such that for every $p \in \mathbb{Z}$, $q \in \mathbb{N}$ we have $|\alpha - p/q| \geq c/q^2$. In fact, for convenience, we will fix α to be the *golden mean* $(\sqrt{5} + 1)/2$, in which case the previous estimate holds with $c = 1/3$.

4.1. A reduction. Theorem 1.4 is a direct consequence of the following result:

Theorem 4.1. *Let B be an arbitrary set of positive integers and α be the golden mean. Then, there exists an increasing sequence (s_n) of integers such that*

- (s1) *For every $b \in B$, the sequence $(s_n^b \alpha)$ has no asymptotic distribution $\pmod{1}$.*
- (s2) *For every $g \in \mathbb{N} \setminus B$ the sequence (s_n^g) is good for the pointwise ergodic theorem.*

Our first goal is to find a more convenient condition to replace (s2). To do this we are going to use the following lemma:

Lemma 4.2. *Let $(w_n)_{n \in \mathbb{N}}$ be a bounded sequence of complex numbers, and $(a_n)_{n \in \mathbb{N}}$ be a sequence of positive integers such that, for all $\gamma > 1$ we have*

$$\sum_{k \geq 0} \sup_{\beta \in \mathbb{R}} \left| \frac{1}{[\gamma^k]} \sum_{n=1}^{[\gamma^k]} w_n e(a_n \beta) \right|^2 < +\infty.$$

Then for every system (X, \mathcal{B}, μ, T) and $f \in L^2(\mu)$, we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N w_n \cdot T^{a_n} f = 0 \quad \mu\text{-almost everywhere.}$$

Sketch of the proof. From the spectral theorem we deduce that

$$\left\| \frac{1}{N} \sum_{n=1}^N w_n \cdot T^{a_n} f \right\|_2^2 \leq \|f\|_2^2 \cdot \sup_{\beta \in \mathbb{R}} \left| \frac{1}{N} \sum_{n=1}^N w_n e(a_n \beta) \right|^2.$$

Combining this with our hypothesis, we obtain that if $\gamma > 1$ then

$$\lim_{k \rightarrow +\infty} \frac{1}{[\gamma^k]} \sum_{n=1}^{[\gamma^k]} w_n \cdot T^{a_n} f = 0 \quad \mu\text{-almost everywhere for } f \in L^2(\mu).$$

Using a classical argument (see for example Lemma 1.5 in [RW]) we get the announced result. \square

Let I_+, I_- be the intervals defined by (2.1). Given a sequence of positive integers (b_j) and functions $\phi, \psi: \mathbb{T} \rightarrow \mathbb{T}$ let $(f_n(\phi, \psi))$ be the sequence defined by (2.8). As in Section 2, we define a sequence (s_n) by taking the elements of the set S given by

$$\mathbb{1}_S(n) = f_n(\mathbb{1}_{I_+}, \mathbb{1}_{I_-})$$

in increasing order.

Proposition 4.3. *Let B be an arbitrary set of positive integers, $\alpha \in \mathbb{R}$, and the sequences $(f_n(\phi, \psi))$ and (s_n) be as above. Suppose that*

- (s1') *For every $b \in B$, the sequence $\left(\frac{1}{N} \sum_{n=1}^N e(s_n^b \alpha)\right)$ diverges.*
- (s2') *For all trigonometric polynomials ϕ and ψ with zero integral, $\gamma > 1$, and $g \in \mathbb{N} \setminus B$, we have*

$$\sum_{k \geq 0} \sup_{\beta \in \mathbb{R}} \left| \frac{1}{[\gamma^k]} \sum_{n=1}^{[\gamma^k]} f_n(\phi, \psi) \cdot e(n^g \beta) \right| < +\infty.$$

Then the sequence (s_n) satisfies conditions (s1) and (s2) of Theorem 4.1.

Proof. It is clear that if (s1') holds then also (s1) holds.

It remains to show that condition (s2') implies that for $g \in \mathbb{N} \setminus B$ the sequence (s_n^g) is good for the pointwise ergodic theorem. We start by observing that condition (s2') combined with Lemma 4.2, guarantees that for every system (X, \mathcal{B}, μ, T) , we have

$$(4.1) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N f_n(\phi, \psi) \cdot T^{n^g} f = 0 \quad \mu\text{-almost everywhere for } f \in L^\infty(\mu).$$

Using the approximation argument of Lemma 2.7 and the fact $\frac{1}{N} \sum_{n=1}^N f_n(\phi, \psi) \rightarrow 0$, (this follows from (4.1)) we conclude that (4.1) remains true if we replace $f_n(\phi, \psi)$ by $f_n(\mathbb{1}_{I_+}, \mathbb{1}_{I_-}) - 1/4 = \mathbb{1}_S(n) - 1/4$. As a consequence, we have

$$(4.2) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N (\mathbb{1}_S(n) - 1/4) \cdot T^{n^g} f = 0 \quad \mu\text{-almost everywhere for } f \in L^\infty(\mu).$$

Using Bourgain's maximal inequality ([Bou]) for the ergodic averages $\frac{1}{N} \sum_{n=1}^N T^{n^g} f$, allows us to replace $L^\infty(\mu)$ by $L^2(\mu)$ in the preceding statement. Finally, using Bourgain's pointwise ergodic theorem ([Bou] or see the Appendix B of [Be2] for a simpler proof) for the ergodic averages along g -th powers, we get that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \mathbb{1}_S(n) \cdot T^{n^g} f \quad \text{exists } \mu\text{-almost everywhere for } f \in L^\infty(\mu).$$

Since the set S has positive density (this follows by setting $f = 1$ in (4.2)), we conclude that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N T^{s_n^g} f \quad \text{exists } \mu\text{-almost everywhere for } f \in L^2(\mu).$$

Therefore, the sequence (s_n^g) is good for the pointwise ergodic theorem. \square

The rest of this section will be devoted to the construction of a sequence (s_n) that satisfies conditions (s1') and (s2').

4.2. Definition of the sequence (s_n) . We remind the reader the set of integers B is given and the irrational number α is the golden mean. Let I_+ and I_- be the intervals defined in (2.1). The sequence (s_n) consists of the elements of a set S , taken in increasing order, that is defined as follows:

$$(4.3) \quad S = \bigcup_{j \geq 1} \{n \in \mathbb{N} \mid [2^{2j-1} \leq n < 2^{2j}, n^{b_j} \alpha \in I_+] \text{ or } [2^{2j} \leq n < 2^{2j+1}, n^{b_j} \alpha \in I_-]\}$$

where the sequence of integers (b_j) satisfies the following conditions

- (b1) We have $b_j \in B$, and all elements of B appear infinitely often in the sequence (b_j) .
- (b2) We have

$$\lim_{j \rightarrow \infty} \sup_{N > 2^{2j-1}} \left| \frac{1}{N} \sum_{n=1}^N \mathbb{1}_{I_\pm}(n^{b_j} \alpha) - \frac{1}{4} \right| = 0.$$

- (b3) For every nonzero $m \in \mathbb{Z}$, and $g \in \mathbb{N} \setminus B$, we have

$$\lim_{j \rightarrow \infty} \sup_{N \geq 2^{2j-1}} \sup_{\beta \in \mathbb{R}} \left| \frac{1}{N^{1-\eta(N)}} \sum_{n=1}^N e(mn^{b_j} \alpha + n^g \beta) \right| < +\infty$$

where $\eta(N) = (\log_2(N))^{-1/2}$.

In the next subsection we construct such a sequence (b_j) .

4.3. Construction of the sequence (b_j) . The key ingredient in the construction is the following exponential sum estimate:

Proposition 4.4. *Let α be the golden mean, and $(\eta(N))$ be a sequence of real numbers which tends to zero at infinity. For every $b, g \in \mathbb{N}$ with $b \neq g$, and nonzero $m \in \mathbb{Z}$, there exists $N_0 = N_0(b, g, m)$, such that if $N > N_0$ then*

$$\sup_{\beta \in \mathbb{R}} \left| \sum_{n=1}^N e(mn^b \alpha + n^g \beta) \right| \leq N^{1-\eta(N)}.$$

Proof. This is an immediate consequence of Lemmas 5.4 and 5.5 in the Appendix. \square

Using that $\frac{1}{N} \sum_{n=1}^N \mathbb{1}_{I_{\pm}}(n^b \alpha) \rightarrow \frac{1}{4}$ for $b \in \mathbb{N}$, and Proposition 4.4, the next result is proved in essentially the same way as Proposition 2.2 (the argument is actually a little bit simpler in this case, since we have already combined the estimates dealing with $b < g$ and $b > g$ into a single estimate).

Proposition 4.5. *There exists a sequence (b_j) that satisfies conditions (b1), (b2), and (b3) of Section 4.2.*

4.4. The sequence (s_n) satisfies conditions (s1) and (s2). The goal of this section is to prove the following proposition, that allows us to immediately deduce Theorem 1.4.

Proposition 4.6. *Suppose that the sequence (b_j) satisfies conditions (b1), (b2), and (b3) of Section 4.2. Then the sequence (s_n) defined by (4.3) satisfies conditions (s1) and (s2) of Theorem 4.1.*

Before starting the proof of Proposition 4.6, let us gather some useful properties that the sequence $\eta(N) = (\log_2(N))^{-1/2}$ satisfies:

- (η 1) The sequence $(N^{-\eta(N)})$ is decreasing.
- (η 2) For every $\gamma > 1$, we have $\sum_{k \geq 0} [\gamma^k]^{-\eta([\gamma^k])} < +\infty$.
- (η 3) If we define

$$\rho(N) = \frac{1}{N} \sum_{i=1}^{\lfloor \log_2 N \rfloor} 2^{i(1-\eta(2^i))},$$

then for every $\gamma > 1$, we have $\sum_k \rho([\gamma^k]) < +\infty$.

The first property is obvious.

To check the second property notice that $\eta([\gamma^k]) \sim c/\sqrt{k}$, so $[\gamma^k]^{-\eta([\gamma^k])} = O(\gamma^{-c\sqrt{k}})$, for some constant $c > 0$.

To check the third property notice that

$$\sum_{i=1}^l 2^{i(1-\eta(2^i))} = \sum_{i=1}^l 2^{i-\sqrt{i}} = \sum_{i=1}^{\lfloor l/2 \rfloor - 1} 2^{i-\sqrt{i}} + \sum_{i=\lfloor l/2 \rfloor}^l 2^{i-\sqrt{i}} \leq 2^{l/2} + 2^{-\sqrt{l/2}} 2^{l+1},$$

hence

$$\sum_{i=1}^l 2^{i(1-\eta(2^i))} \leq 2^{l-\sqrt{l/2}+2}.$$

It follows that

$$\rho([\gamma^k]) \leq \frac{1}{[\gamma^k]} 2^{\log_2([\gamma^k]) - \sqrt{\log_2([\gamma^k])/2} + 2} = O\left(2^{-\sqrt{k(\log_2 \gamma)/2}}\right),$$

which implies $(\eta 3)$.

Lemma 4.7. *Suppose that the sequence (b_j) satisfies condition (b3) of Section 4.2. If ϕ and ψ are two trigonometric polynomials with zero integral, let $(f_n(\phi, \psi))$ be the sequence defined by (2.8). Then for every $\gamma > 1$ and for every $g \in \mathbb{N} \setminus B$, we have*

$$\sum_{k \geq 0} \sup_{\beta \in \mathbb{R}} \left| \frac{1}{[\gamma^k]} \sum_{n=1}^{[\gamma^k]} f_n(\phi, \psi) e(n^g \beta) \right| < +\infty.$$

Proof. For brevity we will write f_n instead of $f_n(\phi, \psi)$. Since (b_j) satisfies condition (b3), there exists a positive constant $C = C(\phi, \psi, g)$ such that for every large enough j and $N \geq 2^{2j-1}$, we have

$$(4.4) \quad \sup_{\beta \in \mathbb{R}} \left| \sum_{n=1}^N \xi(n^{b_j} \alpha) e(n^g \beta) \right| \leq CN^{1-\eta(N)},$$

where ξ is either ϕ or ψ . We will use this estimate to find an upper bound for the averages

$$\frac{1}{N} \sum_{n=1}^N f_n e(n^g \beta).$$

We start by noticing that

$$\frac{1}{2^{2j}} \sum_{n=2^{2j+1}}^{2^{2j+1}} f_n e(n^g \beta) = 2 \frac{1}{2^{2j+1}} \sum_{n=1}^{2^{2j+1}} \psi(n^{b_j} \alpha) e(n^g \beta) - \frac{1}{2^{2j}} \sum_{n=1}^{2^{2j}} \psi(n^{b_j} \alpha) e(n^g \beta).$$

We also get a similar estimate with $2j+1$ in place of $2j$ and ϕ in place of ψ . It follows from (4.4) that there exists $j_0 \geq 0$ (depending on ϕ, ψ and g), such that for every $j \geq j_0$, we have

$$(4.5) \quad \sup_{\beta \in \mathbb{R}} \left| \frac{1}{2^j} \sum_{n=2^{2j+1}}^{2^{2j+1}} f_n e(n^g \beta) \right| \leq C(2^j)^{-\eta(2^j)}.$$

Now consider a large enough integer N , then $N \in (2^j, 2^{j+1}]$ for some $j \geq j_0$. We split the sum between 1 and N into several pieces

$$\sum_{n=1}^N \cdot = \sum_{n=1}^{2^{j_0}} \cdot + \sum_{i=j_0}^{j-1} \sum_{n=2^{i+1}}^{2^{i+1}} \cdot + \sum_{n=2^{2j+1}}^N \cdot,$$

and we obtain the following estimate

$$(4.6) \quad \left| \frac{1}{N} \sum_{n=1}^N f_n e(n^g \beta) \right| \leq C \frac{2^{2j_0}}{N} + \frac{1}{N} \sum_{i=j_0}^{j-1} \left| \sum_{n=2^{i+1}}^{2^{i+1}} f_n e(n^g \beta) \right| + \left| \frac{1}{N} \sum_{n=2^{2j+1}}^N f_n e(n^g \beta) \right|.$$

By (4.5) we have

$$(4.7) \quad \frac{1}{N} \sum_{i=j_0}^{j-1} \left| \sum_{n=2^i+1}^{2^{i+1}} f_n e(n^g \beta) \right| \leq C \frac{1}{N} \sum_{i=1}^{\lfloor \log_2 N \rfloor} 2^{i(1-\eta(2^i))}.$$

Moreover, since

$$\left| \frac{1}{N} \sum_{n=2^j+1}^N f_n e(n^g \beta) \right| \leq \left| \frac{1}{N} \sum_{n=1}^N \xi(n^{b_j} \alpha) e(n^g \beta) \right| + \left| \frac{1}{2^j} \sum_{n=1}^{2^j} \xi(n^{b_j} \alpha) e(n^g \beta) \right|,$$

where ξ is either ϕ or ψ , we get using (4.4) and property ($\eta 1$) that

$$(4.8) \quad \left| \frac{1}{N} \sum_{n=2^j+1}^N f_n e(n^g \beta) \right| \leq C \left(N^{-\eta(N)} + (2^j)^{-\eta(2^j)} \right) \leq 2C (N/2)^{-\eta(N/2)}.$$

Combining equations (4.6), (4.7), and (4.8), and using properties ($\eta 1$), ($\eta 2$), and ($\eta 3$), we get the advertised result. \square

Proof of Proposition 4.6. Let (s_n) be the sequence defined by (4.3). By Proposition 4.3, it suffices to verify properties (s1') and (s2') mentioned there. Using Proposition 2.5, we see that properties (b1) and (b2) give property (s1'). Also, using Lemma 4.7, we see that property (b3) give property (s2'). This completes the proof. \square

5. APPENDIX

We prove some results that were used in the main part of the article.

5.1. Van der Corput's lemma. The following is a Hilbert space version of a classical elementary estimate of van der Corput. It appears in the form stated below in [Be1].

Lemma 5.1. *Let v_1, \dots, v_N be vectors of a Hilbert space with $\|v_i\| \leq 1$ for $i = 1, \dots, N$. Then for every integer H between 1 and N we have*

$$\left\| \frac{1}{N} \sum_{n=1}^N v_n \right\|^2 \leq \frac{2}{H} + \frac{4}{H} \sum_{h=1}^{H-1} \left| \frac{1}{N} \sum_{n=1}^{N-h} \langle v_{n+h}, v_n \rangle \right|.$$

We will also use the following estimate which is an immediate corollary of the previous lemma:

Corollary 5.2. *Let v_1, \dots, v_N be vectors of a Hilbert space with $\|v_i\| \leq 1$ for $i = 1, \dots, N$. Then for every integer H between 1 and N we have*

$$\left\| \frac{1}{N} \sum_{n=1}^N v_n \right\|^2 \leq \frac{4}{H} \sum_{h=1}^H \left| \frac{1}{N} \sum_{n=1}^N \langle v_{n+h}, v_n \rangle \right| + o_{N,H,H \prec N}(1),$$

where $o_{N,H,H \prec N}(1)$ denotes a quantity that goes to zero as $N, H \rightarrow \infty$ in a way that $H/N \rightarrow 0$.

5.2. Dyadic intervals. The next lemma allows us, under suitable assumptions, to concatenate dyadic pieces of sequences and create a new sequence with average zero.

Lemma 5.3. *Let $(u_{n,j})_{n,j \in \mathbb{N}}$ be a family of complex numbers that satisfy*

$$\lim_{j \rightarrow \infty} \sup_{N > 2^j} \left| \frac{1}{N} \sum_{n=1}^N u_{n,j} \right| = 0.$$

Define the sequence (u_n) by

$$u_n = u_{n,j} \text{ if } 2^j \leq n < 2^{j+1}.$$

Then

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N u_n = 0.$$

Proof. Let $\varepsilon > 0$. We define

$$\varepsilon(j) = \sup_{N > 2^j} \left| \frac{1}{N} \sum_{n=1}^N u_{n,j} \right|$$

and

$$\varepsilon'(j) = \sup_{k \geq j} \varepsilon(k).$$

By our assumption, there exists j_0 such that $\varepsilon'(j) < \varepsilon$ for every $j \geq j_0$.

We start by noticing that

$$\frac{1}{2^j} \sum_{n=2^{j+1}}^{2^{j+1}} u_n = 2 \frac{1}{2^{j+1}} \sum_{n=1}^{2^{j+1}} u_{n,j} - \frac{1}{2^j} \sum_{n=1}^{2^j} u_{n,j}.$$

Therefore, for every $j \in \mathbb{N}$ we have

$$(5.1) \quad \left| \frac{1}{2^j} \sum_{n=2^{j+1}}^{2^{j+1}} u_n \right| \leq 3\varepsilon(j).$$

Now suppose that $N > 2^{j_0}$, then $N \in (2^j, 2^{j+1}]$ for some $j \geq j_0$. We split the sum between 1 and N into several pieces

$$\sum_{n=1}^N \cdot = \sum_{n=1}^{2^{j_0}} \cdot + \sum_{i=j_0}^{j-1} \sum_{n=2^{i+1}}^{2^{i+1}} \cdot + \sum_{n=2^{j+1}}^N \cdot,$$

in order to obtain the following upper bound

$$\left| \frac{1}{N} \sum_{n=1}^N u_n \right| \leq \left| \frac{1}{N} \sum_{n=1}^{2^{j_0}} u_n \right| + \frac{1}{N} \sum_{i=j_0}^{j-1} \left| \sum_{n=2^{i+1}}^{2^{i+1}} u_n \right| + \left| \frac{1}{N} \sum_{n=2^{j+1}}^N u_n \right|.$$

Using (5.1), we get

$$\frac{1}{N} \sum_{i=j_0}^{j-1} \left| \sum_{n=2^{i+1}}^{2^{i+1}} u_n \right| \leq \frac{1}{N} \sum_{i=j_0}^{j-1} 3\varepsilon(i) 2^i \leq 3\varepsilon'(j_0) \frac{1}{N} \sum_{i=j_0}^{j-1} 2^i \leq 3\varepsilon'(j_0).$$

We also have

$$\frac{1}{N} \sum_{n=2^j+1}^N u_n = \frac{1}{N} \sum_{n=1}^N u_{n,j} - \frac{1}{N} \sum_{n=1}^{2^j} u_{n,j},$$

which gives

$$\left| \frac{1}{N} \sum_{n=2^j+1}^N u_n \right| \leq \left| \frac{1}{N} \sum_{n=1}^N u_{n,j} \right| + \left| \frac{1}{2^j} \sum_{n=1}^{2^j} u_{n,j} \right| \leq 2\varepsilon'(j).$$

Putting these estimates together, we obtain

$$\left| \frac{1}{N} \sum_{n=1}^N u_n \right| \leq \left| \frac{1}{N} \sum_{n=1}^{2^{j_0}} u_n \right| + 5\varepsilon'(j_0).$$

By taking N large enough we can make sure that the right hand side becomes less than 6ε . This completes the proof. \square

5.3. Exponential sum estimates. We derive two exponential sum estimates. The first gives non-trivial power type savings when one deals with polynomials with leading coefficient an integer multiple of the golden mean. Let us recall the “bad approximation property” of the golden mean α

$$(5.2) \quad \text{For all non-zero integers } q \text{ we have } d(q\alpha, \mathbb{Z}) \geq 1/(3q).$$

Lemma 5.4. *Let α be the golden mean and $b \in \mathbb{N}$. There exists $C = C(b) > 0$ such that, for every $m, N \in \mathbb{N}$, and real polynomial P with degree $< b$, we have*

$$(5.3) \quad \left| \sum_{n=1}^N e(mn^b\alpha + P(n)) \right| \leq C m^{2^{1-b}} N^{1-4^{1-b}}.$$

Proof. We use an induction on b . For $b = 1$, we have

$$\left| \sum_{n=1}^N e(mn\alpha) \right| \leq \frac{2}{|e(m\alpha) - 1|} = \frac{1}{|\sin \pi m\alpha|} \leq \frac{1}{2d(m\alpha, \mathbb{Z})} \leq \frac{3m}{2},$$

by the bad approximation property (5.2).

Suppose that the estimate (5.3) holds for the integer b . We are going to show that it also holds for the integer $b + 1$. Let us define

$$S(m, N, b) = \sup_{P \in \mathbb{R}[X], \deg(P) < b} \left| \sum_{n=1}^N e(mn^b\alpha + P(n)) \right|.$$

From van der Corput’s inequality (Lemma 5.1), we deduce that for every integer H between 1 and N , we have

$$(5.4) \quad S(m, N, b+1)^2 \leq \frac{2N^2}{H} + \frac{4N}{H} \sum_{h=1}^{H-1} S(mh(b+1), N-h, b).$$

The induction hypothesis gives that for some constant $C = C_b$ we have

$$S(mh(b+1), N-h, b) \leq C (mh)^{2^{1-b}} (N-h)^{1-4^{1-b}},$$

and so for h between 1 and $H - 1$ we have

$$(5.5) \quad S(mh(b+1), N-h, b) \leq C m^{2^{1-b}} (H-1)^{2^{1-b}} N^{1-4^{1-b}}.$$

Combining (5.4) and (5.5) and we obtain

$$(5.6) \quad S(m, N, b+1)^2 \leq 4Cm^{2^{1-b}} \left(N^2H^{-1} + N^{2-4^{1-b}}(H-1)^{2^{1-b}} \right).$$

Now choosing $H = \lceil N^{2^{1-2b}} \rceil + 1$ gives

$$N^2H^{-1} \leq N^{2(1-4^{-b})}, \quad N^{2-4^{1-b}}(H-1)^{2^{1-b}} \leq N^{2(1-2\cdot 4^{-b}+2\cdot 8^{-b})} \leq N^{2(1-4^{-b})}.$$

Using this together with (5.6) we find that

$$S(m, N, b+1)^2 \leq 4Cm^{2^{1-b}}N^{2(1-4^{-b})}.$$

Taking square roots establishes (5.3) for $b+1$. This completes the induction and the proof. \square

The second lemma gives non-trivial power type savings for polynomials that have an integer multiple of the golden mean as a non-leading (non-constant) coefficient. Its proof is a simplification of an argument that appears in [BosKoQW].

Lemma 5.5. *Let α be the golden mean and $g \in \mathbb{N}$. For every $\varepsilon > 0$, there exists $C = C(\varepsilon, g) < +\infty$ such that, for every $\beta \in \mathbb{R}$, $b \in \mathbb{N}$ with $b < g$, $N \in \mathbb{N}$, and nonzero $m \in \mathbb{Z}$ with $|m| \leq N^{2^{-g-2}}$, we have*

$$(5.7) \quad \left| \sum_{n=1}^N e(mn^b\alpha + n^g\beta) \right| \leq CN^{1+\varepsilon-2^{-2g-1}}.$$

Proof. The proof proceeds as follows: If β is not well approximated by rationals, then classical estimates of Weyl immediately give the advertised estimate. If β is well approximated by rationals, using partial summation we can replace β by a rational (up to a small error), and reduce the problem to studying an exponential sum involving a polynomial that has an integer multiple of α as leading coefficient. In this case, again the classical estimates of Weyl give the advertised result.

So let us first recall Weyl's classical estimate ([V]). For every $k \in \mathbb{N}$ and $\varepsilon > 0$, there exists a constant C satisfying the following property: for every $\beta \in \mathbb{R}$ and relatively prime $r, s \in \mathbb{N}$ with $|\beta - r/s| < 1/s^2$, and for every real polynomial $P(x)$ with leading coefficient βx^k , we have

$$\left| \sum_{n=1}^N e(P(n)) \right| \leq CN^{1+\varepsilon} \left(\frac{1}{s} + \frac{1}{N} + \frac{s}{N^k} \right)^{1/2^{k-1}}.$$

We fix $\beta \in \mathbb{R}$, $b \in \mathbb{N}$ with $b < g$, $N \in \mathbb{N}$, nonzero $m \in \mathbb{Z}$ with $|m| \leq N^{2^{-g-2}}$, and we define $\gamma = 2^{-g-2}$.

By Dirichlet's principle, there exist $r, s \in \mathbb{N}$, relatively prime, such that $s \leq N^{g-\gamma}$ and

$$\left| \beta - \frac{r}{s} \right| \leq \frac{1}{N^{g-\gamma}s}.$$

We distinguish two cases: either $s > N^\gamma$ (bad approximation) or $s \leq N^\gamma$ (good approximation).

Case 1. Suppose that $s > N^\gamma$. By Weyl's estimate, we have

$$\left| \sum_{n=1}^N e(mn^b\alpha + n^g\beta) \right| \leq C(\varepsilon, g)N^{1+\varepsilon}(N^{-\gamma} + N^{-1} + N^{-\gamma})^{2^{1-g}},$$

which implies the estimate (5.7), since $\gamma = 2^{-g-2}$.

Case 2. Suppose now that $s \leq N^\gamma$. By Dirichlet's principle, there exist $t, u \in \mathbb{N}$, relatively prime, such that $u \leq N^{b-1/2}$ and

$$(5.8) \quad \left| ms^b \alpha - \frac{t}{u} \right| \leq \frac{1}{N^{b-1/2} u}.$$

The bad approximation property of α mentioned in (5.2) gives that $mus^b \geq \frac{1}{3}N^{b-1/2}$. Since $s \leq N^\gamma$ and $|m| \leq N^\gamma$ we have $u \geq \frac{1}{3}N^{b-1/2-\gamma b-\gamma}$.

Consider now an integer M between $N^{1-\gamma}$ and N . We are going to compare the sums $\sum_{n \leq M} e(mn^b \alpha + n^g \beta)$ with the sums $\sum_{n \leq M} e(mn^b \alpha + \frac{r}{s}n^g)$ that are easier to estimate. Let us first estimate the second sum. We have

$$\sum_{n \leq M} e\left(mn^b \alpha + \frac{r}{s}n^g\right) = \sum_{j=1}^s \sum_{i \geq 0, si+j \leq M} e\left(m(si+j)^b \alpha + \frac{r}{s}j^g\right),$$

hence

$$(5.9) \quad \left| \sum_{n \leq M} e\left(mn^b \alpha + \frac{r}{s}n^g\right) \right| \leq \sum_{j=1}^s \left| \sum_{i \geq 0, si+j \leq M} e\left(m(si+j)^b \alpha\right) \right|.$$

By Weyl's estimate and (5.8), we have

$$\left| \sum_{i \geq 0, si+j \leq M} e\left(m(si+j)^b \alpha\right) \right| \leq C(\varepsilon, b) \left(\frac{M}{s}\right)^{1+\varepsilon} \left(\frac{1}{u} + \frac{s}{M} + u \left(\frac{s}{M}\right)^b\right)^{2^{1-b}}.$$

Using that

$$N^{1-\gamma} \leq M \leq N, \quad 1 \leq s \leq N^\gamma, \quad \text{and} \quad \frac{1}{3}N^{b-1/2-\gamma b-\gamma} \leq u \leq N^{b-1/2},$$

we obtain

$$\left| \sum_{i \geq 0, si+j \leq M} e\left(m(si+j)^b \alpha\right) \right| \leq C(\varepsilon, b) \frac{N^{1+\varepsilon}}{s} \left(N^{-b+1/2+\gamma b+\gamma} + N^{-1+2\gamma} + N^{-1/2+2\gamma b}\right)^{2^{1-b}}.$$

The term $N^{-1/2+2\gamma b}$ is dominant, and is of the order less than $N^{-1/4}$. It follows that

$$\left| \sum_{i \geq 0, si+j \leq M} e\left(m(si+j)^b \alpha\right) \right| \leq C(\varepsilon, b) \frac{N^{1+\varepsilon-2^{b-1}}}{s}.$$

Since the integer g is fixed and $b < g$ we have

$$\left| \sum_{i \geq 0, si+j \leq M} e\left(m(si+j)^b \alpha\right) \right| \leq C(\varepsilon, g) \frac{N^{1+\varepsilon-2^{g-1}}}{s}.$$

In conjunction with (5.9) this gives

$$(5.10) \quad \left| \sum_{n \leq M} e\left(mn^b \alpha + \frac{r}{s}n^g\right) \right| \leq C(\varepsilon, g) N^{1+\varepsilon-2^{g-1}}.$$

We come back to our main goal of estimating the sums $\sum_{n \leq M} e(mn^b \alpha + n^g \beta)$. We are going to use summation by parts. We set $S(M) = \sum_{n \leq M} e(mn^b \alpha + \frac{r}{s} n^g)$ and notice that

$$\begin{aligned} \left| \sum_{n \leq N} e(mn^b \alpha + n^g \beta) \right| &\leq N^{1-\gamma} + \left| \sum_{N^{1-\gamma} < n \leq N} e(mn^b \alpha + n^g \beta) \right| \\ &\leq N^{1-\gamma} + \left| \sum_{N^{1-\gamma} < n \leq N} (S(n) - S(n-1)) e\left(n^g \left(\beta - \frac{r}{s}\right)\right) \right|. \end{aligned}$$

We have

$$\left| e\left(n^g \left(\beta - \frac{r}{s}\right)\right) - e\left((n+1)^g \left(\beta - \frac{r}{s}\right)\right) \right| \leq C n^{g-1} \left| \beta - \frac{r}{s} \right|,$$

where the constant C does not depend on β because $\beta - r/s$ is uniformly bounded. We know that for $n \leq N$, we have $n^{g-1} |\beta - r/s| \leq N^{-1+\gamma}$. So using partial summation, we obtain

$$\begin{aligned} \left| \sum_{n \leq N} e(mn^b \alpha + n^g \beta) \right| &\leq \\ &N^{1-\gamma} + |S([N^{1-\gamma}] + 1)| + |S(N)| + C \sum_{N^{1-\gamma} < n \leq N} |S(n)| N^{-1+\gamma}. \end{aligned}$$

Using (5.10) we conclude that

$$\left| \sum_{n \leq N} e(mn^b \alpha + n^g \beta) \right| \leq C(\varepsilon, g) N^{1+\varepsilon-2^{-g-1}+\gamma}.$$

Recalling that $\gamma = 2^{-g-2}$, we derive an estimate stronger than (5.7). This completes the proof. \square

5.4. The PET induction argument. We give the details needed to complete the proof of Lemma 3.4. The next result follows immediately from Corollary 5.2:

Lemma 5.6. *Consider a family of integer polynomials $\{p_1, \dots, p_k\}$ and an integer polynomial p , all of them having zero constant term. Let $\{q_1, \dots, q_{k'}\}$ be the family of distinct integer polynomials that is defined using the following operation: we start with the family of polynomials*

$p_1(n+h) - p_1(h) - p(n), \dots, p_k(n+h) - p_k(h) - p(n), p_1(n) - p(n), \dots, p_k(n) - p(n)$,
and we remove polynomials that are identically zero and repetitions of polynomials. Then for every system (X, \mathcal{B}, μ, T) , and sequence of complex numbers (u_n) with $\|u_n\|_\infty \leq 1$, we have

$$\begin{aligned} \sup_{f_1, \dots, f_k} \left\| \frac{1}{N} \sum_{n=1}^N u_n T^{p_1(n)} f_1 \cdots T^{p_k(n)} f_k \right\|_2^2 &\leq \\ \frac{4}{H} \sum_{h=1}^H \sup_{f_1, \dots, f_{k'}} \left\| \frac{1}{N} \sum_{n=1}^N u_{n+h} \bar{u}_n T^{q_1(n)} f_1 \cdots T^{q_{k'}(n)} f_{k'} \right\|_2 &+ o_{N, H, H \ll N}(1), \end{aligned}$$

where the supremums are taken over families of $L^\infty(\mu)$ functions bounded by 1.

Let \mathcal{P} be a family of non-constant integer polynomials with zero constant term. The maximum degree of the polynomials is called the *degree* of the polynomial family and we denote it by d . Let \mathcal{P}_i be the subfamily of polynomials of degree i in \mathcal{P} . We let w_i denote the number of distinct leading coefficients that appear in the family \mathcal{P}_i . The vector (d, w_d, \dots, w_1) is called the *type* of the polynomial family \mathcal{P} . We use an induction scheme, often called PET induction (Polynomial Exhaustion Technique), on types of polynomial families that was introduced by Bergelson in [Be1]. We order the set of all possible types lexicographically, this means, $(d, w_d, \dots, w_1) > (d', w'_d, \dots, w'_1)$ if and only if in the first instance where the two vectors disagree the coordinate of the first vector is greater than the coordinate of the second vector.

Proposition 5.7 (Bergelson’s PET [Be1]). *Let $\{p_1, \dots, p_k\}$ be a family of non-constant integer polynomials with zero constant term. After applying finitely many times the operation defined in Lemma 5.6 (for good choices of the auxiliary polynomial p at each step) it is possible to obtain the empty family of polynomials.*

Proof. Let p_{min} be any member of the family $\{p_1, \dots, p_k\}$ that has minimal degree. Notice that after applying the operation defined in Lemma 5.6 for $p = p_{min}$, we obtain a new family of polynomials that has type strictly less than the type of the family $\{p_1, \dots, p_k\}$. The result now follows using induction on the type of the polynomial family. \square

The following two examples illustrate how a typical PET induction argument works:

Example 2. Suppose that we start with the family of polynomials $\mathcal{P}_0 = \{n, 2n\}$ that has type $(1, 2)$. Applying the operation defined in Lemma 5.6 with $p(n) = n$ we obtain the family $\mathcal{P}_1 = \{n\}$ that has type $(1, 1)$. After one more application of the operation we obtain an empty family of polynomials.

Example 3. Suppose that we start with the family of polynomials $\mathcal{P}_0 = \{n^2, 2n^2\}$ that has type $(2, 2, 0)$. Applying successively the operation defined in Lemma 5.6 we obtain the following families of polynomials: Using $p(n) = n^2$ we get the family

$$\mathcal{P}_1 = \{2nh_1, n^2 + 4nh_1, n^2\}$$

that has type $(2, 1, 1)$. Using $p(n) = 2nh_1$ we get the family

$$\mathcal{P}_2 = \{n^2 + 2n(h_1 + h_2), n^2 + 2n(h_2 - h_1), n^2 + 2nh_1, n^2 - 2nh_1\}$$

that has type $(2, 1, 0)$. Using $p(n) = n^2$ we get the family

$$\mathcal{P}_3 = \{2n(h_1 + h_2 + h_3), 2n(h_3 + h_2 - h_1), 2n(h_1 + h_3), \\ 2n(h_3 - h_1), 2n(h_1 + h_2), 2n(h_2 - h_1), 2nh_1, -2nh_1\}$$

that has type at most $(1, 8)$ (actually equal to $(1, 8)$ for most values of h_1, h_2, h_3). The last family consists of linear polynomials and can be dealt as in Example 2. After 8 more operations we arrive to an empty family of polynomials.

REFERENCES

[Be1] V. Bergelson. Weakly mixing PET. *Ergodic Theory Dynam. Systems*, **7** (1987), 337–349.
 [Be2] V. Bergelson. Combinatorial and Diophantine applications of ergodic theory. Appendix A by A. Leibman and Appendix B by A. Quas and M. Wierdl. *Handbook of dynamical systems*, Vol. 1B, 745–869, Elsevier B. V., Amsterdam, 2006.

- [BeHKr] V. Bergelson, B. Host and B. Kra. Multiple recurrence and nilsequences, *Inventiones Math.*, **160**, (2005), 261–303.
- [BosKoQW] M. Boshernitzan, G. Kolesnik, A. Quas, M. Wierdl. Ergodic averaging sequences. *J. Anal. Math.*, **95**, (2005), 63–103.
- [Bou] J. Bourgain. On the maximal ergodic theorem for certain subsets of the integers. *Israel J. Math.*, **61**, (1988), 39–72.
- [DES] J. Deshouillers, P. Erdős, A. Sárközy, A. On additive bases. *Acta Arith.*, **30**, (1976), no. 2, 121–132.
- [DFo] J. Deshouillers, E. Fouvry. On additive bases II. *J. London Math. Soc. (2)*, **14**, (1976), no. 3, 413–422.
- [FrLW1] N. Frantzikinakis, E. Lesigne, M. Wierdl. Sets of k -recurrence but not $(k+1)$ -recurrence. *Annales de l'Institut Fourier*, **56**, Issue 4, (2006), 839–849.
- [FrLW2] N. Frantzikinakis, E. Lesigne, M. Wierdl. Powers of sequences and recurrence. To appear in *Proc. Lond. Math. Soc.*
- [HKr1] B. Host, B. Kra. Nonconventional ergodic averages and nilmanifolds. *Annals of Math.*, **161**, (2005), 397–488.
- [HKr2] B. Host, B. Kra. Convergence of polynomial ergodic averages. *Isr. J. Math.*, **149**, (2005), 1–19.
- [HKr3] B. Host, B. Kra. Uniformity seminorms on l^∞ and applications. *Preprint available at <http://www.math.northwestern.edu/~kra/papers/uniformity.pdf>*.
- [RW] J. Rosenblatt, M. Wierdl. Pointwise theorems via harmonic analysis. Ergodic theory and its connections with harmonic analysis (Alexandria, 1993), *London Math. Soc. Lecture Note Ser.*, **205**, 3–151.
- [V] R. Vaughan. The Hardy-Littlewood method. Second edition, *Cambridge Tracts in Mathematics*, **125**, Cambridge University Press, Cambridge, 1997.
- [Z] Universal characteristic factors and Furstenberg averages. *J. Amer. Math. Soc.*, **20**, (2007), 53–97.

(Nikos Frantzikinakis) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MEMPHIS, MEMPHIS, TN, 38152, USA

E-mail address: `frantzikinakis@gmail.com`

(Michael Johnson) DEPARTMENT OF MATHEMATICS & STATISTICS, SWARTHMORE COLLEGE, SWARTHMORE, PA, 19081, USA

E-mail address: `mjohnso3@swarthmore.edu`

(Emmanuel Lesigne) LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE (UMR CNRS 6083), UNIVERSITÉ FRANÇOIS RABELAIS TOURS, FÉDÉRATION DE RECHERCHE DENIS POISSON, PARC DE GRANDMONT, 37200 TOURS, FRANCE

E-mail address: `emmanuel.lesigne@lmpt.univ-tours.fr`

(Máté Wierdl) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MEMPHIS, MEMPHIS, TN, 38152, USA

E-mail address: `mw@csi.hu`