



HAL
open science

Extremal Bases, Geometrically Separated Domains and Applications

Philippe Charpentier, Yves Dupain

► **To cite this version:**

Philippe Charpentier, Yves Dupain. Extremal Bases, Geometrically Separated Domains and Applications. *Algebra i Analiz*, 2014, 26 (1), pp.196-269. hal-00328871v2

HAL Id: hal-00328871

<https://hal.science/hal-00328871v2>

Submitted on 15 Oct 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXTREMAL BASES, GEOMETRICALLY SEPARATED DOMAINS AND APPLICATIONS

PHILIPPE CHARPENTIER & YVES DUPAIN

ABSTRACT. In this paper we introduce the notion of extremal basis of tangent vector fields at a boundary point of finite type of a pseudo-convex domain in \mathbb{C}^n , $n \geq 3$. Using this notion we define the class of geometrically separated domains at a boundary point and we give a description of their complex geometry. Examples of such domains are given, for instance, by locally lineally convex domains, domains with locally diagonalizable Levi form at a point or domains for which the Levi form have comparable eigenvalues near a point. Moreover we show that geometrically separated domains can be localized. We also give an example of a non geometrically separated domains. Next we define what we call “adapted pluri-subharmonic function” and give sufficient conditions, related to extremal bases, for their existence. Then, for these domains, when such functions exist, we prove global and local sharp estimate for the Bergman and Szegő projections. As an application, we strengthen a result by C. Fefferman, J. J. Kohn and M. Machedon ([FKM90]) for the local Hölder estimate of the Szegő projection removing the arbitrary small loss in the Hölder index and giving a stronger non-isotropic estimate.

1. INTRODUCTION

The study of the regularity with sharp estimates for the Bergman and Szegő projections for pseudo-convex domains in \mathbb{C}^n became very active for domains of finite type when D. Catlin proved his fundamental characterization of subelliptic estimates ([Cat87]).

Quite quickly, the case of domains in \mathbb{C}^2 was completely solved by D. Catlin in [Cat89], A. Nagel, J.-P. Rosay, E. M. Stein and S. Wainger in [NRSW89], M. Christ in [Chr88], C. Fefferman and J. J. Kohn in [FK88] and J. McNeal in [McN89].

In higher dimensions, the situation is more complicated and, until now, there are only partial results. One of the main difficulties is the description of the geometry of the domain: there are some special bases of the complex tangent space at the boundary playing an important role in this description and also in the Lipschitz estimates of the projectors. Thus the first results concern domains for which these bases are more or less evident. For example, the class of domains for which the Levi form have rank larger than $n - 2$ was studied by M. Machedon in [Mac88] (see also S. Cho [Cho94, Cho96], [AC99]) and, even in that case, the situation is not so simple. An other example is given by decoupled domains, treated by several authors (see for example [McN91], [CG94]).

A typical example where the choice of special bases is essential, and not evident, is the case of convex domains in \mathbb{C}^n . In [McN94, McN02] J. Mc Neal introduced some special bases (called ε -extremal in [BCD98]) and gave a description of the complex geometry with the construction of a pseudo-distance near the boundary related to these bases. With that geometry, and a construction of a “good” pluri-subharmonic function, he proved sharp point-wise estimates for the Bergman kernel and its derivatives. Using this geometry J. Mc Neal and E. M. Stein ([MS94] and [MS97]) proved sharp estimates for the Bergman and Szegő projections.

More recently similar results were obtained, when the Levi form has comparable eigenvalues, by K. Koenig in [Koe02] and S. Cho in [Cho03], [Cho02b].

In [FKM90] C. L. Fefferman, J. J. Kohn and M. Machedon studied the case where the Levi form is locally diagonalizable near a point p_0 of the boundary. They solved the $\bar{\partial}_b$ -Neuman problem and deduced that if f is a $L^2(\partial\Omega)$ function which is locally in the classical Lipschitz space Λ_α (near p_0) then, for all $\varepsilon > 0$ it's Szegő projection Sf is locally (near p_0) in $\Lambda_{\alpha-\varepsilon}$ (an application of our theory will remove the loss of ε in this estimate and get, in fact, a better non-isotropic estimate).

The main idea of the present paper is to introduce a general notion of “extremal basis” of the complex tangent space at a boundary point of a pseudo-convex domain in \mathbb{C}^n , $n \geq 3$, generalizing the ε -extremal bases of the convex case. With this notion we define a class of pseudo-convex domains, containing all previously studied classes, called “geometrically separated”, for which a good family of extremal bases exists near a point of the boundary. The fundamental properties of an extremal basis allow one to prove that, for these domains, there exists an associated structure of homogeneous space on the boundary (and an extension of that structure inside the domain) which describes the complex geometry of the domain. An important property of domains which are geometrically separated at a boundary point is that this structure can be nicely localized (see the end of Section 2 for more details).

Moreover, when special pluri-subharmonic functions (called “adapted pluri-subharmonic functions” in this paper) exist, this structure is used to obtain sharp global and local estimates for the Bergman kernel, the Bergman and Szegő projection and the classical invariant metrics. The existence of such adapted pluri-subharmonic functions for geometrically separated domains is not evident in general. For example, if the domain is locally convex (or more generally lineally convex), this is

2000 *Mathematics Subject Classification.* Primary 32T25, Secondary 32T27.

Key words and phrases. finite type; extremal basis; complex geometry; adapted pluri-subharmonic function; Bergman and Szegő projections.

done using special support functions (see [DF99, McN02]) which cannot exist, in general, without convexity. Here we prove their existence, under an additional condition (which is satisfied, for example, when the Levi form is locally diagonalizable) on the extremal bases, for the domain and also for the localized one (see the end of Section 2 for more details).

2. NOTATIONS AND ORGANIZATION OF THE PAPER

In all the paper, $\Omega = \{\rho < 0\}$ denotes a bounded domain in \mathbb{C}^n , $n \geq 3$, with a \mathcal{C}^∞ boundary, and $\rho \in \mathcal{C}^\infty(\mathbb{C}^n)$ is a defining function of Ω such that $|\nabla\rho| = 1$ on $\partial\Omega$. We denote by $N = \frac{1}{|\nabla\rho|} \sum \frac{\partial\rho}{\partial\bar{z}_i} \frac{\partial}{\partial z_i}$ the unitary complex normal vector field to ρ (i.e. $N\rho \equiv 1$ and $\|N\| \equiv 1$).

For each point p of the boundary let us denote $T_p^{1,0}(\partial\Omega)$ the subbundle of $T_p(\partial\Omega)$ of tangential complex vectors and $T_p^{0,1}(\partial\Omega)$ its conjugate. As usual, we will say that a family $(L_i)_{1 \leq i \leq n-1}$ of \mathcal{C}^∞ vector fields is a basis of the complex tangent space at $\partial\Omega$ in an open neighborhood $V \subset \partial\Omega$ of a point $p_0 \in \partial\Omega$ if it is a basis of sections of $T^{1,0}(\partial\Omega)$ in V (i.e. $L_i(\rho) \equiv 0$ in V , a condition which is independent of the defining function).

Clearly, every \mathcal{C}^∞ vector field L in an open neighborhood $V \subset \partial\Omega$ can be extended to an open neighborhood $V(p_0) \subset \mathbb{C}^n$ so that $L(\rho) \equiv 0$ in $V(p_0)$. Of course this extension depends on the defining function ρ , but all the stated results will be independent of such a choice. Thus, in all the paper, the tangent vector fields considered in $V(p_0)$ are always supposed to annihilate ρ in $V(p_0)$, and we will use the terminology of “vector fields tangent to ρ ” for this property.

Let L and L' be two $(1,0)$ vector fields tangent to ρ . The bracket $[L, \bar{L}']$ being tangent to ρ , can be written

$$[L, \bar{L}'] = 2\sqrt{-1}c_{LL'}T + L''$$

where T is the imaginary part of N and $L'' \in T_p^{1,0}(\partial\Omega) \oplus T_p^{0,1}(\partial\Omega)$. Thus $c_{LL'} = [L, \bar{L}'](\partial\rho) = \langle \partial\rho; [L, \bar{L}'] \rangle$. The Levi form of $\partial\Omega$ at p is defined as the hermitian form whose value at (L, \bar{L}') is the number $c_{LL'}$. The pseudo-convexity of Ω means that this hermitian form is non-negative. If $(L_i)_{1 \leq i \leq n-1}$ is a local basis of $(1,0)$ vector fields tangent to ρ , then $(c_{L_i L_j})_{i,j}$ is the matrix of the Levi form in the given basis. This matrix will be generally denoted $(c_{ij})_{i,j}$.

Let $p_0 \in \Omega$ and $V(p_0)$ be a neighborhood of p_0 in \mathbb{C}^n . If W is a set of $\mathcal{C}^\infty(V(p_0))$ $(1,0)$ complex vector fields, then $\mathcal{L}(W)$ denotes the set of all lists $\mathcal{L} = (L^1, \dots, L^k)$ such that $L^j \in W \cup \bar{W}$, and, for $l \in \mathbb{N}$, $\mathcal{L}_l(W)$ denotes the set of such lists \mathcal{L} of length $|\mathcal{L}| = k \in \{0, 1, \dots, l\}$. If W contains only one vector field L , then we will write $\mathcal{L}(L)$ and $\mathcal{L}_l(L)$ instead of $\mathcal{L}(\{L\})$ and $\mathcal{L}_l(\{L\})$. Moreover, if $|\mathcal{L}| = k \geq 2$, we denote

$$\mathcal{L}(\partial\rho) = L^1 \dots L^{k-2} \left(\left\langle \partial\rho, [L^{k-1}, L^k] \right\rangle \right).$$

Note that if L^{k-1} and L^k are both $(1,0)$ or both $(0,1)$ then $\langle \partial\rho, [L^{k-1}, L^k] \rangle$ is identically zero. Thus if $\mathcal{L}(\partial\rho)$ is not identically zero, it is equal to \pm a derivative of the value taken by the Levi form on (L^{k-1}, L^k) or (L^k, L^{k-1}) .

Let L be a $\mathcal{C}^\infty(V(p_0))$ $(1,0)$ complex vector field tangent to ρ and $M \geq 2$ be an integer. We define the weight $F_M^\Omega(L, p, \delta) = F^\Omega(L, p, \delta) = F^\Omega(L)$ associated to L at the point $p \in V(p_0)$ and to $\delta > 0$ by

$$F^\Omega(L, p, \delta) = \sum_{\mathcal{L} \in \mathcal{L}_M(L)} \left| \frac{\mathcal{L}(\partial\rho)(p)}{\delta} \right|^{2/|\mathcal{L}|} = 2 \sum_{\tilde{\mathcal{L}} \in \mathcal{L}_{M-2}(L)} \left| \frac{\tilde{\mathcal{L}}(c_{LL})}{\delta} \right|^{2/|\tilde{\mathcal{L}}|+2}.$$

where $\tilde{\mathcal{L}}(c_{LL}) = L^1 \dots L^k(c_{LL})$ if $\tilde{\mathcal{L}} = (L^1, \dots, L^k)$. Moreover, for the complex normal direction N we define $L_n = N$ and $F^\Omega(L_n, p, \delta) = F^\Omega(N, p, \delta) = \delta^{-2}$. When there is no ambiguity (typically when there is only one domain) we will omit the superscript Ω .

Note that, with the conditions on ρ , the functions $\mathcal{L}(\partial\rho)$ restricted to $\partial\Omega$ do not depend on the choice of the defining function ρ . By the finite type hypothesis, for δ small, the weights will be large. Thus if we consider them in δ -strips near the boundary, they are intrinsically attached to the boundary of the domain and do not depend on the choice of the defining function ρ .

In all the paper the defining function ρ of Ω is fixed and the number M also. When we say that some number depends on “ ϑ ” and on “the data”, we mean that it depends on “ ϑ ”, n , M , and ρ but neither on the point p in $V(p_0)$ nor on $\delta \leq \delta_0$.

If $\mathcal{B} = \{L_1, \dots, L_{n-1}\}$ is a \mathcal{C}^∞ basis of $(1,0)$ vector fields tangent to ρ in $V(p_0)$, and $\mathcal{L} \in \mathcal{L}(\mathcal{B} \cup \{N\})$, we denote

$$F(p, \delta)^{\mathcal{L}/2} = \prod_{i=1}^n F(L_i, p, \delta)^{l_i/2},$$

where $l_i = l_i(\mathcal{L})$ is the number of times L_i or \bar{L}_i appears in \mathcal{L} , $i \leq n-1$, and $l_n = l_n(\mathcal{L})$ the number of times N or \bar{N} appears in \mathcal{L} (and thus $|\mathcal{L}| = k = \sum_{i=1}^n l_i$).

The organization of the paper is as follows:

In Section 3 we define the notion of extremal basis and give some examples. Then we give their basic properties and, in Section 3.3, we prove a fundamental property of an extremal basis at a point of finite type: there exists a coordinate system which is adapted to the basis in the sense that all the derivatives of the matrix of the Levi form (in that basis) are controlled

by the weights attached to the basis. We give also some sufficient conditions of extremality for a given basis, useful for some examples. Finally, in Section 3.5 we show how the existence of extremal bases can be localized in the sense that, near a boundary point p_0 of Ω of finite type, if there exist extremal bases at every boundary points near p_0 , then one can construct a small pseudo-convex domain D of finite type inside the original domain, containing a piece of the boundary of Ω in its boundary such that there exist extremal bases at every point of the boundary of D .

In Section 4 we define the notion of geometrically separated domains at a point p_0 of its boundary and give examples. Then we show that a geometrically separated domain is automatically equipped with a local structure of homogeneous space on its boundary. In Section 4.3 we prove that the structure of geometrically separated domain can always be localized (in the sense described above).

In Section 5 we study the existence of pluri-subharmonic functions adapted to a given geometrically separated domain. In particular, we prove their existence when the domain is “strongly” geometrically separated at a point p_0 of its boundary, and we prove that, in this case, such functions exist for the localized domain at every point of its boundary.

In the last Section (6) we show that all the sharp global and local results for Bergman kernel, Bergman and Szegő projections and invariant metrics can be established for geometrically separated domains when there exist adapted pluri-subharmonic functions. The local sharp estimate of the Szegő projection when the Levi form is locally diagonalizable is an example of these results.

3. EXTREMAL BASES

3.1. Definition and examples.

Definition 3.1. Let Ω and $V(p_0)$ defined on Section 2. Let $\mathcal{B} = \{L_1, \dots, L_{n-1}\}$ be a \mathcal{C}^∞ basis of $(1, 0)$ vector fields tangent to ρ in $V(p_0)$ and M an integer. Let $p \in V(p_0)$ and $0 < \delta$. We say that $\mathcal{B} = \{L_1, \dots, L_{n-1}\}$ is (M, K, p, δ) -extremal (or simply (K, p, δ) -extremal or K -extremal) if the \mathcal{C}^{2M} norms, in $V(p_0)$, of all L_i are bounded by K , the Jacobian of \mathcal{B} is bounded from below by $1/K$ on $V(p_0)$, and the two following conditions are satisfied:

EB₁ For any vector field L of the form $L = \sum_{i=1}^{n-1} a_i L_i$, $a_i \in \mathbb{C}$, we have

$$\frac{1}{K} \sum_{i=1}^{n-1} |a_i|^2 F(L_i, p, \delta) \leq F(L, p, \delta) \leq K \sum_{i=1}^{n-1} |a_i|^2 F(L_i, p, \delta).$$

EB₂ For all indexes i, j, k such that $i, j < n$, $k \leq n$ and all lists \mathcal{L} of $\mathcal{L}_M(\mathcal{B} \cup \{N\})$,

$$F(L_k, p, \delta)^{1/2} \left| \mathcal{L} a_{i,j}^k(p) \right| \leq K F(p, \delta)^{\mathcal{L}/2} F(L_i, p, \delta)^{1/2} F(L_j, p, \delta)^{1/2},$$

where $a_{i,j}^k$ is the coefficient of the bracket $\left[\overleftarrow{L}_i, \overleftarrow{L}_j \right]$ in the direction \overleftarrow{L}_k (with $L_n = N$), and \overleftarrow{L}_i means L_i or \overline{L}_i .

Remark. In general this Definition depends of the choice of the defining function ρ . But note that, for $p \in \partial\Omega$, it does not: it depends only on the restriction of \mathcal{B} to $\partial\Omega \cap V(p_0)$.

Example 3.1.

- (1) *Locally lineally convex domains.* A first example of extremal basis concerns the case of a locally convex domain near a point of finite type: it can be easily shown, using the work of Mc Neal [McN94] (see also [Hef04]), that if Ω is convex near a point of finite type $p_0 \in \partial\Omega$, if the canonical coordinate system is chosen so that the last coordinate is the complex normal at p_0 , and, if P is the projection onto the complex tangent space of the defining function of Ω parallel to the last coordinate, then for each point p in a small neighborhood of p_0 , and each $\delta \leq \delta_0$, the P -projection of the first $n - 1$ vectors of the Mc Neal δ -extremal basis at p (c.f. [BCD98, McN94]) is (K, p, δ) -extremal in our sense for a constant K depending only on the data.

More generally, the same thing can be done for *locally lineally convex domains* using the work of Conrad, M. [Con02] (recall that Ω is said lineally convex at a point $p \in \partial\Omega$ if there is a neighborhood U of p such that the intersection of the complex tangent space to $\partial\Omega$ at p with $\Omega \cap U$ is empty; see [Kis98, DF03] for the precise definition and a useful characterization). Some details are given in Section 7.1.

- (2) *Levi form with comparable eigenvalues.* A second example is given by a pseudo-convex domain having a point of finite type $p_0 \in \partial\Omega$ where the eigenvalues of the Levi form are comparable (see [Koe02, Cho02b, Cho03, Cho02a]). Indeed, in [Cho03] it is proved that any (normalized) basis of the complex tangent space is K -extremal for a well controlled constant K .
- (3) *Locally diagonalizable Levi form.* In Section 3.4 we will show that if at a point of finite type $p_0 \in \partial\Omega$ the Levi form is locally diagonalizable then the basis diagonalizing the Levi form is K -extremal for a constant K depending only on the data (in fact, this basis is K -strongly-extremal (see Definition 3.5) for every constant $\alpha > 0$ with $\delta \leq \delta_0$, δ_0 small depending on α).
- (4) *Localization.* Another important example will be given in Section 3.5: for any $\tau > 0$ there exists $M(\tau)$ such that if a family of $(M(\tau), K, p, \delta)$ -extremal bases exists in a neighborhood of a boundary point p_0 , of finite type τ , of Ω , then

one can construct a small smooth pseudo-convex domain D contained in Ω and containing a neighborhood of p_0 in $\partial\Omega$ in its boundary and for which there exists an $(M(\tau), K', q, \delta)$ -extremal basis at every points $q \in \partial D$.

3.2. Basic properties of extremal bases. The first property states that an extremal basis at p can be orthogonalized at the point p :

Proposition 3.1. *For any K there exists a constant K' depending only on K and the data such that, if \mathcal{B} is a basis of complex $(1, 0)$ vector fields tangent to ρ in an open set $V(p_0)$ which is (K, p, δ) -extremal, then there exists a basis \mathcal{B}' , orthonormal at p which is (K', p, δ) -extremal.*

Proof. We can suppose that the vector fields L_i of \mathcal{B} are ordered such that $F(L_{i+1}, p, \delta) \leq F(L_i, p, \delta)$, for $i < n - 1$. Then, using the Gram-Schmidt process, we first define a basis \mathcal{B}_1 by decreasing induction, $L_i^1 = \sum_{j=i}^{n-1} \alpha_i^j L_j$, $\alpha_i^j \in \mathbb{C}$, and $\sum |\alpha_i^j|^2 = 1$. The determinant condition implies that there exists $c > 0$ such that $|\alpha_i^i| > c$. Then

$$F(L_i^1, p, \delta) \simeq_K \sum_{j \geq i} |\alpha_i^j|^2 F(L_j, p, \delta) \simeq_K F(L_i, p, \delta).$$

Now, let $L = \sum_i a_i L_i^1$ be a linear combination, with constant coefficients, of the L_i^1 . Then

$$F(L, p, \delta) \simeq_K \sum_k \left| \sum_{i \leq k} a_i \alpha_i^k \right|^2 F(L_k, p, \delta) \simeq_K \sum_k |a_k|^2 F(L_k, p, \delta),$$

using that $|\sum_{i \leq k} a_i \alpha_i^k| \geq c |a_k| - \sum_{i < k} |a_i|$ and the fact that the $F(L_k, p, \delta)$ are decreasing. This proves EB_1 for \mathcal{B}_1 .

Note now that the decreasing property shows that property EB_2 for \mathcal{B} trivially implies the same property for \mathcal{B}_1 because L_i^1 involves only fields L_j^1 for $j \geq i$.

Finally, define \mathcal{B}' by $L_i' = L_i^1 / \|L_i^1\|$. The condition on the \mathcal{C}^{2M} norm of the vectors L_i immediately implies the result. \square

Let us now prove that the mixed derivatives of the Levi form in the directions of an extremal basis are controlled by the pure ones, that is by the weights associated to the vector fields of the basis:

Proposition 3.2. *Let $\mathcal{B} = \{L_i, 1 \leq i \leq n - 1\}$ be a \mathcal{C}^∞ basis of complex $(1, 0)$ vector fields tangent to ρ in $V(p_0)$ which is (K, p, δ) -extremal for a fixed $\delta > 0$. Let \mathcal{L} be a list of vector fields belonging to $\mathcal{L}_M(\mathcal{B} \cup \{N\})$. Then there exists a constant $C > 0$ depending only on Ω and K such that $|\mathcal{L}(\partial\rho)(p)| \leq C\delta F^{\mathcal{L}/2}(p, \delta)$.*

Proof. Recall the notation $c_{ij} = \langle \partial\rho, [L_i, \overline{L}_j] \rangle$.

Lemma 3.2.1. *With the previous notations (and the definition of the coefficients a_{ij}^s given in Definition 3.1):*

$$\begin{aligned} L_j c_{ik} &= L_i c_{jk} + \sum \alpha_{ki}^s c_{js} - \sum \alpha_{ij}^s c_{sk} - \sum \alpha_{jk}^s c_{is}, \\ \overline{L}_j c_{ik} &= \overline{L}_k c_{ij} + \sum \alpha_{ik}^s c_{sj} + \sum \alpha_{ji}^s c_{sk} - \sum \alpha_{kj}^s c_{is}. \end{aligned}$$

Proof. The first formula is simply obtained considering the coefficient of $\Im m N$ in the Jacobi's identity applied to the bracket $[L_j, [L_i, \overline{L}_k]]$, and the second by the same way using $[\overline{L}_j, [L_i, \overline{L}_k]]$. \square

The proof of Proposition 3.2 is done by induction on the length of the lists. Suppose first $|\mathcal{L}| = 2$. Hypothesis EB_1 imply that, for all numbers a and b and all index i and j ,

$$\left| |a|^2 c_{ii} + |b|^2 c_{jj} + a\overline{b}c_{ij} + \overline{a}bc_{ji} \right| \lesssim \delta \left(|a|^2 F_i + |b|^2 F_j \right).$$

Suppose both F_i and F_j are non zero. Taking $a = F_j^{1/2} F_i^{-1/2} \lambda$ and $b = \mu$, $|\lambda|$ and $|\mu|$ less than 1, the equivalence of norms in finite dimensional spaces gives the result. If $F_i = 0$ or $F_j = 0$ a similar argument gives $c_{ij} = c_{ji} = 0$.

To continue the proof, we need the following notation: if $\mathcal{L} \in \mathcal{L}(\mathcal{B} \cup \{N\})$, we denote by l_i^1 (resp. l_i^2) the number of times L_i (resp. \overline{L}_i) appears in \mathcal{L} (thus $l_i = l_i^1 + l_i^2$).

For lists of greater length, we prove, at the same time, by induction, the estimate and the following Lemma:

Lemma 3.2.2. *Let \mathcal{L} and \mathcal{L}' be two lists of $\mathcal{L}_M(\mathcal{B} \cup \{N\})$, $\mathcal{L}(\partial\rho) = \mathcal{L}_1 c_{ij}$ and $\mathcal{L}'(\partial\rho) = \mathcal{L}'_1 c_{kl}$, such that $l_i^1 = l_i'^1$, $l_i^2 = l_i'^2$. Then $\mathcal{L}(\partial\rho) \simeq \mathcal{L}'(\partial\rho)$ in the sense that*

$$\mathcal{L}(\partial\rho) - \mathcal{L}'(\partial\rho) = \sum_{|\widetilde{\mathcal{L}}| < |\mathcal{L}|} a_{\widetilde{\mathcal{L}}} \widetilde{\mathcal{L}}(\partial\rho),$$

where $a_{\widetilde{\mathcal{L}}}$ satisfies $F^{\widetilde{\mathcal{L}}/2} |\mathcal{L}'' a_{\widetilde{\mathcal{L}}}| \lesssim \delta F^{(\mathcal{L} + \mathcal{L}'')/2}$, $\forall \mathcal{L}'' \in \mathcal{L}_M(\mathcal{B} \cup \{N\})$, the constant depending only on K and the data.

Suppose thus the estimates and the Lemma proved for all lists of length less than or equal to N .

First, we prove Lemma 3.2.2 for lists of length $N + 1$. Let us write $\mathcal{L}(\partial\rho) = \mathcal{L}_1 c_{ij}$ and $\mathcal{L}'(\partial\rho) = \mathcal{L}_2 c_{kl}$. Then three cases can happen:

- (1) $(i, j) = (k, l)$;
- (2) $i \neq k, j \neq l$;
- (3) $i \neq k$ and $j = l$ or $i = k$ and $j \neq l$.

The first case is a trivial consequence of EB₂. For the second, the hypothesis on the length and case (1) imply that there exists a list $\tilde{\mathcal{L}}$ such that $\mathcal{L}_1 c_{ij} \simeq \tilde{\mathcal{L}} L_k \bar{L}_l c_{ij}$ and $\mathcal{L}_2 c_{kl} \simeq \tilde{\mathcal{L}} L_i \bar{L}_j c_{kl}$, in the sense of Lemma 3.2.2. By Lemma 3.2.1 and EB₂, $\bar{L}_j c_{ij} \simeq \bar{L}_j c_{il}$. The result is obtained using another time EB₂, Lemma 3.2.1 and the induction hypothesis. The third case is similar.

Now we prove the estimate of the Proposition for lists of length $N + 1$. Suppose that the vector fields are ordered so that there exists an integer $n_0 \in \{0, \dots, n-1\}$ such that, for $k \leq n_0$, $F_k \neq 0$, and, for $n-1 \geq k > n_0$, $F_k = 0$. Let $L = \sum a_j L_j$, $a_j = \varepsilon \lambda_j F_{n_0}^{1/2} F_j^{-1/2}$ if $j \leq n_0$ and $a_j = \lambda_j$ if $j > n_0$, with $|\lambda_j| \leq 1$. If we apply the extremality property to $F(L)$, then we obtain, for example, for all $k \leq N-1$,

$$\sup_{|\lambda_j| \leq 1} \left| L^k \bar{L}^{N-k-1} c_{LL} \right| \lesssim \delta \varepsilon^{(N+1)/2} F_{n_0}^{(N+1)/2}$$

with the convention $F_{n_0} = 0$ if $n_0 = 0$. Writing $L^k \bar{L}^{N-k-1} c_{LL} = \sum C_{\alpha\beta} \lambda^\alpha \bar{\lambda}^\beta$, the equivalence of norms in finite dimensional spaces gives, when $\varepsilon \rightarrow 0$,

$$(3.1) \quad \begin{aligned} C_{\alpha\beta} &= 0 && \text{if there exists } j > n_0 \text{ such that } \alpha_j + \beta_j > 0, \\ |C_{\alpha\beta}| &\lesssim \delta F^{(\alpha+\beta)/2} && \text{otherwise.} \end{aligned}$$

Let $\mathcal{E}_{\alpha\beta}$ be the set of lists \mathcal{L} such that $l_i^1(\mathcal{L}) = \alpha_i$ and $l_i^2 = \beta_i$. Then $C_{\alpha\beta} = \sum_{\mathcal{L} \in \mathcal{E}_{\alpha\beta}} \mathcal{L}(\partial\rho)$. Now, Lemma 3.2.2 and the induction hypothesis give the required estimation for each list in $\mathcal{E}_{\alpha\beta}$ and finishes the proof of the Proposition. \square

The statement of the last Proposition is not really a statement on the vector fields of an extremal basis but on the linear space generated by an extremal basis. In fact the following Proposition is easily proved:

Proposition 3.3. *In the conditions of Proposition 3.2, there exists a constant C such that, if L'_j , $1 \leq j \leq k$ are vector fields belonging to the linear space generated by the extremal basis $(L_i)_i$ then for every $\mathcal{L} \in \mathcal{L}_M(L'_1, \dots, L'_k)$, if L'_j or \bar{L}'_j appear l'_j times in \mathcal{L} , $|\mathcal{L}(\partial\rho)(p)| \lesssim \delta \prod_j F(L'_j, p, \delta)^{l'_j/2}$.*

3.3. Adapted coordinates system for points of finite 1-type.

3.3.1. *Definition of an adapted coordinate system and statement of the main result.* Let $p_0 \in \partial\Omega$ and $V(p_0)$ a neighborhood of p_0 in \mathbb{C}^n .

Definition 3.2. A basis $\mathcal{B} = (L_1, \dots, L_{n-1})$ of sections of $(1, 0)$ complex tangent vector fields to ρ in $V(p_0)$ and a coordinate system in \mathbb{C}^n , $z = \Phi_p^\delta(Z)$, are called (M, K, δ) -adapted (or simply (K, δ) -adapted) at the point p in $V(p_0)$ if Φ_p^δ and $(\Phi_p^\delta)^{-1}$ are polynomial (of degree less than $(2M)^{n-1}$) diffeomorphisms of \mathbb{C}^n centered at p (i.e. $\Phi_p^\delta(p) = 0$) satisfying (with the notation $F_i = F_i(p, \delta) = F(L_i, p, \delta)$):

- (1) The coefficients of the polynomials of Φ_p^δ and $(\Phi_p^\delta)^{-1}$ (and the Jacobians of Φ_p^δ and $(\Phi_p^\delta)^{-1}$) are bounded by K ;
- (2) For all $|\alpha| \leq 2M$, $\frac{\partial^\alpha (\rho \circ (\Phi_p^\delta)^{-1})(0)}{\partial z^\alpha} = \frac{\partial^\alpha (\rho \circ (\Phi_p^\delta)^{-1})(0)}{\partial \bar{z}^\alpha} = 0$, $z' = (z_1, \dots, z_{n-1})$;
- (3) If $L_i = \sum \alpha_i^j \frac{\partial}{\partial z_j}$, then $\alpha_i^j(0) = \delta_{ij}$ and, for all $\mathcal{L} \in \mathcal{L}_M(\mathcal{B} \cup \{N\})$,

$$\left| \mathcal{L} \alpha_i^j \right| \leq K \text{ in } \Phi_p(V(p_0)) \text{ and } F_j^{1/2} \left| \mathcal{L} \alpha_i^j(0) \right| \leq K F_i^{1/2} F^{\mathcal{L}/2};$$

- (4) For all (α, β) , $|\alpha + \beta| \leq M$, $\left| \frac{\partial^{\alpha+\beta} (\rho \circ (\Phi_p^\delta)^{-1})(0)}{\partial z^\alpha \partial \bar{z}^\beta} \right| \leq K \min \left\{ \delta F^{(\alpha+\beta)/2}, 1 \right\}$.

One of our main goals is to prove the following existence Theorem:

Theorem 3.1. *Suppose p_0 is of finite 1-type τ , and choose an integer M larger than $2 \left(\frac{2(\frac{\tau}{2})^{n-1} + 1}{2} \right)^{n-1}$. For any positive constant K , there exist a constant $\delta_0 > 0$, a neighborhood $V(p_0)$, both depending on the data, and a constant K' depending on K and the data such that if $\mathcal{B} = \{L_i, 1 \leq i \leq n-1\}$ is a \mathcal{C}^∞ basis of $(1, 0)$ complex vector fields tangent to ρ in $V(p_0)$ which is (M, K, p, δ) -extremal at a point $p \in V(p_0) \cap \partial\Omega$, then there exists a coordinate system $(z_i)_{1 \leq i \leq n}$ centered at p which is (K', δ) -adapted to \mathcal{B} .*

The proof is divided in two steps: for the first one, in the next Section, we work without the assumption of finite type and we construct an adapted coordinate system using modified weights; in the second one, which is Section 3.3.3, we use the finite type hypothesis to deduce the Theorem.

3.3.2. *Construction of an adapted coordinate system.* In this Section we suppose that the integer M is fixed. Let $p \in V(p_0)$ and $\delta > 0$. Suppose $\mathcal{B} = (L_1, \dots, L_{n-1})$ is a basis of $(1, 0)$ vector fields tangent to ρ in $V(p_0)$, satisfying the following properties:

- (A) The $\mathcal{C}^{2M}(V(p_0))$ norms of all L_i are bounded by K and \mathcal{B} is ordered so that $F(L_{i+1}, p, \delta) \leq F(L_i, p, \delta)$.
- (B) Let $p \in W(p_0) \Subset V(p_0)$ and $\delta > 0$. Denoting $\tilde{F}_i = F_i + 1 = F(L_i, p, \delta) + 1$:
 - (B₁) For all list $\mathcal{L} \in \mathcal{L}_M(\mathcal{B} \cup \{N\})$, $|\mathcal{L}(\partial\rho)(p)| \leq K\delta\tilde{F}(p, \delta)^{|\mathcal{L}|/2}$;
 - (B₂) \mathcal{B} satisfies condition EB₂ of Definition 3.1 with the $F(L_s, p, \delta)$ replaced by the \tilde{F}_s .

Then under these hypotheses, we have:

Proposition 3.4. *There exists a constant K' depending on K , M and the data (but neither on p nor on δ) such that there exists a (M, K', δ) -adapted coordinate system to \mathcal{B} at p in the sense of Definition 3.2, the weights $F(L_i, p, \delta)$ being replaced by \tilde{F}_i .*

Proof. In [CD06b] (Prop 3.2, p. 85) we proved that hypothesis (A) implies the existence of a coordinate system $\Phi_{p, \delta}$ satisfying conditions (1) and (2) of Definition 3.2 and

$$(3.2) \quad \left\{ \begin{array}{l} \text{For } j < i < n, \text{ and } \alpha = (\alpha_1, \dots, \alpha_{n-1}) \in \mathbb{N}^{n-1} \text{ such that } |\alpha| \leq M, \alpha_p = 0 \text{ if } p > i \text{ or } p \leq j, \\ \frac{\partial^\alpha a_i^j(0)}{\partial z^{\alpha'}} = 0. \end{array} \right.$$

We now prove that under condition (B) the two last properties of Definition 3.2 (with the \tilde{F}_i) are satisfied. This follows quite closely the ideas of p. 87-90 of [CD06b], but, as the context here is more general and as it is a fundamental tool, we write it completely.

Let $\mathcal{L} \in \mathcal{L}(\mathcal{B} \cup \{N\})$ be considered as a differential operator. Denoting by $D^{\alpha\beta}$ the derivative $\frac{\partial^{\alpha+\beta}}{\partial z^\alpha \partial \bar{z}^\beta}$ in the coordinate system $z = \Phi_p^\delta$, it is easy to see that, if $|\mathcal{L}| = S$,

$$\mathcal{L} = \sum_{\substack{m \in \mathbb{N}^n \\ 1 \leq |m| \leq S}} \sum_{\alpha_i + \beta_i = m_i} c_{\alpha\beta}^{\mathcal{L}} D^{\alpha\beta}$$

where

$$c_{\alpha\beta}^{\mathcal{L}} = c_{\alpha\beta} = \sum_{p=1}^S \sum_{i_1, \dots, i_p} * a_{j_1}^{i_1} \cdots a_{j_p}^{i_p} \prod_{k=p+1}^S D^{s_k} \binom{\alpha}{j_k}$$

where the summation in the second formula is taken over the derivatives associated to the multiindex s_k satisfying $\sum_{k=p+1}^S s_k + (m_1, \dots, m_n) = \sum_{k=1}^S \chi(i_k)$, $\sum_{k=1}^S \chi(j_k) = (l_1, \dots, l_{n-1}, l_n)$ and the coefficients $*$ are absolute constants. The following Lemma is then easily established:

Lemma 1. *If for all $s \in \mathbb{N}^n$, $|s| \leq S$, we have $|D^s a_j^i(0)| \lesssim_{K_1} \tilde{F}^{s/2} \tilde{F}_j^{1/2} \tilde{F}_i^{-1/2}$, then*

$$(3.3) \quad |c_{\alpha\beta}(0)| \lesssim \tilde{F}^{|\mathcal{L}|/2} \tilde{F}^{-\frac{\alpha+\beta}{2}}.$$

To fix notations, recall that if f is a \mathcal{C}^2 function and L and L' are two vector fields, then $\langle \partial \bar{\partial} f; L, \bar{L} \rangle = \bar{L}' L f + [L, \bar{L}'](\partial f)$, and, in particular, if $L\rho = 0$, $\langle \partial \bar{\partial} \rho; L, \bar{L} \rangle = [L, \bar{L}'](\partial\rho) = c_{LL}$, where c_{LL} is the coefficient of the Levi form in the direction L . In all the proof that follows, we denote $[L_i, \bar{L}_j](\partial\rho) = c_{ij}$.

To state the second Lemma let us introduce the notation $\tilde{\rho} = \rho \circ (\Phi_p^\delta)^{-1}$:

Lemma 2.

- (1) For every multiindex l , $|l| \leq 2M$, we have $|D^l \tilde{\rho}(0)| \lesssim \delta \tilde{F}^{l/2}$, where D^l is any derivative $\frac{\partial^{|l|}}{\partial z^\alpha \partial \bar{z}^\beta}$ with $|\alpha + \beta| = l$.
- (2) For every multiindex $m \neq (0, \dots, 0)$, $|m| < M$, and every i, j , $|D^m a_i^j(0)| \lesssim \tilde{F}^{m/2} \tilde{F}_i^{1/2} \tilde{F}_j^{-1/2}$.

Proof. Note first that, for (2), it suffices to get the estimate for $D^m a_i^j(0)$ and that the estimate (1) (resp. (2)) is trivial if $l_n > 0$ (resp. $m_n > 0$) (recall $F_n = \delta^{-2}$ and the fact that the \mathcal{C}^{2M} norms of the fields L_i are controlled). We then suppose $l_n = m_n = 0$. The proof is done by induction: the induction hypothesis \mathcal{P}_{k_0} is the two conclusions of the proposition for $|l| \leq k_0$ and $|m| < k_0$.

Remark first that \mathcal{P}_{k_0} and the first property of \mathcal{P}_{k_0+1} imply the second property of \mathcal{P}_{k_0+1} for $j = n$: this is evident if $i = j = n$ and, if $i < j = n$, $L_i r \equiv 0$ implies

$$a_i^n = \left(\frac{\partial \tilde{\rho}}{\partial z_n} \right)^{-1} \sum_{k=1}^{n-1} a_i^k \frac{\partial \tilde{\rho}}{\partial z_k},$$

and the result is clear because $\frac{\partial \tilde{\rho}}{\partial z_k}(0) = 0$ for $k < n$.

Moreover, note also that, the weights \tilde{F}_i , $i \leq n-1$, being “decreasing”, the second inequality of \mathcal{P}_{k_0} is trivial if $i \leq j < n$ and if $i = n$. Thus it suffices to prove this inequality when $j < i < n$.

Let us now prove \mathcal{P}_{k_0} by induction. The case $k_0 = 1$ is trivial. Let us study first the case $k_0 = 2$. By definition of the coordinate system, $\frac{\partial^2 \tilde{\rho}}{\partial z_i \partial \bar{z}_j}(0) = 0$, and, using the notations and remarks stated before the statement of the Lemma, we have

$$(3.4) \quad a_i^j \overline{a_j^i} \frac{\partial^2 \tilde{\rho}}{\partial z_i \partial \bar{z}_j} = c_{i\bar{j}} - \sum_{(k,p) \neq (i,j)} a_i^k \overline{a_j^p} \frac{\partial^2 \tilde{\rho}}{\partial z_k \partial \bar{z}_p}$$

which implies $\frac{\partial^2 \tilde{\rho}}{\partial z_i \partial \bar{z}_j}(0) = c_{i\bar{j}}(0)$ and gives the first inequality by definition of F . To prove the second inequality, let us look at the definition of the functions a_{ij}^k . Writing the bracket $[L_i, \overline{L_p}]$ with the coordinate system and taking the component of $\frac{\partial}{\partial z_j}$, we get

$$(3.5) \quad \sum_{k'=1}^{n-1} a_{i\bar{p}}^{k'} a_{k'}^j = - \sum_{k=1}^n \overline{a_p^k} \frac{\partial}{\partial \bar{z}_k} (a_i^j) - c_{i\bar{p}} a_n^j.$$

Extracting the term $\frac{\partial}{\partial \bar{z}_p} (a_i^j)$ and taking all at zero we obtain $\frac{\partial}{\partial \bar{z}_p} (a_i^j)(0) = a_{i\bar{p}}^j(0)$ and the inequality follows from (B_2) hypothesis.

We have now to consider $\frac{\partial a_i^j}{\partial z_q}$. If $q \leq j$, the inequality comes from the decreasing property of \tilde{F}_k , and, if $j < q \leq i$, this derivative is zero at the origin by the properties of the coordinate system. Suppose then $j < i < q$. Looking at the Lie bracket $[L_i, L_q]$ and taking the component of $\frac{\partial}{\partial z_j}$, we obtain

$$(3.6) \quad a_i^j \frac{\partial}{\partial z_i} (a_q^j) - a_q^j \frac{\partial}{\partial z_q} (a_i^j) = \sum_{k \neq q} a_q^k \frac{\partial}{\partial z_k} (a_i^j) - \sum_{k \neq i} a_i^k \frac{\partial}{\partial z_k} (a_q^j) + \sum_{p=1}^{n-1} a_{i\bar{q}}^p a_p^j,$$

and then, at the origin, $\frac{\partial}{\partial z_q} (a_i^j)(0) = \frac{\partial}{\partial z_i} (a_q^j)(0) - a_{i\bar{q}}^j(0) = -a_{i\bar{q}}^j(0)$, by the properties of the coordinate system, and the conclusion comes again from (B_2) . This proves \mathcal{P}_2 .

Let us now suppose \mathcal{P}_{k_0} verified ($k_0 < 2M$). Let $D^{\bar{l}}$ be a derivative of order $k_0 + 1$. If $D^{\bar{l}}$ is purely holomorphic or anti-holomorphic, then $D^{\bar{l}} \tilde{\rho}(0) = 0$. Then we suppose $D^{\bar{l}} = D^l \frac{\partial}{\partial z_i} \frac{\partial}{\partial \bar{z}_j}$, and we denote by $\tilde{\mathcal{L}} = \mathcal{L} L_i \overline{L_j}$ a list of vectors fields associated to $D^{\bar{l}}$ (in the obvious sense that, if $\partial/\partial z_i$ (resp. $\partial/\partial \bar{z}_i$) appears l_i (resp. \bar{l}_i) times in D^l then L_i (resp. $\overline{L_i}$) appears l_i (resp. \bar{l}_i) times in \mathcal{L}). Applying (3.4), we get

$$(3.7) \quad \begin{aligned} D^l \left(\frac{\partial^2 \tilde{\rho}}{\partial z_i \partial \bar{z}_j} \right) (0) &= \mathcal{L} c_{i\bar{j}}(0) - \sum_{\substack{l_1 \neq 0 \\ l_1 + l_2 = l}} * D^{l_1} \left(a_i^j \overline{a_j^i} \right) D^{l_2} \left(\frac{\partial^2 \tilde{\rho}}{\partial z_i \partial \bar{z}_j} \right) (0) \\ &- \sum_{(k,p) \neq (i,j)} D^l \left(a_i^k \overline{a_j^p} \frac{\partial^2 \tilde{\rho}}{\partial z_k \partial \bar{z}_p} \right) (0) \\ &- \sum_{|\alpha| + |\beta| < k_0 - 1} c_{\alpha\beta} D^{\alpha\beta'} (c_{i\bar{j}})(0), \end{aligned}$$

with $*$ = 0 or 1. The first term of the right hand side of (3.7) satisfies the desired inequality (i.e. $\lesssim \delta \tilde{F}^{l/2} \tilde{F}_i^{1/2} \tilde{F}_j^{1/2}$ in modulus) by (B_1) . For the second, l_1 being non 0, we can apply the induction hypothesis to $D^{l_2} \left(\frac{\partial^2 \tilde{\rho}}{\partial z_i \partial \bar{z}_j} \right) (0)$ to get the right estimate. The third term is of the same nature because, for $(k,p) \neq (i,j)$, $a_i^k \overline{a_j^p}(0) = 0$. If we replace $c_{i\bar{j}}$ by its expression in (3.4), the induction hypothesis \mathcal{P}_{k_0} implies directly (for $s < k_0 - 1$):

$$|D^s c_{i\bar{j}}(0)| \lesssim \delta F^{s/2} F_i^{1/2} F_j^{1/2},$$

and then, using Lemma 1 for $S = k_0$ (whose hypothesis are also verified by the induction hypothesis \mathcal{P}_k), we prove that the last term in (3.7) satisfies also the right estimate.

We finish now proving the second inequality of \mathcal{P}_{k_0+1} . It suffices to consider the case $j < i < n$. Let us first look at a derivative D^m of the form $D^m = D^s \frac{\partial}{\partial \bar{z}_p}$, $|s| = k_0 - 1$. Using formula (3.5), we can write

$$D^m a_i^j = D^s \left(\sum_{t=1}^{n-1} * a_{i\bar{p}}^t a_t^j - \sum_{t \neq p} * \overline{a_p^t} \frac{\partial}{\partial \bar{z}_t} (a_i^j) + * c_{i\bar{p}} a_n^j \right) = D^s(A) - D^s(B) + D^s(C),$$

where $*$ is equal to $\frac{1}{a_p^t}$. In $D^s(B)$, to get a non zero term at 0, $\overline{a_p^t}$ must be derivated because $p \neq t$; this gives derivatives of $\frac{\partial}{\partial \bar{z}_k} (a_i^j)$ of order $< k_0 - 1$ which are well controlled by the induction hypothesis and then $|D^s(B)(0)| \lesssim \tilde{F}^{m/2} \tilde{F}_i^{1/2} \tilde{F}_j^{1/2}$.

Consider now the terms $D^s \left(* a_{i\bar{p}}^t a_t^j \right)$.

Claim. For $|l| \leq k$, $D^l \left(a_{i\bar{p}}^l \right) \lesssim F_i^{1/2} F_p^{1/2} F_t^{-1/2} F^{l/2}$.

Proof of the Claim. We do it by induction on $|l|$. (B₂) proves the result for $|l| = 0$. Assume the claim proved for $|l| < k' \leq k_0 - 1$ and suppose $|l| = k'$. Then,

$$D^l a_{i\bar{p}}^l(0) = \mathcal{L}^l a_{i\bar{p}}^l(0) + \sum_{|s'| < l} c_{s'}(0) D^{s'} a_{i\bar{p}}^l(0).$$

But, by (B₂),

$$\left| \mathcal{L}^l a_{i\bar{p}}^l(0) \right| \lesssim F^{l/2} F_i^{1/2} F_p^{1/2} F_t^{-1/2},$$

and for the second term of the previous identity, we have $|s'| < l$ and we can apply the induction hypothesis and Lemma 1 whose hypotheses are satisfied, using \mathcal{P}_{k_0} , because $|l| \leq k_0$. \square

Then the estimate of $D^s \left(* a_{i\bar{p}}^k a_k^j \right)$ follows from the induction hypothesis \mathcal{P}_{k_0} because $|s| < k_0$. Thus

$$|D^s(A)(0)| \lesssim \tilde{F}^{m/2} \tilde{F}_i^{1/2} \tilde{F}_j^{1/2}.$$

Finally, the terms $D^s \left(* c_{i\bar{p}} a_n^j \right)$ satisfy also the good estimates because $a_n^j(0) = 0$ and, for $|s'| < k_0 - 1$, we have seen that $\left| D^{s'}(c_{i\bar{p}})(0) \right| \lesssim \delta \tilde{F}^{s'/2} \tilde{F}_i^{1/2} \tilde{F}_p^{1/2}$, and, the derivatives of a_n^j are controlled by the induction hypothesis \mathcal{P}_{k_0} .

To finish, we have to consider the case where D^m is a holomorphic derivative. Note that the inequality is trivial if $i \leq j$ or if there exists $k \leq j$ such that $m_k \neq 0$. Suppose then, for all $k \leq j$, $m_k = 0$ and $j < i < n$. Let q be the largest index such that $m_q > 0$. If $q \leq i$, we have $D^m a_i^j(0) = 0$ by the properties of the coordinate system. If $q > i$, then write $D^m = D^s \frac{\partial}{\partial z_i}$. To conclude it suffices then to use (3.6), the first Claim and the fact that $D^s \frac{\partial}{\partial z_i} \left(a_q^j \right)(0) = 0$ also by the properties of the coordinates system. This completes the proof of the Lemma. \square

To finish the proof of Proposition 3.4, it suffices to note that, in addition to the estimates of the coefficients $c_{\alpha\beta}^{\mathcal{L}}$ given by Lemma 1, we also have, for $|\alpha + \beta| \leq 2M$,

$$(3.8) \quad D^{\alpha\beta} = \sum_{1 \leq |\mathcal{L}| \leq |\alpha + \beta|} d_{\mathcal{L}}^{\alpha\beta} \mathcal{L},$$

with $\left| d_{\mathcal{L}}^{\alpha\beta}(0) \right| \lesssim \tilde{F}^{(\alpha + \beta)/2}(p, \delta) \tilde{F}^{-|\mathcal{L}|/2}$. \square

For an extremal basis we have thus proved (using Proposition 3.2):

Corollary. *If \mathcal{B} is (M, K, p, δ) -extremal, for δ small enough, there exists a coordinate system $(M, K'(K), \delta)$ -adapted to \mathcal{B} in the sense of Definition 3.2 with the weights F_i replaced by $\tilde{F}_i = F_i + 1$.*

3.3.3. *Proof of Theorem 3.1.* If p_0 is a point of finite 1-type τ , then, by a Theorem of D'Angelo (see [D'A82, Cat87]) there exists a neighborhood $U(p_0)$ such that, if $p \in \partial\Omega \cap U(p_0)$, then p is of finite 1-type less than $\tau' = 2 \left(\frac{\tau}{2} \right)^{n-1}$. We assume that $V(p_0) \subset U(p_0)$. Then, if \mathcal{B} is a (M, K, p, δ) -extremal basis, by the Corollary of Proposition 3.4 we have a coordinate system $\Phi_{p, \delta}$ adapted to \mathcal{B} in terms of the \tilde{F}_i . Suppose M larger than $2 \left(\frac{\tau'}{2} \right)^{n-1}$. Then, considering the manifold $\zeta \mapsto (0, \dots, 0, \zeta, 0, \dots, 0)$, $|\zeta| \leq \sigma$, Theorem 3.4 of [Cat87] (applied with a suitable constant σ) gives us a derivative of $\tilde{\rho} = \rho \circ \Phi_{p, \delta}$ which is bounded from below by a constant depending only on the data. The last property of Definition 3.2 shows thus that $\tilde{F}_i(p, \delta) \gtrsim \delta^{-2/M}$ with a constant depending only on the data, and, of course, the same is true for $F_i(p, \delta)$.

This proves the following essential Proposition:

Proposition 3.5. *Let $p_0 \in \partial\Omega$ be a point of finite 1-type τ . Let $M = M(\tau) = \left[2 \left(\frac{\tau}{2} \right)^{n-1} \right] + 1$. Then for any integer K there exist a real number $\delta_0 > 0$ and a constant C , depending on K and the data, such that, if there is a coordinate system (M, K, δ) -adapted to a basis $\mathcal{B} = (L_1, \dots, L_{n-1})$ at p_0 , then $F_M(L_i, p_0, \delta) \geq C \delta^{-2/M}$. In particular, if $\tau' = 2 \left(\frac{\tau}{2} \right)^{n-1}$ and $M' = M'(\tau) = \left[2 \left(\frac{\tau'}{2} \right)^{n-1} \right] + 1$, for any integer K there exists a neighborhood $V(p_0)$, a real number $\delta_0 > 0$ and a constant C (depending on τ , Ω and K) such that, for $p \in V(p_0) \cap \partial\Omega$ and $0 < \delta \leq \delta_0$, if there is a coordinate system (M', K, δ) -adapted to a basis $\mathcal{B} = (L_1, \dots, L_{n-1})$ at p , then $F(L_i, p, \delta) \geq C \delta^{-2/M'}$.*

This proves completely Theorem 3.1.

Remark 3.2. Note that the proofs show that if a basis \mathcal{B} satisfies only properties (A) and (B) of the beginning of Section 3.3.2, then, under the assumption of finite 1-type, the conclusions of Proposition 3.5 and Theorem 3.1 are still valid.

A simple consequence (which will be used in Section 3.5) of the minoration of the weights F_i is the following:

Lemma 3.1. *Suppose the point p_0 of finite 1-type τ . For any K , there exist two constants C and δ_0 , depending only on K , τ and the data, such that if $\mathcal{B} = \{L_i^{p,\delta}, i < n\}$ is (K, p, δ) -extremal, $p \in V(p_0) \cap \partial\Omega$, and (α_i) is a family of \mathcal{C}^∞ functions, of \mathcal{C}^{2M} norm $\leq K$ and $1/K \leq |\alpha_i| \leq K$, then the basis $\mathcal{B}_1 = \{L_i\}$, where $L_i = \frac{1}{\alpha_i} L_i^{p,\delta}$, is (C, p, δ) -extremal, and, moreover, $F(\sum a_i L_i, p, \delta) \simeq_C F(\sum a_i L_i^{p,\delta}, p, \delta)$, $a_i \in \mathbb{C}$.*

3.3.4. *Associated polydiscs and pseudo-balls for finite type points.* In this Section we assume p_0 is of finite 1-type τ and we choose $M = M'(\tau)$. Now we will associate to an adapted coordinate system some special ‘‘polydiscs’’ and give some related properties.

Definition 3.3. Let $W(p_0) \Subset V(p_0)$ small enough. Suppose that for some point $p \in W(p_0) \cap \partial\Omega$ and $0 < \delta$ there is a basis $\mathcal{B}(p, \delta) = \{L_i^{p,\delta}\}$ of $(1, 0)$ vector fields tangent to ρ in $V(p_0)$ satisfying conditions (A) and (B) (of Section 3.3.2) and let $\Phi_p^\delta = \Phi_p$ be a coordinate system which is (K, δ) -adapted to $\mathcal{B}(p, \delta)$. Then the functions $F(L_i, p, \delta) = F_i(p, \delta)$ do not vanish and, for $0 < c < 1$, we denote

$$\Delta_c(p, \delta) = \{z \in \mathbb{C}^n \text{ such that } |z_i| < cF_i^{-1/2}, 1 \leq i \leq n\},$$

and

$$B^c(p, \delta) = \Phi_p^{-1}(\Delta_c(p, \delta)) \cap V(p_0).$$

Taylor’s formula, Proposition 3.2 and Theorem 3.1 lead easily to the following properties (denoting $L_i = L_i^{p,\delta}$):

Proposition 3.6. *There exist three constants c_0 , K_0 and δ_0 , depending only on K and the data, such that the following properties hold:*

(1) *If $L_i = \sum \alpha_i^j \frac{\partial}{\partial z_j}$ and $\frac{\partial}{\partial z_j} = \sum b_j^i L_i$, $|\alpha + \beta| \leq M$, for $z \in \Delta_{c_0}(p, \delta)$,*

$$\left| D^{\alpha\beta} a_i^j(z) \right| \leq K_0 F^{(\alpha+\beta)/2}(p, \delta) F_i^{1/2}(p, \delta) F_j^{-1/2}(p, \delta),$$

$$\left| D^{\alpha\beta} b_i^j(z) \right| \leq K_0 F^{(\alpha+\beta)/2}(p, \delta) F_i^{1/2}(p, \delta) F_j^{-1/2}(p, \delta).$$

(2) *If $\mathcal{L} \in \mathcal{L}_M(\mathcal{B}(p, \delta)) \cup \{N\}$, $|\mathcal{L}| = S$, and D^T is a derivative in the coordinate system (z) with $|T| \leq M$, then $\mathcal{L} = \sum_{|s| \leq S} c_s D^s$, $D^T = \sum_{|\mathcal{L}'| \leq |T|} d_{\mathcal{L}'} \mathcal{L}'$, and, for $z \in \Delta_{c_0}(p, \delta)$ and $q = \Phi_p(z)$ we have*

$$|c_s(z)| \leq K_0 F^{(\mathcal{L}-s)/2}(p, \delta),$$

$$|d_{\mathcal{L}'}(q)| \leq K_0 F^{(\mathcal{L}-\mathcal{L}')/2}(p, \delta).$$

(3) *For $L = \sum a_i L_i$, $a_i \in \mathbb{C}$, for all $q \in B^{c_0}(p, \delta)$, $\frac{1}{2}F(L, p, \delta) \leq F(L, q, \delta) \leq 2F(L, p, \delta)$.*

(4) *For all list \mathcal{L} , $|\mathcal{L}| \leq M$ belonging to $\mathcal{L}_M(\mathcal{B})$ and all point $q \in B^c(p, \delta)$,*

(a) $|\mathcal{L}(\partial\rho)(q)| \leq K_0 \delta F(p, \delta)^{|\mathcal{L}|/2}$,

(b) *with the notation introduced in EB₂ in Definition 3.1,*

$$\left| \mathcal{L} a_{ij}^k(q) \right| \leq K_0 F^{|\mathcal{L}|/2}(p, \delta) F_i^{1/2}(p, \delta) F_j^{1/2}(p, \delta) F_k^{-1/2}(p, \delta).$$

(5) $\rho(B^c(p, \delta)) \subset [-\frac{1}{2}\delta, \frac{1}{2}\delta]$.

The proofs are almost straightforward computations.

In Section 4 we will need to use two other kinds of ‘‘pseudo-balls’’ and we will prove that they are closely related to the ‘‘polydisc’’ B^c :

Definition 3.4. Suppose that $\mathcal{B} = (L_1, \dots, L_{n-1})$ is a basis satisfying conditions (A) and (B) (at a point of finite 1-type).

(1) Denote $\mathcal{Y}_i = \Re L_i$ and $\mathcal{Y}_{i+n} = \Im L_i$, $1 \leq i \leq n$ (recall $L_n = N$). Then we denote by $B_{\mathcal{E}}^c(\mathcal{B}, p, \delta)$ the set of points $q \in V(p_0)$ for which there exists a piecewise \mathcal{C}^1 curve $\varphi : [0, 1] \rightarrow V$ such that $\varphi(0) = p$, $\varphi(1) = q$ and $\varphi'(t) = \sum a_i \mathcal{Y}_i(\varphi(t))$, with $\max(|a_i|, |a_{i+n}|) \leq cF^{-1/2}(L_i, p, \delta)$, $0 < c < 1$.

(2) \exp_p denoting the exponential map based at p associated to the vector fields \mathcal{Y}_i (defined in (1)), for $0 < c < 1$, we put

$$B_{\exp}^c(p, \delta) = \left\{ q = \exp_p(u_1, \dots, u_{2n}), \text{ such that } \max(|u_i|, |u_{i+n}|) \leq cF_i(p, \delta)^{-1/2} \right\} \cap V(p_0).$$

The terminology used in Definition 3.1 is justified by the following property:

Proposition 3.7. *Let $\mathcal{B} = \{L_1, \dots, L_{n-1}\}$ be a basis (of $(1, 0)$ complex vector fields, tangent to ρ in $V(p_0)$) satisfying conditions (A) and (B) (for example if it is K -extremal) at $p \in W(p_0) \cap \partial\Omega$. Let $\mathcal{B}^1 = \{L_1^1, \dots, L_{n-1}^1\}$ be another basis in $V(p_0)$ such that, for all i , $L_i^1 = \sum a_i^j L_j$, $a_i^j \in \mathbb{C}$, $\sum |a_i^j|^2 = 1$. Then there exists a constant A depending only on K , τ and the dimension n such that $B_{\mathcal{E}}^c(\mathcal{B}^1, p, \delta) \subset B_{\mathcal{E}}^c(\mathcal{B}, p, \delta)$.*

The proof of this Proposition immediately follows from property (B).

3.4. Sufficient conditions of extremality. In this Section we always assume that p_0 is a point of finite 1-type τ and choose $M = M(\tau)$.

Here and in Section 5.2 we will need a stronger control on certain derivatives of the coefficients of the Levi form. Thus we introduce the following condition: suppose \mathcal{B} is a basis of $(1, 0)$ vector fields tangent to ρ in $V(p_0)$. We say that it satisfies condition $B(\alpha)$, $\alpha > 0$, if for all lists $\mathcal{L} \in \mathcal{L}_{M-2}(\mathcal{B})$ we have

$$B(\alpha) \quad \text{for } i \neq j, 1 \leq i, j \leq n-1, |\mathcal{L}c_{ij}(p)| \leq \alpha \delta F(p, \delta)^{\mathcal{L}/2} F(L_i, p, \delta)^{1/2} F(L_j, p, \delta)^{1/2}.$$

Note that $B(\alpha)$ together with conditions (A) and (B) implies a new condition on the brackets of the vector fields:

Lemma 3.2. *Suppose \mathcal{B} satisfies conditions (A) and (B). Then there exist two constants $K_1 = K_1(K, M, n)$ and δ_0 depending on K , α and the data such that, for all $i \neq k$, $i, k < n$, $j \leq n$ and all $\mathcal{L} \in \mathcal{L}_M(\mathcal{B} \cup \{N\})$, if \mathcal{B} satisfies $B(\alpha)$ at (p, δ) , $p \in W(p_0)$, $0 < \delta \leq \delta_0$, then*

$$\left| \mathcal{L} a_{ik}^{\bar{j}}(p) \right| \leq K_1 \alpha F(p, \delta)^{\mathcal{L}/2} F_i(p, \delta)^{1/2} F_k(p, \delta)^{1/2} F_j(p, \delta)^{-1/2}.$$

Proof. To simplify the notations we write the proof for $a_{ik}^{\bar{j}}$. Choose δ_0 so that $C\delta_0^{-2/M} > \alpha^{-1}$, where C is the constant of Proposition 3.5. Note that the property is trivial if $l_n \neq 0$ or if $l_n = 0$ and $j = n$ ($a_{ik}^{\bar{n}} = \frac{1}{2}c_{ik}$ and $a_{ik}^{\bar{n}} = 0$), thus we suppose $l_n = 0$ and $j < n$. As the property is also trivial if j or k is $\geq i$, we have to study only the case when $j < \min(i, k)$.

To simplify the notations, we introduce the following spaces of functions:

$$*_0 = \{\varepsilon, \varepsilon a_{ij}^{\bar{k}}, \varepsilon c_{ij}^{\bar{k}}, \text{ where } \varepsilon \in \{-1, 0, 1, -\sqrt{-1}, \sqrt{-1}\}\},$$

and

$$\tilde{*}_{k+1} = \bigcup_i \bar{L}_i(*_k) \cup *_k \text{ and } *_k = \left\{ \sum_{i=1}^3 f_i, f_i \in \tilde{*}_{k+1} \right\}.$$

The elements of $*_k$ will be generically denoted by $*_k$.

The Jacobi identity applied to the bracket $[L_j, [L_i, \bar{L}_k]]$ implies

$$a_{ik}^{\bar{j}}c_{jj} + L_jc_{ik} + \sum_{p \neq j} a_{jk}^{\bar{p}}c_{jp} - a_{jk}^{\bar{i}}c_{ii} - L_ic_{jk} - \sum_{p \neq i} a_{jk}^{\bar{p}}c_{ip} - a_{ij}^{\bar{k}}c_{kk} - \sum_{p \neq k} a_{ji}^{\bar{p}}c_{pk} = 0$$

which we write $a_{ik}^{\bar{j}}c_{jj} = *_0c_{ii} + *_0c_{kk} + h$. Then, by induction on the length l of a list $\mathcal{L} \in \mathcal{L}_M(L_j)$, it is easy to show that

$$a_{ik}^{\bar{j}}\mathcal{L}c_{jj} = \mathcal{L}h + \sum_{\mathcal{L}' \in \mathcal{L}_{|\mathcal{L}|-1}(L_j)} (*_{l'}\mathcal{L}'c_{ii} + *_l\mathcal{L}'c_{kk}) + \sum_{\mathcal{L}' \in \mathcal{L}_{|\mathcal{L}|-1}(L_j)} *_l\mathcal{L}'c_{jj},$$

and choosing \mathcal{L} so that $|\mathcal{L}c_{jj}(p)| \gtrsim \delta F(p, \delta)^{(|\mathcal{L}|+2)/2}$, the Lemma is easily proved using the control on the lists and the hypothesis. \square

Now we first prove that conditions $B(\alpha)$, (A) and (B) imply the extremality of the basis and then that Lemma 3.2 implies a better control on mixed lists. This result will be important in Section 5.

Lemma 3.3. *Suppose that $\mathcal{B} = (L_1, \dots, L_{n-1})$ is a basis of $(1, 0)$ vector fields in $V(p_0)$ satisfying conditions (A) and (B) at a point $p \in V(p_0) \cap \partial\Omega$ for a fixed δ .*

Then there exists a function $\alpha(K)$, depending on K and the data, such that, if \mathcal{B} satisfies $B(\alpha)$ for $\alpha \leq \alpha(K)$, there exists a constant K_1 , depending on K , M and n , such that:

If $\mathcal{L}^0 \in \mathcal{L}_M(\mathcal{B})$ satisfies $|\mathcal{L}^0c_{ii}(p)| \geq \frac{1}{K}\delta F_i(p, \delta)F(p, \delta)^{\mathcal{L}^0/2}$ then there exists k_0 , $2k_0 + 2 \leq |\mathcal{L}^0|$, such that $\Re\left((L_i\bar{L}_i)^{k_0}c_{ii}\right)(p) > \frac{1}{K_1}\delta F_i(p, \delta)^{(2k_0+2)/2}$. In particular,

$$F_i(p, \delta) \geq \frac{1}{K'} \sum_{\substack{\Re\left((L_i\bar{L}_i)^k c_{ii}\right)(p) > 0 \\ 2k+1 \leq M}} \left[\frac{\Re\left((L_i\bar{L}_i)^k c_{ii}\right)(p)}{\delta} \right]^{\frac{2}{2k+2}},$$

where K' is a constant depending only on K and the data.

Proof. First we fix the notations used in the proof. We know that there is a coordinate system Φ_p^δ adapted to \mathcal{B} . We denote by (z_i) these coordinates. Let $D^{\alpha\beta}$ denote the derivative $\frac{\partial^{|\alpha+\beta|}}{\partial z^\alpha \partial \bar{z}^\beta}$ with respect to (z_i) , and if \mathcal{L} is a list of vector fields let $D^\mathcal{L}$ be the derivative $D^{\alpha\beta}$ with $\alpha_i = l_i^1(\mathcal{L})$ and $\beta_i = l_i^2(\mathcal{L})$ (notation of Lemma 3.2.2).

In the proof we will use a general result on derivatives of positive functions proved in Section 8.

Suppose $\mathcal{L} \in \mathcal{L}_M(\mathcal{B})$ is such that $\mathcal{L}(\partial\rho) = \mathcal{L}^0 c_{ii}$ and $|\mathcal{L}(\partial\rho)(p)| \gtrsim_K \delta F_i(p, \delta) F(p, \delta)^{\mathcal{L}^0/2}$. Then we can write

$$\mathcal{L}(\partial\rho) = D^{\mathcal{L}^0} c_{ii} + \sum_{|\alpha+\beta| < |\mathcal{L}^0|} c_{\alpha\beta} D^{\alpha\beta} c_{ii}$$

with $|c_{\alpha\beta}| \lesssim_K F^{\mathcal{L}^0/2} F^{-(\alpha+\beta)/2}$.

Thus there exists a derivative $D^{\alpha\beta}$ satisfying $|D^{\alpha\beta} c_{ii}(0)| \gtrsim_K \delta F_i F^{(\alpha+\beta)/2}$ and $|\alpha+\beta| \leq |\mathcal{L}^0|$ and $\alpha_n + \beta_n = 0$ (indeed, if $\alpha_n + \beta_n \geq 1$, $|c_{\alpha\beta}(0)| \lesssim_K F^{\mathcal{L}^0/2} F^{-(\alpha+\beta)/2} \leq \delta F^{\mathcal{L}^0/2}$, and, as $|D^{\alpha\beta} c_{ii}| \lesssim_K 1$, $|c_{\alpha\beta} D^{\alpha\beta} c_{ii}| \ll \delta F^{\mathcal{L}^0/2}$). Then applying Lemma 8.1 to the function $g(z) = \delta F_i^{-1}(p, \delta) c_{ii} \circ \Phi_{p, \delta}^{-1}(z')$, where $z' = (c F_1^{-1/2} z_1, \dots, c F_{n-1}^{-1/2}(p, \delta) z_{n-1}, 0)$ with $c \leq c_0$, c_0 given by Proposition 3.6, we conclude that there exists a derivative $D^{\alpha^1 \beta^1}$, satisfying $\alpha_j^1 = \beta_j^1, \forall j, \alpha_n^1 = \beta_n^1 = 0$, such that $D^{\alpha^1 \beta^1} c_{ii}(0) \geq_K F_i F^{(\alpha^1 + \beta^1)/2}$.

Writing $\mathcal{L}' = (\overline{L_i L_i})^{\alpha_i^1} \prod_{j \neq i, j < n} (\overline{L_j L_j})^{\alpha_j^1}$ and $\mathcal{L}' c_{ii} = D^{\alpha^1 \beta^1} c_{ii} + \sum_{|\alpha+\beta| < |\mathcal{L}'|} c_{\alpha\beta} D^{\alpha\beta} c_{ii}$, by induction we conclude that there exists a differential operator \mathcal{L}^1 of the form $\mathcal{L}^1 = (\overline{L_i L_i})^{\alpha_i^1} \prod_{j \neq i, j < n} (\overline{L_j L_j})^{\alpha_j^1}$ such that $\Re(\mathcal{L}^1 c_{ii})(p) \gtrsim_K \delta F^{\mathcal{L}^1/2} F_i$. Suppose there exists $j \neq i$ such that $\alpha_j \neq 0$. Then

$$\mathcal{L}^1 c_{ii} = \mathcal{L}' \overline{L_j} L_j c_{ii} = \mathcal{L}' \overline{L_j} \left(-\gamma_i^j c_{jj} + L_k c_{jk} + \left(a_{jk}^i - a_{ij}^{\bar{i}} \right) c_{ii} - \sum_{p \neq i} \left(a_{ij}^{\bar{p}} c_{ip} - a_{ji}^p c_{pi} \right) + \sum_{p \neq j} \gamma_i^{\bar{p}} c_{ip} \right).$$

The controls of the coefficients a_{ij}^p and of the lists $\mathcal{L} c_{kp}, k \neq p$ (by condition (B)), imply, for α sufficiently small (depending only on K), that

$$|\mathcal{L}' \overline{L_j} c_{jj}| \gtrsim_K \delta F^{\mathcal{L}'/2} F_j^{3/2} \text{ and } |\gamma_i^j| \gtrsim_K F_i F_j^{-1/2}.$$

Repeating the initial procedure, we conclude that there exists a list $\mathcal{L}'' \in \mathcal{L}(\mathcal{B})$, “completely even”, $|\mathcal{L}''| \leq |\mathcal{L}'|$ such that $|\mathcal{L}'' c_{jj}| \gtrsim_K \delta F^{\mathcal{L}''/2} F_j$. Consider then

$$\mathcal{L}'' \overline{L_j} c_{ii} = \mathcal{L}'' \left(-\gamma_i^j c_{jj} + L_k c_{jk} + \left(a_{jk}^i - a_{ij}^{\bar{i}} \right) c_{ii} - \sum_{p \neq i} \left(a_{ij}^{\bar{p}} c_{ip} - a_{ji}^p c_{pi} \right) + \sum_{p \neq j} \gamma_i^{\bar{p}} c_{ip} \right).$$

Thus $|\mathcal{L}'' c_{jj} \gamma_i^j| \gtrsim \delta F^{\mathcal{L}''/2} F_j^{1/2} F_i$, and, by similar arguments, for α sufficiently small, we conclude that there exists a list \mathcal{L}^2 , $|\mathcal{L}^2| < |\mathcal{L}^0|$ such that $\mathcal{L}^2 c_{ii} \gtrsim_K \delta F^{\mathcal{L}^2/2} F_i$, and we can repeat the procedure. The Lemma is thus proved by induction. \square

Proposition 3.8. *There exist constants α_0 and K' depending on K and the data such that if the basis \mathcal{B} satisfies (A), (B) and $B(\alpha)$ for $\alpha \leq \alpha_0$ at (p, δ) , $p \in V(p_0)$, then \mathcal{B} is (K', p, δ) -extremal.*

Proof. We may suppose the basis ordered so that the weights $F_i = F(L_i, p, \delta)$ are ordered decreasingly. Let $L = \sum_{i=1}^{n-1} a_i L_i$, $a_i \in \mathbb{C}$, $\sum |a_i|^2 = 1$ so that $c_{LL} = \sum_{i=1}^{n-1} |a_i|^2 c_{ii}$. Denote $F(L) = F(L, p, \delta)$. By hypothesis (B) it is clear that $F(L) \lesssim_K \sum |a_i|^2 F_i$.

To show the converse inequality, we prove the following assertion:

Claim. For every constant $K > 0$, there exists a constant K_1 , depending on K and the data, such that:

if $i_0 \in \{1, \dots, n-1\}$ and $k_0 \in \{1, \dots, M\}$ are such that $|a_{i_0}|^2 F_{i_0}(p) \geq \frac{\sum |a_i|^2 F_i(p)}{K}$ and $\Re(L_{i_0} \overline{L_{i_0}})^{k_0} c_{i_0 i_0}(p) > \delta \frac{F_{i_0}^{k_0+1}(p)}{K}$, then:

- either $\Re(L \overline{L})^{k_0} c_{L \overline{L}} > \delta \frac{(\sum |a_i|^2 F_i(p))^{k_0+1}}{K_1}$,
- or there exist i_1 and $k_1 < k_0$ such that $|a_{i_1}|^2 F_{i_1}(p) \geq \frac{\sum |a_i|^2 F_i(p)}{K_1}$ and $\Re(L_{i_1} \overline{L_{i_1}})^{k_1} c_{i_1 i_1}(p) > \delta \frac{F_{i_1}^{k_1+1}(p)}{K_1}$.

Proof of the Claim. We have

$$(3.9) \quad (L \overline{L})^{k_0} c_{L \overline{L}} = \sum |a_i|^{2k_0+2} (L_i \overline{L_i})^{k_0} c_{ii} + \sum \alpha_{\mathcal{L}} \mathcal{L}(\partial\rho),$$

where the second sum contains lists of length $2k_0 + 2$ containing L_i or $\overline{L_i}$ for, at least, two different values of i . As

$$|a_{i_0}|^{2k_0+2} \Re(L_{i_0} \overline{L_{i_0}})^{k_0} c_{i_0 i_0}(p) > \delta \frac{(\sum |a_i|^2 F_i(p))^{k_0+1}}{K^{k_0+2}},$$

the conclusion is clear except in the two following cases:

- in the first sum of (3.9), there is a term whose real part is $< -A = -\delta \frac{(\sum |a_i|^2 F_i(p))^{k_0+1}}{C K^{k_0+2}}$;
- in the second sum of (3.9), there is a term which is, in modulus, bigger than A , with a constant C depending only on M and the coefficients a_i .

Suppose first that there exists an index $i \neq i_0$ such that $|a_i|^{2k_0+2} \Re(L_i \bar{L}_i)^{k_0} c_{ii}(p) < -A$. This implies first $|a_i|^2 F_i(p) \geq \frac{\sum |a_i|^2 F_i(p)}{K_1}$ and secondly $\Re(L_i \bar{L}_i)^{k_0} c_{ii}(p) < -\delta \frac{1}{K_1} F_i^{k_0+1}$. By Lemma 3.3 there exists $k_1 < k_0$ such that $\Re(L_i \bar{L}_i)^{k_1} c_{ii}(p) > \delta \frac{1}{K_1} F_i^{k_1+1}$. Thus the second assertion of the Claim is verified.

Suppose now that there is a term $\alpha_{\mathcal{L}} \mathcal{L}(\partial\rho)$ in the second sum of (3.9) satisfying $|\alpha_{\mathcal{L}} \mathcal{L}(\partial\rho)| > A$. Denote by l_i the number of times the vector fields L_i and \bar{L}_i appear in \mathcal{L} . If $l_k \neq 0$, hypothesis (B) implies immediately $|a_k|^2 F_k \gtrsim \sum |a_i|^2 F_i$ and $|\mathcal{L}(\partial\rho)| \gtrsim \delta \prod F_i^{l_i/2}$. \square

Corollary. *Suppose that $p_0 \in \partial\Omega$ is a point of finite type τ where the Levi form is locally diagonalizable. Then there exists a neighborhood $V(p_0)$ of p_0 and constants K and $\delta_0 > 0$ such that at every point p of $V(p_0) \cap \partial\Omega$ and for every $0 < \delta \leq \delta_0$, the basis diagonalizing the Levi form is (M, p, δ) -extremal (with $M = M'(\tau)$).*

Proof. Properties (A) and (B) were proved in [CD06b], and, by definition the basis diagonalizing the Levi form satisfies B(α) for all $\alpha > 0$. \square

Definition 3.5. \mathcal{B} is called (K, α, p, δ) -strongly-extremal if it is (K, p, δ) -extremal and, if, it satisfies B(α) at (p, δ) .

Note that the first part of Proposition 3.2 says that every (K, p, δ) -extremal basis is (K, α, p, δ) -strongly-extremal for some large positive number α depending on K and Ω . Thus this is an extra hypothesis only for small α .

The next Proposition shows that for a strongly extremal basis some derivatives of the diagonal terms of the Levi matrix satisfy a better control:

Proposition 3.9. *Suppose p_0 is of finite 1-type τ and let $M = M'(\tau)$. Then there exists a neighborhood $V(p_0)$ of p_0 with the following property:*

for $\alpha > 0$, there exist constants $\delta_0 = \delta_0(\alpha, \text{data})$ and $K' = K'(K, \text{data})$ such that:

if \mathcal{B} is a (K, α, p, δ) -strongly-extremal basis, ordered so that F_i are decreasing, then for all lists $\mathcal{L} \in \mathcal{L}_{2M}(\mathcal{B})$ such that there exists $j > i$ with $l_j \neq 0$ we have $|\mathcal{L}c_{ii}(p)| \leq K' \alpha F(p, \delta)^{|\mathcal{L}|/2} F_i(p, \delta)$.

Proof. Let $\mathcal{L} = \mathcal{L}' \overleftarrow{L}_j \overleftarrow{L}_p \mathcal{L}''$ with $j \leq i$ and write

$$\mathcal{L}c_{ii} = \mathcal{L}' \overleftarrow{L}_p \overleftarrow{L}_j \mathcal{L}'' c_{ii} + \sum \mathcal{L}' \left(a_{j\bar{p}}^k L_k + a_{j\bar{p}}^{\bar{k}} \bar{L}_k \right) \mathcal{L}'' c_{ii}.$$

Then successive application of Lemma 3.2 show that there exists a list $\tilde{\mathcal{L}} = \tilde{\mathcal{L}}' L_j$ such that, for all $k, \tilde{l}_k = l_k$ and $|\tilde{\mathcal{L}}c_{ii} - \mathcal{L}c_{ii}| \leq K_2 \alpha F^{\tilde{\mathcal{L}}/2} F_i$.

Now the result is trivial, applying once again Lemma 3.2, Lemma 3.2.1 and the hypothesis B(α). \square

Proposition 3.10. *If the basis \mathcal{B} is (K, α, p, δ) -strongly extremal, the conclusion of Proposition 3.9 is still valid at each point $q \in B^{c_0}(p, \delta)$ with α replaced by 2α for $\delta \leq \delta(\alpha)$ ($\delta(\alpha)$ depending on α, K and the data).*

3.5. Localization of extremal bases.

3.5.1. Definition of the local domain.

Definition 3.6. Let Ω be a bounded pseudo-convex domain in \mathbb{C}^n . Suppose that P_0 is a boundary point of Ω and $W(P_0) \Subset V(P_0)$ are neighborhoods of P_0 . Let O be a point of the real normal to $\partial\Omega$ at P_0 and denote by d the distance from O to P_0 . Let us denote by $(z_i)_{1 \leq i \leq n}$ the coordinate system obtained translating the origin at O .

Let $\mu > 0$ and $\psi(z) = \varphi(|z|^2)$ where

$$\varphi(x) = \begin{cases} 0 & \text{if } x \leq \mu^2, \\ K_0 e^{-1/(x-\mu^2)} & \text{if } x \geq \mu^2, \end{cases}$$

with $\frac{4}{3}d \leq \mu \leq 2d$.

Let us denote $r(z) = \rho(z) + \psi(z)$. Then d is chosen small enough and K_0 large enough such that, in particular:

- $D = \{r(z) < 0\} \subset W(P_0)$ and r is a defining function of D ;
- D have a \mathcal{C}^∞ boundary and is pseudo-convex;
- At each point of $\partial\Omega \setminus \partial D$, the boundary of D is strictly pseudo-convex;
- In the closure of $B(0, 2\mu)$ the vector z (in the coordinate system centered at 0) is not tangent to ρ (i.e. $\sum_{i=1}^n \frac{\partial \rho}{\partial z_i} z_i \neq 0$ everywhere in the closure of $B(0, 2\mu)$).

The fact that such a domain always exists for any $d > 0$ small and $K_0 > 0$ large is based on the construction of R. Gay and A. Sebbar in [GS85] (Théorème 2.1). Simply, note that, on $\partial D \setminus \partial\Omega$, the function r is strictly pluri-subharmonic if K_0 is large enough and μ small enough (the hessian of ρ is $O(\varphi(|z|^2))$). Moreover, if P_0 is of finite type, then all the boundary points of D are of finite type because the order of contact of $\partial\Omega$ with ∂D is infinite at the points of $\partial(\partial\Omega \cap \partial D)$.

The goal of this Section is to prove the following:

Theorem 3.3. *Suppose that P_0 is a point of finite 1-type τ of $\partial\Omega$ and choose $M'(\tau)$ (c. f. Proposition 3.5). Let $\delta > 0$ and $K > 0$. If at every point of $\partial\Omega \cap V(P_0)$ there is a (K, p, δ) -extremal basis then one can construct the domain D contained in $V(P_0)$ so that, at every point p' of its boundary there exists a (K', p', δ) -extremal basis with K' depending only on K and the data.*

The proof of this theorem is done in the two following sections.

3.5.2. *Preliminary remarks.* We fix now some general notations.

Let π be the \mathcal{C}^∞ projection of $V(P_0) \cap \bar{\Omega}$ onto $\partial\Omega$ defined with the integral curves of the real normal to ρ . It is clear that there exists a neighborhood \mathcal{U} of $\partial\Omega \cap V(P_0)$ such that π is a \mathcal{C}^∞ diffeomorphism of $\partial D \cap \mathcal{U}$ onto an open set of $\partial\Omega \cap \mathcal{U}$.

If L is a \mathcal{C}^∞ vector field, defined on an open set U of $\partial D \cap \mathcal{U}$, tangent to ∂D , we associate to it a vector field L^ρ , defined in the open set $\pi(U) \subset \partial\Omega$, tangent to $\partial\Omega$ using π (considered as a \mathcal{C}^∞ diffeomorphism of U onto $\pi(U)$) as follows: if $L = \sum a_i \frac{\partial}{\partial z_i}$, considering it as an application of U into \mathbb{C}^n , we denote by $L \circ \pi^{-1}$ the vector field in $\pi(U)$ defined by $L \circ \pi^{-1} = \sum a_i \circ \pi^{-1} \frac{\partial}{\partial z_i}$, and

$$(3.10) \quad L^\rho = L \circ \pi^{-1} - \beta N,$$

where N is the complex unitary normal to ρ and $\beta = L \circ \pi^{-1}(\rho)$.

Clearly, $L \mapsto L^\rho$ is an isomorphism from $T_{\partial D \cap U}^{1,0}$ onto $T_{\partial\Omega \cap \pi(U)}^{1,0}$ ($V(P_0)$ and \mathcal{U} sufficiently small), and thus, we also consider L associated to L^ρ by $L = L^\rho \circ \pi + (\beta \circ \pi)N \circ \pi$. As L is tangent to ∂D and $(L^\rho \circ \pi)(\rho)$ is identically zero on $\partial\Omega$, we have

$$(3.11) \quad \beta \circ \pi(z) = \frac{-\langle L^\rho \circ \pi, z \rangle \varphi'(|z|^2)}{(N \circ \pi)(\rho) + \langle N \circ \pi, z \rangle \varphi'(|z|^2)} + k,$$

where k is a \mathcal{C}^∞ function whose derivatives of order less than M are $O(\varphi(|z|^2))$ with constants controlled by the \mathcal{C}^{2M} norm of L , and, if $L = \sum a_i \frac{\partial}{\partial z_i}$ (in the coordinate system of Definition 3.6), $\langle L, z \rangle$ denotes the usual scalar product $\sum a_i \bar{z}_i$, and $\langle L, L' \rangle = \sum a_i a'_i$.

With the previous notations, let P be a point of ∂D such that $\psi(P) = 0$ (thus $P \in \partial D \cap \partial\Omega$) and let $V(P) \subset \mathcal{U}$ be a neighborhood of P such that π is a diffeomorphism of $V(P) \cap \partial D$ onto $V(P) \cap \partial\Omega$.

Let $p \in \partial D \cap V(P)$. Essentially, the construction of the extremal basis \mathcal{B} at p for D is done using a suitable basis \mathcal{B}^ρ of the tangent space of $\partial\Omega$ near the point $\pi(p)$ translated at p (using π) then projected onto the tangent space of ∂D , to get a basis $\tilde{\mathcal{B}}$ which will be used (in the next section) to define the basis \mathcal{B} .

Currently, we only look at the relation between the weights of the basis $\tilde{\mathcal{B}}$ and \mathcal{B}^ρ .

Thus, if $\tilde{\mathcal{B}} = \{\tilde{L}_1, \dots, \tilde{L}_{n-1}\}$ is a basis of $T_{\partial D}^{1,0}$ in $V(P) \cap \partial D$, with our notations, the basis $\mathcal{B}^\rho = \{L_1^\rho, \dots, L_{n-1}^\rho\}$ of $T_{\partial\Omega}^{1,0}$ (in $V(P) \cap \partial\Omega$) is given by

$$(3.12) \quad L_i^\rho = \tilde{L}_i \circ \pi^{-1} - \beta_i N,$$

with $\beta_i = \tilde{L}_i \circ \pi^{-1}(\rho)$, and

$$(3.13) \quad \tilde{L}_i = L_i^\rho \circ \pi + (\beta_i \circ \pi)N \circ \pi.$$

with

$$(3.14) \quad \beta_i \circ \pi = \frac{-\langle L_i^\rho \circ \pi, z \rangle \varphi'(|z|^2)}{(N \circ \pi)(\rho) + \langle N \circ \pi, z \rangle \varphi'(|z|^2)} + k.$$

Let us calculate the weights $F(\tilde{L}_i, p, \delta)$ in terms of the weights $F(L_i^\rho, \pi(z), \delta)$ and the derivatives of φ . We suppose that L_i^ρ are normalized. Writing $\tilde{c}_{ij} = [\tilde{L}_i, \tilde{L}_j](\partial r)$ and $c_{ij}^\rho = [L_i^\rho, L_j^\rho](\partial \rho)$, using that $(N \circ \pi)(\rho)$ is identically 1 on $\partial\Omega$, a simple computations shows

$$(3.15) \quad \begin{aligned} \tilde{c}_{ij} &= c_{ij}^\rho \circ \pi + \langle L_i^\rho \circ \pi, L_j^\rho \circ \pi \rangle \varphi'(|z|^2) + \langle L_i^\rho \circ \pi, z \rangle \overline{\langle L_j^\rho \circ \pi, z \rangle} \varphi''(|z|^2) + \\ &+ \varphi'(|z|^2) \sum_{k=1}^{n-1} \left(* \langle L_k^\rho \circ \pi, z \rangle + * \overline{\langle L_k^\rho \circ \pi, z \rangle} \right) + k, \\ &= c_{ij}^\rho \circ \pi + \varphi'(|z|^2) \left(\langle L_i^\rho \circ \pi, L_j^\rho \circ \pi \rangle + h \right) + \langle L_i^\rho \circ \pi, z \rangle \overline{\langle L_j^\rho \circ \pi, z \rangle} \varphi''(|z|^2) + k, \end{aligned}$$

where all the derivatives of k are $O(\varphi(|z|^2))$ and the functions $*$ have a bounded \mathcal{C}^M norm, the constants depending only on Ω and the \mathcal{C}^{2M} norms of the \tilde{L}_i .

As L_i^p are normalized, we also have

$$(3.16) \quad \begin{aligned} \tilde{c}_{ii} &= c_{ii}^p \circ \pi + \varphi'(|z|^2) + |\langle L_i^p \circ \pi, z \rangle|^2 \varphi''(|z|^2) + \varphi'(|z|^2) \sum_{k=1}^{n-1} \left(* \langle L_k^p \circ \pi, z \rangle + * \langle \overline{L}_k^p \circ \pi, z \rangle \right) + k \\ &= c_{ii} \circ \pi + \varphi'(|z|^2)(1+h) + |\langle L_i^p \circ \pi, z \rangle|^2 \varphi''(|z|^2) + k \end{aligned}$$

and d is chosen small enough such that the \mathcal{C}^M norm of h is small.

Now we need to introduce a new notation. Let L be a $\mathcal{C}^\infty(\partial D \cap V(P))$ vector field tangent to ∂D . For $z \in \partial D \cap V(P)$ let us define

$$\widetilde{F^\varphi}(L, z, \delta) = \sum_{k=1}^{M/2} \left(\frac{\varphi^{(k)}(|z|^2)}{\delta} \right)^{1/k} + |\langle L^p \circ \pi(z), z \rangle|^2 \sum_2^M \left| \frac{\varphi^{(k)}(|z|^2)}{\delta} \right|^{2/k} + \delta^{-1/M}.$$

Lemma 3.4. *For δ and $V(P)$ small enough and for $z \in \partial D \cap V(P)$, we have:*

$$\widetilde{F^\varphi}(L, z, \delta) \simeq \frac{\varphi'(|z|^2)}{\delta} + |\langle L^p \circ \pi, z \rangle|^2 \frac{\varphi''(|z|^2)}{\delta} + \delta^{-1/M}.$$

Proof. It suffices to consider the case when $|z|^2 = \mu^2 + x > \mu^2$. Note that, for $V(P)$ small, $\varphi^{(k)}(\mu^2 + x) \simeq K e^{-1/x} x^{-2k}$ and $(\frac{1}{x})^{2k} \leq e^{1/Mx}$, for $k \leq M$.

Suppose $\left(\frac{\varphi^{(k)}(\mu^2 + x)}{\delta} \right)^{1/k} > \delta^{-1/M}$ and $e^{-1/x} < \delta$. Then

$$\left(\frac{\varphi^{(k)}(\mu^2 + x)}{\delta} \right)^{1/k} \simeq \left(\frac{K_0 e^{-1/x}}{\delta} \right)^{1/k} \frac{1}{x^2} \lesssim K_0^{1/2} \delta^{-1/kM} \leq \delta^{-1/M},$$

for δ small. Thus, for $\delta \leq \delta_0(K_0)$, $\left(\frac{\varphi^{(k)}(\mu^2 + x)}{\delta} \right)^{1/k} > \delta^{-1/M}$ implies $e^{-1/x} > \delta$ and $\sum_1^{M/2} \left(\frac{\varphi^{(k)}(\mu^2 + x)}{\delta} \right)^{1/k} \simeq \frac{\varphi'(\mu^2 + x)}{\delta}$.

Similarly, $\left(\frac{\varphi^{(k)}(\mu^2 + x)}{\delta} \right)^{2/k} > \delta^{-1/M}$ implies $e^{-1/x} > \delta$ and $\sum_2^M \left(\frac{\varphi^{(k)}(\mu^2 + x)}{\delta} \right)^{2/k} \simeq \frac{\varphi''(\mu^2 + x)}{\delta}$. \square

Thus, we denote

$$F^\varphi(L, z, \delta) = \frac{\varphi'(|z|^2)}{\delta} + |\langle L^p \circ \pi, z \rangle|^2 \frac{\varphi''(|z|^2)}{\delta} + \delta^{-1/M},$$

$F_i^\varphi = F_i^\varphi(z, \delta) = F^\varphi(\tilde{L}_i, z, \delta)$, $1 \leq i \leq n-1$ and $F_n^\varphi = \delta^{-2}$. Let \tilde{L}_n denotes the unitary complex normal to r (the defining function of D) and L_n^p the unitary complex normal to ρ .

Proposition 3.11. *Let $\tilde{\mathcal{L}}$ be a list of $\mathcal{L}_M(\tilde{\mathcal{B}} \cup \{\tilde{L}_n\})$ and \mathcal{L}^p be the list obtained replacing \tilde{L}_i in $\tilde{\mathcal{L}}$ by \overline{L}_i^p . Then, reducing $V(P)$ if necessary, on $\partial D \cap V(P)$ we have (\tilde{l}_i denoting the number of times the vector fields \tilde{L}_i or \overline{L}_i^p appears in $\tilde{\mathcal{L}}$):*

- (1) $\left| \tilde{\mathcal{L}}(c_{ij}^p \circ \pi) - (\mathcal{L}^p c_{ij}^p) \circ \pi \right| \lesssim \delta \prod_{k=1}^n (F_k^\varphi)^{\tilde{l}_k/2}$, for $|\tilde{\mathcal{L}}| \geq 2$,
- (2) $\left| \tilde{\mathcal{L}}\varphi(|z|^2) \right| \lesssim \delta \prod_{i=1}^n (F_i^\varphi)^{\tilde{l}_i/2}$, $|\tilde{\mathcal{L}}| \geq 2$,

the constants depending only on Ω and the \mathcal{C}^{M+2} norms of the \tilde{L}_i .

Proof. These properties are trivially satisfied if $\tilde{l}_n \neq 0$, thus we suppose $\tilde{l}_n = 0$. Using (3.15) and the fact that if f is a \mathcal{C}^∞ function on $\partial \Omega \cap V(P)$ and if $L^p \rho \equiv 0$ then $(L^p \circ \pi)(f \circ \pi) - (L^p f) \circ \pi = O_f(\varphi)$ on $\partial D \cap V(P)$, the Proposition is an easy consequence of (3.14) and the following Lemma:

Lemma 3.5. *Let $\mathcal{L}^{\rho, \pi}$ be a list of $\mathcal{L}_M\{L_i^p \circ \pi, i \leq n-1\}$ of length ≥ 1 . Then $|\mathcal{L}^{\rho, \pi} \psi| \lesssim \delta \prod_{i=1}^{n-1} (F_i^\varphi)^{\tilde{l}_i/2}$.*

Proof of Lemma 3.5. By induction, we have

$$\mathcal{L}^{\rho, \pi} \psi = \mathcal{L}^{\rho, \pi} \left(\varphi(|z|^2) \right) = \sum_{l=1}^{\lfloor \frac{m-1}{2} \rfloor} * \varphi^{(l)}(|z|^2) + \sum_{l=\lfloor \frac{m+1}{2} \rfloor}^m \alpha_l \varphi^{(l)}(|z|^2),$$

with

$$\alpha_{m-k} = \sum_{\substack{\mathcal{L}^* = \{W_1^*, \dots, W_m^*\} \subset \mathcal{L}^{\rho, \pi} \\ m^* \leq m-2k}} * \prod_{W_i^* \in \mathcal{L}^*} \langle W_i^*, \tilde{z} \rangle,$$

where $\langle W_i^*, \tilde{z} \rangle$ denotes $\langle W_i^*, z \rangle$ if W_i^* is of type $(0, 1)$ and $\langle W_i^*, \bar{z} \rangle$ if not, and the \mathcal{C}^{2M} norms of the functions $*$ are controlled by the \mathcal{C}^{2M} norms of the vector fields \tilde{L}_i . Now, the proof of Lemma 3.4 shows that

$$\sum_{l=1}^{\lfloor \frac{m-1}{2} \rfloor} \frac{* \varphi^{(l)}(|z|^2)}{\delta} \lesssim \left(\delta^{-1/M} + \frac{\varphi'(|z|^2)}{\delta} \right)^{m/2},$$

and it is enough to see that $|\alpha_l \varphi^{(l)}(|z|^2)| \lesssim \delta (F^\varphi)^{\mathcal{L}/2}$, for $l \in \{[\frac{m+1}{2}], \dots, m\}$. If $l = m$, this follows from Lemma 3.4; suppose $l = m - k$, $k \geq 1$.

Suppose $\left| \frac{\varphi^{(m-k)}(|z|^2)}{\delta} \right|^{2/(m-k)} \geq \delta^{-1/M}$. By Lemma 3.4 $\left| \frac{\varphi^{(m-k)}(|z|^2)}{\delta} \right|^{2/(m-k)} \leq \frac{\varphi''(|z|^2)}{\delta}$. Let $\mathcal{L}^* \subset \mathcal{L}$ of length $m^* = m - 2k = \sum_{i=1}^{n-1} l_i^*$. The corresponding term in α_{m-k} is bounded by

$$\begin{aligned} * \left(\frac{\varphi''(|z|^2)}{\delta} \right)^{(m-k)/2} \prod_i |\langle L_i^p \circ \pi, z \rangle|^{l_i^*} &= * \left(\frac{\varphi''(|z|^2)}{\delta} \right)^{k/2} \prod_{i=1}^{n-1} \left(\frac{\varphi''(|z|^2)}{\delta} |\langle L_i^p \circ \pi, z \rangle|^2 \right)^{l_i^*/2} \\ &\lesssim * \left(\frac{\varphi'(|z|^2)}{\delta} \right)^k \prod_{i=1}^{n-1} (F_i^\varphi)^{l_i^*/2}, \end{aligned}$$

because the hypothesis implies $\left(\frac{\varphi''(|z|^2)}{\delta} \right)^{1/2} \lesssim \frac{\varphi'(|z|^2)}{\delta}$. \square

To finish the proof of Proposition 3.11 note that, for $|\tilde{\mathcal{L}}| \geq 1$,

$$|\tilde{\mathcal{L}}(\beta_i \circ \pi)(z)| \leq F^\varphi(z, \delta)^{|\tilde{\mathcal{L}}|/2} F_i^\varphi(z, \delta)^{1/2},$$

and use (3.14). \square

Finally the relations between the weights associated to $\tilde{\mathcal{B}}$ and to \mathcal{B}^p are as follows.

Let \tilde{L} a holomorphic vector field on ∂D tangent to ∂D near p and L^p the associated vector field tangent to $\partial \Omega$. Then

Proposition 3.12. *For $V(P)$ sufficiently small, we have, if $\frac{1}{K} \leq \|\tilde{L}\| \leq K$,*

$$F(\tilde{L}, z, \delta) \simeq F(L^p, \pi(z), \delta) + F^\varphi(\tilde{L}, z, \delta),$$

with constants depending on the \mathcal{C}^{2M} norm of \tilde{L} , K and the data.

Proof. From Proposition 3.11 it easily follows that $F(\tilde{L}, z, \delta) \lesssim F(L^p \circ \pi, z, \delta) + F^\varphi(\tilde{L}, z, \delta)$. Let us then see that there exists a list $\tilde{\mathcal{L}}$ composed of \tilde{L} and \bar{L} such that $\tilde{\mathcal{L}} \tilde{c}_{\tilde{L}\tilde{L}} \simeq \delta \left(F(L^p \circ \pi, z, \delta) + F^\varphi(\tilde{L}, z, \delta) \right)^{(|\tilde{\mathcal{L}}|+2)/2} \stackrel{\text{def}}{=} \delta F^{(|\tilde{\mathcal{L}}|+2)/2}$. If $\frac{\varphi'}{\delta} + |\langle L^p \circ \pi(z), z \rangle|^2 \frac{\varphi''}{\delta} \simeq F$, then $c_{\tilde{L}\tilde{L}}$ do it. Suppose $\frac{\varphi'}{\delta} + |\langle L^p \circ \pi(z), z \rangle|^2 \frac{\varphi''}{\delta} \ll F$. Then, there exists a list \mathcal{L}^p such that $|\mathcal{L}^p c_{L^p L^p}(\pi(z))| \simeq \delta F^{(|\mathcal{L}^p|+2)/2}$. Then calculating $\tilde{\mathcal{L}} \tilde{c}_{\tilde{L}\tilde{L}}$ in term of $\mathcal{L}^p (c_{L^p L^p}) \circ \pi$, the result follows Proposition 3.11, (3.16) and the properties of the functions h and k . \square

3.5.3. Extremal bases on D . In this Section, we assume that p_0 is of finite type τ , $M = M'(\tau)$ and that, at all points q of $V(P_0) \cap \partial \Omega$ and for all $\delta > 0$, $0 < \delta \leq \delta_0$, there exists a (K, q, δ) -extremal basis. Then we will show that at all points p of ∂D and for all $\delta > 0$ there exists a (K', p, δ) -extremal basis (for D) with a constant K' controlled by K and the data.

If P is a point of ∂D such that $|P| > \mu$ then ∂D is strictly pseudo-convex near P and the construction of extremal basis in $V(P) \cap \partial D$ is trivial (for $V(P)$ small). If $|P| < \tau$ then $V(P) \cap \partial D$ is contained in $\partial \Omega$ and the existence of extremal basis is the hypothesis. Thus, we have only to consider neighborhood of points $P \in \partial D$ such that $|P| = \mu$ (that is points P in the boundary of $\partial \Omega \cap \partial D$).

As we said before, the final extremal basis for D , at $p \in V(P) \cap \partial D$, will be obtained extending a basis $\tilde{\mathcal{B}}$ defined on $V(P) \cap \partial D$ which is a projection onto the tangent space to r of a translation of a basis \mathcal{B}^p , at $\pi(p)$, tangent to p .

Formula (3.16) shows that the expressions $\langle L_i^p \circ \pi, z \rangle$ plays an important role: we have to take into account the vector fields which are orthogonal to z . In particular, to construct an extremal basis on ∂D , we cannot simply translate an extremal basis on $\partial \Omega$ and project it onto the tangent space to ∂D , because, even if the basis (L_i^p) is extremal, we may have $\langle L_i^p \circ \pi, z \rangle \neq 0$, for all i , and there are linear combinations of the $L_i^p \circ \pi$ which are orthogonal to z .

From now the point p and the positive number δ are fixed. We suppose we have a $(K, \pi(p), \delta)$ -extremal (for ρ) basis $\mathcal{B}^\Omega = \{L_1^\Omega, \dots, L_{n-1}^\Omega\}$ at the point $\pi(p)$ (the L_i^Ω being \mathcal{C}^∞ in $V(P)$), such that the vectors $L_i^\Omega(\pi(p))$ are orthogonal (c.f. Proposition 3.1) and we construct the basis $\mathcal{B}^p = \{L_1^p, \dots, L_{n-1}^p\}$ using it. The weights associated to \mathcal{B}^Ω are denoted $F_i^\Omega = F_i^\Omega(\pi(p), \delta) = F^\Omega(L_i^\Omega, \pi(p), \delta)$, and we suppose $F_{i+1}^\Omega \leq F_i^\Omega$, for $i \leq n-2$, changing the order of L_i^Ω if necessary.

Recall that the canonical coordinate system is centered at the point O of Definition 3.6, thus $|z(P)| = \mu$.

For simplicity of notations, we denote $q = \pi(p)$ (thus $p = \pi^{-1}(q)$, π being considered as a diffeomorphism between open sets of the boundaries of Ω and D).

Let

$$\mathcal{H}_{n-1} = \left\{ W = \sum a_i L_i^\Omega, a_i \in \mathbb{C}, \sum |a_i|^2 = 1, \text{ such that } \langle W(q), p \rangle = 0 \right\}.$$

Let $W_{n-1} = \sum a_i^{n-1} L_i^\Omega \in \mathcal{H}_{n-1}$ such that $\sum_{i=1}^{n-2} |a_i^{n-1}|^2 F(L_i^\Omega, q, \delta) = \inf_{W \in \sum a_i L_i^\Omega \in \mathcal{H}_{n-1}} \sum_{i=1}^{n-2} |a_i|^2 F(L_i^\Omega, q, \delta)$, and define

$$(3.17) \quad L_{n-1}^\rho = \begin{cases} L_{n-1}^\Omega & \text{if } \sum_{i=1}^{n-2} |a_i^{n-1}|^2 F(L_i^\Omega, q, \delta) \geq \frac{\varphi''(|p|^2)}{\delta} |\langle L_{n-1}^\Omega(q), p \rangle|^2, \\ W_{n-1} & \text{otherwise.} \end{cases}$$

Suppose L_{n-l}^ρ are defined for $1 \leq l \leq k-1 < n$. Let $\mathcal{H}_{n-k} = \mathcal{H}_{n-1} \cap [\mathcal{E}(L_{n-1}^\rho, \dots, L_{n-k+1}^\rho)]^\perp$, $\mathcal{E}(L_{n-1}^\rho, \dots, L_{n-k+1}^\rho)$ being the linear space spanned by $L_{n-1}^\rho, \dots, L_{n-k+1}^\rho$, the orthogonality being taken at q . Let $W_{n-k} = \sum_{i=1}^{n-1} a_i^{n-k} L_i^\Omega$ a vector in \mathcal{H}_{n-k} minimizing $\sum_{i=1}^{n-k-1} |a_i|^2 F(L_i^\Omega, q, \delta)$ for vectors $\sum_{i=1}^{n-1} a_i L_i^\Omega \in \mathcal{H}_{n-k}$. Let T_{n-k} be a vector field, of norm 1 at q , in $\mathcal{G}_{n-k} = \mathcal{E}(L_{n-1}^\Omega, \dots, L_{n-k}^\Omega) \cap [\mathcal{E}(L_{n-1}^\rho, \dots, L_{n-k+1}^\rho)]^\perp$. Then L_{n-k}^ρ is defined by

$$L_{n-k}^\rho = \begin{cases} T_{n-k} & \text{if } \sum_{i=1}^{n-k-1} |a_i^{n-k}|^2 F(L_i^\Omega, q, \delta) \geq \frac{\varphi''(|p|^2)}{\delta} |\langle T_{n-k}(q), p \rangle|^2, \\ W_{n-k} & \text{otherwise.} \end{cases}$$

Note that $\{L_i^\rho(q), 1 \leq i \leq n-1\}$ is orthonormal. We will note later that if the dimension of \mathcal{G}_{n-k} is strictly greater than 1 then $F^{\rho\varphi}(L_{n-k}^\rho)$ (see below) is, up to a multiplicative constant, independent of the choice of T_{n-k} .

The next two Lemmas prove some important properties of the vector fields L_i^ρ . Let us denote $\mathcal{B}^\rho = \{L_i^\rho, i < n\}$ and L_n^ρ the unitary complex normal to ρ .

For $L = \sum_{i=1}^{n-1} a_i L_i^\rho$, $a_i \in \mathbb{C}$, let us denote

$$F^{\rho\varphi}(L) = F(L, q, \delta) + \frac{\varphi'(|p|^2)}{\delta} + |\langle L(q), p \rangle|^2 \frac{\varphi''(|p|^2)}{\delta},$$

$F_i^{\rho\varphi} = F^{\rho\varphi}(L_i^\rho)$, $1 \leq i \leq n-1$, $F_n^{\rho\varphi} = \frac{1}{\delta^2}$ and $(F^{\rho\varphi})^{\mathcal{L}/2} = \prod_i (F_i^{\rho\varphi})^{l_i/2}$, if \mathcal{L} is a list of $\mathcal{L}_M(\mathcal{B}^\rho \cup \{L_n^\rho\})$, with the usual notation for l_i .

We will show that, up to constants, the vector fields L_i^ρ give the successive minima of the functions $F^{\rho\varphi}(L)$ for $L = \sum a_i L_i^\Omega$, $\sum |a_i|^2 = 1$.

Lemma 3.6. *There exists a constant K' depending only on K such that:*

- (1) If $L = \sum a_i L_i^\Omega$, $\sum |a_i|^2 = 1$, is orthogonal, at q , to the space generated by L_j^ρ , $i+1 \leq j \leq n-1$, $i \leq n-1$, then $F^{\rho\varphi}(L) \geq \frac{1}{K'} F^{\rho\varphi}(L_i^\rho)$;
- (2) $F^{\rho\varphi}(L_i^\rho) \geq \frac{1}{K'} F^{\rho\varphi}(L_{i+1}^\rho)$, $i < n-1$;
- (3) $F^{\rho\varphi}(L_i^\rho) \geq \frac{1}{K'} F(L_i^\Omega, q, \delta)$, $i < n$.

Proof. Note first that if property (2) is satisfied for $i \geq k$ then property (3) is also satisfied for $i \geq k$. Indeed, more generally, if L is orthogonal to the vectors L_j^ρ , $i+1 \leq j \leq n-1$, and if property (2) is satisfied for $i+1, \dots, n-1$, then

$$(3.18) \quad F^{\rho\varphi}(L) \gtrsim \max \left\{ F^\Omega(L), F^\Omega(L_{i+1}^\rho), \dots, F^\Omega(L_{n-1}^\rho) \right\} \gtrsim F^\Omega(L_i^\Omega, q, \delta) = F(L_i^\Omega, q, \delta) = F_i^\Omega,$$

because the L_j^ρ and L are orthogonal and the basis $(L_i^\Omega)_i$ is extremal.

Now we show that if $L = \sum a_i L_i^\Omega$, $\sum |a_i|^2 = 1$, then $F^{\rho\varphi}(L) \gtrsim F_{n-1}^{\rho\varphi}$.

If $L_{n-1}^\Omega \in \mathcal{H}_{n-1}$, then $L_{n-1}^\rho = L_{n-1}^\Omega$ and $F_{n-1}^{\rho\varphi} = F(L_{n-1}^\Omega, q, \delta) + \frac{\varphi'(|p|^2)}{\delta}$ which gives the result. Suppose thus $L_{n-1}^\Omega \notin \mathcal{H}_{n-1}$. We separate the two cases of (3.17):

Suppose we are in the first case ($L_{n-1}^\rho = L_{n-1}^\Omega$). If $L \in \mathcal{H}_{n-1}$, then the inequality is an immediate consequence of the extremality (EB₁) of \mathcal{B}^Ω . Suppose $L \notin \mathcal{H}_{n-1}$. Then we can write $L = \alpha (L_{n-1}^\rho + \gamma H)$ with $H \in \mathcal{H}_{n-1}$. Writing $H = \sum a'_i L_i^\Omega$, we have

$$F^{\rho\varphi}(L) \simeq |\alpha|^2 \left[\sum_{i=1}^{n-2} |\gamma a'_i|^2 F(L_i^\Omega, q, \delta) + |1 + \gamma a'_{n-1}|^2 F(L_{n-1}^\Omega, q, \delta) \right] + \frac{\varphi'(|p|^2)}{\delta} + |\alpha|^2 \frac{\varphi''(|p|^2)}{\delta} |\langle L_{n-1}^\Omega(q), p \rangle|^2,$$

and as $\sum_{i=1}^{n-2} |a'_i|^2 F(L_i^\Omega, q, \delta) \geq \sum_{i=1}^{n-2} |a_i^{n-1}|^2 F(L_i^\Omega, q, \delta) \geq \frac{\varphi''(|p|^2)}{\delta} |\langle L_{n-1}^\Omega(q), p \rangle|^2$, we obtain

$$F^{\rho\varphi}(L) \gtrsim |\alpha|^2 \left(1 + |\gamma|^2 \right) \frac{\varphi''(|p|^2)}{\delta} |\langle L_{n-1}^\Omega(q), p \rangle|^2 \gtrsim_K \frac{\varphi''(|p|^2)}{\delta} |\langle L_{n-1}^\Omega(q), p \rangle|^2,$$

because, by equivalence of norms in finite dimensional spaces, $|\alpha|^2 (1 + |\gamma|^2) \gtrsim_K 1$. The extremality of \mathcal{B}^Ω implies $F(L, q, \delta) \gtrsim F(L_{n-1}^\Omega, q, \delta)$, and the inequality is proved.

Let us now look to the case $L_{n-1}^\rho = W_{n-1}$. The result is trivial if $L \in \mathcal{H}_{n-1}$, thus we suppose $L \notin \mathcal{H}_{n-1}$. Using the same decomposition as before, we get

$$F^{\rho\varphi}(L) \gtrsim |\alpha|^2 \left(1 + |\gamma|^2\right) \sum_{i=1}^{n-2} |a_i^{n-1}|^2 F(L_i^\Omega, q, \delta) \gtrsim \sum_{i=1}^{n-2} |a_i^{n-1}|^2 F(L_i^\Omega, q, \delta),$$

and, as $F^{\rho\varphi}(L) \gtrsim F(L, q, \delta) \gtrsim F(L_{n-1}^\Omega, q, \delta)$, we have $F^{\rho\varphi}(L) \gtrsim F(W_{n-1}, q, \delta)$.

The induction is as follows. Suppose $F^{\rho\varphi}(L_{i+1}^\rho) \gtrsim F^{\rho\varphi}(L_{i+2}^\rho) \gtrsim \dots \gtrsim F^{\rho\varphi}(L_{n-1}^\rho)$ and that for all $L = \sum a_i L_i^\Omega$, $\sum |a_i|^2 = 1$, orthogonal to the L_j^ρ , $i+2 \leq j \leq n-1$, then $F^{\rho\varphi}(L) \gtrsim F^{\rho\varphi}(L_{i+1}^\rho)$. Let $L = \sum a_i L_i^\Omega$, $\sum |a_i|^2 = 1$, orthogonal to the L_j^ρ , $i+1 \leq j \leq n-1$. Suppose T_i is chosen. If $T_i \in \mathcal{H}_i$ then $L_i^\rho = T_i$, $F^{\rho\varphi}(T_i) \lesssim F_i^\Omega + \frac{\varphi'}{\delta}$ and, using (3.18), $F^{\rho\varphi}(L) \gtrsim F^{\rho\varphi}(L_i^\rho)$. Suppose now $T_i \notin \mathcal{H}_i$. If $L_i^\rho = T_i$ then, decomposing $L = \alpha(T_i + \gamma H)$ as in the first step, we obtain $F^{\rho\varphi}(L) \gtrsim \frac{\varphi''}{\delta} |\langle T_i(q), p \rangle|^2$ and we use (3.18). Otherwise $L_i^\rho = W_i$ and, again, the same decomposition gives $F^{\rho\varphi}(L) \gtrsim \sum_{j=1}^{i-1} |a_j|^2 F(L_j^\Omega, q, \delta)$ and we conclude with (3.18).

Finally we obtain $F^{\rho\varphi}(L) \gtrsim F^{\rho\varphi}(L_i^\rho)$ (which proves the statement about the choice of T_{n-k}), and, as L_i^ρ is orthogonal to $L_{i+1}^\rho, \dots, L_{n-1}^\rho$, the induction hypothesis imply $F^{\rho\varphi}(L_i^\rho) \gtrsim F^{\rho\varphi}(L_{i+1}^\rho)$ and finishes the proof. \square

We now estimate the brackets of the vector fields L_i^ρ , $i < n$, at the point q .

Lemma 3.7. Let $\left[\begin{smallmatrix} \overleftarrow{L}_k^\rho \\ \overleftarrow{L}_s^\rho \end{smallmatrix} \right] = \sum_{t=1}^n b_{ks}^t L_t^\rho + \sum_{t=1}^n \overline{b_{ks}^t} \overline{L}_t^\rho$. For all lists \mathcal{L} , of $\mathcal{L}_M(\mathcal{B}^\rho \cup \{L_n^\rho\})$, we have

$$\left| \mathcal{L} \left(\begin{smallmatrix} \overleftarrow{b}_{ks}^t \end{smallmatrix} \right) (q) \right| < K' (F^{\rho\varphi})^{\mathcal{L}/2} (F_k^{\rho\varphi})^{1/2} (F_s^{\rho\varphi})^{1/2} (F_t^{\rho\varphi})^{-1/2}$$

with K' depending only on K and the data.

Proof. Note that the Lemma is trivial if $l_n(\mathcal{L}) \geq 1$ and if $F_t^{\rho\varphi} \lesssim \frac{\varphi''(|p|^2)}{\delta}$ (because $F_k^{\rho\varphi}$ and $F_s^{\rho\varphi}$ are both \geq to $\frac{\varphi'(|p|^2)}{\delta}$ and $\frac{\varphi''(|p|^2)}{\delta} \geq \delta^{-2/M}$ implies $\left| \frac{\varphi'(|p|^2)}{\delta} \right|^2 \geq \left| \frac{\varphi''(|p|^2)}{\delta} \right|$). Moreover, we also have $F_t^{\rho\varphi} \lesssim F(L_t^\rho, q, \delta) + \frac{\varphi''(|p|^2)}{\delta}$, and, if $L_t^\rho = T_t$, then, by the definition of T_t and the extremality of \mathcal{B}^Ω , $F(L_t^\rho, q, \delta) \lesssim F(L_t^\Omega, q, \delta)$, and, if $L_t^\rho = W_t$, then $F(L_t^\rho, q, \delta) \lesssim F(L_t^\Omega, q, \delta) + \frac{\varphi''(|q|^2)}{\delta}$.

Thus, it suffices to prove that if $l_n = 0$

$$\left| \mathcal{L} \left(\begin{smallmatrix} \overleftarrow{b}_{ks}^t \end{smallmatrix} \right) (q) \right| \lesssim (F^{\rho\varphi})^{\mathcal{L}/2} (F_k^{\rho\varphi})^{1/2} (F_s^{\rho\varphi})^{1/2} \left(F(L_t^\Omega, q, \delta) \right)^{-1/2}.$$

Let us write $L_k^\rho = \sum \alpha_k^i L_i^\Omega$ and $L_s^\rho = \sum \beta_s^j L_j^\Omega$. Using the notation $\left[\begin{smallmatrix} \overleftarrow{L}_i^\Omega \\ \overleftarrow{L}_j^\Omega \end{smallmatrix} \right] = \sum_{m=1}^n a_{ij}^m L_m^\Omega + \sum_{m=1}^n \overline{a_{ij}^m} \overline{L}_m^\Omega$, a computation gives, if $t < n$,

$$b_{ks}^t = \sum_m \left(\sum_{i,j} \overleftarrow{\alpha}_k^i \overleftarrow{\alpha}_s^j a_{ij}^m \right) \beta_m^t,$$

with $\beta_m^t = \frac{1}{\det(\alpha)} \sum_\sigma \varepsilon_\sigma \prod_i \alpha_i^{\sigma(i)}$, where σ describes the set of permutations from $\{1, \dots, n-1\} \setminus \{t\}$ onto $\{1, \dots, n-1\} \setminus \{m\}$, and

$$b_{ks}^n = \sum_{i,j} \overleftarrow{\alpha}_k^i \overleftarrow{\alpha}_s^j C_{ij}^{\overleftarrow{c}}$$

with $C_{ij}^{\overleftarrow{c}} = \left[\begin{smallmatrix} \overleftarrow{L}_i^\Omega \\ \overleftarrow{L}_j^\Omega \end{smallmatrix} \right] (\partial\rho)$ (note that this notation gives $c_{ij} = C_{ij}$).

First, we prove that, if $t < m$, then $|\beta_m^t| \lesssim (F_k^{\rho\varphi})^{1/2} (F_s^{\rho\varphi})^{1/2} (F(L_t^\Omega, q, \delta))^{-1/2}$ for any k and s . In that case, there exists an index $i > t$ such that $\sigma(i) \leq t$; if $L_i^\rho = T_i$ then $\alpha_i^{\sigma(i)} = 0$, and if $L_i^\rho = W_i$ then

$$\left| \alpha_i^{\sigma(i)} \right| \leq \left[\frac{\varphi''(|p|^2)}{\delta} \left(F(L_{\sigma(i)}^\Omega, q, \delta) \right)^{-1} \right]^{1/2} \leq \left(\frac{\varphi''(|p|^2)}{\delta} \right)^{1/2} \left(F(L_t^\Omega, q, \delta) \right)^{-1/2} \leq (F_k^{\rho\varphi})^{1/2} (F_s^{\rho\varphi})^{1/2} \left(F(L_t^\Omega, q, \delta) \right)^{-1/2},$$

because $F_m^{\rho\varphi} \gtrsim \delta^{-1/M} + \frac{\varphi'(|p|^2)}{\delta}$ and $\frac{\varphi''(|p|^2)}{\delta} \geq \delta^{-2/M}$ implies $\left(\frac{\varphi'(|p|^2)}{\delta} \right)^2 \geq \frac{\varphi''(|p|^2)}{\delta}$.

To finish the proof, it suffices to remark that the extremality of \mathcal{B}^Ω implies

$$\left| \alpha_k^i \right| \lesssim F(L_k^\rho, q, \delta)^{1/2} F(L_i^\Omega, q, \delta)^{-1/2},$$

and

$$\begin{aligned} \left| \mathcal{L} \left(a_{ij}^m \right) \right| &\lesssim \prod F(L_k^\Omega, q, \delta)^{k/2} F(L_i^\Omega, q, \delta)^{1/2} F(L_j^\Omega, q, \delta)^{1/2} F(L_m^\Omega, q, \delta)^{-1/2} \\ &\lesssim (F^{\rho\varphi})^{\mathcal{L}/2} F(L_i^\Omega, q, \delta)^{1/2} F(L_j^\Omega, q, \delta)^{1/2} F(L_t^\Omega, q, \delta)^{-1/2}, \end{aligned}$$

by Lemma 3.6, for $t \geq m$. □

Then, with the notations introduced before, we consider the basis at p (for D)

$$\widetilde{\mathcal{B}} = \{\widetilde{L}_1, \dots, \widetilde{L}_{n-1}\} \text{ with } \widetilde{L}_i = \frac{1}{\|L_i^\rho \circ \pi\|} (L_i^\rho \circ \pi + (\beta_i \circ \pi)N^\rho \circ \pi).$$

Note that Lemma 3.6 and Lemma 3.7 are proved for the vector fields L_i^ρ but it is easy to see that they are also valid for the vector fields $L_i^\rho / \|L_i^\rho\|$.

To simplify the notations, in the remainder of the proof, the vector fields $\frac{L_i^\rho}{\|L_i^\rho\|}$ will be denoted by L_i^ρ , and the function $\frac{\beta_i}{\|L_i^\rho\|}$ will be denoted β_i so that $\widetilde{L}_i = (L_i^\rho \circ \pi + (\beta_i \circ \pi)N^\rho \circ \pi)$.

Proposition 3.13. *The basis $\widetilde{\mathcal{B}}$ is (K', p, δ) -extremal for a constant K' depending only on K and the data.*

Proof. We first prove condition EB₁, that is, if α_i are complex numbers then

$$F \left(\sum_{i=1}^{n-1} \alpha_i \widetilde{L}_i, p, \delta \right) \simeq \sum_{i=1}^{n-1} |\alpha_i|^2 F(\widetilde{L}_i, p, \delta).$$

By induction, it suffices to see that, for all k ,

$$F \left(\sum_{i=1}^{n-k} \alpha_i \widetilde{L}_i, p, \delta \right) \simeq F \left(\sum_{i=1}^{n-k-1} \alpha_i \widetilde{L}_i, p, \delta \right) + |\alpha_{n-k}|^2 F(\widetilde{L}_{n-k}, p, \delta).$$

To simplify notations we write $\widetilde{X} = \sum_{i=1}^{n-k-1} \alpha_i \widetilde{L}_i$ and $X^\rho = \sum_{i=1}^{n-k-1} \alpha_i L_i^\rho$. By Proposition 3.12, we have to prove that

$$\begin{aligned} (3.19) \quad &F(X^\rho + \alpha_{n-k} L_{n-k}^\rho, q, \delta) + \frac{\varphi''(|p|^2)}{\delta} |\langle (X^\rho + \alpha_{n-k} L_{n-k}^\rho) \circ \pi(p), p \rangle| + \frac{\varphi'(|p|^2)}{\delta} \\ &\simeq F(X^\rho, q, \delta) + |\alpha_{n-k}|^2 F(L_{n-k}^\rho, q, \delta) + \frac{\varphi''(|p|^2)}{\delta} \left(|\langle (X^\rho \circ \pi)(p), p \rangle|^2 + |\alpha_{n-k}|^2 |\langle (L_{n-k}^\rho \circ \pi)(p), p \rangle|^2 \right) + \frac{\varphi'(|p|^2)}{\delta}. \end{aligned}$$

Indeed, if $\beta(q) = \frac{\|\sum_{i=1}^t \alpha_i L_i^\rho\|(q)}{\|\sum_{i=1}^t \alpha_i L_i^\rho\|(p)}$, then the \mathcal{C}^M norm of β^{-1} is controlled by K and $F(\beta^{-1} \sum_{i=1}^t \alpha_i \widetilde{L}_i, p, \delta) \simeq_K F(\sum_{i=1}^t \alpha_i \widetilde{L}_i, p, \delta)$.

Note that if Y and Z are two linear combinations (with constant coefficients) of the L_i^Ω , by extremality, $F(Y + Z, q, \delta) \leq K^2 [F(Y, q, \delta) + F(Z, q, \delta)]$, and then

$$(3.20) \quad F(Y + Z, q, \delta) \geq \frac{1}{K^2} F(Y, q, \delta) - F(Z, q, \delta).$$

This implies that the left hand side of (3.19) is \lesssim than the right hand side, and we have only to prove the converse inequality. To do it, we consider separately the two possibilities for L_{n-k}^ρ .

Suppose first $L_{n-k}^\rho = T_{n-k}$.

If the right hand side of (3.19) is equivalent to $F(X^\rho, q, \delta) + |\alpha_{n-k}|^2 F(L_{n-k}^\rho, q, \delta)$, by (3.20), we have only to consider the case when $F(X^\rho, q, \delta) \simeq |\alpha_{n-k}|^2 F(L_{n-k}^\rho, q, \delta)$. Using that $F(T_{n-k}, q, \delta) \lesssim F(L_{n-k}^\rho, q, \delta)$, Lemma 3.6 gives the result.

Suppose now that the right hand side of (3.19) is equivalent to

$$\frac{\varphi''(|p|^2)}{\delta} \left(|\langle (X^\rho \circ \pi)(p), p \rangle|^2 + |\alpha_{n-k}|^2 |\langle (L_{n-k}^\rho \circ \pi)(p), p \rangle|^2 \right).$$

Then, we only have to consider the case when $\langle (X^\rho \circ \pi)(p), p \rangle = -(1 + \varepsilon) \alpha_{n-k} \langle (L_{n-k}^\rho \circ \pi)(p), p \rangle$, with ε small. Then if W is the vector field $X^\rho + (1 + \varepsilon) \alpha_{n-k} L_{n-k}^\rho$ normalized at q , $W \in \mathcal{H}_{n-k}$ and thus $F(W, q, \delta) \geq \frac{\varphi''}{\delta} |\langle T_{n-k}(q), p \rangle|^2 = \frac{\varphi''}{\delta} |\langle L_{n-k}^\rho(q), p \rangle|^2$. Then $F(X^\rho, q, \delta) \gtrsim \frac{1}{K^2} \left(\frac{\varphi''}{\delta} |\langle L_{n-k}^\rho(q), p \rangle|^2 \right) - 2F(L_{n-k}^\rho, q, \delta)$, and the conclusion follows.

To finish, suppose that $L_{n-k}^\rho = W_{n-k}$.

If the right hand side of (3.19) is equivalent to $\frac{\varphi''(|p|^2)}{\delta} \left(|\langle (X^\rho \circ \pi)(p), p \rangle|^2 + |\alpha_{n-k}|^2 |\langle (L_{n-k}^\rho \circ \pi)(p), p \rangle|^2 \right)$, there is nothing to do because $\langle L_{n-k}^\rho \circ \pi(p), p \rangle = 0$.

Suppose then that the right hand side of (3.19) is equivalent to $F(X^p, q, \delta) + |\alpha_{n-k}|^2 F(L_{n-k}^p, q, \delta)$. As before, the conclusion is evident except if $F(X^p, q, \delta) \simeq |\alpha_{n-k}|^2 F(L_{n-k}^p, q, \delta)$. Suppose

$$F(X^p + \alpha_{n-k} L_{n-k}^p, q, \delta) + \frac{\varphi'(|p|^2)}{\delta} \ll |\alpha_{n-k}|^2 F(W_{n-k}, q, \delta).$$

Note that $\langle T_{n-k}(q), p \rangle \neq 0$, and we can define $W = X^p + \alpha_{n-k} L_{n-k}^p + \mu T_{n-k}$ such that $\langle W(q), p \rangle = 0$. Then by Lemma 3.6,

$$|\alpha_{n-k}|^2 F(W_{n-k}, q, \delta) \gg F(L_{n-k}^\Omega, q, \delta),$$

and (extremality of \mathcal{B}^Ω) $|\langle T_{n-k}(q), p \rangle|^2 \frac{\varphi'(|p|^2)}{\delta} > \frac{1}{K} (F(W_{n-k}, q, \delta) - KF(L_{n-k}^\Omega, q, \delta))$. From this we deduce $|\mu| \ll |\alpha_{n-k}|$ and W is of norm almost 1 at q . Then

$$\begin{aligned} F(W, q, \delta) &\leq K^2 \left(F(X^p + \alpha_{n-k} L_{n-k}^p, q, \delta) + |\mu|^2 F(T_{n-k}, q, \delta) \right) \\ &\ll |\alpha_{n-k}|^2 \left(F(W_{n-k}, q, \delta) + F(L_{n-k}^\Omega, q, \delta) \right), \end{aligned}$$

because $T_{n-k} \in \mathcal{E}(L_{n-k}^\Omega, \dots, L_{n-1}^\Omega)$, and thus $F(W, q, \delta) \ll F(W_{n-k}, q, \delta)$ which contradicts the definition of W_{n-k} .

To see that $\tilde{\mathcal{B}}$ satisfy EB₂, a simple computation shows that it suffices to apply Lemma 3.7 and Proposition 3.11. \square

Then, by Lemma 3.1 we conclude:

Proposition 3.14. *The basis \mathcal{B} previously defined by $\mathcal{B} = \{L_i, \dots, L_{n-1}\}$, with $L_i = L_i^p \circ \pi + (\beta_i \circ \pi) N^p \circ \pi$ is (K', p, δ) -extremal for a constant K' depending on the constant K of extremality of \mathcal{B}^Ω and the data.*

Now the proof of Theorem 3.3 is complete.

4. GEOMETRICALLY SEPARATED DOMAINS

4.1. Definition and examples.

Definition 4.1. Let $\Omega = \{\rho < 0\}$ be a bounded pseudo-convex domain with \mathcal{C}^∞ boundary ($\nabla \rho \neq 0$ in a neighborhood of $\partial\Omega$). We say that Ω is K -geometrically separated at $p_0 \in \partial\Omega$ if p_0 is a point of finite 1-type τ and there exist two neighborhoods of p_0 , $W(p_0) \Subset V(p_0)$, a constant $\delta_0 > 0$, a constant $K > 0$, an integer M larger than $\tau + 1$ and a basis $\mathcal{B}^0 = \{L_1^0, \dots, L_{n-1}^0\}$ of $(1, 0)$ vector fields tangent to ρ in $V(p_0)$, whose \mathcal{E}^{2M} norm are bounded by K and their ‘‘determinant’’ bounded from below by $1/K$, and a positive real number δ_0 such that:

For each point $p \in W(p_0) \cap \partial\Omega$ and each δ , $0 < \delta < \delta_0$, there exists a (M, K, p, δ) extremal basis $\mathcal{B}(p, \delta) = \{L_1^{p, \delta}, \dots, L_{n-1}^{p, \delta}\}$ such that, for each i , the vector field $L_i^{p, \delta}$ can be written (on $V(p_0)$) $L_i^{p, \delta} = \sum_j a_i^j L_j^0$ with $a_i^j \in \mathbb{C}$, $\sum |a_i^j|^2 = 1$. In other words, the $L_i^{p, \delta}$ are normalized vector fields belonging to the vector space E_0 generated by \mathcal{B}^0 .

A notable property (that will not be used later) of these domains is that the weights F_i satisfy a better estimate than the one given in Proposition 3.5:

Proposition. *Suppose Ω is geometrically separated at p_0 (of type τ). Then for $V(p_0)$ and δ_0 sufficiently small, there exists a constant $C > 0$ depending only on K and Ω , such that the extremal basis $\mathcal{B}(p, \delta) = \{L_i^{p, \delta}, 1 \leq i \leq n-1\}$, $p \in W(p_0) \cap \partial\Omega$, $0 < \delta < \delta_0$, satisfies $F_M(L_i^{p, \delta}, p, \delta) \geq C\delta^{-2/\tau+1}$, for all i and all $\delta \in [0, \delta_0]$, with $M = [\tau] + 1$.*

Proof. Suppose there exists a sequence of points p_m converging to p_0 , a sequence δ_m in $]0, \delta_0[$ and an integer $i \leq n-1$ such that, denoting $\mathcal{B}(p_m, \delta_m) = (L_1^m, \dots, L_{n-1}^m)$ the (M, K, p_m, δ_m) -extremal basis at p_m , we have $\sum_{\mathcal{L} \in \mathcal{L}_M(L_i^m)} |\mathcal{L}(\partial\rho)(p_m)| \leq 1/m$. Then $L_i^m = \sum a_i^j(p_m) L_j^0$, $\sum |a_i^j(p_m)|^2 = 1$, and we may suppose that the sequences $n \mapsto a_i^j(p_m)$ converge to complex numbers a^j satisfying $\sum |a^j|^2 = 1$. Then, by uniform convergence, the vector field $L = \sum a^j L_j^0$ satisfies $F_M(L, p_0, \delta) = 0$, for all δ . But, we have $L = \sum b_k L_k^{p_0}$, $\sum |b_k|^2 \geq K$, and, by extremality $F(L, p_0, \delta) \simeq_K \sum |b_k|^2 F_M(L_k^{p_0}, p_0, \delta)$, thus there exists k such that $F_M(L_k^{p_0}, p_0, \delta) = 0$, i. e. $\sum_{\mathcal{L} \in \mathcal{L}_M(L_k^{p_0})} |\mathcal{L}(\partial\rho)(p_0)| = 0$. Then, by (4) of Definition 3.2 this contradicts the definition of the 1-type. \square

Thus, in all the paper, for a geometrically separated domain at a boundary point p_0 , the integer M could be changed to $[\tau] + 1$. As this change gives no advantage, we will keep $M = M'(\tau)$ and then we can apply directly the results of the preceding Sections.

Remark 4.1. Suppose Ω is geometrically separated at $p_0 \in \partial\Omega$. Let p be a point of $\overline{\Omega} \cap W(p_0)$. If π is the projection onto $\partial\Omega$ defined in Section 3.5.2 let $q = \pi(p)$. Then, reducing $W(p_0)$ and δ_0 if necessary, if $-\frac{1}{3}\rho(p) < \delta < \delta_0$, the basis $\mathcal{B}(q, \delta) = (L_1^{q, \delta}, \dots, L_{n-1}^{q, \delta})$ is clearly $(2K, p, \delta)$ -extremal, and $F_M(L_i^{q, \delta}, p, \delta) \geq C'\delta^{-2/\tau+1}$ for a constant C' depending only on K and the data. Thus we will always assume that a geometrically separated domain is equipped, by definition, with extremal bases of the form given in the definition, at every point of $V(p_0) \cap \overline{\Omega}$ for $-\frac{1}{3}\rho(p) < \delta < \delta_0$.

This is clear, because if $\mathcal{L} \in \mathcal{L}_M(\mathcal{B})$, then $|\mathcal{L}(\partial\rho)(p) - \mathcal{L}(\partial\rho)(\pi(p))| = O(\delta)$, where O depends only on K and Ω . Then EB_1 is satisfied because $F_i(p, \delta) \geq C\delta^{-2/M}$ with C depending only on Ω and EB_2 is also satisfied because $F_k(p, \delta) \leq \delta^{-2}$ (δ_0 small enough).

Example 4.1.

- (1) The three first examples of extremal basis given in Example 3.1 immediately show that, if p_0 is a point of finite type of $\partial\Omega$ then Ω is geometrically separated at p_0 , under one of the following four conditions:
 - (a) $\partial\Omega$ is convex near p_0 , or, more generally, linearly convex near p_0 (see Section 7.1);
 - (b) The eigenvalues of the Levi form are comparable at p_0 ;
 - (c) The Levi form is locally diagonalizable at p_0 .
 - (d) Near p_0 , $\partial\Omega$ belongs to the class introduced by M. Derridj in [Der99].
- (2) Moreover, we will see in Section 4.3 that, if Ω is geometrically separated at p_0 then the local domain D defined in Section 3.5.1 is geometrically separated at every point of its boundary.

Example 4.2. The domain $\Omega = \{z \in \mathbb{C}^3 \text{ such that } \Re z_1 + |z_2|^6 + |z_3|^6 + |z_2 z_3|^2 < 0\}$ studied by G. Herbort in [Her83] is not geometrically separated at $(0, 0)$ (see Section 7.2 for details).

4.2. Structure of homogeneous space. First recall that we define in Section 3.3.4 the ‘‘polydisc’’ $B^c(\mathcal{B}, p, \delta)$ (Definition 3.3) and the ‘‘pseudo-balls’’ $B_{\text{exp}}^c(\mathcal{B}, p, \delta)$ and $B_{\mathcal{C}}^c(\mathcal{B}, p, \delta)$ (Definition 3.4).

In general, we will just denote by $B_{\text{exp}}^c(p, \delta)$ and $B_{\mathcal{C}}^c(p, \delta)$ the pseudo-balls $B_{\text{exp}}^c(\mathcal{B}, p, \delta)$ and $B_{\mathcal{C}}^c(\mathcal{B}, p, \delta)$ omitting \mathcal{B} , but recall that, if $\delta_1 \neq \delta_2$, the balls $B_{\text{exp}}^c(p, \delta_1)$ and $B_{\text{exp}}^c(p, \delta_2)$ are not necessarily constructed with the same basis.

Then by the methods used in [CD06b] (based on the Campbell-Hausdorff formula and the ideas of [NSW85]), reducing $W(p_0)$ if necessary, one can prove the following properties of the balls:

Proposition 4.1. *There exist constants $c_0, \delta_0, \alpha, \beta$ and γ such that, for $p \in W(p_0) \cap \partial\Omega$, $\delta \leq \delta_0$ and $c \leq c_0$, we have $B_{\text{exp}}^{\alpha c}(p, \delta) \subset B^c(p, \delta) \subset B_{\text{exp}}^{\beta c}(p, \delta)$ and $B_{\text{exp}}^c(p, \delta) \subset B_{\mathcal{C}}^c(p, \delta) \subset B_{\text{exp}}^{\gamma c}(p, \delta)$.*

The importance of this Proposition to construct the structure of homogeneous space is the following: to be able to use Taylor’s formula, we have to work with a coordinates system, which is easy in the sets $B^c(p, \delta)$; the hypothesis of geometric separation and Proposition 3.6 imply that the sets associated to curves are associated to a pseudo-distance; and, finally, the sets associated to the exponential map are used to prove that all these sets are equivalent.

Ideas of the proof of Proposition 4.1. It is similar to the proofs of Proposition 3.4 (p. 96) and Lemma 3.16 (p. 101) of [CD06b]. Thus we will only give the main articulations.

The first inclusion comes easily from the control of the coefficients of the vector fields L_i in the coordinate system (z_i) in the polydisc (Proposition 3.6). The second one is more complicated.

Let exp_p be the exponential map based at p relatively to the vector fields \mathcal{Y}_i (real and imaginary parts of the L_i). Let $\Psi^p = (\Psi_i^p)_{i=2, \dots, 2n} = (\text{exp}_p)^{-1}$. We establish the following estimate on the derivatives of the functions Ψ_i^p : there exist constants β and K_1 , depending on K and the data, such that

$$(4.1) \quad \text{if } q = \text{exp}_p(u), \max\{|u_i|, |u_{i+n}|\} \leq \beta F_i(p, \delta)^{-1/2} \text{ then } \left| \mathcal{Y}_k \Psi_j^p(q) \right| \leq K_1 F_k(p, \delta)^{1/2} F_j(p, \delta)^{-1/2},$$

with the notation of Definition 3.4.

To prove this, we estimate the derivatives of the exponential map. Considering, for $u \in \mathbb{R}^n$, the vector field $\mathcal{Y}_u = \sum u_i \mathcal{Y}_i$, the derivatives of exp_p are estimated via the Campbell-Hausdorff formula. Let $q = q(u) = \text{exp}_p(u)$, $|u| \leq u_0$,

$$\left| d\text{exp}_p \left(\frac{\partial}{\partial u_i} \right) (u) - \mathcal{Y}_i(q) + \sum_{k=2}^M \alpha_k [\mathcal{Y}_u, \dots [\mathcal{Y}_u, \mathcal{Y}_i] \dots] (q) \right| \leq C |u|^{M+1},$$

where α_k are universal constants corresponding to brackets of length k (see Lemma 1 (p. 97) of [CD06b]). The brackets are then estimated with Proposition 3.6 and thus (4.1) is easily obtained. The second inclusion of the Proposition is then easily proved.

The equivalence between the sets defined with the exponential map and the curves is a quite simple consequence of (4.1). \square

Proposition 4.2. *Let Ω be a bounded pseudo-convex domain K -geometrically separated at $p_0 \in \partial\Omega$. Let B denote one of the sets $B_{\mathcal{C}}^c, B_{\text{exp}}^c$ or B^c . Then there exists a constant $c_0 > 0$, depending on K and the data such that, for all $c \leq c_0$, the sets $B(\mathcal{B}(p, \delta), p, \delta)$ are associated to a pseudo-distance in the following sense: there exists a constant C depending on K and the data (but not on c) such that, if $p \in W(p_0) \cap \partial\Omega$ and $\delta \leq \delta_0$, and if $q \in B(\mathcal{B}(p, \delta), p, \delta) \cap \partial\Omega$, then*

$$B(\mathcal{B}(q, \delta), q, \delta) \subset B(\mathcal{B}(p, \delta), p, C\delta).$$

Remark. If we define γ , on $W(p_0) \cap \partial\Omega$, by

$$(4.2) \quad \gamma(p, q) = \inf \{ \delta \text{ such that } q \in B(\mathcal{B}(p, \delta), p, \delta) \},$$

then γ is a real pseudo-distance.

Lemma.

(1) For all $A > 0$ there exists B depending on A and K such that

$$B_{\mathcal{C}}^{Ac}(\mathcal{B}(q, \delta), q, \delta) \subset B_{\mathcal{C}}^c(\mathcal{B}(q, B\delta), q, B\delta).$$

(2) For all $B > 0$ there exists C depending on B such that

$$B_{\mathcal{C}}^c(\mathcal{B}(q, B\delta), q, B\delta) \subset B_{\mathcal{C}}^{Cc}(\mathcal{B}(q, \delta), q, \delta).$$

Proof of the Lemma. Let us denote by $L_i(q, \delta)$ (resp $L_i(q, B\delta)$) the vector fields of $\mathcal{B}(q, \delta)$ (resp. $\mathcal{B}(q, B\delta)$). By the hypothesis on Ω , we have $L_i(q, \delta) = \sum_k \beta_i^k L_k(q, B\delta)$, with β_i^k constants. By extremality,

$$\begin{aligned} |\beta_i^k| &\leq KF(L_i(q, \delta), q, B\delta)^{1/2} F(L_k(q, B\delta), q, B\delta)^{-1/2} \\ &\leq KB^{-1/M} F(L_i(q, \delta), q, \delta)^{1/2} F(L_k(q, B\delta), q, B\delta)^{-1/2}, \end{aligned}$$

which proves the first part of the Lemma with $B = (AK(n-1))^M$. The second part is proved similarly with $C = (BK(n-1))^M$. \square

Proof of Proposition 4.2. To prove the assertion on the pseudo-distance in the Proposition, by Proposition 4.1, it is enough to prove that, there exists a constant K_0 such that if $q, q' \in B_{\mathcal{C}}^c(\mathcal{B}(p, \delta), p, \delta)$ then $q' \in B_{\mathcal{C}}^{K_0c}(\mathcal{B}(q, \delta), q, \delta)$. But there exists φ , \mathcal{C}^1 piecewise smooth, such that $\varphi(0) = q$, $\varphi(1) = q'$ and, almost everywhere, $\varphi'(t) = \sum_{i=1}^{2n} a_i(t) \mathcal{Y}_i(\varphi(t))$, with $\max\{|a_i(t)|, |a_{i+n}(t)|\} \leq 2cF(L_i(p, \delta), p, \delta) \leq 4cF(L_i(p, \delta), q, \delta)$, if we choose c small enough (Proposition 3.6). Now, as in the Lemma, writing $L_i(p, \delta) = \sum \alpha_i^k L_k(q, \delta)$ (with α_i^k constants) and using extremality, we easily conclude $q' \in B_{\mathcal{C}}^{K_0c}(\mathcal{B}_1^{q, \delta}, q, \delta)$. \square

Let us now define the ‘‘pseudo-balls’’ centered at points of $\Omega \cap W(p_0)$, denoted ${}^{\pi}B^c(q, \delta)$ (resp. ${}^{\pi}B_{\mathcal{C}}^c(q, \delta)$, ${}^{\pi}B_{\text{exp}}^c(q, \delta)$) by

$${}^{\pi}B^c(q, \delta) = \{q' \in V(p_0) \text{ such that } \pi(q') \in B^c(\mathcal{B}(\pi(q), \delta), \pi(q), \delta) \text{ and } \rho(q') \in [\rho(q) - c\delta, \rho(q) + c\delta]\}.$$

Then:

Theorem 4.1. *Let Ω be a pseudo-convex domain geometrically separated at $p_0 \in \partial\Omega$. There exists a constant $c_0 > 0$, depending on K and the data, such that, for all $c \leq c_0$, the sets $B(q, \delta)$ define a structure of ‘‘homogeneous space’’ on $W(p_0) \cap \bar{\Omega}$ in the following sense: there exists a constant C , depending only on K and the data (not on c) such that, if $q_1 \in W(p_0) \cap \bar{\Omega}$, $\delta < \delta_0$, and $q_2 \in B(q_1, \delta)$, we have*

$$B(q_2, \delta) \subset B(q_1, C\delta)$$

and

$$\text{Vol}(B(q, 2\delta)) \leq C \text{Vol}(q, \delta),$$

B denoting one of the sets ${}^{\pi}B_{\mathcal{C}}^c$, ${}^{\pi}B_{\text{exp}}^c$ or ${}^{\pi}B^c$.

Proof. The first assertion follows immediately the Proposition. To prove the second assertion, we use the fact that both $B_{\mathcal{C}}^c(\mathcal{B}(p, \delta), p, \delta)$ and $B_{\text{exp}}^c(\mathcal{B}(p, \delta), p, \delta)$ are equivalent to $B^c(\mathcal{B}(p, \delta), p, \delta)$, the fact that the coordinate system associated to the extremal basis have a Jacobian uniformly bounded from above and below and the preceding Lemma. \square

Remark 4.2.

- (1) For $p \in \partial\Omega$, the sets ${}^{\pi}B^c(q, \delta) \cap \partial\Omega$ (for each definition) are the pseudo-balls of a structure of homogeneous space on $\partial\Omega \cap W(p_0)$.
- (2) On $\partial\Omega$, as in [NRSW89], we could define equivalent pseudo-balls using complex tangent curves.
- (3) It is not difficult to see that the pseudo-balls of the structure of homogeneous space can also be defined directionnally: they are equivalent to the sets $B_{\text{dir}}^c(p, \delta)$ defined as to be the set of points q of the form $q = \exp_p(a, b)$, where \exp_p is the exponential map associated to the vector field $a\Re L + b\Im L$, L being a vector field of the linear space E , the coefficients a and b satisfying $\max(|a|, |b|) \leq cF(L, p, \delta)^{-1/2}$.

4.3. Localization. Suppose that Ω is K -geometrically separated at $p_0 \in \partial\Omega$, and consider the domain D constructed in Section 3.5.1 near that point. Then D is K -geometrically separated at each point of $\partial\Omega \cap \partial D$, and, by strict pseudo-convexity, the same is true on $\partial D \setminus \overline{\partial\Omega} \cap \partial D$.

Suppose that P is a point of the boundary of $\partial\Omega \cap \partial D$, and let p be a point of $V(P) \cap \partial D$ and δ small enough (with the notations of the previous Section). Let us denote by $\mathcal{B}(p, \delta) = \{L_1^{p, \delta}, \dots, L_{n-1}^{p, \delta}\}$ the extremal basis given by Proposition 3.14 and by $\mathcal{B}^{0, \Omega} = \{L_1^{0, \Omega}, \dots, L_{n-1}^{0, \Omega}\}$ the basis denoted \mathcal{B}^0 in Definition 4.1. Then, by the construction of $\mathcal{B}(p, \delta)$ made in the previous Section, we have $L_i^{p, \delta} = L_i^p \circ \pi - \beta(L_i^p)N^p \circ \pi$ with $L \mapsto \beta(L)$ linear. Thus, if we define $\mathcal{B}^{0, D} = \{L_1^{0, D}, \dots, L_{n-1}^{0, D}\}$ by $L_i^{0, D} = L_i^{0, \Omega} \circ \pi - \beta(L_i^{0, \Omega})N^p \circ \pi$, then we see that the vector fields of $\mathcal{B}(p, \delta)$ are linear combinations (with constant coefficients) of the vector fields of $\mathcal{B}^{0, D}$. Thus, we have proved the following result:

Theorem 4.2. *If Ω is K -geometrically separated at $p_0 \in \partial\Omega$, then the domain D defined in Definition 3.6 is K' -geometrically separated (at every point of its boundary) for a constant K' depending only on K and the data.*

Remark. Recall that every point of ∂D is of finite 1-type.

5. ADAPTED PLURI-SUBHARMONIC FUNCTION FOR GEOMETRICALLY SEPARATED DOMAINS

5.1. Definition and examples.

Definition 5.1. Let Ω be geometrically separated at p_0 . Let E be the vector space generated by $\mathcal{B}^0 \cup \{N\}$, and, if $L = \sum_{i=1}^{n-1} b_i L_i^0 + b_n N = L_\tau + b_n N \in E$ denotes, for $\delta \leq \delta_0$, $F(L, q, \delta) = F(L_\tau, q, \delta) + \frac{|b_n|^2}{\delta^2}$.

A \mathcal{C}^3 pluri-subharmonic function in Ω , H_δ , is said to be β -adapted to \mathcal{B}^0 at p_0 if there exists a constant β such that the following properties hold:

- (1) $|H_\delta| \leq 1$ on Ω ;
- (2) For $q \in W(p_0) \cap \Omega \cap \{\rho \geq -2\delta\}$ and for all vector fields $L \in E$,

$$\langle \partial \bar{\partial} H_\delta; L, \bar{L} \rangle (q) \geq \frac{1}{\beta} F(L, q, \delta);$$

- (3) For $q \in W(p_0) \cap \Omega \cap \{\rho \geq -2\delta\}$ and for all lists $\mathcal{L} \in \mathcal{L}_3(E)$,

$$|\mathcal{L} H_\delta| (q) \leq \beta \prod_{L \in \mathcal{L}} F(L, q, \delta)^{1/2}.$$

Remark 5.1. Note that (3) implies in particular that, for all $\mathcal{L} \in \mathcal{L}_3(\mathcal{B}(\pi(q), \delta) \cup \{N\})$,

$$|\mathcal{L} H_\delta| (q) \lesssim F(\mathcal{B}(\pi(q), \delta), q, \delta)^{\mathcal{L}/2}.$$

Definition 5.2. A bounded pseudo-convex domain Ω is called “ K -completely geometrically separated” at p_0 if it is K -geometrically separated and, there exists $\delta_0 > 0$ such that, for all $0 < \delta \leq \delta_0$, there exists a pluri-subharmonic function H_δ which is K -adapted to \mathcal{B}^0 at p_0 .

Example 5.1.

- (1) If the boundary of Ω is locally convex near p_0 (a point of finite type), then it is proved in [McN94, McN02] that it is completely geometrically separated at p_0 . More generally, using the results of [DF03] it can be shown that if Ω is locally linearly convex near p_0 (see [Kis98]) then it is completely geometrically separated at p_0 (see Section 7.1 for some details on the construction). Moreover, when the boundary of Ω is locally convex, resp. locally linearly convex, near p_0 , the local domain D can be chosen convex, resp. linearly convex, (choosing d small enough and K_0 large enough) and thus, in both cases, it is completely geometrically separated at every point of its boundary.
- (2) In [Cho02b, Cho02a, Cho03], it is proved that, at a point of finite type, if the eigenvalues of the Levi form are comparable at p_0 then it is also completely geometrically separated at p_0 .
- (3) In the next Section, we prove that geometrically separated domains whose extremal bases are strongly extremal with a sufficiently small α are completely geometrically separated, and, moreover that, for those domains, the local domain defined in Section 3.5 is completely geometrically separated at every point of its boundary. In particular, this applies when the Levi form is locally diagonalizable at p_0 .
- (4) It can also be proved that if a domain is of the type considered by M. Derridj in [Der99] near a boundary point p_0 then it is completely geometrically separated at p_0 .

5.2. The case of geometrically separated domains with strongly extremal bases. In this Section we prove the two following Theorems:

Theorem 5.1. *Suppose Ω is K -geometrically separated at $p_0 \in \partial\Omega$. Then there exists a constant α_0 , depending on K and the data, such that, if for all $p \in W(p_0) \cap \partial\Omega$ and $\delta \leq \delta_0$, the bases $\mathcal{B}(p, \delta)$ are (K, α, p, δ) -strongly extremal (c.f. Definition 3.5) with $\alpha \leq \alpha_0$ then it is completely geometrically separated at p_0 .*

The second theorem deals with the local domain D constructed in Section 3.5.1, and, in fact contains the first one:

Theorem 5.2. *Suppose that Ω is K -geometrically separated at $p_0 \in \partial\Omega$. There exists a constant α_1 , depending on K and the data such that, if for all $p \in W(p_0) \cap \partial\Omega$ and $\delta \leq \delta_0$, the bases $\mathcal{B}(p, \delta)$ are (K, α, p, δ) -strongly extremal with $\alpha \leq \alpha_1$, then the local domain constructed in Section 3.5.1 is K' -completely geometrically separated at every point of its boundary for a constant K' depending only on K and Ω .*

We will prove in details the first Theorem and only give the modifications needed to obtain the second one.

5.2.1. *Proof of Theorem 5.1.* Here we suppose that the bases $\mathcal{B}(p, \delta)$, $p \in W(p_0) \cap \partial\Omega$, $\delta \leq \delta_0$, are (K, α, p, δ) -strongly extremal for a constant α not yet fixed. During the proof, we will impose successive conditions on α (depending on K , M and n) to be able to construct the good pluri-subharmonic function. The existence of α will be clear at the end of the proof but we will not give an explicit value. Now, we fix $\delta > 0$.

The ideas of construction are comparable to those developed in [CD06b] (following ideas of [Cat87]) but the technical proofs are slightly different. On the one hand the basis are local instead of global and we have to construct local ‘‘almost pluri-subharmonic’’ functions and then add them using the structure of homogeneous space instead of constructing directly a global function. On the other hand, the control of lists following our hypothesis are weaker than those following the local diagonalizability of the Levi form. Thus, for reader’s convenience, we will write the proof with enough details.

Let us first introduce some new notations: δ being fixed, we denote by $Q^c(p, \delta)$ the points q in $W(p_0)$ such that $\pi(q)$ belongs to the polydisc $B^c(p, \delta)$, associated to the extremal basis $\mathcal{B}(p, \delta) = (L_i^{p, \delta})_i$ (see Definition 3.3). If L is a vector field in E (the vector space generated by \mathcal{B}^0 and N), we write it $L = L_\tau + a_n N$, where L_τ is tangent to ρ . Ω being geometrically separated we can write $L_\tau = \sum_{i=1}^{n-1} a_i^p L_i^{p, \delta}$ ($a_i^p \in \mathbb{C}$). As usual, c_{ii}^p will denote the coefficient of the Levi form associated to the vector field $L_i^{p, \delta} \in \mathcal{B}(p, \delta)$, and $\Omega_\varepsilon = \{-\varepsilon < \rho < 0\}$.

With these notations, we now we state a local result and show how it leads to Theorem 5.1. For the proof we need only estimates in the strip $\Omega_{3\delta} = \{-3\delta \leq \rho \leq 0\}$, but in Section 5.2.3 we will need corresponding results in a larger domain, and thus we state the local result for the sets $Q^c(p, \delta)$:

Proposition 5.1. *For all constants $C > 1$ there exist constants α_0 (depending only on K , c , C and the data), β and γ_1 such that if the bases $\mathcal{B}(p, \delta)$ are (K, α, p, δ) -extremal with $\alpha \leq \alpha_0$, then for all $\delta \leq \delta(\alpha_0)$ (depending on α_0 , K and the data) and all point $p \in W(p_0) \cap \partial\Omega$, there exists a function $H_{p, \delta} = H$ with support in $Q^c(p, \delta)$ satisfying, for every vector field $L \in E$, the following conditions:*

- (1) $|H| \leq 1$;
- (2) $\langle \partial \bar{\partial} H; L, \bar{L} \rangle(q) \geq \beta F(L_\tau, q, \delta) - \gamma_1 \left(\sum_{i=1}^{n-1} |a_i^p|^2 \frac{c_{ii}}{\delta} + \frac{|a_n|^2}{\delta^2} + 1 \right)(q)$, for $q \in Q^{c/2}(p, \delta) \cap \Omega_{3\delta}$.
- (3) $\langle \partial \bar{\partial} H; L, \bar{L} \rangle(q) \geq -\frac{\beta}{C} F(L_\tau, q, \delta) - \gamma_1 \left(\sum_{i=1}^{n-1} |a_i^p|^2 \frac{c_{ii}}{\delta} + \frac{|a_n|^2}{\delta^2} + 1 \right)(q)$, for $q \in Q^{c/2}(p, \delta) \cap \Omega_{3\delta}$.
- (4) For $\mathcal{L} \in \mathcal{L}_3(\mathcal{B}(p, \delta) \cup \{N\})$, $|\mathcal{L}H|(q) \leq \gamma_2 \prod_{L \in \mathcal{L}} F(L, q, \delta)^{1/2}$, for $q \in Q^{c/2}(p, \delta) \cap \Omega_{3\delta}$.

We will prove this Proposition in the next Section. Now we show how Theorem 5.1 follows this Proposition:

Proof of Theorem 5.1. We cover $\partial\Omega \cap W(p_0)$ with a minimal system of pseudo-balls $\pi B^{c/2}(p_k, \delta) \cap \partial\Omega$, $p_k \in \partial\Omega$. As the pseudo-balls are associated to a structure of homogeneous space, there exists an integer S , independent of δ , such that each point of $W(p_0)$ belongs to at most S sets $Q^c(p_j, \delta)$. Applying Proposition 5.1 with $C = 2SC_1$ we get a function $H_{p_k, \delta}$.

For all point $q \in V(P_0) \cap \Omega_{3\delta}$ there exists j_0 such that $q \in Q^{c/2}(p_{j_0}, \delta)$ and thus (denoting c_{ii}^k the coefficient of the Levi form in the direction $L_i^{p_k}$ and $a_i^k = a_i^{p_k}$), by Proposition 5.1,

$$(5.1) \quad \left\langle \partial \bar{\partial} \sum_k H_{p_k, \delta}; L, \bar{L} \right\rangle(q) \geq \frac{\beta}{2} F(L_\tau, q, \delta) - \gamma_1 \sum_{k \text{ s.t. } q \in Q^c(p_k, \delta)} \left(\sum_{i=1}^{n-1} |a_i^k|^2 \frac{|c_{ii}^k(q)|}{\delta} + \frac{|a_n|^2}{\delta^2} + 1 \right).$$

Let us consider the function

$$H = \sum_k H_{p_k, \delta} + A e^{-\rho/\delta} + B |Z|^2,$$

for suitable constants A and B and α small enough:

Claim. There exist constants A , B , γ and α'_0 depending only on K and the data such that if $\alpha \leq \alpha'_0$:

- (1) H is uniformly bounded, independently of $\delta \leq \delta_0$, on $\Omega_{3\delta}$;
- (2) For any vector field $L \in E$ and any $q \in \Omega_{3\delta} \cap W(p_0)$, $\langle \partial \bar{\partial} H; L, \bar{L} \rangle(q) \geq \frac{\beta}{2} F(L_\tau, q, \delta) + \frac{|a_n|^2}{\delta^2}$;
- (3) For $q \in \Omega_{3\delta} \cap W(p_0)$ and all lists $\mathcal{L} \in \mathcal{L}_3(E)$, $|\mathcal{L}H|(q) \leq \gamma_2 \prod_{L \in \mathcal{L}} F(L, q, \delta)^{1/2}$.

Proof of the Claim. For every k such that $q \in Q^c(p_k, \delta)$,

$$\left\langle \partial \bar{\partial} e^{\rho/\delta}; L, \bar{L} \right\rangle(q) = e^{\rho/\delta} \left[\frac{1}{\delta} \left(\frac{1}{2} \sum_{i,j=1}^{n-1} a_i^k \bar{a}_j^k c_{ij}^k + 2\Re \left(\sum_{i=1}^{n-1} a_i^k \bar{a}_n^k \langle \partial \bar{\partial} \rho; L_i^{p_k}, \bar{N} \rangle \right) + |a_n|^2 \langle \partial \bar{\partial} \rho; N, \bar{N} \rangle \right) + \frac{|a_n|^2}{\delta^2} \right].$$

Then, we use the hypothesis of strong extremality and Taylor's formula to estimate $|c_{ij}^k|$, $i \neq j$, in the set $Q^c(p_k, \delta) \cap \Omega_{3\delta}$. Using the fact that $c_{ii} = |c_{ii}| + O(\delta)$ (recall Ω is pseudo-convex), this gives a constant K_0 depending on K and the data such that

$$\langle \partial \bar{\partial} e^{\rho/\delta}; L, \bar{L} \rangle(q) \geq -K_0 + \frac{e^{-3}}{2\delta^2} |a_n|^2 + \frac{e^{-3}}{2\delta} \sum_{i=1}^{n-1} |a_i^k|^2 |c_{ii}^k(q)| - 4n^2 K \alpha F(L_\tau^q, q, \delta),$$

because, by definition of c , in the sets $Q^c(p_k, \delta)$, we have $F(L_\tau, q) \leq 3F(L_\tau, p_k)$ (see Proposition 3.6).

Now if we choose $A = 2Se^3\gamma_1 + 1$ and $B = K_0A + \gamma_1$, the Claim follows easily (5.1). \square

To finish the proof of Theorem 5.1, we truncate H to adapt it to good neighborhoods $V(p_0)$ and $W(p_0)$ and to the strip $\{\delta_\Omega(p) < 2\delta\}$, and we add $D|z|^2$ with a large constant D . More precisely, the cutting functions are defined as follows:

Let $\vartheta = \vartheta_1 \vartheta_2$ where $\vartheta_1(q) = \chi_1\left(\frac{1}{2} \frac{|q-p_0|}{r}\right)$, with χ_1 a \mathcal{C}^∞ increasing function equal to 0 on $]-\infty, 0]$, 1 on $[1/4, +\infty[$ and $\chi_1(t) = t^4$ on $[0, 1/8]$, and $\vartheta_2(q) = \chi_\delta(\rho(q))$ with $\chi_\delta(t) = \chi(t/\delta)$, χ being even, increasing on $]-\infty, 0]$, equal to 0 on $]-\infty, -4]$, to 1 on $]-2, 0[$ and to $\frac{(t+4)^4}{16}$ for $t \in [-4, -8/3]$.

Then, remarking that $\langle \partial \bar{\partial} \vartheta; L, \bar{L} \rangle \geq -O(1)$ the final computation is made as in [CD06b, Section 4.2.3]. \square

5.2.2. *Proof of Proposition 5.1.* The proof uses essentially the ideas developed in Section 4.1 of [CD06b], except that we have to work locally around the point p . Thus the technique is more complicated (it needs to use the structure of homogeneous space) and we will give it with some details.

For $p \in W(p_0) \cap \partial\Omega$ and $\delta \leq \delta_0$ fixed, let $\mathcal{B}(p, \delta) = \{L_i^{p, \delta} = L_i, 1 \leq i \leq n-1\}$ be the (K, α, p, δ) -strongly extremal basis and $\Phi = \Phi_p^\delta$ be the adapted change of coordinates at (p, δ) .

For $i = 1, \dots, n-1$ and $l = 3, \dots, M$, let us define

$$\mathcal{E}_l^i = \{\Re(\mathcal{L}(\partial\rho)), \Im(\mathcal{L}(\partial\rho)), |\mathcal{L}| = l-1, \mathcal{L} = \{L^1, \dots, L^{l-1}\}, L^k \in \{L_i, \bar{L}_i\}\},$$

$$\mathcal{E}^i = \bigcup_l \mathcal{E}_l^i.$$

For $\varphi \in \mathcal{E}^i$, if $\varphi \in \mathcal{E}_l^i$, we denote $\tilde{l}(\varphi) = l$.

Note that $F_i(\cdot, \delta) = F(L_i, \cdot, \delta) \simeq \frac{|c_{ii}|}{\delta} + \sum_{\varphi_i \in \mathcal{E}^i} \left| \frac{L_i \varphi_i}{\delta} \right|^{2/\tilde{l}(\varphi_i)}$. The functions $\frac{|c_{ii}|}{\delta}$ and $\left| \frac{L_i \varphi_i}{\delta} \right|^{2/\tilde{l}(\varphi_i)}$ are called the *components* of F_i and are denoted generically f_i . We also define $l(c_{ii}) = 2$, and, for the other functions f_i , $l(f_i) = \tilde{l}(\varphi_i)$. In the following proof, these components cannot be considered individually. Thus, we introduce the terminology of “ $(n-1)$ -uplet” of components: $f = (f_1, \dots, f_{n-1})$, where f_i are component of F_i , is called a $(n-1)$ -uplet of components of the weights F_i . The set of all such $(n-1)$ -uplet is denoted by \mathcal{H} . \mathcal{H} is ordered by the lexicographic order.

First we define a cutoff function with support in $Q^c(p, \delta)$ and in the set where a component is “dominant”. More precisely, if B is a positive number and $f = (f_i)$ a $(n-1)$ -uplet of components of F_i , we define, for fixed $c \leq c_0$,

$$\chi_{f, B} = \prod_i \chi_B \left(\frac{f_i \circ \pi}{F_i(p, \delta)} \right) \chi_0 = \chi'_{f, B} \chi_0,$$

where $\chi_B(t) = \chi(Bt)$, $\chi : [0, +\infty[\mapsto [0, 1]$, being a \mathcal{C}^∞ function equal to 0 on $[0, 1/2]$ and 1 on $[1, +\infty[$, and $\chi_0(q) = \chi_1\left(\left(\frac{F_i(p, \delta)^{1/2}}{c} \Phi_p(\pi(q))\right)_i\right)$, with χ_1 a \mathcal{C}^∞ function identically 1 on $B(0, 1/2)$ and with compact support in $B(0, 1)$.

We say that f is B dominant if $\chi'_{f, B} = 1$.

Then, to each $(n-1)$ -uplet $f = (f_i)$ and to each i such that $f_i = \left| \frac{L_i^p \varphi_i}{\delta} \right|^{2/l(f_i)}$, we associate, for $\lambda > 1$, the function

$$H_i(f, \lambda, B) = \lambda^{-3/2} e^{\lambda \psi_i} \chi_{f, B},$$

where $\psi_i(q) = \frac{\varphi_i(\pi(q))}{\delta} F_i(p, \delta)^{\frac{1-\tilde{l}(\varphi_i)}{2}}$.

Lemma 5.1. *For each constant $B > 0$, there exists a constant K_0 depending only on B, c, K and the data such that, for each i , if $q \in Q^c(p, \delta) \cap \Omega_{3\delta}$, for each $L = \sum_{j=1}^n a_j L_j$, $\sum |a_j|^2 = 1$, we have the following estimates:*

- (1) $|L\psi_i(q)| \leq K_0 \left(F(L_\tau, q, \delta)^{1/2} + \frac{|a_n|}{\delta} \right)$, and $|\bar{L}L(\psi_i)(q)| \leq K_0 \left(F(L_\tau, q, \delta) + \frac{|a_n|^2}{\delta^2} \right)$;
- (2) $|L\chi_{f, B}(q)| \leq K_0 \left(F(L_\tau, q, \delta)^{1/2} + \frac{|a_n|}{\delta} \right)$, and $|\bar{L}L\chi_{f, B}| \leq K_0 \left(F(L_\tau, q, \delta) + \frac{|a_n|^2}{\delta^2} \right)$;
- (3) $|\bar{L}L(\partial(H_i(f, \lambda, B)))| \leq K_0 \lambda^{-1/2} e^{\lambda \psi_i} \left(F(L_\tau, q, \delta)^{1/2} + \frac{|a_n|}{\delta} \right)$.

Proof. If $q \in \partial\Omega$, the inequality $|L\psi_i(q)| \leq K_0 \left(F(L_\tau, q, \delta)^{1/2} + \frac{|a_n|}{\delta} \right)$ follows immediately from Proposition 3.6 and the extremality of the basis (L_i) at (p, δ) . The general case for (1) follows.

Property (2) is obtained using the fact that, if (z) is the coordinates system associated to Φ and $L_i = \sum a_i^j \frac{\partial}{\partial z_j}$, then

$$|a_i^j| \lesssim F_i^{1/2}(p, \delta) F_j^{-1/2}(p, \delta)$$

for $q \in Q^c(p, \delta) \cap \partial\Omega$ (Proposition 3.6), and techniques similar to those used for (1).

Finally (3) is proved similarly, using the estimates of the coefficients of the brackets $[L_i \bar{L}_j]$ in $Q^c(p, \delta) \cap \partial\Omega$ (Proposition 3.6). \square

For $f = (f_1, \dots, f_{n-1})$, a $(n-1)$ -uplet of components of the weights F_i , let us denote by I the set of indexes i such that $f_i = \left| \frac{L_i^p \varphi_i}{\delta} \right|^{2/l(f_i)}$. Then we consider the function

$$H(f, \lambda, B) = \sum_{i \in I} H_i(f, \lambda, B).$$

The next Lemma gives some properties of the function $H(f, \lambda, B)$. To state it we need to introduce the following set:

For f a $(n-1)$ -uplet of components of the weights F_i and B' a positive number, we set

$$U_{B', f} = \{q \in Q^c(p, \delta) \text{ for which there exists } f' < f \text{ such that } f'(q) \text{ is } B' \text{ dominant}\}.$$

Lemma 5.2. *Let f be a $(n-1)$ -uplet of component, A, B and ε three positive fixed real numbers. Then there exist positive constants $\alpha_0, \lambda, A', B', A' > A, B' > B, \varepsilon' \leq \varepsilon$ and K_1 , depending only on A, B, ε, K and the data, such that, if the constant α of strong extremality is $\leq \alpha_0$, then the function $H(f, A, B, \varepsilon) = H(f, \lambda, B) = H$ satisfies, on $Q^c(p, \delta) \cap \Omega_{3\delta}$:*

- (1) $|H| \leq K_1$;
- (2) If $L = \sum_{i=1}^n a_i L_i^p = L_\tau + a_n N$, $a_i \in \mathbb{C}$, $\sum |a_i|^2 = 1$, then $|\langle \partial \bar{\partial} H; L, \bar{L} \rangle|(q) \leq A' \left(F(L_\tau, q, \delta) + \frac{|a_n|^2}{\delta^2} \right)$;
- (3) If $L = \sum_{i=1}^n a_i L_i^p = L_\tau + a_n N$, $a_i \in \mathbb{C}$, $\sum |a_i|^2 = 1$, $q \notin U_{B', f}$, $\chi'_{f, B}(q) = 1$ and $\chi_0(q) \geq \varepsilon$, then

$$\langle \partial \bar{\partial} H; L, \bar{L} \rangle(q) \geq AF(L_\tau, q, \delta) - K_2 \left(\sum_{i=1}^{n-1} |a_i|^2 \frac{|c_{ii}(q)|}{\delta} + \frac{|a_n|^2}{\delta^2} + 1 \right);$$

- (4) If $L = \sum_{i=1}^n a_i L_i^p = L_\tau + a_n N$, $a_i \in \mathbb{C}$, $\sum |a_i|^2 = 1$, the condition $\langle \partial \bar{\partial} H; L, \bar{L} \rangle(q) \leq - \left(F(L_\tau, q, \delta) + \frac{|a_n|^2}{\delta^2} \right)$ implies $q \in U_{B'}$ and $\chi_0(q) \geq \varepsilon'$.
- (5) For all lists $\mathcal{L} \in \mathcal{L}_3(\mathcal{B}(p, \delta) \cup \{N\})$, $|\mathcal{L}H(q)| \leq K_2 \left(\prod_{L \in \mathcal{L}} F(L, q, \delta) \right)^{1/2}$.

Proof. Recall that $H = \sum_{i \in I} H_i$, thus the properties are trivially satisfied if $I = \emptyset$ and we suppose $I \neq \emptyset$. The functions $|\psi_i|$ being bounded by 2 (see Proposition 3.6), (1) is satisfied with a constant K_1 depending only on λ and n .

Let $i \in I$. Then $\langle \partial \bar{\partial} H_i; L, \bar{L} \rangle = \bar{L} L H_i + [L, \bar{L}] (\partial H_i)$, and as

$$\bar{L} L H_i = \lambda^{-3/2} e^{\lambda \psi_i} \left[\left(\lambda^2 |L \psi_i|^2 + \lambda \bar{L} L \psi_i \right) \chi_{f, B} + \lambda (L \psi_i \bar{L} \chi_{f, B} + \bar{L} \psi_i L \chi_{f, B}) + \bar{L} L \chi_{f, B} \right],$$

Lemma 5.1 implies $\langle \partial \bar{\partial} H_i; L, \bar{L} \rangle(q) \geq \lambda^{-3/2} e^{\lambda \psi_i} \left(\lambda^2 |L \psi_i|^2 \chi_{f, B} - K'_0 \lambda F(L_\tau, q, \delta) + \frac{|a_n|^2}{\delta^2} + 1 \right)$ and thus shows the existence of a constant A' , depending only on the choice of λ, B, c, K and the data, satisfying (2).

Now, if for all $i \in I$, $|\lambda \psi_i| \leq 1$, then, for λ large enough, we have $\langle \partial \bar{\partial} H; L, \bar{L} \rangle \geq -F(L)$. Thus we suppose that there exists an $i \in I$ such that $|\lambda \psi_i(q)| = \frac{\lambda |\varphi_i(\pi(q))|}{\delta} F_i(p, \delta)^{(1-\tilde{l}(\varphi_i))/2} \geq 1$. Then there exists a constant $B' > B$, depending on λ , such that $\left| \frac{\varphi_i(\pi(q))}{\delta} \right|^{2/(\tilde{l}(\varphi_i)-1)} > \frac{4}{B'} F_i(p, \delta)$, and this implies that there exists a $(n-1)$ -uplet $f' < f$ which is B' -dominant at the point q . In other words, to each choice of λ we can associate B' such that the first conclusion in (4) is true. Moreover, λ, B and c being fixed, χ_1 being \mathcal{C}^∞ , there exists ε' , depending on λ, B, c and χ_1 , such that the hypothesis of (4) implies the second conclusion (i. e. $\chi_0(q) \geq \varepsilon'$).

Let us now show that we can choose λ (thus A', B', K_1 and ε' will be fixed) such that (3) is satisfied if α is small enough. Suppose then $\chi'_{f, B}(q) = 1$ and $\chi_0(q) > \varepsilon$. The hypothesis of strong extremality and the invariance of the $F_i(q)$ and the a_i^k in $B^c(p, \delta)$ (Propositions 3.6 and 3.10) give, if $\delta \leq \delta(\alpha)$,

$$|L \psi_i(q)|^2 \geq \frac{1}{4} \left| \sum_{j \leq i} a_j (L_j \psi_i)(q) \right|^2 - 4nC(K) \left(2\alpha \sum_{n-1 \geq j > i} |a_j|^2 F_j(p) + \frac{|a_n|^2}{\delta^2} \right),$$

and then, by extremality at p ,

$$|L \psi_i(q)|^2 \geq \frac{1}{4} \left| \sum_{j \leq i} a_j L_j \psi_i(q) \right|^2 - C_1(K) \left(\alpha^2 F(L_\tau, q, \delta) + \frac{|a_n|^2}{\delta^2} + 1 \right).$$

Now we make use of the following Lemma:

Lemma. Let β_i^j be complex numbers, $i = 1, 2, \dots, n-1$, $j \leq i$, verifying $|\beta_i^i| \geq c\alpha_i$ and $|\beta_i^j| \leq C\alpha_j$ for $j < i$. Then there exists a constant $W = W(c, C, n)$ such that $\sum_{i=1}^{n-1} \left| \sum_{j=1}^i \beta_i^j \right|^2 \geq W \sum_{i=1}^{n-1} (\alpha_i)^2$.

The above lemma implies, using the invariance of $F_i(q)$ and $F(L, q)$ in the ball and the extremality of the basis at p , that there exist constants W , K_3 and K_4 , depending on B , M , K and the data, such that:

$$\sum_{i \in I} |L\psi_i(q)|^2 + \sum_{i \notin I} \frac{|c_{ii}(q)|}{\delta} \geq \frac{W}{2K} F(L_\tau, q, \delta) - \alpha K_3 (F(L_\tau, q, \delta) + 1) - K_4 \frac{|a_n|^2}{\delta^2},$$

and thus, for $\alpha_0 = W/4KK_3$ (depending only on the data M , K , B , c and n),

$$\sum_{i \in I} |L\psi_i(q)|^2 + \sum_{i \notin I} \frac{|c_{ii}(q)|}{\delta} \geq W' F(L_\tau, q, \delta) - K_4 \left(\frac{|a_n|^2}{\delta^2} + 1 \right).$$

This finishes the proof of Lemma 5.2 for a choice of λ depending on A , ε , B , M , K and c , c depending itself only on M , K and the data, the property (5) being trivial. \square

Proof of Proposition 5.1. First, note that there exists a constant D , depending on M and n , such that, for $p \in W(p_0)$ and $\delta \geq \frac{1}{3} |\rho(p)|$, there exists a component f_i of $F_i(p, \delta)$ verifying $f_i(q) \geq \frac{1}{D} F_i(p, \delta)$ for all points $q \in B^c(p, \delta)$, $c \leq c_0$ (Proposition 3.6).

To define completely our function H , we have to define, for each $(n-1)$ -uplet of component $f \in \mathcal{H}$ (the set of $(n-1)$ -uplets of components of the weights $F_i(p, \delta)$), the constants A_f , B_f and ε_f from which $\lambda(f)$ is constructed. Let f^0 be the largest element of \mathcal{H} for the lexicographic order. Define $A_{f^0} = C4^{Mn+1}$, $B_{f^0} = D$ and $\varepsilon_{f^0} = 1$. Suppose we have constructed the constants A_f , B_f and ε_f for $f \geq f^1$. Consider the constants A'_{f^1} , B'_{f^1} and ε'_{f^1} obtained applying Lemma 5.2 for the constants A_{f^1} , B_{f^1} and ε_{f^1} , and define, for f^2 preceding f^1 , $A_{f^2} = 3C \sum_{f > f^2} A'_{f^1}$, $B_{f^2} = B'_{f^1}$ and $\varepsilon_{f^2} = \varepsilon'_{f^1}$. Thus $H = \sum_{f \in \mathcal{H}} H(f, A_f, B_f \varepsilon_f)$ is well defined.

For $q \in Q^c(p, \delta)$ define the following subsets of \mathcal{H} :

$$\begin{aligned} E_1(q) &= \left\{ f \in \mathcal{H} \text{ such that there exists } f' < f, \text{ such that } f'(q) \text{ is } B'_{f'}\text{-dominant and } \chi_0(q) \geq \varepsilon'_{f'} \right\}, \\ E_3(q) &= \left\{ f \in \mathcal{H} \text{ such that } \chi'_{f, B_f}(q) = 1 \text{ and } \chi_0(q) \geq \varepsilon_f \right\}, \\ E_2(q) &= \mathcal{H} \setminus \{E_1(q) \cup E_3(q)\}. \end{aligned}$$

Note that if $E_1(q)$ is not empty, and if f is its smallest element, then there exists $f' < f$ such that $f'(q)$ is $B'_{f'}$ dominant, that is $\chi'_{f', B_{f'}}(q) = 1$, and, as $\varepsilon_{f'} \leq \varepsilon'_f$, we also have $\chi_0(q) \geq \varepsilon_{f'}$ which means $f' \in E_3(q)$, f being the smallest element of $E_1(q)$.

Now suppose first that $q \in Q^{c/2}(p, \delta)$. Then, by definition of D , $E_3(q)$ is not empty, and, if $E_1(q)$ is also not empty there exists in $E_3(q)$ some strict minorant of $E_1(q)$. Then, by Lemma 5.2

$$\langle \partial \bar{\partial} H; L, \bar{L} \rangle(q) \geq \left(\sum_{f \in E_3(q)} A_f - \sum_{f \in E_1(q)} A'_f - \#E_2(q) \right) F(L_\tau, q, \delta) - \sum K_2(A_f, B_f, A'_f) \left(\sum_{i=1}^{n-1} |a_i|^2 \frac{c_{ii}(q)}{\delta} + \frac{|a_n|^2}{\delta^2} + 1 \right),$$

for α small enough, depending only on M , K and n ($\#E_2(f)$ denotes the number of elements of $E_2(f)$). Then, the preceding remark and the fact that $\#E_2(q) \leq 4^{Mn} \leq \frac{1}{4C} A_{f^0}$ imply

$$\langle \partial \bar{\partial} H; L, \bar{L} \rangle(q) \geq C4^{Mn} F(L_\tau, q, \delta) - \gamma_1 \left(\sum_{i=1}^{n-1} |a_i|^2 \frac{|c_{ii}(q)|}{\delta} + \frac{|a_n|^2}{\delta^2} + 1 \right).$$

Finally, if q is any point in $B^c(p, \delta)$ then $E_3(q)$ may be empty, but then $E_1(q)$ is also empty, and thus

$$\langle \partial \bar{\partial} H; L, \bar{L} \rangle(q) \geq -4^{Mn} F(L_\tau, q, \delta) - \gamma_1 \left(\sum_{i=1}^{n-1} |a_i|^2 \frac{|c_{ii}(q)|}{\delta} + \frac{|a_n|^2}{\delta^2} + 1 \right).$$

This finishes the proof of Proposition 5.1, property (4) being trivial. \square

5.2.3. Proof of Theorem 5.2. If P is a point of the boundary of D then, by the definition of D and Theorem 5.1, to prove that there exists a pluri-subharmonic function adapted to the structure of geometrically separated domain near P , we have only to consider the case where P is in the boundary of $\partial\Omega \cap \partial D$. Thus, with the notations introduced just before, we prove the following reformulation of Theorem 5.2:

Proposition 5.2. Let P be a point of the boundary of $\partial\Omega \cap \partial D$, and $V(P)$ the neighborhood considered in the previous Section. For all $K > 0$, there exist constants α_1 and δ_1 depending on K and the data such that if Ω is K -geometrically separated at $p_0 \in \partial\Omega$ and if the extremal bases of Ω are (K, α, p, δ) -strongly extremal with $\alpha \leq \alpha_1$, then, for $0 < \delta \leq \delta_1$, there exists a pluri-subharmonic function H_δ on the local domain D which is (δ, K') -adapted to $\mathcal{B}^{0,D}$ at P .

Proof. The proof is a modification of the proof of Theorem 5.1 and we will only indicate the differences. Here we have to consider both weights associated to the domains Ω and D , denoted F^Ω and F^D , which are constructed respectively with the defining functions ρ and $\rho + \varphi$.

We fix δ small enough and then we will omit the subscript δ in the notations of the vector fields. Consider, as in Section 5.2.1 the covering of $\partial D \cap V(P)$ by the pseudo-balls $B^{c/2}(q_k, \delta) \cap \partial\Omega$ (note that here P plays the role of p_0 in the previous Sections).

We denote $p_k = \pi(q_k)$ and fix k . Let $(L_i^k)_i = (L_i)_i$ be the δ -extremal basis (for D) at the point q_k . Let L_i^ρ be the vector field tangent to ρ associated to L_i (i. e. $L_i = L_i^\rho \circ \pi + (\beta \circ \pi)N_\Omega \circ \pi$). We saw (in Section 3.5.3) that the weights $F^D(L_i, \cdot, \delta)$ associated to the vector fields L_i are equivalent to

$$F^\Omega(L_i^\rho, \pi(\cdot), \delta) + \frac{\varphi'(|\cdot|)}{\delta} + \frac{\varphi''(|\cdot|^2)}{\delta} |\langle L_i^\rho(\pi(\cdot)), \cdot \rangle|^2.$$

Let $(L_i^{\Omega, k})_i = (L_i^\Omega)_i$ be the δ -extremal (for Ω) basis at p_k so that the vector fields L_i^ρ are linear combinations of the L_i^Ω . Let

$$I_k = \left\{ i \text{ such that } F^\Omega(L_i^\Omega, p_k, \delta) > \frac{\varphi'(|q_k|^2)}{\delta} \right\}.$$

We suppose I_k non empty. As the vector fields L_i^Ω are ordered so that their weights are decreasing, I_k is a segment of \mathbb{N} , $\{1, 2, \dots, n_k\}$. Then, we consider the n_k -uplets of components of the weights $F_\Omega(L_i^\Omega, p_k, \delta)$, $i \leq n_k$, $f = (f_1, \dots, f_{n_k})$ and the function

$$\chi_{f,B} = \prod_{i \leq n_k} \chi_B \left(\frac{f_i \circ \pi}{F_\Omega(L_i^\Omega, p_k, \delta)} \right) \chi_0,$$

where $\chi_0(q) = \chi_1 \left(\frac{F_{D,i}(q, \delta)}{c} \tilde{\Phi}_{q_k}(\pi_D(q)) \right)$, π_D being the projection onto ∂D associated to the real normal to D .

To obtain the good estimates of the derivatives of $\chi_{f,B}$ with respect to the vector fields L_i , we first estimate the derivatives of the functions $f_i \circ \pi$ at the point q_k :

Lemma 5.3. For $i \in I_k$, if $|f_i(p_k)| \geq \frac{1}{2B} F^\Omega(L_i^\Omega, p_k, \delta)$, for $\mathcal{L} \in \mathcal{L}_M(L_1, \dots, L_{n-1})$, we have

$$|\mathcal{L}(f_i \circ \pi)(q_k)| \lesssim_B F^\Omega(L_i^\Omega, p_k, \delta) F^D(q_k, \delta)^{\mathcal{L}/2}.$$

Proof. Let us consider the case $|\mathcal{L}| = 1$. As $L_j = L_j^\rho \circ \pi + (\beta_j \circ \pi)N_\Omega \circ \pi$, for $p = \pi(q)$,

$$|L_j(f \circ \pi)(q)| \leq |L_j^\rho(p)(f_i \circ \pi)(q)| + O(|\beta_j(p)|).$$

By (3.11), $|\beta_j(p)| \lesssim \varphi'(|p|^2) \lesssim \varphi'(|q|^2) + O(\varphi(|q|^2)) = O(\varphi'(|q|^2))$, thus

$$|\beta_j(p_k)| \lesssim F^{\Omega 1/2}(L_i^\Omega, p_k, \delta) F^{D 1/2}(L_j, q_k, \delta)$$

because $i \in I_k$. As L_j^ρ are tangent to ρ , $L_j^\rho(p)(f_i \circ \pi)(q) = L_j^\rho(p)(f_i)(p)$, and, as L_j^ρ are in the space spanned by the L_i^Ω , by Proposition 3.3, we have

$$\left| (L_j^\rho(p_k)(f_i))(p_k) \right| \lesssim F^\Omega(L_i^\Omega, p_k, \delta) F^\Omega(L_j^\rho, p_k, \delta)^{1/2},$$

and thus

$$\left| (L_j^\rho(p_k)(f_i))(q_k) \right| \lesssim F^\Omega(L_i^\Omega, p_k, \delta) F^\Omega(L_j^\rho, p_k, \delta)^{1/2} + \frac{\varphi(|q_k|^2)}{\delta}.$$

Derivatives of higher order are treated similarly. □

Corollary. Under the same hypothesis, for $q \in \mathcal{Q}_D^c(q_k, \delta) \cap D_{3\delta}$ and $\mathcal{L} \in \mathcal{L}_M(L_1, \dots, L_{n-1})$, we have

$$|\mathcal{L}(f_i \circ \pi)(q)| \lesssim_B F^\Omega(L_i^\Omega, p_k, \delta) F^D(q_k, \delta)^{\mathcal{L}/2}.$$

The derivatives of χ_0 being trivial, we deduce from (3.8) and Taylor's formula:

Lemma 5.4. For $i \in I_k$ and $q \in \mathcal{Q}_D^c(q_k, \delta) \cap D_{3\delta}$, for $\mathcal{L} \in \mathcal{L}_M(L_1, \dots, L_{n-1})$, we have

$$|\mathcal{L}\chi_{f,B}(q)| \lesssim_B F^D(q_k, \delta)^{\mathcal{L}/2}.$$

We now define the basic functions $H(f, \lambda, B)$ used here. Let $I = \left\{ i \leq n_k \text{ such that } f_i \neq \frac{c_{ii}}{\delta} \right\}$, and, for $i \in I$, if $f_i = \left| \frac{L_i \Phi_i}{\delta} \right|^{2/l(f_i)}$, we put

$$H_i(f, \lambda, B) = \lambda^{-3/2} e^{\lambda \psi_i} \chi_{f,B} \text{ where } \psi_i = \frac{\Phi_i}{\delta} \left| \frac{L_i \Phi_i}{\delta} \right|^{\frac{1}{l(\Phi_i)-1}},$$

and we define H by

$$H = H(f, \lambda, B) = \sum_{i \in I} H_i(f, \lambda, B).$$

If $L = \sum_{i=1}^n a_i L_i = L_\tau + a_n L_n$ ($\sum |a_i|^2 = 1$), as in the proof of Lemma 5.2, using the last Lemma we get

$$\langle \partial \bar{\partial} H_i; L, \bar{L} \rangle (q) \geq \lambda^{-3/2} e^{\lambda \psi_i} \left(\lambda^2 |L \psi_i|^2 \chi_{f,B} - K_0 \left(\lambda F^D(L_\tau, q_k, \delta) + \frac{|a_n|^2}{\delta} + 1 \right) \right),$$

for $q \in Q_D^c(q_k, \delta) \cap D_{3\delta}$. The estimate of $|L \psi_i|^2$ has now to be done more carefully.

The formula $L_\tau(q) = L_\tau^\rho(p) + \beta(p) N_\Omega$ gives

$$|L_\tau \psi_i|^2(q) \geq \frac{1}{4} |L_\tau^\rho(p) \psi_i|^2(q) - C \left(\frac{\varphi'(|q|^2)}{\delta} + \frac{\varphi(|q|^2) + \delta}{\delta} \right).$$

Then, decomposing L_τ^ρ on the δ -extremal basis at p_k ($(L_i^\Omega)_i$), $L_\tau^\rho = \sum_{j=1}^{n-1} b_j L_j^\Omega$, we obtain, using the strong extremality hypothesis,

$$|L_\tau^\rho(p) \psi_i|^2(q) \geq \frac{1}{4} \left| \sum_{j \leq i} b_j L_j^\Omega \psi_i \right|^2(q) - C \left(\alpha^2 F^\Omega(L_\tau^\rho, p_k, \delta) + \frac{\varphi(|q|^2) + \delta}{\delta} \right).$$

Using the same method, we sum all these inequality to get (writing $c_{ii}^\Omega = [L_i^\Omega, \bar{L}_i^\Omega](\partial \rho)$)

$$\begin{aligned} \sum_{i \in I} |L_\tau^\rho \psi_i|^2(q) &\geq \beta F^\Omega \left(\sum_{j=1}^{n_k} b_j L_j^\Omega, p_k, \delta \right) - C \left(\sum_{i \notin I, i \leq n_k} |b_i|^2 \frac{|c_{ii}^\Omega|}{\delta}(q) + \alpha^2 F^\Omega(L_\tau^\rho, p_k, \delta) + \frac{\varphi(|q|^2)}{\delta} + 1 \right) \\ &\geq \beta F^\Omega(L_\tau^\rho, p_k, \delta) - C \left(\sum_{i \notin I, i \leq n_k} |b_i|^2 \frac{|c_{ii}^\Omega|}{\delta}(q) + \alpha^2 F^\Omega(L_\tau^\rho, p_k, \delta) + \frac{\varphi'(|q|^2)}{\delta} + 1 \right), \end{aligned}$$

and, as $|L \psi_i|^2 \geq \frac{1}{4} |L_\tau \psi_i|^2 - C \frac{|a_n|^2}{\delta^2}$, we finally obtain

$$\sum_{i \in I} |L \psi_i|^2 \geq \beta F^\Omega(L_\tau^\rho, p_k, \delta) - C \left(\sum_{i \notin I, i \leq n_k} |b_i|^2 \frac{|c_{ii}^\Omega|}{\delta}(q) + \frac{|a_n|^2}{\delta} + \alpha^2 F^\Omega(L_\tau^\rho, p_k, \delta) + \frac{\varphi'(|q|^2)}{\delta} + 1 \right).$$

Then the proof is finished as in the previous Section using that, in $Q_D^c(q_k, \delta) \cap D_{3\delta}$, we have

$$\langle \partial \bar{\partial} e^{r/\delta}; L, \bar{L} \rangle (q) \geq \beta \left(\sum_{i=1}^{n-1} |b_i|^2 \frac{|c_{ii}^\Omega|}{\delta}(q) + \frac{|a_n|^2}{\delta^2} + \frac{\varphi'(|q|^2) + \varphi''(|q|^2)}{\delta} |\langle L_\tau^\rho(p), q \rangle|^2 \right) - K \left(\alpha F^\Omega(L_\tau^\rho, p_k, \delta) + 1 \right).$$

Indeed, a direct calculation gives

$$(5.2) \quad \langle \partial \bar{\partial} e^{r/\delta}; L, \bar{L} \rangle = e^{r/\delta} \left(\frac{2\Re(e^{a_n} \langle \partial \bar{\partial} r; L_\tau, \bar{N} \rangle)}{\delta} + \frac{|a_n|^2}{\delta^2} \right) (q) + \langle \partial \bar{\partial} e^{r/\delta}; L_\tau, \bar{L}_\tau \rangle (q).$$

For $q \in \{r \geq -3\delta\}$, the first term of (5.2) is $\geq \frac{1}{2e^3} \frac{|a_n|^2}{\delta^2} - K_0$. Let us look at the second term of (5.2).

$$\langle \partial \bar{\partial} e^{r/\delta}; L_\tau, \bar{L}_\tau \rangle = \frac{e^{r/\delta}}{\delta} \left(\langle \partial \bar{\partial} \rho; L_\tau, \bar{L}_\tau \rangle + \|L_\tau\|^2 \varphi'(|q|^2) + |\langle L_\tau^\rho(p), q \rangle|^2 \varphi''(|q|^2) \right).$$

But

$$\langle \partial \bar{\partial} \rho; L_\tau, \bar{L}_\tau \rangle (q) = \langle \partial \bar{\partial} \rho; L_\tau^\rho(p), \bar{L}_\tau^\rho(p) \rangle (q) + O \left(\varphi'(|q|^2) |\langle L_\tau^\rho(p), q \rangle|^2 \right)$$

and, first we can choose V small enough so that $\varphi'(|q|^2) |\langle L_\tau^\rho(p), q \rangle|^2 \ll \varphi'(|q|^2) + \varphi''(|q|^2) |\langle L_\tau^\rho(p), q \rangle|^2$, and secondly

$$\begin{aligned} \langle \partial \bar{\partial} \rho; L_\tau^\rho(p), \bar{L}_\tau^\rho(p) \rangle (q) &= \sum b_i b_j c_{ij}^\Omega(q) \\ &\geq \sum |b_i|^2 |c_{ii}^\Omega|(p) + O \left[\alpha \delta F^\Omega(L_\tau^\rho(p)) + \varphi(|q|^2) + \delta \right]. \end{aligned}$$

The proof of Proposition 5.2 is now complete. \square

6. APPLICATIONS TO COMPLEX ANALYSIS

6.1. Statements of the results for geometrically separated domains. In [CD06b] and [CD06a] we proved that the methods introduced, for the study of the Bergman and Szegő projection, by A. Nagel, J. P. Rosay, E. M. Stein and S. Wainger in \mathbb{C}^2 ([NRSW89]) and by J. McNeal and E. M. Stein for convex domains ([MS94, MS97]) can be adapted to pseudo-convex domains having an “adapted geometry”. The work made in the previous Sections shows that it is the case for completely geometrically separated domains and thus we have the following sharp estimates:

Theorem 6.1. *Suppose Ω is completely geometrically separated at $p_0 \in \partial\Omega$. Let $K_B(z, w)$ be the Bergman kernel of Ω . There exists a neighborhood $W(p_0)$ of p_0 such that:*

- (1) *For $p \in W(p_0) \cap \Omega$, $K_B(p, p) \simeq \prod_{i=1}^n F(L_i^{p, \delta(p)}, p, \delta_{\partial\Omega}(p))$, where $\delta_{\partial\Omega}(p)$ is the distance from p to $\partial\Omega$.*
- (2) *For $p_1, p_2 \in W(p_0) \cap \Omega$, for all integer N , there exists a constant C_N depending on Ω and N , such that for all lists $\mathcal{L}_{Z_1} = \{L_1^1, \dots, L_1^k\}$ (resp. $\mathcal{L}_{Z_2} = \{L_2^1, \dots, L_2^{k'}\}$) of length $k \leq N$ (resp. $k' \leq N$) with $L_1^j \in \mathcal{B}(\pi(p_1), \tau) \cup \{N\}$ (resp. $L_2^j \in \mathcal{B}(\pi(p_1), \tau) \cup \{N\})$, we have*

$$|\mathcal{L}_{Z_1} \overline{\mathcal{L}_{Z_2}} K_B(Z_1, Z_2)(p_1, p_2)| \leq C_N \prod_{i=1}^n F(L_i^{\pi(p_1), \tau}, \pi(p_1), \tau)^{1+l_i/2},$$

where $\tau = \delta_{\partial\Omega}(p_1) + \delta_{\partial\Omega}(p_2) + \gamma(\pi(p_1), \pi(p_2))$, $\gamma(\pi(p_1), \pi(p_2))$ is the pseudo-distance from $\pi(p_1)$ to $\pi(p_2)$ associated to the structure of homogeneous space and l_i is the number of times the vector fields $L_i^{\pi(p_1), \tau}$ or $\overline{L_i^{\pi(p_1), \tau}}$ appear in the union of the lists \mathcal{L}_{Z_1} and \mathcal{L}_{Z_2} .

Corollary. *Suppose Ω satisfies the hypothesis of Theorem 5.2. Let D be the local domain considered in Theorem 5.2. Then the Bergman kernel $K_D(z, w)$ of D satisfy all the estimates stated in the Theorem at any point of its boundary.*

Using the methods of Section 5 of [CD92] the following result on invariant metrics is easily proved:

Theorem 6.2. *Suppose Ω is completely geometrically separated at $p_0 \in \partial\Omega$. Let us denote by $B_\Omega(z, L)$ (resp. $C_\Omega(z, L)$, resp. $K_\Omega(z, L)$) the Bergman (resp. Caratheodory, resp. Kobayashi) metric of Ω at the point $z \in \Omega$. Then there exists a neighborhood $V(p_0)$ such that, for all vector fields $L \in E$ (E being the vector space spanned by the basis \mathcal{B}^0 (see Definition 4.1), $L = L_\tau + a_n N$, we have, for $q \in V(p_0) \cap \Omega$,*

$$B_\Omega(q, L) \simeq C_\Omega(q, L) \simeq K_\Omega(q, L) \simeq F(L_\tau, q, \delta(q)) + \frac{|a_n|}{\delta_{\partial\Omega}(q)},$$

where the constants in the equivalences depend only on the constant of geometric separation and the data.

Remark. The last point of Remark 4.2 and this Theorem show that the structure of homogeneous space we associate to a completely geometrically separated domain is essentially unique.

Theorem 6.3. *Suppose Ω is completely geometrically separated at every point of its boundary. Then the following results hold:*

- (1) *Let P_B be the Bergman projection of Ω . Then:*
 - (a) *for $1 < p < +\infty$ and $s \geq 0$, P_B maps continuously the Sobolev space $L_s^p(\Omega)$ into itself;*
 - (b) *for $0 < \alpha < +\infty$, P_B maps continuously the Lipschitz space $\Lambda_\alpha(\Omega)$ into itself;*
 - (c) *for $0 < \alpha < 1/M$, P_B maps continuously the Lipschitz space $\Lambda_\alpha(\Omega)$ into the non-isotropic Lipschitz space $\Gamma_\alpha(\Omega)$.*
- (2) *Let P_S be the Szegő projection of Ω . Then:*
 - (a) *for $1 < p < +\infty$ and $s \in \mathbb{N}$, P_S maps continuously the Sobolev space $L_s^p(\partial\Omega)$ into itself;*
 - (b) *for $0 < \alpha < +\infty$, P_S maps continuously the Lipschitz space $\Lambda_\alpha(\partial\Omega)$ into itself;*
 - (c) *for $0 < \alpha < 1/M$, P_S maps continuously the Lipschitz space $\Lambda_\alpha(\partial\Omega)$ into the non-isotropic Lipschitz space $\Gamma_\alpha(\partial\Omega)$.*

Note.

- (1) Statements (1) (c) and (2) (c) can be extended to all $\alpha > 0$ with convenient definitions of the spaces $\Gamma_\alpha(\Omega)$ and $\Gamma_\alpha(\partial\Omega)$.
- (2) In view of Example 5.1, the previous theorem applies in particular for all lineally convex domains of finite type.

Corollary. *Suppose Ω satisfy the hypothesis of Theorem 5.2. Let D be the local domain considered in Theorem 5.2. Then all the results stated for Ω in the previous Theorem are valid for D .*

Using an idea of M. Machedon [Mac88] we deduce local estimates for the Szegő projection:

Theorem 6.4. *Suppose Ω satisfies the hypothesis of Theorem 5.2. Let P_S be the Szegő projection of Ω . Then if f is a $L^2(\partial\Omega)$ function which is locally near p_0 in the Sobolev space L_s^p , $1 < p < +\infty$ and $s \in \mathbb{N}$, (resp. in the Lipschitz space Λ_α , $0 < \alpha < 1/M$) then its projection $P_S(f)$ is locally near p_0 in L_s^p (resp. in the non-isotropic Lipschitz space Γ_α). In particular this applies if the Levi form of Ω is locally diagonalizable at p_0 .*

Proof. if $f \in L^2(\Omega)$ and if $\chi \in \mathcal{C}^\infty(\partial\Omega)$ has compact support in a sufficiently small neighborhood of p_0 and $\chi = 1$ in a neighborhood of p_0 , then the subelliptic estimates for \square_b and Kohn's theory ([Koh85, KN65]) implies $P_S((1-\chi)f)$ is \mathcal{C}^∞ near p_0 , and, denoting P_S^D the Szegő projection of D , $(P_S - P_S^D)(\chi f)$ is \mathcal{C}^∞ in a neighborhood of p_0 (see also [Kan90]); the result follows thus the previous Corollary. \square

6.2. A guide for the proofs of the results of Section 6.1. Let U be a neighborhood of $\partial\Omega$ where we can define a projection π onto $\partial\Omega$ using the integral curve of the real normal to ρ . We will always suppose that $V(p_0) \subset U$.

The two notions of “weak homogeneous space” and “adapted pluri-subharmonic function” plays a crucial role in [CD06b, CD06a]:

Definition 6.1. We say that the domain Ω satisfies the hypothesis of “weak homogeneous space” at a boundary point p_0 of finite type τ if there exist two neighborhoods $V(p_0)$ and $W(p_0) \Subset V(p_0)$ and a constant K such that:

- (1) There exists $\delta_0 > 0$ such that, for every $p \in W(p_0)$, $\forall \delta \in [-\frac{1}{3}\rho(p), \delta_0]$, there exists a basis of vector fields tangent to ρ in $V(p_0)$, $\mathcal{B}(p, \delta)$, for which there exists a K -adapted coordinate system
- (2) There exists two constants C and c_0 , depending on K and τ , such that, for $c \leq c_0$, the sets $B^c(\mathcal{B}(p, \delta), p, \delta)$ (associated to the coordinate system), $B_{\mathcal{C}}^c(\mathcal{B}(p, \delta), p, \delta)$ and $B_{\text{exp}}^c(\mathcal{B}(p, \delta), p, \delta)$ satisfy, for all $p \in W(p_0) \cap \bar{\Omega}$ and all $\delta \in [-\frac{1}{3}\rho(p), \delta_0]$, the following conditions:
 - (a) for $q \in B_0^c(p, \delta)$, $B_0^c(\mathcal{B}(q, \delta), q, \delta) \subset B_1^c(\mathcal{B}(p, \delta), p, C\delta)$, where B_0^c and B_1^c denotes one of the sets B^c , $B_{\mathcal{C}}^c$ or B_{exp}^c .
 - (b) $\text{Vol}(B_0^c(\mathcal{B}(p, 2\delta), p, 2\delta)) \leq C \text{Vol}(B_0^c(\mathcal{B}(p, \delta), p, \delta))$.

Note that, in this Definition the weights F_i are defined with $M = M'(\tau)$.

Definition 6.2. Let $\mathcal{B} = \{L_1, \dots, L_{n-1}\}$ be a basis of vector fields tangent to ρ in a neighborhood $V(p_0)$ of a boundary point p_0 and $0 < \delta \leq \delta_0$. We say that a pluri-subharmonic function $H \in \text{PSH}(\Omega)$ is (p_0, K, c, δ) -adapted to this basis \mathcal{B} if the following properties are satisfied:

$|H| \leq 1$ in Ω , and, for all point $p \in W(p_0) \cap \bar{\Omega}$, $\rho(p) \geq -3\delta$, the two following inequalities are verified for points $q \in B_{\mathcal{C}}^c(\mathcal{B}, p, \delta) \cap \Omega$:

- (1) For all $L = \sum_{i=1}^n a_i L_i$, $a_i \in \mathbb{C}$,

$$\langle \partial \bar{\partial} H, L, \bar{L} \rangle \geq \frac{1}{K} \sum_{i=1}^n |a_i|^2 F(L_i, p, \delta).$$

- (2) For $\mathcal{L} \in \mathcal{L}_3(\mathcal{B} \cup \{N\})$,

$$|\mathcal{L}H| \leq K \prod_{L \in \mathcal{L}} F(L, p, \delta)^{1/2}.$$

Note that this Definition depends on the values of the vector fields L_i^p at points q in Ω . But, in the situation of the applications below (i.e. with a finite type hypothesis) it can be shown that it depends only (up to uniform constants) on the restriction of the basis on $\partial\Omega$.

The following Proposition follows from the work in [CD06b, CD06a]:

Proposition 6.1. *Let Ω be a bounded pseudo-convex domain and p_0 be a boundary point of finite type (resp. a bounded pseudo-convex domain of finite type). Then, if Ω satisfies the hypothesis of “weak homogeneous space” at p_0 (resp. at every point of its boundary) and if there exists a pluri-subharmonic function \mathcal{H}_δ adapted to $\mathcal{B}(p, \delta)$ for all $p \in W(p_0) \cap \bar{\Omega}$ and all $\delta \in [-\frac{1}{3}, \delta_0]$ (resp. if this property holds at every point p_0 of $\partial\Omega$) then the conclusions of Theorem 6.1 (resp. Theorem 6.3) hold.*

To prove Theorems 6.1 and 6.3 it suffices then to use the properties of extremal bases and to note the two following facts:

- (1) The existence of extremal bases and adapted coordinate systems for points of $\partial\Omega \cap W(p_0)$ allows us to define bases and coordinate systems for points inside Ω (see Remark 4.1);
- (2) if $p_1 \in W(p_0) \cap \Omega$, $p = \pi(p_1)$, the sets $\widetilde{B}_0^c(\mathcal{B}(p, \delta), p_1, \delta)$, $-\frac{1}{3}\rho(p_1) < \delta \leq \delta_0$, defined by $q \in \widetilde{B}_0^c(\mathcal{B}(p, \delta), p_1, \delta)$ if $\pi(q) \in B_0^c(\mathcal{B}(p, \delta), p, \delta)$ and $|\rho(q) - \rho(p_1)| < c\delta$ induce a structure of “weak homogeneous space”.

6.3. Main articulations of the proof of Proposition 6.1. In Section 2 of [CD06b] we showed that if the Levi form is locally diagonalizable then the local hypothesis of the Proposition is satisfied, and in [CD06a, CD06b], even if the statements are given in the case of a locally diagonalizable Levi form, the proofs of the estimates on the Bergman and Szegő projections are done only using the hypothesis of the Proposition. We just give here the main articulations of the proofs:

- The Bergman kernel estimates on the diagonal is done using Theorem 6.1 of [Cat89] and the change of coordinates Φ_p adapted to the basis $\mathcal{B}(p, \delta(p))$.
- The estimates on the derivatives of the Bergman kernel outside the diagonal follow the methods developed by A. Nagel, J. P. Rosay, E. M. Stein and S. Wainger [NRSW89] and J. Mc Neal [McN89] for the pseudo-convex domains of finite type in \mathbb{C}^2 , and used for some generalizations (see the introduction) in particular by J. Mc Neal [McN94] in the case of convex domains. It consists to obtain uniform local estimates for the Neumann operator \mathcal{N} and then

to apply the ideas developed by N. Kerzman [Ker72] in the study of the strictly pseudo-convex case. This requires scaling.

The starting point is to write the Bergman kernel K_B^Ω using the Bergman projection. More precisely, if ψ_ζ is a radial function centered at ζ with compact support in Ω and of integral 1, and P_B^Ω is the Bergman projection of Ω , then $D^\mu \bar{D}^\nu K_B^\Omega(w, \zeta) = D_w^\mu P_B^\Omega(\bar{D}_\zeta^\nu \psi_\zeta)(w)$. Then, P_B^Ω being related to the $\bar{\partial}$ -Neumann problem by the formula $P_B^\Omega = \text{Id} - \vartheta \mathcal{N} \bar{\partial}$, where ϑ is the formal adjoint to $\bar{\partial}$ and \mathcal{N} the inverse operator of $\bar{\partial} \bar{\partial}^* + \bar{\partial}^* \bar{\partial}$, the estimates on P_B^Ω are obtained via estimates on \mathcal{N} . To obtain these estimates, we use the theory developed by J. J. Kohn and L. Nirenberg [KN65] which gives local Sobolev estimates for \mathcal{N} if there exists a local sub-elliptic estimates for the $\bar{\partial}$ -Neumann problem and the famous work of D. Catlin ([Cat87]), where it is proved that the existence of an adapted pluri-subharmonic function implies the existence of a sub-elliptic estimates for the $\bar{\partial}$ -Neumann operator.

The study of the Bergman kernel is not directly done in Ω but in $\Phi_p(\Omega)$, where Φ_p is a coordinate system adapted to the basis $\mathcal{B}(p, \delta_{\partial\Omega}(p) + \delta_{\partial\Omega}(q) + \gamma(\pi(p), \pi(q)))$, where γ is the pseudo-distance on $\partial\Omega$. One difficulty is to see that all the constants appearing in the estimates and all the domains where the estimates are done are uniformly controlled.

- The estimates for the Bergman and Szegö projectors are obtained adapting the methods developed by J. Mc Neal and E. M. Stein in [MS94, MS97] (and also [NRSW89]), related, in particular, to the theory of non isotropic smoothing operators, to non convex domains.

Remark. The results on the Szegö projection are thus obtained adapting the theory of NIS operators to our settings. The Λ_α estimates, for example, for the domains considered by M. Derridj in [Der99] can also be obtained using the estimate for \square_b of Derridj's paper, the estimate on the Bergman projection derived from the fact that these domains are completely geometrically separated and the results on the comparison of the Bergman and Szegö projection obtained by K. D. Koenig in [Koe07].

7. EXAMPLES AND ADDITIONAL REMARKS

7.1. The lineally convex case. In this Section we show, with some details the statements made on lineally convex domains in Example 3.1, Example 4.1 and Example 5.1.

Suppose $\Omega = \{\rho < 0\}$ is lineally convex near $p_0 \in \partial\Omega$, a point of finite type, and W is a small neighborhood of p_0 . $(Z_i)_i$ is a coordinate system centered at p_0 such that Z_n is the complex normal to $\partial\Omega$ at p_0 , and $\frac{\partial\rho}{\partial Z_n} \simeq 1$ in W .

We begin with the statement in Example 3.1 (1). Let $p \in \partial\Omega \cap W$ and $\delta > 0$. Let $(z_i)_i$ be the δ -extremal basis (considered as a coordinate system) at p defined by M. Conrad in [Con02] (the main results concerning this basis are summarized in [DF06]), which is centered at p . To be coherent with our previous notations, we suppose that the complex normal to $\partial\Omega$ at p is z_n (in M. Conrad paper this normal is z_1).

To each vector $v = (a_1, \dots, a_{n-1}, 0) \in \mathbb{C}^n$ we associate the $(1, 0)$ -vector field, tangent to ρ ,

$$(7.1) \quad L_v = \sum_{i=1}^{n-1} a_i \frac{\partial}{\partial z_i} + \beta_v \frac{\partial}{\partial Z_n} := V + \beta_v \frac{\partial}{\partial Z_n}$$

(thus $\beta_v = -V(\rho) \left(\frac{\partial\rho}{\partial Z_n} \right)^{-1}$).

If $v_i = (\delta_k^i)_{1 \leq k \leq n}$, $1 \leq i \leq n-1$, we denote $L_i = L_{v_i} = \frac{\partial}{\partial z_i} + \beta_i \frac{\partial}{\partial Z_n}$. Note that the vector fields L_i depend on p and δ ($L_i = L_i(p, \delta)$) and are the vector fields of a basis of the complex tangent space to ρ in W .

Proposition 7.1. *There exists a constant K such that, for all $p \in \partial\Omega \cap W$ and all $\delta \leq \delta_0$, δ_0 small enough, the basis $(L_i(p, \delta))_i$ is (K, δ) -extremal at p .*

Proof. p and δ being fixed, we drop them in the notations. First we express the weights $F(L_v, p, \delta)$ in terms of the vector field V of (7.1).

Lemma. *Let \mathcal{L} be a list composed of αL_v and $\beta \bar{L}_v$, $\|v\| \leq 1$. Then*

$$\mathcal{L}(\partial\rho) = 2V^\alpha \bar{V}^\beta(\rho) + \sum_{\alpha' + \beta' < \alpha + \beta} *V^{\alpha'} \bar{V}^{\beta'}(\rho)$$

where $*$ are functions of $\mathcal{C}^{2m - (\alpha + \beta)}$ norm uniformly bounded in p and δ .

Proof. Look first at $c_{v\bar{v}} = 2[L_v, \bar{L}_v](\partial\rho)$: $c_{v\bar{v}} = -2\bar{L}_v(\beta_v) \frac{\partial\rho}{\partial Z_n} = 2\bar{L}_v V(\rho) + *V(\rho) = 2\bar{V}V(\rho) + *V(\rho)$. The Lemma is then proved by induction. \square

Corollary. $F(L_v, p, \delta) \simeq \sum |a_i|^2 F(L_i, p, \delta)$ uniformly in p and δ .

Proof. It suffices to prove this formula when $\|v\| = 1$. By the Lemma

$$\begin{aligned} F(L_v, p, \delta) &\simeq \sum_{2 \leq \alpha + \beta \leq m} \left| \frac{V^\alpha \bar{V}^\beta(\rho)}{\delta} \right|^{\frac{2}{\alpha + \beta}} \\ &= \sum_{2 \leq \alpha + \beta \leq m} \left| \frac{\frac{\partial^{\alpha + \beta}}{\partial \lambda^\alpha \partial \bar{\lambda}^\beta}(\rho)(p + \lambda v)|_{\lambda=0}}{\delta} \right|^{\frac{2}{\alpha + \beta}} \\ &\simeq \left(\frac{2}{\tau(p, v, \delta)} \right)^2, \end{aligned}$$

where $\tau(p, v, \delta)$ is Conrad's notation.

Using properties (iii) and (iv) of Proposition 3.1 of [DF06] we get

$$F(L_v, p, \delta) \simeq \sum_2^m \frac{|a_i|^2}{\tau(p, v_i, \delta)^2}.$$

As all constants are uniform in p and δ the Corollary is proved. \square

To finish the proof of Proposition 7.1 we have to prove property EB₂ of Definition 3.1. For example, let us look at the bracket $[L_i, \bar{L}_j]$:

$$[L_i, \bar{L}_j] = \left(-\frac{\partial}{\partial \bar{z}_j} + \bar{\beta}_j \frac{\partial}{\partial \bar{Z}_n} \right) (\beta_i) \frac{\partial}{\partial Z_n} + \left(\frac{\partial}{\partial z_i} + \beta_i \frac{\partial}{\partial Z_n} \right) (\bar{\beta}_i) \frac{\partial}{\partial \bar{Z}_n} = a \frac{\partial}{\partial Z_n} + b \frac{\partial}{\partial \bar{Z}_n}.$$

Let $\mathcal{L} \in \mathcal{L}_M(L_1, \dots, L_{n-1})$. As, for all k , $F_k^{-1/2} \geq \delta$ and $\frac{\partial}{\partial z_n} = \sum \alpha_i \frac{\partial}{\partial z_i}$ with α_i uniformly bounded in \mathcal{C}^M norm, it is enough to show that

$$(|\mathcal{L}a| + |\mathcal{L}b|)(p) \lesssim \delta F^{\alpha/2}(p, \delta) F_i^{1/2}(p, \delta) F_j^{1/2}(p, \delta).$$

If $|\mathcal{L}| = 0$, $a(p) = \frac{\partial^2 \rho}{\partial z_i \partial \bar{z}_j}(0) \left(\frac{\partial \rho}{\partial Z_n} \right)^{-1}(p)$ and, if $|\mathcal{L}| = 1$, $\bar{L}_k a(p) = \frac{\partial^3 \rho}{\partial z_k \partial z_i \partial \bar{z}_j}(0) \left(\frac{\partial \rho}{\partial Z_n} \right)^{-1}(p) + * \frac{\partial^2 \rho}{\partial z_k \partial \bar{z}_j}$. Thus, in those cases, the result follows from Lemma 3.2 of M. Conrad's paper [Con02] which states

$$(7.2) \quad \left| \frac{\partial^{\alpha + \beta} \rho}{\partial z^\alpha \partial \bar{z}^\beta}(p) \right| \lesssim \delta \prod \left(\frac{1}{\tau(p, v_i, \delta)} \right)^{\alpha + \beta_i} \simeq \delta F(p, \delta)^{(\alpha + \beta)/2},$$

the last equivalence resulting of the proof of the previous Corollary. The case of a general \mathcal{L} is easily done similarly. \square

Now let us prove the statement made in Example 5.1 (1). The construction of the adapted plurisubharmonic function is inspired by the McNeal's construction for convex domains, using support function, written in [McN02]. We use the support function for lineally convex domains constructed by J. E. Fornæss and K. Diederich in [DF03]. The right behavior in the normal direction is obtained, as in Section 5.2, adding the functions $Ke^{\rho/\delta}$ and $K|z|^2$.

Consider the support function constructed by K. Diederich and J. E. Fornæss in [DF03] at the point p :

$$\begin{aligned} S_p(z_1, \dots, z_n) &= -\varepsilon \sum_{j=2}^{2m} M^{2j} \sigma_j \sum_{|\alpha|=j, \alpha_n=0} \frac{1}{\alpha!} \frac{\partial^j \rho(0)}{\partial z^\alpha} z^\alpha \\ &\quad + z_n \left(\frac{1}{1 - A_p(z)} \right) + K_0 \left(\frac{z_n}{1 - A_p(z)} \right)^2, \end{aligned}$$

where A_p is a \mathcal{C}^∞ function, uniformly bounded (in p), such that $A_p(0) = 0$. Shrinking $W(p_0)$ if necessary, S_p is uniformly bounded on $W(p_0)$.

Then there exists a constant $M_0 (> 8n$ and independent of p and δ) such that, if $S = \frac{M_0 \Re(S_p)}{\delta}$, we have:

- (1) $\Re(S) \leq 0$;
- (2) $\Re(S(z)) \leq -n$, if there exists $i < n$ such that $|z_i| \geq F_i(p, \delta)^{-1/2}$;
- (3) $-1/4 \leq \Re(S(z))$ if $z \in cP(p, \delta) = \{z \text{ such that } |z_i| \leq cF_i(p, \delta)^{-1/2}, i = 1, \dots, n\}$.

Let F be the function defined by $F(z) = \sum_{i=1}^{n-1} F_i(p, \delta) |z_i|^2$, and χ be the convex function such that $\chi \equiv 0$ on $] -\infty, -1[$ and $\chi(x) = e^{x-1} - 1$ on $] -1, +\infty[$. Define then

$$H_1 = \chi \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right).$$

Clearly $H_1 \equiv 0$ on a neighborhood of the boundary of $P(p, \delta)$. Thus we denote by H the function equal to H_1 in $P(p, \delta)$ and 0 outside. Then:

- (1) $\text{Supp}(H) \subset P(p, \delta)$;
- (2) $|H| \leq C_0$;

(3) On $P(p, \delta)$ we have

$$\left\langle \partial \bar{\partial} H; \sum_{i=1}^{n-1} a_i \frac{\partial}{\partial z_i}, \overline{\sum_{i=1}^{n-1} a_i \frac{\partial}{\partial z_i}} \right\rangle \geq \chi' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \sum_{i=1}^{n-1} |a_i|^2 F_i(p, \delta).$$

We now estimate $\langle \partial \bar{\partial} H; L, \bar{L} \rangle$ in $P(p, \delta)$ for a vector field $L = \sum_{i=1}^n b_i L_i^0$ where $L_i^0 = \frac{\partial}{\partial z_i} - \beta_i^0 \frac{\partial}{\partial z_n}$, for $i < n$, and $L_n^0 = N$ the complex normal vector (recall that the extremal bases are linear combinations of these L_i^0). Denote $L = L_\tau + b_n N$, so that L_τ is tangent to ρ . Then

$$(7.3) \quad \langle \partial \bar{\partial} H; L, \bar{L} \rangle = \langle \partial \bar{\partial} H; L_\tau, \overline{L_\tau} \rangle + 2\Re \left(\overline{b_n} \langle \partial \bar{\partial} H; L_\tau, \bar{N} \rangle \right) + |b_n|^2 \langle \partial \bar{\partial} H; N, \bar{N} \rangle.$$

The last term of the right hand side of this equality is $\geq -O \left(|b_n|^2 \chi' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \frac{1}{\delta^2} \right)$, and, if (L_i) is the δ -extremal basis at p and $L_\tau = \sum_{i=1}^{n-1} a_i L_i$, we have

$$\langle \partial \bar{\partial} H; L_\tau, \bar{N} \rangle = \left\langle \partial \bar{\partial} H; \sum_{i=1}^{n-1} a_i \frac{\partial}{\partial z_i}, \bar{N} \right\rangle + \left\langle \partial \bar{\partial} H; \left(\sum a_i \beta_i \right) \frac{\partial}{\partial z_n}, \bar{N} \right\rangle.$$

Using (7.2) the first term of the right hand side of this equality is $O \left(\frac{1}{\delta} \chi' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \right)$, and, using also the fact that $\beta_i(p) = 0$ implies $\beta_i = O \left(\delta F_i(p, \delta)^{1/2} \right)$, the second term is $O \left(\frac{1}{\delta} \chi' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \sum |a_i| F_i(p, \delta)^{1/2} \right)$. Notice that, by extremality, $\sum |a_i| F_i(p, \delta)^{1/2} \simeq F(L_\tau, p, \delta)^{1/2}$, thus, there exists a constant K_1 such that

$$2\Re \left(\overline{b_n} \langle \partial \bar{\partial} H; L_\tau, \bar{N} \rangle \right) + |b_n|^2 \langle \partial \bar{\partial} H; N, \bar{N} \rangle \geq -K_1 \chi' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \left(\frac{|b_n|^2}{\delta^2} + \frac{|b_n|}{\delta} F(L_\tau, p, \delta)^{1/2} \right).$$

Let us now look at the first term of the right hand side of (7.3):

$$\begin{aligned} \langle \partial \bar{\partial} H; L_\tau, \overline{L_\tau} \rangle &= \chi' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \left\langle \partial \bar{\partial} \left(F + S - n \frac{|z_n|^2}{\delta^2} \right); L_\tau, \overline{L_\tau} \right\rangle + \\ &\quad + \chi'' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \left| L_\tau \left(F + S - n \frac{|z_n|^2}{\delta^2} \right) \right|^2 \\ &= A + B. \end{aligned}$$

Shrinking $W(p_0)$ if necessary, we have

$$\begin{aligned} A &\geq \chi' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \left[\frac{1}{2} \sum_{i=1}^{n-1} |a_i|^2 F_i(p, \delta) - \frac{n}{\delta^2} \left| \langle \partial \bar{\partial} |z_n|^2; L_\tau, \overline{L_\tau} \rangle \right| \right] \\ &\geq \chi' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \left[\frac{1}{2} \sum_{i=1}^{n-1} |a_i|^2 F_i(p, \delta) - \frac{2n}{\delta^2} \left| \sum_{i=1}^{n-1} a_i \beta_i \right|^2 \right]. \end{aligned}$$

To estimate B , write

$$L_\tau \left(F + S - n \frac{|z_n|^2}{\delta^2} \right) = \sum_{i=1}^{n-1} a_i \frac{\partial}{\partial z_i} \left(F + S - n \frac{|z_n|^2}{\delta^2} \right) + \left(\sum_{i=1}^{n-1} a_i \beta_i \right) \frac{\partial}{\partial z_n} \left(F + S - n \frac{|z_n|^2}{\delta^2} \right).$$

Then the first term of the right hand side of this equality is $O \left(F(L_\tau, p, \delta)^{1/2} \right)$ by extremality (use (7.2)), and

$$\frac{\partial}{\partial z_n} \left(F + S - n \frac{|z_n|^2}{\delta^2} \right) = O \left(F(L_\tau, p, \delta)^{1/2} \right) + \frac{M_0}{\delta} \frac{\partial}{\partial z_n} \left(\frac{z_n}{1 - A_p(z)} \right) + O(1) - \frac{n}{\delta^2} \frac{\partial}{\partial z_n} (z_n \bar{z}_n).$$

But, if $W(p_0)$ is small enough, $\left| \frac{\partial}{\partial z_n} \left(\frac{z_n}{1 - A_p(z)} \right) \right| \in [1/2, 3/2]$, and, in $P(p, \delta)$, $\frac{1}{\delta^2} \left| \frac{\partial}{\partial z_n} (z_n \bar{z}_n) \right| \leq \frac{2n}{\delta}$.

Thus, as $\chi'' = \chi'$, for δ small enough, we have, by the choose of M_0 ,

$$\langle \partial \bar{\partial} H; L_\tau, \overline{L_\tau} \rangle \geq \chi' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \sum_{i=1}^{n-1} |a_i|^2 F_i(p, \delta).$$

Using again the extremality of the basis (L_i) , we conclude that

$$\langle \partial \bar{\partial} H; L, \bar{L} \rangle \geq \alpha \chi' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \left[F(L_\tau, p, \delta) - K_2 \frac{|b_n|^2}{\delta^2} + K_2 \frac{|b_n|}{\delta} F(L_\tau, p, \delta)^{1/2} \right],$$

and, using Cauchy-Schwarz inequality, we get

$$\langle \partial \bar{\partial} H; L, \bar{L} \rangle \geq \beta \chi' \left(F + S - \frac{n}{\delta^2} |z_n|^2 \right) \left[F(L_\tau, p, \delta) - K_3 \frac{|b_n|^2}{\delta^2} \right].$$

In particular, on $cP(p, \delta)$, we have

$$\langle \partial \bar{\partial} H; L, \bar{L} \rangle \geq \gamma F(L_\tau, p, \delta) - K \frac{|b_n|^2}{\delta^2},$$

and $\langle \partial \bar{\partial} H; L, \bar{L} \rangle \geq -K \frac{|b_n|^2}{\delta^2}$ on $P(p, \delta)$.

If we note that choosing c sufficiently small we have $F(L_\tau, p, \delta) \simeq F(L_\tau, q, \delta)$, we get:

Proposition 7.2. *There exist two constants γ and K depending only on the data such that, if $L = L_\tau + b_n N = \sum_{i=1}^{n-1} b_i L_i^0 + b_n N$, we have $\langle \partial \bar{\partial} H; L, \bar{L} \rangle \geq -K \frac{|b_n|^2}{\delta^2}$, and, if $q \in cP(p, \delta)$, we have $\langle \partial \bar{\partial} H; L, \bar{L} \rangle(q) \geq \gamma F(L_\tau, q, \delta) - K \frac{|b_n|^2}{\delta^2}$.*

To finish the construction of the plurisubharmonic function adapted to the structure of Ω at p , as in the proof of Theorem 5.1, we have to add functions of the precedent type to get a local function. Thus, we cover $\partial\Omega \cap W(p_0)$ with a minimal system of polydiscs $\frac{c}{2}P(p_k, \delta)$, $p_k \in \partial\Omega \cap W(p_0)$ and, then, there exists an integer J , independent of δ , such that every point of Ω belongs to at most J polydiscs $P(p_k, \delta)$. Indeed, there exists a constant C such that

$$P\left(p, \frac{c}{C}\delta\right) \subset \frac{c}{2}P(p, \delta) \subset P(p, cC\delta)$$

and the polydiscs $P(p, \delta)$ are associated to a structure of homogeneous space.

Consider $H = \sum H_{p_k}$ where the function H_{p_k} is the one considered in the previous Proposition relatively to the point $p = p_k$ (notice that $\|H\| \leq JC_0$). Then, shrinking eventually $W(p_0)$ and choosing ρ equivalent to the distance to the boundary with a constant close to 1, for all point $q \in W(p_0) \cap \{0 > \rho > -\frac{c}{2}\delta\}$ there exists k_0 such that $q \in P(p_{k_0}, \delta)$ and the set $E(q)$ of index k so that $q \in cP(p_k, \delta)$ has at most J elements and we have

$$\langle \partial \bar{\partial} H; L, \bar{L} \rangle(q) \geq \gamma F(L_\tau, q, \delta) - KJ \frac{|b_n|^2}{\delta^2}.$$

Moreover, without conditions on q , we have

$$\langle \partial \bar{\partial} H; L, \bar{L} \rangle(q) \geq -KJ \frac{|b_n|^2}{\delta^2},$$

and $\partial \bar{\partial} H(q) = 0$ if $\rho(q) < -2\delta$.

We now evaluate $\langle \partial \bar{\partial} e^{\rho/\delta}; L, \bar{L} \rangle$ in $W(p_0)$:

$$\langle \partial \bar{\partial} e^{\rho/\delta}; L, \bar{L} \rangle \geq e^{\rho/\delta} \left[\frac{1}{\delta} \left(\frac{1}{2} \sum_{i,j=1}^{n-1} b_i \bar{b}_j c_{ij}^0 + \Re \sum_{i=1}^{n-1} b_i \bar{b}_n \langle \partial \bar{\partial} \rho; L_i^0, \bar{N} \rangle \right) + \frac{|b_n|^2}{\delta^2} \right]$$

c_{ij}^0 being the coefficient of the Levi form in the direction (L_i^0, \bar{L}_j^0) . As the level set of ρ are pseudo-convex (in $W(p_0)$), we get

$$\langle \partial \bar{\partial} e^{\rho/\delta}; L, \bar{L} \rangle \geq e^{\rho/\delta} \left(\frac{1}{2} \frac{|b_n|^2}{\delta^2} - K_1 \right).$$

Consider now $\tilde{H} = H + K_1 e^{\rho/\delta} + K_2 |z|^2$, for K_1 and K_2 large enough (independent of δ). Then \tilde{H} is plurisubharmonic on $\Omega \cap W(p_0)$, uniformly bounded (with respect to δ) and satisfies, on $W(p_0) \cap \{0 > \rho > -\frac{c}{2}\delta\}$

$$\langle \partial \bar{\partial} \tilde{H}; L, \bar{L} \rangle \geq \gamma F(L_\tau, \cdot, \delta) + \frac{|b_n|^2}{2\delta^2}.$$

To change $\frac{c}{2}\delta$ in 2δ it suffices to apply the relations between $F(\cdot, \cdot, \alpha\delta)$ and $F(\cdot, \cdot, \delta)$.

Finally, we extend \tilde{H} to a bounded plurisubharmonic function in Ω using the function ϑ_1 of the end of the proof of Theorem 5.1.

7.2. Example of non geometrically separated domain. The example presented here is the domain of \mathbb{C}^3 introduced by G. Herbort in [Her83]:

$$\Omega = \left\{ z \in \mathbb{C}^3 \text{ such that } \Re z_1 + |z_2|^6 + |z_3|^6 + |z_2|^2 |z_3|^2 < 0 \right\}.$$

Let $L_i^0 = \frac{\partial}{\partial z_i} + \beta_i \frac{\partial}{\partial \bar{z}_i}$, $i = 2, 3$, $\beta_2 = -\left(6|z_2|^4 \bar{z}_2 + 2|z_3|^2 \bar{z}_2\right)$ and $\beta_3 = -\left(6|z_3|^4 \bar{z}_3 + 2|z_2|^2 \bar{z}_3\right)$ so that (L_2^0, L_3^0) is a basis of $(1, 0)$ tangent vector fields in a neighborhood of the origin.

The fact that this domain is not geometrically separated at the origin is a consequence of the stronger following result:

Proposition 7.3. *For all real constants K and C , there exists $\delta_0 > 0$ such that for all δ , $0 < \delta \leq \delta_0$, there does not exist a basis (L_1^δ, L_2^δ) of $(1,0)$ tangent vector fields in a neighborhood of the origin of \mathcal{C}^6 norm bounded by C satisfying property EB_1 of Definition 3.1, for the constant K , at the origin.*

Proof. Let L be a $(1,0)$ tangent vector field in a neighborhood of the origin, and, with our usual notations, $F(L,0,\delta) = \sum_{\mathcal{L} \in \mathcal{L}_6(L)} \left| \frac{\mathcal{L}(\partial\rho)}{\delta} \right|^{2/|\mathcal{L}|} (0)$. We write $L = aL_2^0 + bL_3^0$.

Lemma. $F(L,0,\delta) \simeq \frac{|a(0)b(0)|}{\delta^{1/2}} + \left(\frac{1}{\delta}\right)^{1/3}$.

Proof. Because $c_{L\bar{L}} = 2[L, \bar{L}](\partial\rho)$, it is easy to see that:

- $c_{L\bar{L}}(0) = Lc_{L\bar{L}}(0) = \bar{L}c_{L\bar{L}}(0) = 0$;
- $LLc_{L\bar{L}}(0) = \bar{L}\bar{L}c_{L\bar{L}}(0) = 0$ and $L\bar{L}c_{L\bar{L}}(0) = \bar{L}Lc_{L\bar{L}}(0) = 4|a(0)b(0)|^2$;
- There exists a constant C_0 depending only of the \mathcal{C}^6 norm of a and b (i.e. of L) such that, if $|\mathcal{L}| = 3$, $|\mathcal{L}c_{L\bar{L}}(0)| \leq C_0|a(0)b(0)|$;
- There exists a constant α_0 depending only of the \mathcal{C}^6 norm of L , such that $F(L,0,\delta) \geq \alpha_0\delta^{-1/3}$. Indeed, the origin being of type 6, this follows from a result of T. Bloom [Blo81] and a compactness argument.

Then, the Lemma follows the fact that, for all $x \geq 0$, $\left(\frac{x}{\delta}\right)^{2/5} \leq \frac{x}{\delta^{1/2}} + \left(\frac{1}{\delta}\right)^{1/3}$. \square

We now finish the proof of the Proposition. Let δ be small enough. Suppose that there exists a (K, δ) - extremal basis at the origin, (L_1^δ, L_2^δ) , the \mathcal{C}^6 norms of the vector fields bounded by C . Let $L = \alpha L_1^\delta + \beta L_2^\delta$ and $L' = \alpha' L_1^\delta + \beta' L_2^\delta$ with $\alpha, \beta, \alpha', \beta' \in \mathbb{C}$ chosen so that $L(0) = L_2^0(0)$ and $L'(0) = L_3^0(0)$. Then, by extremality of (L_1^δ, L_2^δ) and the Lemma, we get

$$|\alpha|^2 F(L_1^\delta, 0, \delta) + |\beta|^2 F(L_2^\delta, 0, \delta) \simeq_K F(L, 0, \delta) \simeq_{C,K} \left(\frac{1}{\delta}\right)^{1/3}$$

and

$$|\alpha'|^2 F(L_1^\delta, 0, \delta) + |\beta'|^2 F(L_2^\delta, 0, \delta) \simeq_K F(L', 0, \delta) \simeq_{C,K} \left(\frac{1}{\delta}\right)^{1/3}.$$

Similarly, the extremality would imply

$$F(L+L', 0, \delta) \simeq_K |\alpha + \alpha'|^2 F(L_1^\delta, 0, \delta) + |\beta + \beta'|^2 F(L_2^\delta, 0, \delta) \lesssim_{C,K} \left(\frac{1}{\delta}\right)^{1/3}.$$

But the Lemma gives $F(L+L', 0, \delta) \simeq_C \frac{1}{\delta^{1/2}}$ which is a contradiction for δ small. \square

7.3. Additional remarks. Let Ω be geometrically separated at $p \in \partial\Omega$. In Definitions 3.3 and 3.4 we defined the pseudo-balls $B^c(p, \delta)$, $B_{\mathcal{C}}^c(p, \delta)$ and $B_{\text{exp}}^c(p, \delta)$, which are equivalent by Proposition 4.1, and we expressed the Bergman kernel at (p, p) with their volumes.

Let (z_i) be the coordinate system adapted to the extremal basis $(L_i)_{1 \leq i \leq n-1} = (L_i^{p,\delta})$ at p . $B(p, \delta)$ is defined (in the coordinate system) using only the directions of the extremal basis. Let us now define a new pseudo-ball using all the directions of the linear space generated by the vector fields L_i (i.e. the space E_0) (compare to the last point of Remark 4.2):

For $|Z| = 1$, $Z \in \mathbb{C}^n$, define $L_Z = \sum_{i=1}^{n-1} Z_i L_i + Z_n N$ and (in the coordinate system (z_i))

$$D_Z(p, \delta) = \left\{ \alpha Z \text{ such that } |\alpha| < cF(L_Z, p, \delta)^{-1/2} \right\}$$

and

$$D(p, \delta) = \bigcup_{|Z|=1} D_Z(p, \delta).$$

Then, property EB_1 of extremality for (L_i) , implies that these pseudo-balls $D(p, \delta)$ are equivalent (in the sense that they define the same structure of homogeneous space) to the previous ones. Indeed, if $z \in D_Z(p, \delta)$,

$$|z_i| = |\alpha z_i| \lesssim \left| Z_i \left(\sum |Z_i|^2 F(L_i, p, \delta) \right)^{-1/2} \right| \leq F(L_i, p, \delta),$$

and use Propositions 4.1 and 4.2. Conversely, if $z \in B^c(p, \delta)$, $z \neq 0$, and $Z = z/\|z\|$, then $F(L_Z, p, \delta) \simeq \sum \frac{|z_i|^2}{\|z\|^2} F(L_i, p, \delta) \leq \frac{nc^2}{\|z\|^2}$, thus $\|z\|^2 \lesssim nc^2 F(L_Z, p, \delta)$ and we conclude as before.

Note that this shows that, if Ω is completely geometrically separated at $p_0 \in \partial\Omega$ then the Bergman kernel $K(p, p)$ at a point p near p_0 is equivalent to the inverse of the volume of $D(p, \delta)$.

If Ω is not geometrically separated at p_0 choosing a coordinate system and a basis of tangent $(1,0)$ vector fields conveniently associated (in a sense to be defined), one can always define a ‘‘pseudo-ball’’ $D(p, \delta)$.

Let us do this, for example, for the domains considered in the previous Section, at the origin with the canonical coordinate system (z_i) and the vector fields L_i^0 (note that (z_i) is not adapted to the basis (L_i^0) in the sense of Definition 3.2 because, even if condition (3) is satisfied, the conditions on the derivatives of ρ are not).

A direct computation shows that, at the point $p_\delta = (-\delta, 0, 0)$, the volume of $D(p_\delta, \delta)$ is $\text{Vol}(D(p_\delta, \delta)) \simeq (\delta^3 \log(\frac{1}{\delta}))^{-1}$ (uniformly in δ), thus $B(p_\delta, \delta)$ and $D(p_\delta, \delta)$ are not equivalent, and the result of G. Herbot ([Her83]) shows that the Bergman kernel K of the domain satisfies $K(p_\delta, p_\delta) \simeq \text{Vol}(D(p_\delta, \delta))^{-1}$.

Then it is natural to ask if, for that example, the ‘‘pseudo-balls’’ $D(p, \delta)$ define a structure of homogeneous space. Unfortunately this is absolutely not the case. Indeed, in $D(0, \delta)$ consider the two points $p = (0, -\alpha\delta^{1/4}, 0)$ and $q = (0, 0, \alpha\delta^{1/4})$ (for α small enough, these points are in $D(0, \delta)$ for all δ , $0 < \delta \leq \delta_0$) and estimate a constant K so that $q \in D(p, K\delta)$. In the coordinate system centered at p , we have $q = (0, \alpha\delta^{1/6}, \alpha\delta^{1/6}) = \sqrt{2}\alpha\delta^{1/6} (0, 1/\sqrt{2}, 1/\sqrt{2})$; then calculating $c_{L\bar{L}}$ for $L = \frac{1}{\sqrt{2}}L_1^0 + \frac{1}{\sqrt{2}}L_2^0$ we see that

$$F(L, p, K\delta) \gtrsim \frac{\alpha^2 \delta^{-2/3}}{K} \text{ i.e. } F(L, p, K\delta)^{-1/2} \lesssim \sqrt{K} \delta^{1/3}.$$

Then q belongs to $D(p, K\delta)$ implies $K \gtrsim \delta^{-1/3}$.

8. APPENDIX

The following Lemma is an improvement of Lemma 3.9 of [CD06b]:

Lemma 8.1. *Let B_j be the unit ball in \mathbb{C}^j . Let K_1 be a positive real number, M and n two positive integers. There exists a constant $C(K_1)$ depending on K_1 , M and n such that, for $j = 1, \dots, n-1$, if g is a non negative function of class \mathcal{C}^M on B_j satisfying $\sup_{B_j} \{|D^{\alpha\beta} g(w)|, |\alpha + \beta| \leq M\} \leq K_1$, where $D^{\alpha\beta} = \frac{\partial^{|\alpha+\beta|}}{\partial w^\alpha \partial \bar{w}^\beta}$, then, for all $(\alpha^0, \beta^0) \in (\mathbb{N}^j)^2$, $|\alpha^0 + \beta^0| < M$, there exists $a \in \mathbb{N}^j$, $2|a| \leq |\alpha^0 + \beta^0|$ such that, denoting Δ^a the differential operator $\prod_{i=1}^j \Delta_i^{a_i}$, where $\Delta_i = \frac{\partial^2}{\partial z_i \partial \bar{z}_i}$ is the Laplacian in the z_i coordinate,*

$$\Delta^a g(0) \geq \frac{1}{C(K_1)} \left| D^{\alpha^0 \beta^0} g(0) \right|^{2|\alpha^0 + \beta^0|}.$$

Note that there is no absolute value in the left hand side of the inequality.

Proof. We only indicate how the proof of Lemma 3.9 of [CD06b] has to be modified.

Without loss of generality, we can suppose $|D^{\alpha^0 \beta^0} g(0)| = \max_{|\alpha+\beta|=|\alpha^0+\beta^0|} |D^{\alpha\beta} g(0)|$. By induction, it is enough to prove that there exist two constants c and C , depending on M and n , such that one of the following two cases holds:

First case there exists $a \in \mathbb{N}^j$, $2|a| = |\alpha^0 + \beta^0|$ such that $\Delta^a g(0) \geq c |D^{\alpha^0 \beta^0} g(0)|$;

Second case there exists $(\tilde{\alpha}, \tilde{\beta}) \in (\mathbb{N}^j)^2$, $|\tilde{\alpha} + \tilde{\beta}| < |\alpha^0 + \beta^0|$ such that

$$\left| D^{\tilde{\alpha} \tilde{\beta}} g(0) \right| \geq \frac{1}{C} \left| D^{\alpha^0 \beta^0} g(0) \right|^{-|\tilde{\alpha} + \tilde{\beta}| + |\alpha^0 + \beta^0| + 1}.$$

Let $p = |\alpha^0 + \beta^0|$, $\xi = \mu\varepsilon$, $\mu \in]0, 1[$, $\varepsilon = (\varepsilon_i)$, $|\varepsilon_i| \leq 1$, and, as in the proof of Lemma 3.9 of [CD06b] let us write Taylor formula:

$$\begin{aligned} g(\xi) &= \sum_{k=0}^{p-1} \mu^k \sum_{|\alpha+\beta|=k} *D^{\alpha\beta} g(0) \varepsilon^\alpha \bar{\varepsilon}^\beta + \mu^p \sum_{|\alpha+\beta|=p} *D^{\alpha\beta} g(0) \varepsilon^\alpha \bar{\varepsilon}^\beta + \mu^{p+1} R(\varepsilon, \mu) \\ &= A_1(\xi) + \mu^p A_2(\xi) + \mu^{p+1} R(\varepsilon, \mu), \end{aligned}$$

where $*$ are multinomial coefficients and $|R| \leq K_1 K_2$, K_2 depending only on M and n .

Remark now that, g being non negative,

(*) $\left\{ \begin{array}{l} \text{If there exists } \mu \simeq \left| D^{\alpha^0 \beta^0} g(0) \right| \text{ such that } A_2(\xi) + \mu R(\varepsilon, \mu) < -c_1 \left| D^{\alpha^0 \beta^0} g(0) \right|, \\ \text{case hold.} \end{array} \right. c_1 > 0$, then the *Second*

In the proof of Lemma 3.9 of [CD06b] we introduced a multi-index c ($|c| = p$), depending on g , and complex numbers ε_i ($\forall i, |\varepsilon_i| \geq c(M, n)$), depending on g and $K(M, n)$, such that

$$(8.1) \quad \sum_{\substack{|\alpha+\beta|=p \\ \alpha+\beta \neq c}} \left| *D^{\alpha\beta} g(0) \varepsilon^\alpha \bar{\varepsilon}^\beta \right| \leq \frac{\left| D^{\alpha^0 \beta^0} g(0) \right|}{K}$$

and

$$(8.2) \quad \left| \sum_{\alpha+\beta=c} *D^{\alpha\beta} g(0) \varepsilon^\alpha \bar{\varepsilon}^\beta \right| \geq 4 \frac{|D^{\alpha^0\beta^0} g(0)|}{K}.$$

To finish the proof, we show now that, either we can find ε and μ satisfying the hypothesis of (*), or we are in the *First case*.

We take $\mu = \frac{|D^{\alpha^0\beta^0} g(0)|}{KK_1K_2}$. Then $|A_2(\xi) + \mu R(\xi)| \geq \frac{|D^{\alpha^0\beta^0} g(0)|}{K}$ and $A_2(\xi) + \mu R(\xi)$ has the sign of $\sum_{\alpha+\beta=c} *D^{\alpha\beta} g(0) \varepsilon^\alpha \bar{\varepsilon}^\beta$.

If $\sum_{\alpha+\beta=c} *D^{\alpha\beta} g(0) \varepsilon^\alpha \bar{\varepsilon}^\beta < 0$, then (*) is satisfied, thus we consider the case where $\sum_{\alpha+\beta=c} *D^{\alpha\beta} g(0) \varepsilon^\alpha \bar{\varepsilon}^\beta > 0$.

If there is an index i such that c_i is odd, taking ε' defined by $\varepsilon'_j = \varepsilon_j$ if $j \neq i$ and $\varepsilon'_i = -\varepsilon_i$, then

$$\sum_{\alpha+\beta=c} *D^{\alpha\beta} g(0) \varepsilon'^\alpha \bar{\varepsilon}'^\beta \leq -\frac{4}{K} |D^{\alpha^0\beta^0} g(0)|,$$

and, by (8.1), (*) is verified.

So we suppose that for all i , $c_i = 2c'_i$, and we write

$$\sum_{\alpha+\beta=c} *D^{\alpha\beta} g(0) \varepsilon'^\alpha \bar{\varepsilon}'^\beta = \sum_{k=0}^{c_1} \varepsilon_1^k \bar{\varepsilon}_1^{c_1-k} A_k^1(\varepsilon'_2, \dots, \varepsilon'_n),$$

with $|\varepsilon'_i| = |\varepsilon_i|$, and choose $c \ll 4/K$. We separate two cases.

First suppose that $A_{c'_1}^1(\varepsilon_2, \dots, \varepsilon_n) \leq c |D^{\alpha^0\beta^0} g(0)|$. If $c_1 = 0$ then (8.2) implies

$$\sum_{\alpha+\beta=c} *D^{\alpha\beta} g(0) \varepsilon^\alpha \bar{\varepsilon}^\beta \leq -c' |D^{\alpha^0\beta^0} g(0)|$$

which gives (*). Thus suppose $c_1 \neq 0$. Let

$$\mathcal{E}_0 = \{\varepsilon', \text{ such that } \varepsilon'_i = \varepsilon_i, i > 1, \varepsilon'_1 = \vartheta \varepsilon_1, \text{ with } \vartheta^{c_1} = 1\}.$$

Thus

$$\sum_{\varepsilon' \in \mathcal{E}_0} \sum_{\alpha+\beta=c} *D^{\alpha\beta} g(0) \varepsilon'^\alpha \bar{\varepsilon}'^\beta = c_1 A_{c'_1}^1 |\varepsilon_1|^{c_1}.$$

Then, by (8.2), there exists $\varepsilon' \in \mathcal{E}_0$ such that

$$\sum_{\alpha+\beta=c} *D^{\alpha\beta} g(0) \varepsilon'^\alpha \bar{\varepsilon}'^\beta \leq -c'' |D^{\alpha^0\beta^0} g(0)|,$$

(recall $|\varepsilon_1| > c(M, n)$) and (*) is verified as before.

Suppose now $A_{c'_1}^1(\varepsilon_2, \dots, \varepsilon_n) > c' |D^{\alpha^0\beta^0} g(0)|$. Write

$$A_{c'_1}^1 = \sum_{k=0}^{c_2} \varepsilon_2^k \bar{\varepsilon}_2^{c_2-k} A_k^2(\varepsilon_3, \dots, \varepsilon_n).$$

As before, if $c_2 = 0$ or if $A_{c'_2}^2(\varepsilon_3, \dots, \varepsilon_n) \leq c''' |D^{\alpha^0\beta^0} g(0)|$ then we can change ε_2 such that we obtain

$$A_{c'_1}^1(\varepsilon'_2, \dots, \varepsilon'_n) \leq -c'''' |D^{\alpha^0\beta^0} g(0)|,$$

and we conclude that (*) is satisfied. If $A_{c'_2}^2(\varepsilon_3, \dots, \varepsilon_n) \geq c''' |D^{\alpha^0\beta^0} g(0)|$, we do another time the same thing, on the third variable. Then, by induction, if the process does not stop, the last step shows that if (*) is not satisfied, then the inequality on $D^{c'c'} g(0)$ implies that we are in the *First case*. \square

REFERENCES

- [AC99] H. Ahn and S. Cho, *In the mapping properties of the Bergman projection on pseudoconvex domains with one degenerate eigenvalue*, Complex Variable Theory Appl. **39** (1999), no. 4, 365–379.
- [BCD98] J. Bruna, Ph. Charpentier, and Y. Dupain, *Zeros varieties for the Nevanlinna class in convex domains of finite type in \mathbb{C}^n* , Ann. of Math. **147** (1998), 391–415.
- [Blo81] T. Bloom, *On the contact between complex manifolds and real hypersurfaces in \mathbb{C}^3* , Trans. Amer. Math. Soc **263** (1981), no. 2, 515–529.
- [Cat87] D. Catlin, *Subelliptic estimates for $\bar{\partial}$ -Neumann problem on pseudoconvex domains*, Annals of Math **126** (1987), 131–191.
- [Cat89] ———, *Estimates of invariant metrics on pseudoconvex domains of dimension two*, Math. Z. **200** (1989), 429–466.
- [CD92] P. Charpentier and Y. Dupain, *Une estimation des coefficients tangents d'un courant positif fermé dans un domaine de \mathbb{C}^3* , Publications Mathématiques **36** (1992), 319–349.
- [CD06a] ———, *Estimates for the Bergman and Szegő projections for pseudo-convex domains of finite type with locally diagonalizable Levi form*, Publications Mathématiques **50** (2006), no. 2, 413–446.
- [CD06b] ———, *Geometry of Pseudo-convex Domains of Finite Type with Locally Diagonalizable Levi Form and Bergman Kernel*, Jour. Math. Pures et Appl. **85** (2006), 71–118.

- [CG94] Der-Chen Chang and S. Grellier, *Estimates for the Szegő Kernel on Decoupled Domains*, Jour. of Math. Anal. and App. **187** (1994), 628–649.
- [Cho94] S. Cho, *Boundary behavior of the Bergman kernel function on some pseudoconvex domains in \mathbb{C}^n* , Trans. Amer. Math. Soc. **345** (1994), no. 2, 803–817.
- [Cho96] ———, *Estimates of the Bergman kernel function on certain pseudoconvex domains in \mathbb{C}^n* , Math Z. **222** (1996), 329–339.
- [Cho02a] ———, *Estimates of invariants metrics on pseudoconvex domains with comparable Levi form*, J. Math. Kyoto Univ. **42** (2002), no. 2, 337–349.
- [Cho02b] ———, *Estimates of the Bergman kernel function on pseudoconvex domains with comparable Levi form*, J. Korean Math. Soc. **39** (2002), no. 3, 425–437.
- [Cho03] ———, *Boundary behavior of the Bergman kernel function on pseudoconvex domains with comparable Levi form*, J. Math. Anal. Appl. **283** (2003), 386–397.
- [Chr88] M. Christ, *Regularity peoperties of the $\bar{\partial}_b$ equation on weakly pseudoconvex CR manifold of dimension 3*, J. Amer. Math. Soc. **1** (1988), 587–646.
- [Con02] M. Conrad, *Anisotrope optimale Pseudometriken für lineal konvex Gebeite von endlichem Typ (mit Anwendungen)*, PhD thesis, Berg.Universität-GHS Wuppertal (2002).
- [D'A82] J. P. D'Angelo, *Real hypersurfaces, orders of contact, and applications*, Annals of Math. **115** (1982), 615–637.
- [Der99] M. Derridj, *Régularité Hölderienne pour \square_b , sur des Hypersurfaces de \mathbb{C}^n , à Forme de Levi Décomposable en Blocs*, J. Geom. Anal. **9** (1999), no. 4, 627–652.
- [DF99] K. Diederich and J. E. Fornæss, *Support functions for convex domains of finite type*, Math. Z. **230** (1999), 145–164.
- [DF03] ———, *Lineally convex domains of finite type: holomorphic support functions*, Manuscripta Math. **112** (2003), 403–431.
- [DF06] K. Diederich and B. Fischer, *Hölder estimates on lineally convex domains of finite type*, Michigan Math. J. **54** (2006), no. 2, 341–452.
- [FK88] C. L. Fefferman and J. J. Kohn, *Hölder estimates on domains of complex dimension two and on three dimensional CR manifold*, Adv. Math. **69** (1988), 233–303.
- [FKM90] C. L. Fefferman, J. J. Kohn, and M. Machedon, *Hölder Estimates on CR Manifold with a diagonalizable Levi Form*, Advances in Math **84** (1990), no. 1, 1–90.
- [GS85] R. Gay and A. Sebbar, *Division et extension dans l'algèbre $A^\infty(\Omega)$ d'un ouvert pseudoconvexe à bord lisse de \mathbb{C}^n* , Math. Z. **189** (1985), 421–447.
- [Hef04] T. Hefer, *Extremal bases and Hölder Estimates for $\bar{\partial}$ on Convex Domains of Finite Type*, Michigan Math. J. **52** (2004), 573–602.
- [Her83] G. Herbort, *Logarithmic growth of the Bergman kernel for weakly pseudoconvex domains in \mathbb{C}^n of finite type*, Manuscripta Math. **45** (1983), no. 1, 69–76.
- [Kan90] H. Kang, *An Approximation Theorem for Szegő Kernels and Applications*, Michigan Math. J. **37** (1990), 447–458.
- [Ker72] N. Kerzman, *The Bergman kernel function. Differentiability at the boundary*, Math. Ann. **195** (1972), 149–158.
- [Kis98] C. O. Kiselman, *A differential inequality characterizing weak lineal convexity*, Math. Annalen **311** (1998), 1–10.
- [KN65] J. J. Kohn and L. Nirenberg, *Non coercive boundary value problems*, Comm. Pure Appl. Math. **18** (1965), 443–492.
- [Koe02] K. D. Koenig, *On maximal Sobolev and Hölder estimates for the tangential Cauchy-Riemann operator and boundary Laplacian*, Amer. J. Math. **124** (2002), 129–197.
- [Koe07] ———, *Comparing the Bergman and Szegő projections on domains with subelliptic boundary Laplacian*, Math. Annalen **339** (2007), 667–693.
- [Koh85] J. J. Kohn, *Estimates for $\bar{\partial}_b$ on Pseudoconvex CR Manifolds*, Proc. Symposia Pure Math. **43** (1985), 207–217.
- [Mac88] M. Machedon, *Szegő kernels on pseudoconvex domains with one degenerate eigenvalue*, Ann. of Math. **128** (1988), 619–640.
- [McN89] J. McNeal, *Boundary behavior of the Bergman kernel function in \mathbb{C}^2* , Duke Math. Journal **58** (1989), no. 2, 499–512.
- [McN91] ———, *Local geometry of decoupled pseudoconvex domains*, Aspekte der Math. **E17** (1991), 223–230.
- [McN94] ———, *Estimates on Bergman Kernels of Convex Domains*, Advances in Math (1994), 108–139.
- [McN02] ———, *Uniform subelliptic estimates on scaled convex domains of finite type.*, Proc. Amer. Math. Soc. **130** (2002), no. 1, 39–47.
- [MS94] J. D. McNeal and E. M. Stein, *Mapping properties of the Bergman projection on convex domains of finite type*, Duke Math. J. **73** (1994), no. 1, 177–199.
- [MS97] ———, *The Szegő projection on convex domains*, Math. Z. **224** (1997), no. 4, 519–553.
- [NRSW89] A. Nagel, J. P. Rosay, E. M. Stein, and S. Wainger, *Estimates for the Bergman and Szegő kernels in \mathbb{C}^2* , Annals of Math. **129** (1989), 113–147.
- [NSW85] A. Nagel, E. Stein, and S. Wainger, *Balls and metrics defined by vectors fields: basic properties*, Acta Mathematica **155** (1985), 103–147.

PHILIPPE CHARPENTIER & YVES DUPAIN: UNIVERSITÉ BORDEAUX 1, INSTITUT DE MATHÉMATIQUES, 351, COURS DE LA LIBÉRATION, 33405 TALENCE, FRANCE

E-mail address: philippe.charpentier@math.u-bordeaux1.fr, yves.dupain@math.u-bordeaux1.fr